

H. NOUVEL

Report on the Crustacea. By the Rev. A. Merle Norman, M.A.

One hundred and one Crustacea were obtained. To this number two or three more Entomostraca may yet be added, when the gatherings of the drift net shall have been more minutely examined. The sectional distribution of the species may be thus shown:—

Brachyura . . . 9.	Amphipoda . . . 34.
Anomoura . . . 7.	Isopoda . . . 5.
Macroura . . . 7.	Entomostraca . . 16.
Stomapoda . . . 8.	Pycnogonoidea . . 10.
	Cirripedia . . . 5.

The object of the present paper is to give a catalogue of the species obtained; a few notes on the rarer forms, and descriptions of the new Stomapods. The description of the new Amphipoda and Entomostraca must be deferred for the present. Mr. Hodge, who is especially devoting himself to the study of the Pycnogonoidea, will report upon the species of that very interesting order.

In the following table the three columns represent the same localities as those in the report on the Mollusca:—

<i>Stenorhynchus rostratus</i> , Lin.	r.c.		
<i>Inachus Dorsettensis</i> , Penn.	r.		A few young specimens obtained off Berwick Bay. New to our local Fauna.
— <i>dorhynchus</i> , Leach	r.		One small example only.
<i>Hyas coarctatus</i> , Leach	c. c. c.		The most abundant of the larger Crustacea, in deep water. Specimens small.
<i>Portunus holsatus</i> , Fabr.	r.c.	r.c.	
— <i>pusillus</i> , Leach	r.	r.	
<i>Ebalia tuberosa</i> , Penn.	r.c.	r.c.	
— <i>Cranchii</i> , Leach	r.	r.	
<i>Atelecyclus septemdentatus</i> , Mont.		r.	Only very young.
<i>Pagurus Bernhardus</i> , Lin.	c. c. c.		
— <i>Hyndmanni</i> , Thompson	v.r.		One only, east of Tynemouth.
— <i>laevis</i> , Thompson	r.c. c. c.		Sandy ground.

1. Colonne . . . off Tynemouth
 2. — . . . Dogger Bank
 3. — . . . Coquet and Berwick Bay

<i>Pagurus Thompsoni</i> , Bell	a. c. c.	With the last, and also on harder ground.
<i>Porcellana longicornis</i> , Penn.	r.	
<i>Galathea dispersa</i> , Bate	r.c. c.	Widely diffused.
— <i>Andrewsii</i> , Kinahan	r. c.	
<i>Crangon Altmanni</i> , Kinahan	c. c. c.	Abundant.
— <i>spinosus</i> , Leach	v.r. r.c.	A very large example taken off Berwick.
— <i>bispinosus</i> , Westwood	v.r.	Two specimens of this rare species dredged in 40 fathoms, 40-50 miles east of Tynemouth.
<i>Hippolyte pusiola</i> , Kroyer	v.c. r.	
— <i>securifrons</i> , Norman	r. r.c.	
<i>Pandalus annulicornis</i> , Leach	c. c. c.	Abundant everywhere.
— <i>Thompsoni</i> , Bell	c. r.c.	
<i>Myas spiritus</i> , Norman	v.c. c. r.	
— <i>Didelphys</i> , n. sp.	v.c.	A new species.
<i>Diastylis Rathkii</i> , Kroyer	r. r.	
<i>Vaunthompsonia cristata</i> , Bate	r.	50-60 miles east of Tynemouth.
— <i>rosea</i> , n. sp.	v.r.	A new species.
<i>Eudora truncatula</i> , Bate	r.	50-60 miles east of Tynemouth, on a muddy bottom.
<i>Cyrianassa ciliata</i> , n. sp.	v.r.	A new species.
— <i>elegans</i> , n. sp.	v.r.	A new species.
<i>Montagna Alderii</i> , Bate	r.	40-50 miles off Tynemouth.
— <i>pollexiana</i> , Bate	c. r.c.	
<i>Callisoma crenata</i> , Bate	v.r.	One specimen, 7 miles off Tynemouth, 25 fathoms, and a second off Berwick.
<i>Anonyx Edwardsii</i> , Kroyer	r. v.r.	40 and 100 miles east of Tynemouth.
— <i>ampulla</i> , Kroyer	v.r.	Only a single example.
— <i>denticulatus</i> , Bate	r.c. r.	
— <i>longipes</i> , Bate	v.r.	One specimen only.
<i>Ampelisca Belliana</i> , Bate	c. r. r.c.	
— <i>Gaimardi</i> , Kroyer	c. c. c.	
<i>Haploops tubicola</i> , Lillj.	v.r. v.r.	A genus new to Britain.
<i>Phoxus plumosus</i> , Holbüll	r. r.	
<i>Edicerus parvimanus</i> , B. and W.	v.r.	

