

XXIII. *A Catalogue of Land, Fresh-Water, and Marine Crustacea found in the Basin of the River Forth and its Estuary.* By THOMAS SCOTT, LL.D., F.L.S., Mem. Soc. Zool. de France, etc.

(Read 26th March 1906.)

PART II.—THE OSTRACODA, COPEPODA, AND CIRRIPIEDIA.

(Part I., published in the *Proceedings*, Vol. XVI. pp. 97-190, contained the MALACOSTRACA, CLADOCERA, and BRANCHIURA.)

Order OSTRACODA.

The work that has been mainly followed in the arrangement and nomenclature of the Ostracoda is that by Professor G. S. Brady and the Rev. Canon A. M. Norman, entitled "A Monograph of the Marine and Fresh-Water Ostracoda of the North Atlantic and North-Western Europe." The work was published in two separate portions in the scientific *Transactions of the Royal Dublin Society*. The first portion, which formed Part II. of Vol. IV. (Series 2), appeared in 1889, and the second, which formed Part II. of Vol. V. of the same series, in 1896.

In this work the Ostracoda are divided into four sections, viz.—(1st) the Podocopa, which comprises all the fresh-water and most of the marine species; (2nd) the Myodocopa, which is represented in this Catalogue by only two species, *Asterope Mariæ* and *Philomedes interpuncta*; (3rd) the Cladocopa, which is represented by *Polycope orbicularis* and *Polycopsis compressa*; and (4th) the Platycopa—a section not represented in this Catalogue.

As this work has been, with few exceptions, followed throughout, I have not considered it necessary to mention it under every species, and where it is mentioned it is referred to as the Monograph (or *l. c.*), Part I. or Part II., as the case may be.

Various other works have been consulted, two of which

may be mentioned here, viz., one by Dr G. W. Müller on "The Fresh-Water Ostracoda of Germany," and the other by Dr A. Kaufmann on "The Fresh-Water Ostracoda of Switzerland."

Section I. PODOCOPA.

Family C Y P R I D I D Æ.

Genus (1) *Cypria*, Zenker, 1854.

1. *Cypria exculpta* (S. Fischer).

1855. *Cypris exculpta*, Fischer, Beitrag zur Kenntniss der Ostracoden, Abhdlg. der math.-phys. Klasse der k. bayr. Akad. d. Wiss., Bd. vii. p. 652, taf. xix. figs. 36-38.

This species, though widely distributed, is not very common in the district under consideration. It has been obtained in Loch Leven,¹ Lochgelly Loch, Lochs Lurg and Dow, Kinross; Loch Achray, Perthshire; and Goldenhoof Dam, near Howietoun, Stirlingshire. Loch Ard, July 1906 (W. Evans).

2. *Cypria ophthalmica* (Jurine).

1820. *Monoculus ophthalmicus*, Jurine, Hist. des Monocles, p. 178, taf. xix. figs. 16, 17.

This is one of the most common and widely distributed species of our fresh-water Ostracoda.

Genus (2) *Cyclocypris*, G. S. Brady and A. M. Norman, 1889.

3. *Cyclocypris globosa* (G. O. Sars).

1863. *Cypris globosa*, G. O. Sars, Om en i Somm., 1862, foretagen Zool. Reise i Christianias og Trondhjems Stifter, p. 27.

Has been obtained in a pond near Musselburgh in August 1894; Loch Leven, Loch Fitty, Black Loch, Loch Dow, Loch Katrine, Loch Achray, Loch Vennachar, and Loch Coulter. Teith at Callander (Evans).

¹ It will be understood that the Loch Leven mentioned in this Catalogue is Loch Leven, Kinross.

4. *Cyclocypris serena* (Koch).1838. *Cypris serena*, Koch, Deutschlands Crustaceen, Heft xxi. p. 22.1896. *Cyclocypris serena*, B. and N., *l. c.*, Part II. (Appendix) p. 718.

A common and widely distributed species. It has been found in Loch a Chroin, at an altitude of 2500 feet, by Mr Evans.

5. *Cyclocypris lævis* (O. F. Müller).1896. *Cyclocypris lævis*, Brady and Norman, *l. c.*, Part II. (Appendix) p. 718.

This, which is not such a common species as the last, has an elliptical instead of an ovate form when looked at from above. It has been observed in Duddingston Loch, Lochgelly Loch, pools on Luffness Links, etc.

Dr G. W. Müller appears to think that the species referred by G. S. Brady and Norman to *Cyclocypris serena*, Koch, is the true *C. lævis* of O. F. Müller, and that their *C. lævis* is probably identical with a form described by Croneberg under the name of *Cyclocypris pygmæa*.¹

Genus (3) *Cypris*, O. F. Müller, 1785.6. *Cypris fuscata*, Jurine.1820. *Cypris fuscata*, Jurine, Hist. des Monocles de Geneva, p. 174, pl. xix. figs. 1-16.

Hab.—Duddingston Loch; Upper Elf Loch, Braid Hills; Linlithgow Loch; Loch Leven. Roslin Curling Pond, March 1906 (W. Evans).

7. *Cypris incongruens*, Ramdohr.1806. *Cypris incongruens*, Ramdohr, Mag. der Gesells. naturf. Freunde in Berlin, Bd. ii. p. 86, taf. iii. figs. 1-12 *et seq.*

Hab.—Abundant in pools in the brickfield at Portobello in the autumn of 1888. In pools on May Island in September 1890. Common in pools on the side of the Union Canal at

¹ *Scottia Browniana* (Jones), which up till 1887, when living specimens were obtained near Rothesay, was only known as a fossil, was collected in a post-Tertiary deposit at Elie, Fifeshire, by the late James Bennie of the Geological Survey (cf. *Proc. Roy. Phys. Soc. Edin.*, vol. x. pp. 339-341, 1890-91). This form, which is placed between *Cyclocypris* and *Cypris*, has not yet been found *living* on the east side of Scotland.

Kingsknowe in May 1888. Pond at House-o'-Hill Farm, Midlothian, common, July 1900 (Dr and Miss Sprague).

8. *Cypris virens*, Jurine.

1820. *Cypris virens*, Jurine, Hist. des Monocles, p. 174, taf. xviii. figs. 15, 16.

Hab.—Duddingston Loch; pools on Luffness Links; pools at Slateford; Kilconquhar Loch; Loch Leven. Marchfield Pond, July 1900, very common (Dr and Miss Sprague). Roslin Curling Pond, March 1906 (Evans).

9. *Cypris affinis*, Fischer.

1851. *Cypris affinis*, Fischer, Ueber das genus *Cypris*, p. 32, pl. x. figs. 9-11.

1889. ,, *reticulata*, Brady and Norman, *l. c.*, Part I. p. 76, pl. viii. figs. 1, 2; pl. xi. figs. 5-7.

1900 ,, *affinis*, Kaufmann, Cypriden u. Darwinuliden der Schweiz, Revue Suisse de Zool., T. 8, p. 272, taf. xv. figs. 17-20 *et seq.*

Hab.—Camilla Loch, Fifeshire, and Duddingston Loch. This species does not appear to be very common. (I have recorded *C. affinis* from Linlithgow Loch, but the dissections show that the specimens belong to *C. fuscata*.)

10. *Cypris obliqua*, G. S. Brady.

1868. *Cypris obliqua*, Brady, Mon. rec. Brit. Ostrac., p. 364, pl. xxiii. figs. 33-38.

Hab.—Lurg Loch (near L. Glow), Kinross-shire; Lochgelly Loch and Loch Achray. Craigleith Quarry, near Edinburgh, February 1901, one specimen (Dr and Miss Sprague).

Genus (4) **Eurycypris**, G. W. Müller, 1898.

11. *Eurycypris pubera* (O. F. Müller).

1785. *Cypris pubera*, O. F. Müller, Entomostraca, p. 56, taf. v. figs. 1-5.

1900. *Eurycypris pubera*, G. W. Müller, Deutschl. Süßw.-Ostrac., p. 54, taf. xiii. figs. 1-11.

Hab.—Duddingston Loch (Baird, Norman, and myself); Kilconquhar Loch; Kinghorn Loch (common); Lochgelly

Loch. Townhill Loch, Dunfermline (D. Robertson). Otterston Loch, Fife, October 1905 (W. Evans).

Genus (5) *Cyprinotus*, G. S. Brady, 1886.

12. *Cyprinotus prasina* (Fischer).

1889. *Cypris prasina*, Brady and Norman, Monograph, Part I. p. 78.

1896. *Cyprinotus prasina*, id., Part II. (Appendix) p. 772.

Hab.—Pools in an old brickfield at Seafield, near Dunbar, August 1890; in pools at the mouth of the Peffer Burn, near Aberlady, August 1893. *Cyprinotus prasinus* was obtained by the late James Bennie, of the Geological Survey, in material excavated from the bed of the Old "Nor Loch," Edinburgh (cf. *Proc. Roy. Phys. Soc. Edin.*, vol. x. p. 139, 1889).

Genus (6) *Herpetocypris*, G. S. Brady and A. M. Norman, 1896.

Erpetocypris, idem, 1889.

13. *Herpetocypris reptans* (Baird).

1835. *Candona reptans*, Baird, Trans. Berw. Nat. Club, vol. i. p. 99, pl. iii. fig. 11.

Hab.—Duddingston Loch, Loch Leven, Linlithgow Loch, Loch Lubnaig, pools on Gullane Links, and various other places throughout the district. Loch Ard (Evans).

14. *Herpetocypris strigata* (O. F. Müller).

1785. *Cypris strigata*, O. F. Müller, Entomostraca, p. 54, pl. iv. figs. 4-6.

Hab.—Duddingston Loch, Loch Leven, Black Loch near L. Glow; pools at Kingsknowe near Edinburgh, May 1888; Loch Coulter; and about Howietoun, Stirlingshire, 1889.

15. *Herpetocypris tumefacta*, G. S. Brady and D. Robertson.

1870. *Cypris tumefacta*, B. and R., Ann. and Mag. Nat. Hist. (4), vol. vi. p. 13, pl. iv. figs. 4-16.

Hab.—Duddingston Loch, Loch Leven, pools on Luffness Links, Loch Achray, Loch Vennachar, Loch Coulter, and pools near Howietoun, etc. Ben Ledi, at 2500 feet (Evans).

Genus (7) *Ilyodromus*, G. O. Sars, 1894.16. *Ilyodromus olivacea* (G. S. Brady and A. M. Norman).

1889. *Erpetocypris olivacea*, B. and N., *l. c.*, Part I. p. 89, pl. viii. figs. 3, 4.

1896. *Ilyodromus olivacea*, *l. c.*, Part II. (Appendix) p. 724.

Hab.—Duddingston Loch, Kinghorn Loch, Black Loch near L. Glow.

17. *Ilyodromus Robertsoni*, G. S. Brady and A. M. Norman.

1889. *Erpetocypris Robertsoni*, B. and N., *l. c.*, Part I. p. 88 (with two text figures).

1896. *Ilyodromus Robertsoni*, *l. c.*, Part II. (Appendix) p. 724.

Hab.—Linlithgow Loch; Black Loch near L. Glow, 14th September 1889.

Genus (8) *Cypridopsis*, G. S. Brady, 1867.18. *Cypridopsis villosa* (Jurine).

1820. *Monoculus villosus*, Jurine, *Hist. des Monocles*, p. 178, pl. xix. figs. 14, 15.

1868. *Cypridopsis villosa*, Brady, *Mon. rec. Brit. Ostrac.*, p. 377, pl. xxiv. figs. 11-15; pl. xxxvi. fig. 9.

Hab.—Duddingston Loch, Loch Leven, and in various other lochs and ponds throughout the district

19. *Cypridopsis aculeata* (Lilljeborg).

1853. *Cypris aculeata*, Lillj., *De Crust. ex Ord. trib.*, p. 117, taf. xi. figs. 15, 16.

1868. *Cypridopsis aculeata*, Brady, *l. c.*, p. 376, pl. xxiv. figs. 16-20; pl. xxxvi. fig. 10.

Hab.—Pools in an old brickfield at Seafield, near Dunbar; Gullane, near Aberlady, August 1893. Cramond Island, July 1901 (Dr and Miss Sprague).

Genus (9) *Pionocypris*, G. S. Brady and A. M. Norman, 1896.20. *Pionocypris vidua* (O. F. Müller).

1785. *Cypris vidua*, O. F. Müller, *Entomotraca*, p. 55, taf. iv. figs. 7-9.

1889. *Cypridopsis vidua*, B. and N., *l. c.*, Part I. p. 89.

1896. *Pionocypris vidua*, *id.*, *ibid.*, Part II. (Appendix) p. 726.

Hab.—Duddingston Loch, Loch Leven, Loch Coulter, Loch

Achray; Loch Rusky (Evans); and generally throughout the district, especially in the smaller bodies of water.

Objection is taken to the separation of this species from the genus *Cypridopsis*, to which it was ascribed by Dr G. S. Brady in 1867, but, as pointed out by Brady and Norman in the Appendix to their valuable *Monograph*, pp. 725 and 726, its removal became necessary when it was found to differ materially in the principal character on which the genus was founded, viz., the caudal rami. These were described as "being quite rudimentary, consisting of two slender setiform processes springing from a common base." Careful examination has shown that while *C. villosa*, *C. aculeata*, and one or two others agree so far with this definition, the caudal rami in *C. vidua* consists of four setiform processes, and also that the basal part of each pair of processes differs to some extent. Dr Kaufmann does not appear to attach much value to this difference, and therefore replaces *C. vidua* under *Cypridopsis*, but he at the same time removes *the others* to a new genus—*Cypridopsella*—the chief distinguishing character of which is that while the first has five the other has only two hairs on the branchial plate of the first maxilla.¹ Though *C. vidua* was made the type of the genus *Cypridopsis* by Dr Brady, it was doubtless because he considered its furca to be similar to that of the two species associated with it, and as these two species agree with the definition of *Cypridopsis*, I think *they* ought to be retained in that genus, and *C. vidua* removed, as has been done by Brady and Norman.

Genus (10) *Potamocypris*, G. S. Brady, 1870.

21. *Potamocypris fulva*, G. S. Brady.

1868. *Bairdia fulva*, Brady, Mon. rec. Brit. Ostrac., p. 474, pl. xxviii. fig. 21.

Hab.—Duddingston Loch, Loch Leven, Raith Lake, Loch Gelly, and several other places. Ben Ledi, at 2500 feet (Evans).

¹ *Cypridopsis*—Branchial platte des Kiefer fusses mit fünf Borsten.
Furka verkümmert mit geisselförmiger Borste.

Cypridopsella—Branchial platte des Kiefer fusses mit zwei Borsten.
Furke verkümmert mit geisselförmiger Borste.

Genus (14) *Candona*, Baird, 1845.25. *Candona candida* (O. F. Müller).

1785. *Cypris candida*, Müller, Entomostraca, p. 62, taf. vi. figs. 7-9.

Hab.—This appears to be a moderately common species throughout the whole district.

26. *Candona neglecta*, G. O. Sars.

1887. *Candona neglecta*, G. O. Sars, Mittelh. Invert.—fauna IV., Ostrac. Mediterranea, p. 279, taf. xv. figs. 5-7; taf. xix. figs. 1-21.

1889. „ *candida* (Part), B. and N., *l. c.*, Part I. p. 99, pl. x. figs. 20, 21.

Hab.—I find this species generally distributed but not very plentiful, and the males appear to be more frequent and larger than the females. The following are some of the places where it has been obtained:—Duddingston Loch (♂); Loch Leven (♂); Linlithgow Loch (♀); Cocklemill Burn, near Largo (♂ and ♀); Lochgelly Loch (♂); Camilla Loch (♂ and ? ♀); Kinghorn Loch (♂).

27. *Candona lactea*, Baird.

1850. *Candona lactea*, Baird, Proc. Zool. Soc. Lond. (1850), p. 255, pl. xviii. figs. 25-27.

Hab.—Duddingston Loch, Loch Leven, Camilla Loch, Linlithgow Loch, Loch Coulter, and other places within the area.

28. *Candona compressa* (S. Fischer).

1851. *Cypris compressa*, Fischer, Abhandl. über das genus *Cypris*, Mém. des Sav. étrang. des Sci. de St. Pétersb., t. 7, p. 144, taf. ii. figs. 7-12; taf. iii. figs. 1-5.

1889. *Candona pubescens*, B. and N., *l. c.*, Part I. p. 101, pl. xii. figs. 32-37.

1900. „ *compressa*, Kaufmann, Cyprid. u. Darwinul., p. 371, taf. xxvii. figs. 4-6 *et seq.*

Hab.—Duddingston Loch (Brady and Norman), Lochgelly Loch, Loch Leven, and several other places within the district.

29. *Candona stagnalis*, G. O. Sars.

1890. *Candona stagnalis*, G. O. Sars, Christ. Videns-Selsk. Forhandl.,
p. 69.

1891. ,, *ambigua*, T. Scott, Ninth Rept. F. B. S., pt. iii.
p. 277, pl. iv. fig. 7 a-c.

Hab.—Lochgelly Loch and Loch Fitty, Fifeshire; not common.

30. *Candona rostrata*, Brady and Norman.

1889. *Candona rostrata*, B. and N., *l. c.*, Part I. p. 101, pl. ix.
figs. 11, 12, 12a, b; pl. xii. figs. 22-31.

Hab.—Duddingston Loch, Lurg Loch, Raith Lake, pond at Howietoun; not common. The larva of the tapeworm, *Tænia gracilis*, has been found encysted in a specimen of this species collected in Duddingston Loch.¹

31. *Candona fabæformis* (Fischer).

1851. *Cypris fabæformis*, Fischer, Über das genus *Cypris*, p. 146,
pl. iii. figs. 6-16.

Hab.—Duddingston Loch; marl-pit, near Davidson's Mains, Edinburgh; pools on Luffness Links; Loch Fitty, and other places in Fifeshire; Linlithgow Loch; and in a pool near Howietoun.

32. *Candona acuminata* (Fischer).

1851. *Cypris acuminata*, Fischer, Über das genus *Cypris*, p. 148,
taf. iv. figs. 12-16.

Hab.—Ditch beside Harelaw Dam, Balerno; Threipmuir Reservoir; Loch Fitty; ponds at Howietoun; and Loch Coulter.

The specimens recorded here agree closely with the female of *Candona caudata*, figured by Kaufmann in *Cypriden u. Darwinuliden der Schweiz*, p. 365, taf. xxiv. figs. 16-20; taf. xxvi. figs. 17-23.

¹ *Proc. Roy. Phys. Soc.*, vol. x. p. 314 (with text-figure), 1890-91.

33. *Candona hyalina*, G. S. Brady and D. Robertson.

1870. *Candona hyalina*, B. and R., Ann. and Mag. Nat. Hist. (4),
vol. vi. p. 18, pl. ix. figs. 5-8; pl. v.
figs. 4-11.

1889. „ „ B. and N., *l. c.*, Part I. p. 247, with text-
figures 1-5.

Hab.—Threipmuir Reservoir, near Balerno (♂); Loch Fitty and Loch Gelly, Fifeshire; Loch Dow, Kinross-shire (♀); Loch Katrine. The shell is much compressed, and of a white colour in living specimens. The dorsal edge is obtusely angular.

Genus (15) *Candonopsis*, Vavra, 1891.34. *Candonopsis Kingsleii* (Brady and Robertson).

1870. *Candona Kingsleii*, B. and R., Ann. and Mag. Nat. Hist. (4),
vol. vi. p. 17, pl. ix. figs. 9-12.

1891. *Candonopsis kingsleii*, Vavra, Mon. der Ostrac. Bohmens,
p. 54, fig. 16.

Hab.—Duddingston Loch, pools on Luffness Links, Loch Leven, Camilla Loch, Loch Coulter, Loch Katrine, and others. This is a well-marked and widely distributed species. The shell in living specimens is white and much compressed.

Genus (16) *Paracandona*, Hartwig, 1899.35. *Paracandona euplectella* (Robertson).

1880. *Candona euplectella*, Robertson, Fresh and Brackish-Water
Ostrac. of Clydesdale, p. 23. (See also Proc. Nat. Hist.
Soc. Glasgow, vol. iv. pt. i., 1880, p. 77.)

1900. *Paracandona euplectella*, G. W. Müller, Deutschl. Süßw.-
Ostrac., p. 37, taf. ix. figs. 1-9, 14.

Hab.—Loch Dow, near Loch Glow, Kinross-shire, 14th September 1889. This is still the only record I have of *P. euplectella* from the district under consideration. It is the most beautiful of the species belonging to the British *Candonæ*. The shell is cylindrical and very tumid, and its structure is suggestive of the glass-sponge *Euplectella*, as implied by the specific name.

Genus (17) *Ilyocypris*, Brady and Norman, 1889.

36. *Ilyocypris gibba* (Ramdohr).

1808. *Cypris gibba*, Ramdohr, Mag. der Gesells. naturf. Freunde in Berlin, ii. p. 91, taf. iii. figs. 13-17.

Hab.—Union Canal, near Edinburgh, 4th September 1889. I obtained a number of specimens here—the only place where I have observed it within the district. The shell of this species is ornamented with prominent tubercles.

37. *Ilyocypris bistrigata* (Jurine).

1820. *Monoculus bistrigatus*, Jurine, Hist. Nat. des Monocles, p. 177, pl. xix. figs. 12, 13.

1838. *Cypris biplicata*, Koch, Deutschlands Crustaceen, H. 21 (161), fig. xvi.

Hab.—Duddingston Loch, Loch Leven, etc. This species is generally distributed throughout the district. I have occasionally observed specimens distinctly larger than some of the others, but could find no other difference sufficient to permit of their separation even as a variety.

Genus (18) *Pontocypris*, G. O. Sars, 1865.

38. *Pontocypris mytiloides* (Norman).

1862. *Cythere mytiloides*, Norman, Ann. and Mag. Nat. Hist. (3), vol. ix. p. 50, pl. iii. figs. 1-3.

Hab.—Off Aberlady, off Musselburgh, and various other parts of the Forth estuary. Living specimens are usually moderately dark in colour.

39. *Pontocypris acupunctata*, G. S. Brady.

1866. *Pontocypris acupunctata*, Brady, Brit. Assoc. Report (1866), p. 209.

1868. „ „ Brady, Mon. rec. Brit. Ostrac., p. 386, pl. xxiv. figs. 53-56.

Hab.—Off St Monans and one or two other places in the estuary, 1890; rare. Living specimens are brownish in colour, and the surface is minutely punctate.

40. *Pontocypris trigonella*, G. O. Sars.

1865. *Pontocypris trigonella*, G. O. Sars, Oversigt af Norges mar. Ostrac., p. 16.

Hab.—Largo Bay and other parts of the estuary, but not very common.

Genus (19) *Argillœcia*, G. O. Sars, 1865.

41. *Argillœcia cylindrica*, G. O. Sars.

1865. *Argillœcia cylindrica*, G. O. Sars, *l. c.*, p. 18.

Hab.—Firth of Forth (Brady and Robertson). Off St Monans, rare.

Family BAIRDIIDÆ.

Genus (20) *Bairdia*, M'Coy (? 1849).

42. *Bairdia inflata* (Norman).

1862. *Cythere inflata*, Norman, Ann. and Mag. Nat. Hist. (3), vol. ix. p. 49, pl. iii. figs. 6-8.

1868. *Bairdia inflata*, Brady, Mon. rec. Brit. Ostrac., p. 388, pl. xxvii. figs. 9-17; pl. xxxviii. fig. 5.

Hab.—Dredged off St Monans, 1889; rare. The specimens were of a brownish colour, and tumid, as indicated by the name.

Family DARWINULIDÆ.

Genus (21) *Darwinula*, Brady and Robertson, 1885.

43. *Darwinula Stevensoni*, Brady and Robertson.

1870. *Polycheles Stevensoni*, B. and R., Ann. and Mag. Nat. Hist. (4), vol. vi. p. 25, pl. vii. figs. 1-7; pl. ii. figs. 4-14.

Hab.—Loch Lubnaig, Perthshire, rare; obtained near the east shore by hand-net on 29th September 1894. This somewhat rare species was obtained by the late James Bennie in a lacustrine deposit in the Meadows, Edinburgh. (Cf. "The Ancient Lakes of Edinburgh," by James Bennie and T. Scott, *Proc. Roy. Phys. Soc. Edin.*, vol. x. p. 131.)

Family CYTHERIDÆ.

Genus (22) *Cythere*, O. F. Müller, 1785.

44. *Cythere lutea*, O. F. Müller.

1785. *Cythere lutea*, Müller, Entomostraca, p. 65, taf. vii. figs. 3, 4.

Hab.—Forth estuary, especially inshore—a common British species.

45. *Cythere pellucida*, Baird.1850. *Cythere pellucida*, Baird, Brit. Entom., p. 173, pl. xxi. fig. 7.

Hab.—Cramond Island, taken with hand-net at low water in 1889, and subsequently in other parts of the estuary, chiefly in brackish water.

46. *Cythere confusa*, Brady and Norman.1865. *Cythere pellucida*, G. O. Sars (not Baird), Oversigt af Norges mar. Ostrac., p. 311889. ,, *confusa*, B. and N., *l. c.*, Pt. I. p. 127, pl. xiv. figs. 16-18.

Hab.—Largo Bay, off Musselburgh, and other places; moderately frequent.

47. *Cythere porcellanea*, G. S. Brady.1889. *Cythere porcellanea*, B. and N., *l. c.*, Pt. I. p. 127, pl. xiv. figs. 22-24.

Hab.—South Bay, off Musselburgh, and other places, but for the most part in shallow water that is slightly brackish.

48. *Cythere tenera*, G. S. Brady.1868. *Cythere tenera*, Brady, Mon. rec. Brit. Ostrac., p. 599, pl. xxviii. figs. 29-32.

Hab.—South Bay, and deep water west of May Island; not common.

49. *Cythere semipunctata*, G. S. Brady.1868. *Cythere (?) semipunctata*, Brady, Mon. rec. Brit. Ostrac., p. 411, pl. xxix. figs. 33-37.

Hab.—Aberlady Bay and Largo Bay; not common.

50. *Cythere crispata*, G. S. Brady.1889. *Cythere crispata*, B. and N., *l. c.*, Pt. I. p. 131, pl. xv. figs. 1, 2.

Hab.—Off the west side of Inchkeith; scarce.

51. *Cythere gibbosa*, Brady and Robertson.1869. *Cythere gibbosa*, B. and R., Ann. and Mag. Nat. Hist. (4), vol. iii. p. 368, pl. xxi. figs. 1-3.

Hab.—Brackish pools at the mouth of the Cocklemill Burn, Largo Bay, 1890.

52. *Cythere semiovata*, T. Scott.

1890. *Cythere semiovata*, T. Scott, Eighth Rept. F. B. S., pt. iii.
p. 321, pl. xii. figs. 1, 2.

1896. „ „ B. and N., *l. c.*, Pt. II. (Appendix) p. 732,
pl. lxiii. figs. 14, 15.

Hab.—Off St Monans, frequent. I had at first some doubt concerning the validity of this species, but having as yet seen no described form with which it could agree, either as adult or young, it is retained in this Catalogue. In the original description of the species, “anterior” end should read “posterior” end, and *vice versa*.

53. *Cythere albomaculata*, Baird.

1850. *Cythere albomaculata*, Baird, Brit. Entomotraca, p. 169,
pl. xx. fig. 7.

Hab.—Off Bo’ness and throughout the estuary, but apparently not very plentiful.

54. *Cythere Robertsoni*, G. S. Brady.

1868. *Cythere Robertsoni*, Brady, Ann. and Mag. Nat. Hist. (4),
vol. ii. p. 33, pl. iv. figs. 5, 8-10.

Hab.—Dredged in the neighbourhood of Inchkeith, as well as in other parts of the estuary, but not very common.

55. *Cythere convexa*, Baird.

1850. *Cythere convexa*, Baird, Brit. Entomotraca, p. 174, pl. xxi.
fig. 3.

Hab.—Firth of Forth; distribution similar to that of *C. Robertsoni*, but it appears to be even scarcer. I have observed it only sparingly.

56. *Cythere limicola*, A. M. Norman.

1865. *Cythere limicola*, Norman, Nat. and Mag. Hist. Trans.
Northumb. and Durham, vol. i. p. 20, pl. vi.

Hab.—South Bay, and in deep water west of May Island (1888). I find this to be a moderately rare species in the Firth of Forth.

57. *Cythere cuneiformis*, G. S. Brady.

1868. *Cythere cuneiformis*, Brady, Mon. rec. Brit. Ostrac., p. 404, pl. xxxi. figs. 47-54.

Hab.—Aberlady Bay in 3 fathoms, bottom muddy sand; rare.

58. *Cythere navicula* (A. M. Norman).

1869. *Cytherura navicula*, Norman, Brit. Assoc. Rept. for 1868, p. 292.

Hab.—Off St Monans; not common.

59. *Cythere villosa*, G. O. Sars.

1865. *Cythere villosa*, G. O. Sars, Oversigt af Norges marine Ostracoder, p. 42.

Hab.—South Bay, off Inchkeith, and at various other places; frequent.

60. *Cythere pulchella*, G. S. Brady.

1868. *Cythere pulchella*, Brady, Mon. rec. Brit. Ostrac., p. 404.

Hab.—Firth of Forth (Brady and Norman). Off St Monans; not common. Without careful examination this species might be mistaken for the more common *C. villosa*.

61. *Cythere quadridentata*, Baird.

1850. *Cythere quadridentata*, Baird, Brit. Entom., p. 173, pl. xxi. fig. 2.

Hab.—Off the west side of Inchkeith (1888), and subsequently, but very sparingly, in other parts of the estuary.

62. *Cythere emaciata*, G. S. Brady.

1867. *Cythere emaciata*, Brady, Brit. Assoc. Rept. for 1866, p. 210.

Hab.—Firth of Forth, taken from the stomach of a Sharp-tailed *Lumpenus*, *L. lampretiformis*, captured at Station III. (east of Inchkeith), 13th July 1901. This is the only record I have of *C. emaciata* for the Forth estuary.

63. *Cythere tuberculata*, G. O. Sars.

1865. *Cytheris tuberculata*, G. O. Sars, Oversigt af Norges marine Ostracoder, p. 37.

Hab.—South Bay; deep water west of May Island and other parts of the estuary; frequent.

64. *Cythere concinna*, Rupert Jones.1856. *Cythere concinna*, Jones, Tert. Entom., p. 29, pl. iv. fig. 7 a-f.

Hab.—South Bay, Largo Bay, west of May Island; not very common. Firth of Forth (Brady and Norman).

65. *Cythere finmarchica* (G. O. Sars).1865. *Cythereis finmarchica*, G. O. Sars, Oversigt af Norges marine Ostracoder, p. 41.

Hab.—Off St Monans, frequent (1890).

66. *Cythere angulata* (G. O. Sars).1865. *Cythereis angulata*, G. O. Sars, *l. c.*, p. 40.

Hab.—Firth of Forth (Brady and Norman). Dredged in deep water west of May Island, not common; and subsequently in one or two other parts of the estuary.

67. *Cythere dunelmensis* (Norman).1865. *Cythereis dunelmensis*, Norman, Nat. Hist. Trans. Northumb. and Durh., vol. i. p. 22, pl. vii. figs. 1-4.

Hab.—Firth of Forth (Brady and Norman). Dredged in deep water west of May Island, 1888.

68. *Cythere antiquata* (Baird).1857. *Cythereis antiquata*, Baird, Brit. Entom., p. 176, pl. xx. fig. 2.

Hab.—Firth of Forth (Brady and Norman). Dredged near Inchkeith and in South Bay; not common.

69. *Cythere Whitei* (Baird).1850. *Cythereis Whitei*, Baird, *l. c.*, p. 175, pl. xx. figs. 3, 3a.

Hab.—Largo Bay; not common.

70. *Cythere Jonesii* (Baird).1850. *Cythereis Jonesii*, Baird, *l. c.*, p. 175, pl. xx. fig. 1.

Hab.—South Bay, deep water west of May Island, and other places in the estuary; not common.

Genus (23) *Limnocythere*, G. S. Brady, 1867.

71. *Limnocythere inopinata* (Baird).

1850. *Cythere inopinata*, Baird, *l. c.*, p. 172, pl. xxi. figs. 1, 1 *a-e*.

Hab.—Duddingston Loch, Loch Leven, Linlithgow Loch, Kilconquhar Loch, Lochgelly Loch, and other places. It is a small species that lives amongst the mud, and is easily missed.

72. *Limnocythere Sancti-Patricii*, G. S. Brady and D. Robertson.

1869. *Limnocythere Sancti-Patricii*, B. and R., *Ann. and Mag. Nat. Hist.* (4), vol. iii. p. 17, pl. xviii. figs. 8-11; pl. xxi. fig. 4.

Hab.—Loch Leven, moderately frequent. This is the only loch within the district where this species has been observed.

Genus (24) *Cytheridea*, Bosquet, 1852.

73. *Cytheridea elongata*, Brady.

1868. *Cytheridea elongata*, Brady, *Mon. rec. Brit. Ostrac.*, p. 421, pl. xxviii. figs. 13-16; pl. xl. fig. 6.

Hab.—Firth of Forth (G. S. Brady). Off the west side of Inchkeith, and a few other parts of the estuary; not common.

74. *Cytheridea papillosa*, Bosquet.

1862. *Cytheridea papillosa*, Bosq., *Entom. fossil des terrains de la France*, p. 42, pl. ii. fig. 5 *a-d*.

Hab.—South Bay, off North Berwick, and some other places; not common.

75. *Cytheridea punctillata*, Brady.

1865. *Cytheridea punctillata*, Brady, *Ann. and Mag. Nat. Hist.* (3), vol. xvi. p. 189, pl. ix. figs. 9-11.

Hab.—Aberlady Bay, Largo Bay, and other places.

76. *Cytheridea torosa* (Jones).

1850. *Candona torosa*, Jones, *Ann. and Mag. Nat. Hist.* (2), vol. vi. p. 27, pl. iii. fig. 6.

Hab.—Granton Harbour (D. Robertson); brackish-water

pools on the shore at Aberlady; pools in an old brickfield at Seafield, near Dunbar; mouth of the Cocklemill Burn, Largo Bay.

77. *Cytheridea lacustris* (G. O. Sars).

1862. *Cythere lacustris*, G. O. Sars, Zoolgisk Reise i. Sommeren, 1862, p. 30.

Hab.—Union Canal, near Edinburgh (D. Robertson); Loch Leven, moderately frequent.

Genus (25) **Eucythere**, G. S. Brady, 1868.

78. *Eucythere declivis* (Norman).

1864. *Cythere declivis*, Norman, Nat. Hist. Trans. Northumb. and Durh., vol. i. p. 16, pl. v. figs. 9-12.

Hab.—Off the west side of Inchkeith and various other places; frequent.

Genus (26) **Krithe**, G. S. Brady, Crosskey, and Robertson, 1874.

79. *Krithe bartonensis* (T. R. Jones).

1856. *Cytherideis bartonensis*, Jones, Mon. Ter. Entom., p. 50, pl. v. figs. 2*a*, *b*; 3*a*, *b*.

Hab.—Near the mouth of the Forth estuary; moderately common.

Genus (27) **Loxoconcha**, G. O. Sars, 1862.

80. *Loxoconcha impressa* (Baird).

1850. *Cythere impressa*, Baird, Brit. Entom., p. 173, pl. xxi. fig. 9.

Hab.—Aberlady Bay, Largo Bay, and other parts of the estuary; frequent.

81. *Loxoconcha guttata* (Norman).

1864. *Cythere guttata*, Norman, Nat. Hist. Trans. Northumb. and Durh., vol. i. p. 19, pl. vi. figs. 9-12.

Hab.—Firth of Forth (Brady and Norman), South Bay and off North Berwick; frequent.

82. *Loxoconcha viridis* (O. F. Müller).1785. *Cythere viridis*, Müller, Entomostraca, p. 64, taf. vii. figs. 1, 2.

Hab.—Granton Harbour (D. Robertson); Largo Bay, rather rare.

83. *Loxoconcha multifora* (Norman).1864. *Cythere multifora*, Norman, Nat. Hist. Trans. Northumb. and Durh., vol. i. p. 18, pl. vi. figs. 13-16.

Hab.—Granton Harbour (D. Robertson). This is the only record for the Forth known to me. The species is, however, widely distributed, and is likely to occur in other parts of the estuary.

84. *Loxoconcha pusilla*, Brady and Robertson.1870. *Loxoconcha pusilla*, B. and R., Ann. and Mag. Nat. Hist. (4), vol. vi. p. 23, pl. viii. figs. 1-3.

Hab.—Taken in the vicinity of Cramond Island where the water is somewhat brackish; apparently rare. Firth of Forth (Brady and Norman).

85. *Loxoconcha tamarindus* (T. R. Jones).1856. *Cytherideis tamarindus*, Jones, Tert. Entom., p. 49, pl. iii. figs. 4a, 4b.

Hab.—Near Inchkeith, South Bay, and off North Berwick; frequent.

86. *Loxoconcha fragilis*, G. O. Sars.1865. *Loxoconcha fragilis*, G. O. Sars, Oversigt af Norges marine Ostracoder, p. 65.

Hab.—Firth of Forth (Brady and Norman). This appears to be a rare species in the estuary. I have not myself observed it.

Genus (28) *Xestoleberis*, G. O. Sars, 1865.

87. *Xestoleberis aurantia* (Baird).1835. *Cythere aurantia*, Baird, Mag. Zool. and Bot., vol. ii. p. 143, pl. v. fig. 26.

Hab.—Off St Monans, Aberlady Bay, Largo Bay, etc., but not very plentiful.

88. *Xestoleberis depressa*, G. O. Sars.

1865. *Xestoleberis depressa*, G. O. Sars, Oversigt af Norges marine Ostracoder, p. 68.

Hab.—Off the west side of Inchkeith; rather rare.

Genus (29) *Cytherura*, G. O. Sars, 1865.89. *Cytherura gibba* (O. F. Müller).

1785. *Cythere gibba*, Müller, Entomotraca, p. 66, pl. vii. figs. 7, 8.

1889. *Cytherura gibba*, B. and N., *l. c.*, Part I., p. 190, pl. xviii. figs. 13-16; pl. xxii. figs. 6-12; pl. xxiii. fig. 8.

Hab.—Granton Harbour (D. Robertson), Largo Bay, and off Aberlady.

90. *Cytherura cornuta*, G. S. Brady.

1838. *Cytherura cornuta*, Brady, Mon. rec. Brit. Ostrac., p. 445, pl. xxxii. figs. 12-15.

Hab.—Near Fidra Island; off Musselburgh, and Burnt-island.

91. *Cytherura sella*, G. O. Sars.

1865. *Cytherura sella*, G. O. Sars, *l. c.*, p. 73.

Hab.—Largo Bay, South Bay, and other parts of the estuary; moderately frequent.

92. *Cytherura acuticostata*, G. O. Sars.

1865. *Cytherura acuticostata*, G. O. Sars, *l. c.*, p. 76.

Hab.—Dredged off the west side of Inchkeith. This is said to be one of the commonest species belonging to the *Cytheruræ*.

93. *Cytherura bodotria*, T. Scott.

1890. *Cytherura bodotria*, T. Scott, Eighth Rept. F. B. S., part iii. p. 232, pl. xii. figs. 6, 7.

1896. „ „ B. and N., *l. c.*, Part II. (Appendix) p. 736, pl. xiv. figs. 16, 17.

Hab.—Dredged off St Monans; rare. This species has also been obtained in the Firth of Clyde.

The authors of the Monograph referred to above are inclined to think that this form "must be regarded as the young of

Cytherura acuticostata," and in support of this view they state that though they do not have very young examples of *that* species to compare it with, a very fine series of the young of the *C. cornuta* shows that "the beak occupies a much larger proportion of shell, and is thus more prominently conspicuous" in the young of the *Cytheruræ*, and "that the lateral projections are also more acute than in the adult."¹ These remarks are supported by an indifferent outline side-view of young forms of *C. cornuta* and *C. nigrescens*, but I think for the purpose of comparison a dorsal view should also have been given. I am, however, not extremely anxious whether the "species" stands or not, and therefore will not discuss its merits here. There are one or two points, however, that may be referred to—(1st) If a careful comparison of the *original* figures of *C. bodotria* be made with those of *C. acuticostata* given on plate xxxii. (figs. 12-15) of Dr Brady's excellent "Monograph of recent British Ostracoda," published in the *Transactions of the Linnean Society*, 1868, it will be seen that *C. bodotria* is proportionally more elongated and more depressed, and that, when seen from above, the lateral angles are remarkably prominent and not like the species it is said to be the young of; but (2nd), and leaving *these* differences meanwhile out of account, and taking for granted that *Cytherura bodotria* is the young of *Cytherura acuticostata*, then from what happens with other species one might expect these young to be smaller than the adult, but they are not so. The length of the specimen of *C. bodotria* represented by our drawings in Part III. of the *Eighth Annual Report of the Fishery Board for Scotland* was .5 mm., and the size of *C. acuticostata* as given in the Monograph already referred to at page 446, and which I suppose represents the adult size, is the $\frac{1}{50}$ th of an inch, so that this supposed young form is as big as the one full grown.

But I had even a better and larger specimen of *C. bodotria* than the one figured, though it was similar in form and sculpture, and it would have been figured instead, but it was

¹ I have two smaller (younger) specimens of *C. bodotria*, and these have the lateral projections *less* developed than in the one figured, though otherwise similar to it.

unfortunately accidentally crushed by the object-glass of the microscope having been lowered down on it. I have examined a considerable number of *C. acuticostata* and allied forms, but have failed to observe any with which *C. bodotria* could be incorporated.

94. *Cytherura striata*, G. O. Sars.

1865. *Cytherura striata*, G. O. Sars, *l. c.*, p. 74.

Hab.—Off Musselburgh, Largo Bay, and other parts of the estuary; moderately frequent.

95. *Cytherura angulata*, G. S. Brady.

1868. *Cytherura angulata*, Brady, Monograph recent British Ostracoda, p. 440, pl. xxxii. figs. 22-25.

Hab.—Dredged off the west side of Inchkeith and other parts of the estuary, but not very plentiful.

96. *Cytherura undata*, G. O. Sars.

1865. *Cytherura undata*, G. O. Sars, *l. c.*, p. 75.

Hab.—Dredged off Musselburgh, and off the west side of Inchkeith and other places. It appears to be generally distributed, but being very small (about the $\frac{1}{6\frac{1}{5}}$ th of an inch in length) it is easily overlooked.

97. *Cytherura producta*, G. S. Brady.

1868. *Cytherura producta*, Brady, Mon. rec. Brit. Ostrac., p. 443, pl. xxxii. figs. 60, 61.

1890. ,, *mucronata*, T. Scott, Eighth Rept. F. B. S., pt. iii. p. 323, pl. xii. figs. 3-5 (? juv.).

Hab.—Dredged in Largo Bay and off St Monans, but not very common.

98. *Cytherura nigrescens* (Baird).

1850. *Cythere nigrescens*, Baird, Brit. Entom., p. 171, pl. xxi. figs. 4, 4a.

Hab.—South Bay, off Musselburgh, common. It has also been observed in other parts of the estuary.

99. *Cytherura simplex*, Brady and Norman.

1889. *Cytherura simplex*, B. and N., *l. c.*, Pt. I. p. 200, pl. xviii. figs. 1, 2.

Hab.—Off St Monans, frequent, in 12 to 15 fathoms, bottom partly clean sand and partly gravel.

100. *Cytherura similis*, G. O. Sars.

1865. *Cytherura similis*, G. O. Sars, Oversigt af Norges marine Ostracoder, p. 72.

Hab.—Dredged off Musselburgh and in Largo Bay; not very common.

101. *Cytherura clathrata*, G. O. Sars.

1865. *Cytherura clathrata*, G. O. Sars, *l. c.*, p. 77.

Hab.—Bo'ness, Firth of Forth (Brady and Robertson). Dredged off Bo'ness, off the west side of Inchkeith, and other parts of the Forth; not common.

102. *Cytherura cellulosa*, A. M. Norman.

1865. *Cytherura cellulosa*, Norman, Nat. Hist. Trans. Northumb. and Durh., vol. i. p. 22, pl. v. figs. 17-20; pl. vi. fig. 17.

Hab.—South Bay, and in other parts of the estuary, but apparently not common; but as it only measures $\frac{1}{70}$ of an inch (about .36 mm.) it may have frequently been overlooked.

Genus (30) *Cytheropteron*, G. O. Sars, 1865.

103. *Cytheropteron latissimum* (Norman).

1865. *Cythere latissima*, Norman, *l. c.*, vol. i. p. 19, pl. vi. figs. 5-8.

Hab.—Dredged in South Bay, and in deep water west of May Island; frequent.

104. *Cytheropteron nodosum*, G. S. Brady.

1868. *Cytheropteron nodosum*, Brady, Mon. rec. Brit. Ostrac., p. 448, pl. xxxiv. figs. 31-34.

Hab.—Firth of Forth (Brady and Robertson). South Bay, off Musselburgh; not unfrequent.

105. *Cytheropteron punctatum*, G. S. Brady.

1868. *Cytheropteron punctatum*, Brady, *l. c.*, p. 449, pl. xxxiv.
figs. 45-48.

Hab.—Dredged off St Monans; rather rare.

106. *Cytheropteron angulatum*, Brady and Robertson.

1872. *Cytheropteron angulatum*, B. and R., *Ann. and Mag. Nat. Hist.* (4), vol. ix. p. 62, pl. ii. figs. 7, 8.

Hab.—Largo Bay and off Aberlady; rare.

107. *Cytheropteron depressum*, Brady and Norman.

1889. *Cytheropteron depressum*, B. and N., *l. c.*, Pt. I. p. 218, pl. xx.
figs. 22, 23.

Hab.—Largo Bay and off Aberlady; not common.

108. *Cytheropteron humile*, Brady and Norman.

1889. *Cytheropteron humile*, B. and N., *l. c.*, Pt. II. p. 220, pl. xx.
figs. 4-7.

Hab.—Dredged off Limekilns and in Largo Bay. This small species, which measures only .33 mm., has also been found fairly plentiful in the Clyde in the crevices of partly decayed pieces of wood brought up in the dredge, and usually associated with an interesting species of Copepoda, *Harrietella simulans*, T. Scott.

Genus (31) *Bythocythere*, G. O. Sars, 1865.109. *Bythocythere turgida*, G. O. Sars.

1865. *Bythocythere turgida*, G. O. Sars, *l. c.*, p. 84.

Hab.—Dredged off Musselburgh and other places; not common. This moderately large species has sometimes been obtained in the stomachs of small fishes.

110. *Bythocythere constricta*, G. O. Sars.

1865. *Bythocythere constricta*, G. O. Sars, *l. c.* p. 85

Hab.—Largo Bay and off Aberlady; frequent. This is a moderately common species in the Forth.

111. *Bythocythere simplex* (Norman).

1865. *Cythere simplex*, Norman, Nat. Hist. Trans. Northumb. and Durh., vol. i. p. xvii, pl. v. figs. 1-4.

Hab.—South Bay, in deep water west of May Island, and other places, but not very common. This is a fine large species, conspicuous from its white colour; it forms part of the food of young Gadoids, Rocklings, *Lumpenus*, Long Rough Dabs, and other fishes.

112. *Bythocythere recta*, G. S. Brady.

1868. *Cytheropteron rectum*, Brady, Mon. rec. Brit. Ostrac., p. 476.

Hab.—Largo Bay; rare.

Genus (32) *Pseudocythere*, G. O. Sars, 1865.113. *Pseudocythere caudata*, G. O. Sars.

1865. *Pseudocythere caudata*, G. O. Sars, Oversigt af Norges marine Ostracoder, p. 88.

Hab.—Firth of Forth (Brady and Robertson). Off St Monans; frequent.

Genus (33) *Sclerochilus*, G. O. Sars, 1865.114. *Sclerochilus contortus* (Norman).

1862. *Cythere contorta*, Norman, Ann. and Mag. Nat. Hist. (3), vol. ix. p. 48, pl. ii. fig. 15.

Hab.—South Bay, and in deep water west of May Island; frequent.

Genus (34) *Cytherideis*, T. R. Jones, 1856.115. *Cytherideis subulata*, G. S. Brady.

1868. *Cytherideis subulata*, Brady, Mon. rec. Brit. Ostrac., p. 454, pl. xxxv. figs. 43-46.

Hab.—Dredged off the west side of Inchkeith and other parts in the estuary; not very common.

Genus (35) *Cytherois*, W. Müller, 1884.116. *Cytherois Fischeri* (G. O. Sars).

1865. *Paradoxostoma Fischeri*, G. O. Sars, *l. c.*, p. 96.

Hab.—Off Aberlady and St Monans; frequent.

Genus (36) *Paradoxostoma*, Fischer, 1851.117. *Paradoxostoma variabile* (Baird).

1835. *Cythere variabilis*, Baird, Trans. Berw. Nat. Club, vol. i.
p. 98, pl. iii. fig. 7 *a, b*.

Hab.—Largo Bay, and generally throughout the estuary, especially in the littoral and laminarian zones.

118. *Paradoxostoma ensiforme*, G. S. Brady.

1868. *Paradoxostoma ensiforme*, Brady, Mon. rec. Brit. Ostrac.,
p. 460, pl. xxxv. figs. 8-11.

Hab.—Dredged off the west side of Inchkeith; off Musselburgh and elsewhere.

119. *Paradoxostoma abbreviatum*, G. O. Sars.

1865. *Paradoxostoma abbreviatum*, G. O. Sars, Oversigt af Norges
marine Ostracoder, p. 94.

Hab.—Off Musselburgh; not very common. Easily distinguished from its being remarkably short in comparison to the height; there is also a perceptible difference in the contour of the shell when seen from the side.

120. *Paradoxostoma obliquum*, G. O. Sars.

1865. *Paradoxostoma obliquum*, G. O. Sars, *l. c.*, p. 97.

Hab.—Off Fidra, Musselburgh, and Burntisland; moderately rare. This species is more tumid as viewed from above when compared with most of the other British species.

121. *Paradoxostoma pulchellum*, G. O. Sars.

1865. *Paradoxostoma pulchellum*, G. O. Sars, *l. c.*, p. 95.

Hab.—Dredged off Musselburgh; moderately rare.

122. *Paradoxostoma hibernicum*, G. S. Brady.

1868. *Paradoxostoma hibernicum*, Brady, *l. c.*, p. 460, pl. xxxv.
figs. 35, 36; pl. xl. fig. 7.

Hab.—Largo Bay; rather rare.

123. *Paradoxostoma arcuatum*, G. S. Brady.

1868. *Paradoxostoma arcuatum*, Brady, *l. c.*, p. 461, pl. xxxv.
fig. 37.

Hab.—Granton (Brady and Robertson). Off St Monans, Largo Bay, and vicinity of Inchkeith; rare.

124. *Paradoxostoma orcadense*, G. S. Brady and D. Robertson.

1872. *Paradoxostoma orcadense*, B. and R., Ann. and Mag. Nat. Hist. (4), vol. ix. p. 53, pl. i. figs. 5-7.

Hab.—Off St Monans; rare.

125. *Paradoxostoma Hodgei*, G. S. Brady.

1870. *Paradoxostoma Hodgei*, Brady, Nat. Hist. Trans. Northumb. and Durh., vol. iii. p. 371, pl. xii. figs. 12, 13.

Hab.—Off St Monans, frequent; also near Fidra.

126. *Paradoxostoma flexuosum*, G. S. Brady.

1868. *Bythocythere* (?) *flexuosa*, Brady, Brit. Assoc. Report (1866), p. 211.

1889. *Paradoxostoma flexuosum*, B. and N., *l. c.*, Part I. p. 236, pl. xxi. figs. 11, 12.

Hab.—Firth of Forth (Brady and Robertson). Off Bo'ness, South Bay, and other parts of the estuary; frequent.

127. *Paradoxostoma affine*, T. Scott.

1890. *Paradoxostoma affine*, T. Scott, Eighth Rept. F. B. S., pt. iii. p. 325, pl. xii. figs. 8, 9.

1896. „ „ B. and N., *l. c.*, Part II. (Appendix) p. 739, pl. lxiii. figs. 10, 11.

Hab.—Off St Monans; not common. The species which this form most closely resembles is the *Paradoxostoma arcuatum* of G. S. Brady, but it is not so narrow posteriorly, and its greatest breadth is nearer the posterior extremity. It has also been taken by Canon Norman near Inveraray, Loch Fyne.

Genus (37) *Machærina*, Brady and Norman, 1889.128. *Machærina tenuissima* (Norman).

1869. *Bythocythere tenuissima*, Norman, Brit. Assoc. Rept. for 1868, p. 294.

Hab.—Dredged in deep water west of May Island; not common. Also observed in the stomach of a Five-bearded Rockling captured in the Forth on 13th May 1901. This species is remarkably compressed, and though moderately high in the middle, both ends taper gradually to a narrow point, imparting to the shell a very slender appearance.

Section II. MYODOCOPA.

Family ASTEROPIDÆ, Brady and Norman.

Genus (38) *Asterope*, Philippi, 1840.129. *Asterope maricæ* (Baird).1850. *Cypridina maricæ*, Baird, Proc. Zool. Soc. Lond., pt. xviii. p. 257, pl. xvii. figs. 5-7.1896. *Asterope maricæ*, B. and N., *l. c.*, Part II. p. 630, pl. 1. figs. 1-6 *et seq.**Hab.*—Dredged near the Bass Rock; not common.

Family CYPRINIDÆ, Baird, 1850.

Genus (39) *Philomedes*, Lilljeborg, 1853.130. *Philomedes interpuncta* (Baird).1850. *Cypridina interpuncta*, Baird, *l. c.*, p. 257, pl. xvii. figs. 8-10.*Hab.*—South Bay, off west side of Inchkeith, and other parts of the estuary; sometimes taken with the tow-net, and often with the dredge.

Section III. CLADOCOPA, G. O. Sars, 1865.

Family POLYCOPIDÆ.

Genus (40) *Polycope*, G. O. Sars, 1868.131. *Polycope orbicularis*, G. O. Sars.1865. *Polycope orbicularis*, G. O. Sars, Oversigt af Norges marine Ostracoder, p. 122.*Hab.*—Dredged near Fidra, Firth of Forth; rare.Genus (41) *Polycopsis*, G. W. Müller, 1894.132. *Polycopsis compressa* (Brady and Robertson).1869. *Polycope compressa*, B. and R., Ann. and Mag. Nat. Hist. (4), vol. iii. p. 20, pl. xxi. figs. 5-11.*Hab.*—Off St Monans; frequent.

It will be observed that the Ostracoda enumerated here amount to 132 species, and belong to 41 genera. There are

some others which, judging from their distribution, may also occur within the Forth area, though hitherto they have escaped notice. I have already mentioned that *Scottia Browniana* has been observed in a lacustrine deposit at Elie, Fifeshire, and as it appears to be very local in its distribution, it may yet be found living somewhere within the district. *Cypridopsis Newtoni* is another that may be expected to occur in some of the inland waters, and there are also a few marine forms that are likely to be met with.

About 170 species have, I think, been recorded from Scotland (including the Orkney and Shetland Islands), and if those which have hitherto been only observed in the neighbourhood of Shetland be excepted, it will be found that fully 84 per cent. of the Scottish species are represented within the Forth area.

Order COPEPODA.

The Copepoda comprise a much larger number of species than any of the other Crustacean orders. The species enumerated here amount to 306, and include both free-swimming and parasitic forms.

Various methods have been adopted for the scientific arrangement of these organisms. In that used by Professor G. O. Sars in his great work on the *Crustacea of Norway*, now in course of publication, and which for the sake of uniformity I propose to follow generally, the Copepoda are separated into seven divisions, viz.:—1st, the Calanoida, most of which are free-living and pelagic; 2nd, the Harpacticoida, which for the most part are free-living but demersal; 3rd, the Cyclopoida, some of which are free-living fresh-water species, while others are “semi-parasites,” and live as commensals or messmates with various other organisms; 4th, the Noto-delphyoida, which, for the most part, are “semi-parasites,” associated with various Ascidians; 5th, the Monstrilloida, a small but curious group, which appear to live a partly parasitic and partly free life; 6th, the Caligoida, chiefly parasitic on fishes, but some of which possess a certain freedom of movement; and 7th, the Lernæoida, fish para-

sites, which, in the adult stage, are more or less permanently fixed on some part of the fish.

This arrangement does not differ greatly from that of Professor G. S. Brady's excellent *Monograph of the Free and Semi-Parasitic Copepoda of the British Islands*, except that the Cyclopidae and Notodelphydæ are in that work placed between the Calanoids and Harpacticoids.

The species recorded in the sequel are distributed among the seven divisions as follows:—

1st.	The Calanoida,	represented by	30	species.
2nd.	The Harpacticoida,	„ „	170	„
3rd.	The Cyclopoida,	„ „	59	„
4th.	The Notodelphyoida,	„ „	9	„
5th.	The Monstrilloida,	„ „	5	„
6th.	The Caligoida,	„ „	12	„
7th.	The Lernæoida,	„ „	21	„
			306	„
	Total number,		306	„

Division CALANOIDA.

In the arrangement and nomenclature of the species under this division, Professor G. O. Sars' work—*An Account of the Crustacea of Norway*, vol. iv., Copepoda Calanoida—is generally followed. See also Professor G. S. Brady's *Monograph of the Free and Semi-Parasitic Copepoda of the British Islands*, vol. i.

Family CALANIDÆ.

Genus (1) *Calanus*, Leach, 1816.

1. *Calanus septentrionalis* (Goodsir).

1843. *Cetochilus septentrionalis*, Goodsir, Edin. New Phil. Jour., vol. xxxv. p. 339, pl. vi. figs. 1-11.

1863. *Calanus helgolandicus*, Claus, Die frei-lebenden Copepoden, p. 171, pl. xxvi. figs. 2-9.

Hab.—Common in the Firth. Previously recorded as *C. finmarchicus* (Gunn.), but, as indicated by G. O. Sars, that species is slightly larger, and differs in some structural details, and its distribution appears for the most part to

be confined to the arctic or subarctic seas. There seems to be little doubt that *Cetochilus septentrionalis* is identical with *Calanus helgolandicus*, Claus, and having priority in publication, it should have preference over *C. helgolandicus*.

Family PARACALANIDÆ, G. O. Sars, 1902.

Genus (2) *Paracalanus*, Boeck, 1864.

2. *Paracalanus parvus* (Claus).

1863. *Calanus parvus*, Claus, Die frei-lebenden Copepoden, p. 173, pl. xxvi. figs. 10-14; pl. xxvii. figs. 1-4.

Hab.—Occasionally taken with the tow-net near the seaward limits of the Forth estuary, sometimes in moderate abundance, but usually it is a rather scarce species.

Family PSEUDOCALANIDÆ.

Genus (3) *Pseudocalanus*, Boeck, 1872.

3. *Pseudocalanus elongatus*, Boeck.

1864. *Clausia elongata*, Boeck, Forhandl. Videnskabs-Selskabet, Christiania (1864), p. 9.

Hab.—Common throughout the estuary. Boeck described this species as *Clausia elongata*, but as that generic name had been previously used by Claparède for a genus of parasitic Copepods, he withdrew it and substituted the name *Pseudocalanus*.

Family ÆTIDEIDÆ.

Genus (4) *Bradyidius*, Giesbrecht, 1897.

4. *Bradyidius armatus* (G. S. Brady).

1878. *Pseudocalanus armatus*, G. S. Brady, Monogr. Brit. Copep., vol. i. p. 46, pl. iv. figs. 1-11.

1897. *Bradyidius armatus*, Giesbrecht, Zool. Anzeiger, No. 536.

1898. „ „ T. Scott, Sixteenth F.B. Rept., pt. iii. p. 264, pl. xii. figs. 1-19.

1902. *Undinopsis Bradyi*, G. O. Sars, Crustacea of Norway, vol. iv. p. 32, pls. xix., xx.

Hab.—Off St Monans and a few other places near the

seaward limits of the estuary. I have given several synonyms for this species, as there appears to be some doubt about the name that should be used for it.

Family STEPHIDÆ.

Genus (5) *Stephos*, T. Scott, 1892.

5. *Stephos minor*, T. Scott.

1892. *Stephos minor*, T. Scott, Tenth F. B. Rept., pt. iii. p. 245, pl. vii. figs. 1-13.

Hab.—Off St Monans and other parts of the estuary; not common. Dredged in 7 fathoms off the east side of Inchkeith, 23rd May 1901.

6. *Stephos Scotti*, G. O. Sars.

1897. *Stephos gyrans*, T. Scott (not *S. gyrans*, Giesb.), Fifteenth F. B. Rept., pt. iii. p. 146; pl. ii. figs. 2-9; pl. iii. figs. 17, 18.

1902. ,, *Scotti*, G. O. Sars, Crustacea of Norway, vol. iv. p. 63, pl. xliii.

1903. ,, ,, T. Scott, Twenty-first F. B. Rept., pt. iii. p. 110, pl. ii. figs. 1-4.

Hab.—Obtained in a gathering collected in 1892, but not examined till 1902; also in a gathering from an old quarry near Granton to which the sea has access, collected in 1894; rather rare.

Family PSEUDOCYCLOPIIDÆ.

Genus (6) *Pseudocyclopia*, T. Scott, 1892.

7. *Pseudocyclopia crassicornis*, T. Scott.

1892. *Pseudocyclopia crassicornis*, T. Scott, Tenth F. B. Rept., pt. iii. p. 246, pl. vii. figs. 15-29.

Hab.—Off St Monans, 1891; off the east side of Inchkeith in May 1901, and at a few other places; not common.

8. *Pseudocyclopia minor*, T. Scott.

1892. *Pseudocyclopia minor*, T. Scott, Tenth F. B. Rept., pt. iii. p. 247, pl. viii. figs. 1-10.

Hab.—Off St Monans, 1891; not common. This is a smaller, and apparently a rarer species than the last.

9. *Pseudocyclopia caudata*, T. Scott.

1894. *Pseudocyclopia caudata*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 236, pl. v. figs. 1-8.

Hab.—Off St Monans in 1893; off the east side of Inchkeith, in about 5 fathoms, in May 1901.

10. *Pseudocyclopia Giesbrechti*, Wolfenden.

1902. *Pseudocyclopia Giesbrechti*, Wolfenden, Jour. M. B. A. Plymouth, January 1902, p. 370, pl. iv.

Hab.—West of Queensferry; rare. A single male was dredged to the west of Queensferry, 17th November 1893, but was only recently identified. A female (the one he describes and figures) was taken by Dr Wolfenden with a surface tow-net off the island of Bressay, Shetland, in March 1900.

Family CENTROPAGIDÆ.

Genus (7) *Centropages*, Kröyer, 1848.11. *Centropages typicus*, Kröyer.

1848. *Centropages typicus*, Kröyer, Naturh. Tidsskr. (N.S.), vol. ii. p. 588, pl. vi. figs. 22-26.

Hab.—Not uncommon in the seaward portion of the estuary, especially during summer; but it appears to become rarer in the winter months.

12. *Centropages hamatus* (Lilljeborg).

1853. *Ichtyophorba hamata*, Lillj., De Crust. ex ord. tribus in Scania occurrentibus, p. 185, pl. xxi.; pl. xxvi. figs. 9-12.

Hab.—This species, which is rather more common than the last, has a seasonal distribution somewhat similar to it.

Family DIAPTOMIDÆ.

Genus (8) *Diaptomus*, Westwood, 1836.13. *Diaptomus castor* (Jurine).

1820. *Monoculus castor*, Jurine, Hist. des Monocles, p. 50, pl. iv. figs. 1-6 et seq.

Hab.—In an old quarry on the Braid Hills, Edinburgh, in April 1888: this quarry is now included within the public

park. Roslin Curling Pond (♂ and ♀), 3rd March 1906 (Evans).

14. *Diaptomus gracilis*, G. O. Sars.

1862. *Diaptomus gracilis*, G. O. Sars, Oversigt af de indenlandske Ferskvandscopepoder, Chr. Vid.-Selsk. Forh., 1862, p. 9.

Hab.—Duddingston Loch, Loch Leven, and in most of the fresh-water lochs within the area; common. Loch a Chroin, 2500 feet, September 1906 (W. Evans).

15. *Diaptomus hircus*, G. S. Brady.

1891. *Diaptomus hircus*, Brady, Nat. Hist. Trans. Northumb. and Durham, vol. ii. p. 35, pl. x. figs. 2-4.

Hab.—Loch Katrine, Trossachs, 16th March 1898; rather rare.

16. *Diaptomus Wierzejskii*, J. Richard.

1888. *Diaptomus Wierzejskii*, Richard, Bull. Soc. Zool. de France, vol. xiii. p. 53.

1891. „ *serricornis*, Lillj., Brady, *l. c.*, p. 36, pl. ix. figs. 3-10.

Hab.—Loch Achray, Trossachs, 27th November 1897. Though this is the only loch within the area of the Forth basin in which I have obtained *Diaptomus Wierzejskii*, it is a widely distributed species in the north of Scotland, Shetland, and the Outer Hebrides.

Family TEMORIDÆ.

Genus (9) *Temora*, Baird, 1830.

17. *Temora longicornis* (O. F. Müller).

1792. *Cyclops longicornis*, Müller, Entomostraca Daniae et Norvigiae, p. 115.

Hab.—Common, and generally distributed throughout the estuary.

Genus (10) *Eurytemora*, Giesbrecht, 1881.

18. *Eurytemora velox* (Lilljeborg).

1853. *Temora velox*, Lillj., De Crust. ord. tribus in Scania occurrentibus, p. 177, pl. xx. figs. 2-9 (♀).

Hab.—In the estuary near South Queensferry.

19. *Eurytemora affinis* (Poppe).

1881. *Temora affinis*, Poppe, Abhandl. d. Naturh. ver. Bremen, vol. vii. p. 55, pl. iii.

1891. *Eurytemora affinis*, G. S. Brady, Nat. Hist. Trans. Northumb., Durh., and Newcastle-upon-Tyne, vol. ix. p. 42, pl. xiii. figs. 6-9.

Hab.—Taken in the vicinity of Culross, and between Kincardine-on-Forth and Alloa, in July 1891 and February 1892; moderately common.

Family METRIDIIDÆ.

Genus (11) *Metridia*, Boeck, 1864.20. *Metridia lucens*, Boeck.

1864. *Metridia lucens*, Boeck, Chr. Vid.-Selsk. Forh., p. 238.

1878. „ *armata*, G. S. Brady, Monograph, vol. i. p. 42, pl. ii. figs. 1-12.

Hab.—Obtained sparingly near the mouth of the estuary, and from there west to near Inchkeith.

Family PSEUDOCYCLOPIDÆ.

Genus (12) *Pseudocyclops*, Brady, 1872.21. *Pseudocyclops crassiremis*, G. S. Brady.

1872. *Pseudocyclops crassiremis*, Brady, Nat. Hist. Trans. Northumb. and Durham, vol. iv. p. 431, pl. xvii. figs. 1-8.

Hab.—Off St Monans, 1893; rare.

22. *Pseudocyclops obtusatus*, G. S. Brady and D. Robertson.

1873. *Pseudocyclops obtusatus*, B. and R., Ann. and Mag. Nat. Hist. (4), vol. xii. p. 128, pl. viii. figs. 4-7.

Hab.—Taken sparingly with the dredge off St Monans. Also off the north-west end of Inchkeith, in about 5 fathoms, on 23rd May 1901.

Family CANDACIIDÆ.

Genus (13) *Candacia*, Dana, 1846.23. *Candacia armata*, Boeck.

1872. *Candace armata*, Boeck, Nye Slægter og Arter Saltvands-Copepoder, Chr. Vid.-Selsk. Forh., p. 39.

1878. „ *pectinata*, Brady, Monograph, vol. i. p. 49, pl. viii. figs. 14, 15; pl. x. figs. 1-12.

1902. *Candacia armata*, G. O. Sars, *l. c.*, vol. iv. p. 135, pl. xci.

Hab.—Occasionally captured with the tow-net between Inchkeith and May Island. This species appears to be more frequent in winter and spring than during the summer months.

Family PONTELLIDÆ.

Genus (14) *Anomalocera*, Templeton, 1837.24. *Anomalocera Patersoni*, Templeton.

1837. *Anomalocera Patersoni*, Templ., Trans. Ent. Soc. Lond., vol. ii. p. 35, pl. v. figs. 1-3.

Hab.—This species is, at times, moderately frequent in the Forth estuary, especially during summer and autumn.

Genus (15) *Labidocera*, Lubbock, 1853.25. *Labidocera Wollastoni*, Lubbock.

1857. *Pontella Wollastoni*, Lubbock, Ann. and Mag. Nat. Hist. (2), vol. xx. p. 406, pls. x., xi.

Hab.—Off the east side of Inchkeith; collected 8th June 1891, but not examined till 1899.

Family PARAPONTELLIDÆ.

Genus (16) *Parapontella*, G. S. Brady, 1878.26. *Parapontella brevicornis* (Lubbock).

1857. *Pontellina brevicornis*, Lubbock, Ann. and Mag. Nat. Hist. (2), vol. xx. p. 407, pl. xi. figs. 4-8.

Hab.—This species has been observed above Queensferry, off Musselburgh, in the neighbourhood of Inchkeith, off the Wemyss, and in the neighbourhood of Dunbar.

Family ACARTIIDÆ.

Genus (17) *Acartia*, Dana, 1846.27. *Acartia longiremis* (Lilljeborg).

1853. *Dias longiremis*, Lillj., De Crust. ex ord. tribus in Scania
occurrentibus, p. 181, pl. xxiv.

Hab.—Generally distributed throughout the estuary; not usually uncommon, but at some seasons it appears to be more numerous than at others.

28. *Acartia Clausi*, Giesbrecht.

1889. *Acartia Clausi*, Giesb., Atti Accad. Lincei Rend. Roma (4),
vol. v. sem. 2, p. 25.

Hab.—The distribution and frequency of this species is somewhat similar to the last.

29. *Acartia bifilosa*, Giesbrecht.

1881. *Dias bifilosus*, Giesb., Zool. Anzeiger, vol. iv. p. 257.
1882. „ „ Giesb., Die frei-leb. Copep. der Kieler Förde,
p. 147, pl. iii. figs. 4, 22, 23 *et seq.*

Hab.—Taken in the vicinity of Culross, near the head of the estuary, in 1891.

30. *Acartia discaudata*, Giesbrecht.

1882. *Dias discaudatus*, Giesb., Die frei-leb. Copep. der Kieler
Förde, p. 148, pl. iii. figs. 4, 22, 23 *et seq.*

Hab.—Collected with a tow-net between Portobello and Cockenzie in 1890, and on several subsequent occasions, chiefly in the same part of the estuary. The female of this species is readily distinguished by the character of the furcal joints, which are of the form of broad flattened plates fringed with stout and not very long setæ.

Division HARPACTICOIDA.

I have decided to omit the separating of the Harpacticoida into families, because Professor G. O. Sars, in his work now in course of publication, is creating an entirely new arrangement of these Copepoda. The reader is therefore referred to vol. v. of *The Crustacea of Norway*, part of which is already published, for the systematic distribution of this group.

Genus (18) *Misophria*, Boeck, 1864.31. *Misophria pallida*, Boeck.

1864. *Misophria pallida*, Boeck, Chr. Vidensk.-Selskab. Forhandl.,
p. 248.
1904. „ „ Sars, Crustacea of Norway, vol. v. p. 6,
pls. i., ii.

Hab.—Dredged off St Monans, west of Queensferry, and other parts of the estuary, but not very common.

Genus (19) *Longipedia*, Claus, 1863.32. *Longipedia Scotti*, G. O. Sars.

1893. *Longipedia coronata*, T. and A. Scott (not *L. coronata*, Claus),
Ann. Scot. Nat. Hist., vol. ii. pt. ii. p. 91,
pl. ii. figs. 4-6.
1904. „ *Scotti*, G. O. Sars, *l. c.*, vol. v. p. 11, pl. v. fig. 1.

Hab.—Moderately common throughout the estuary.

33. *Longipedia minor*, T. and A. Scott.

1893. *Longipedia coronata*, var. *minor*, T. and A. Scott, Ann. Scot.
Nat. Hist., vol. ii. p. 93.
1893. „ „ „ „ T. Scott, Eleventh F. B.
Rept., pt. iii. p. 200,
pl. ii. figs. 14-20.
1904. „ *minor*, G. O. Sars, *l. c.*, vol. v. p. 12, pl. v. fig. 2.

Hab.—Not uncommon in various parts of the estuary, as off Musselburgh, Largo Bay, etc.

Genus (20) *Canuella*, T. and A. Scott, 1893.34. *Canuella perplexa*, T. and A. Scott.1893. *Canuella perplexa*, T. and A. Scott, *l. c.*, p. 92, pl. ii. figs. 1-3.1893. „ „ T. Scott, Eleventh F. B. Rept., pt. iii.
p. 201, pl. ii. figs. 21-35.1904. „ „ G. O. Sars, *l. c.*, vol. v. p. 17, pls. viii., ix.

Hab.—Frequent in material dredged off Musselburgh, Largo Bay, Aberdour Bay, and other places.

Genus (21) *Neobradya*, T. Scott, 1892.35. *Neobradya pectinifer*, T. Scott.1892. *Neobradya pectinifer*, T. Scott, Tenth F. B. Rept., pt. iii.
p. 249, pl. xiii. figs. 19-32.

Hab.—Dredged off the north end of Inchkeith in November 1889, and also off St Monans in 1891. This species appears to have a fairly wide distribution, as it has been found in the Clyde by myself, and off the Isle of Man by the late I. C. Thompson of Liverpool.

Genus (22) *Zosime*, Boeck, 1872.36. *Zosime typica*, Boeck.1872. *Zosime typica*, Boeck, Chr. Vid.-Selsk. Forhandl. (1872), p. 46.1880. „ „ G. S. Brady, Monogr. Brit. Copep., vol. ii.
p. 15, pl. xxxix. figs. 1-12.

Hab.—Taken off Musselburgh; frequent.

Genus (23) *Ectinosoma*, Boeck, 1864.37. *Ectinosoma Sarsi*, Boeck.1872. *Ectinosoma Sarsi*, Boeck, Chr. Vid.-Selsk. Forhandl. (1872),
p. 45.1880. „ „ *spinipes*, G. S. Brady, *l. c.*, vol. ii. p. 9, pl. xxxvi.
figs. 1-10.1904. „ „ *Sarsi*, G. O. Sars, Crust. of Norway, vol. v. p. 30,
pl. xvi.

Hab.—Moderately common throughout the estuary.

38. *Ectinosoma melaniceps*, Boeck.

1864. *Ectinosoma melaniceps*, Boeck, Chr. Vid.-Selsk. Forhandl. (1864), p. 30.

1880. „ „ G. S. Brady, *l. c.*, vol. ii. p. 11, pl. xl. figs. 17-20.

Hab.—Moderately common, especially among weeds in shallow inshore water.

39. *Ectinosoma propinquum*, T. and A. Scott.

1896. *Ectinosoma propinquum*, T. and A. Scott, Trans. Linn. Soc. Zool. (S. 2), vol. vi. p. 428, pl. xxxvi. figs. 19, 27, 46 *et seq.*

Hab.—Taken off Musselburgh, not very common; but as it is somewhat similar to *E. Sarsi* in size and general appearance, it is easily overlooked.

40. *Ectinosoma Herdmani*, T. and A. Scott.

1896. *Ectinosoma Herdmani*. T. and A. Scott, *op. cit.*, p. 432, pl. xxxvi. figs. 16, 44 *et seq.*

Hab.—Moderately common in material dredged in Aberdour Bay and off Musselburgh; taken also sparingly off St Monans.

41. *Ectinosoma Normani*, T. and A. Scott.

1896. *Ectinosoma Normani*, T. and A. Scott, *op. cit.*, p. 435, pl. xxxvi. figs. 21, 29, 39 *et seq.*

1904. „ „ G. O. Sars, *op. cit.*, p. 35, pl. xix. fig. 2.

Hab.—Dredged sparingly off Burntisland.

42. *Ectinosoma curticorne*, Boeck.

1872. *Ectinosoma curticorne*, Boeck, Chr. Vid.-Selsk. Forhandl. (1872), p. 45.

1896. „ „ T. and A. Scott, *op. cit.*, p. 430, pl. xxxvi. figs. 22, 30, 34 *et seq.*

Hab.—Taken off Burntisland and Musselburgh, and a few other inland parts of the estuary.

43. *Ectinosoma erythrops*, G. S. Brady.

1880. *Ectinosoma erythrops*, Brady, Monogr. Brit. Copep., vol. ii. p. 12, pl. xxxvi. figs. 11-17.

1896. „ „ T. and A. Scott, *op. cit.*, p. 431, pl. xxxvi. figs. 24, 31, 36 *et seq.*

Hab.—Off St Monans; moderately rare.

44. *Ectinosoma gothiceps*, Giesbrecht.

1882. *Ectinosoma gothiceps*, Giesb., Die frei-leb. Copep. d. Kieler
Föhrde, p. 106, pl. i. figs. 3, 12 *et seq.*

1896. „ „ *pygmæum*, T. and A. Scott, *op. cit.*, p. 433,
pl. xxxvi. figs. 15, 41 *et seq.*

Hab.—Off St Monans ; not common.

45. *Ectinosoma armiferum*, T. and A. Scott.

1896. *Ectinosoma armiferum*, T. and A. Scott, *op. cit.*, p. 434,
pl. xxxvi. figs. 20, 43 *et seq.*

Hab.—Obtained off the west side of May Island in moderately deep water ; rare.

46. *Ectinosoma tenuipes*, T. and A. Scott.

1896. *Ectinosoma tenuipes*, T. and A. Scott, *op. cit.*, p. 436,
pl. xxxvi. figs. 25, 32, 35 *et seq.*

Hab.—Dredged off St Monans ; rare.

47. *Ectinosoma longicorne*, T. and A. Scott.

1896. *Ectinosoma longicorne*, T. Scott, *op. cit.*, p. 439, pl. xxxvi.
figs. 23, 38 *et seq.*

Hab.—Dredged off St Monans ; rare.

48. *Ectinosoma tenuireme*, T. and A. Scott.

1896. *Ectinosoma tenuireme*, T. and A. Scott, *op. cit.*, p. 439,
pl. xxxvi. fig. 33 *et seq.*

Hab.—Dredged off St Monans ; rare.

49. *Ectinosoma gracile*, T. and A. Scott.

1896. *Ectinosoma gracile*, T. and A. Scott, *op. cit.*, p. 429, pl. xxxvi.
figs. 18, 37 *et seq.*

1904. „ „ „ G. O. Sars, Crust. of Norway, vol. v.
p. 39, pl. xxii. fig. 1.

Hab.—Dredged off St Monans ; not very common.

Genus (24) **Microsetella**, Brady and Robertson, 1873.

50. *Microsetella norvegica* (Boeck).

1864. *Setella norvegica*, Boeck, Chr. Vid.-Selsk. Forhandl. (1864), p. 281.

1873. *Microsetella atlantica*, Brady and Robertson, Ann. and Mag. Nat. Hist. (4), vol. xii. p. 130, pl. ix. figs. 11-16.

1904. *Microsetella norvegica*, G. O. Sars, *l. c.*, vol. v. p. 44, pl. xxiv.

Hab.—Taken near Inchkeith, with the surface tow-net, in November 1890, and subsequently in various parts of the estuary, sometimes in considerable numbers.

Genus (25) **Pseudobradya**, G. O. Sars, 1904.

51. *Pseudobradya minor* (T. and A. Scott).

1896. *Bradya minor*, T. and A. Scott, Trans. Linn. Soc. Zool. (S. 2), vol. vi. p. 425, pl. xxxv. figs. 5, 9, 13, 21 *et seq.*

1904. *Pseudobradya minor*, G. O. Sars, *l. c.*, vol. v. p. 41, pl. xxii. fig. 2.

Hab.—Taken off St Monans and near Musselburgh, but apparently not very common.

52. *Pseudobradya elegans* (T. and A. Scott).

1896. *Bradya elegans*, T. and A. Scott, *op. cit.*, vol. v. p. 422, pl. xxxv. figs. 4, 10, 15 *et seq.*

Hab.—Taken sparingly in Largo Bay and off Musselburgh.

53. *Pseudobradya hirsuta* (T. and A. Scott).

1896. *Bradya hirsuta*, T. and A. Scott, *op. cit.*, p. 423, pl. xxxv. figs. 2, 8, 17 *et seq.*

Hab.—Taken in Largo Bay in 1891, and afterwards in other parts of the estuary, but always very sparingly.

54. *Pseudobradya similis* (T. and A. Scott).

1896. *Bradya similis*, T. and A. Scott, *op. cit.*, p. 424, pl. xxxv. figs. 3, 7, 16, 27 *et seq.*

1904. *Pseudobradya similis*, G. O. Sars, Crustacea of Norway, vol. v. p. 42, pl. xxiii. fig. 2.

Hab.—The distribution of this species is similar to the last, but it is a smaller species, being little more than half the

size. It resembles that species in having the furcal joints comparatively long and spreading.

55. *Pseudobradya fusca* (T. and A. Scott).

1896. *Bradya fusca*, T. and A. Scott, *op. cit.*, p. 424, pl. xxxv. figs. 6, 12, 18, 20 *et seq.*

Hab.—Taken very sparingly in Largo Bay. This is a moderately stout species, but the abdominal part of the body tapers more gradually towards the furcal joints than does some of the others.

Genus (26) *Bradya*, Boeck, 1872.

56. *Bradya typica*, Boeck.

1872. *Bradya typica*, Boeck, Chr. Vid.-Selsk. Forhandl. (1872), p. 47.
1904. „ „ G. O. Sars, *l. c.*, vol. v. p. 46, pl. xxv.

Hab.—Obtained off the west side of May Island, and in some other parts of the estuary; not very rare.

Genus (27) *Harpacticus*, M.-Edwards, 1838.

57. *Harpacticus chelifera* (O. F. Müller).

1785. *Cyclops chelifera*, O. F. Müller, Entomotraca. p. 114, pl. xix. figs. 1-3.
1904. *Harpacticus chelifera*, G. O. Sars, *l. c.*, p. 49, pls. xxvii., xxviii.

Hab.—This species appears to be moderately common throughout the estuary, especially in the littoral and laminarian zones.

58. *Harpacticus uniremis*, Kröyer.

1838-40. *Harpacticus uniremis*, Kröyer, in Gaimard's Voy. en Scand., pl. xliii. fig. 1 *a-p.*
1904. „ „ „ G. O. Sars, *l. c.*, vol. v. p. 51, pl. xxix.

Hab.—Taken in the vicinity of Dunbar on 26th April 1894, but not previously recorded.

59. *Harpacticus flexus*, Brady and Robertson.

1873. *Harpacticus flexus*, B. and R., Ann. and Mag. Nat. Hist. (4),
vol. xii. p. 134, pl. ix. figs. 17-21.

1904. ,, ,, G. O. Sars, *l. c.*, vol. v. p. 53. pl. xxx. fig. 2.

Hab.—Obtained off St Monans, in Largo Bay, and off Musselburgh, but not very common.

60. *Harpacticus obscurus*, T. Scott.

1895. *Harpacticus obscurus*, T. Scott, Thirteenth F. B. Rept, pt. iii.
p. 170, pl. iv. figs. 4-12.

Hab.—Taken sparingly in an old quarry at Granton, open to the sea. This species has a superficial resemblance to the last, but is smaller, and the posterior foot-jaws (second maxillipedes) are different.

Genus (28) *Tigriopus*, Norman, 1868.61. *Tigriopus fulvus* (Fischer).

1860. *Harpacticus fulvus*, Fischer, Abhandl. d. König. Bayer.
Akad., vol. viii. p. 656, pl. i. figs. 30-33; pl. ii. figs. 34-39.

1904. *Tigriopus fulvus*, G. O. Sars, *l. c.*, vol. v. p. 54, pls. xxxi.,
xxxii.

Hab.—Moderately common in pools about high-water mark on Cramond Island.

Genus (29) *Zaus*, Goodsir, 1845.62. *Zaus spinatus*, Goodsir.

1845. *Zaus spinatus*, Goodsir, Ann. and Mag. Nat. Hist., vol. xvi.
p. 326, pl. xi. figs. 1-8.

1880. ,, ,, Brady, Monograph, vol. ii. p. 153, pl. lxvi.
figs. 1-9.

1904. ,, ,, G. O. Sars, *l. c.*, vol. v. p. 57, pl. xxxi.

Hab.—Frequent in dredged material collected off Musselburgh, in Largo Bay, and various other places.

63. *Zaus Goodsiri*, G. S. Brady.

1880. *Zaus Goodsiri*, Brady, Monograph, vol. ii. p. 156, pl. lxvi.
figs. 10-13.

1904. ,, ,, G. O. Sars, *l. c.*, vol. v. p. 59, pl. xxxv.

Hab.—Frequent in dredged material collected off St Monans, and also occasionally off the east side of Inchkeith.

Genus (30) *Alteutha*, Baird, 1845.64. *Alteutha interrupta* (Goodsir).

1845. *Sterope interrupta*, Goodsir, Ann. and Mag. Nat. Hist., vol. xvi. p. 326, pl. xi. fig. 10.

1863. *Alteutha bopyroides*, Claus, Die frei-lebenden Copepoden, p. 143, pl. xxii. figs. 10-17.

1904. ,, *interrupta*, G. O. Sars, *l. c.*, vol. v. p. 62, pls. xxxvi. and xxxvii.

Hab.—Frequent in tow-net gatherings collected amongst *Laminaria*, and also in inshore dredgings.

Genus (31) *Eupelte*, Claus.65. *Eupelte purpurocineta* (Norman).

1868. *Alteutha purpurocineta*, Norman, Brit. Assoc. Rept., p. 298.

1880. *Peltidium depressum*, Brady (not Baird), Monograph, vol. ii. p. 160, pl. lxxii. figs. 1-5.

1904. *Alteutha depressa*, G. O. Sars, *l. c.*, vol. v. p. 64, pl. xxxviii.

Hab.—Taken off Musselburgh, off the north end of Inchkeith, and at other parts of the estuary, but not very common.

Genus (32) *Tegastes*, Norman, 1903.66. *Tegastes falcata*, Norman.

1869. *Amymone falcata*, Norman, Brit. Assoc. Rept. for 1868, p. 296.

1880. ,, *sphærica*, Brady (not Claus), Brit. Copep., vol. ii. p. 28, pl. xlix. figs. 1-11.

1904. *Tegastes falcatus*, G. O. Sars, *l. c.*, vol. v. p. 69, pl. xli.

Hab.—Found sparingly throughout the estuary. This curious species was first recorded for the Forth in Part III. of the *Sixth Annual Report of the Fishery Board for Scotland* (1888). The *A. sphærica* of Claus (*A. nigrans*, T. and A. Scott) has not yet been observed in the Forth.

Genus (33) *Porcellidium*, Claus, 1860.67. *Porcellidium fimbriatum*, Claus

1863. *Porcellidium fimbriatum*, Claus, Die frei-lebenden Copepoden, p. 140, pl. xxii. fig. 1.

1904. ,, ,, G. O. Sars, *l. c.*, vol. v. p. 76, pls. xlv., xlv.

Hab.—Obtained in Largo Bay on seaweed brought up

with the dredge. This species frequents the fronds of *Laminaria* and other seaweeds, to which it can adhere very firmly; and from its colour and very flattened form it is, though not uncommon, easily missed unless the weed be carefully examined.

Genus (34) *Aspidiscus*, Norman, 1868.

68. *Aspidiscus littoralis*, G. O. Sars.

1880. *Scutellidium fasciatum*, G. S. Brady (not Norman), Monograph, vol. ii. p. 178, pl. lxxviii. fig. 11; pl. lxxix. figs. 1-9.

1904. *Aspidiscus littoralis*, G. O. Sars, *l. c.*, vol. v. p. 79, pls. xlvi., xlvii.

Hab.—This species was obtained very sparingly in a shore gathering near the mouth of the estuary.

Genus (35) *Tisbe*, Lilljeborg, 1853.

69. *Tisbe furcata* (Baird).

1837. *Cyclops furcatus*, Baird, Mag. Zool. and Bot., vol. i. p. 330, pl. ix. figs. 26-28.

1850. *Canthocamptus furcatus*, idem, Brit. Entomostraca, p. 210, pl. xxv. figs. 1 and 2; pl. xxx. figs. 1-6.

1863. *Tisbe furcata*, Claus, Die frei-lebenden Copepoden, p. 116, pl. xv. figs. 1-10.

1880. *Idya furcata*, G. S. Brady, *l. c.*, vol. ii. p. 172, pl. lxxvii. figs. 1-11.

Hab.—Various parts of the estuary. Common, especially within the littoral and laminarian zones.

70. *Tisbe gracilis* (T. Scott).

1895. *Idya gracilis*, T. Scott, Thirteenth F. B. Rept., pt. iii. p. 171, pl. iv. figs. 13-21.

Hab.—In an old quarry at Granton, open to the sea; rare. Collected by hand-net near low-water.

Genus (36) *Thalestris*, Claus, 1863.

71. *Thalestris longimana*, Claus.

1863. *Thalestris longimana*, Claus, Die frei-lebenden Copepoden, p. 130, pl. xviii. figs. 1-11.

Hab.—In rock pools near high-water on Cramond Island;

in the neighbourhood of Inchkeith; at Station VII. and other parts of the estuary; usually not very plentiful.

Genus (37) *Parathalestris*, G. O. Sars, 1905.

72. *Parathalestris Clausi* (Norman).

1869. *Thalestris Clausi*, Norman, Brit. Assoc. Report for 1868, p. 297.

1888. „ „ G. S. Brady, Monograph vol. ii. p. 128, pl. lxii. figs. 1-12.

Hab.—Largo Bay, washed from seaweed brought up by the dredge, frequent; dredged also in shallow water off Musselburgh.

73. *Parathalestris harpactoides* (Claus).

1863. *Thalestris harpactoides*, Claus, *l. c.*, p. 133, pl. xix. figs. 2-12.

1905. *Parathalestris harpacticoides*, G. O. Sars, Crust. of Norway, vol. v. p. 112, pl. lxvii.

Hab.—Dredged off St Monans in 1891, but only a few specimens were observed.

Genus (38) *Phyllothalestris*, G. O. Sars, 1905.

74. *Phyllothalestris mysis* (Claus).

1863. *Thalestris mysis*, Claus, *l. c.*, p. 130, pl. xviii. figs. 12-16.

1880. „ „ G. S. Brady, *l. c.*, vol. ii. p. 121, pl. lviii. figs. 1-13.

Hab.—Firth of Forth, 1894,—a fine species, very sparingly but widely distributed.

Genus (39) *Halithalestris*, G. O. Sars, 1905.

75. *Halithalestris Croni* (Kröyer).

1849. *Harpacticus Croni*, Kröyer, in Gaimard's Voyages en Scand., Zool., pl. xliii fig. 3 a-n.

1880. *Thalestris serrulatus*, G. S. Brady, *l. c.*, vol. ii. p. 133, pl. lix. figs. 2-11.

Hab.—East of Inchkeith, several taken with surface tow-net in 1889; and in surface tow-net in June 1891 at Station IX. This appears to be a truly pelagic species, and widely distributed, but which only occasionally enters the Forth

estuary. As the colour of these Copepods is bright red they are easily noticed, but the colour quickly disappears when they are preserved in alcohol.

Genus (40) *Rhynchothalestris*, G. O. Sars, 1905.

76. *Rhynchothalestris rufocincta* (Norman).

1880. *Thalestris rufocincta*, Norman (MS. name), in G. S. Brady, *l. c.*, vol. ii. p. 125, pl. lvii. figs. 1-9.

1905. *Rhynchothalestris rufocincta*, G. O. Sars, *Crust. of Norway*, vol. v. p. 120, pls. lxxiii., lxxiv.

Hab.—In pools between tide-marks about Joppa and Granton, 1887, frequent, and subsequently in various other parts of the estuary, but more common in the littoral zone than in deep water.

77. *Rhynchothalestris helgolandica* (Claus).

1863. *Thalestris helgolandica*, Claus, *l. c.*, p. 131, pl. xvii. figs. 12-21.

1880. „ „ G. S. Brady, *l. c.*, vol. ii. p. 123, pl. lxi. figs. 9-14.

1905. *Rhynchothalestris helgolandica*, G. O. Sars, *Crust. of Norway*, vol. v. p. 121, pl. lxxv.

Hab.—Washed from some seaweed obtained while dredging in Largo Bay in April 1891. Also obtained in dredged material collected off the north-east end of Inchkeith in May 1901.

• Genus (41) *Microthalestris*, G. O. Sars, 1905.

78. *Microthalestris forficula* (Claus).

1863. *Thalestris forficula*, Claus, *l. c.*, p. 131, pl. xvii. figs. 7-12.

1894. „ „ *forficuloides*, T. and A. Scott, *Ann. and Mag. Nat. Hist.* (6), vol. xii. p. 142, pl. ix. figs. 4-9.

1905. *Microthalestris forficula*, G. O. Sars, *l. c.*, vol. v. p. 123, pl. lxxvi.

Hab.—In pools near low-water between Leith and Portobello, not very common. The part of the shore where the specimens were chiefly obtained was opposite Seafield, where there is a good deal of mud.

Genus (42) *Amenophia*, Boeck, 1864.79. *Amenophia peltata*, Boeck.

1864. *Amenophia peltata*, Boeck, Oversigt af Norges Copepoder, Chr. Vid.-Selsk. Forhandl., p. 45.

1880. *Thalestris peltata*, G. S. Brady, *l. c.*, vol. ii. p. 138, pl. liii. figs. 11-19.

1895. „ „ T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xiv. p. 35, pl. xv. figs. 11-15; pl. xvi. figs. 1-8.

Hab.—Dredged off Musselburgh; not common. The species differs from the typical *Thalestris* in the body being very depressed and in the structure of some of the appendages, as shown by the drawings in the works referred to.

Genus (43) *Westwoodia*, Dana, 1855.80. *Westwoodia nobilis* (Baird).

1845. *Arpacticus nobilis*, Baird, Trans. Berw. Nat. Club, vol. ii. p. 155.

1880. *Westwoodia nobilis*, Brady, Brit. Copep., vol. ii. p. 141, pl. lxiii. figs. 1-13.

Hab.—Taken at Cramond Island in rock-pools between tide-marks, and in dredged material collected off Musselburgh in 3 to 4 fathoms; not common.

Genus (44) *Pseudothalestris*,¹ Brady, 1883.81. *Pseudothalestris Andrewi* (T. Scott).

1894. *Pseudowestwoodia Andrewi*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 257, pl. ix. figs. 21-29.

Hab.—Dredged off Burntisland in 3 to 4 fathoms water; frequent.

This group of Copepods have so close a general resemblance to *Westwoodia*, Dana, that I ascribed the first species to a new genus, *Pseudowestwoodia*, which recognised that resemblance, but afterwards I discovered that they were identical

¹ The genus *Pseudothalestris* is closely related to *Westwoodia*, Dana, but as it contains a group of species which differ distinctly in the structure of the first pair of thoracic feet, I prefer to retain it as a separate genus.

with the genus *Pseudothalestris*, Brady, described by that author in his "Report on the Challenger Copepoda."¹

82. *Pseudothalestris pygmæa* (T. and A. Scott).

1895. *Pseudothalestris pygmæa*, T. and A. Scott, *Ann. and Mag. Nat. Hist.* (6), vol. xv. p. 55, pl. vi. figs. 8-16 (January 1895).

Hab.—Dredged sparingly in the neighbourhood of Dunbar.

83. ²*Pseudothalestris major* (T. and A. Scott).

(?) 1863. *Westwoodia minuta*, Claus, *Die frei-lebenden Copepoden*, p. 118, taf. xxi. figs. 10-14.

1895. *Pseudowestwoodia major*, T. and A. Scott, *loc. cit.*, p. 56, pl. vi. figs. 17-20.

Hab.—Dredged in the neighbourhood of Granton, and near Dunbar.

Genus (45) **Dactylopusia**, Norman, 1903.³

84. *Dactylopusia tisboides* (Claus).

1863. *Dactylopus tisboides*, Claus, *Die frei-lebenden Copepoden*, p. 127, pl. xvi. figs. 24-28.

Hab.—Shore-pools at Cramond Island; dredged off Musselburgh and other parts of the estuary; frequent in the littoral and laminarian zones.

85. *Dactylopusia neglecta*, G. O. Sars.

1880. *Dactylopus tisboides*, G. S. Brady (brackish-water var.), *l. c.*, vol. ii. p. 108, pl. liv. figs. 14-16.

1905. *Dactylopusia neglecta*, G. O. Sars, *Crust. of Norway*, vol. v. p. 127, pl. lxxviii. fig. 2.

Hab.—This brackish-water form was observed in the lagoon at the mouth of the Cocklemill Burn (east end of Largo Bay).

¹ Cf. paper by T. and A. Scott in *Ann. and Mag. Nat. Hist.* (6), vol. xv. p. 463 (June 1895).

² See remarks on this species and also on the genus in Part III. of the *Twenty-fourth Annual Report of the Fishery Board for Scotland*, at page 277 (1906).

³ The name *Dactylopus*, Claus, being preoccupied by Gill for a genus of fishes, was by Canon Norman changed to *Dactylopusia*.

86. *Dactylopusia vulgaris*, G. O. Sars.

1863. *Dactylopus Strömi*, Claus, *l. c.*, p. 126, pl. xvi. figs. 1-6.

Hab.—Forth, west of Queensferry, washed from lumps of hardened mud; not very common. According to G. O. Sars, this is not the *Canthocamptus Strömi*, Baird, as Claus supposed it to be.

87. *Dactylopusia littoralis* (T. Scott).

1903. *Dactylopus littoralis*, T. Scott, Twenty-first F. B. Rept., pt. iii. p. 124, pl. iii. figs. 2-8.

Hab.—Obtained in pools between tide-marks at Musselburgh in 1894, but not recorded till 1903. This seems to be a rare species.

88. *Dactylopusia finmarchica*, T. Scott.

1903. *Dactylopus longirostris*, Claus, var. *finmarchicus*, T. Scott, Ann. and Mag. Nat. Hist. (7), vol. xi. (Jan. 1903), p. 21, pl. ii. figs. 4-8.

1903. „ „ *mixtus*, T. Scott, Twenty-first F. B. Rept., pt. iii. (pub. 20th July 1903), p. 126, pl. iii. figs. 9-16.

Hab.—This *Dactylopusia* was first described from specimens obtained in East Finmark, within the Arctic Circle, in a collection made by Canon Norman in the summer of 1890. The species was also observed in a gathering collected in 1894 in the old quarry at Granton, where several other interesting Copepoda have been captured. This gathering was not thoroughly examined till 1903, when the *Dactylopusia* was by an oversight redescribed as a “new species,” under the name of *D. mixtus*.

89. *Dactylopusia debilis* (Giesbrecht).

1882. *Dactylopus debilis*, Giesb., Frei-leb. Copep. d. Kieler Föhrde, p. 122, pl. i. figs. 7, 19 *et seq.*

1903. „ „ T. Scott, Twenty-first F. B. Rept., pt. iii. p. 128, pl. v. figs. 20-31.

Hab.—This species occurred very sparingly in a gathering dredged off Musselburgh in 4 to 5 fathoms.

90. *Dactylopusia brevicornis* (Claus).

1868. *Dactylopus brevicornis*, Claus, Die Copepoden-fauna von Nizza, p. 29, pl. iii. figs. 20-25.

1906. *Dactylopusia brevicornis*, T. Scott, Twenty-fourth F. B. Rept., pt. iii. p. 276, pl. xiv. figs. 10-18.

Hab.—Dredged in Largo Bay and in the old quarry at Granton; not common.

Genus (46) *Dactylopodella*, G. O. Sars, 1905.

91. *Dactylopodella flava* (Claus).

1866. *Dactylopus flavus*, Claus, *l. c.*, p. 28, pl. iii. figs. 13-16.

Hab.—Taken very sparingly with the dredge in Largo Bay and at Station III.

Genus (47) *Idomene*, Philippi, 1843.

92. *Idomene forficata*, Philippi.

1843. *Idomene forficata*, Philippi, Archiv. f. Naturgesch., 1843, p. 65, taf. iii. fig. 4.

1880. *Dactylopus flavus*, Brady, Monograph, vol. ii. p. 116, pl. lvi. figs. 1-11 (♂).

Hab.—I have occasionally dredged this species in the Forth estuary, but it was usually passed over as the male of *Dactylopodella flava*.

Genus (48) *Idomenella*, T. Scott, nov. gen., 1896.

This genus has a general resemblance to *Idomene*, Philippi. The antennules (anterior antennæ) short, six- or seven-jointed.

Posterior antennæ tolerably well developed, and composed of two joints.

Mandibles, with masticatory edge, moderately broad, and armed with several stout but irregular teeth. Mandible-palp well developed, and furnished with two uniarticulate branches of moderate size, the outer one wanting the strong spines that appear to be characteristic of species belonging to *Idomene*.

Maxillæ similar in structure to those of the genus mentioned.

The first and second maxillipeds are also similar to those of the same species.

The first pair of thoracic feet resemble those of *Idomene*, but the first joint of the inner branch is not nearly so robust, and the terminal setæ of the end joint are not so strongly clawed.

The second, third, and fourth pairs in the female are similar to those of *Idomene*, but the second pair in the male has the inner branch armed with a strong claw-like spine.

The fifth pair comparatively small, lamelliform, and the primary and secondary joints are sub-equal, and only moderately developed (cf. *Twelfth Annual Report Fishery Board for Scotland*, pt. iii. p. 255, pl. lx. fig. 19, etc.).

Furcal joints short. One ovisac.

The differences that separate this genus from *Idomene* are comparatively small; but in the following two species, which I have ascribed to it, the armature of the mandible-pulp, and the structure of the inner branch of the first pair of thoracic feet, preclude them from a place in Philippi's genus.

93. *Idomenella rostrata*, T. Scott.

1893. *Dactylopus rostratus*, T. Scott, Eleventh F. B. Rept., pt. iii. p. 205, pl. iii. figs. 7-20.

Hab.—The specimens from which this species was described were obtained by carefully washing shells inhabited by the common hermit crab, *Eupagurus*, Bernhardus, which were dredged off the west side of Inchkeith.

94. *Idomenella coronata*, T. Scott.

1894. *Dactylopus coronatus*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 255, pl. ix. figs. 12-20.

Hab.—Dredged near the Bass Rock and also in Largo Bay; not common. It has also been obtained very sparingly off Musselburgh, and at the north end of Inchkeith.

Genus (49) *Amphiascus*, G. O. Sars, 1905.95. *Amphiascus minutus* (Claus).

1863. *Dactylopus minutus*, Claus, Die frei-lebenden Copepoden,
p. 126, taf. xvi. figs. 14, 15.

1880. „ „ Brady, *l. c.*, vol. ii. p. 119, pl. lxvii.
figs. 12-14.

Hab.—In dredged material from Largo Bay,—the only place within the estuary where this species has been noticed.

96. *Amphiascus Catharinæ*, T. Scott.¹

1906. *Amphiascus Catharinæ*, T. Scott, Twenty-fourth F. B. Rept.,
pt. iii. p. 275, pl. xiv. figs. 10-18.

Hab.—Collected by hand-net in an old quarry at Granton, open to the tide, where many interesting forms have been obtained.

97. *Amphiascus tenuiremis* (Brady and Robertson).

1876. *Dactylopus tenuiremis*, B. and R., Brit. Assoc. Report for
1875, p. 197.

1880. „ „ Brady, *l. c.*, vol. ii. p. 115, pl. lvi.
figs. 12-18.

Hab.—Dredged in Largo Bay, where it occurred very sparingly.

98. *Amphiascus similis* (Claus).

1866. *Dactylopus similis*, Claus, Die Copepoden-fauna von Nizza,
p. 25, taf. ii. figs. 29, 30.

1880. „ „ Brady, *l. c.*, vol. ii. p. 110, pl. lv.
figs. 14-16.

Hab.—Dredged west—off Queensferry, 17th November 1893; not common.

99. *Amphiascus imus* (G. S. Brady).

1872. *Canthocamptus imus*, Brady, Nat. Hist. Trans. Northumb.
and Durham, vol. iv. p. 436, pl. xix. figs. 1-5.

1880. *Stenhelia ima*, idem, Monograph, vol. ii. p. 35, pl. xliii.
figs. 1-14.

Hab.—Collected near Cramond and North Berwick in 1887, and near Inchkeith in 1901, but apparently not very common.

¹ Named in compliment to a friend of the author.

Genus (50) *Stenhelia*, Boeck, 1864.100. *Stenhelia hispida*, G. S. Brady.

1880. *Stenhelia hispida*, Brady, Monograph, vol. ii. p. 32, pl. xlii. figs. 1-14.

Hab.—This Copepod was observed sparingly in some material dredged off St Monans in 1893, and off North Berwick in July 1901.

101. *Stenhelia hirsuta*, I. C. Thompson.

1893. *Stenhelia hirsuta*, I. C. Thompson, Revised Rept. on the Copepoda of L—pool Bay, p. 20, pl. xxxi.

Hab.—Dredged off St Monans in 1893, and in the neighbourhood of Inchkeith in 1901. The female of this species carries two ovisacs instead of one, and thus differs from some of the other forms grouped under this genus.

102. *Stenhelia dispar*, T. and A. Scott.

1894. *Stenhelia dispar*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xii. p. 141, pl. viii. figs. 8-12.

Hab.—Dredged in the neighbourhood of the Bass Rock in 1893; apparently rare.

103. *Stenhelia denticulata*, I. C. Thompson.

1893. *Stenhelia denticulata*, I. C. Thompson, *l. c.*, p. 20, pl. xxx. figs. 1-11.

Hab.—This very distinct species was dredged off St Monans in 1893, and it has also been obtained near Inchkeith, but appears to be somewhat rare in the estuary.

104. *Stenhelia reflexa*, T. Scott.

1895. *Stenhelia reflexa*, T. Scott, Thirteenth F. B. Rept., pt. iii. p. 166, pl. iii. figs. 1-9.

Hab.—This species was obtained in pools between tide-marks on the shore north-east of Dunbar in 1894. Only a few specimens were observed. It resembles *Amphiascus imus* in some respects, but differs in the structure of the anterior antennæ and of the first and fifth pairs of thoracic feet, as pointed out in the description.

105. *Stenhelia Blanchardi*, T. and A. Scott.

1895. *Stenhelia Blanchardi*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xvi. p. 353, pl. xv. figs. 1-10.

Hab.—This tolerably distinct form was dredged off St Monans in 1896. The species was described from specimens obtained in some material dredged off Arisaig, Argyleshire, in 1892. Although this form appears to be widely distributed, it does not seem to be very common.

106. *Stenhelia pygmæa*, A. M. Norman and T. Scott.

1905. *Stenhelia pygmæa*, N. and S., Ann. and Mag. Nat. Hist. (7), vol. xv. p. 284.

1906. „ „ idem, Crust. of Devon and Cornwall, p. 142, pl. x. figs. 1-3; pl. xi. figs. 1, 2 *et seq.*

Hab.—Station II., Forth, dredged 26th December 1894. This species, which is very small, appears to be widely distributed, as the type specimens were collected by the Rev. Canon Norman near Eddystone Lighthouse.

107. *Stenhelia confusa*, T. Scott.

1902. *Stenhelia confusa*, T. Scott, Twentieth F. B. Rept., pt. iii. p. 458, pl. xxii. figs. 17-25.

Hab.—Dredged at Station III., 7th June 1901. Apparently rare.

Genus (51) *Ameira*, Boeck, 1864.108. *Ameira longipes*, Boeck.

1864. *Ameira longipes*, Boeck, Oversigt af Norges Copepoder, Chr. Vid.-Selsk. Forhandl., p. 55.

1880. „ „ Brady, Monograph, vol. ii. p. 37, pl. liii. figs. 1-10.

Hab.—Dredged off St Monans and in other parts of the estuary, but not very common.

109. *Ameira longicaudata*, T. Scott.

1892. *Ameira longicaudata*, T. Scott, Tenth F. B. Rept., pt. iii. p. 250, pl. ix. figs. 1-18.

Hab.—Taken in various parts of the estuary. First noticed

in material dredged off St Monans in 1891. This species appears to be moderately frequent in the Firth.

110. *Ameira exilis*, T. and A. Scott.

1894. *Ameira exilis*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xii. p. 139, pl. viii. figs. 18-20; pl. ix. figs. 1-3.

1894. „ „ T. Scott, Twelfth F. B. Rept., pt. iii. p. 242, pl. ix. fig. 30; pl. x. figs. 1-12.

Hab.—This moderately large and distinct species was taken in shore-pools, near low-water, at Seafield, Leith; it was subsequently obtained at Musselburgh and near Dunbar, and in a somewhat similar situation.

111. *Ameira longiremis*, T. Scott.

1894. *Ameira longiremis*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 241, pl. v. figs. 29-32; pl. vi. figs. 1-5.

Hab.—This *Ameira* was taken very sparingly off St Monans in 1893, and this is the only record of it that I have for the Forth estuary, but it has since been obtained in the Clyde in Kilbrannan Sound.

112. *Ameira exigua*, T. Scott.

1894. *Ameira exigua*, T. Scott, *op. cit.*, p. 243, pl. vi. figs. 15-23.

Hab.—Off St Monans, not very common. This species, like the last, has also been obtained in the Clyde.

113. *Ameira tenuiremis*, T. Scott.

1902. *Ameira tenuiremis*, T. Scott, Twentieth F. B. Rept., pt. iii. p. 459, pl. xxiv. figs. 1-9.

Hab.—Dredged off St Monans; rather rare.

114. *Ameira propinqua*, T. Scott.

1902. *Ameira propinqua*, T. Scott, *op. cit.*, p. 460, pl. xxii. figs. 36-42; pl. xxiii. fig. 1; pl. xxiv. figs. 10-18.

Hab.—Dredged off St Monans; rare.

115. *Ameira pusilla*, T. Scott.

1903. *Ameira pusilla*, T. Scott, Twenty-first F. B. Rept., pt. iii. p. 114, pl. v. figs. 1-10.

Hab.—This, which appears to be the smallest member of the genus, was taken sparingly, in shallow water, off Musselburgh; it has not, so far, been noticed anywhere else.

116. *Ameira ambigua*, T. Scott.

1903. *Ameira ambigua*, T. Scott, *op. cit.*, p. 114, pl. v. figs. 11-19.

Hab.—Taken off Musselburgh, in shallow water; apparently rare.

117. *Ameira elegans*, T. Scott.

1905. *Ameira elegans*, T. Scott, Twenty-third F. B. Rept., pt. iii. p. 144, pl. x. figs. 18, 19; pl. xi. figs. 1-9.

Hab.—Collected with hand-net in pools on the shore near low-water, north-west from Dunbar, and also in a shore gathering collected the same year near Musselburgh.

Genus (52) *Robertsonia*, G. S. Brady, 1880.118. *Robertsonia tenuis* (G. S. Brady and Robertson).

1876. *Ectinosoma tenue*, Brady and Robertson, British Assoc. Rept. for 1875, p. 196.

1880. *Robertsonia tenuis*, Brady, Brit. Copep., vol. ii. p. 25, pl. xli. figs. 1-14.

Hab.—This species, which was first noticed in material dredged off North Berwick in 1887, appears to be sparingly distributed throughout the estuary.

Genus (53) *Heteropsyllus*, T. Scott, 1894.119. *Heteropsyllus curticaudatus*, T. Scott.

1894. *Heteropsyllus curticaudatus*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 252, pl. viii. figs. 27-34; pl. ix. fig. 1.

Hab.—Dredged off Musselburgh and near Aberdour; frequent. This species, though extensively distributed, having been observed in the Firth of Clyde and on the south

coast of England, is readily passed over from its resemblance to more widely-known forms.

Genus (54) *Delavalia*, G. S. Brady, 1868.

120. *Delavalia palustris*, G. S. Brady.

1868. *Delavalia palustris*, Brady, Nat. Hist. Trans. Northumb. and Durham, vol. iii. p. 134, pl. v. figs. 10-15.

1880. „ „ idem, Monograph, vol. ii. p. 43, pl. 1. figs. 1-8.

Hab.—Taken in the neighbourhood of Culross, west of Queensferry, where the water is brackish, and also in brackish-water pools at the mouth of the Peffer Burn, Aberlady Bay, 1895.

121. *Delavalia robusta*, Brady and Robertson.

1875. *Delavalia robusta*, B. and B., Brit. Assoc. Rept. (1875), p. 196.

1880. „ „ Brady, *l. c.*, vol. ii. p. 46, pl. li. figs. 15-21.

Hab.—In pools at the mouth of the Cocklemill Burn at the east end of Largo Bay, 16th August 1890; rare.

122. *Delavalia reflexa*, Brady and Robertson.

1875. *Delavalia reflexa*, B. and R., Brit. Assoc. Rept. (1875), p. 196.

1880. „ „ Brady, *l. c.*, vol. ii. p. 45, pl. li. figs. 1-14.

Hab.—Dredged off Burntisland in November 1893; moderately rare.

Genus (55) *Beatricella*,¹ T. Scott, 1905.

123. *Beatricella mimica*, T. Scott.

1897. *Delavalia mimica*, T. Scott, Fifteenth F. B. Rept., pt. iii. p. 150, pl. i. figs. 1-9.

1905. *Beatricella mimica*, T. Scott, Ann. and Mag. Nat. Hist. (7), vol. xvi. p. 568.

Hab.—Dredged in the neighbourhood of Granton and off

¹ This genus has been named in compliment to Miss Beatrice Sprague, daughter of Dr T. B. Sprague, Edinburgh, whose names are so frequently referred to in this Catalogue as successful students of Scottish fresh-water Crustacea.

Musselburgh; not very rare. Dredged also off the north-west end of Inchkeith.

124. *Beatricella œmula*, T. Scott.

1893. *Delavalia œmula*, T. Scott, Eleventh F. B. Rept., pt. iii. p. 204, pl. iv. figs. 36-47.

1905. *Beatricella œmula*, T. Scott, Ann. and Mag. Nat. Hist. (7), vol. xvi. p. 569 (footnote).

Hab.—Dredged in Largo Bay, and subsequently in one or two other places.

Genus (56) **Tachidius**, Lilljeborg, 1853.

125. *Tachidius discipes*, Giesbrecht.

1853. *Tachidius brevicornis*, Lilljeborg (non *Cyclops brevicornis*, O. F. Müller, 1776), De Crust. ex ordinibus tribus, Clad. Ostrac. Copep., p. 196.

1880. „ „ Brady (non *C. brevicornis*, Müller), Monograph, vol. ii. p. 20, pl. xxxvii.

1881. „ *discipes*, Giesb., Die frei-leb. Copep. der Kieler Förhde, p. 108, pl. ii. fig. 4; pl. iv. figs. 25, 28 *et seq.*

Hab.—Brackish-water pools at the mouth of the Cockle-mill Burn at the east end of Largo Bay, 1890. This is a moderately common species in places such as that referred to.

126. *Tachidius littoralis*, Poppe.

1885. *Tachidius littoralis*, Poppe, Die frei-leb. Copep. des Jadebusens, Abhandl. d. nat. Ver. zu Bremen, vol. xi. p. 167, pl. vii. figs. 10-20.

1891. „ *crassicornis*, T. Scott, Tenth F. B. Rept., pt. iii. p. 250, pl. viii. figs. 14-27.

Hab.—Dredged near Culross, west of Queensferry, 1892; not very rare. This, like *T. discipes*, is a brackish-water species, and appears to be generally distributed where the conditions are favourable.

Genus (57) *Pontopolites*, T. Scott, 1894.127. *Pontopolites typicus*, T. Scott.

1894. *Pontopolites typicus*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 251, pl. viii. figs. 9-17.

Hab.—This species was described from specimens dredged off Musselburgh in 1893, and was at that time regarded as rare; subsequently, however, it was found to be moderately frequent in gatherings collected in shallow inshore waters. Like *Tachidius discipes*, this species has the fifth pair of thoracic feet composed each of a single lamelliform joint, but it differs very markedly in the structure of the other thoracic legs, so much so that it is with some hesitation I have placed it under this family.

Genus (58) *Canthocamptus*, Westwood, 1836.128. *Canthocamptus minutus* (Müller).

1785. *Cyclops minutus*, O. F. Müller, Entomostraca, p. 101, pl. xvii. figs. 1-7.

1820. *Monoculus staphylinus*, Jurine, Hist. des Monocles, p. 74, pl. vii. figs. 1-19.

1880. *Canthocamptus minutus*, G. S. Brady, Monograph, vol. ii. p. 48, pl. xlv. figs. 1-17.

Hab.—Moderately common, and generally distributed in lochs, ponds, etc., throughout the district.

129. *Canthocamptus horridus*, S. Fischer.

1860. *Canthocamptus horridus*, Fischer, Akad. d. Wissensch., 8ten Bd. 3te Abth., p. 760, pl. ii. figs. 57-59, 59A.

1880. *Canthocamptus northumbrius*, Brady, *l. c.*, vol. ii. p. 57, pl. xlv. figs. 1-14.

Hab.—Duddingston Loch, 1892; Lochgelly Loch, Fifeshire, 19th August 1896. "Vicinity of Edinburgh" (Dr and Miss Sprague). This species does not appear to be very common in Scotland.

130. *Canthocamptus gracilis*, G. O. Sars.

1863. *Canthocamptus gracilis*, G. O. Sars, Vidensk. i Christiania Forhandl., 1862 (Aftr.), p. 22.

1897. ,, *inornatus*, T. Scott, Fifteenth F. B. Rept., pt. iii. p. 323, pl. ix. figs. 1-12.

1902. ,, *gracilis*, Lillj., Synopsis Spec. hucusque in aquis dulc. Suec. observ. Fam. Harpactic., p. 26, pl. ii. figs. 8-13.

Hab.—Linlithgow Loch, Upper Elf Loch (near Edinburgh),

Loch Achray (Trossachs); not very rare. It appears to be more frequent in small lakes or ponds than in large bodies of water.

131. *Canthocamptus lucidulus*, Rehberg.

1863. *Canthocamptus minutus*, Claus, Die frei-lebenden Copep., p. 122, pl. xii. figs. 1-3 (name preoccupied by O. F. Müller).
 1880. „ „ *lucidulus*, Rehberg, Beitrag. z. Kenntn., p. 551.
 1895. „ „ *minutus*, T. and A. Scott, Ann. Scot. Nat. Hist. (Oct. 1895), p. 236, pl. iv. figs. 14-20.

Hab.—Duddingston Loch, Upper Elf Loch, Loch Leven, Loch Katrine, and others; Humbie Reservoir, near Winchburgh (Evans); moderately frequent, and generally distributed.

132. *Canthocamptus hirticornis*, T. Scott.

1895. *Canthocamptus hirticornis*, T. Scott, Thirteenth F. B. Rept., pt. iii. p. 251, pl. ix. figs. 13-26.
 1902. „ „ *megalops*, Lillj., l. c., p. 30, pl. ii. figs. 14-19.
 1903. „ „ *hirticornis*, T. Scott, Ann. and Mag. Nat. Hist. (7), vol. xi. p. 188.

Hab.—In brackish-water pools at the mouth of the Cocklemill Burn at the east end of Largo Bay. This species is sometimes not uncommon where the conditions are favourable.

133. *Canthocamptus palustris*, G. S. Brady.

1880. *Canthocamptus palustris*, Brady, Monograph, vol. ii p. 53, pl. xxxix. figs. 13-23.
 1895 „ „ „ var. *elongatus*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xv. p. 459, pl. xvi. figs. 7-17.

Hab.—In pools on May Island, 1889.

134. *Canthocamptus parvus*, T. and A. Scott.

1896. *Canthocamptus parvus*, T. and A. Scott, l. c. (6), vol. xviii. p. 6, pl. ii. figs. 14-22.

Hab.—Between tide-marks at Aberlady, and dredged in shallow water off Musselburgh; not common.

135. *Canthocamptus inconspicuus*, T. Scott.

1900. *Canthocamptus inconspicuus*, T. Scott, Eighteenth F. B. Rept., pt. iii. p. 390, pl. xiv. figs. 1-8.

Hab.—Off Musselburgh, 1894; rare. This species was described from specimens found in the Moray Firth.

136. *Canthocamptus Schmeili*, Mrazek.

1893. *Canthocamptus Schmeili*, Mrazek, Zool. Jahrb. Sieb. Bd., p. 116, pl. vii. figs. 107-117.

1895. „ „ T. and A. Scott, Ann. Scot. Nat. Hist. (Oct. 1895), p. 234, pl. iv. figs. 1-13.

Hab.—Loch Leven, Kinross; moderately frequent.

137. *Canthocamptus crassus*, G. O. Sars.

1863. *Canthocamptus crassus*, G. O. Sars, Chr. Vidensk.-Selsk. Forhandl., 1862 (Aft.), p. 23.

1880. *Attheyella spinosa*, Brady, Monograph, vol. ii. p. 58, pl. xliii. figs. 15-18; pl. xlvi. figs. 13-18.

Hab.—Frequent in lochs and ponds throughout the district.

Genus (59) *Attheyella*, G. S. Brady, 1880.138. *Attheyella pygmæa* (G. O. Sars).

1863. *Canthocamptus pygmæus*, G. O. Sars, *op. cit.*, p. 21.

1880. *Attheyella cryptorum*, Brady, Monograph, vol. ii. p. 60, pl. lii. figs. 1-18.

1893. „ „ T. Scott, Eleventh F. B. Rept., pt. iii. p. 225, pl. vi. figs. 21-31.

Hab.—Moderately common, and generally distributed throughout the district.

139. *Attheyella Zschokkei* (Schmeil).

1893. *Canthocamptus Zschokkei*, Schmeil, Copep. des Rhätikon-Geberges, p. 31, pl. iii.

1893. *Attheyella propinqua*, T. Scott, Eleventh F. B. Rept., pt. iii. p. 227, pl. vii. figs. 1-11.

Hab.—Loch Leven, Duddingston Loch, Upper Elf Loch; not very rare, but easily overlooked. Ben Ledi, at about 2500 feet (Evans).

140. *Attheyella Duthiei*, T. and A. Scott.

1895. *Attheyella Duthiei*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xviii. p. 4, pl. ii. figs. 1-13.

1902. *Canthocamptus Duthiei*, Lillj., Synops. sp. huc usque in aquis dulcibus Sueciae observ. Fam. Harpact., p. 41, pl. iii. figs. 5-10.

Hab.—Loch Leven is the only locality within the district where this species has been observed; it was obtained in gatherings collected by hand-net in 1890, 1897, and 1898.

141. *Attheyella cuspidata* (Schmeil).

1893. *Canthocamptus cuspidatus*, Schmeil, *op. cit.*, p. 36, pl. iv.

1897. „ „ „ T. Scott, Fifteenth F. B. Rept., pt. iii. p. 323, pl. ix. figs. 21, 22.

Hab.—Loch Vennachar, Perthshire; not common. In spring at 2500 feet on Ben Ledi, September 1906 (W. Evans).

Genus (60) *Nitocra*, Boeck, 1864.142. *Nitocra tau*, Giesbrecht.

1882. *Nitocra tau*, Giesb., Die frei-leben. Copep. der Kieler Foehrde, p. 117, pl. i. figs. 9, 13; pl. iii. fig. 13; pl. iv. figs. 2, 11, 29 *et seq.*

Hab.—In pools overflowed by the tide at the mouth of the Cocklemill Burn at the east end of Largo Bay; not common.

Genus (61) *Moraria*, T. and A. Scott, 1893.143. *Moraria brevipes* (G. O. Sars).

1863. *Canthocamptus brevipes*, G. O. Sars, Vidensk. i Christiania Forhandl., 1862 (Aftr.), p. 64.

1893. *Moraria Anderson-Smithi*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. vi. p. 213, pl. viii.

Hab.—Loch Vennachar, Loch Leven, Duddingston Loch, and the Upper Elf Loch are some of the places where this species has been obtained.

144. *Moraria Mrazeki*, T. Scott.

1893. *Ophiocamptus brevipes*, Mrazek (not *Canthocamptus brevipes*, G. O. Sars), Zool. Jahrb. 7ter Bd., p. 116, pl. v. fig. 66; pl. vi. figs. 67-70.

1903. *Moraria Mrazeki*, T. Scott, Ann. and Mag. Nat. Hist. (7), vol. xi. p. 194.

Hab.—Loch Lubnaig, Loch Achray, Loch Vennachar.

Genus (62) *Maraenobiotus*, Mrazek, 1893.145. *Maraenobiotus Vejdovskyi*, Mrazek.

1893. *Maraenobiotus Vejdovskyi*, Mrazek, *op. cit.*, p. 103, pl. iv. figs. 17-32; pl. v. figs. 33-37.

1896. „ „ „ T. and A. Scott, *Ann. and Mag. Nat. Hist.* (6), vol. xviii. p. 3, pl. i. figs. 13-21; pl. ii. fig. 23.

Hab.—The only locality within the district where I have found this species is Loch Vennachar.

Genus (63) *Mesochra*, Boeck, 1864.146. *Mesochra Lilljeborgi*, Boeck.

1864. *Mesochra Lilljeborgi*, Boeck, *Oversigt Norges Copepoder*, p. 51.

1873. *Paratachidius gracilis*, Brady and Robertson, *Ann. and Mag. Nat. Hist.* (4), vol. xii. p. 131, pl. viii. figs. 8-16.

1880. *Mesochra Lilljeborgi*, Brady, *Brit. Copep.*, vol. ii. p. 62, pl. xli. figs. 15-21; pl. xlvii. figs. 16-21.

Hab.—Frequent in rock-pools near high-water, Cramond Island (1888); in pools at the mouth of the Cocklemill Burn, Largo Bay, 1890; and subsequently in other parts of the estuary.

147. *Mesochra spinicaudata*, T. and A. Scott.

1895. *Mesochra spinicaudata*, T. and A. Scott, *Ann. and Mag. Nat. Hist.* (6), vol. xv. p. 52, pl. v. figs. 12-25.

Hab.—Shore at Musselburgh, in pools near low-water; frequent.

148. *Mesochra MacIntoshi*, T. and A. Scott.

1895. *Mesochra Mac-Intoshi*, T. and A. Scott, *l. c.*, p. 53, pl. vi. figs. 1-7.

Hab.—Shore at Musselburgh, in pools near low-water, not uncommon, 1894; and also in a gathering dredged off St Monans in July 1901.

149. *Mesochra propinqua*, T. Scott.

1896. *Mesochra propinqua*, T. Scott, *Fourteenth F. B. Rept.*, pt. iii. p. 162, pl. iii. figs. 11-22.

Hab.—In pools between tide-marks at Aberlady Bay, collected by hand-net.

Genus (64) *Danielssenia*, Boeck, 1872.150. *Danielssenia typica*, Boeck.

1872. *Danielssenia typica*, Boeck, Nye Slægter og arter af Saltvands Copepoder, Vid. Selsk. Forhandl., p. 55.

1876. *Zosime spinulosa*, Brady and Robertson, British Assoc. Rept. for 1875, p. 196.

1880. *Jonesiella spinulosa*, G. S. Brady, *l. c.*, vol. ii. p. 41, pl. xlvi. figs. 14-17; pl. xlix. figs. 14, 15.

Hab.—Largo Bay, 1890, and afterwards in other parts of the estuary, but nowhere very plentiful. The food in the stomachs of a sample of small plaice, *Pleuronectes platessa*, measuring off and on about 2 inches in length, sent from Annan on the Solway in 1900, consisted entirely of this species.

Genus (65) *Thompsonula*, T. Scott, 1905.151. *Thompsonula hyæncæ* (I. C. Thompson).

1889. *Jonesiella hyæncæ*, I. C. Thompson, Proc. Biol. Soc. L—pool, vol. viii. p. 193, pl. ix. figs. 1-10.

1893. „ „ T. Scott, Eleventh F. B. Rept., pt. iii. p. 202, pl. iii. figs. 1-6.

1905. *Thompsonula hyæncæ*, T. Scott, Ann. and Mag. Nat. Hist. (7), vol. xvi. p. 570.

Hab.—Dredged near the island of Fidra in February 1893, and later off Musselburgh; not common. The species appears to be widely distributed, but not very plentiful.

Genus (66) *Laophonte*, Philippi, 1840.152. *Laophonte lamellifera* (Claus).

1863. *Cleta lamellifera*, Claus, Die frei-lebenden Copepoden, p. 123, pl. xv. figs. 21-25.

1880. *Laophonte lamellifera*, Brady, Monograph, vol. ii. p. 83, pl. lxxv. figs. 15-23.

Hab.—Dredged off Musselburgh; collected by hand-net in rock-pools between tide-marks at Cramond Island; and dredged at various other parts of the estuary; frequent.

153. *Laophonte curticauda*, Boeck.

1864. *Laophonte curticauda*, Boeck, Oversigt af Norges Copepoder,
p. 65.

1880. „ „ Brady, *l. c.*, vol. ii. p. 80, pl. lxxiii.
figs. 15-18; pl. lxxvi. figs. 1-9.

Hab.—Rock-pool on the shore, Cramond Island; dredged off North Berwick and in other parts of the estuary.

154. *Laophonte longicaudata*, Boeck.

1864. *Laophonte longicaudata*, Boeck, Oversigt af Norges Copepoder,
p. 55.

1880. „ „ Brady, *l. c.*, vol. ii. p. 82, pl. lxxiv.
figs. 12-15; pl. lxxvi. figs. 10-15.

Hab.—Dredged off St Monans in 1889; scarce.

155. *Laophonte longiremis*, T. Scott.

1905. *Laophonte longiremis*, T. Scott, Twenty-third F. B. Rept.,
pt. iii. p. 145, pl. xi. figs. 10-20.

Hab.—In an old quarry at Granton which is open to the sea, collected 25th August 1894. Apparently rare.

156. *Laophonte thoracica*, Boeck.

1864. *Laophonte thoracica*, Boeck, Oversigt af Norges Copepoder,
p. 54.

1880. „ „ Brady, *l. c.*, vol. ii. p. 76, pl. lxxvii.
figs. 1-8.

Hab.—Dredged in the neighbourhood of Inchkeith; off Musselburgh and other parts of the estuary; not very common.

157. *Laophonte similis*, Claus.

1866. *Laophonte similis*, Claus, Copepoden-Fauna von Nizza, p. 23,
pl. v. figs. 13, 14.

1880. „ „ Brady, *l. c.*, vol. ii. p. 78, pl. lxxv. figs. 1-14.

Hab.—In rock-pools between tide-marks at Cramond Island and in one or two other places, but not common.

158. *Laophonte horrida* (Norman).

1876. *Cleta horrida*, Norman, Rept. of the "Valorous" Exped.,
Proc. Roy. Soc. London, 1876, p. 206.

1880. *Laophonte horrida*, Brady, *l. c.*, vol. ii. p. 74, pl. lxxiv.
figs. 1-12.

Hab.—Dredged off St Monans, and at Station V., to the west of May Island; not common. Dredged at the west end of Station VI., 22nd May 1901.

159. *Laophonte denticornis*, T. Scott.

1890. *Laophonte serrata*, T. Scott (not Claus), Eighth F. B. Rept.,
pt. iii. p. 318.

1894. „ *denticornis*, idem, Twelfth F. B. Rept., pt. iii.
p. 246, pl. vii. figs. 13-23.

Hab.—Dredged off St Monans, 1889; dredged off Musselburgh, 30th May 1891; and at Station III. (near the west end, in 5 fathoms), 23rd May 1901.

160. *Laophonte inopinata*, T. Scott.

1892. *Laophonte inopinata*, T. Scott, Tenth F. B. Rept., pt. iii.
p. 256, pl. xi. figs. 1-12.

Hab.—Off the west side of May Island. Several specimens were washed from a large "root" of sea-weed brought up in the trawl-net of the fishery cruiser "Garland" while at work in the neighbourhood of May Island in 1891; males and females carrying ovisacs were obtained. *L. inopinata* appears to be a rare species in the Forth.

161. *Laophonte intermedia*, T. Scott.

1895. *Laophonte intermedia*, T. Scott, Thirteenth F. B. Rept.,
pt. iii. p. 168, pl. iii. figs. 10-20.

Hab.—Shore at Musselburgh, in pools near low-water; dredged off the same place in 3 to 4 fathoms. Obtained also near Granton in an old quarry open to the sea. This is a distinct and easily recognised species.

162. *Laophonte littorale*, T. and A. Scott.

1893. *Laophonte littorale*, T. and A. Scott, Ann. and Mag. Nat.
Hist. (6), vol. xii. p. 238, pl. xi. figs. 7-14.

Hab.—In brackish pools at the mouth of the Peffer Burn, near Aberlady.

163. *Laophonte gracilis*, T. Scott.

1903. *Laophonte gracilis*, T. Scott, Twenty-first F. B. Rept., pt. iii.
p. 118, pl. vi. figs. 6-12.

Hab.—Collected by hand-net near Granton, in an old quarry open to the sea.

164. *Laophonte depressa*, T. Scott.

1894. *Laophonte depressa*, T. Scott, Twelfth F. B. Rept., pt. iii.
p. 245, pl. vi. figs. 24-31; pl. vii. figs. 1-3.

Hab.—Off St Monans and Musselburgh; very sparingly in dredged material from both places.

165. *Laophonte hispida* (Brady and Robertson).

1873. *Asellopsis hispida*, B. and R., Ann. and Mag. Nat. Hist. (4),
vol. xii. p. 137, pl. ix. figs. 6-10.

1880. *Laophonte hispida*, Brady, Monograph, vol. ii. p. 85,
pl. lxxxii. figs. 1-11.

Hab.—Largo Bay, frequent; dredged also off Musselburgh and at various other parts of the estuary.

Genus (67) *Harrietella*,¹ T. Scott, 1906.166. *Harrietella simulans*, T. Scott.

1894. *Laophonte simulans*, T. Scott, Twelfth F. B. Rept., pt. iii.
p. 248, pl. vii. figs. 24-32; pl. viii. fig. 1.

1906. *Harrietella simulans*, T. Scott, Ann. and Mag. Nat. Hist. (7),
vol. xvii. p. 464, pl. xi. figs. 9, 10.

Hab.—Off West Wemyss and other places. The first specimens were obtained inside the valves of a dead *Cyprina*, among trawl refuse; but specimens were found afterwards to be moderately frequent in the crevices of partly decayed pieces of wood brought up in the dredge or trawl-net.

Genus (68) *Laophontodes*, T. Scott, 1894.167. *Laophontodes typicus*, T. Scott.

1894. *Laophontodes typicus*, T. Scott, Twelfth F. B. Rept., pt. iii.
p. 249, pl. viii. figs. 2-8.

Hab.—Dredged at the north end of Inchkeith. This

¹ This genus has been named in complement to Miss Harriet Richardson, M.A., D.Phil., Washington, U.S.A., author of *A Monograph on the Isopods of North America*.

species, which is very small—scarcely $\frac{1}{60}$ of an inch in length—but quite distinct, is apparently very rare in the Firth of Forth. I have only met with it in the neighbourhood of Inchkeith; it seems, however, to have a fairly extensive distribution. Frequent in a gathering of small Crustacea from an old quarry at Granton collected in 1894.

Genus (69) *Normanella*, G. S. Brady, 1880.

168. *Normanella dubia* (Brady and Robertson).

1875. *Laophonte dubia*, B. and R., Brit. Assoc. Report for 1874, p. 196.

1880. *Normanella dubia*, Brady, Monograph, vol. ii. p. 87, pl. lxxviii. figs. 12-22.

Hab.—Dredged off Musselburgh; not very rare, but easily overlooked.

169. *Normanella attenuata*, A. Scott.

1896. *Normanella attenuata*, A. Scott, Lancashire Sea-Fish Lab. Report for 1895, p. 47, pl. iv. figs. 8-20.

1902. „ „ T. Scott, Twentieth F. B. Rept., pt. iii. p. 464, pl. xxiii. figs. 2-4.

Hab.—Dredged off St Monans, in about 10 fathoms; rare. This species was described from specimens dredged off Spanish Head, Isle of Man, in 1895.

Genus (70) *Cletodes*, Brady, 1872.

170. *Cletodes limicola*, G. S. Brady.

1872. *Cletodes limicola*, Brady, Nat. Hist. Trans. Northumb. and Durham, vol. iv. p. 438, pl. xxi. figs. 10-17.

1880. „ „ idem, Monograph, vol. ii. p. 90, pl. lxxix. figs. 1-12.

Hab.—Dredged off North Berwick, off Musselburgh, and off the east side of Inchkeith, but not very common. It seems to be sparingly distributed throughout the estuary.

171. *Cletodes propinqua*, Brady and Robertson.

1876. *Cletodes propinqua*, B. and R., Brit. Assoc. Report for 1875,
p. 196.

1880. „ „ Brady, Monograph, vol. ii. p. 94, pl.
lxxvii. figs. 9-17.

Hab.—In pools between tide-marks at Newhaven and Cramond Island; also dredged off Musselburgh; not very common.

172. *Cletodes Sarsi*, T. Scott.

1905. *Cletodes Sarsi*, T. Scott, Twenty-third F. B. Rept., pt. iii.
p. 146, pl. xii. figs. 1-9.

Hab.—Firth of Forth, 1901; rare. This species was one of several in a bottle containing specimens from various parts of the estuary collected during 1901.

173. *Cletodes curvirostris*, T. Scott.

1894. *Cletodes curvirostris*, T. Scott, Twelfth F. B. Rept., pt. iii.
p. 250, pl. viii. figs. 18-26.

Hab.—Dredged in Largo Bay; dredged off the east side of Inchkeith, and a few other places; not very rare. In this species the rostrum is distinctly, though not strongly, recurved.

174. *Cletodes irrasa*, T. and A. Scott.

1894. *Cletodes irrasa*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6),
vol. xii. p. 141, pl. viii. figs. 8-12.

Hab.—Dredged in 1893 in the neighbourhood of the Bass Rock. It seems to be a rare species, as I have observed it on only one or two occasions since; but it has also been taken in the Clyde and the Moray Firth.

175. *Cletodes tenuipes*, T. Scott.

1897. *Cletodes tenuipes*, T. Scott, Fifteenth F. B. Rept., pt. iii.
p. 170, pl. i. figs. 19-27.

Hab.—Off Musselburgh, dredged in 3 to 4 fathoms, in 1891; rare. The species was described from specimens

taken in the Clyde in 1896. The Forth examples, which had been put aside when collected, were not identified till later.

176. *Cletodes lata*, T. Scott.

1892. *Cletodes lata*, T. Scott, Tenth F. B. Rept., pt. iii. p. 257, pl. x. figs. 10-18.

Hab.—Dredged off St Monans, off Musselburgh, and other parts of the estuary; not common.

177. *Cletodes similis*, T. Scott.

1895. *Cletodes similis*, T. Scott, Thirteenth F. B. Rept., pt. iii. p. 168, pl. iii. figs. 12-26; pl. iv. figs. 1-3.

Hab.—In the same gatherings with the last, as well as in pools between tide-marks, but always sparingly distributed. This species has a somewhat close resemblance to *C. lata*, and was at first regarded as a variety of it, but afterwards it was found to be quite distinct.

178. *Cletodes longicaudata*, Brady and Robertson.

1876. *Cletodes longicaudata*, B. and R., Brit. Assoc. Report for 1875, p. 196.

1880. „ „ Brady, Monograph, vol. ii. p. 92, pl. lxxix. figs. 13-19.

1902. „ „ T. Scott, Twentieth F. B. Rept., pt. iii. p. 465, pl. xxiii. figs. 26-33.

Hab.—Dredged off St Monans in 1889; dredged off the east side of Inchkeith in June 1901; rare.

179. *Cletodes neglecta*, T. Scott.

1903. *Cletodes neglecta*, T. Scott, Twenty-first F. B. Rept., pt. iii. p. 120, pl. iv. figs. 20-31.

Hab.—Dredged in Aberlady Bay in 1895; not common. This species, which has been known to me for a considerable time, is somewhat intermediate between *C. longicaudata* and *C. limicola*; the furcal joints are about half as long as those of *C. longicaudata*; it thus differs from both the species named. Though collected in 1895, this is the first time it has been recorded for the Forth. The species was described from Moray Firth specimens.

Genus (71) *Itunella*, Brady, 1894.180. *Itunella tenuiremis* (T. Scott).

1893. *Cletodes tenuiremis*, T. Scott, Thirteenth F. B. Rept., pt. iii. p. 204, pl. iii. figs. 21-28.

1894. *Itunella subsalsa*, G. S. Brady, Nat. Hist. Trans. Northumb., Durh., and Newcastle-upon-Tyne, vol. xiii. p. 6, pl. i. (separate reprint).

Hab.—Dredged in the neighbourhood of Inchkeith in February 1893; and subsequently, but very sparingly, in several other places.

Genus (72) *Fultonia*, T. Scott, 1902.181. *Fultonia hirsuta*, T. Scott.

1902. *Fultonia hirsuta*, T. Scott, Twentieth F. B. Rept., pt. iii. p. 466, pl. xxiii. figs. 5-12.

Hab.—Dredged very sparingly off St Monans, in 14 to 15 fathoms, on 22nd May 1901.

Genus (73) *Enhydrosoma*, Boeck, 1872.182. *Enhydrosoma curvatum* (Brady and Robertson).

1876. *Rhizothrix curvata*, B. and R., Brit. Assoc. Report for 1875, p. 197.

1880. *Enhydrosoma curvatum*, Brady, Monograph, vol. ii. p. 98, pl. lxxxii. figs. 12-15; pl. lxxxii. figs. 11-19.

Hab.—Largo Bay and other parts of the estuary; moderately frequent.

183. *Enhydrosoma gracile*, T. Scott.

1903. *Enhydrosoma gracile*, T. Scott, Twenty-first F. B. Rept., pt. iii. p. 122, pl. ii. figs. 16-26; pl. iii. fig. 1.

Hab.—Musselburgh, pools on the shore near low-water; moderately rare. This is a smaller species than *E. curvatum*.

Genus (74) *Nannopus*, G. S. Brady, 1880.184. *Nannopus palustris*, G. S. Brady.

1880. *Nannopus palustris*, Brady, *l. c.*, vol. ii. p. 101, pl. lxxvii. figs. 18-20.

1902. ,, ,, T. Scott, Twentieth F. B. Rept., pt. iii. p. 466, pl. xxiii. figs. 13-25.

Hab.—Brackish-water pools at the mouth of the Cocklemill

Burn at the east end of Largo Bay, also in shore-pools at Musselburgh, and in an old quarry near Granton which is open to the sea; not common.

Genus (75) *Platychelipus*, G. S. Brady, 1880.

185. *Platychelipus littoralis*, G. S. Brady.

1880. *Platychelipus littoralis*, Brady, *l. c.*, vol. ii. p. 103, pl. lxxix. figs. 20-23; pl. lxxx. figs. 15-19.

1893. ,, ,, T. Scott, Eleventh F. B. Rept., pt. iii. p. 205.

Hab.—Sparingly in a gathering of dredged material collected near Culross, and in an old quarry at Granton; dredged in shallow water off Musselburgh and in Aberlady Bay; scarce.

Genus (76) *Cylindropsyllus*, G. S. Brady, 1880.

186. *Cylindropsyllus lævis*, G. S. Brady.

1880. *Cylindropsyllus lævis*, Brady, *l. c.*, vol. iii. p. 30, pl. lxxxiv. figs. 1-8.

1892. ,, ,, T. Scott, Tenth F. B. Rept., pt. iii. p. 258, pl. xiii. figs. 1-18.

Hab.—Dredged off St Monans; frequent. I have found this species in other parts of the estuary, but much less frequently than in the "Fluke Hole" off St Monans.

187. *Cylindropsyllus minor*, T. Scott.

1892. *Cylindropsyllus minor*, T. Scott, *op. cit.*, p. 260, pl. xi. figs. 17-24.

1900. ,, ,, idem, Eighteenth F. B. Rept., pt. iii. p. 349, pl. xiv. figs. 23-32.

Hab.—Dredged off St Monans along with *C. lævis*, but not so common.

Genus (77) *Leptopontia*, T. Scott, 1902.

188. *Leptopontia curvicauda*, T. Scott.

1902. *Leptopontia curvicauda*, T. Scott, Twentieth F. B. Rept., pt. iii. p. 463, pl. xxii. figs. 26-35.

Hab.—Dredged off St Monans; not very common.

Genus (78) *Leptastacus*, T. Scott, 1906.189. *Leptastacus macronyx*, T. Scott.1892. *Tetragoniceps macronyx*, T. Scott, Tenth F. B. Rept., pt. iii.
p. 253, pl. x. figs. 19-28.1906. *Leptastacus macronyx*, T. Scott, Ann. and Mag. Nat. Hist. (7),
vol. xvii. p. 461.*Hab.*—Dredged off St Monans in 1891, and again in May 1901; not very common.Genus (79) *Evansula*, T. Scott, nov. nom.(Syn. *Evansia*, T. Scott, preoccupied for a genus of Spiders.¹)190. *Evansula incerta*, T. Scott.1892. *Tetragoniceps incertus*, T. Scott, Tenth F. B. Rept., pt. iii.
p. 254, pl. xii. figs. 1-17.1906. *Evansia incerta*, T. Scott, Ann. and Mag. Nat. Hist. (7),
vol. xvii. p. 461.*Hab.*—Dredged off St Monans in 1891, and again in May 1901; not common.191. *Evansula pygmæa*, T. Scott.1903. *Tetragoniceps pygmæus*, T. Scott, Twenty-first F. B. Rept.,
pt. iii. p. 117, pl. iv. figs. 11-19.1906. *Evansia pygmæa*, T. Scott, Ann. and Mag. Nat. Hist. (7),
vol. xvii. p. 461.*Hab.*—Collected near Musselburgh in 1894. This differs from the last by its smaller size, and in the structure of fifth thoracic feet and caudal joints.

¹ In the *Annals and Magazine of Natural History* for May 1906 I instituted a genus of the Copepoda under the name of *Evansia*, in compliment to William Evans, Edinburgh, a successful investigator in various departments of Scottish natural history. I find, however, that the name *Evansia* is preoccupied for a genus of spiders, described by the Rev. O. Pickard Cambridge in *Proc. Dorset Nat. Hist. and Antiquarian Field Club*, vol. xxi. p. 37, and I therefore adopt the modified form *Evansula* for the Copepod genus referred to.

Genus (80) *Tetragoniceps*, G. S. Brady, 1880.

192. *Tetragoniceps* (?) *maleolata*, G. S. Brady.

1880. *Tetragoniceps maleolata*, Brady, Monograph, vol. ii. p. 66,
pl. lxxviii. figs. 1-11.

1892. „ (?) „ T. Scott, Tenth F. B. Rept., pt. iii.
p. 252, pl. viii. figs. 11, 12.

1900. „ „ idem, Eighteenth F. B. Rept.,
pt. iii. p. 391, pl. xiv. figs. 9-17.

Hab.—Dredged off St Monans in 1892, apparently not very rare; and again on 22nd May 1901, several specimens were obtained. This form differs from the *T. maleolata* described by Dr Brady in the structure of fifth pair of thoracic feet, but is otherwise identical with it.

193. *Tetragoniceps brevicauda*, T. Scott.

1900. *Tetragoniceps brevicauda*, T. Scott, Eighteenth F. B. Rept.,
pt. iii. p. 392, pl. xiv. figs. 18-22.

Hab.—This species was obtained somewhat sparingly in material dredged off St Monans in 1896, and again in 1901. It has a general resemblance to *T. maleolata*, but the furcal joints are distinctly shorter.

Genus (81) *Phyllopodopsyllus*, T. Scott, 1896.

194. *Phyllopodopsyllus Bradyi*, T. Scott.

1892. *Tetragoniceps Bradyi*, T. Scott, Tenth F. B. Rept., pt. iii.
p. 253, pl. lix. figs. 19-32.

1906. *Phyllopodopsyllus Bradyi*, T. Scott, Ann. and Mag. Nat.
Hist. (7), vol. xvii. p. 459.

Hab.—Dredged off St Monans in 1891, and again in 1901; apparently rare. This species and the next are easily distinguished by the large size and leaf-like form of the fifth pair of thoracic feet of the female.

Genus (82) *Pteropsyllus*, T. Scott, 1906.

195. *Pteropsyllus consimilis*, T. Scott.

1894. *Tetragoniceps consimilis*, T. Scott, Twelfth F. B. Rept., pt. iii.
p. 244, pl. vii. figs. 4-12.

1906. *Pteropsyllus consimilis*, T. Scott, Ann. and Mag. Nat. Hist. (7),
vol. xvii. p. 459, pl. xi. figs. 7, 8.

Hab.—Dredged off St Monans; rare. This species is

similar in its general appearance to *T. Bradyi*, but there are a few structural differences, one of the most important being the three-jointed inner branches of the first pair of thoracic feet: in *T. Bradyi* these branches are only two-jointed.

Genus (83) *Leptosyllus*, T. Scott, 1894.

196. *Leptosyllus typicus*, T. Scott.

1894. *Leptosyllus typicus*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 254, pl. ix. figs. 2-11.

Hab.—West of Queensferry, washed from lumps of hardened mud, which were composed for the most part of the agglutinated tubes of a species of *Sabella*, collected 25th January 1894; rare.

197. *Leptosyllus Robertsoni*, T. and A. Scott.

1895. *Leptosyllus Robertsoni*, T. and A. Scott, Ann. Scot. Nat. Hist. (January 1895), p. 30, pl. ii. figs. 1-14.

Hab.—In pools between tide-marks at Musselburgh, 20th August 1894; rare.

198. *Leptosyllus minor*, T. and A. Scott.

1895. *Leptosyllus minor*, T. and A. Scott, *op. cit.*, p. 31, pl. ii. figs. 15-22.

Hab.—In pools between tide-marks at Musselburgh, taken at the same time as *L. Robertsoni*; moderately rare.

199. *Leptosyllus intermedius*, T. and A. Scott.

1895. *Leptosyllus intermedius*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xv. p. 51, pl. v. figs. 1-11.

Hab.—In pools between tide-marks at Musselburgh, in company with *L. minor*, *L. Robertsoni*, and a few of the other rare things mentioned elsewhere in this Catalogue. This species appeared to be rather more frequent than the other two.

Genus (84) *Paramesochra*, T. Scott, 1892.

200. *Paramesochra dubia*, T. Scott.

1892. *Paramesochra dubia*, T. Scott, Tenth F. B. Report, pt. iii. p. 252, pl. xii. figs. 18-32.

Hab.—Dredged off the west side of May Island in February

1892, and off St Monans, in 13 fathoms, on 22nd May 1901. This curious form was also taken near Port Erin, Isle of Man, by the late I. C. Thompson of Liverpool.

Division CYCLOPOIDA.

Family CYCLOPIDÆ.

Genus (85) *Oithona*, Baird, 1843.201. *Oithona similis*, Claus.

1866. *Oithona similis*, Claus, Copepoden-Fauna von Nizza, p. 14.

Hab.—Generally distributed in the Firth of Forth, and sometimes common. This is the form which, in my earlier papers on "Scottish Marine Copepoda," is recorded under the name of *Oithona spinifrons*, Boeck.

202. *Oithona* (?) *setigera*, Dana.

1849. *Oithona setigera*, Dana, U.S. Explor. Exped. [Amer. Jour. Sci. (2), vol. viii.].

1892. ,, ,, Giesb., Fauna u. Flora Golfes v. Neapel; Monogr. xix.—Pelag. Copep., p. 548, pl. xxxiv. figs. 3, 14, 41 *et seq.*

Hab.—Obtained in a bottom tow-net gathering collected east of Inchkeith in March 1891, and again on 22nd April 1901.

Genus (86) *Cyclopina*, Claus.203. *Cyclopina gracilis*, Claus.

1863. *Cyclopina gracilis*, Claus, Die frei-lebenden Copepoden, p. 104, pl. x. figs. 9-15.

1878. ,, ,, Brady, Monograph, vol. i. p. 93, pl. xxiv. figs. 1-9; vol. ii. pl. xci. figs. 10, 11.

1900. ,, ,, Giesb., Mitth. über Copep., Mitth. a. d. Zool. Station z. Neapel, Bd. 14, p. 45.

Hab.—Generally, though somewhat sparingly, distributed in the Firth of Forth. I have taken it as far west as Charlestown, as well as near the outside limits of the estuary.

204. *Cyclopina littoralis* (G. S. Brady).

1872. *Cyclops littoralis*, Brady, Nat. Hist. Trans. Northumb. and Durham, vol. iv. p. 429, pl. xvii. figs. 9-14.

1878. *Cyclopina littoralis*, idem, Monograph, vol. i. p. 92, pl. xv. figs. 1-9.

1900. „ „ Giesb., Mitth. über Copep., Mitth. a. d. Zool. Station z. Neapel, Bd. 14, p. 43, pl. iii. figs. 1-13.

Hab.—Collected at Cramond Island in rock-pools; dredged off Musselburgh and other parts of the estuary, especially in the littoral and laminarian zones; not uncommon.

205. *Cyclopina elegans*, T. Scott.

1894. *Cyclopina elegans*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 237, pl. v. figs. 9-19.

1900. „ „ Giesb., Mitth. über Copep., Mitth. a. d. Zool. Station z. Neapel, Bd. 14, p. 44, pl. iii. figs. 14-22.

Hab.—This species has been taken off St Monans, but not common.

Genus (87) *Pterinopsyllus*, G. S. Brady, 1880.

(Syn. *Lophophorus*, Brady, 1878, a name preoccupied by Temminck in 1815 for a genus of Birds.)

206. *Pterinopsyllus insignis*, G. S. Brady.

1878. *Lophophorus insignis*, G. S. Brady, Monograph, vol. i. p. 122, pl. xiii. figs. 1-10; pl. xv. fig. 10.

1880. *Pterinopsyllus insignis*, idem, *ibidem*, vol. iii. p. 23.

Hab.—This species was on several occasions observed moderately frequent in dredged material collected to the west of Queensferry, but appeared to be rare near the seaward limits of the estuary.

Genus (88) *Cyclops*, O. F. Müller, 1776.

The members of this genus are, with few exceptions, fresh-water or brackish-water species. Those recorded here

may be conveniently arranged according to the number of joints in the antennules.

(a) SPECIES WITH SEVENTEEN-JOINTED ANTENNULES.

207. *Cyclops strenuus*,¹ Fischer.

1851. *Cyclops strenuus*, Fischer, Bull. Soc. Imp. des Nat. Moscou, vol. xxiv. p. 419, taf. ix. figs. 12-21.

Hab.—Duddingston Loch, and most of the other lochs within the district, where it occurs nearly all the year round. *Cyclops pulchellus* and *Cyclops abyssorum* are other names by which the species is known.

208. *Cyclops Leuckarti*, Claus.

1857. *Cyclops Leuckarti*, Claus, Das gen. Cycl., Archiv. f. Naturg., Jahrg. xxiii. Bd. 1, p. 35, taf. ii. figs. 13, 14.

1897. „ „ T. Scott, Fifteenth F. B. Rept., pt. iii. p. 322, pl. ix. figs. 23-25.

Hab.—Loch Vennachar and Loch Voil, Perthshire; frequent. (See also note on this species in Appendix, p. 380.)

209. *Cyclops bicuspidatus*, Claus.

1857. *Cyclops bicuspidatus*, Claus, *op. cit.*, p. 209, taf. xi. figs. 6 and 7.

1892. „ „ Schmeil, Deutschl. freileb. Süßw.-Copep. (Cyclopidae), p. 75, taf. ii. figs. 1-3.

Hab.—Loch Achray (Trossachs), Loch Leven, Duddingston Loch, ponds on the Braid Hills, Edinburgh (April 1888), as well as in other lochs and ponds throughout the district, but usually not very common.

210. *Cyclops vernalis*, Fischer.

1853. *Cyclops vernalis*, Fischer, Bull. Soc. Imp. des Nat. Moscou, vol. xxvi. p. 90, taf. iii. figs. 1-5.

1891. „ *elongatus*, G. S. Brady, Revis. Brit. Cyclopidae and Calanidae, p. 5, pl. i. figs. 1-5.

1892. „ *vernalis*, Schmeil, *l. c.*, p. 88, taf. ii. figs. 4-7.

Hab.—The distribution of this species is somewhat similar

¹ *Cyclops Ewarti*, Brady, was described in the *Sixth Annual Report of the Fishery Board for Scotland*, p. 232, pl. viii. figs. 1-6, from specimens obtained above Queensferry. I am now inclined to consider this as representing a scarcely mature stage of *C. strenuus*, Fischer.

to that of *C. bicuspidatus*. It has been obtained in several localities near Edinburgh, *i.e.*, Upper Elf Loch, Braids (Scott and Lindsay), and pond at Bonaly Golf Course (Dr and Miss Sprague). Ben Ledi, at 2500 feet (W. Evans).

211. *Cyclops bisetosus*, Rehberg.

1880. *Cyclops bisetosus*, Rehberg, Beitrag. z. Kenntnis der freileb. Süßw.-Copep., Abhand. d. Natur. Ver. zu Bremen, vol. vii. p. 533, pl. iv.

1901-1902. „ „ Dr and Miss Sprague, Entom. of Midlothian, Trans. Edin. Field Natur., vol. iv. p. 255, pl. xxxi. figs. 18-22.

Hab.—Loch Achray, Loch Leven, and Duddingston Loch. Pond on House-o'-Hill Farm, Midlothian (Dr and Miss Sprague).

212. *Cyclops viridis* (Jurine).

1820. *Monoculus quadricornis* var. *viridis*, Jurine, Hist. des Monocles, p. 46, pl. iii. fig. 1.

1878. *Cyclops gigas*, G. S. Brady, Monograph, vol. i. p. 105, pl. xx. figs. 1-16.

Hab.—Loch Katrine, Loch Leven, Duddingston Loch, etc. Burntisland Reservoir and Loch a Chroin (Evans). A common and generally distributed species.

213. *Cyclops signatus*, Koch.

1820. *Monoculus quadricornis fuscus*, Jurine, *l. c.*, p. 47, pl. ii. fig. 2.

1838. *Cyclops signatus*, Koch, Deutschlands Crustaceen, Myriapoden, und Arachniden, Heft. 21, fig. 8.

Hab.—Loch Katrine, Loch Achray, Loch Lubnaig, Perthshire. Ravelston Cottage Quarry, May 1900 (Dr and Miss Sprague). Side of river Teith, near Callander (W. Evans).

214. *Cyclops annulicornis*, Koch.

1820. *Monoculus quadricornis albidus*, Jurine, *l. c.*, p. 44, pl. ii. figs. 10 and 11.

1838. *Cyclops annulicornis*, Koch, Deutschlands Crustaceen, Myriapoden, und Arachniden, Heft. 21, pl. vi.

Hab.—This is a common and generally distributed species throughout the entire area.

(b) SPECIES WITH SIXTEEN-JOINTED ANTENNULES.

215. *Cyclops languidus*, G. O. Sars.

1863. *Cyclops languidus*, G. O. Sars, Forhandl. Vidensk.-Selsk. i
Christiania, p. 249.

1901-1902. ,, ,, Dr and Miss Sprague, *l. c.*, p. 255, pl. xxxi.
figs. 1-7.

Hab.—This species, which was observed for the first time in Scotland in Loch Doon, in Ayrshire,¹ was obtained in a pond near Tynehead, Midlothian, on 30th March 1901 (Dr and Miss Sprague).

(c) SPECIES WITH TWELVE-JOINTED ANTENNULES.

216. *Cyclops serrulatus*, Fischer.

1851. *Cyclops serrulatus*, Fischer, Bull. Soc. imp. des Natur.
Moscou, vol. xxiv. (2) p. 423, pl. x. figs: 22, 23, 26-31.

Hab.—The *Cyclops*, generally referred to as "*Cyclops serrulatus*, Fischer," is common in lochs and ponds throughout the district. A form with short furcal joints, which appears to be the var. *brachyurus* of *Cyclops varius*, Lilljeborg, has been observed in Duddingston Loch and other places. Another form with the furcal joints elongated, and which may be the var. *speratus* of the same species, has also been observed, but they approximate so closely to the species described by Fischer, that I prefer, for the present, to regard them as varieties of that species.

217. *Cyclops macrurus*, G. O. Sars.

1863. *Cyclops macrurus*, G. O. Sars, *l. c.*, p. 254.

1878. ,, ,, G. S. Brady, Monograph, vol. i. p. 111,
pl. xxiv. figs. 1-5.

Hab.—Loch Vennachar, Loch Katrine, Loch Leven, Black Loch near Loch Glow (Kinross-shire), and Loch Lubnaig.

¹ Cf. *Seventeenth Annual Report of the Fishery Board for Scotland*, pt. iii. p. 187 (1899).

(d) SPECIES WITH ELEVEN-JOINTED ANTENNULES.

218. *Cyclops affinis*, G. O. Sars.1863. *Cyclops affinis*, G. O. Sars, *l. c.*, p. 256.

1878. „ „ Brady, Monograph, vol. i. p. 112, pl. xv. figs. 11-14; pl. xxiv. figs. 10-15.

Hab.—Raith Lake, near Kirkcaldy, Fifeshire, 1890. Elf Loch, and ponds in Penicuik grounds, 1900 (Dr and Miss Sprague). Humbie Reservoir, near Winchburgh, June 1906 (W. Evans).

219. *Cyclops diaphanus*, Fischer.1853. *Cyclops diaphanus*, Fischer, *l. c.*, vol. xxvi. p. 93, pl. iii. figs. 6-12.1863. „ *nanus*, G. O. Sars, *l. c.*, p. 42.

1899. „ „ T. Scott, Seventeenth F. B. Rept., pt. iii. pp. 141, 172, 187.

1901. „ *diaphanus*, Lillj., Kongl. Sv. Vet.-Handlingar, vol. xxxv. No. 4, p. 63, pl. iv. figs 13-15.1901-1902. „ *nanus*, Dr and Miss Sprague, Trans. Edin. Field Naturalists, vol. iv. p. 256, pl. xxxia. figs. 7-12.

Hab.—This small species, first added to the British fauna in 1899, was obtained at Auchencorth Moss, Midlothian, in April 1901, by Dr and Miss Sprague.

(e) SPECIES WITH TEN-JOINTED ANTENNULES.

220. *Cyclops phaleratus*, Koch.1838. *Cyclops phaleratus*, Koch, Deutschl. Crustaceen, Heft 21, pl. ix.

1878. „ „ Brady, Monograph, vol. i. p. 116, pl. xxiii. figs. 7-13.

Hab.—Duddingston Loch; Lochgelly Loch and Raith Lake, Fifeshire. Marl-pit, Davidson's Mains, near Edinburgh, June 1900 (Dr and Miss Sprague). This species also occurs in some material from the marl-pit collected by myself in May 1888. Teith at Callander (Evans).

(f) SPECIES WITH EIGHT-JOINTED ANTENNULES.

221. *Cyclops fimbriatus*, Fischer.

1853. *Cyclops fimbriatus*, Fischer, *l. c.*, p. 94, pl. iii. figs. 19-28 and 30.

1878. „ *crassicornis*, Brady, Monograph, vol. i. p. 118, pl. xxiii. figs. 1-6.

Hab.—Moderately common in ponds and lakes throughout the district.

Genus (89) *Halicyclops*, Norman, 1903.222. *Halicyclops æquoreus* (Fischer).

1860. *Cyclops æquoreus*, Fischer, Abhandl. Math. Classe der Königl. Bayer. Akad. der Wiss. München, vol. viii. (3) p. 654, pl. xx. figs. 26-29.

1878. „ „ Brady, Monograph, vol. i. p. 119, pl. xix. figs. 8-10; pl. xxi. figs. 10-17.

Hab.—Cramond Island, in pools above high-tide mark, 1888; and in brackish-water pools at Aberlady in May 1895. Dr and Miss Sprague have also collected this species in a rock-pool on Cramond Island in July 1901.

Genus (90) *Euryte*, Philippi, 1843.223. *Euryte longicauda*, Philippi.

1843. *Euryte longicauda*, Philippi, Archiv. für Naturh., Jahrg. 9, p. 63, pl. iii. fig. 3, a-d.

1864. *Thorellia brunnea*, Boeck, Översigt Norges Copepoder, p. 26.

1872. „ „ Brady, *op. cit.*, vol. i. p. 95, pl. xvi. figs. 1-10.

Hab.—Moderately frequent throughout the estuary.

Family LICHOMOLGIDÆ.

For further information concerning the Lichomolgidæ and Hersiliidæ, the reader is referred to Dr Canu's work, *Les Copepodes du Boulonnais*.

Genus (91) *Lichomolgus*, Thorell, 1859.224. *Lichomolgus fucicolus*, G. S. Brady.

1880. *Lichomolgus fucicolus*, Brady, Monograph, vol. iii. p. 41,
pl. lxxxv. figs. 1-11.

Hab.—Firth of Forth; moderately rare. It has usually been obtained in shallow inshore water, amongst *Laminaria*, etc.

225. *Lichomolgus furcillatus*, Thorell.

1859. *Lichomolgus furcillatus*, Thorell, Om Krustaceer i Ascidier,
p. 74, taf. 13, fig. 30.

1880. „ „ Brady, *l. c.*, vol. iii. p. 49,
pl. lxxxviii. figs. 10-14.

Hab.—Collected near May Island and in the vicinity of Inchkeith, in the branchial cavity of large Ascidians; not common.

226. *Lichomolgus hirsutipes*, T. Scott.

1893. *Lichomolgus hirsutipes*, T. Scott, Eleventh F. B. Rept.,
pt. iii. p. 206, pl. iv. figs. 1-12.

Hab.—Taken with the dredge a short distance north of the Bass Rock, and off the North Craig; rather rare. On one occasion I found a considerable number of specimens adhering to the outside surface of the tubes of a large species of *Sabella* brought up on the hooks when line-fishing in 1895, in 15 to 20 fathoms, where the bottom consisted of mud. The Copepods were white, and therefore easily noticed on the mud-tubes made by the Annelids.¹

227. *Lichomolgus agilis* (Leydig).

1853. *Doridicola agilis*, Leydig, Zeitschr. f. Wiss. Zool., Bd. 4,
p. 377, taf. 14.

1892. *Lichomolgus agilis*, Canu, Les Copep. du Boulonnais, p. 228,
pl. xxii. figs. 1-14.

1892. „ „ *concinus*, T. Scott, Tenth F. B. Rept., pt. iii.
p. 261, pl. ii. figs. 25-33.

Hab.—Dredged off St Monans in 1891; rare. Taken also

¹ *Ann. Scot. Nat. Hist.* (October 1895), p. 238.

in the neighbourhood of Granton in 1893, on the branchial appendages of *Doris* (?) *tuberculatus*.¹

Genus (92) *Pseudanthessius*, Claus, 1889.

228. *Pseudanthessius gracilis*, Claus.

1889. *Pseudanthessius gracilis*, Claus, Arb. Zool. Inst. Wien., t. viii.
p. 344, taf. 4, figs. 1-7.

1893. „ „ T. and A. Scott, Ann. and Mag. Nat.
Hist. (6), vol. xii. p. 241, pl. xii.
figs. 15-20.

Hab.—Dredged off Musselburgh in 1891; not common.

229. *Pseudanthessius liber* (Brady and Robertson).

1876. *Lichomolgus liber*, B. and R., Brit. Assoc. Rept. for 1875,
p. 197.

1880. „ „ Brady, Monograph, vol. iii. p. 44,
pl. lxxxvi. fig. 197.

Hab.—Dredged very sparingly off the north end of Inchkeith in 1894. Also dredged in the neighbourhood of Inchkeith on 23rd May, and off North Craig on 4th July 1901.

230. *Pseudanthessius Thorelli* (Brady and Robertson).

1876. *Lichomolgus Thorelli*, B. and R., Brit. Assoc. Rept. for 1875,
p. 197.

1880. „ „ Brady, *l. c.*, vol. iii. p. 47, pl. lxxxviii.
figs. 1-9.

Hab.—Dredged very sparingly off St Monans in 1893-1894. Also off the east side of Inchkeith on 23rd May 1901; rare.

231. *Pseudanthessius Sauvagei*, Canu.

1891. *Pseudanthessius Sauvagei*, Canu, Bull. Sci. France et Belgique,
vol. xxiii. p. 481.

1894. „ „ T. and A. Scott, Ann. and Mag.
Nat. Hist. (6), vol. xii. p. 146.

Hab.—Dredged off St Monans in 1894; rare. During 1895 the fishery steamer "Garland" was engaged in some line-fishing experiments, when various things were brought up on the hooks, and the common Sea-Urchin, *Echinus*

¹ *Ann. and Mag. Nat. Hist.* (6), vol. ii. (March 1893) p. 212, pl. vii. figs. 12-15.

esculentus, among others. Some of these urchins I put into a bottle containing strong methylated spirit; after washing them in this spirit, an examination of the residue revealed a considerable number of specimens of this rare Copepod, and of little else.

Genus (93) *Modiolicola*, Aurivilius, 1883.

232. *Modiolicola insignis*, Aurivilius.

1883. *Modiolicola insignis*, Aur., Akad. Afhandl. Stockholm (1883), pp. 10 and 39, taf. 2, figs. 1-10; taf. 4, figs. 13-16.
 1893. „ „ T. Scott, Eleventh F. B. Rept., pt. iii. p. 207, pl. iv. figs. 13-24.

Hab.—Frequent in the shells of living “Horse Mussels,” *Mytilus modiolus*. The species was obtained in most of the large mussels examined.

Genus (94) *Herrmannella*, Canu, 1891.

233. *Herrmannella rostrata*, Canu.

1891. *Herrmannella rostrata*, Canu, Bull. Sci. France et Belgique, t. xxiii. p. 480.
 1892. „ „ idem, Les Copep. du Boulonnais, p. 236, pl. xxiv. figs. 1-13.
 1892. *Lichomolgus agilis*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. x. p. 201, pl. xv. figs. 1-14.

Hab.—Frequent in the shells of living Cockles, *Cardium edule*, found in the cockle-beds at Cramond.

234. *Herrmannella maxima* (I. C. Thompson).

1893. *Lichomolgus maximus*, I. C. T., Trans. L—pool Biol. Soc., vol. vii. p. 34, pl. xxxv.

Hab.—Frequent in the shells of the Clam, *Pecten opercularis*, dredged on the clam-beds to the east of Inchkeith, but this Copepod was first observed by I. C. Thompson in the shell of living *Pecten maximus*, hence its name. This is not a true *Lichomolgus*, and as it agrees very closely with *Herrmannella*, I place it meanwhile under that genus.

235. *Herrmannella arenicola* (G. S. Brady).

1872. *Boeckia arenicola*, Brady, Nat. Hist. Trans. Northumb. and Durham, vol. iv. p. 430.

1880. *Lichomolgus arenicolus*, idem, Monograph, vol. iii. p. 46, pl. lxxxvii. figs. 1-7.

1892. „ „ T. and A. Scott, Ann. Scot. Nat. Hist. (July 1892), p. 151, pl. vii. figs. 1-10.

Hab.—Dredged off St Monans in 1891; rare.

Genus (95) *Sabelliphilus*, M. Sars, 1862.236. *Sabelliphilus Sarsi*, Claparède.

1870. *Sabelliphilus Sarsi*, Clap., Ann. des Sci. Nat. (5), vol. xiii. p. 6, pl. vii.

1887. *Lichomolgus sabellæ*, I. C. Thompson, Proc. L—pool Biol. Soc., vol. ii. p. 68, pl. ii.

Hab.—Collected off St Monans, 7th July 1897, on the plumes of *Sabella* sp. (? *S. pavonina*); rare.

Family CLAUSIIDÆ.

Genus (96) *Hersiliodes*, Canu, 1888.237. *Hersiliodes aberdonensis* (T. and A. Scott).

1892. *Lichomolgus aberdonensis*, T. and A. Scott, Ann. Scot. Nat. Hist. (July 1892), p. 149, pl. vi. figs. 1-12.

Hab.—Dredged off the east side of Inchkeith on 23rd May 1901. The specimens from which the species was described in 1892 were obtained in a tow-net gathering collected in Aberdeen Bay.

238. *Hersiliodes littoralis* (T. Scott).

1892. *Lichomolgus littoralis*, T. Scott, Tenth F. B. Rept., pt. iii. p. 260, pl. x. figs. 1-9.

Hab.—Collected in the neighbourhood of Culross in 1891. This appears to be a rare species in the Forth estuary, and is probably parasitic on some other invertebrate.

Family ASTEROCHERIDÆ.¹Genus (97) *Asterocheres*, Boeck, 1859.239. *Asterocheres Lilljeborgi*, Boeck.

1859. *Asterocheres Lilljeborgi*, Boeck, Forhandl. Vid.-Selsk. Chr. (1859), p. 6, pl. ii. figs. 1-11.

1880. *Artotrogus Lilljeborgi*, Brady, Monograph, vol. iii. p. 64.

Hab.—Firth of Forth; rare (cf. Giesbrecht, *Asterocheridæ*, pp. 70 and 73).

240. *Asterocheres echinicola* (Norman).

1869. *Ascomyzon echinicola*, Norman, Brit. Assoc. Rept. for 1868, p. 300.

1880. *Cyclopicera lata*, Brady, *l. c.*, vol. iii. p. 56, pl. lxxxix. fig. 12; pl. xc. figs. 11-14.

1893. „ „ T. Scott, Eleventh F. B. Rept., pt. iii. p. 210, pl. iii. figs. 41 and 42.

Hab.—Obtained, in 1889, by washing a number of Sponges in methylated spirit; the sponges were dredged in Aberlady Bay.

241. *Asterocheres Boeckii* (G. S. Brady).

1880. *Artotrogus Boeckii*, Brady, *l. c.*, vol. iii. p. 60, pl. xci. figs. 1-9.

Hab.—Obtained at Granton Harbour in the water-passages of Sponges (*Chalina oculata*) growing on the walls of the pier; frequent.

¹ In *Fauna und Flora des Golfes von Neapel*, Monograph 25, the *Asterocheridæ*, by Dr W. Giesbrecht (1899), the Forth species belonging to this family are for the most part described and figured by the author; and he also refers to their Scottish habitats in his remarks on the distribution of the various forms. The reader should consult this fine work, which contains a full synonymy and other information.

Genus (98) *Dermatomyzon*, Claus, 1889.242. *Dermatomyzon nigripes* (Brady and Robertson).

1876. *Cyclopicera nigripes*, B. and R., Brit. Assoc. Rept. for 1875,
p. 197.

1880. „ „ Brady, *l. c.*, vol. iii. p. 54, pl. lxxxix.
figs. 1-11.

1892. „ „ T. Scott, Tenth F. B. Rept., pt. iii.
p. 267.

Hab.—Collected in the neighbourhood of May Island in 1892. Dredged at Station I. on 30th August 1894, and subsequently captured at a few other places, but always very sparingly.

Genus (99) *Rhynchomyzon*, Giesbrecht, 1895.243. *Rhynchomyzon purpurocinctum* (T. Scott).

1893 *Cyclopicera purpurocincta*, T. Scott, Eleventh F. B. Rept.,
pt. iii. p. 209, pl. iii. figs. 29-40.

Hab.—Dredged on the “Rath ground,” to the north of the Bass Rock, on 20th November 1889, and off the east side of Inchkeith on 23rd May 1901. This species, which appears to be rare in the Forth estuary, has a wide distribution, having been recorded not only from various parts of the British seas, but also from the Gulf of Naples.

Genus (100) *Collocheres*, Canu, 1893.244. *Collocheres gracilicauda* (G. S. Brady).

1880. *Cyclopicera gracilicauda*, Brady, *l. c.*, vol. iii. p. 58, pl.
lxxxiii. figs. 1-10.

Hab.—Dredged off St Monans; rare.

Genus (101) *Scottomyzon*, Giesbrecht, 1897.245. *Scottomyzon gibberum* (T. and A. Scott).

1894. *Dermatomyzon gibberum*, T. and A. Scott, Ann. and Mag.
Nat. Hist. (6), vol. xii. p. 144,
pl. ix. figs. 10-14.

1894. „ „ T. Scott, Twelfth F. B. Rept.,
pt. iii. p. 260, pl. x. figs. 26-34.

Hab.—Dredged in the neighbourhood of the Bass Rock in

1893. In 1895 a number of specimens—adults and young—were obtained by washing specimens of the common Starfish, *Asterias rubens*, in a bottle containing methylated spirit, and afterwards examining the residue. These specimens were of a brick-red colour on the back, but nearly white beneath. Their normal habitat is apparently on this kind of starfish.

Genus (102) *Acontiophorus*, Brady, 1880.

246. *Acontiophorus scutatus* (Brady and Robertson).

1873. *Solenostoma scutatum*, B. and R., Ann. and Mag. Nat. Hist. (4), vol. xii. p. 141.

1880. *Acontiophorus scutatus*, Brady, *l. c.*, vol. iii. p. 69, pl. xc. figs. 1-10.

Hab.—Collected near Inchkeith in 1888. Off Musselburgh, dredged 30th May 1891, and subsequently collected in various other parts of the estuary; moderately frequent.

247. *Acontiophorus ornatus* (Brady and Robertson).

1876. *Ascomyzon ornatum*, B. and R., Brit. Assoc. Rept. for 1875, p. 197.

1880. *Acontiophorus armatus*, Brady, *l. c.*, vol. iii. p. 71, pl. lxxxvii. figs. 8-15.

Hab.—Dredged off North Craig on 4th July 1901; rare.

Genus (103) *Scottocheres*, Giesbrecht, 1897.

248. *Scottocheres elongatus* (T. and A. Scott).

1894. *Acontiophorus elongatus*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xii. p. 145, pl. ix. figs. 15-20.

1898. *Scottocheres elongatus*, T. Scott, Sixteenth F. B. Rept., pt. iii. p. 278, pl. xiii. figs. 10-21.

Hab.—Obtained in 1893 in material dredged in the neighbourhood of the Bass Rock; frequent.

Genus (104) **Cribropontius**, Giesbrecht, 1899.

249. *Cribropontius Normani* (Brady and Robertson).

1876. *Dyspontius Normani*, B. and R., Brit. Assoc. Rept. for 1875, p. 197.

1880. *Artotrogus Normani*, Brady, *l. c.*, vol. iii. p. 63, pl. xci. figs. 12-15; pl. xcii. fig. 10.

1897. *Bradypontius Normani*, T. Scott, Fifteenth F. B. Rept., pt. iii. p. 154, pl. ii. figs. 1 and 2; pl. iii. figs. 1-11.

Hab.—Taken off the North Craig in material dredged on 4th July 1901; rare. This large and sometimes highly-coloured species appears to be rare in Firth of Forth.

Genus (105) **Bradypontius**, Giesbrecht, 1895.

250. *Bradypontius magniceps* (G. S. Brady).

1880. *Artotrogus magniceps*, Brady, *l. c.*, vol. iii. p. 61, pl. xciii. figs. 1-9.

Hab.—Dredged west of Queensferry in 1888, and at the north-west end of Inchkeith on 23rd May 1901. Several fine specimens were obtained in material dredged off the North Craig on 4th July 1901, when a few of the females were found with ovisacs attached.

251. *Bradypontius papillatus* (T. Scott).

1888. *Artotrogus papillatus*, T. Scott, Sixth F. B. Rept., pt. iii. p. 232, pl. viii. figs. 7-12.

1899. *Bradypontius chelifera*, Giesbrecht, Fauna u. Flora Golfes v. Neapel, Monogr. 25 (Asterocheridæ), p. 88, pl. vi. figs. 15-25.

1899. *Bradypontius papillatus*, T. Scott, Seventeenth F. B. Rept., pt. iii. p. 262, pl. xi. fig. 21; pl. xii. figs. 7-15

Hab.—Dredged in the neighbourhood of Inchkeith in 1888; apparently very rare. The specimen from which the description and drawings were prepared is the only one that has yet been observed in the Forth estuary. This specimen is, I think, certainly identical with the *Bradypontius chelifera* described by Dr Giesbrecht in his fine work on the Asterocheridæ.

Genus (106) *Dyspontius*, Thorell, 1859.252. *Dyspontius striatus*, Thorell.

1859. *Dyspontius striatus*, Thorell, Om Krustaceer i Acidier, p. 81,
pl. xiv. fig. 22.

1880. „ „ Brady, *l. c.*, vol. iii. p. 66, pl. xcii. figs.
1-12.

Hab.—Dredged at the north-west end of Inchkeith on 23rd May 1901. *D. striatus* has not previously been noticed within the limits of the Forth estuary, but it is a widely distributed species in our seas, though not very plentiful.

253. *Dyspontius curticaudatus*, T. Scott.

1905. *Dyspontius curticaudatus*, T. Scott, Twenty-third F. B. Rept.,
pt. iii. p. 148, pl. xiii. figs. 1-10.

Hab.—Dredged in the vicinity of Culross, a few miles above Queensferry; rare.

Genus (107) *Parartotrogus*, T. and A. Scott, 1893.254. *Parartotrogus Richardi*, T. and A. Scott.

1893. *Parartotrogus Richardi*, T. and A. Scott, Ann. and Mag. Nat.
Hist. (6), vol. xi p. 211, pl. vii. figs. 1-11.

Hab.—Dredged near Fidra in 1889, but not recorded till 1893. It has subsequently been dredged off St Monans, in Largo Bay, and one or two other places, but nowhere very common.

Family CORYCÆIDÆ.

Genus (108) *Corycæus*, Dana, 1845.255. *Corycæus anglicus*, Lubbock.

1857. *Corycæus anglicus*, Lubb., Ann. and Mag. Nat. Hist. (2),
vol. xx., pl. xi. figs. 16-19.

1880. „ „ Brady, *l. c.*, vol. iii p. 34, pl. lxxxii.
figs. 16-19 *et seq.*

1900. „ „ T. Scott, Eighteenth F. B. Rept., pt. iii.
p. 397, pl. xiii. figs. 1-14.

Hab.—Dredged in the “Fluke Hole” off St Monans; also washed from trawl refuse collected off West Wemyss,

October 1895. This species appears to be of rare occurrence in the Forth estuary.

Family NICOTHOIDÆ.

Genus (109) *Nicothoë*, Audouin and Milne-Edwards, 1826.

256. *Nicothoë astaci*, Aud. and M.-Edwards.

1826. *Nicothoë astaci*, Aud. and M.-Edwards, Ann. Sci. Nat., 1st ser., vol. ix. p. 345, taf. 49, figs. 1-9.

Hab.—Found adhering to the gills of a Lobster sent from Dunbar. I am indebted to my colleague, Dr H. C. Williamson, for the specimens of the *Nicothoë*. The distribution of the parasite appears to be co-extensive with that of its host.

Family ERGASILIDÆ.

Genus (110) *Bomolochus*, Nordmann, 1832.

257. *Bomolochus soleæ*, Claus.

1864. *Bomolochus soleæ*, Claus, Zeitschrift für Wissenschaft, Zool., vol. xiv. p. 374, pl. xxxv.

1893. „ „ T. Scott, Eleventh F. B. Rept., pt. iii. p. 212, pl. v.

1902. „ „ idem, Twentieth F. B. Rept., pt. iii. p. 288, pl. xiii. figs. 13-18.

Hab.—Firth of Forth; found on the back of a Black Sole, *Solea vulgaris*, and among dredged material, but in this case the specimens must, in some way, have been detached from the fish. Found in the nostrils of a Cod-fish by John Lindsay, Edinburgh. This *Bomolochus* has been obtained in the nasal fossæ of several kinds of fishes, but it is most frequent in those of the cod.

258. *Bomolochus onosi*, T. Scott.

1902. *Bomolochus onosi*, T. Scott, Twentieth F. B. Rept., pt. iii. p. 289, pl. xiii. figs. 19-22.

Hab.—Firth of Forth, on the inside of the gill-covers of a Five-bearded Rockling, *Onos mustelus* (Linn.), captured in the estuary in May 1901. This *Bomolochus* has also been

obtained on a specimen of the same kind of fish captured off Kinnaird Head in July 1901.

Genus (111) *Thersitina*, Norman, 1905.

(Syn. *Thersites*, Pagenstecher, 1861, preoccupied by Spence Bate in 1857 for a genus of Amphipods.)

259. *Thersitina gasterostei* (Pagenstecher).

1861. *Thersites gasterostei*, Pagenst., Arch. f. Naturh., vol. xvii. p. 118, pl. vi. figs. 1-9.

1899. *Ergasilus gasterostei*, Bassett-Smith, Proc. Zool. Soc. Lond. (April 1899), p. 444.

1900. *Thersites gasterostei*, T. Scott, Eighteenth F. B. Rept., pt. iii. p. 146, pl. v. figs. 1-7.

Hab.—On the inside of the gill-covers of a Three-spined Stickleback, *Gasterosteus aculeatus*, captured in the river Forth, near Alloa, in February 1896. I have also taken this minute Copepod on the same species of fish captured in a small loch in Barra, Outer Hebrides; and in brackish-water ditches at Aberdeen, as well as on a Fifteen-spined Stickleback, *Gasterosteus spinachia*, captured in Loch Etive, on the west of Scotland.

Division NOTODELPHYOIDA.

Family NOTODELPHYDÆ.

Genus (112) *Notodelphys*, Allman, 1847.

260. *Notodelphys Allmani*, Thorell.

1860. *Notodelphys Allmani*, Thorell, Bidrag til Känned. Krust. Art. Slägtet Ascidia, p. 31, pl. i., and pl. ii. fig. 1.

1878. „ „ „ Brady, Monograph, vol. i. p. 126, pl. xxv. figs. 1-10.

Hab.—In the branchial cavity of Ascidiæ found adhering to the walls of Granton Harbour, and dredged in the neighbourhood of Inchkeith and elsewhere; moderately frequent.

261. *Notodelphys agilis*, Thorell.

1860. *Notodelphys agilis*, Thorell, *l. c.*, p. 40, pl. iv.; pl. v. fig. 6.

1878. „ „ „ Brady, *l. c.*, vol. i. p. 130, pl. xxvi. figs. 1-10.

Hab.—In the branchial cavity of Ascidiæ found adhering to the pier at Granton.

Family DOROPYGIDÆ.

Genus (113) *Doropygus*, Thorell, 1859.262. *Doropygus Normani*, G. S. Brady.1878. *Doropygus Normani*, Brady, *l. c.*, vol. i. p. 136, pl. xxxii. figs. 1-14.

Hab.—In the branchial cavity of Ascidians dredged off Musselburgh, not uncommon, but apparently rare in other parts of the estuary.

263. (?) *Doropygus porcicauda*, G. S. Brady.1878. *Doropygus porcicauda*, Brady, *l. c.*, vol. i. p. 138, pl. xxvii. figs. 1-9; pl. xxxiii. figs. 14-16.

1888. (?) ,, ,, T. Scott, Sixth F. B. Rept., pt. iii. p. 239.

Hab.—A specimen that appeared to belong to this species was obtained among some material dredged near Inchkeith in 1887, but as it is not now in my collection, I am unable to verify the record.

Family ASCIDICOLIDÆ.

Genus (114) *Ascidicola*, Thorell, 1859.264. *Ascidicola rosea*, Thorell.1859. *Ascidicola rosea*, Thorell, *l. c.*, p. 59, pl. ix.; pl. x. fig. 13.1878. ,, ,, Brady, *l. c.*, vol. i. p. 145, pl. xxx. figs. 1-10.

Hab.—Obtained occasionally in the branchial chambers of Ascidians dredged in the neighbourhood of Inchkeith, off Musselburgh, and at a few other places in the estuary.

Genus (115) *Aplostoma*, Canu, 1886.

Syn. *Enterocola*, T. Scott, 1892 (not *Enterocola*, P. J. v. Beneden).

265. *Aplostoma affinis*, T. Scott, nov. nom.1869. (?) *Enterocola eruca*, Norman, Last Report of Dredging among the Shetland Isles (Brit. Assoc. Rept. for 1868), p. 300.

1892. ,, ,, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. x. p. 203, figs. 1-11.

Hab.—A few specimens of this curious species were

obtained within the intestines of Ascidians found frequently adhering to the inside of the valves of dead mussels and other shells, dredged in various parts of the estuary. These specimens were mistakenly ascribed to *Enterocola*, van Beneden, but they more nearly resemble *Aplostoma brevicauda*, Canu. They differ, however, in some respects from that species, as shown by the description and drawings. I have therefore named the species as above.

Family LAMIPPEIDÆ.

Genus (116) *Lamippe*, Bruzelius, 1859.

266. *Lamippe proteus*, Claparède.

1867. *Lamippe proteus*, Claparède, Ann. des Sci. Nat., ser. 5, vol. viii. p. 23, pl. v.

1895. *Alcyoncola fusiformis*, T. and A. Scott, Ann. and Mag. Nat. Hist. (6), vol. xvi. p. 357, pl. xvi. figs. 10-14; pl. xvii. fig. 13.

1896. *Lamippe* (?) *proteus*, T. Scott, Fourteenth F. B. Rept., pt. ii. p. 164.

Hab.—On *Alcyonium digitatum*, dredged in various parts of the estuary; frequent. This curious species appears to live in the cells of the polyps. If an *Alcyonium* be broken into pieces, and the fragments washed in a bottle containing methylated spirit, numbers of the *Lamippe* will generally be found amongst the sediment.

267. *Lamippe Forbesi*, T. Scott.

1896. *Lamippe* sp., T. Scott, Fourteenth F. B. Rept., pt. iii. p. 164, pl. iv. figs. 9-13.

1901. *Lamippe Forbesi*, T. Scott, Nineteenth F. B. Rept., pt. iii. p. 256.

Hab.—This, which is a larger and apparently a much rarer species than *L. proteus*, has been found along with it on *Alcyonium digitatum*, dredged in the Firth of Forth. It has been observed on the same *Alcyonium* from other parts of the Scottish coast, and my son has also obtained it in Liverpool Bay.¹

¹ Cf. Tenth Ann. Rept. Liverpool Marine Biological Committee, p. 21 (1897).

Genus (117) *Jeanella*, T. Scott, *F. B. Rept.*, 1904,
pt. iii. p. 258.

(Syn. *Platypsyllus*, T. Scott, 1902, a name preoccupied
by Dr Ritsema and Prof. Westwood in 1869.)

268. *Jeanella minor*, T. Scott.

1902. *Platypsyllus minor*, T. Scott, Twentieth F. B. Rept., pt. iii.
p. 455, pl. xxv. figs. 15, 16.

Hab.—This somewhat remarkable species was obtained in a gathering dredged in about 8 fathoms in the neighbourhood of the North Craig on 4th July 1901. Only a few specimens—all females, and one or two carrying ovisacs—were observed.

Division MONSTRILLOIDA.

Family MONSTRILLIDÆ.

Genus (118) *Monstrilla*, Dana, 1848.

269. *Monstrilla anglica*, Lubbock.

1857. *Monstrilla anglica*, Lubbock, Ann. and Mag. Nat. Hist. (2),
vol. xx. p. 409, pl. x. figs. 7 and 8.

1904. „ „ T. Scott, Twenty-second F. B. Rept.,
pt. iii. p. 246, pl. xiii. fig. 13; pl. xiv.
figs. 12-14.

Hab.—Captured to the west of May Island, with a tow-net, on 26th July 1901, and also to the east of May Island on 20th August 1903, but only females were observed in both gatherings.

270. *Monstrilla longicornis*, I. C. Thompson.

1890. *Monstrilla longicornis*, I. C. Thompson, Trans. L—pool Biol.
Soc., vol. iv. p. 119, pl. iv. figs. 1, 2, and 4 (♂).

1892. „ *longiremis*, Giesb., Pelag. Copep. des Golfes v.
Neapel, p. 589, pl. xlvi. figs. 10, 14, 22, 37,
and 41 (♀).

1904. „ *longicornis*, T. Scott, *l. c.*, p. 244, pl. xiii. figs. 1-7
(♂ and ♀).

Hab.—This species was taken in the estuary very sparingly in 1890, 1891, 1893, 1894, and 1901.

271. *Monstrilla gracilicauda*, Giesbrecht.

1892. *Monstrilla gracilicauda*, Giesb., *l. c.*, p. 587, pl. xlvi. figs. 9, 16, 18 *et seq.*

1904. „ „ „ T. Scott, Twenty-second F. B. Rept., pt. iii. p. 245, pl. xiii. figs. 8-10; pl. xiv. fig. 15.

Hab.—Taken with tow-net above Queensferry, 26th June 1890; and off Musselburgh, 29th September 1892. Also captured east of May Island on 20th August 1903. All the specimens were females.

272. *Monstrilla dubia*, T. Scott.

1904. *Monstrilla dubia*, T. Scott, Twenty-second F. B. Rept., pt. iii. p. 247, pl. xiii. fig. 14; pl. xiv. figs. 16-18.

Hab.—This species was captured east of Inchkeith on 14th August 1891, and is apparently rare. It resembles *Monstrilla Danæ*, Claparède, but that author shows only three setæ instead of four on each furcal joint; there are also one or two other differences.

Genus (119) *Thaumaleus*, Kröyer, 1849.273. *Thaumaleus rigidus* (I. C. Thompson).

1888. *Cymbasoma rigidum*, I. C. Thompson, Jour. Linn. Soc. (Zool.), vol. xx. p. 154, pl. xiii. figs. 1-4.

1892. *Thaumaleus Claparèdei*, Giesb., *op. cit.*, p. 381, pl. xlvi. figs. 5, 15, 21, 26.

1904. „ „ *rigidus*, T. Scott, Twenty-second F. B. Rept., pt. iii. p. 248, pl. xiii. figs. 15-17; pl. xiv. fig. 19.

Hab.—Captured off St Monans, by tow-net, on 6th September 1890; apparently rare.

Division CALIGOIDA.

Family CALIGIDÆ.

Genus (120) *Caligus*, O. F. Müller, 1785.274. *Caligus curtus*, Müller.

1785. *Caligus curtus*, Müller, Entomostraca, p. 130, pl. xxi. fig. 1.

1850. „ *diaphanus*, Baird, British Entomostraca, p. 269, pl. xxxii. fig. 1 (♂).

1850. „ *Mülleri*, idem, *l. c.*, p. 271, pl. xxxii. figs. 4 and 5.

Hab.—Frequent on different kinds of fishes, especially

Gadoids. The male *Caligus*, which Dr Baird describes and figures in his *British Entomostraca* under the name of *Caligus diaphanus*, Nordmann, is not Nordmann's species, but is the male of *C. curtus*, Müller.

275. *Caligus rapax*, M.-Edwards.

1840. *Caligus rapax*, M.-Edw., Hist. Nat. Crust., vol. iii. p. 453, pl. xxxviii. fig. 9.

Hab.—Common on several kinds of fishes, especially Gadoids, as well as on flat fishes, Skate, Dogfishes, and others. From Short Sunfish, North Berwick, September 1905 (W. Evans). It is also frequently captured by tow-net in the open sea.

276. *Caligus minimus*, Otto.

1828. *Caligus minimus*, Otto, Beschreib. neuer Crust., Nov. Act. Acad. Cæs. Leop., vol. xiv. p. 354, pl. xxii.

1901. „ „ A. Scott, Trans. L—pool Biol. Soc., vol. xv. p. 349, pl. i. figs. 1-8.

Hab.—Found inside the throat of a Bass, *Labrax lupus*, Cuv. and Valenc., captured in the Forth, above Queensferry, in February 1903.

277. *Caligus diaphanus*, Nordmann.

1832. *Caligus diaphanus*, Nordm., Mikrog. Beiträge, vol. ii. p. 26.

1894. „ *isonyx*, T. Scott, Twelfth F. B. Rept., pt. iii. p. 310 (not *C. isonyx*, Stp. and Lützk.).

1900. „ *diaphanus*, idem, Eighteenth F. B. Rept., pt. iii. p. 149, pl. v. figs. 20-25.

Hab.—Taken in the gill-cavity of Grey Gurnards, *Trigla gurnardus*, captured in the estuary; not very rare.

Genus (121) *Pseudocaligus*, A. Scott, 1900.

278. *Pseudocaligus brevipedis* (Bassett-Smith).

1896. *Caligus brevipedis*, Bassett-Smith, Ann. Nat. Hist. (6), vol. xviii. p. 11, pl. iii. fig. 1.

1901. *Pseudocaligus brevipedis*, A. Scott, Trans. L—pool Biol. Soc., vol. xv. p. 350, pl. ii. figs. 1-6.

1902. „ „ T. Scott, Twentieth F. B. Rept., pt. iii. p. 291.

Hab.—Found inside the throat of a Three-bearded Rockling,

Onos tricirratatus, captured at Dunbar in 1892. In this species the fourth pair of feet are more rudimentary than in typical *Caligus*.

Genus (122) *Lepeophtheirus*, Nordmann, 1832.

279. *Lepeophtheirus pectoralis* (O. F. Müller).

1776. *Lernæa pectoralis*, Müller, Zool. Dan. Prodr., vol. i. p. 41, pl. xxxiii. fig. 7.

1850. *Lepeophtheirus pectoralis*, Baird, Brit. Entom., p. 275, pl. xxxii. fig. 10.

1900. „ „ T. Scott, Eighteenth F. B. Rept., pt. iii. p. 150, pl. v. figs. 26-31.

Hab.—Frequent under the pectoral fins of Plaice, Flounders, and Dabs.

280. *Lepeophtheirus Thompsoni*, Baird.

1850. *Lepeophtheirus Thompsoni*, Baird, Brit. Entom., p. 278, pl. xxxiii. fig. 2.

1900. „ „ T. Scott, *l. c.*, p. 152, pl. v. figs. 43-45.

1900. „ „ *obscurus*, idem, *l. c.*, p. 153, pl. vi. figs. 16-19.

Hab.—Frequent on the gills of Turbot, *Bothus maximus*, captured off St Monans. The form I referred doubtfully to *L. obscurus*, Baird, appears to be identical with *L. Thompsoni*; it is sometimes common on the gills of the Brill, *Bothus rhombus*.

281. *Lepeophtheirus pollachii*, Bassett-Smith.

1896. *Lepeophtheirus pollachii*, Bassett-Smith, Ann. and Mag. Nat. Hist. (6), vol. xviii. p. 12, pl. iv. fig. 1.

1900. „ „ T. Scott, *op. cit.*, p. 153, pl. vi. figs. 9-15.

Hab.—Firth of Forth, on the gills of Lythe, *Gadus pollachius*, 8th February 1895; apparently not very rare.

282. *Lepeophtheirus salmonis* (Kröyer).

1837. *Caligus salmonis*, Kröyer, Naturhist. Tidssk., vol. i., pl. vi.
fig. 7 a-c; vol. ii. (1838) pp. 13, 18.

1847. ,, *Stromi*, Baird, Trans. Berw. Naturalists Club.

1850. *Lepeophtheirus Stromi*, Baird, Brit. Entom., p. 274, pl. xxxii.
figs. 8 and 9.

1850. ,, *obscurus*, idem, ibidem, p. 277, pl. xxxii. fig. 11.

Hab.—On Salmon caught in the salmon nets in Largo Bay and other parts of the estuary; frequent.

Genus (123) *Trebius*, Kröyer, 1838.283. *Trebius caudatus*, Kröyer.

1838. *Trebius caudatus*, Kröyer, Naturhist. Tidsskr. (1838), R. i. B. ii.
p. 30, pl. i. fig. 4.

1900. ,, ,, T. Scott, Eighteenth F. B. Rept., pt. iii.
p. 155, pl. vi. figs. 20-26.

Hab.—Taken on the backs of grey (or blue) Skates, *Raia batis*; moderately frequent.

Genus (124) *Cecrops*, Leach, 1816.284. *Cecrops Latreillei*, Leach.

1816. *Cecrops Latreillei*, Leach, Encycl. Brit., Suppl. i., pl. xx.
figs. 1-5.

1850. ,, ,, Baird, Brit. Entom., p. 293, pl. xxxiv.
fig. 1.

Hab.—Found on the gills of a Short Sunfish, captured in the estuary in October 1890 (A. Scott). *Cecrops* appears to be a common parasite on the gills of this species of Sunfish.

Genus (125) *Orthagoriscola*, Franz Poche, 1902.

(Syn. *Læmargus*, Kröyer, 1838, name preoccupied for a genus of Fishes.)

285. *Orthagoriscola muricata* (Kröyer).

1850. *Læmargus muricatus*, Baird, Entomotraca, p. 295, pl. xxxiv.
figs. 3 and 4.

1900. ,, ,, T. Scott, Eighteenth F. B. Rept., pt. iii.
p. 158, pl. vi. figs. 39-42.

1902. *Orthagoriscola muricata*, Franz Poche, Zool. Anzeiger,
vol. xxvi. p. 15.

Hab.—On the same Sunfish with the *Cecrops* just recorded

not, however, attached to the gills, but burrowing in hollows formed in the flesh of the fish, behind the anal fin (A. Scott). Off a Short Sunfish captured at North Berwick, 28th September 1905 (W. Evans).¹ This appears to be a rarer species than the *Cecrops*.

Division LERNÆOIDA.

Family LERNÆIDÆ.

Genus (126) *Lernæa*, Linné, 1767.

286. *Lernæa branchialis*, Linné.

1767. *Lernæa branchialis*, Linn., Systema Naturæ, ed. 12, vol. i. pt. ii. p. 1092.

1850. ,, ,, Baird, Brit. Entom., p. 344, pl. xxxv. fig. 12.

Hab.—Frequent on the gills of Whitings and some other Gadoids.

Genus (127) *Lernæenicus*, Lesueur, 1824.

287. *Lernæenicus spratta* (Sowerby).

1806. *Lernæa spratta*, Sowerby, Brit. Miscell., vol. ii. p. 17, pl. lxxviii.

1900. *Lernæenicus sprattæ*, T. Scott, Eighteenth F. B. Rept., pt. iii. p. 161, pl. vii. figs. 7-10.

Hab.—On the eye of a Sprat, *Clupea spratta*, seen in one of the Leith Docks (J. Scott, August 1890). This Copepod is apparently seldom met with in the Firth of Forth, and this is the only record of its occurrence in the estuary known to me.

Genus (128) *Hæmobaphes*, Steenstrup and Lütken, 1861.

288. *Hæmobaphes cyclopterinus* (Fabricius).

1780. *Lernæa cyclopterina*, Fabr., Fauna Grönlandica, p. 337.

1900. *Hæmobaphes cyclopterinus*, T. Scott, Eighteenth F. B. Rept., pt. iii. p. 162, pl. vii. fig. 14.

Hab.—Found on the gills of a Pogge, *Agonus cataphractus*, captured at Dunbar; and on a Butterfish, *Pholis gunnellus*, taken near Fidra in 1901.

¹ *Ann. Scot. Nat. Hist.*, 1906, p. 57.

Family CHONDRA CANTHIDÆ.

Genus (129) *Oralien*, Bassett-Smith, 1899.289. *Oralien asselinus* (Linné).1761. *Lernæa asselina*, Linn., Fauna Suecica, 2101.1850. *Lernentoma asselina*, Baird, Brit. Entom., p. 329, pl. xxxv.
fig. 4.

Hab.—Found on the gills of the Grey Gurnard, *Trigla gurnardus*; not uncommon. What appears to be the same species is also found on the gills of Plaice and one or two other fishes.

Genus (130) *Chondracanthus*, De la Roche, 1811.290. *Chondracanthus cornutus* (O. F. Müller).1776. *Lernæa cornuta*, Müll., Zool. Dan. Prodr., vol. i., pl. xxxiii.
fig. 6.1850. *Lernentoma cornuta*, Baird, *l. c.*, p. 328, pl. xxxv. fig. 2.

Hab.—Found on the gills of Plaice, *Pleuronectes platessa*; not uncommon.

291. *Chondracanthus clavatus*, Bassett-Smith.1896. *Chondracanthus clavatus*, Bassett-Smith, Ann. and Mag. Nat.
Hist. (6), vol. xviii. p. 13, pl. v. fig. 6.

Hab.—Found on the gills of Lemon Dabs, *Pleuronectes microcephalus*, especially large fishes.

292. *Chondracanthus fluræ*, Krøyer.1864. *Chondracanthus fluræ*, Krøyer, Naturh. Tidsskr., R. iii. B. ii.
p. 323, pl. xiii. fig. 6.1900. „ „ T. Scott, Eighteenth F. B. Rept.,
pt. iii. p. 166, pl. vii. figs. 32-34.

Hab.—Found on the gills of Long Rough Dabs, *Drepanopsetta platessoides*, captured in the estuary in 1891, and at Station III. in May 1901.

293. *Chondracanthus soleæ*, Kröyer.

1839. *Chondracanthus soleæ*, Kröyer, Naturh. Tidsskr. (1838-39), p. 139, pl. iii. fig. 4.

Hab.—Found on the gills of the Black Sole, *Solea vulgaris*, captured in the estuary.

294. *Chondracanthus depressus*, T. Scott.

1905. *Chondracanthus depressus*, T. Scott, Twenty-third F. B. Rept., pt. iii. p. 114, pl. vi. figs. 7-13.

Hab.—Found on the gills of Dabs, *Pleuronectes limanda*, Linn., captured in the estuary; not very common.

295. *Chondracanthus merluccii* (Holten).

(?) 1802. *Lernæa merluccii*, Holten, Mem. Soc. Hist. Nat. Copenhagen, vol. v., pl. viii. fig. 2.

1837. *Chondracanthus merluccii*, Kröyer, Naturh. Tidsskr., vol. i. p. 278, pl. iii. fig. 9a-d.

Hab.—Found in the gill-cavity of a Hake, *Merluccius vulgaris*, landed at Newhaven in February 1885. This is a moderately common parasite on the Hake; it may frequently be found clinging to the roof and sides of the mouth, and sometimes on the underside of the tongue, as well as inside the gill-covers.

296. *Chondracanthus lophii*, Johnston.

1836. *Chondracanthus lophii*, Johnston, Lond. Mag. Nat. Hist., p. 81, fig. 16.

1850. *Lernentoma lophii*, Baird, Brit. Entom., p. 330, pl. xxxv. fig. 3.

Hab.—Common in the gill-chambers of the Angler-fish, *Lophius piscatorius*, captured in the estuary. This is one of the most common species of the Chondracanthi I have observed in the Firth of Forth.

297. *Chondracanthus zeii*, De la Roche.

1811. *Chondracanthus zeii*, De la Roche, Nouv. Bull. de Soc. Philom., vol. ii. p. 270, pl. ii. fig. 2.

1850. *Lernentoma zeii*, Baird, Brit. Entom. p. 327, pl. xxxv. fig. 1.

Hab.—Found on the gills of a John Dory, *Zeus faber*, captured in the estuary in 1891.

Family LERNÆOPODIDÆ.

Genus (131) *Charopinus*, Kröyer, 1863.298. *Charopinus Dalmanni* (Retzius).

1829. *Lernæa Dalmanni*, Retz., Kongl. Vetensk. Acad. Handl. för 1829; Frorieps Notizen, Bd. 29, N. 617, p. 6.

1900. *Charopinus Dalmanni*, T. Scott, *op. cit.*, p. 169, pl. viii. figs. 6-10.

Hab.—In the nasal fossæ of a large Grey Skate, *Raia batis*, captured near May Island in 1891. This curious parasite is frequent in the nasal fossæ of large skates, but seldom in small specimens.

Genus (132) *Brachiella*, Cuvier, 1817.299. *Brachiella insidiosa*, Heller.

1865. *Brachiella insidiosa*, Heller, Reise der Novara, p. 239, pl. xxiv. fig. 1.

1896. „ „ Bassett-Smith, Ann. and Mag. Nat. Hist. (6), vol. xviii. p. 14, pl. vi. fig. 2.

Hab.—Found on the gills of a Hake captured in the estuary. This species is less common than the *Chondranchus merluccii*, which is also found on the Hake.

300. *Brachiella triglæ*, Claus.

1860. *Brachiella triglæ*, Claus, Zur Morp. der Copepoden, pl. i. fig. 6.

1896. „ „ Bassett-Smith, Jour. M. B. Assoc. Plymouth, p. 163.

Hab.—Found on *Trigla lineata* captured at Station VIII. in September 1897.

Genus (133) *Anchorella*, Cuvier, 1817.301. *Anchorella emarginata*, Kröyer.

1837. *Anchorella emarginata*, Kröyer, Naturh. Tidsskr., R. i. B. i. p. 287, pl. iii. fig. 7.

1900. „ „ T. Scott, Eighteenth F. B. Rept., pt. iii. p. 176, pl. viii. figs. 49-51.

Hab.—Found on the gills of a Twaite Shad, *Clupea finta*, captured near Dunbar in February 1897. This appears to be a rare species.

302. *Anchorella rugosa*, Kröyer.1837. *Anchorella rugosa*, Kröyer, *l. c.*, p. 294, pl. iii. fig. 6.

1850. „ „ Baird, Brit. Entom., p. 338, pl. xxxv. fig. 8.

Hab.—Found on gills and gill-covers and inside the throat of the Cat- or Wolf-fish, *Anarrhichas lupus*; not uncommon.

303. *Anchorella uncinata* (O. F. Müller).1776. *Lernæa uncinata*, Müll., Zool. Dan. Prodr., vol. i., pl. xxxiii. fig. 2.1850. *Anchorella uncinata*, Baird, *l. c.*, p. 337, pl. xxxv. fig. 9.

Hab.—Found on the gills and inside the throat of Cod-fish, and probably on other Gadoids, not uncommon; but the *Anchorellæ* of these other Gadoids require further study, as they may not all belong to the same species.

304. *Anchorella brevicola*, M.-Edwards.1840. *Anchorella brevicola*, M.-Edw., Hist. Nat. Crust., vol. iii. p. 418.

1901. „ „ T. Scott, Nineteenth F. B. Rept., pt. iii. p. 135, pl. viii. figs. 11-16.

Hab.—Found on a Haddock. The parasite was attached on the under side of the fish, and near the anal fin. Only one specimen was observed.

I add here two other interesting parasitic Copepods, each of which represents a different family, viz.:—

Family HERPYLLOBIIDÆ.

Genus (134) *Salenskya*, Giard and Bonnier, 1893.305. *Salenskya tuberosa*, Giard and Bonnier.1893. *Salenskya tuberosa*, G. and B., Compt.-rend. de l'Acad. des Sci. (25th Sept. 1893).

1895. „ „ idem, Bull. Sci. de la France et de la Belgique, vol. xxv. p. 472, pl. xiii.

1902. „ „ T. Scott, Twentieth F. B. Rept., pt. iii. p. 474, pl. xxv. figs. 17-22.

Hab.—Found in the marsupium of specimens of *Ampelisca*

spinipes, Boeck; several examples of this species of Amphipod were infested with the parasite.

Family CHONIOSTOMATIDÆ.

Genus (135) *Aspidœcia*, Giard and Bonnier, 1889.

306. *Aspidœcia Normani*, Giard and Bonnier.

1889. *Aspidœcia Normani*, G. and B., Compt.-rend. de l'Acad. des Sci., 29th April.

1897. „ „ H. J. Hansen, The Choniostomatidæ, p. 187, pl. xii. figs. 3a, 3b.

1898. „ „ T. Scott, Sixteenth F. B. Rept., pt. iii. p. 279.

Hab.—Found adhering to a specimen both of *Erythroops elegans* (G. O. Sars) and *Erythroops erythrophthalmus* (Goës), captured in 1901 to the west of May Island; rare.

THE THYROSTRACA OR CIRRIPEDIA.

Scarcely any addition has been made to the number of Forth Crustacea belonging to this group since the publication of Leslie and Herdman's work on the *Invertebrate Fauna of the Estuary*.

THORACICA.

Family LEPADIDÆ.

Genus (1) *Lepas*, Linné, 1767.

1. *Lepas anatifera*, Linné.

1767. *Lepas anatifera*, Linn., Systema Naturæ, ed. xii., vol. i. pt. ii. p. 1109.

1851. „ „ Darwin, Monogr. of the Cirripedia (The Lepadidæ), p. 73, pl. i. fig. 1.

Hab.—Attached to floating timber (Edin. Mus., Leslie and Herdman). Aberlady Bay, on piece of stranded timber, September 1893 (Evans).

Genus (2) *Conchoderma*, Olfers, 1814.2. *Conchoderma aurita* (Linné).

1767. *Lepas aurita*, Linn., Systema Naturæ, ed. xii., vol. i. part ii.
p. 1110.

1851. *Conchoderma aurita*, Darwin, *l. c.*, p. 41, pl. iii. fig. 4.

Hab.—Attached to floating timber (Edin. Mus., Leslie and Herdman).

3. *Conchoderma virgata* (Spengler).

1790. *Lepas virgata*, Spengler, Skrift. Naturh. Selbsk., vol. i., pl. vi.
fig. 9.

1851. *Conchoderma virgata*, Darwin, *l. c.*, p. 146, pl. iii. fig. 2;
pl. ix. fig. 4.

Hab.—Attached to floating timber (Edin. Mus., Leslie and Herdman).

Family BALANIDÆ.

Genus (3) *Balanus*, Da Costa, 1778.4. *Balanus porcatus*, Da Costa.

1778. *Balanus porcatus*, Da Costa, Hist. Nat. Test. Brit., p. 249.

1854. „ „ Darwin, Monograph: Balanidæ, p. 256,
pl. vi. figs. 4a-4e.

Hab.—Not uncommon attached to stones, etc. (Leslie and Herdman).

5. *Balanus crenatus*, Bruguière.

1789. *Balanus crenatus*, Bruguière, Encycl. Method. (des vers).

1854. „ „ Darwin, *l. c.*, p. 261, pl. vi. figs. 6a-6g.

Hab.—Shore at Portobello (Leslie and Herdman). East side of Inchkeith.

6. *Balanus balanoides* (Linné).

1767. *Lepas balanoides*, Linn., Systema Naturæ, vol. i. part ii.
p. 1108.

1854. *Balanus balanoides*, Darwin, *l. c.*, p. 267, pl. vii. figs. 2a-2d.

Hab.—Common on rocks and stones between tide-marks, and also in deep water.

7. *Balanus Hameri* (Ascanius).

1767. *Lepas Hameri*, Ascan., Icones rerum Naturalium, Tab. 10.

1854. *Balanus Hameri*, Darwin, *l. c.*, p. 277, pl. vii. figs. 5a-5c.

Hab.—Largo Bay, and off the west side of Inchkeith, single specimens, and occasionally a cluster attached to pieces of wood.¹

Genus (4) *Verruca*, Schumacher, 1817.

8. *Verruca Strömia* (O. F. Müller).

1776. *Lepas Strömia*, Müll., Zool. Dan. Prodr., No. 3025.

1854. *Verruca Strömia*, Darwin, *l. c.*, p. 518, pl. xxi. figs. 1a-1f.

Hab.—Frequent on stones, dead shells, etc., brought up in the dredge.

RHIZOCEPHALA.

Family SACCULINÆ.

Genus (5) *Sacculina*, Thompson.

9. *Sacculina carcini*, Thompson.

Hab.—Attached to the abdomen of Shore Crabs, *Carcinus mænas*, at Joppa (J. Anderson²); not very rare.

10. *Sacculina triangularis*, J. Anderson.²

Hab.—Firth of Forth (J. Anderson). Dunbar, attached to the abdomen of *Cancer pagurus*; rare (H. C. Williamson).

11. *Sacculina* sp.

Hab.—Attached to the abdomen of *Portunus holsatus*; this Sacculine has been observed very sparingly in different parts of the estuary.

¹ *Balanus tintinnabulum* (Linn.), from Leith Dock, in Leslie and Herdman's *Invertebrate Fauna*, is, as stated by these authors, "a foreign importation," and is not included with the others here.

² Cf. *Proc. Roy. Phys. Soc.*, vol. ii.

Genus (6) *Peltogaster*, Rathke.12. *Peltogaster paguri*, Rathke.

Hab.—On the abdomen of the common Hermit (or Soldier) Crab, *Eupagurus bernhardus* (J. Anderson¹). This Cirriped does not appear to be rare in the Forth estuary.

13. *Peltogaster carcini* (Anderson¹).

Hab.—Firth of Forth, at Joppa (J. Anderson).

 APPENDIX.

The following are some alterations in names, and additions to the species and localities, recorded in Part I. The total number of species enumerated in this Catalogue is now 794.

Genus *Eupagurus*, Brandt, 1851 (p. 112).

The genus name *Pagurus*, Fabricius, 1793, has been restored, and included under it are the four species recorded in Part I. under *Eupagurus*. See *Crustacea of Devon and Cornwall*, by Canon A. M. Norman and T. Scott, p. 8 (1906).

Leptognathia Lilljeborgia, Stebbing (p. 132).

This species has been removed to another genus, viz., *Tanaissus*, Norman and Scott. See *Crustacea of Devon and Cornwall*, p. 34.

Genus *Idothea*, Fabricius (p. 135).

To the species recorded under *Idothea*, add *Idothea granulosa*, Rathke, *Beiträge zur Fauna Norwegens*, p. 23. In March 1906 Mr William Evans obtained near Aberdour, Fife, two

¹ Cf. *Proc. Roy. Phys. Soc.*, vol. ii.

specimens of an *Idothea* that appear to me to belong to this species.

After *Porcellio pictus* (p. 140), add

The genus *Metoponorthus*, Budde-Lund, 1879, and the species *M. pruinosus* (Brandt). This Oniscoidan species has been captured (in April this year) at Slateford, near Edinburgh, by Mr William Evans, and is recorded by him in the *Annals of Scottish Natural History* for July 1906, p. 187.

Gammarus Duebeni, Lillj. (p. 163).

To the localities for this species add: Brackish pools, Luffness, near Aberlady, July 1906 (W. Evans).

Genus *Jassa*, Leach (p. 170).

The two species, *Jassa falcata* (Montagu) and *Jassa pusilla* (G. O. Sars), have been removed to the new genus *Bruzeliella*, Norman. See *Crustacea of Devon and Cornwall*, p. 92.

Corophium grossipes, Linné (p. 172).

This species, under the name of *Cancer grossipes*, was described by Linné in 1767 in *Systema Naturæ*, vol. i. part 2, p. 1055; but as Pallas had already, in 1766, described under the name of *Oniscus volutator*, what appears to be the same form (cf. *Miscellanea Zoologica*, p. 102, pl. xiv. fig. 20), the name of the species should be *Corophium volutator* (Pallas), with *C. grossipes* as a synonym.

Protella phasma, Montagu (p. 174).

Professor Mayer in his work on the "Caprellidæ of the Bay of Naples," *Caprelliden des Golfes von Neapel, Nachtrag*, p. 19, refers this species to the new genus *Pseudoprotella*.

Scapholeberis mucronata, O. F. Müller (p. 179).

To the localities given for this species add:—Side of river Teith, at Callander, September 1906 (W. Evans).

Cyclops Leuckarti, Claus (Part II. p. 347).

While this part of the Catalogue was passing through the press, Mr W. Evans sent me numerous specimens of a *Cyclops* which he had recently captured at the Falls of Leny, by holding a net so that some of the falling water would pass through it. The identification of the specimens somewhat puzzled me at first, but on carefully dissecting one or two of them, I found them to be the young of *Cyclops Leuckarti*; they were in the penultimate or antipenultimate stage; the segmentation of the body being slightly incomplete, while the antennules were furnished with only eleven articulations. Loch Voil, one of the two lochs from which the species is recorded, is situated to the north of Ben Ledi, and the river Balvaig carries its overflow water to Loch Lubnaig; this, in turn, is the source of the river Leny. It is probable, therefore, that *Cyclops Leuckarti* occurs in this loch also, though I have not as yet found it there. The loch and river were in flood when Mr Evans visited the Falls, and the specimens were very likely carried from the loch by the increased force of the current.

The normal habitat of *Cyclops*, as of most fresh-water Entomostraca, is in the comparatively still waters of lochs, ponds, and other similar places, and their presence in the rapid currents of rivers is usually accidental. The occurrence of this *Cyclops* in the Leny is, therefore, of interest from its bearing on the question of distribution, and as showing that the species is probably present in Loch Lubnaig as well as in the other two lochs from which it has been recorded.

ADDITIONAL NOTE.—As the second part of this Catalogue was ready to go to press, I received a copy of an important work on the Amphipoda Gammaridea by the Rev. T. R. R. Stebbing, M.A., F.R.S., etc. This work, which has just been published, forms the twenty-first of the series of Monographs that are being issued under the general title *Das Tierreich*. The Amphipoda are here divided into three legions, viz., the Gammaridea, the Hyperiidea, and the Caprellidea, and it is the first that forms the subject of Mr Stebbing's volume. The number of known species belonging to the Gammaridea as given in it is 1333. Of these, 1076 are "accepted species," while the remaining 257 are regarded as doubtful. Their classification differs to some extent from that of the Monograph by G. O. Sars. There are also a considerable number of changes in the nomenclature, and I take the opportunity to indicate the more important of these, so far as they concern the species recorded in this Catalogue.

In the work referred to—*Talitrus locusta* (Pall.) becomes *T. saltator* (Mont.), and *Orchestia littorea* (Mont.) *O. gammarellus* (Pall.), while *Hyale Nilssoni* (Rathke) takes the name of *H. Prevostii* (M.-E.). The genus *Callisoma*, A. Costa, becomes *Scopelocheirus*, Bate, and *Orchomene humilis* (A. Costa) becomes *O. Batei*, O. Sars. *Tryphosella*, Bonnier, is replaced by *Tryphosa*, Boeck, but *Tryphosa nana* (Kröy.) becomes *Orchomenella nanus* (Kröy.). *Anonyx nugax* (Phipps) becomes *A. lagena*, Kröy., and the genus *Tmetonyx*, Stebb., replaces *Hoplonyx*, O. Sars. *Phoxocephalus Fultoni*, T. Scott, becomes *Paraphoxus Fultoni* (T. S.), *Harpinia neglecta*, O. Sars, becomes *H. antennaria*, Meinart, and *Amphilochoides serratipes* (Norm.) resumes the name of *A. Boeckii*, O. Sars. *Metopa Bruzelii* (Göes) becomes *M. sinuata*, O. Sars, and *M. robusta* is restored to its place in the same genus. *Synchelidium brevicarpum* (B. & W.) becomes *S. haplocheles* (Grube), and *Halimедon parvimanus* (B. & W.) *Westwoodilla cæcula*, Bate. The genus *Paramphithoë*, Bruz., becomes *Neopleustes*, Stebb., and *Paratylus*, O. Sars, *Nototropis*, A. Costa; *N. uncinatus*, O. Sars, is also made a synonym to *N. falcatus* (Metzg.). *Iphimedia minuta*, O. Sars, becomes *Panoploea minuta* (O. Sars). The genus *Amathilla*, Bate, is replaced by *Gammarella*, Herbst, and *Gammaropsis*, Lillj., by *Eurystheues*, Bate, while *Gammaropsis nana*, O. Sars, becomes *E. palmatus* (Stebb. and Rob.). *Podoceropsis excavata* (Bate) becomes *P. nitida*, Stimps., *Jassa Herdmani* (A. O. Walker) *J. dentex* (Czern.), and *Jassa pelagica*, Leach, *Parajassa pelagica* (Leach). *Erichthonius abditus* (Templ.) becomes *E. brasiliensis* (Dana), and *Siphonoëcetes Whitei* (Gosse) *E. Colletti*, Boeck.

The following two species are additions to my list, viz., *Peltocoxa Marioni*, Catta, and *Cressa minuta*, Boeck, both being recorded from the Forth. This raises the total to 796.

I need scarcely add that this important work, by one of the foremost of living Carcinologists, is indispensable to those engaged in the study of the Amphipoda.

Before closing I desire to take this opportunity to express my indebtedness to Mr William Evans, F.R.S.E., President of the Royal Physical Society, for the whole-hearted assistance he has given me while this Catalogue was passing through the press. I am also not forgetful of other friends who have shown an interest in my efforts to make this contribution to Scottish Natural History, unpretentious though it be, of some use to students of the Crustacea of Scotland.

ERRATA IN FIRST PART.

- Page 99. Near bottom; for *Homarus vulgarus* read *Homarus vulgaris*, and for *Crangon vulgarus* read *Crangon vulgaris*.
- „ 103. Near top; for *Paramphithoë monocupis* read *Paramphithoë monocuspis*.
- „ 107. Near top; for “Stalked-eyed Crustacea” read “Stalk-eyed Crustacea.”
- „ 110. Near bottom; in the reference “1767, *Cancer rostratus*, Linn., Syst. Nat., ed. xii., vol. ii. p. 1014,” for vol. ii. p. 1014 read “vol. i. part ii. p. 1045”; and in the subsequent references to the twelfth edition of *Systema Naturæ*, Linné, for “vol. ii.” read “vol. i. part ii.”
- „ 113. Near top; for *Turretella* read *Turritella*.
- „ 120. Near top; for *Nyctiphanus* read *Nyctiphanes*.
- „ 122. Near bottom; in the reference “Reise i Chr. oy,” etc., “oy” should be “og.”
- „ 125. In footnote; for “Dr Welley” read “Dr Willey.”
- „ 137. In footnote; for *Leptospidia* read *Leptaspidia*.
- „ 144. Near bottom; *Calisoma* should be *Callisoma* in each of the four lines where it occurs.
- „ 165. Near top; the name *Platessoides limandoides* is a synonym of *Drepanopsetta platessoides*. (See also p. 150.)
- „ 165. Middle; for *Cheirocrates* read *Cheirocratus* in four places.
- „ 189. Near the middle; for “Sonntagshlatt.” read “Sonntagsblatt.”

Part of the cost of publication of the foregoing Catalogue has been defrayed by a Grant made to the Author by the Carnegie Trust for the Universities of Scotland.

INDEX

TO

GENERA AND PRINCIPAL LARGER DIVISIONS.

(Pages 108 to 190 are in Part I.)

	PAGE		PAGE
Acantholeberis,	182	Beatricella,	326
Acartia,	304	Bodotria,	126
Acidostoma,	144	Bomolochus,	361
Acontiophorus,	358	Bosmina,	180
Acroperus,	182	Brachiella,	373
Aglaia,	274	Brachyura,	108
Alona,	183	Bradya,	310
Alonella,	185	Bradyidius,	298
Alonopsis,	182	Bradypontius,	359
Alteutha,	312	Branchiura,	190
Amathilla,	162	Bruzeliella,	379
Ameira,	323	Byblis,	150
Amenophia,	316	Bythocythere,	291
Ampelisca,	149	Bythotrephes,	188
Amphiascus,	321		
Amphilochoides,	151	Calanus,	297
Amphilocheus,	150	Caligus,	366
Amphipoda,	142	Calliopius,	160
Amphithoë,	169	Callisoma,	144
Anapagurus,	113	Calocaris,	115
Anchorella,	373	Camptocercus,	182
Anomalocera,	303	Campylaspis,	130
Anonyx,	146	Cancer,	108
Aora,	166	Candacia,	303
Apherusa,	159	Candona,	275
Aplostoma,	363	Candonopsis,	277
Arcturella,	136	Canthocamptus,	328
Argillœcia,	279	Canuella,	306
Argissa,	158	Caprella,	174
Argulus,	190	Carcinus,	108
Armadillidium,	141	Cecrops,	369
Ascidicola,	363	Centropages,	300
Asellus,	136	Cerapus,	171
Aspidiscus,	313	Ceriodaphnia,	179
Aspidœcia,	375	Charopinus,	373
Astacilla,	135	Cheirocratus,	165
Astacus,	116	Chondracanthus,	371
Asterocheres,	356	Chydorus,	187
Asterope,	295	Cirripedia,	375
Atelecyclus,	110	Cladocera,	175
Athelges,	141	Cletodes,	337
Attheyella,	330	Collocheres,	357
		Conchoderma,	376
Bairdia,	279	Copepoda,	296
Balanus,	376	Corophium,	172, 379
Bathyporeia,	147	Corycæus,	360

	PAGE		PAGE
Corystes,	109	Gastrosaccus,	122
Crangon,	116	Gitana,	152
Cressa,	155, 382	Gnathia,	133
Cribrapontius,	359	Graptoleberis,	185
Cryptothir,	142	Guernea,	162
Cumella,	130		
Cumopsis,	126	Hæmobaphes,	370
Cyclocypris,	268	Halicyclops,	351
Cyclopina,	345	Halimedon,	156
Cyclops,	346, 380	Halithalestris,	314
Cylindropsyllus,	341	Haploops,	150
Cylisticus,	140	Harpacticus,	310
Cypria,	268	Harpinia,	148
Cypridopsis,	272	Harrietella,	336
Cyprinotus,	271	Haustorius,	147
Cypris,	269	Hemimysis,	124
Cyprois,	274	Herpetocypris,	271
Cythere,	279	Herrmannella,	354
Cytheridea,	284	Hersiliodes,	355
Cytherideis,	292	Heteromysis,	122
Cytherois,	292	Heteropsyllus,	325
Cytheropteron,	290	Hippolyte,	118
Cytherura,	287	Hippomedon,	145
		Holopedium,	177
Dactylopodella,	319	Hoplonyx,	146
Dactylopusia,	317	Hyale,	144
Danielssenia,	333	Hyas,	111
Daphnia,	177	Hyperia,	142
Darwinula,	279	Hyperoche,	142
Delavalia,	326		
Dermatomyzon,	357	Idomene,	319
Dexamine,	161	Idomenella,	319
Diaphanosoma,	176	Idothea,	134, 378
Diaptomus,	300	Ilyocryptus,	181
Diastylis,	128	Ilyocypris,	278
Doropygus,	363	Ilyodromus,	272
Drepanothrix,	181	Inachus,	111
Dulichia,	173	Iphimedia,	158
Dyspontius,	360	Iphinoë,	126
		Ischyrocerus,	169
Ebalia,	112	Isopoda,	130, 379
Ectinosoma,	306	Itunella,	340
Enhydrosoma,	340		
Epimeria,	158	Jæra,	136
Erichthonius,	171	Janira,	136
Erythrops,	122	Jassa,	170, 379
Eucythere,	285	Jeanella,	365
Eudorella,	127		
Eudorelloopsis,	128	Krithe,	285
Eupagarus,	112, 378		
Eupelte,	312	Labidocera,	303
Eurycercus,	182	Læmargus (see Orthagoriscola).	
Eurycope,	138	Lamippe,	364
Eurycypris,	270	Lamprops,	127
Eurydice,	133	Laophonte,	333
Eurynome,	111	Laophontodes,	336
Euryte,	351	Laphystius,	159
Eurytemora,	301	Lathonura,	181
Euthemisto,	143	Latona,	176
Evadne,	189	Leander,	119
Evansula,	342	Ledigia,	185
		Lembos,	166
Fultonina,	340	Lepas,	375
		Lepeophtheirus,	368
Galathea,	114	Lepidepecreum,	147
Gammaropsis,	167	Leptastacus,	342
Gammarus,	162, 379	Leptocheirus,	167

	PAGE		PAGE
Leptodora,	189	Orchestia,	143
Leptognathia,	131, 378	Orchomene,	145
Leptomysis,	123	Orthagoriscola,	369
Leptopontia,	341	Ostracoda,	267
Leptopsyllus,	344		
Lernæa,	370	Pagurus,	378
Lernæenicus,	370	Pandalina,	119
Leucon,	127	Pandalus,	119
Leucothoë,	155	Paracalanus,	298
Lichomolgus	352	Paracandona,	277
Ligia,	138	Paradoxostoma,	293
Lilljeborgia,	165	Paramesochra,	344
Limnocythere,	284	Paramphithoë,	157
Limnoria,	133	Paramunna,	137
Lithodes,	112	Parapontella,	303
Longipedia,	305	Parartotrogus,	360
Loxoconcha,	285	Paratanais,	131
		Parathalestris,	314
Machærina,	294	Parathemisto,	143
Macromysis,	124	Paratylus,	160
Macropodia,	110	Pariambus,	174
Macropsis,	124	Peltocoxa,	382
Macrothrix,	181	Peltogaster,	378
Macrura,	112	Peratacantha,	186
Mæra,	164	Periocolodes,	156
Maraenobiotus,	332	Petalosarsia,	130
Megaluropus,	164	Philomedes,	295
Megamphopus,	168	Philoscia,	140
Melita,	163	Photis,	168
Melphidippella,	162	Phoxocephalus,	148
Mesochra,	332	Phryxus,	141
Metopa,	153	Phtisica,	174
Metopella,	154	Phyllopodopsyllus,	343
Metoponorthus,	379	Phyllothalestris,	314
Metridia,	302	Pinnotheres,	110
Microdeutopus,	166	Pionocypris,	272
Microprotopus,	168	Platyarthrus,	140
Microsetella,	309	Platychelipus,	341
Microthalestris,	315	Pleonexes,	169
Misophria,	305	Pleurogonium,	137
Modiolicola,	354	Pleuroxus,	186
Monocolodes,	155	Podoceropsis,	168
Monospilus,	188	Podon,	188
Monstrilla,	365	Polycope,	295
Moraria,	331	Polycopsis,	295
Munida,	115	Polyphemus,	188
Munna,	137	Pontocrates,	156
Mysidopsis,	123	Pontocypris,	278
		Pontopolites,	328
Næsa,	134	Porcellana,	114
Nannopus,	340	Porcellidium,	312
Nematosceli,	121	Porcellio,	140
Neobradya,	306	Portumnus,	109
Neomysis,	125	Portunus,	108
Nephrops,	116	Potamocypris,	273
Nicothoë,	361	Protella,	174, 379
Nika,	117	Protomedeia,	166
Nitocra,	331	Pseudanthessius,	353
Normanella,	337	Pseudobradya,	309
Notodelphys,	362	Pseudocalanus,	298
Notodromas,	274	Pseudocaligus,	367
Nyctiphanes,	120	Pseudocuma,	129
		Pseudocyclopia,	299
Oithona,	345	Pseudocyclops,	302
Oniscus,	139	Pseudocythere,	292
Oralien,	371	Pseudoprotella,	379
		Pseudothalestris,	316

	PAGE		PAGE
Pterinopsyllus,	346	Tanais,	131
Pteropsyllus,	343	Tanaissus,	378
Rhoda,	120	Tanaopsis,	132
Rhynchomyzon,	357	Tegastes,	312
Rhynchotalona,	185	Temora,	301
Rhynchothalestris,	315	Tetragoniceps,	343
Robertsonia,	325	Thalestris,	313
Sabelliphilus,	355	Thaumaleus,	366
Sacculina,	377	Thersites,	362
Salenskya,	374	Thersitina,	362
Scapholeberis,	179, 380	Thompsonula,	333
Schistomysis,	125	Thysanoessa,	120
Schizopoda,	119	Tigriopus,	311
Sclerochilus,	292	Tisbe,	313
Scottocheres,	358	Trebius,	369
Scottomyzon,	357	Trichoniscus,	139
Sida,	176	Tritæta,	161
Simosa,	179	Tryphosa,	145
Siphonoëcetes,	172	Tryphosella,	145
Siriella,	121	Tryphosites,	146
Socarnes,	144	Typhlotanais,	131
Sphæroma,	134	Unciola,	173
Spirontocaris,	118	Urothoë,	148
Stenhelia,	322	Vaunthompsonia,	127
Stenothoe,	152	Verruca,	377
Stephos,	299	Westwoodia,	316
Sthenometopa,	154	Xestoleberis,	286
Sympleustes,	157	Zaus,	311
Sympoda,	125	Zosime,	306
Synchelidium,	156		
Tachidius,	327		
Talitrus,	143		

For some recent changes in nomenclature, see p. 381.