

## Review SUPPLEMENTARY MATERIAL

## Alien species in British brackish and marine waters

Dan Minchin<sup>1,2</sup>, Elizabeth J. Cook<sup>3\*</sup> and Paul F. Clark<sup>4</sup>

**1** Marine Organism Investigations, Caragh, Ballina, Killaloe, Co Clare, Ireland

**2** Coastal Research and Planning Institute, Klaipeda University, Klaipeda, Lithuania

**3** Scottish Association for Marine Science, Scottish Marine Institute, Oban, Argyll PA37 1QA, UK

**4** Natural History Museum, Cromwell Road, London, SW7 5BD, UK

E-mail: [moiireland@yahoo.ie](mailto:moiireland@yahoo.ie) (DM), [ejc@sams.ac.uk](mailto:ejc@sams.ac.uk) (EJC), [p.clark@nhm.ac.uk](mailto:p.clark@nhm.ac.uk) (PFC)

\*Corresponding author

Received: 20 April 2012 / Accepted: 3 December 2012 / Published online: 10 January 2013

Handling editor: Vadim Panov

Online article is available at [http://www.aquaticinvasions.net/2013/AI\\_2013\\_1\\_Minchin\\_et al.pdf](http://www.aquaticinvasions.net/2013/AI_2013_1_Minchin_et al.pdf)

### Supplementary material

The following supplementary material is available for this article:

**Appendix 1.** List of alien species recorded from British brackish and marine waters.

**Appendix 1.** List of alien species recorded from British brackish and marine waters. Their order does not relate to date of discovery. Region: C = Celtic Seas; E = English Channel; N = North Sea; region of first introduction if known underlined. Abbreviations for Vector/ Pathway: AQ = Aquaculture; V = Vessels; D = Natural Drift; PL = Planting; ET = Eel Transport; LF = Live Food; AF = Anthropogenic Flotsam; TI = Timber importation; BB = Bait Buckets; AQU = Aquarium.

Species	Region	Date of first record placed within the salinity range of the recipient zone 1-30psu >30psu	Native Range	Likely Vector/ Pathway	Current status	Impact (known or expected in Britain)	References
<b>Magnoliophyta</b>							
Haplosporididae							
<i>Bonamia ostreae</i> Pichot, Comps, Tigé, Grizel & Rabouin, 1980	E C N	1982	1982	NE Pacific	AQ, V	widespread, established	native oyster mortalities Howard 1994; Hudson and Hill 1991; Laing 2006; Brown et al. 2006
<b>Ocrophyta, Phaeophyceae</b>							
Alariaceae							
<i>Undaria pinnatifida</i> (Harvey) Suringar, 1872	E N	1994	1994	NW Pacific	V	regionally established	fouling, possible cultivation, competition Fletcher and Manfredi 1995; Fletcher and Farrell 1999
Scytoniphonaceae							
<i>Colpomenia peregrina</i> (Sauvageau) Hamel, 1927	E C N		1907	Pacific	D, AQ	widespread, established	possible competition Mineur et al. 2008; Arenas et al. 2006; Cotton 1908; Norton 1976; Wilkinson 1975
Sargassaceae							
<i>Sargassum muticum</i> (Yendo) Fensholt, 1955	E C		1971	N Pacific	AQ, D	regionally established	habitat modification, affecting navigation, competition Farnham et al. 1981; Fletcher and Farrell 1999; Withers et al. 1975; Harries et al. 2007; Critchley et al. 1983; Critchley et al. 1986; Farnham et al. 1973b
<b>Rhodophyta, Florideophyceae</b>							
Bonnemaisoniaceae							
<i>Bonnemaisonia hamifera</i> Hariot, 1891	E C N		1890	N Pacific	unknown	widespread, established	competition Holmes 1897; Maggs and Stegenga 1999; Irvine 1982; Mineur et al. 2008; Dixon and Irvine 1977

**Appendix 1 (continued).**

Species	Region	Date of first record placed within the salinity range of the recipient zone 1-30psu >30psu	Native Range	Likely Vector/ Pathway	Current status	Impact (known or expected in Britain)	References
<b>Ceramiaceae</b>							
<i>Antithamnionella spirographidis</i> (Schiffner) Wollaston, 1968	E C	1931	N Pacific	V	regionally established	unknown	Westbrook 1934; Maggs and Stegenga 1999; Brodie et al. 2007
<i>Antithamnionella ternifolia</i> (C. Agardh) Nageli, 1847	E C	1906	Indo-Pacific	unknown	regionally established	unknown	Maggs and Hommersand 1993; Maggs and Stegenga 1999; Brodie et al. 2007
<i>Asparagopsis armata</i> Harvey, 1855	C E N	1949	S Pacific	V	widespread, established	competition	Drew 1950; Irvine et al. 1975; Maggs and Stegenga 1999; Ni Ó Cuáin et al. 2004
<b>Caulacanthidae</b>							
<i>Caulacanthus ustulatus</i> (Mertens ex Turner) Kützing, 1843	E	2004	NW Pacific	unknown	one locality, established	unknown	Mineur et al. 2006; ICES 2006
<b>Dasyaceae</b>							
<i>Heterosiphonia japonica</i> Yendo, 1920	N E C	2004	NW Pacific	unknown	widespread, established	competition	Bjaerke and Rueness 2004; ICES 2005b, 2006; C. Moore (pers.comm.)
<b>Dumontiaceae</b>							
<i>Pikea californica</i> Harvey, 1853	E	1967	N Pacific	V	localities, established	unknown	Maggs and Guiry 1987; Maggs and Ward 1996; Brodie et al. 2007
<b>Rhodomeliaceae</b>							
<i>Polysiphonia harveyi</i> J.W. Bailey, 1848	E C N	1908	N Pacific	unknown	widespread, established	unknown	Maggs and Hommersand 1993; Maggs and Stegenga 1999; McIvor et al. 2001; Brodie et al. 2007; Mineur et al. 2008
<b>Halymeniaceae</b>							
<i>Cryptonema hibernica</i> Guiry & L. Irvine, 1974	E	2000	S Atlantic	V	one locality, no recent record	unknown	Hardy and Guiry 2003
<i>Grateloupia subpectinata</i> Holmes, 1912	E	1947	Indo-Pacific	AQ	one locality, established	unknown	Farnham and Irvine 1968; Irvine 1983; Arenas et al. 2006; Guiry 2011
<i>Grateloupia turuturu</i> Yamada, 1941	E C	1969	NW Pacific	AQ	regionally established	unknown	Farnham and Irvine 1973b; Farnham 1980; Maggs and Stegenga 1999; Gavio and Fredericq 2002

**Appendix 1 (continued).**

Species	Region	Date of first record placed within the salinity range of the recipient zone		Native Range	Likely Vector/Pathway	Current status	Impact (known or expected in Britain)	References
		1-30psu	>30psu					
<b>Solieriaceae</b>								
<i>Sarcodiotheca gaudichaudi</i> , (Montagne) P.W. Gabrielson, 1982	E	1974	E Pacific	V, AQ	localities, established	unknown		Farnham and Irvine 1979; Hardy and Guiry 2003; Brodie et al. 2007
<i>Solieria chordalis</i> (C. Agardh) J. Agardh, 1842	E C	1976	Unclear	V, D	localities, established	unknown		Parke and Dixon 1976; Farnham and Irvine 1979; Farnham 1980; Hiscock and Maggs 1984; Brodie et al. 2007; C. Maggs (pers. comm.)
<b>Chlorophyta, Bryopsidophyta</b>								
<b>Codiaceae</b>								
<i>Codium fragile</i> ssp <i>fragile</i> (Proven, Booth, Todd, Beatty, & Maggs, 2008)	C E N	1891	N Pacific	unknown	widespread, established	competition		Silva 1955; Fletcher et al. 1989; Trowbridge 1998; Chapman 1999; Provan et al. 2008
<b>Anthophyta, Poaceae</b>								
<i>Spartina townsendii</i> var. <i>anglica</i> C.E. Hubbard	E C N	1850s	NW Atlantic	PL	widespread, established	habitat engineer		Hubbard and Stebbings 1967; Marchant 1967; Gray et al. 1991; Frid et al. 1999
<b>Cnidaria, Anthozoa</b>								
<b>Diadumenidae</b>								
<i>Diadumene lineata</i> (Verrill, 1869)	C E N	1899	1899	NW Pacific	AQ, V	widespread, established	unknown	Manuel 1981; Fautin et al. 2007
<b>Olindiasidae</b>								
<i>Gonianemus vertens</i> A. Agassiz, 1862	N C E	1913	1913	NW Pacific	V, AQ	widespread, established	unknown	Edwards 1976
<b>Nematoda, Dracunculoidea</b>								
<b>Anguillicolidae</b>								
<i>Anguillicoloides crassus</i> Kuwahara, Niimi & Itagaki, 1974	N C E	1987	1987	NW Pacific	ET, V	widespread, established	endoparasite of <i>Anguilla anguilla</i>	Kennedy and Fitch 1990; Kennedy 1993

**Appendix 1 (continued).**

Species	Region	Date of first record placed within the salinity range of the recipient zone		Native Range	Likely Vector/Pathway	Current status	Impact (known or expected in Britain)	References
		1-30psu	>30psu					
<b>Trematoda, Cercarida</b>								
<i>Cercaria sensifera</i> Stunkard & Shaw, 1981	N	1939	1939	NW Atlantic	AQ	one locality, no recent record	endoparasite of <i>Urosalpinx cinerea</i>	Cole 1942
<b>Annelida, Polychaeta</b>								
Serpulidae								
<i>Ficopomatus enigmaticus</i> (Fauvel, 1923)	E C N	1922	1922	Indo-Pacific	V	widespread, established	fouling	Monro 1924; Harris 1970; Hiscock and Hoare 1975; Dixon 1981; Davies et al. 1989; Zibrowius and Thorp 1989
<i>Hydroides dianthus</i> (Verrill, 1873)	E		1970	NW Atlantic	V, AQ	localities, no recent record	unknown	Zibrowius and Thorp 1989
<i>Hydroides elegans</i> (Haswell, 1883)	C		1937	Indo-Pacific	V	one locality, no recent record	unknown	Naylor 1965; ten Hove 1974; Hayward and Ryland 1995
<i>Hydroides ezoensis</i> , Okuda, 1934	E		1976	NW Pacific	V	localities, established	fouling	Thorp 1980; Thorp et al. 1987; Zibrowius and Thorp 1989
Spirorbidae								
<i>Janua brasiliensis</i> (Grube, 1872)	E	1974	1974	SW Atlantic	V, D	one locality, no recent record	unknown	Knight-Jones et al. 1975; Knight Jones and Knight-Jones 1977, 1980; Critchley and Thorp 1985; Zibrowius and Thorp 1989
<i>Pileolaria berkeleyana</i> (Rioja, 1942)	E		1974	NE Pacific	V, D	localities, established	unknown	Knight-Jones et al. 1975; Knight Jones and Knight-Jones 1977, 1980; Zibrowius and Thorp 1989; Thorp 1991; Thorp et al. 1986
Goniadidae								
<i>Goniadella gracilis</i> Verrill, 1873	C		1970	NW Atlantic	V	localities, established	unknown	Walker 1972; Mackie et al. 1995

**Appendix 1 (continued).**

Species	Region	Date of first record placed within the salinity range of the recipient zone 1-30psu >30psu	Native Range	Likely Vector/ Pathway	Current status	Impact (known or expected in Britain)	References
<b>Sabellidae</b>							
<i>Desdemona ornata</i> Banse, 1957	E	1997	Indo-Pacific	unknown	localities, established	unknown	Smith et al. 1999
<b>Maldanidae</b>							
<i>Clymenella torquata</i> , (Leidy, 1855)	N	1936	1936	NW Atlantic	AQ	localities, established	unknown
<b>Spionidae</b>							
<i>Marenzelleria viridis</i> (Verrill, 1973)	N	1979	1979	NW Atlantic	V, D	localities, established	unknown
<b>Mollusca, Gastropoda</b>							
<b>Hydrobiidae</b>							
<i>Potamopyrgus antipodarum</i> (J.E. Gray, 1843)	N C E	1859	S Pacific (New Zealand)	V, Birds	widespread, established	competition	McMillan 1968; Ponder 1988
<b>Calypttraeidae</b>							
<i>Crepidula fornicata</i> (Linnaeus, 1758)	C E N	1872	1872	NW Atlantic	AQ	widespread, established	fouling, trophic competition
<i>Calyptraea chinensis</i> Linnaeus, 1758	C	1957	NE Atlantic	AQ	one locality, no recent record	unknown	McMillan 1968; Minchin et al. 1987; Hayward and Ryland 1995
<b>Muricidae</b>							
<i>Rapana venosa</i> (Valenciennes, 1846)	N	1991	N Pacific	V, LF	single records, not established	likely bivalve predation, selective fishery	Kerckhof et al. 2006; ICES 2004a, 2007
<i>Urosalpinx cinerea</i> (Say, 1822)	N E	1920s	NW Atlantic	AQ	regionally established	oyster predator	Orton and Winkworth 1928; Orton 1930; Cole 1942; Hancock 1954, 1959; Gibbs et al. 1991

**Appendix 1 (continued).**

Species	Region	Date of first record placed within the salinity range of the recipient zone 1-30psu    >30psu	Native Range	Likely Vector/Pathway	Current status	Impact (known or expected in Britain)	References
<b>Bivalvia</b>							
Mytilidae							
<i>Brachidontes exustus</i> Linnaeus, 1758	C	1977	W Atlantic	AF	single record, not established	none	J Light (pers. comm.)
<i>Aulacomya ater</i> (Molina, 1782)	N	1994	S Atlantic, SE Pacific	V	one locality, no recent record	none	McKay 1994; B Leyshon (pers. comm.)
<i>Choromytilus chorus</i> (Molina, 1782)	C E N	1967	SW Atlantic, SE Pacific	AQ	not established	unknown	ICES 1972; Utting and Spencer 1992
Ostreidae							
<i>Crassostrea gigas</i> (Thunberg, 1793)	E C N	1926	1926	N Pacific	AQ	widespread for cultivation, regionally established in wild (C, E)	fouling, trophic competition, aquaculture
Walne and Helm 1979; Utting and Spencer 1992; Spencer et al. 1994; Drinkwaard 1998; Eno 1998; O'Foighil et al. 1998; Couzens 2006; ICES 2009							
<i>Crassostrea rhizophorae</i> Guilding, 1828	C	1980	W Atlantic	AQ	not established	experimental culture	Utting and Spencer 1992
<i>Crassostrea virginica</i> (Gmelin, 1791)	N C E	1872	1872	NW Atlantic	AQ	not established	once cultivated
Utting and Spencer 1992							
<i>Lopha folium</i> , (Linnaeus, 1758)	E	1986	NW Atlantic	AF	not established	none	Turk 1988
<i>Tiostrea chilensis</i> (Philippi, 1845)	C	1963	S Pacific	AQ	one locality, established	experimental culture	Walne 1979; Buroker et al. 1983; Utting and Spencer 1992; Richardson et al. 1993; Jeffs et al. 1997; Eno 1998
Atrinidae							
<i>Atrina rigida</i> (Lightfoot, 1786)	E	1986	NW Atlantic	AF	not established	none	Turk 1988

**Appendix 1 (continued).**

Species	Region	Date of first record placed within the salinity range of the recipient zone		Native Range	Likely Vector/Pathway	Current status	Impact (known or expected in Britain)	References
		1-30psu	>30psu					
<b>Dreissenidae</b>								
<i>Dreissena polymorpha</i> (Pallas, 1771)	N E	1824		Ponto-Caspian	TI	regionally established	fouling, trophic competition	Aldridge et al. 2004; Kerney 1976; Kerney and Morton 1970
<i>Mytilopsis leucophaeta</i> Conrad, 1831	C	1998		W Atlantic	V	one locality, no recent record	unknown	Oliver et al. 1998; Eno 1998; Bamber and Taylor 2002
<b>Corbiculidae</b>								
<i>Corbicula fluminea</i> (O.F. Müller, 1774)	N	1998		NW Pacific	BB, V, AQU	regionally established	trophic competition	Elliott and zu Ermgassen 2008; Baker et al. 1999; Morton and Tong 1985; Aldridge and Müller 2001
<b>Petricolidae</b>								
<i>Petricola pholadiformis</i> Lamark, 1818	N E	1890	1890	NW Atlantic	AQ	regionally established	borer	Purchon 1955; Tebble 1966; Eno 1998; ICES 1972; Duval 1963
<b>Pharidae</b>								
<i>Ensis directus</i> (Conrad, 1843)	N E C		1989	NW Atlantic	D	regionally established	competition, potential fishery	Howlett 1990; Eno 1998; Palmer 2004; Dansey 2011
<b>Pinctadidae</b>								
<i>Pinctada radiata</i> , (Leach, 1814)	E		1986	NW Atlantic	AF	not established	unknown	Turk 1988
<b>Pteriidae</b>								
<i>Pteria colymbus</i> (Röding, 1798)	E		1986	NW Atlantic	AF	not established	unknown	Turk 1988
<b>Veneridae</b>								
<i>Mercenaria mercenaria</i> (Linnaeus, 1758)	N E C		1860	NW Atlantic	V	regionally, established	localised fishery	Heppel 1961; Ansell 1963; Tebble 1966; Mitchell 1974; Cox 1991
<i>Venerupis philippinarum</i> (Adams & Reeve, 1850)	C E N	1980	1980	NW Pacific	AQ	widespread established, recruits in wild (E)	in cultivation	Utting and Spencer 1992; Kaiser et al. 1996; Ashelby et al. 2004; Jensen et al. 2004; Caldow et al. 2007; ICES 2007

**Appendix 1 (continued).**

Species	Region	Date of first record placed within the salinity range of the recipient zone 1-30psu    >30psu	Native Range	Likely Vector/Pathway	Current status	Impact (known or expected in Britain)	References
<b>Arthropoda, Maxillopoda, Copepoda, Calanoida</b>							
<b>Acartiidae</b>							
<i>Acartia tonsa</i> Dana, 1849	E N	1956	N Pacific	V	regionally, established	unknown	Conover 1957; Brylinski 1981; Taylor 1987
<b>Temoridae</b>							
<i>Eurytemora americana</i> L.W. Williams, 1906	E	1931	NW Atlantic	V	localities, no recent record	unknown	Lowndes 1931, 1932; Gurney 1933
<b>Ostracoda</b>							
<b>Sarsiellidae</b>							
<i>Eusarsiella zostericola</i> Cushman, 1906	N E	>1870	>1870	NW Atlantic	AQ	regionally, established	Bamber 1987a, 1987b; ICES 2006
<b>Thecostraca, Cirripedia, Thoracica</b>							
<b>Archaeobalanidae</b>							
<i>Austrominius modestus</i> (Darwin, 1854)	E C N	1943	1943	S Pacific	V	widespread, established	fouling, competition
							Bishop 1947; Stubbings 1950; Crisp 1958; Barnes and Barnes 1960; Hiscock et al. 1978; Muxagata et al. 2004; O'Riordan 2009
<b>Balanidae</b>							
<i>Balanus amphitrite</i> Darwin, 1854	E C	1937	circum-tropical	V	not established	fouling	Bishop 1950; Pitombo 2004
<i>Balanus trigonus</i> Darwin, 1854	E	1986	W Atlantic	AF	not established	unknown	Turk 1988

## Appendix 1 (continued).

Species	Region	Date of first record placed within the salinity range of the recipient zone 1-30psu    >30psu	Native Range	Likely Vector/ Pathway	Current status	Impact (known or expected in Britain)	References
<b>Malacostraca, Amphipoda</b>							
Caprellidae							
<i>Caprella mutica</i> Schurin, 1935	C E N	2000	NW Pacific	V	widespread, established	fouling	Willis et al. 2004; Cook et al. 2007; Ashton et al. 2008 ; Ashelby 2005
Aoridae							
<i>Grandidierella japonica</i> Stephensen, 1938	E	1997	NW Pacific	V	one locality, no recent record	unknown	Smith et al. 1999
<i>Gammarus tigrinus</i> Sexton, 1939	C	1939	NW Atlantic	V	localities, established	competitor	Hynes 1954; Lincoln 1979
<b>Decapoda</b>							
Palaemonidae							
<i>Palaemon macrodactylus</i> Rathbun, 1902	N	2001	2001	NW Pacific	V	localities, established	Ashelby et al. 2004; Worsfold and Ashelby 2008
Penaeidae							
<i>Marsupenaeus japonicus</i> (Bate, 1888)	E C	1989	Indo-Pacific	AQ	individual records, not established	unknown	Clark 1990; Lockley 2004; D Quigley (pers. comm.); A Crosnier (pers. comm.)
Nephropidae							
<i>Homarus americanus</i> (H. Milne Edwards, 1837)	N	1988	NW Atlantic	LF, escapes and deliberate introduction	individual records, not established	possible hybridisation	Stebbing et al. 2012
Portunidae							
<i>Callinectes sapidus</i> Rathbun, 1896	E N	1975	NW Atlantic	V	four records, not established	unknown	Ingle 1980; Clark 1984
Varunidae							
<i>Eriochier sinensis</i> (H. Milne Edwards, 1853)	N C E	1935	1935	NW Pacific	V	many localities, established	bank erosion
<i>Brachynotus sexdentatus</i> (Risso, 1827)	C	1957	Mediterranean	V	one locality, no recent record	unknown	Naylor 1957, 1965

## Alien species in British brackish and marine waters

## Appendix 1 (continued).

Species	Region	Date of first record placed within the salinity range of the recipient zone 1-30psu    >30psu	Native Range	Likely Vector/Pathway	Current status	Impact (known or expected in Britain)	References
<b>Pilumnoididae</b>							
<i>Pilumnoides perlatus</i> (Poepigg, 1836)	E	1898	E Pacific	V	single record, not established	unknown	Valentin 1900; Ingle 1980; Clark 1986;
<b>Panopeinae</b>							
<i>Dyspanopeus sayi</i> (S.I. Smith, 1869)	C	1950s	W Atlantic	V	single record, not established	none	Naylor 1965
<i>Rhithropanopeus harrisii</i> (Gould, 1841)	C	1996	NW Atlantic	V	one locality, established	unknown	Naylor 1965; Ingle 1980; D. Minchin (pers. obs.)
<b>Pycnogonida</b>							
<b>Ammotheidae</b>							
<i>Ammothea hilgendorfi</i> (Böhm, 1879)	E	<1912	S Pacific	V	localities, established	unknown	Bamber 1985; S Trewella (pers. comm.)
<b>Bryozoa</b>							
<b>Bugulidae</b>							
<i>Bugula neritina</i> (Linnaeus, 1758)	E C	<1911	unclear	V, AQ	regionally, established	fouling	Ryland 1960; Hayward and Ryland 1995; Arenas et al. 2006; Ryland et al. 2011
<i>Bugula simplex</i> Hinks, 1886	C E N	<1900	W. Atlantic	V, AQ	localities,	unknown	Ryland 1958; Ryland et al. 2011
<i>Bugula stolonifera</i> Ryland, 1960	CE	<1960	Atlantic	V	localities established	unknown	Ashelby 2005; Ryland and Hayward 1997; Ryland et al. 2011
<b>Candidae</b>							
<i>Tricellaria inopinata</i> d'Hondt & Occhipinti Ambrogi, 1985	E C	1998	NW Pacific	V	regionally, established	fouling	Dyrynda et al. 2000
<b>Watersiporidae</b>							
<i>Watersipora subtorquata</i> (d'Orbigny, 1952)	E	2008	unclear	V	localities, established	unknown	Ryland et al. 2009

## Appendix 1 (continued).

Species	Region	Date of first record placed within the salinity range of the recipient zone 1-30psu    >30psu	Native Range	Likely Vector/ Pathway	Current status	Impact (known or expected in Britain)	References
<b>Cordata, Ascidiacea</b>							
Botryllidae							
<i>Botrylloides violaceus</i> Oka, 1927	E C	2004	NW Pacific	V	regionally, established localities, established	fouling	Arenas et al. 2006; JS Ryland (pers. comm.)
<i>Botrylloides c.f. diegensis</i> (Ritter & Forsyth, 1917)	E	2004	N Pacific	V	localities, established	unknown	Faasse 2006; A. Gittenberger (pers. comm.)
Corellidae							
<i>Corella eumyota</i> Traustedt, 1882	E C	2004	circum sub-Antarctic	V	regionally, established	fouling	Arenas et al. 2006; Dupont et al. 2007; Collin et al. 2010; J Nunn (pers. comm.); JDD Bishop & JS Ryland (unpub.)
Perophoridae							
<i>Perophora japonica</i> Oka, 1927	E	1999	NW Pacific	V	localities, established	unknown	Nishikawa et al. 2000; Baldock and Bishop 2001, JDD Bishop (pers. comm.)
Styelidae							
<i>Styela clava</i> Herdman, 1882	E C N	1953	1953	NW Pacific	V, AQ	widespread, established	fouling
Didemnidae							
<i>Didemnum vexillum</i> Kott, 2002	E C	2005	NW Pacific	V, AQ	regionally, established	fouling	Holt et al. 2008; Griffith et al. 2009; Stefaniak et al. 2009
<b>Chordata, Pisces</b>							
Salmonidae							
<i>Oncorhynchus gorbuscha</i> (Walbaum, 1792)	C N	1960	N Pacific	AQ	individuals, not established	unknown	Wheeler 1969; Svetovidov 1986;
<i>Oncorhynchus mykiss</i> (Walbaum, 1792)	C	1884	1884	NE Pacific	AQ	localities, established	unknown
<i>Oncorhynchus keta</i> (Walbaum in Artedi, 1792)			1960s	N Pacific	AQ	individual, not established	unknown
<i>Oncorhynchus kisutch</i> (Walbaum, 1792)	E	1975	NE Pacific	AQ	individual, not established	unknown	Eno et al. 1997; Svetovidov 1986