

***Anticoma filicauda* n. sp.**

Fig. 3 a, b

Localities and material. — Falkland Islands, St. 59: 1♂; South Georgia, St. 22: 1♀, St. 25: 3♂♂, 3♀♀, 1 juv.

Dimensions: ♂ L = 2,720 mm.,  $a = 34,0$ ,  $\beta = 5,23$ ,  $\gamma = 7,22$   
♀ L = 2,775 mm.,  $a = 34,69$ ,  $\beta = 5,55$ ,  $\gamma = 7,4$ , V = 46,85 %  
♀ L = 3,0 mm.,  $a = 29,4$ ,  $\beta = 5,0$ ,  $\gamma = 6,0$ , V = 43,33 %.

In its general shape this species also is very similar to *A. acuminata* from which it seems to differ only in the *more elongated* and *evidently more strongly thinned tail*, the conical part of which makes 100  $\mu$  and the thinned part 277  $\mu$ . It contains only 2 glands. The spicules are very slender, strongly curved, evenly thick and measure  $1,6 \times$  the anal body diameter. Supplementary organ seems to be lacking.




Fig. 3. *Anticoma filicauda* n. sp. a. Anterior end,  $\times 450$ . b. Tail and spicules,  $\times 270$

St. 22. South Georgia. Off the May-Bay.  $54^{\circ} 17' S.$  —  $36^{\circ} 28' W.$  Clay with some algae. 75 m. Bottom temp. + 1,5 C. 14. 5. 1902.

Number of species found: 26; Number of specimens found: 85.

St. 25. South Georgia. Off the "Kochtopf"-Bay  $54^{\circ} 22' S.$  —  $36^{\circ} 27' W.$  Grey clay with some algae. 24—52 m. 21. 5. 1902.

Number of species found: 29; Number of specimens found: 83.

St. 59. Falkland Islands. S. W. West Falkland. On the Burdwood-Bank.  $53^{\circ} 45' S.$  —  $61^{\circ} 10' W.$  Crushed shells with stones 137—150 m. 12. 9. 1902.

Number of species found: 20; Number of specimens found: 70.