

***Anticoma longissima* ALLGÉN**

Fig. 2 a, b

ALLGÉN 1958, p. 208, fig. 2 a, b

Localities and material. — Coast of North Argentina, St. 2: 2♂♂, 2♀♀, 1 juv.; Falkland Islands, St. 58: 1♀

Dimensions: ♂ L = 2,835 mm.,  $\alpha = 74,61$ ,  $\beta = 5,15$ ,  $\gamma = 15,32$

♀ L = 2,500 mm.,  $\alpha = 50,0$ ,  $\beta = 4,55$ ,  $\gamma = 12,5$ , V = 58 %

From the same localities from which *Anticoma tenuis* is dating also another species is present, characterized by the strongly elongated and very slender body, being absolutely evenly thick from the posterior end of the oesophageal region until the anus and tapering only scarcely noticeably in the oesophageal region.

Diagnose: Cuticle thin and smooth; head rounded, provided with very short, submedian bristles. (Cervical hairs were not found.) Oesophagus of mediate length, posteriorly gradually and only very little increasing. Tail tapering in the first third conically and in the 2 posterior thirds uniformly more strongly thinned. Spicules short, slender, only slightly curved, rounded at the proximal end. Supplementary auxilliary organ situated immediately in front of the spicules. Vulva situated caudad to the middle of the body. Female organs paired-symmetrical, with reflexed ovaries.

St. 2. Off the Coast of North Argentina. 37° 15' S. — 56° 8' W. Sand-mixed gravel. 100 m. 23. 12. 1901.  
Number of species found: 12; Number of specimens found: 26.

St. 58. Falkland Islands. S. W. West Falkland. 52° 29' S. — 60° 36' W. Sand and gravel. 197 m. 11. 9. 1902.

Number of species found: 23; Number of specimens found: 93.




Fig. 2. *Anticoma longissima* ALLGÉN a. Anterior end,  $\times 450$ , b. Tail and spicules,  $\times 270$