

## **Synonchus fasciculatus** COBB

*S. fasciculatus* COBB 1893, p. 411, fig. 20; *Cylicolaimus armatus* DITLEVSEN 1923, p. 192, fig. 15—19; *Cylicolaimus armatus* ALLGÉN 1933, p. 16—17, fig. 2.

Locality and material. — Falkland Islands, St. 55: 1 young ♂

Dimensions:  $L = 5,900$  mm.,  $\alpha = 57,84$ ,  $\beta = 5,36$ ,  $\gamma = 77,6$

From the Falklands a single male of a *Leptosomatid* is present, which in all respects seems to agree rather well with *Synonchus armatus*, described by DITLEVSEN (1923) from the Atlantic Coast of France and refound by the author in Norway (1933). This species, indeed, later on has been regarded as synonymous with the old *Synonchus fasciculatus* COBB.

Geographical distribution. — Norway: Stappen (ALLGÉN 1943), Stjernesund and Reis (ALLGÉN 1940 (1)), Bud (ALLGÉN 1939), Trondheimsfjord (ALLGÉN 1933), France: Atlantic Coast, Bretagne, rivière de Trinité (DITLEVSEN 1923), Mediterranean: Bay of Naples (COBB 1893).

St. 55. Falkland Islands. Port Albemarle.  $52^{\circ} 11' S$ . —  $60^{\circ} 26' W$ . Sandy bottom with algae. 40 m. 8. 9. 1902.

Number of species found: 33; Number of specimens found: 113.