

HARPACTICOIDA - AMEIRIDAE

FILEXINIA CONROY-DALTON + HUYS, 1997

- CONROY-DALTON + HUYS, 1997:

= FILEXILIA

Canthocamptidae - Harpacticoida.

Ameira attenuata Thompson, ♀. ①

— I. C. Thompson, 1893, p. 195: pl. 32, f. 1-11.

— Sars, 1911, p. 397.

— tenella Sars, 1907, p. 220: pl. 145.

—, LANG, 1948, p. 795.

Habitat. British Isles: Norway Coast.

(2)

FILEXILIA ATTENUATA (THOMPSON I. C., 1893)

E- CONROY-DALTON + HUYS, 1997: 318, 322-327,

FIGS. 3 F, 4, 5 A, D, REDESCRIPTION; SYNS. AMEIRA

BREVIPIES ♂, A. TENELLA SARS. N. COMB.

= FILEXILIA ATTENUATA Canthocamptidae - Harpacticoida

Ameira tenella G.O. Sars, ♀.

— — Sars, 1907, p. 220: pl. 145.

— — Klie, 1949 (Harp. v. Helgoland)

A synonym of Ameira attenuata.
(Sars, 1911, p. 397).

— — Kunz, 1954, pp. 225 - 226.

— —, LANG, 1948, p. 793,

HABITAT: ENGLISH WATERS, IRELAND, MAYO.

— — —, Wells, 1970, p. 263

~ Moore, 1976: 42 - 43, fig. 7

--, Jakobi, 1959, p. 278

--, Klie, 1950, p. 89, fig. 82

--, **SOYER, 1974d: 390**

-- KONDALARAO 1984: 82.

-- ARUNACHALAM + BALAKRISHNAN NAIR 1988: 517, INDIA

-- KUNZ 1983: 120, 162 - 165, FIGS. 25, 26, AZORES

-- BODIN 1964: 161

FILEXILIA AZORICA CONROY-DALTON & HUYS, 1997

-- CONROY-DALTON & HUYS 1997: 318, 327-328, NOM.

NOV. FOR AMEIRA TENELLA SENSU KUNZ, 1983.

= FILEXILIA

Ameira brevipes Kunz

①

1954---Kunz, Von Helmut, pp. 224-225, figs. 3-11.

-- --, Wells, 1963. p. 90

-- --, Wells, 1970, p. 263

--, SOYER, 1974 d: 400, 401

--, Kunz, 1974, p. 264

-- KUNZ 1983:164-, s. STR.

-- APOSTOLOV 1977:15, 17, FIG. 6, NORTH SEA, BULGARIAN
COAST.

= FILEXILIA

AMEIRA BREVIPEDES KUNZ

(2)

FR. CONROY-DALTON & HUYS 1997: 322, 327, 331, 335;

♂ OF A. ATTENUATA THOMPSON = A. BREVIPEDES;

IN FILEXILIA, N. COMB. - GERMANY + ISLE OF MAN

FILEXILIA BREVIPES (KUNZ, 1954)

③

FILEXILIA BREVIPES PONTICA (APOSTOLOV, 1969)

AMEIRA --, APOSTOLOV 1969:

A-7, APOSTOLOV 1977:15

A.--, KUNZ 1983:164-

F.--, N.COMB., CONROY-DALTON & HUYS 1997:333,335;

SP. INQUIRENDAE IN FILEXILIA N.G.

FLEXILIA GRAVELLICOLA (Guilke + Soyfer, 1966)

ANEIRA -, Guilke + Soyfer 1966: 369-373, fig. 12, N.S.F.

A. -, Soyfer 1974: 400, 401

F. -, N. comb., Conroy-Darmon & Hays 1997: 318, 335, Tab. I

FILEXILIA INTERMEDIA (GALHANO, 1970)

AMEIRA -, GALHANO 1970:

F. -, N. COMB., CONROY-DALTON & Huys 1997: 318, 335, 336,
TAB. I

FILEXILIA

Ameira longifurca (Bodin, 1964)

A.- Bodin 1964: 140, 159-161, Pl. XVI, n. sp. female +
- male, Gulf of Marseilles, Mediterranean

A.--- Kunz 1983: 164-

A.- CONROY-DALTON & HUYS 1997: 327, 335; NOT =

A. TENELLA

F.-, N. COMB., CONROY-DALTON & HUYS 1997: 318, 328-331,
FIG. 5 B, E, 6, 7 B-E

FILE X141A MARINOV CONROY-DALTON & HUYS, 1997
-- CONROY-DALTON & HUYS, 1997: 318, 335, 336, TAB I,
NOMEN NOVUM FOR AMEIRA BARBIPES PESTAE
SENSU MARINOV

FILEXILIA PESTAE (PETKOVSKI, 1955)

AMEIRA PESTAE, PETKOVSKI 1955: 132-134, FIGS. 21-35, N. SP.

A. P., RAO + GANAPATI 1969: 13

A. P., KUNZ 1974: 264

A. BREVIPEDES P., APOSTOLOV 1977: 15

A. P., KUNZ 1983: 164-

FILEXILIA PESTAE, N. COMB., CONROY-DALTON + HUYS 1997:

~~318-323, FIGS. 13A, E, 14A, B, 333-335, TAB. I~~