

ⁱⁿ Siphonostomatoida Catigoidae Galigidae.
Achtheinus - valid Pandaridae

Wilson, 1908. ^(M.) A. oblongus.

Syn. Achtheinus Wilson, 1908, p. 450.
= Achtheinus Stebbing, 1918, p. 40.

'Ax θεινός, burdensome, troublesome

First thorax seg't. fused with head - 2nd
+ 3rd seg'ts fused into re and carrying
but a single pair of dorsal plates - 4th.
seg't. free + carrying a pair of plates -
Gen seg't. smaller than carapace -
Abdomen wholly concealed - frontal
plates fused with carapace - no first
anx. 2nd mspds. with terminal claws.
all swimming legs biramous - rami of first
3 pairs 2-jointed, of 4th pair 1-jointed all
rami rudimentary with spines only.

Achtheinus chinensis Thomsen, 1949, pp. 22-23.

Philodopus (Achtheinus) intermedius Kurtz. -Yu, 1933,
pp. 120-123, pl. 3, fig. 1-8. Host: Squalus
mitsukuri Jordan & Snyder, on skin. China.

Achtheinus dentatus. n. sp. ♀ Caligidæ - Caligoidæ.

- . Wilson, 1911, p. 630 = pl. 67, figs. 22-31.
— Stebbing, 1918, p. 41; pl. 97; 17 figs.
— Wilson, 1921, p. 6: pl. 3, figs. 20-27.
— ♂ Wilson 1944, p. 537, pl. 33, figs. 196-208
— , Brian (A.), 1944, pp. 203-205, pl. 2, fig. 15-17; pl. 3, fig. 18-29.

Host. Body and fins of Soup-fin Shark (Galeus Zyopterus). Atopias vulpes, Mustelus lumulatus Poecimaya, Peru.
Mustelus asterias, Heptanchias rectirostris, C.

— — , Nunes-Ruivo, 1956, pl. 12-15, pl. 3. Fins of Squalus
acanthias L., Angolo.

Achtheinus dentatus Wilson 1911

-- Kensley & Grindley 1973: 124

Pandariidae - Caligoidea

Achtheinus galeorhini Yamaguti ♀

— Yamaquti, 1936, p. 11; pl. 7, figs. 67-78.

--, Yamaguti & Yamasu, 1959, p. 90

Host. Gills, Galeorhinus manazo (Bleeker),
Japan.

Achtheinus galeorhini Yamaguti, 1936

-- Oldewage 1992: 49-

Achtheinus impudens Shen + Wang

— — Shen + Wang, 1959. Acta Zool. Sinica 10(1): 27-31.

From Triakis scyllium, Peitaiho, Hopei Province, China.

Caligidae - Caligoidea

Achtheinus intermedius (Kurtz).

Pholidopus (Achtheinus) i - Kurtz, 1924, p. 614: pl. I, f. 1-12

— — Yu, 1933, p. 120: pl. 3, figs. 1-8

Achtheinus - Wilson, 1927, p. 397.

Host, Skin of Triakis semifasciatum and
Squalus mitsukurii. China

Caligidae Caligidae

Achtheinus japonicus n. sp. ♀

— Wilson, 1922, p. 4: pl. 2, f. 9-17

Host Fin of Acanthias sp.

Japan.

Achtheinus oblongus -- see Perissopus oblongus

Pandarinae - Caligoidea.

Achtheinus parvidens New species ♀.

— Wilson, 1923, p. 7: pl. 1, figs. 6-12.

Host. Outside surface of Acanthia vulgaris.

South Africa.

Achtheinus pinguis, new species, ♂. ♀.

— — Wilson, 1912, p. 235: pl. 31, figs. 8-14:
pl. 32, figs. 15-21.

— — Wilson, 1923, p. 5.

- -, BASHIRULLAH, 1975: 248

Host. Pectoral fin of Pliotrema warreni:
Dogfish Shark.

Cape-of-Good Hope.

- Capart, 1959: 97-98

Achtheinus cf. pinguis Wilson

-- Kensley & Grindley 1973: 124

Caligidae-Caligoidea

Achtheinus platensis Thomsen

Thomsen, Ricardo, 1949, pp. 20-25, pl. VII