

HARPACTICOIDA

AMEIRIDAE MONARD, 1927, LANG, 1936

- FIERS 1996: 71, CF. CANCRINOLIDAE N. FAM.
- ROUCH 1992: 152, PSAMMONITOCRELLA N.G.
- MIELKE 1994c: 252, PSAMMOLEPTOMESOCYTRA N.G.

AMEIRIDAE 3P.

-- MOORE, C. G. + O'REILLY 1993: 653, TAB. 1, SCOTLAND

-- COTTARELLI, BRUNO + FORNIZ 1996: 263, TAB. 2,
SARDINIA

Harpacticoida

Superfam. AMEIRIDIMORPHA, nov.

_____ Lang 1948, p. 779

Ameiridae-Harpacticoida

Subfam. AMEIRINAE, nov.

_____ Lang 1948, p. 786

in) Canthocamptidae - Harpacticoida.
Ameiridae

Ameira, valid

Boeck, 1865,

A. longipes ^(M)

Syn. Ameira Boeck, 1865, p. 273.

• Ameira Sars, 1907, p. 214.

Ameira Brady, 1880. vol. 2. p. 36.

Ameira Pesta, 1921, p. 614.

Nicholls 1939b & 1940, p. 270-272 Key

Monard 1935a, p. 46

Lang 1935

Anterior antennae 8-jointed; endopod of posterior antennae 1-jointed, small. Mandible-palp small, 1- or 2-branched. Rami of all the swimming legs 3-jointed; exopod of first pair short, endopod long, the first joint much elongated.

Canthocamptidae - Harpacticoida:

Ameira ambigua T. Scott,

— T. Scott, 1903, p. 114: pl. 5, fig. 11-19.

— T. Scott, 1906, b, p. 325.

A Synonym of Ameira minuta.
(Sars, 1907, p. 216).

Canthocamptidae - Harpacticoida -
Ameira amphibia Brady, ♀.

— — Brady,

A synonym of Nitocra typica.
(Sars, 1907, p. 212).

Canthocamptidae - Harpacticoida

Ameira arenicola Liang, ♀.

— — Liang, 1935, p. 3: text-figs. 1-13.

— — Liang, 1938, p. 800.

Habitat, Öresund, 3m. depth.

Ameira ariana (Monard) - see Ameiropsyllus

Ameira atlantica Noodt

-- -- Noodt, 1958. pp. 72-77, fig. 72-86

--, Kunz, 1974, p. 264, 266 (tbl. 6)

Ameira atlantica Noodt f. mediterranea forma n.

--, Kunz, 1974, p. 264 (figs. 34-44, 47-52), 266 (tbl. 6)

AMEIRA ATTENUATA - SEE FLEXILIA

Ameira bathyalis

BECKER

1979

-- Schriever 1982b: 34

-- Kunz 1984d: 52

AMEIRA BENGALENSIS - SEE GLABRAMEIRA

Canthocamptidae-Harpacticoida

Ameira boeckii Lang ♀

— — Lang, 1936, p. 32 - 33 : figs. 45 - 47
-- KRISHNASWAMY 1957d: 129

Habitat. Spitzbergen.

Ameira brevicornis

-- Bodin 1964: 161

AMEIRA BREVIPEDES - SEE FILEXILIA

AMEIRA BREVIPEDES PESTAE - SEE FILEXICIA PESTAE

AMEIRA BREVIPEDES PONTICA - SEE FILEXILIA

Canthocamptidae - Harpacticoida

Ameira breviremis Brady,

- - Brady, 1905, p. 214: pl. 3, fig. 11-14; pl. 4, fig. 1-6.
- - Norman & Brady, 1909, p. 138.
- - Sars, 1911, p. 392 (= *Stenkelia gibba*).

Habitat, British Seas,

Ameira confluens Ranga Reddy, 1984

--Ranga Reddy, 1984: 95-99, 103, figs 1-12 n. sp. ♀♂

-- CONROY- DALTON + HUYS 1997: 318

Canthocamptidae - Harpacticoida

Ameira curviseta Willey, ♀

— — Willey, 1929, p. 538: figs. 29-30

— —, LANG, 1948, p. 798.

Habitat. St. Andrews, N.B. CANADA.

~~Diosaccidae~~-Harpacticoida
Ameiridae

Ameira divagans Nicholls

Nicholls 1939^b, p. 276, fig. 11, St. Lawrence ♀
Nicholls 1940, p. 276, fig. 11

--, Scheibel, 1974, p. 1-10, figs. 1-18

-- KRISHNASWAMY 1957d: 129

-- LANG 1965: 334, 335

Ameira divagans Nicholls africana Kunz

n. forma, Kunz, 1963. pp. 43-46, fig. 57-68

--, Scheibel, 1974, p. 1, 10

-- LANG 1965: 334, 335

Ameira divagans Nicholls, 1939 pontica Marinov, 1973

-- --, Marinov, 1973, p. 231-233

Amelra elegans T. Scott, ♀

syn.

Pl. 11, fig. 1-9

— — T. Scott, 1905.7, p. 144; pl. 10, fig. 18-19; ^

— — T. Scott, 1906.7, p. 325.

=Nitocra elegans, Monard 1935a, p. 56

Habitat. British Seas.

Ameira exigua T. Scott

— — T. Scott, 1894, p. 243, pl. 6, fig. 15-23.

— — T. Scott, 1906, b, p. 324; 1899^c, p. 97.

— —, LANG, 1948, p. 745.

— Thompson, 1895: 101

Habitat. British Seas.

-- CHISLENKO 1977b: 272, FRANZ JOSEF LAND

Ameira exilis T. & A. Scott, ♀.

— — T. & A. Scott, 1894, p. 139: pl. 8, fig. 18-20; pl. 9, fig. 1-3.

— — Thompson & Scott, 1900, p. 140

— — T. Scott, 1894, p. 242: pl. 9, fig. 30; pl. 10, fig. 1-12.

— — T. Scott, 1906, 6, p. 324.

--, Lang, 1944, p. 25, a syn. of ~~*~~ Pseudosarsameira e.

Habitat. British Seas.

Ameira faroerensis Schriever 1982

-- Schriever 1982b: 34, fig. 5, n. sp. ♀ only; North
Atlantic

-- Kunz 1984d: 52

AMEIRA GRACILE - SEE SICAMBIRA

Canthocamptidae - Harpacticoida.

Ameira gracilimana (Giesbrecht)

Nitocra - Giesbrecht, 1902

Ameira - Gurney, 1920, p. 220.

Habitat. Belgium -

Ameira grandis Nicholls

Nicholls, 1939 b, p. 274-276, fig. 10, St. Lawrence ♀ ♂
Nicholls 1940, p. 274, fig. 10

- KRISHNASWAMY 1957d: 129
- BODIN 1977: 332
- LANG 1965: 334, 335, 336, "SYSTEMATIC POSITION IS PROBLEMATIC"

AMEIRA GRAVELLICOLOA - SEE FILEXILIA

PROAMERIA MIDDEN SOENSIS (SCHAFER)

AMERIA ——— SCHÄFER, 1936 S. 16.

PRO ———, LANG, 1948, P. 802.

HABITAT: GERMANY; MIDDENSEE.

Psammameira hyalina (Wood, 1952)

- Mielke, 1975: 63-65, fig. 40.

[syn. Psammameira hyalina]

AMEIRA INTERMEDIA - SEE FILIXILIA

Ameira leptoderma (Klie)

Sicameira leptoderma Klie, 1950.

Ameira leptoderma, Wells, 1963. pp. 90-92

— Wells, 1967: 193

Armeira listensis Mielke

- Mielke, 1973: 4-10, fig. 1-3A, B.
- Mielke, 1975: 62-63, fig 39.

Canthocamptidae - Harpacticoida
ameiridae

Ameira longicaudata T. Scott, ♀♂.

— T. Scott, 1892, p. 250: pl. 9, figs. 1-18.

— Thompson, 1895: 101

-- A. Scott 1896: 140

A synonym of Stenocopia longicaudata.
Sars, 1907, p. 228.

AMEIRA LITTORALIS - SEE NITOKRA TYPICA
Nicholls 1949

Selmincola thymalli

Shelf 30

AMEIRA LONGICAUDATA NICHOLLS - SEE FILEXILIA
TRISSETOSA

AMEIRA LONGIFURCA - SEE FILEXILIA

Amelra longipes Boeck, ♀.

— — Boeck, 1865, p. 273 (49).

— — Sara, 1916, p. 7.

Not Brady, 1880, vol. 2, p. 37: pl. 53, fig. 1-10.

— — T. Scott, 1906, b, p. 323; 1903, p. 12.

— — Chappuis, 1958 p. 411

Habitat. British Seas; Mediterranean.

CANADA, TUNISIA, SWEDEN, NORWAY.

— Scott & Scott, 1901: 343

- — Sars, 1907, p. 215: pl. 140.
— — Pesta, 1921, p. 615: text-fig. N 55.
— — T. Scott, 1949^c, p. 97.
— — Thompson, 1893, p. 495: pl. 19. f. 7, a-d.

— — Ameira longipes, Monk 1941, p. 98, pl. 2, figs.
12-13 ♀ California

— — Monard 1935a, p. 52, Roscoff, France

- —, LANG, 1948, p. 788,
— —, Wells, 1963, p. 15
— —, Wells, 1964, p. 454
— —, Wells, 1970, pg. 262

Ameira longipes Boeck, 1865

- , Klie, 1950, p. 88
- , Lang, 1944, p. 24
- , WELLS, 1979: 3
- , CHOLENKO, 1977: TAB. 1
- , Coffin, 1978-79, p. 593, NEW ENGLAND
- Mielke, 1974: 31, fig 17.
- Pallares, 1975: 219-221, fig 8 (#1-11)
- Knatz, 1986: passim
- Lang, 1965: 340-343, fig. 184-185, pl. V, fig. a.
CALIFORNIA
- Wells, 1967: 193
- Willey, 1923: 330

Ameira longipes Boeck, 1865

card 3

-- Schriever 1982b: 34

-- SHIH ET AL. 1971: 209, CANADA-ARCTIC

-- KASK ET AL. 1982: 36, NANAIMO, BRITISH COLUMBIA

-- CHISLENKO 1977b: 255, FIG. 11 (4-7), FRANZ
JOSEF LAND

-- PESTA 1959: 140-, FIGS. 108-114, OFF NAPLES, ITALY

-- WILSON, C.B. 1936: 371, FROZEN STRAIT, CANADIAN ARCTIC

-- CONROY-DALTON & HUYS 1997: 336

Ameira, longipes- Group

---Ranga Reddy 1984: 99

Canthocamptidae - Harpacticoida.

Ameira longipes Brady, ♀.

— — Brady, 1880, vol. 2, p. 37: pl. 53, figs. 1-10.

— — Thompson

-- --, Chappuis, 1957. Puget Sound.

a synonym of Ameiropsis brevicornis,
Norman & Brady, 1909, p. 137.

--, Vervoort, 1964, p. 5, 6, 251-253, 255, 260, 264,
figs. 100-102

--, Roe, 1958: ²³⁰250; Fig. 130

--, Jakobi, 1959, p. 137

-- Hicks 1980: 164-, BRITAIN

-- GRANDORI 1913: 166, VENICE LAGOON

Canthocamptidae - Harpacticoida -

Ameira longiremis T. Scott, ♀.

- - T. Scott, 1894, p. 241: pl. 5, f. 29-32: pl. 6, f. 1-5.
- - T. Scott, 1906, b, p. 324: 1899^c, p. 97.
- - var. *intermedia*, n. var. T. Scott, 1894, p. 242: pl. 6, f. 6-14.
- Thompson, 1895: 101

Habitat. British Seas.

Canthocamptidae - Harpacticoida.

Ameira minor Thompson & Scott, ♀.

— — Thompson & Scott, 1903, p. 264: pl. 5, f. 23-29.

— —, LANG, 1948, p. 797.

Habitat. Pearl oyster washings, Ceylon.

Canthocamptidae-Harpacticoida.

Ameira minuta Boeck, ♀. ①

— — Boeck, 1865, p. 273.

— — Sars, 1907, p. 216: pl. 141.

— — ambigua T. Scott, 1903, p. 114: pl. 5, f. 11-19

— — T. Scott, 1906, b, p. 325.

— —, LANG, 1948, p. 789,

Habitat. British Seas: Norway Coast.

--, KUE, 1950: 88

--, ROE 1958: 230

- — Klie, 1927, p. 14:
— —, Wells, 1963. p. 15
— —, Wells, 1970, p. 263
— —, Pesta, 1932, p. 72, fig. 1
— —, Vervoort, 1964, p. 5, 6, 257-259, 262, fig. 103, 104 -
— —, CHISLENKO, 1977: TAB. 1 IFALUK ATOLL
— —, HICKS, 1977a: 455, 457; TAB. 3, 4
— —, Cull, et al., 1983: passim.
— — SCHRIEVER 1982b: 34
— — CRANDELL 1967: 38
— — KASK ET AL. 1982: 36, NANAIMO, BRITISH COLUMBIA
— — HICKS 1980: 165-, BRITAIN

AMEIRA MINUTA BOECK, 1865

(2)

-- CONROY - DALTON & HUYS 1997: 336

Ameira parascotti Chislenko, 1977

-- Chislenko 1977b: 255-257, Fig. 12(5-7), n. sp.

female only, Franz Josef Land

Ameira parasimulans Lang

—Lang, 1965: 343-347, fig. 186-188.

Canthocamptidae - Harpacticoida.

Ameira parva Boeck, ♀

— — Boeck, 1873, p. 49

Habitat. Norway-

Ameira parvula (Claus), ♀

— — Klie, 1929, p. 351:

— — Willey, 1935, p. 52:

— — Willey, 1932, p. 90: fig. 20.

— — Nicholls 1939b & 1940, p. 272

— — Gurney 1927, p. 545 syn.

Habitat. Baltic Sea; Kieler Bay; Bermuda.

Amphiascus parvulus - syn. Monard 1928,

Nicholls 1941b, p. 85

— — Monard 1935a, p. 52, Roscoff, France

Ameira parvula, Sewell 1940, p. 301, ♀, Maldives Arch.

— —, *NOODT, 1955, P.P. 207,*

— —, *NOODT, 1956. P.P. 52, N. SEA AREA*

— —, *HANG, 1949, P.P. 790. FJORDS NORWAY.*

— —, *Petkovski, 1964. pp. 12-13, fig. 30-32 (s. str.)*

— —, *Vervoort, 1962. pp. 304-451, fig. 22b, 24a, b, 25*

— —, *Wells, 1963. p. 15*

— —, *Wells, 1964 p. 454*

— —, *Wells, 1965 p. 4*

— —, *Por, 1960, pg. 117*

— —, *Wells, 1970, p. 263*

— —, *Coull, 1971, p. 208, 218*

— —, *Pesta, 1932, p. 72-3, fig. 2*

— —, *SOYER, 1974 d: 400, 401*

Ameira parvula

card 2

--, Dolzonolskaya, 1970, p. 86, pl. XXIII, fig. 2, syn.:

Canthocamptus p.; Nitocra tau

--, Vervoort, 1964, p. 5, 261-263, figs. 105ab, 106a-f

--, Jakobi, 1959, p. 135

--, Klie, 1950, p. 88-9, figs. 81

--, CHASLENKO, 1977) TAB. 1

— Kunz, 1975: 186 —

— Pallares, 1975: 221 — , figs 53-58

, fig. 8 (12-19)

~ Wells, 1967: 193

Ameira parvula (Claus)

--, Stock & Vos, 1960, p. 206

--, Klie, 1950, p. 88-89, fig. 81, Canthocamptus p.
is a syn. ; A. tau is a syn.

--, MARCUS, 1965: 91

--, HICKS, 1977a: TAB. 3, 4

--, ANGER & SCHIEBEL 1976: pp. 4-6, Tab. 1.

--, BRIAN 1926: p. 248 - makes a synonym of Canthocamp-
tus parvulus & Nitocra tau.

--, BOZIC 1954: 8; Pl. 4, figs. 4-6.

--, BRIAN 1923b: 870.

- Mielke, 1975: 62, fig. 38.
- Noodt, 1952: 109
- Michailova - Neikova, 1968: 217
- .. MASONARI 1985: 53, SANTOS BAY, SÃO PAULO, BRAZIL

Ameira parvula (Claus)

card 4

- , Becker, Von K., 1970, p. 59
- , Moore, C.G., 1976, p. 455 (fig. 3)
- , Coffin, 1978-79, p. 593, NEW ENGLAND
- , Coull, et al, 1983: passim
- Mielke, 1974: 33-34, fig 18.
- Moore, 1976: 39-42, fig 6
- CECCHERELLI + ROSSIN 1979: 115-116, 122
- MASUNARI 1985: 53, SANTOS BAY, SÃO PAULO, BRAZIL
- HOCKIN 1984: 320, ENGLAND

- Serban, 1959: 271-272
- Vorobjeva, 1984: 47
- CARLI & CRISAFI 1983: 106, PL. 24 FIG 1, ITALY
- SCHRIEVER 1982b: 34
- DRZYCIŃSKI 1986: 552-, BALTIC
- POR & MARCUS, 1972: 269 SUEZ CANAL

Ameira parvula (Claus)

card 5

- Por, 1979: 13, 14 Gulf of Elat, Red Sea

-- CECCHERELLI + FERRARI 1982: 345, LAGOONS, ITALY

-- CARRADA ET AL. 1987: 155-, LAGOONS, ITALY

-- SHIH ET AL. 1988: 321-, BRAS D'OR LAKES, NOVA

SCOTIA

-- SHIH ET AL. 1971: 209, CANADA - ARCTIC (= A.TAU)

-- GOMOU 1971: 158, BLACK SEA

-- COULL 1968: 59-, BERMUDA

-- HICKS 1980: 165-, BRITAIN

-- REID 1990g: 195

Ameira parvula (Claus)

card 6

- cf. -, Eckman & Thistle 1991:1405-, off California

-- CHISLENKO 1977b: 255, FIG. 12(1-4), FRANZ
JOSEF LAND

-- BODIN 1964: 139-, GULF OF MARSEILLES

-- PESTA 1959: 142-, FIGS. 115(a-d)-117, OFF NAPLES,
ITALY

-- APOSTOLOV 1977: 15, 16, FIG. 5, NORTH SEA-BULGARIAN
COAST

-- CARMAN & DOBBS 1997: 121, TAB. 1

AMEIRA PARVULA (CLAUS)

(7)

-- POR 1962: 31, 32, BLACK SEA

-- CONROY-DALTON + HUYS 1997: 331, 336

= A. PARVULA

Canthocamptidae - Harpacticoida.

Ameira tau (Giesbrecht). ♀.

←

Nitocra tau Giesbrecht, 1882, p. 117: pl. 1, f. 9, 13:
pl. 3, f. 13: pl. 4, f. 2, 11, 29.

— — T. Scott, 1906, b, p. 331.

Ameira tau Sars, 1907, p. 218: pl. 143

Habitat. Scottish tide-pools (Scott): Bay of Keel
(Giesbrecht): Norway (Sars): Grinnel
Land: Adriatic.

- — Pesta, 1921, p. 616: text-fig. N56.
— — Klie, 1913, p. 28.
Ameira Parvula — Gurney, 1927, p. 545.
— tau - Monard, 1928, p. 405: fig. 36.
— — Wilson, 1932, p. 245: fig. 163
-- , Pesta, 1932, p. 72
-- , Vervoort, 1964, p. 264
-- , Klie, 1958, p. 88, a sym. of A. parvula
— Wiborg, 1944: 63

AMEIRA PARVULA TENUSETA (CHAPUIS)

--- APOSTOLOV 1977:16

Ameira parvula f. nana Willey, 1935

--- Dinet 1971: 756, Pl. 5 - Marseilles

--- BODIN 1964: 139; GULF OF MARSEILLES

Ammeira parvuloides Lang

→ Lang, 1965: 347-352, fig. 189-191

-- SCHRIEVER 1982b: 39

AMEIRA PESTAE - SEE FILEXILIA

Ameiridae-Harpacticoida

Ameira phaedra Monard

————— Monard 1935a, p. 53, fig. 101-107 ♀, Roscoff
Atlantic coast, France.

PRO —, LANG, 1948, P. 801.

Canthocamptidae - Harpacticoida.

Ameira propinqua T. Scott, ♀, ♂.

— — T. Scott, 1902, b, p. 460: pl. 24, fig. 10-18, ♀,
pl. 22, fig. 36-42: pl. 23, fig. 1, ♂.

— — T. Scott, 1906, b, p. 324.

Habitat, British Seas.

Canthocamptidae - Harpacticoida.

Ameira pusilla T. Scott.

— — T. Scott, 1903, p. 114: pl. 5, fig. 1-10.

— — T. Scott, 1906, b, p. 325,

— — Monard 1935a, p. 52, Roscoff, France. First record
since discovery

— —, LANG, 1948, p. 797,

Habitat. British Seas, SCOTLAND, FIRTH OF
FORTH.

-- KRISHNASWAMY 1957d: 129

-- --, Wells, 1970, p. 263

-- BODIN 1977: 329-332, FIG. 8, 9, RIVEDOUX, FRANCE

Ameira reducta Petkovski

---1954. Petkovski, Trajan K., pp. 102-105, figs. 32-41

--, Kunz, 1974, p. 264, 266 (tbl. 6)

Canthocamptidae - Harpacticoida -
Ameira reflexa T. Scott. ♀

— T. Scott, 1894, p. 240; pl. 5, f. 20-28,

— T. Scott, 1899^e, p. 97,

-- A. SCOTT 1896: 140, OFF ISLE OF MAN

Habitat Franz-Joseph Land -

Canthocamptidae - Harpacticoida.

Ameira reflexa, var. T. Scott, ♀.

— T. Scott, 1902, a, pl. 23, fig. 34-42.
(without description).

A synonym of Pseudameira crassicornis.
(Sars, 1911, p. 400).

Canthocamptidae - Harpacticoida.

Ameira scotti G. O. Sars, ♀.

Ameira tenuicornis Sars, 1907, p. 217: pl. 142

A — *scotti* Sars, 1911, p. 397.

— — Gurney, 1927, p. 545.

— — Monard, 1928, p. 404: fig. 36

Habitat. Norway Coast; Bermuda - Suez,
TUNISIA,

--, ROE, 1958: 230

— — Willey, 1932, p. 90: figs. 18-19-

— — Monard 1935a, p. 52, Roscoff, France

— —, LANG, 1948, P. 793,

— —, Wells, 1963, p. 15

--, Vervoort, 1964, p. ²⁵⁶~~5~~, ~~261-263~~, figs. 105a, b, 106a-f

--, Jakobi, 1959, p. 135

--, CHISLENKO, 1977: TAB. 1

card 2
Ameira scotti Sars

--, Hammond, 1968, p. 176

--, Klie, 1950, p. 88

--, WELLS, 1979: 3

--, SOYER, 1974d: 400, 402

--, BODIN 1970: p. 409; Tab. B.

--, BODIN 1972: p. 665; Tab. 2.

—, Hammond, 19—: p. 240, tab. 2.

-- PESCE 1983c: 63, 65, 79

— Serban, 1959: 272 Romania

-- SCHRIEVER 1982a: 34

Ameira scotti G. O. Sars

card 3

-- Hicks 1980: 164-, Britain

-- REID 1990g: 195

-- BODIN 1964: 139-, Pl. V FIG 36, GULF OF MARSEILLES

Ameira scotti brasiliensis

-- -- Jakobi 1954 pp. 228 fig. 13 .

Ameira scotti brevicornis Monard, 1928

---, Dolzonolskaya, 1970, p. 87, pl. XXIII, fig. 3,
syn.: A. tenuicornis

Canthocamptidae - Harpacticoida

Ameira sibogae A. Scott, ♀.

— — A. Scott, 1909, p. 222: pl. 60, f. 1-7.

— —, LANG, 1948, p. 797.

Habitat. Paternoster Islands.

Ameira simplex T. Scott, ♀.

— — Norman & Scott, 1905, p. 291.

— — Sars, 1907, p. 219: pl. 144.

— — Wilson, 1932, p. 244: fig. 162.

Proameira, Wells, 1965. p. 5

--, Lang, 1944, p. 25, a syn. of Proameira s.

Habitat. Norway Fjords.

Canthocamptidae - Harpacticoida

Ameira simulans T. Scott ♀.

— — T. Scott, 1912, p. 561: pl. 7, figs. 23-28

— —, LANG, 1948, p. 798.

Habitat. South Orkneys, Scotia Bay.

Ameiridae-Harpacticoida

Ameira speciosa Monard

Monard 1935a, p. 53, fig. 108-118 ♀♂, Roscoff,
Atlantic coast, France

---, Klie, 1950, p. 91
---, LANG, 1948, p. 795,

HABITAT: FRANCE, ROSCOFF.

-- SCHRIEVER 1982b: 34

-- CONROY - DALTON & HUYS 1997: 327

Diosaccidae-Harpacticoida

Ameira spinipes Nicholls

Nicholls 1939, b, p. 278, fig. 12, St. Lawrence, ♀ ♂
Nicholls 1940, p. 278, fig. 12

-- KRISHNASWAMY 1957d: 129

-- LANG 1965: 334, 335, 336, "SYSTEMATIC POSITION IS
PROBLEMATIC"

Ameira tau (Giesbrecht) - see A. parvula

AMEIRA TENELLA - SEE FILEXILIA ATTENUATA

Ameira tenuicornis T. Scott, ♀.

— — T. Scott, 1902.3, p. 459: pl. 24, fig. 1-9.

not — — Sars, 1907, p. 217: pl. 142.

— — Sars, 1911, p. 397: Suppl. pl. 27.

— — Pesta, 1921, p. 616: text-fig. N 57.

Habitat. British Seas: Norway Coast:
Adriatic.

— Makkaveeva, 1961: 154, Pl. 10-11

— — Wilson, 1932, p. 247: fig. 164.

— —, Lang, 1948, p. 792.

— —, Wells, 1970, p. 263

—, Dolzonolskaya, 1970, p. 87, a syn. of A. scotti
brevicornis

--, Jakobi, 1959, p. 142, 278

--, Klie, 1950, p. 91

-- SCHRIEVER 1982b: 34

-- GRANDORI 1913: 166, VENICE LAGOON

Canthocamptidae - Harpacticoida

Ameira tenuipes Thompson & Scott, ♀.

— — Thompson & Scott, 1903, p. 264: pl. 6, fig. 15-20
— —, LANC, 1948, p. 797.

Habitat. Pearl oyster washings, Ceylon.

Canthocamptidae - Harpacticoida.
Ameira tenuiremis T. Scott.

- — T. Scott, 1902, a, p. 459: pl. 24, f. 1-9.
— — T. Scott, 1906, b, p. 324.

Habitat. British Seas.

Ameira trisetosa Krishnaswamy, 1957

1957---pp. 230-232, text-fig. 1., n. sp., MADRAS

-- KRISHNASWAMY 1957d: 127-129, FIG. 33, MADRAS

D.D.

Ameira usitata Klie 1950.

Kiel Meeresforsch., VII: 89-92, fig. 83-84.

— Kunz, 1975: 187-188, figs 59-73.

Canthocamptidae - Harpacticoida.

Ameira sp.

— — Stephensen, 1913. b, p. 73

Habitat, Greenland.

AMEIRA SP.

- , RAO, G.C. 1967: 131, INDIA
- MURPHY & HOLMES 1990: 225, DUBLIN BAY, IRELAND
- CRANDELL 1967: 38-, YAQUINA BAY, OREGON
- HARTZBAND 1971: 12-, ST. THOMAS
- COTTARELLI 1971C: 29, ITALY
- FLEEGER ET AL. 1983: 940-941, 945, TABS. 1, 2,
CHANGJIANG ESTUARY, CHINA