

in

Canthocamptidae - Harpacticoida,
AMEIRIDAE

Ameiropsis valid.

Sars, 1907

A. brevicornis [⊕]

Syn. Ameiropsis Sars, 1907, p. 223.

Ameira and opsis, likeness.

Body stouter than in Ameira; caudal rami short. Anterior antennae 8-jointed, first 2 joints the largest. Exopod of posterior antennae 2-jointed. Mandible palp distinctly biramous, both rami 1-jointed. First mxpds. with 2 digitiform lobes inside the claw-bearing joint. First legs prehensile, endopod longer than exopod, 3-jointed; endopod of 3rd. legs not transformed in ♂.

AMEIROPSIS SARS, 1907

CARD 2

- BODIN 1977: 339

Canthocamptidae - Harpacticoida.
Ameiropsis abbreviata G. O. Sars, ♀.

— — Sars, 1911, p. 404: Suppl. pl. 33.

— —, LANG, 1919, p. 832.

— —, Jakobi, 1959, p. 278.

Habitat. Norway Coast.

Canthocamptidae - Harpacticoida.

Ameiropsis angulifera G. O. Sars, ♀.

— — Sars, 1911, p. 403: Suppl. pl. 32.

— —, LANG, 1948, p. 832.

--, Jakobi, 1959, p. 278

Habitat. Norway Coast.

Ameiopsis ariana Monard - see Ameiopsyllus

Ameiropsis australis Kunz

— Kunz, 1975: 193—196, fig 113-137.

Canthocamptidae - Harpacticoida

Ameiropsis brevicornis Sars, ♀.

— Sars, 1907, p. 224: pl. 148.

— Norman & Brady, 1909, p. 137.

Ameira longipes Brady, 1880, vol. 2, p. 37: pl. 53, fig. 1-10

— Monard 1935a, p. 52, Roscoff, France

— —, LANG., 1948, p. 830.

Habitat, British Seas; FRANK PERCIN, ROSCOFF, FR.

— —, Chappuis 1958 p 411

--, Jakobi, 1959, p. 278

--, ROE, 1958: 231

A. brevicornis, Chappuis, 1957. Puget Sound
--, Klie, 1950, p. 94-5, figs. 93, 94

Canthocamptidae-Harpacticoida:

Ameiropsis longicornis G.O. Sars, ♀.

— — Sars, 1907, p. 225: pl. 149.

— —, LANG, 1948, p. 830.

-- --, Wells, 1965. p. 5

-- --, Wells, 1970, p. 263

-- --, Por, 1964, p. 95, pl. 15 figs. 164-165

Habitat. Norway Coast.

--, Klie, 1950, p. 93-4, fig. 92

--, SOYER, 1974d: 406*

--, ROE, 1958: 281

--, Dolzonolskaya, 1970, p. 90, pl. XXV, fig. 4

-- KASK ET AL. 1982:36, NANAIMO, BRITISH COLUMBIA

.....
Ameiropsis minor

--, Jakobi, 1959, p. 278

Canthocamptidae-Harpacticoida

Ameiropsis mixta G.O. Sars. ♀.

— — Sars, 1907, p. 226: pl. 150.

— — Monard 1935 a, p. 51 ♀, Roscoff. First record since discovery

— —, *hanc*, 1918, p. 831.

— —, Wells, 1970, p. 263

Habitat. Norway Coast.

— —, ROE, 1958: ^{23,} 25 ♀; FIG. 137, 138

— — HICKS 1980: 165. BRITAIN

Canthocamptidae - Harpacticoida.

Ameiropsis nobilis G. O. Sars, ♀. /

— — Sars, 1911, p. 402: Suppl. pl. 31.

— —, LANG, 1948, p. 832.

— —, Wells, 1970, p. 263

— —, Guille & Soyer, 1966, p. 375

Habitat. Norway Coast. GRENAND.

—, Jakobi, 1959, p. 278

—, SOYER, 1974 d: 406

- CP. - DAHMS 1988 b. 17 - FIGS. HELGOLAND

Ameiropsis nobilis G.O. Sars, 1911

card 2

-- Crandell 1967:38

-- COULL 1968: 59-, BERMUDA

-- REID 1990g: 195

-- BODIN 1964: 140-, Pl. VII FIG. 39, 40, Pl. VI FIG. 37,
GULF OF MARSEILLES

Canthocamptidae-Harpacticoida

Ameiropsis robinsoni Gurney, ♀.

— — Gurney, 1927f, p. 549: fig. 155, A-G.

— —, LANG, 1948, p. 834.

Habitat. Suez Canal.

AMEIROPSIS SP.

- GRANDALL 1967:38-, YAQUINA BAY, OREGON
- ? SP. NOV., HICKS 1980:164-, BRITAIN