

Kernaeopodidae - Kernaeopodoidea

Anchorella = synonym.

Cuvier, 1830.

A. adunca (M)

Syn. Anchorella, Cuvier, 1830, III, p. 257.
Clavella, Oken, 1815, p. 358.

Anchorella

-, Kabata, 1968, p. 1413

-, ACLOQUE 1899: 203-4, key to genera.

- NERESH FIMER 1909b: 80

Ternaeopodidae. Ternaeopodoidea.

Anchorella agilis Krøyer, ♀.

— Krøyer, 1863, p. 300; pl. 16. f. 2, a-d.

A synonym of Clavella agilis.
(Wilson, 1915, p. 668).

Anchorella adunca, Ström. ♀.

Lernaea a—, Ström, 1762, p. 167; pl. I, figs. 7+8.

Anchorella a—, Cuvier, 1830, p. 257.

— —, M-Edwards, 1840, p. 520.

— — Wilson, 1915, p. 667.

Clavella —, Nunes-Riivo, 1957, pp. 230-245, fig. 1-51

A synonym of Clavella uncinata.

(over)

an "Anchorella", but not well enough
described to fix the species, made the
type of the genus by Cuvier -
The figures and description given by Ström
show this to be a Clavella uncinata
almost without doubt, and so Cuvier's
genus Anchorella has nothing upon
which to stand.

Lernaeopodidae.

Anchorella anomala, Abildgaard, ♀.

Lernaea. Abildgaard, 1794, p. 57: pl. VI, fig. 2.

Anchorella, Milne-Edwards, 1840, p. 520.

Anchorella Wilson, 1915, p. 668.

Host. Unknown.

Milne Edwards says "Abildgaard's Ternaea
anomala appears to belong also to this genus
(Anchorella), but differs from all preceding
species in the considerable length of the organ
of adhesion - and in the existence of a swelling
like a knob at the anterior end of the thorax.

This is probably a Clavella, but had better
be left where it is until further identification

Leuconaeidae.

Anchorella appendiculosa Kröyer ♀

Anchorella —, Kröyer 1863, p. 306: pl. 16, fig. 6a-c.

— — . Vogt, 1877b, p. 65.

— — . B. Smith, 1899, p. 506.

Transferred to Genus Brachiella 1914.

NROBRACHIELLA —, CASTRO & BAEZA 1987: 21

Host, Corvina sp. Pagellus sp.

probably Aplodinotus grunniens, Croaker.

New Orleans, La.

Anchorella appendiculata Kröyer ♀
^{hemalopoda}

Anchorella —, Kröyer, 1863, p. 305; pl. 16, fig. 7a-d.

— —, Vogt, 1877b, p. 65; pl. 3, fig. 10.

— —, I. C. Thompson, 1895^b, p. 321

— —, B. Smith, 1899, p. 506.

— Thompson, 1895: 103

Host, Gills of lake —

Valparaiso (over)

H. Scott, 1904, b. p. 124; 1904, a. p. 44.

Transferred to genus Brachiella 1914.

NEOBRACHIELLA -, CASTRO + ROMERO 1987: 22

Anchorella brevicollis

--, Beneden & Bessels, 1868?, p. 43-

Anchorella canthari Heller, 1865 - see Alella canthari

Lernaeopodidae. Lernaeoidea.

Anchorella centrodonti T. J. v. Beneden, ♀.

— — Beneden, 1870, p. 43: pl. 2. fig. 5.

— — Wilson, 1915, p. 669.

A synonym of Clavella pagelli.
(Wilson, 1915, p. 669)

Anchorella clara Richiardi ♀ ^{Ternaeopodidae.}

Anchorella —, Richiardi, 1880, p. 152.

— —, Carus, 1885, p. 378.

Clavella —, Brian, 1906, p. 121.

— — Wilson, 1915, p. 668.

Host, Gills of Maena vulgaris.

Mediterranean.

Zernaeopodidae -
Anchorella crassa 'Richardi', ♀

— — Richardi, 1880, p. 152 nomen nudum.

A probable synonym of Clavella crassa.
(Wilson, 1915, p. 668).

ANCHORELLA DILATATA - SEE CLAVELLOTIS

Lernaeopoda.

Anchorella elongata Richiardi, ♀.

— — Richiardi, 1880, p. 152 nomen nudum.

A probable synonym of Clavella elongata
(Wilson, 1915, p. 668).

Lernaeopodidae. Lernaeoidea.

Anchorella emarginata Krøyer, ♂, ♀.

— — Krøyer, 1837, p. 287: pl. 3, fig. 7, a-e.

-- QUIDOR 1906:11

A synonym of Clavellisa emarginata.
(Wilson, 1915, p. 669)

Lernaeopodidae. Lernaeoidea

Anchorella gallax Heller, ♀.

— — Heller, 1865, p. 241; pl. 24, figs. 4-5.

A synonym of Clavelloopsis gallax.
(Wilson, 1915, p. 669)

-> TRANSFERRED TO CLAVELLOTIS, KABATA 1996:
2565.

Anchorella fasciculata Leidy, ♀.

— . Leidy, 1889, p. 138, 1 text figure.

A synonym of Brachiella fasciculata

Lernaeopodidae. — Lernaeopodoidea

Anchorella hostilis Heller, ♀

-- Heller, 1865, p. 243: pl. 24, f. 7.

A synonym of Brachiella hostilis
(Wilson, 1915, p. 669)

-- BEN HASSINE + RAIBAUT 1978: 169, =

BRACHIELLA HOSTILIS

NEOBACHIELLA --, CASTRO + BAEZA 1987: 22

Anchorella intermedia Quidor ♀♂

— —, Quidor, 1906, p. 29: pl. 3. figs. 37-44.

Transferred to new genus Clavellodes. 1914.
(Wilson, 1915, p. 669).

Host. Mouth cavity of a Nothostenia.

Antarctic Ocean.

Achorella lagenula Guérin ♀ ^{Veruatozoidae}

— —, Guérin, 1817, pl. 9, fig. 5.

— —, Enc. Brit. 7^o Edit. Vol. 24, pl. 502, fig. 14.

(Synonym of Clavella uncinata)
Wilson, 1915, p. 669.

Anchorella lichiae Richiardi, ♀.

— — Richiardi, 1880, p. 152, nomen nudum.

A probable synonym of Clavella lichiae.
Wilson, 1915, p. 668.

Anchorella lizae Kröyer ♀ ^{Ternaeopodidae.}

— —, Kröyer, 1863, p. 294: pl. 16, fig. 11 a-c.

— —, Vogt, 1877, p. 432.

Synonym for Naobranchia lizae.
(Wilson, 1915, p. 669).

Anchorella microcephala ^{Ternaeopodidae.} Nordmann ♀.

Anchorella —, Nordmann, 1832, p. 107.

— — Wilson, 1915, p. 669.

(never described or figured)

Host. Gills of a Sargus sp.

Anchorella ovalis Krøyer. ♀. Lernaeopodidae.

- —, Krøyer, 1837, p. 289: pl. 3, fig 6. a + b.
— —, Krøyer, 1863, p. 383.
— —, M. Edwards, 1840, p. 519.

Transferred to new genus Clavellisa. 1914.

Host. Gills of Trigla gurnardus.

--, Beneden & Bessels, 1868?, p. 43-

Lernaeopodoida

Anchorella pagelli Krøyer, ♀.

— — Krøyer, 1863, p. 295: pl. 16, f. 3, a-b.

A synonym of Clavella pagelli.
(Wilson, 1915, p. 672).

—, Kabata, 1972, p. 217, = Alella pagelli (Krøyer, 1833),
type of Alella.

Anchorella pagri Kroyer - see Clavellotis

ANCHORELLA PARADOXA - SEE ADUENA

Lernaeopodidae. Lernaeopodoidea.

Anchorella quadrata Bassett-Smith, ♀.

— — B-Smith, 1896. b, p. 15: pl. 4, §. 5.

A synonym of Clavella quadrata.
(Wilson, 1915, p. 670).

Lernaeopodidae. Lernaeoidea.

Anchorella rugosa Krøyer, ♂.♀.

— — Krøyer, 1837, p. 284: pl. 2, f. 7: pl. 3, f. 14, a-e.

Made type of new genus Clavellodes.
(Wilson, 1915, p. 690).

— Tempere, 1898: 225, pl. 29, fig 10

Lernaeopodidae.

Lernaeoidea

Anchorella sargi Kurz, ♀

— — Kurz, 1877, p. 393: pl. 25, fig. 5-6; pl. 26, fig. 29; pl. 27, fig. 51-52

A synonym of Clavellopsis sargi
(Wilson, 1915, p. 670).

-- TRANSFERRED TO CLAVELLOTH, KABATA
1936: 2565

Anchorella sciaenophila Heller, ♀ ^{Ternaeopodidae}

Anchorella —, Heller, 1865, p. 243: pl. 24, fig. 8.

— —. B-Smith, 1899, p. 506.

(See C. sciaenae).

Transferred to ~~the~~ genus Bachiella, 1914.

—, Pillai, 1968, p. 133

Host. Gills of Sciaena sp.

Mediterranean Indian Ocean.

Lernaeopodidae. Lernaeoidea,

Anchorella scombrici Kurz, ♀.

Fig. 41

— — Kurz, 1877, p. 403: pl. 25, fig. 12: pl. 26, fig. 35: pl. 27,

A synonym of Clavellisa scombrici
(Wilson, 1915, p. 670)

Lernaeopodidae. Lernaeopodoidea.
Anchorella stichaei Krøyer, ♀.

-- Krøyer, 1863, p. 372 pl. 16, fig. 1.

A synonym of Clavella stichaei,
--, Stephensen, 1940, a syn. of " "

1 p. 13

Anchorella subtilis Richiardi, 1880

-- Ben Hassine & Raibaut 1978: 173, nomen nudum

Lernaeopodidae.

Lernaeoidea.

Anchorella triglae (Claus), ♂, ♀.

Fig. 46-47.

— — Kurz, 1877, p. 404: pl. 26, fig. 13-15, 22-23; pl. 27,

A synonym of Brachiella triglae.
(Wilson, 1915, p. 670).

Lernaeopodidae. Lernaeopodoida.

Anchorella uncinata (Müller), ♀.

— — Nordmann, 1832, p. 102: pl. 8, f. 8-12.

— — Timm, 1894, p. 397.

A synonym of Clavella uncinata.
(Wilson, 1915, p. 680).

— — Tempere, 1898: 225, pl. 29, fig. 11

— — Thompson, 1893, p. 213: pl. 27, f. 2.

--, Beneden & Bessels, 1868?, p. 43-, pl. 5, figs. 1-7

--, Belloc —, : 249-250, fig. 42.

--, ACHOQUE, 1899: 205

Anchorella urolophi Kröyer ♀ Ternaeopodidae

- Anchorella —, Kröyer, 1863, p. 304; pl. 16, fig. 10, a-d
— —, Vogt, 1877b, p. 65.
— —, B. Smith, 1899, p. 506.
— —, Wilson, 1915, p. 670.

Host, Gills of Urolophus oerstedii.

Gulf of Mexico