

Caligodes, valid.

Caligoida
Caligidae.

Heller, 1865,

C. laciniatus.

This had been called Chondracanthus L. by Kollar, and Sciaenophilus L. by Kröyer, but Heller established it clearly as a new genus.


Caligodes Heller, 1865, p. 180

Caligodes = Caligus and εἶδος, likeness
or similarity.

Caligodes Wilson, 1905, p. 608.

Carapace orbicular, minute: Frontal plates with very small lunules: Fourth thoracic segment free, elongate: Genital segment flask-shaped, produced forwards as a ^{part of the} long neck, and backwards as two divergent processes, one on either side abdomen.

Abdomen itself large, broad, with minute anal laminae - First & fourth swimming legs uniramous, second & third biramous.

In other respects like Caligus.
First mx. small with swollen base: 2nd pair simple, Furca large & prominent & bent so as to appear like a grapnel anchor in side view.
Egg-cases long & narrow,  Eggs as in Caligus.

Caligodes Heller

—, Yamaguti, 1963, p. 63

— Dojiri, 1983: 133-136, vi

-- PRABHA 1983: 52-

Caligodes alatus Heegaard

_____ Heegaard 1945a, p. 5, figs. 16-28, Host: Cardus
sp. Java. ★ ♀

Caligodes carangis Bassett-Smith - see Parechetus

CALIGODES LACINIATUS - SEE CALIGAS

Caligodidae - Caligoida.

Caligodes lamarcki Quidor, ♀.

= carangis

— —, Quidor, 1913^b, p. 192 - 7 text figures.

Lamarckina caligusa Quidor, 1913^a, p. 1096.

Name substituted for Lamarckina caligusa.

Host. unknown.

Djibouti.

Caligodidae - Caligoida

Caligodes megacephalus, Wilson, ♀.

— —, Wilson, 1905, p. 609: pl. XVI.

— —, M. J. Rathbun, 1905, p. 91.

— —, Wilson, 1932, p. 409: fig. 256.

— —, Kabata, 1970, fig. 17

Host

Silver Gair (*Tylosurus marinus*
(*Lepisosteus osseus*)
mouth.

Woods Hole