

Canvella valid.

T. Scott, 1893 -

Harpacticorda - longipediidae
CANUELLIDAE

C. perplexa (M).

Syn. Canvella T. Scott, 1893, p. 92.

Canvella G. O. Sars, 1903, p. 16.

Longipedia (pars) Brady, not Claus.

Canvella Pesta, 1921, p. 561.

To Eugen Canu.

Body nearly cylindrical: head separate
from 1^o segt. Gen. segt. in ♀ with dorsal
transverse suture, dilated in ♂ with 2 acute
projections - 1^o antennae like those of Suu-
aristes. Exopod of 2^o antenna 7-jointed!
Endopod of 2^o legs normal: 5^o legs re-
dimentary - Ovipos double -

Canuella

~~Longipediidae~~-Harpacticoida
CANUELLIDAE

Sewell 1940, p. 131, review, new subgenera

—, Noodt, 1971, p. 99

—, Monchenko 1986: 337

—, Vincx + Heip 1979: 296

— IZAWA 1987: 165—

— BOXSHALL 1986 (IN KABATA-ED.): 201

— KRISHNASWAMY 1957 d: 9

— TIEMANN 1984: 56—

— MIELKE 1994: 55—

Canuella T. Scott, sensu restricto Por, 1984

- Por 1984:14, diagnosis, species list

- GOTTO 1993:207- , DIAGNOSIS

Canuella brevifurca Wells

— Wells, 1967: 214-216, fig 13.

Canuella bulbifera Chislenko, 1971

Chislenko 1971: 151-157, Figs. 1-4, n. sp. female + male,

Possjet Bay, Sea of Japan

Canuella canadensis Willey - see Coullana

INTERSUNARISTES CURTICAUDATA - SEE

INTERSUNARISTES CURTICAUDATUS

SEE ALSO C. SCOTTI

Canuella curticaudata sensu A. Scott 1909

-- A. Scott 1909: 197, pl. 64, figs. 1-6.

-- Fiers 1982: 11, C. curticaudata sensu Scott 1909 is
a Scottolana.

-- UMMERKUTTY 1966d: 306

-- POR 1984: 16, CANNOT BE ASSIGNED TO GENUS.

Canuella elanítica Por - see Elanella

Longipediidae - Harpacticoida

Canvella furcigera G. O. Sars, ♀, ♂, 1903

— — Sars, 1904, p. 18; pl. 10.

— — Sewell, 1924, p. 807; pl. 49, fig. 1

— — Wilson, 1932, p. 172; fig. 117; MASSACHUSETTS

— — Lang, 1936, p. 4; fig. 1.

— — Monard 1935a, p. 9, Roscoff, Atlantic coast France

Habitat, Norway Fjords; Chilka Lake.

— — Sewell 1940, p. 134, Nicobar Ids. ♀

— —, NOODT, 1956, p. 44, N. Sea.

— —, LANG, 1948, p. 164, Pl. 102:2

- - Jespersen, 1940, p. 71-72. Iceland.
- - --, Wells, 1963. p. 11
- - --, Por, 1960, pg. 102, table 2 fig. 1
- -, Stock & Vos, 1960, p. 207
- -, Dolzonolskaya, 1970, p. 59, pl. VI, fig. 3
- -, Jakobi, 1959, p. 141
- -, Klie, 1941, p. 4
- -, MARCUS, 1965: 91
- -, BODIN 1968: p. 283.
- - DINET 1971: 748, 758 - MARSEILLES
- Willey, 1923: 33, CHRISTIANA FJORD, NORWAY

Canuella furcigera Sars

card 2

- Glatzel, 1988:105-112 pl I figs 5-8, pl II figs 13-16, pl III fig 18 Wadden Sea
- KRISHNAJWAMY 1957d: 9-10, PORTO NOVO, INDIA;
11-12
- POR 1984: 14
- GOMOLU 1971: 157, BLACK SEA
- VOROB'YEVA & KULAKOV 1990: 14, SEA OF AZOV
- BODIN 1964: 119, GULF OF MARSEILLES
- POR 1962: 11, 17, 27, 31, 33, MEDITERRANEAN COAST OF ISRAEL, & BLACK SEA

Canuella (Canuella) indica Krishnaswamy, 1957

--- Krishnaswamy 1957d:10-12, Text-Fig. 1, n. sp., ♀
only, Madras, India (unpublished thesis)

- SEE INDICANUELLA

CANUELLA (INDICANUELLA) -, MIELKE 1984: 63

CANVELLA INOPINATA (THOMPSON & SCOTT)

— —, LANE, 1948, P. 165, PL. 103:1.

Scottolana inopinata. — POR, 1967, p. 105.

CANVELLA —, POR 1984: 16, CANNOT YET BE ASSIGNED TO
GENUS.

SCOTTOLANA —, POR 1977: 88

SCOTTOLANA —, HUYS 1995: 237, 238, TAB. 1

C. —, POR 1962: 27, 42

CANUELLA (ELLUCANA) LONGICAUDA SEWELL

— — SEWELL, 1940, P.P. 136-139, FIGS. 3 A-J.

HAB: NICOBAR IS. (WEED WASHINGS)

~~Canuella longicauda Sewell 1940, p. 136, fig. 3, Nic~~

Canuella longicauda Sewell 1940, p. 136, fig. 3,
Nicobar Ids.

--, = ELLUCANA L., COULL, 1971: 203

Canuella longipes (Thompson & Scott) - see Scottolana

Canuella paenelantica Fiers 1982

-- Fiers 1982: 1, 16-19, Pl. X, figs. 1-7; Pl. XI, figs. 1-4; Pl. XII, figs. 1-5; n. sp. ♀♂, Papua New Guinea.

-- Wells, 1985: 2

-- MIELKE 1994: 61

Longipediidae - Harpacticoida.

Canuella perplexa T. & A. Scott, ♀.

— T. & A. Scott, 1893, p. 92: pl. 2, f. 1-3.

— T. Scott, 1893, p. 201: pl. 2, f. 21-35.

— Sars, 1904, p. 17: pls. 8, & 9.

— T. Scott, 1906, p. 306.

— Thompson, 1895: 100.

Habitat. British Seas; Indian Ocean.

— Wilkey, 1923: 312-313

— — Thompson & Scott, 1903, p. 257,
Longipedia coronata ♀ Brady (not Claus), 1880,
vol. 2, p. 6; pls. 34 & 35, fig. 1-9, ♀.

Canuella perplexa Pesta, 1921, p. 561: text f. H3.

— — Klie, 1913, p. 8

— — Gurney, 1927f, p. 487; fig. 125, A-B.

— — Monard, 1928, f. p. 289: fig. 2.

— — Klie, 1927, p. 3:

— — Monard 1935 a, p. 8, Roscoff, Atlantic coast

— —, ^{France}NOODT, 1956, p. 44.

— —, BORUTSKY, 1952, p. 82-83 Fig. 289 ^{gussie}

— —, LANG, 1948, p. 163, (p. 1, 102:1)

— —, Dussart, in Aguessé + Dussart, 1957, p. 492. Camargue.

— —, Petkovski, 1964, p. 4

Canuella perplexa-2

-- --, Por, 1960, pg. 102, table 2, fig. 2

- Noodt, 1952:107

-- --, Bodin, 1972, p. 845, 859

--, Stock & Vos, 1960, p. 207

--, Pesta, 1932, p. 10-11, fig. 3

--, Dolzonolskaya, 1970, p. 59, pl. VI, fig. 2

--, Vervoort, 1964, p. 61

--, Jakobi, 1959, p. 142

--, MARCUS, 1965: 91

--, BODIN 1970: p. 391, Pl. 3, TAB. B.

--, BODIN 1968: pp. 279, 283.

--, BODIN 1971: pp. 416, 428-Tab:

- , BODIN 1972: pp. 657, 675; Tabs. 2, 3, 6.
- , BOZIC 1965: 352, 355.
- Mielke, 1975: 17, fig. 5
- CECCHERELLI + ROSSIN 1979: 99
- KONDALARAO 1984: 82.
- ONBE' 1984: 630
- Dussart, 1982: 98, pl. 28-A
- Michailova-Neikova, 1968: 211
- Mielke, 1975: 17-22, fig. 6-8
- Damian-Georgescu, 1970: 52-53, fig. 14a-f.
- VINCE + HEIP 1979: 281-299, FIGS. 1-9, DEVELOPMENT
- Serban, 1959: 260-261

Canuella perplexa T & A. Scott, 1893

card 3

- Raibaut 1967: 559 -, Thau Basin, France
- CARLI + CRISAFI 1983: 87, PL. 18 FIG. 1, ITALY
- PERPLEX, IZAWA 1987: 194
- HEIP + HERMAN 1986: 630-631, BELGIUM
- Por e' Marcus, 1972: 259 N. Red Sea (Suez Canal)
- Dahms, 1988: 505 - fig 2A Helgoland
- Glatzel, 1988: 105-112 pl I figs 1-7, pl II figs 9-12
pl. III fig 17 Wadden Sea, North Sea
- Huys 1988: 17
- CECCHERELLI + CORTESI 1985: 243-245, LAGOON, ITALY

Canuella perplexa T. & A. Scott, 1893

card 4

- Ceccherelli & Rossin 1979:99-, lagoons, Italy
- CARRADA ET AL. 1987:155-, LAGOONS, ITALY
- BODIN 1977:335, ÎLE DE RÉ, FRANCE
- DAHMS 1988b:14-, FIGS. 5,6, HELGOLAND-NORTH SEA
- POR 1984:6-, FIG. 16
- GOMOIK 1971:157-, BLACK SEA
- VOROB'YEVA & KULAKOV 1990:14-, SEA OF AZOV
- DAHMS 1991:88-, FIGS. 2B, HELGOLAND-NORTH SEA
- GOTTO 1993:207-, FIGS. 526-I

Canuella perplexa T. & A. Scott, 1893

card 5

- Bodin 1964: 119-, Gulf of Marseilles
- POR 1973: 91-, SINAI (MEDITERRANEAN)
- BOXSHALL 1990: 211-, FIGS. 1, 2
- DAHMS 1992a: 231-, TABLE 1
- CHAMPEAU 1966: 1289
- HUYS 1995: 237, 238, TAB. 1, DUTCH WADDENSEA
- A. SCOTT 1896: 134
- ITO, TAT. 1985: 153, 156, 157, FIG. 1F
- POR 1962: 11, 16-17, 27, 32, MEDITERRANEAN COAST OF ISRAEL

CANUELLA PERPLEXA T. + A. SCOTT, 1893

(6)

-- DÜRBAUM 1997: 213

-- HOSFELD + SCHMINKE 1997: 13-19, FIGS. 3, 4, 9, 10,
NORTH SEA

Canuella pontica Apostolov

-- Por 1984:14

Canuella reichi Por - see Nathaniella

Canuella scotti Sewell

Canuella scotti, Sewell 1940, p. 136, fig. 2, ♀,
Nicobar Ids., new name for C. curticauda, A.
Scott 1909

--, Hamond, 1972, p. 178.

- (CANUELLA)-, UMMERKUTTY 1966d: 305-308, FIGS. 4-6,
S E INDIA COAST

- - POR 1984: 14, POSSIBLY A. SCOTTOLANA

CANUEHA TUBA - SEE CANUELLINA

CANUELLA SP. CARVALHO, 1939

-- CARVALHO, J.P. 1939: 39, 40, FIG. 8a, b, SÃO PAULO,
BRAZIL.

Fam. CANUELLIDAE nov.

Harpacticoida

Lang 1948, p. 160

- MOESCHLER & ROUCH 1988: 12-
- POR 1984: 1-24
- DAHMS 1991: 87-, COMPARATIVE MORPHOLOGY
- FERRARI 1992a: 130
- GOTTO 1993: 207, DIAGNOSIS, KEY TO SOME GENERA
- MIELKE 1994: 61-62
- HUYS 1995: 225-, INTERSUNARISTES N.G.; ECHINOSUNARISTES N.G.; KEY TO GENERA

CANUELLIDAE

- BOXSHALL & HUYS 1989: 133, FIG. 3a - HOST OF
TANTULOCARID