

12
Cecrops, valid.

Leach, 1816,

Caligoida.
Cecropidae Caligidae.

C. latreillii (M)

Syn. Cecrops, Leach, 1816, p. 405.

Cecrops, M-Edwards, 1840, p. 472.

Cecrops Baird, 1850, p. 289.

Cecrops = The fabulous first king of Athens.

Cecrops T. & A. Scott, 1913, p. 97.

Cecrops Wilson, 1908, p. 466.

not

Cecrops - H.übner - Lepidoptera - 1816,

Carapace oval, stout, deeply notched anteriorly.
Frontal plates indistinct, fused with v carapace.
First antennae small, hidden - First thorax
segt only fused with head, others free: 4th segt.
with a pair of short dorsal plates.

Gen. segt. as long or longer than carapace -
2nd + 3rd thorax segments fused inter se, with no
dorsal or lateral plates. Abdomen small; anal
lamina very small.

All v thorax legs biramous, increasing in size
backward: terminal joints with short, hook-like
setae - Egg tubes very long, convoluted and com-
pletely hidden.

CECROPIDAE - SIPHONOSTOMATOIDA

CECROPS LEACH, 1816

2

- KABATA 1988: 31, 33

- KABATA 1988b: 3, ORIGIN OF NAME

- KABATA 1992: 33-, DIAGNOSIS

Caligoida - Caligidae.

Cecrops achantii-vulgaris Hesse, ♀.

— Hesse, 1883, p. 26: pl. 5, figs. 15-20.

— acanthiae — B-Smith, 1903, p. 108.

Host. Body of Acanthis vulgaris

French Coast. (over)

Hesse evidently transposed the "h" in the specific name, and his genitive is wrong also. B. Smith rearranged the "h" correctly, but his genitive is spelled incorrectly = acanthiuv is the correct spelling.

Cecrops desmaresti Risso, ♀. 1826. Caligidae.

— — Risso, 1826: vol. 5: 138

— — Cuvier, 1829, p. 363.

Host. At the surface far from shore.

Mediterranean.

Cecropinae Caligoida
Cecrops exiguus, n. sp.

— — Wilson, 1923, p. 1: pl. unnn., figs. 1-15.
-- KENSLEY & GRINDLEY 1973: 90, FIG. 14a, b, ♀[♂]
FROM MOLA LANCEOLATA, BANTRY BAY, SOUTH AFRICA;
p. 123.

Host. Gills of large shark, Florida.

Cecropidae: Siphonostomatoida

Cecrops exiguus Wilson

(Leach 1816?)

-- --, Shiino, 1965, p. 381-390,

CECROPS FULGENS (TELESÍUS, 1819)

ONISCUS FULGENS TELESÍUS 1814-1819: 10, pl.
213, FIG. 24

CARCINIUM FULGENS, MEYER 1832: 154, N. COMB; EX
ONISCUS.

Cecrops latreillii, Leach, ♂. ♀.

Caligidae

- , Leach, 1816, p. 405, pl. XX, figs. 1-5.
—, Lamark, 1818, p. 138; Leach, 1819, p. 534.
—, Guérin, 1817, pl. XXXV, fig. 8.
—, Audouin, 1823, p. 297.

Host. Gills of Orthogoriscus mola; Mola aspera
Thynnus vulgaris (Heller); Oblong Diodon
(white); Pleuro- nectus maximus and
P. rhombus. N. Atlantic

- — —, Desmarest, 1825, p. 338; pl. 20, fig. 271a-e.
 — — —, Nordmann, 1832, p. 39, [1829, p. 199, vol. 4
 — — —, Latreille, 1835, pl. 335, figs. 3-9. ^
 — — —, M-Edwards, 1840, p. 474. ^{1849, pl. LXXXVIII} fig. 4.
 — — —, Baird, 1850, p. 293; pl. XXXIV, fig. 1-2.
 — — —, White, 1850, p. 123; 1857, p. 321.
 — — —, van Beneden, 1855, p. 523; pl. n.n. figs. 1-14.
 — — —, — — —, 1861, p. 149; Bosc, 1830, vol. 2, p. 221.
 — — —, Kröyer, 1863, p. 190; Tamarck, 1838, p. 206.
 — — —, Heller, 1866, p. 754 (32).
 — — —, Olsson, 1869, p. 22.

Cecrops latreilli (2)

Caligidae.

- , van Beneden, 1870, p. 86.
—, Rathbun, 1884, p. 489.
—, Carus, 1885, p. 363.
—, Lucas, 1887, p. 31. (xxx1).
—, Hesse, 1888, p. 340; pl. XIV, figs. 1-22.
—, Thomson, 1889, p. 362; pl. xxvi, fig. 3a-f.
—, A. Scott, 1892, p. 266.
—, Brian, 1898, p. 14.
—, Brian, 1899, p. 200; 1908, p. 4.
—, Oken, Alg. Naturgesch. Zool. Bd. II 2, p. 626; taf. XX, fig. 3.

—, B. Smith, 1899, p. 465: 1896, a, p. 159.

—, T. Scott, 1900, p. 154.

—, —, 1901, p. 126.

—, Bose, 1830, p. 221. — Hoerer, 1857, 21 pp,

—, Hoerer, 1857, p. 67. — pls. III & IV.
(original paper).

—, Richiardi, 1880, p. 3.

—, Valle, 1880, p. 60. — Stussich, 1880, p. 258.

—, Cuvier, 1849, p. 199. — Brian, 1908, p. 4.

—, Brian, 1905, p. 6. — Stebbing, 1910, p. 558.

—, Latreille, 1816, p. 566.

—, " " , 1817, p. 65.

—, Teach, 1819, p. 534.

- —. M. J. Rathbun. 1905, p. 96.
— —. Wilson. 1907, p. 468: pls. 38 & 39.
— —. Brian, 1912, p. 13: Wilson, 1908, p. 453
— —. Pearson, 1906, p. 27.
— —. Fowler, 1912, p. 107: pl. 27.
— —. Elwes, 1909, p. 20.
— —. Latreille, 1817, III, p. 65: 1829, IV, p. 199.
— —. Leidy, 1890, p. 230.
— —. Norman + Brady, 1909, p. 155.
— —. T. Scott, 1904, b, p. 369.

Cecrops latreilli

- Hewitt & Hine, 1972, pg. 90
- , ACLOQUE, 1899: 200-201, pl. 95 (as CECROPS LATREILLE)
- KENSLEY & GRINDLEY 1973: 91, 123
- KABATA 1988: 33, FIG. 5. 14D-F; MOLA MOLA
- ZWERNER & LAWLER 1972: 89, CHESAPEAKE BAY,
HOST MOLA MOLA
- KABATA 1992: 33; FIG. 5 C-G, SYNONYMY, HOST LIST
- ~~KABATA~~

Cecrops latreillei

--, Boer, 1970-2, tbl. 1

--, Threlfall, 1967, p. 171, tbl. 2

— Tempere, 1898: 222, pl. 29, fig 6, 7