

in) Lernaepodoida - Drecheleskidae
Chondrocarpus. valid.
Bassett-Smith, 1903 - C. reticulosus^①.

Syn. Chondrocarpus B-Smith, 1903, p. 105.

- Hays, 2001: 136, INCERTAE SEDIS IN PHILOBLENNIDAE

Χόνδρος; cartilage & καρπός, wrist.

Female, Cephalothorax coriaceous, elongated, with
4 lateral short lobe-like processes on either
side; abdomen one-fourth length of whole, bi-
articulate, tapering; no distinct antennae or
thoracic limbs: mouth placed on under sur-
face, with minute maxillae: external ovaries
as elongated filiform sacs containing
ova in a single series -

Male, a Pigny.

(from B. Smith, 1903, pp. 104 & 5).

Chondrocanthiidae

Chondrocarpus reticulosus, F. S. B-Smilk.

— — —, B-Smilk, 1903, p. 105, Text-fig. 11, A-G.

— — — Monod & Dollfus, 1932, p. 174.

— — — JENSEN 1987: 76—

Host. Kidneys of Pleurobranchids.

Zanzibar.

Chondrocarpus sp. ♀ B-Smith.

— — B-Smith, 1903, p. 106; text fig. 11-H.

— — Monod & Dollfus, 1932, p. 174.

Host. Kidneys of a Pleurobranchid.

Zanzibar.