

Herpyllobiidae Lernaeopodida

Crypsidomus = synonym -

Levinson, 1878.

C. terebellae<sup>M.</sup>

not Crypsidomus - Günther - Reptile - 1864.

not Crypsidromus - Ausserer - Arachnids - 1871.

Crypsidomus Levinson, 1878, p. 375.

Changed to Aphanodomus Wilson.

Kρυψίδομος = dwelling in secret places.

Body of ♀ semilunar, not jointed, very soft and with no rudiments of appendages, the central portion fastened to the intestine of the host, the lateral portions, diminishing in size, adjacent to the intestine. Egg sacks filiform, longer than the body, perforating the skin of the host and hanging free on its back.  
Where it is fastened to the intestine of the host there is an aperture, but there is no anus.

Herpyllobiidae - Monstrillidae.

Cryptidomus terebellae Levinson. ♀.

— . Levinson, 1877, p. 375: pl. 6, figs. 19-20.

— . Stephensen, 1913, p. 343.

— . Stephensen, 1916, p. 305.

Host. Terebella cirata or Thelepus circumatus  
(Vernes)

Greenland.

Changed to Aphanodomus terebellae.

— Wilson, 1920, p. 95.

Cryptidomus — Hansen, 1923, p. 83.