

Notodelphyidae: Cyclopoida

Doroixys Kerschner, 1879

- Stock 1967e: 30; only one species, D. uncinata K.
- GOTTO 1993: 42, DIAGNOSIS

in) Notodelphyoidea — Doropygidae.
Doroixys, Kerschner.
Kerschner, 1879 — D. uncinata ^(M).

Syn. Doroixys Kerschner, 1879, p.
Doroixys Schellenberg, 1921, b, p. 254.

Δορός, a bag and ἱξύς, small of the back.

Body bent, convex in thorax - abdomen 4-jointed
short, cylindrical: incubatory chamber covering v
4° + 5° thorax somites, convex dorsally = 1° ant, short,
7-8 jointed, setiferous: 2° ant, 3-jointed, short,
solid, prehensile, without setae on 1° joint.
Mds, biramous, masticatory, rami subequal.
armed with setae: 1° mx, masticatory at base.
vibratile at tip - with sessile epipod: 2° mx. seg
4-joints, with prehensile spine on 2° joint: 1° joint
with large tubercle like that in Doropygus: 2°
mx endop, much reduced - 4 thorax legs bira-
mous - rami 3-jointed, except 2-jointed endopod
of 4° pair, converted in 2°-4° pairs into creeping
organs: 5° legs in form of a claw. anal laminae
with small hairs but without setae.

Ascidicolidae - Notodelphyoidea.

Doroixys uncinata Kerschner, ♀.

— — Kerschner, 1879, p.

— — Gournet, 1888, p. 2.

— — T. Scott, 1907, p. 366.

— — Carus, 1885, p. 343. Mediterranean.

Host. Diazona violacea: Aplidium cris-
tallinum: A. gibbulosum: Perophora
listeri. Morche llium argus M-Edwards,
Circinalium concrescens Giard.

-- Schellenberg, 1921³, p. 254.

-- --, Illg & Dudley, 1961. pp. 72-73

--, ~~Kabata, 197~~

--, Gotto, 1966, p. 192

-- Stock 1967e: 30

Doroixys uncinata Kerschner, 1879

card 2

--, Ooishi, 1972, p. 305-311, figs. 1-3; pl XIV, 2

-- IZAWA 1987: 155 -

-- OOISHI 1987: 95, JAPAN

-- GOTTO 1993: 42, FIG. 8A, B

? DOROIXYS SP.

-- Ooishi 1987: 96, JAPAN, ASCIDIAN