

Echinolaophonte Nicholls

Nicholls 1941b, p. 95. Genotype-E. (Laophonte) horrida
(Norman).

- Lang, 1965: 508-509 key
- FIERs 1988: 76
- FIERs 1992: 400
- FIERs 1991c: 287 -
- SCHIZAS + SHIRLEY 1994: 132-, cf. ELAPHOLAOPHONTE N.G.

Echinolaophonte armiger-group, Cottarelli, Forniz &

Bascherini, 1992

--- Cottarelli et al. 1992: 298, 3 taxa

Echinolaophonte horrida-group, Cottarelli, Forniz &

Bascherini, 1992.

--- Cottarelli et al. 1992: 298, 6 species

= ECHINOLAOPHONTE Laophontidae - Harpacticoida

Laophonte armiger Gurney, ♀. /

— — Gurney, 1927f, p. 554: fig. 159, A-E.

— — Willey, 1932, p. 107: figs. 65-67.

Echinolaophonte ____, Nicholls 1941, p. 95

Habitat. Suez Canal: Bermuda

— —, Carvalho, 1952, p. 159-160, pl. 2, fig. 68-71. Baía de Santos, Brazil.

--, Vervoort, 1964, p. 357, 359, 362

= ECHINOLAOPHONTE

Onychocamptus armiger (Gurney, 1927)

--, Vervoort, 1964, p. 5, 358, 359-364, figs. 142, 143, 144, 146c-e - IFALUK ATOLL

ONYCHOCAMPTUS -, REID 1990g: 200

ECHINOLAOPHONTE -, FIEBS 1991c: 289

E. ARMIGER TYPICA, COTTARELLI, FORNIZ ET AL. 1992:

289-, NAPLES, ITALY, ~~IN~~ TYPE OF "ARMIGER

GROUP" NOV.

O. -, POR 1962: 12, 14, 24, MEDITERRANEAN COAST OF ISRAEL

Echinolaophonte armiger (Gurney)

Laophonte armiger Gurney 1927, p. 554, fig. 159, a-e
♀, Suez Canal

_____, Willey, 1932, p. 107, figs. 65-67, Bermuda

Echinolaophonte _____ Nicholls 1941, p. 95

_____ Nicholls 1945a, p. 11, fig. 7, Western Australia

--, WELLS, 1979: 5

- Knatz, 1986: passim

-- KRISHNASWAMY 1937d: 64-65, FIG. 13, RECORDS, MADRAS
typica.

-- COTTARELLI & FORNIZ 1931: 665, SPARGI X DEI CAVOLI
ISLETS, NEAR SARDINIA

Echinolaophonte armiger (Gurney)

card 4

Onychocamptus -, Pesta 1959: 160-, fig. 147, off Naples,
Italy

E. -, WILLEN 1996: 1322, TAB. 1

Echinolaofonte armingier f. briani

—Lang, 1965: 509. — SIS, fig. 280-283,

N. FORMA, CALIFORNIA

--- COTTARELLI, FORNIZ ET AL. 1992: 298, IN
"ARMIGER-GROUP" NOV.

3 ECHINOLAOPHONTE Laophontidae - Harpacticoida.

Laophonte brevispinosa Sars, ♂. ♀. 1

L. — Sars, 1908, p. 247; pl. 168.

L., Vervoort, 1964, p. 357, 359

Habitat. Norway.

Echinolaophonte brevispinosa (Sars) 1908

2

--, Montagna & Carey, 1978, p. 119, BEAUFORT SEA

- COTTARELLI + FORNIZ 1991: 674

- COTTARELLI, FORNIZ ET AL. 1992: 298-, IN HORRIDA-
GROUP

Echinolaophonte gladiator (Vervoort, 1964)

-- Cottarelli & Forniz 1991: 674

-- COTTARELLI, FORNIZ ET AL. 1992: 297.

^{Laophontidae — Harpacticoida}
Laophonte horrida (Norman), ♀. ♂. 1

Cleta minuticornis Buchholz, 1874, p. 393: pl. 15, f. 3.
— horrida Norman, 1876, p. 206.

Laophonte — Brady, 1880, p. 74: pl. 74, figs. 1-11.

— — Sars, 1908, p. 246: pls. 166 & 167,
—, Lang, 1944, p. 38, a syn. of Onychocamptus h.

A synonym of Laophonte minuticornis.

Stephensen, 1913, p. 333.

— Scott & Scott, 1901: 345

Arctic Seas.

- — T. Scott, 1906, b, p. 335; 1903, p. 17.
— — T. Scott, 1899c, p. 101.
— — Brian, 1921, p. 94
— — Thompson, 1893, p. 199: pl. 20, f. 7, a-d.
— — Klie, 1927, p. 15:
— — Wilson, 1932, p. 279: fig. 173.
— — Lang, 1936, p. 34-35:
— — Nicholls 1939 b & 1940, p. 302, ST. LAWRENCE RIVER
— — Jespersen 1940, p. 86. Iceland.
— —, Pesta, 1932, p. 133-4, fig. 61
— —, Klie, 1941, p. 89
— —, Klie, 1941, p. 35
— Willey, 1923: 330

Onychocamptus horridus

= ECHINOLAOPHONTE

2

- , Jakobi, 1959, p. 139
- , Klie, 1950, p. 126
- , Lang, 1944, p. 38, Laophonte h. is a syn.
- , CHISLENKO, 1977: TAB. 1
- , ROE, 1958: 233
- , Becker, Von, K., 1970, p. 59
- HORRIDA, CITARELLA 1989: 136, ST. LAWRENCE GULF,
CANADA
- HORRIDA, SHIH ET AL. 1971: 211, CANADA - ARCTIC
- HORRIDA, CITARELLA 1984: 139, SHÉDIAC BAY, NEW
BRUNSWICK, CANADA

ECHINOLAOPHONTE HORRIDA (Norman, 1876) 3

- REID 1990g: 199
- , ITO, 1969: 72
- , Coffin, 1978-79, p. 593
- Fiers, 1986c: 146-147
- KRISHNASWAMY 1957d: 63, MADRAS, INDIA
- KASK ET AL. 1982: 37, NANAIMO, BRITISH COLUMBIA
- COTTARELLI + FORNIZ 1991: 674

ONYCHOCAMPTUS HORRIDUS, CHISHENKO 1977b: 269,
FIG. 17(5), FRANZ JOSEF LAND

Echinolaophonte horrida (Norman, 1876)

card 4

-- Cottarelli, Forniz & Bascherini 1992: 298, type of
"horrida-group" novo.

LAOPHONTE HORRIDA, NICHOLLS 1940: 244-, ST. LAWRENCE
RIVER, CANADA (MARINE)

L. HORRIDA, C.B. WILSON 1936: 372, CANADIAN ARCTIC

L. -, MCINTYRE 1969: 278, KIEL BAY, GERMANY

Laophonte horrida trispinosa

---, Vervoort, 1964, p. 357, 359

Echinolaophonte minuta Cottarelli & Forniz, 1991

-- Cottarelli & Forniz 1991: 666-676, Figs. 1-3, n.sp.

female + male, Li Nibani 2 islet, near Sardinia

ECHINOLAOPHONTE sp., COTTARELLI, FORNIZ ET AL. 1992:

297, 298

ECHINOCAOPHONTE MIRABILIS (GURNEY, 1927)

- , WELLS, 1979: 5
- FIERs 1988: 76, INDIAN OCEAN, PAPUA NEW GUINEA
- COTTARELLI + FORNIZ 1991: 674
- FIERs 1991c: 289
- MIRABILIS, COTTARELLI, FORNIZ ET AL. 1992: 297-

ECHINOCLAOPHORITE OSHOROENSIS ITO, 1969

- ITO, 1969: 68-73 ♀♂; FIG. 6-9
- KITO 1977: 692-4
- COTTARELLI & FORNIZ 1991: 674
- COTTARELLI, FORNIZ ET AL. 1992: 298, IN
HORRIDA-GROUP

Echinolaophonte tetracheir n. sp.

--, Mielke, 1981, p. 53, figs. 29-34

- COTTARELLI & FORNIZ 1991: 674-675

- COTTARELLI, FORNIZ ET AL. 1992: 297

Echinolaophonte tropica Ummerkutty, 1970

-- Cottarelli & Forniz 1991: 674

-- FIEBS 1991c: 289-

-- COTTARELLI, FORNIZ ET AL. 1992: 298, IN HORRIDA-
GROUP

Echinolaophone[†] veniliae Cottarelli, Forniz & Bascherini,
1992

-- Cottarelli et al. 1992: 289-, figs. 4-7, n. sp. male +
female, Cetara seashore, Ischia, Naples, Italy; *IN*

HORRIDA-GROUP

Echinolaophonte sp.

-- Masunari 1985: 54.

-- COTTARELLI, FORNIZ ET AL. 1992: 297, 298; = F.

MINUTA