

Lernaeomyzon = synonym.  
Blainville, 1822.

Lernaeopoda  
Lernaeopodidae.

L. uncinata.

A synonym for Clavella (Oken, 1815)

Δέρνη and μύζειν = to suck.

Lernaeopodidae: Siphonostomatoida

Lernaeomyzon Blainville 1822

Blainville, 1822: 439, fig. 9.9a

Lernaeomyzon canthari (Heller, 1865) — SEE ALELLA 91

~~see~~ Anchorella canthari Heller, 1865, Kabata, 1972, p. 216

Lernaeomyzon elongata, Blainville ♀

—, Blainville, 1822, p. 439; pl. III., figs 9, 9a.

(This is probably Brachiella thynni)

Attached to v cellular masses containing  
v intestinal worms ("vers") in a "Cheilo-  
diptère-aigle" at Håvre.

Lerneomyzon incisa Blainville, ♀.

— — Blainville, 1822, p. 447: pl. n.n. fig. 10.

Host. unknown.

(over)

Blainville does not mention this species in his text, but in the Explanation of the figures he says -

"10. To. incisa, after a manuscript figure of Dr. Marion de Procé - although incomplete this figure indicates a new species related to the preceding and to that of Pernetty."

The fig. given is probably a Brachiella.

Lernaeomyzon pernettiana Blainville, ♀

"A Lernaeon" Pernetty, 1770, vol. 1, p. 93: pl. 1, f. 5+6.  
— — Blainville 1822, p. 439.

A synonym of Brachiella pernettiana

Lernaeopodidae.

Lernaeomyzon pinnarum, Blainville, ♀.

— —, Blainville, 1822, p. 438.

— —, Desmarest, 1825, p. 247. (footnote).

— —, M. Edwards, 1840, p. 520.

A synonym for Clavella uncinata.

See Fabricius, 1779, p. 282.



Blainville says - p. 439.

"I report this species, which I have never seen, in this section with some doubt; in fact Fabricius says <sup>organ</sup> that the ~~proboscis~~ <sup>which I make the sucker</sup> is dorsal, which is remarkable. He adds that it is attached to a fin with a whole anterior part body buried under the skin, which is unlike the true Ternatomyzon!"

Loercomyzon pyriformis, Blainville. ♀.

——. Blainville, 1822, p. 439; pl. un. fig. 8.

a synonym for Clavella uncinata.

LERNAEOMYZON UNCINATA - SEE CLAVELLA