

NOTODIAPTONUS KIEFER, 1936

- , DUSSART 1985, 208-214, ERECTED SUBGENERA
- , REID 1987a: 372, 378, "NORDESTINUS-GROUP"
- D., S. WRIGHT 1937: 76, "NORDESTINUS-GROUP"
- FERRARI 1991: 117-, DEVELOPMENT
- BOWMAN 1986: 297
- GAVIRIA 1989: 116, CR. COLOMBODIAPTONUS
- DIAPTONUS (N.), LIGHT 1939: 473 -

NOTODIAPTONUS (NOTODIAPTONUS) DUSSART 1985

- DUSSART 1985: 208-214, n. subgen. - N. DEITTERS',
TYPE-SPECIES
- REID 1987a: 372, 378

NOTODIAPTONUS (AMAZONIUS) DUSSART 1985

- DUSSART 1985: 208-214, N. SUBGEN.;
N. (A.) OHLEI, TYPE-SPECIES.
- REID 1987a: 378, NOT AVAILABLE

NOTODIAPTOMUS (CALEODIAPTOMUS) DUSSART 1985

-- DUSSART 1985: 208-214, N. SUGEN.;

N. (C.) CONIFEROIDES, TYPE-SPECIES

-- REID 1987a: 378, NOT AVAILABLE

NOTODIAPTONUS (WRIGHTIUS) DUSSART 1985

- DUSSART 1985: 208-214, N. SUBGEN.;
N. (W.) HIBBER
- REID 1987a: 378, NOT AVAILABLE
- REID 1989b: 682

[Noto-]

Diaptomidae - Calanoida

Diaptomus amazonicus Wright ♂♀.

— henseni - Wright, 1927, p. 96 : pl. 8. figs. 7-11

— amazonicus Wright, 1935, p. 221 : pl. 1. figs. 2, 5, 9, 14, 16.

-- Brehm, 1960, pp. 50.

-- WRIGHT 1936 : 80

D. -, S. WRIGHT 1937 : 78

Habitat. Northeastern Brazil.

D. -, S. WRIGHT 1938 : 562

NOTODIAPTONUS AMAZONICUS (WRIGHT, 1935) 2
~~, 1927.~~

--, ANDRADE & BRANDORFF, 1975: 97

--, BRANDORFF, 1976: 614

--, BRANDORFF, 1973: 205, 206

-- Brandorff & de Andrade, 1978: passim

— Hardy, 1980: 594—595, fig. 12

-- DUSSART 1984: 34, 35, 39, 48—49, FIG. 5, 63

-- DUSSART 1985: 208, IN NOTODIAPTONUS s. str.

— Matsumura-Tundisi, 1986: 537, fig. 22—25

Notodiaptomus amazonicus (Wright, 1935)

3

- Braendorff 1972: 25-, figs. 29-32, Amazon
- CARVALHO, M. L. 1981: 37-, LAGO GRANDE, AMAZONAS
- MONTEIRO 1980b: 57-, LAGOA DOS PATOS, BRAZIL
- MONTEIRO & LOEFDEN 1986: 83; FIG. 25, "
- SILVA ET AL. 1989: 727; FIGS 47-68, PARA', BRAZIL
- CICCHINO ET AL. 1989: 101
- BOZELLI 1992: 254-, LAKES, PARA', BRAZIL
- DUSSART & FRUTOS 1985(1986): 307
- SANTOS SILVA & ROBERTSON 1993: 101, RONDÔNIA, BRAZIL

NOTODIAPTONUS AMAZONICUS (S. WRIGHT, 1935) 4

-- LIMA ET AL. 1996: 114-115, MATO GROSSO DO SUL,
BRAZIL; AS "CF".

^{CF.} -- LANSAC-TÔHA ET AL. 1997: 140, 141, 143 TAB. 3, PARANA,
BRAZIL

Notodiaptomus amazonicus occidentalis

--Loeffler, 1963, p. 208 (fig 8a-b).
-- DUSSART 1984: 63

Notodiaptomus anceps Brehm, 1958

card 1

-- Brehm 1958: 139-147, fig. 1, n. sp.

-- Brehm 1960: 50, figs. 103-108

D. -, Brehm 1965: 7, Argentina

-- Brandorff 1972: 43-, distribution

D. -, Brandorff 1976: 618, fig. 3

-- Dussart 1984: 64, as N. anceps (?)

--?, Dussart & Robertson 1984: 391

--?, Dussart 1985: 210, in n. subgen. Wrightius

N.-, DUSSART & FRUTOS 1985(1986): 302

Diaptomidae - Calanoida

Diaptomus anisitsi Daday ♂♀.

1

150

— — Daday, 1905, p. 149: pl. 9, figs. 16-22.

— — Tollinger, 1911, p. 65

— — Wright, 1927, p. 77: pl. 1, figs. 4-6.

— — Kiefer, 1928, p. 172: fig. 2.

--, Pesta, 1927, p. 80

Habitat. Paraguay.

D. -, WRIGHT 1937: 76

- S. lat. -, BREHM 1958a: 166 - Key.

D. -, S. WRIGHT 1938: 562, 563, syn. D. INFLEXUS BRIAN

NOTODIAPTOMUS AWISITSI (DADAY) CARD 2

-- BRANDORFF 1972: 43-, DISTRIBUTION

-- BRANDORFF, 1976: 614, 622; FIG. 2; SYN:

DIAPTOMUS INFLEXUS BRIAN, 1925; BREHM, 1965;

D. AWISITSI WRIGHT, 1927; KIEFER, 1928; WRIGHT,

1938; BREHM, 1958; NOTODIAPTOMUS INFLEXUS

BREHM, 1958

--, BREHM 1958: 575-582; FIGS. 1-4; TAB. I,
COMPARISON w/ N. perelegans.

NOTODIAPTONUS ANISITSI (DADAY)

CARD 3

-- DUSSART 1984:63 (= INFLexus?)

-- DASSART 1985: 201, SWN. D. INFLexus BRIAN; 208, IN

NOTODIAPTONUS S. STR.

-- REID 1987a: 377

N. INFLexus, BREHM 1938: 29

D. ANISITSI, FORRO' 1986: 560

D. -, S. WRIGHT 1939: 647, ARGENTINA

V. -, PAGGI & PAGGI 1990: 690-, RIO PARANA' MÉDIO, ARGENTINA

V. -, DUSSART & FRUTOS 1985 (1986): 306

NOTODIAPTONUS ANISITSI (DADAY)

(4)

-- LOPES ET AL. 1997: 45-46, TAB. 1c, PARANA, BRAZIL

Diaptomidae - Calanoida -

Diaptomus inflexus Brian ♂♀.

— Brian, 1925, p. 180; text figs. 4-6

— — Brehm, 1965, p. 7, Argentina.

- s. lat. -, BREHM 1958a: 166 - Key.

Habitat. Río Uruguay, Argentina.

-- DESSART 1984: 63, syn. of Noro. ANISITSI (DADAY)

-- DESSART 1985: 201, syn. of N. ANISITSI

Noctodiaptomus BIDIGITATUS

- see D. BIDIGITATUS

-- DUSSART 1984: 64 - AS N. (BIDIGITATUS)?

-- Dussart 1985: 214, not assigned to subgenus

Notodiaptomus brandorffi Reid, 1987

-- Reid 1987a:364, 372-377, figs. 32-59, n. sp. ♀♂,
Brazil (northeast)

-- REID & TURNER 1988: 489-, MARANHÃO, BRAZIL
Notodiaptomus sp., BARBIERI ET AL. 1989: 406-;
MARANHÃO, BRAZIL

①

Notodiaptomus Caperatus Bowman

--, Bowman, 1979: 219-225, figs 2-4.

--, Dussart, 1982: 316-318 (= Pectenodiaptomus
caperatus)

- Bowman, 1986: 297

PECTENODIAPTONUS --, Dussart 1985: 214

-- Reid 1987a: 377-378, RETURNED TO NOTODIAPTONUS.

PECTENODIAPTONUS --, DASSART & FERNANDO 1986a: 41

N., Reid 1990g: 179, DISTRIBUTION

NOTODIAPTONUS CAPERATUS BOWMAN, 1979 CARD 2
-- BOWMAN 1990: 293

Noto-

Diaptomidae - Calanoida

Diaptomus carteri Towner ♂♀.

①

D. — Towner, 1934, p. 98: fig. 3, a-d.

D. —, WRIGHT 1937: 76

D. —, S. WRIGHT 1938: 562

Habitat. Brazil.

Notodiaptomus carteri (Lowndes)

(2)

- , Bowman, 1973, p.199
--, BRANDORFF, 1976: 614; FIG. 2
-- DUSSART 1984: 63
-- DUSSART 1985: 208, IN *NOTODIAPTOMUS* S. STR.
-- DUSSART 1984d: 264, FIG. 7C
-- REID 1987a: 377
-- MONTÚ 1980b: 57-, RG5, BRAZIL
-- MONTÚ + GLOEDEN 1986: 82-, FIG. 25, "
-- MATVEEV ET AL. 1992: 1-18, LAGUNA AEROCLUB,
CORRIENTES, N. ARGENTINA
-- MATVEEV 1991: 53-68, " " " "

Notodiaptomus carteri (Lowndes, 1934)

card 3

-- Dussart & Frutos 1985(1986): 306

Notodiaptomus cearensis (Wright)

1

--, Bowman, 1973, pp.193-199, 200, 201, figs.1-21

-- DUSSART 1984: 26-28, 34-36, 38, 39, 49, FIG. 6, 63

DIAPTONUS -, PEREIRA ET AL. 209-226, Pls. 32-33, as
intermediate host of nematode; L. TAUÁPE, FORTALEZA;
L. SORRÉ, L. PORANGABA - BRAZIL.

DIAPTONUS -, WRIGHT 1936: 80-81, Pl. I FIG. 2, BRAZIL

-- DUSSART 1985: 208, in *NOTODIAPTOMUS* S. STR.

— Matsumura-Tundisi, 1986: 542, fig. 61-66

Notodiaptomus cearensis (Wright, 1936)

card 2

-- Brandorff 1972: 44-, distribution

-- NOGUEIRA 1987: 77-85, FIGS. I, II - PERNAMBUCO,
BRAZIL

-- NEUMANN-LEITÃO & NOGUEIRA 1987: 87-118,
FIG. 12, PERNAMBUCO, BRAZIL

D., WRIGHT 1937: 76

D., WRIGHT 1938: 563

N., CICCHINO ET AL. 1989: 101

N., PARANAGUA, SILVA ET AL. 1990: 87, RIO CAPIBARIBE, PE, BRAZIL

Notodiaptomus cearensis (Wright, 1936)

card 3

-- Lima & Watanabe 1990:108, Paraiba, Brazil

- COSTA 1990: 112, BAHIA, BRAZIL

[Noto-]

Diaptomidae - Calanoidae

Diaptomus conifer G.O. Sars, ♂♀.

— — Sars, 1901, p. 13: pl. 3. figs. 1-8.

— — Lowndes, 1934, p. 93.

— — Daday, 1905, p. 147: pl. 9. fig. 10

— — Tollinger, 1911, p. 68.

D. —, WRIGHT 1937: 76

Habitat. Brazil. Paraguay. Venezuela.

--, BREHM 1949: 416, 420.

- s. lat. - , BREHM 1958a: 167 - Key.

- -- Pearse, 1921, p. 459.
- -- Wright, 1927, p. 91: pl. 6, figs. 10-12.
- -- Brehm, 1965, p. 5, Argentina. also p 8
- - , Bowman, 1973, p. 201
- , Pesta, 1927, p. 70, 76
- , Kiefer, 1956, p. 236. see *Notodiaptomus venezolanus*
- , Kiefer, 1979, p. 224-225
- - WRIGHT 1936: 79

Notodiaptomus conifer (SARS, 1901)

2

- , Sendacz & Kubo, 1982, p. 51, 66-68.
- De Gouyea, 1980: 1047-1059, fig 1-41.
- Ringuet, 1959: 314
- DUSSART 1984: 63
- DUSSART 1985: 208, IN Notodiaptomus s. str.
- DUSSART 1984d: 264, FIG. 7B
- REID 1987a: 372
- Matsumura - Tundisi, 1986: 542, fig. 4b-50
- D. -, BRANDORFF 1972: 47-48, DISTRIBUTION

Notodiaptomus conifer (Sars, 1901)

card 3

D. -, Forró 1986: 560

D. -, S. WRIGHT 1937: 66, 75-77, Pl. III FIGS. 1-4,
SÃO PAULO, BRAZIL

D. -, S. WRIGHT 1938: 562

N. -, DEFAYE + DUSSART 1988(89): 123

N. -, PAGGI + PAGGI 1990: 690; RIO PARANA' MÉDIO, ARGENTINA

N. -, DUSSART + FRUTOS 1985(1986): 306

N. CONIFERA, FERRARI 1993 b: 766, TAB. 1

[Noro-J]

Diaptomidae - Calanoida

Diaptomus coniferoides Stillman Wright. ♂♀.

— — Wright, 1927, p. 92: pl. 7, figs. 1-4.

— — Loawdes, 1934, p. 94: fig. 1, a-d.

— — Brehm, 1965, p. 7, Argentina, also p 8.

-- Brehm, 1960, p. 50

--, Brehm, 1957: 60-62; FIG. 72-76

Habitat. Brazil.

D., WRIGHT 1937: 77

--, BREHM 1958: 147-149; Fig. 2. = Diaptomus lobifer Pesta, 1927

- s. lat. - , BREHM 1958a: 167- Key.

D. -, S. WRIGHT 1938: 562, RIO PARANA' DELTA,
ARGENTINA

NOTODIAPTONUS CONIFEROIDES (WEIGHT, 1957) ②

--, ANDRADE & BRANDORFF, 1975: 97

--, BRANDORFF, 1976: 616; FIG. 2

--, BRANDORFF, 1973: 206

--, Herbst, 1967, p. 96

-- Brandorff, & de Andrade: 1978: passim

-- Hardy, 1980: 594-595, fig. 12.

-- DUSSART 1984: 34, 35, 38, 39, 54-55, FIG. 9, 63

-- DUSSART 1985: 201, NOT = *N. LOBIFER*

Notodiaptomus coniferoides (Wright, 1927)

card 3

-- Dussart 1985: 214, in subgen. Caleodiaptomus; ^{TYPE-SPECIES}

-- DUSSART 1984d: 264, FIG. 7A - NOT = N.
LOBIFER

-- REID 1987a: 372

— Matsumura-Tundisi, 1966: 542, fig. 51-54

D. —, BRANDORFF 1972: 24-, FIGS. 27, 28 - AMAZON; 98

—, CARVALHO, M.L., 1981: 37-, LAGO GRANDE, AMAZONAS

D. —, S. WRIGHT 1939: 647, ARGENTINA

N. —, REID & MORENO 1990: 726; PANTANAL, BRAZIL

Notodiaptomus coniferoides (Wright, 1927) card 4

-- Paggi & Jose de Paggi 1990:685-, Rio Parana Medio,
Argentina

-- DUSSART & FRUTOS 1985(1986): 306, 309, FIGS. 14-18

-- SANTOS SILVA & ROBERTSON 1993:101, RONDONIA, BRAZIL

= *NOTO DIAPTOMUS CONIFEROIDES*

Diaptomus lobifer Pesta, 1927

n. sp. Pesta, 1927, pp. 76-7, fig. a-e

--, BRANDORFF, 1976: 622

--, BREHM 1949: 416.

--, BREHM 1958: 147-149. = *Diaptomus coniferoides* Wright, 1927.

-♀. lat.-, BREHM 1958a: 167 - Key.

-- BRANDORFF 1972: 50, DISTRIBUTION

-- DUSSART 1984: 54, = *NOTO DIAPTOMUS CONIFEROIDES*

D.L., FORRO' 1986: 560

D. LOBIFER, S. WRIGHT 1938: 562

D. LOBIFER, S. WRIGHT 1939: 647, ARGENTINA

? NOTODIAPTONUS CORDEROI (WRIGHT)

-- DUSSART 1984:64: as N. (CORDEROI) (?)

Notodiaptomus corderoi

- Dussart 1985: 208, in Notodiaptomus s. str.
- DASSART 1984 : 64, as N. CORDEROI?
- REID 1987a: 365, 372 : = Scodiaptomus
CORDEROI.

NOTODIAPTONUS CORONATUS (SARS, 1901)

= TRICHOEDIAPTONUS BRANDORFF ~~CORONATUS~~ SARS

-- DUSSART 1984:34, 3963 (apparently moved to genus
Notodiaptomus without test; Fieckodiaptomus
Brandorff et al. 1983 has priority.)

NOTODIAPTONUS DAHLI (WRIGHT)
DIAPTONUS - , WRIGHT 1936: 79-80, Pl. I FIG. 1, N.S.P.-
BRAZIL

--, BRANDORFF, 1976 : 616 ; FIG. 2

-- DUSSART 1984: 64

-- DUSSART 1985: 208, IN NOTODIAPTONUS J. STR.

-- REID 1987 a: 377

D. -, ANDRADE + BRANDORFF 1975: 97

D. -, BRANDORFF 1972: 48

D. -, WRIGHT 1937: 76

D. -, S. WRIGHT 1938: 562

Notodiaptomus deeveyorum - see hensi

Diaptomidae - Calanidea.
Diaptomus deitersi Poppe, ♀. ♂. /

- Poppe, 1891^d, p. 248: 3 text-figs.
— Giesbrecht & Schmeil, 1898, p. 81.
— Lowndes, p. 96: fig. 2, a-b. 1934
— Wright, 1927, p. 95: pl. 8, figs. 5-6.
—, Pesta, 1927, p. 80

Habitat. Brasil.

- , Chappuis, 1927: 41

NOTODIAPTONUS DETERSSI (POPPE) ②

D., SPANDL 1926: 104-105, FIG. 7, CEARÁ, BRAZIL

-- BRANDORFF 1972: 44-, DISTRIBUTION

--, BRANDORFF, 1976: 616; FIG. 2

-- DUSSART 1984: 63

-- DUSSART 1985: 208, TYPE-SPECIES OF NEW
SUBGENUS NOTODIAPTONUS S. STR.; 210.

-- REID 1987a: 377

— Matsumura-Tundisi, 1966: 537, fig. 26-33

D., WRIGHT 1938: 562

-- DUSSART & FRUTOS 1985 (1986): 306

NOTODIAPTONUS DEITERSI (POPPE)

(3)

-- LANSAC-TÔ HA ET AL. 1997: 140, TAB. 3, PARANÁ, BRAZIL

NOTODIAPLOMUS DILATATUS DUSSART, 1984

- DUSSART 1984: 35, 38, 39, 51, 53-54, FIG. 8. N. SP. ♂♀,
-- DUSSART 1985: 210, IN SUBGEN. WRIGHTIUS VENEZUELA
- DUSSART & ROBERTSON 1984: 391

Notodiaptomus dubius Dussart 1985

①

-- Dussart 1985: 208-210, fig. 6, n. sp. ♀; in subgen.

Wrightius; article has priority over article of
Dussart & Matsumura-Tundisi

— Dussart & Matsumura-Tundisi, 1986:
250-254, fig. 2

-- REID 1987a: 378

— Matsumura-Tundisi 1986: 537, fig. 16-21

-- DEPAYE & DUSSART 1988 (89): 114

-- MATSUMURA-TUNDISI & TUNDISI 1986: 35-37, 39, 42,
TABS. 1, 2, LAGOA AMARELA, MINAS GERAIS, BRAZIL

NOTODIAPTOMUS DUBIUS DUSSART, 1985

-- HAMER ET AL. 1996: 92

(2)

Diaptomidae - Calanoida

Noto-

Diaptomus echinatus Lowndes ♂♀.

— — Lowndes, 1934, p. 100: fig. 4, a-d.

--, BRANDORFF, 1976: 618; FIG. 3

- s. lat. -, BREHM 1958a: 168 - Key.

-- DUSSART 1984: 65

-- BRANDORFF 197a: 49, DISTRIBUTION

Habitat. Brazil, Paraguay, ARGENTINA

-- WRIGHT 1937: 77, SÃO PAULO, BRAZIL

-- J. WRIGHT 1938: 562

— — — BREHMI 1955: 7 ARGENTINA

Notodiaptomus echinatus (Lowndes, 1934)

card 2

D. echinulatus, Brehm 1965:7, Argentina

Notodiaptomus -, n. comb., Defaye & Dussart 1988(89):

110, 111, 113, figs. 6-10, French Guiana; syn. N.
kieferi Brandorff

NOTODIAPTONUS FALCIFER

-see under DIAPTONUS FALCIFER

[Noto-

Diaptomidae - Calanoida.

Diaptomus gibber Poppe, ♂·♀. 1

1527

- — Poppe, 1889, p. 95: pl. 2, fig. 2, 14: pl. 3, fig. 1: pl. 4,
— — DeGuerne & Richard, 1889, pl. 18: table.
— — Herrick, 1895, p. 63: pl. 8, fig. 1,
— — Schmeil, 1898, p. 172: pl. 14, fig. 4-5.

-- BEANDORFF, 1976: 613

Habitat. Brazil, ARGENTINA

D. -, WRIGHT 1938: 562-

-- BREHM 1949: 416.

--, BREHM 1938: 30.

- Giesbrecht & Schmeil, 1898, p. 82.
— Daday, 1905, p. 150
— Tollinger, 1911, p. 70
— Wright, 1927, p. 89: pl. 6. figs. 4-6.
—, Pesta, 1927, p. 80
- s. lat. -, BREHM 1958a : 167 - Key.

NOTODIAPTONUS GIBBER (POPPÉ) 1889 2

N., N. COMB., PALLARES 1963: 39-42, LAM. II, BUENOS AIRES

D., BRANDORFF 1972: 49, DISTRIBUTION

--, BRANDORFF, 1976: 616; FIG. 2

-- DUSSART 1984: 63

-- DUSSART 1985: 210, IN SUBGEN. WRIGHTIUS - TYPE-SPECIES

-- DUSSART & ROBERTSON 1984: 391

-- DUSSART & FRUTOS 1985 (1986): 306

NOTODIAPTOMUS HANDELI - SEE

NEODIAPTOMUS SCHMACKERI

Diaptomidae - Calanoida.
Diaptomus hensenii Dahl, ♂. ♀. ,

— — Dahl, 1894, p. 19: pl. I, fig. 1-5, a.

— — Giesbrecht & Schmeil, 1898, p. 78

— — Daday, 1905, p. 151

— — Tollinger, 1911, p. 70

D. -, WRIGHT 1937: 76

Habitat. Mouth of the Amazon.

D. -, WRIGHT 1938: 562

— — Wright, 1927, p. 96: pl. 8, figs. 1-11.
 (= Diaptomus amazonicus Wright - Wright, 1935, pp. 221-222,
 Pl. I, fig. 2, 5, 9, 14, 16)

— — . - Wright, 1935, pp. 222-223, pl. 1, fig. 3.

— , Pesta, 1927, p. 80

— Wright 1936: 79

NOTODIAPTONUS HENSENI (DAHL, 1894) 2

-- BRANDORFF 1972: 44-, DISTRIBUTION

-- ANDRADE & BRANDORFF, 1975: 97

--, BRANDORFF, 1976: 616; FIG. 2

-- DUSSART 1984: 34, 39, 43, 46, FIG. 3 (= N. VENEZOLANUS KIEFER,
p. 63)

-- DUSSART 1985: 208, IN NOTODIAPTONUS S. STR.

— Matsumura-Tundisi, 1986: 542-543, fig. 81-85

Notodiaptomus hensenii (Dahl, 1894)

card 3

N. venezolanus deeveyorum, Bowman 1973:193-202, figs.
22-34, 36-39, n. ssp., Venezuela

N. deeveyorus, Dussart 1984:25, 34, 35, 38, 39, 46, 64, fig.
4, "nov. nom.", Venezuela (ACTUALLY NEW RANK)

N. (Notodiaptomus) deeveyorus, Dussart 1985:208

N. venezolanus deeveyorum, Matsumura-Tundisi 1986:542,
fig. 43-45

N. deeveyorum, Reid 1987a:378, CORRECTED NAME

N. DEEVEYORUS, GONZALEZ DE INFANTE 1988: 14-, 10⁸, FIGS. 5(9),
36(5a, b), VENEZUELA

= N. HENSENI

Notodiaptomus venezolanus Kiefer, 1956

n. sp., Kiefer, 1956, p. 233, 236-239, figs. 5-8

--, BRANDORFF, 1976: 616; FIG. 2

--, Kiefer, 1979, p. 224-232

--, Infante, 1978, passim

— Montiel & Zoppi, 1979: 114 - passim.

-- DUSSART 1984: 46, SYN. OF N. HENSENI (DAHL)

-- BRANDORFF 1972: 46-, DISTRIBUTION

Notodiaptomus hensenii (Dahl, 1894)

card 5

--, Cicchino et al. 1989:98-105, fig. 1, records,

Venezuela; synonyms N. venezolanus, N. deeveyorum

-- PAIVA 1990:109, PARA', BRAZIL

N. DEEVEYORUM, DEFAYE & DASSART 1988(89): 110,

111, 113, FIGS. 11-15, FRENCH GUIANA

CF.
N. DEEVEYORUM, DUSSART & FRUTOS 1985-(1986): 307-

308, FIGS. 7, 8, PARANA' RIVER, ARGENTINA

N. DEEVEYORUS, SAUNDERS & LEWIS 1988: 339-349,
LAKE VALENCIA, VENEZUELA

[Noto- Diaptomidae - Calanoidae

Diaptomus iheringi Wright, ♂♀.

1

— — Wright, 1935, p. 223: pl. I, fig. 4; pl. 2, figs. 3,
5-11.

-- Brehm, 1960, pp. 50.

-- WRIGHT 1936: 80, 81

D.-, S. WRIGHT 1937: 76

Habitat. Açude Puxinâna, Parahyba, NE Brazil.

D.-, S. WRIGHT 1938: 562

NOTODIAPTONUS IHERINGI (WRIGHT) 1935 2

-- BRANDORFF 1972: 45-, DISTRIBUTION

--, BRANDORFF, 1976: 616, 621; FIG. 2

--, REID + ESTEVES, 1984: 310-311, 317, 321-322.

-- DUSSART 1984: 63

-- REID 1987a: 377, 378

-- SENDACZ + KUBO, 1982: 51, 69-71

-- DUSSART 1985: 210, IN N. SUBGEN. WRIGHTIUS

Notodiaptomus iheringi (Wright) 1935

- Reid, 1985: 574-579, fig 1-28
— Matsumura-Tundisi, 1986: 542, fig 67-72

NOTODIOPTONUS -, ROLLA ET AL. 1990: 241-,
VOLTA GRANDE RESERVOIR, MG/SP, BRAZIL

- FONSECA & TUNDISI 1990: 106, BARRA BONITA RES, SP, BRAZIL
-- COSTA 1990: 112, BAHIA, BRAZIL
-- O. ROCHA ET AL. 1990: 94-, BARRA BONITA, SP, BRAZIL
-- MATSUMURA-TUNDISI, RIETZLER ET AL. 1990: 149-,
BRAZIL

Notodiaptomus iheringi (Wright, 1935)

card 4

- Tomm et al. 1992: 57-, Itaipu Reservoir, Paraná, Brazil
- LIMA ET AL. 1996: 115, MATO GROSSO DO SUL, BRAZIL
- REID + PINTO-COELHO 1994: 93-108, FURNAS RESERVOIR,
MINAS GERAIS/SÃO PAULO, BRAZIL
- LANSAC-TÔHA ET AL. 1997: 140, TAB. 3, PARANÁ, BRAZIL

[NOTE -

Diaptomidae Calanoida

Diaptomus incompositus Brian, ♂♀.

1

B

— Brian, 1925, p. 182; text figs. 7-9.

— Brehm, 1965, p. 8, Argentina; D. PARANAENSIS PESTA, p. 7, 8

--, BRANDORFF, 1976: 621

D., WRIGHT 1937: 76

Habitat. Río de la Plata: Río Uruguay: Río
Trocadero, Argentina.

D., WRIGHT 1938: 562, 56³; D. PARANAENSIS A SYNONYM.

--, BREHM 1949: 415, 416.

Notodiaptomus incompositus (Brian) 1925

2

--, Bowman, 1973, p. 199

--, BRANDORFF, 1976: 616, 620; FIG. 2; SYN: DIAPTOMUS
PARANAENSIS PESTA, 1927; BREHM, 1965;
D. incompositus BRIAN, 1927

--, BREHM 1949: 416.

--, BREHM 1938: 27-29.

--, Ringuet, 1959, p. 316

-- DUSSART 1984: 63

-- DUSSART 1985: 201, SYN. D. PARANAENSIS PESTA, 1927;
208, IN NOTODIAPTOMUS S. STR.

— Bello F., 1985: 5, FIG. 1-3

Notodiaptomus incompositus (Brian, 1925)

card 3

-- Brandorff 1972: 44-, distribution

-- Kontá 1980b: 57-, BRAZIL - RGS

-- Kontú + GLOEDEN 1986: 80-, FIG. 25, "

D. PARANAENSIS, PESTA 1927: 68-71, FIG. 1, N.sp.

D., S. WRIGHT 1939: 647, ARGENTINA; SYN. D.

PARANAENSIS

-- REID + MORENO 1990: 732

-- PAGGI + PAGGI 1990: 690-, RIO PARANA' MÉDIO, ARGENTINA

-- DUSSART + FRUTOS 1985(1986): 306

[Noto-] Diaptomidae - Calanoida

Diaptomus inflatus (Kiefer) ♂, ♀ 1

-- Kiefer, 1933, p. 38: figs. 1 - 7.

--, ANDRADE & BRANDORFF, 1975: ~~ILL~~ 102

--, BRANDORFF, 1976: 618; FIG. 3

- s. lat. - , BREHM 1958a: 166 - Key.

--, WRIGHT 1936: 79

Habitat Brazil

D., WRIGHT 1937: 76

D., WRIGHT 1938: 562

NOTO DIAPLOMUS INELATUS (KIEFER, 1933) 2

- BRANDORFF 1972: 45-, DISTRIBUTION
- ANDRADE & BRANDORFF, 1975: 97
- DUSSART 1984: 63
- DUSSART 1985: 210, IN SUBGEN. WRIGHTIUS
- DUSSART & ROBERTSON 1984: 391

Notodiaptomus inflexus Brian

- SEE ALSO N. ANISITSI

--, BREHM 1938: p. 29.

-- DUSSART 1984:63 (a possible syn. of anisitsi)

Notodiaptomus isabelae (Wright, 1936)

card 1

Diaptomus -, Wright 1936: 81-82, Pl. II fig. 5, n. sp.,

Brazil

D. -, Brehm 1949: 416

D. -, Brandorff 1972: 50, distribution

N. -, Bowman 1973: 199

N. -, Brandorff 1976: 616, fig. 2

N. -, Dussart 1984: 64

N. -, Dussart 1985: 208, in Notodiaptomus s. str.

N. -, Matsumura-Tundisi 1986: 542, figs. 55-60

Notodiaptomus isabelae (Wright, 1936)

card 2

--, Reid 1987a: 377

D., WRIGHT 1937: 76

D., S. WRIGHT 1938: 563

N., DUSSART & FRUTOS 1985 (1986): 307, FIGS. 3-6,
PARANA' RIVER TRIBUTARIES, ARGENTINA

N., TUNDISI & MATSUMURA TUNDISI 1981: 206-
RIO DORE SYSTEM, MINAS GERAIS, BRAZIL

N., LANSAC TOURA ET AL. 1993: 48-, PARANA' RIVER, BRAZIL

N., LIMA ET AL. 1996: 115, MATO GROSSO DO SUL, BRAZIL

NOTODIAPTOMUS ISABELAE (S. WRIGHT, 1936) ③

-- LANSAC-TÔHA ET AL. 1997: 140, 141, TAB. 3, PARANÁ,
BRAZIL

NOTODIAPTONUS STATOENSIS (WRIGHT) 1936
DIAPTONUS —, WRIGHT 1936: 82, PL. II FIG. 4, BRAZIL - N. SP.

D. —, BRANDORFF 1972: 50, DISTRIBUTION

--, BRANDORFF, 1976: 616; FIG. 2

-- DUSSART 1984: 64

-- DUSSART 1985: 208, IN NOTODIAPTONUS S. STR.

-- REID 1987a: 377

— Matsumura-Tundisi 1986: 542, fig. 73-77, LAKES
PARANOÁ, DF, + ITAIPU, BRAZIL

D. —, WRIGHT 1937: 76

P. —, J. WRIGHT 1938: 563

NOTODIAPROMUS KIEFERI BRANDORFF, 1973 ①

- , BRANDORFF, 1973: 205, 206-208 ♀ ♂; PL. 1-2
- , ANDRADE & BRANDORFF, 1975: 97
- , BRANDORFF, 1976: 616; FIG. 2
- DUSSART 1984: 35, 38, 39, 49-51, 64, FIG. 7
- DUSSART 1985: 210, IN SUBGEN. WRIGHTIUS
- DUSSART & ROBERTSON 1984: 391
- BRANDORFF 1972: 30-, FIGS. 40-48, AMAZON (THESIS);
ALSO D. -, p. 50
- SILVA ET AL. 1989: 728-, FIGS. 116-135, PARA, BRAZIL

Notodiaptomus kieferi Brandorff, 1973

card 2

-- Magalhães et al. 1988:271, types at INPA

-- DEFAYE & DASSART 1988(89): 113, SYN. OF

N. ECHINATUS

NOTODIAPROMUS LOBIFER (PESTAL)

--, BRANDORFF, 1976: 616; FIG. 2

-- DUSSART 1984: 63

-- DUSSART 1985: 201, NOT = *N. CONIFEROIDES*;
210, IN SUBGEN. WRIGHTIUS

-- DUSSART 1984d: 264, FIG. 6, NOT = *N. CONIFEROIDES*

-- DUSSART & ROBERTSON 1984: 391

-- DUSSART & FRUTOS 1985 (1986): 306

Notodiaptomus maracaibensis Kiefer, 1956

--, F. Kiefer, 1954, pp. 172 fig. 5 (Venezuela. S. Am.)

n. sp. Kiefer, 1956, p. 233, 235, 239-243, figs. 9-18

--, BRANDORFF, 1976: 616 s FIG. 3

-- DUSSART 1984: 64

-- DUSSART 1985: 208, IN NOTODIAPTONUS S. STR.

-- BRANDORFF 1972: 45-, DISTRIBUTION

-- CICCHINO ET AL. 1989: 101

-- HAMER ET AL. 1996: 92

[Noro- Diaptomidae - Calanoidae

Diaptomus nordestinus Wright. ♂♀

— — Wright, 1935, p. 214: pl. 1, figs. 1, 6-8, 10-14.

pl. 2, figs. 1, 2, 4: text-fig. 1.

-- Brehm, 1960, p. 50

-- WRIGHT 1936: 80

D., S. WRIGHT 1937: 76

Habitat. Açude Simão, Paraíba, NE Brazil.

D., S. WRIGHT 1938: 562

NOTODIAPTOMUS JORDANUS (WRIGHT) - 2-

- BRANDORFF 1972: 45-, DISTRIBUTION
- , BRANDORFF, 1976: 616, 621; FIG. 2
- DUSSART 1984: 63
- DUSSART 1985: 208, IN NOTODIAPTOMUS S. STR.

Notodiaptomus ohlei (BRANDORFF, 1978)

-- Dussart 1985: 214, in new subgen. Amazonius; TYPE-SPECIES
-, REID 1987a: 378, SUBGENUS NOT AVAILABLE

Notodiaptomus orellanai, n.sp.

--, Dussart, 1979, p. 5-6, fig. 3

-- DUSSART 1974: 64

-- DUSSART 1985: 210, IN SUBGEN. WRIGHTIUS

-- DUSSART & ROBERTSON 1984 : 391

-- DUSSART & FRUTOS 1985 (1986) : 307

Notodiaptomus paraensis Dussart & Robertson 1984

-- Dussart & Robertson 1984:389-394, figs. 1-3, n. sp. ♀♂

-- Dussart 1985: 210-212, fig. 7, in subgen. Wrightius

-- REID 1987a: 378

-- SILVA ET AL. 1989: 728-, FIGS. 69-93, PARÁ, BRAZIL

-- MAGALHÃES ET AL. 1988: 271, TYPES AT INPA

Notodiaptomus paraensis f. similis Dussart & Robertson

-- Dussart & Robertson 1984: 390-391, figs. 3C,D

Notodiaptomus paulistanus (WRIGHT, 1936)

-- Dussart 1985: 214, not assigned to subgenus

Notodiaptomus perelegans (Wright) - see Calodiaptomus

Notodiaptomus pseudodubius Defaye & Dussart 1989

-- Defaye & Dussart 1988(89):110, 111, 113-114, Figs.
16-24, n. sp. ♂♀, French Guiana

Notodiaptomus santafesinus - see Diaptomus

[Noto - Diaptomidae - Calanoida
Diaptomus santaremensis Wright, ♂♀.

— — Wright, 1927, p. 82: pl. 3, figs. 6-9.

— — Brehm, 1965, p. 8, Argentina

—, BRANDORFF, 1976: 618; FIG. 3

D., S. WRIGHT 1937: 76

D., S. WRIGHT 1938: 562, ILHA DE MARAJO, BRAZIL

Habitat. Santarem, Brazil.

2

NOTODIAPTONUS SANTAREMENSIS (WRIGHT, 1927)

- BRANDORFF 1972: 45-, DISTRIBUTION
- ANDRADE & BRANDORFF, 1975: 97
- DUSSART 1984: 63
- DUSSART 1985: 208, IN *NOTODIAPTONUS* S. STR.
- SILVA ET AL. 1989: 728-, FIGS. 94-115, PARÁ, BRAZIL

Notodiaptomus spiniger - see Diaptomus

Notodiaptomus spinuliferus Dussart

(1)

-- Dussart 1985: 208-210, fig. 6♀; in Notodiaptomus

s. str.; N.S.P. - WITH PRIORITY OVER ARTICLE

OF DUSSART & MATSUMURA-TUNDISI

— Dussart & Matsumura-Tundisi, 1986: 250, fig. 1

-- REID 1987a: 377

— Matsumura-Tundisi, 1986: 537, fig. 34-37

- c.f., REID & MORENO 1990: 726; PANTANAL, BRAZIL

- DUSSART & FRUTOS 1985(1986): 307, 308; PARANÁ RIVER, ARGENTINA

NOTODIATOMUS SPINULIFERUS DUSSART, 1985

(2)

-- LANSAC - TÔ HA ET AL. 1997: 140, TAB. 3, PARANÁ,
BRAZIL

Diaptomidae - Calanoida

[Noto-

Diaptomus transitans (Kiefer) ♀, ♂

— — Kiefer, 1929, p. 307 : figs. 6 — 8

— s. lat. — , BREHM 1958a: 167 - Key.

"D" - , DASSART 1985: 201, "D" MILDREDAE BREHM
POSSIBLY A SYN.

-- DASSART 1984d: 255, D. MILDREDAE A SYN.

Habitat. South America

D. - , S. WRIGHT 1938: 562

-- BRANDORFF 1972: 52, DISTRIBUTION

-- FORRÓ 1986: 560

NOTODIAPTONUS PRAEQUITANS (KIEFER) CARD 2

D., S. WRIGHT 1939: 648, ARGENTINA

N., BRANDORFF, 1976 ~~1975~~: 616; FIG. 2

N. DUSSART 1984: 63

N. DUSSART 1985: 214, IN SUBGEN. CALÉODIAPTONUS

N. Matsumura-Tundisi, 1966: 537, 542, fig. 38-42

N., DUSSART & FREITOS 1985 (1986): 306

Notodiaptomus venezolanus - see N. hensenii

Notodiaptomus venezolanus deeveyorum - see hensenii

NOTODIAPTONUS SP.

- DUSSART, 1984: 26
- BARBIERI ET AL. 1989: 406, MARANHÃO, BRAZIL;
= N. BRANDORFFI REID
- REID & MORENO 1990: 726-, PANTANAL, BRAZIL
- BOZELLI, 1992: 254; LAKES, PARÁ, BRAZIL
- REID 1993(1994) b: 71-, CENTRAL BRAZIL
- PARANAGUA' 1985: 774, PERNAMBUCO, BRAZIL
- LOPES ET AL. 1997: 45, TAB. 1c, PARANA', BRAZIL

NOTODIAPTONUS spp.

- TUNDISI + MATSUURA TUNDISI, 1981: 206-
RIO DOCE SYSTEM, MINAS GERAIS, BRAZIL
N. spp. 1+2, LIMA ET AL. 1996: 114-115, MATO GROSSO DO
SUL, BRAZIL
N. spp. 1+2, LANSAC-TÔHA ET AL. 1997: 140, 147, TAB. 3,
PARANÁ, BRAZIL