

in

Ceratohoeamptidae - Harpacticoida.
TETRAGONICIPITIDAE

Phyllopodopsyllus. valid.

T. Scott, 1906.

P. bradyi. ^(M)

Syn. Phyllopodopsyllus T. Scott, 1906,

Phyllopodopsyllus G.O. Sars, 1907, p. 230.

Tetragoniceps T. Scott, pars, not Brady.

φυλλον = leaf - πους = foot - ψύλλη = a flea.

- Lang, 1965: 386-389 Key

Body slender, constricted at center; caudal rami with lamellar expansion near the base. First antennae 9-jointed, first joint $\frac{1}{3}$ the whole length. First legs slender, exopod shorter than 1st. joint of endopod. Fourth legs with exopod twice as long as that of 2 preceding pairs; last legs more than $\frac{1}{3}$ the whole length of the animal, with tooth-like projections at the postero-superior corner.

Phyllopodopsyllus

TETRAGONICIPITIDAE

(2)

Pteropsyllidae-new fam., Nicholls 1944, p. 492

Review of genus and key to species: Nicholls 1944

- , VERVOORT, 1964: 276, 277, 278 - KEY

- DAHMS + BERGMANS 1988: 369

- POR & MARCUS, 1972: 272, Suez Canal

- HAYS 1992: 71-

- KUNZ 1984c: 33-, IN FAMILY REVIEW

PHYLLOPODOPSYLLUS AEGYPTICUS - GROUP KUNZ, 1984

-- KUNZ 1984 C: 38-

--- FIERS 1986 F: 110

PHYLLOPODOPSYLLUS BORUTZKYI - Group KUNZ, 1984

-- KUNZ 1984c: 38-

PHYLLPODOSYLLUS BRADYI - GROUP KUNZ, 1984

-- KUNZ 1984c: 37 -

PHYLLPODOPSYLLUS FURCIGER - GROUP KUNZ, 1984
-- KUNZ 1984c: 38-

PHYLLOPODOPSYLLUS LONGIPALPATUS - GROUP KUNZ, 1984
-- KUNZ 1984c: 43 -

PYLLPODOPSYLLUS MOSSMANI-GROUP KUNZ, 1984

-- KUNZ 1984: 41--

PHYLLOPODOPSYLLUS OPISTOCERATUS - G-ROUM KUNZ, 1984

-- KUNZ 1984 c: 41 -

PHYLLPODOPSYLLUS PAUKI - GROUP KUNZ, 1984

-- KUNZ 1984c: 38--

-- KUNZ 1985: 97, LIST; ADDS P. HARTMANNORUM

PHYLLPODOPSYLLUS XENUS-GROUP KUNZ, 1984

-- KUNZ 1984c: 41-

Phyllopodopsyllus aegypticus Nicholls

Nicholls- __, Nicholls 1944, p. 493, fig. 4, ♀, Red Sea

--, COULL, 1973a: 20; TAB. 2

--, FIERS 1986F: 110

-- BRICKMAN 1972: 135-, NEW JERSEY

-- MIELKE 1992b: 144

-- KUNZ 1984a: 27

-- KUNZ 1984b: 10

-- KUNZ 1984c: 38-

Phyllopodopsyllus alatus Fiers 1986f

-- Fiers 1986f:99, 106-110, figs. 7-8, n. sp. ♀♂,
Papua New Guinea

-- MIELKE 1989b:141, 144

Phyllopodopsyllus ancylus Mielke, 1992

-- Mielke 1992b: 101, 126-133, Figs. 18-21, n. sp.,
female + male, Punta Morales, Costa Rica (Pacific
side)

-- KUNZ 1995: 97

Phyllopodopsyllus angolensis Kunz, 1984

- KUNZ 1984b: 5- , FIG. 5-10, N.SP. ♀♂, ANGOLA
- Fiers 1986f:110
- MIELKE 1992b: 144
- MIELKE 1989b: 127-135, FIGS. 20-24, GALAPAGOS
- KUNZ 1984c: 38-

Canthocamptidae - Harpacticoida

Phyllopodopsyllus armatus Willey, ♂♀

— — Willey, 1935, p. 84: figs. 127-148.

Laophontella —, Sewell 1940, p. 336

— —, hans. 1948, p. 888.

Willeyella armata: Por, 1964, p. 105.

--, VERVOORT, 1964: 277

Habitat. Bermuda

-- BODIN 1964: 145 -

PHYLLOPODOPSYLLUS BAHAMENSIS GEDDES, 1968

- GEDDES, 1968a: 23-25 ♀; FIGS. 1-4
- , COLL, 1971: 201
- , COLL, 1973a: 15, 18; TAB. 2
- REID 1990g: 202

Tetragonicepsidae-Harpacticoida
Phyllopodopsyllus bermudae n. sp. LANG, 1948

— longicaudatus: Willey 1935, p. 88, figs 149-162.

— Lang 1948, p. 887

--, GEDDES, 1968a: 34

--, VERVOORT, 1964: 277, 278, 284

--, COULL, 1973a: 19; TAB. 1, 2

-- REID 1990g: 202

-- KITAZIMA 1981: 403-

-- KUNZ 1984b: 10

-- KUNZ 1984c: 38-

Phyllopodopsyllus berrieri Monard, 1936

Monard 1936, p. 27, fig. 14-15♀, Algeria, n. sp.

— —, LANG, 1948, p. 890.

--, COULL, 1973a: 20; TAB. 2

-- MIELKE 1992a: 98

-- MIELKE 1992b: 141

- KUNZ 1983: 120, 188-189, FIG. 39, AZORES

-- KUNZ 1984c: 41-

PHYLLOPODOPSYLLUS BIARTICULATUS WELLS, 1967

- Wells, 1967: 313-316, fig 61-62A-G
- , GEDDES, 1968a: 22
- , COULL, 1971: 201
- , COULL, 1973a: 15, 19; TAB. 2
- KUNZ 1984c: 43 -

PHYLLOPODOPSYLLUS BORUTSKYI LANG

- Lang, 1965: 395 - 399, fig. 217-219
- , GEDDES, 1968a: 25
- , COOK, 1973a: 18
- KUNZ 1984c: 38-

Canthocamptidae - Harpacticoida

Phyllopodopsyllus bradyi (T. Scott), ♂. ♀.

Tetragoniceps — T. Scott, 1892, p. 253: pl. 9, fig. 19-32

— — Brady, 1905, p. 217: pl. 5, figs. 1-10.

Phyllopodopsyllus — Sars, 1908, p. 231: pl. 155, ♀

— — Norman & Brady, 1909, p. 138.

— —, Wells, 1970, p. 264

Habitat, British Seas: Norway:

--, GEDDES, 1968a: 34

--, COUL, 1973a: 15, 18; TAB. 2; SYN: TETRAGONICEPS B.

— — T. Scott, 1906, b. p. 343.

— — Sars, 1911, p. 409: suppl. pl. 37, f. 2, ♂

^{PO}
Phyllodopsyllus bradyi (T. Scott,

- , Vervoort, 1964, p. 278, 279, 284
- , Klie, 1950, p. 101-2, figs. 111-114
- , Lang, 1944, p. 28, Tetragoniceps b. is a syn.
- MENZEL 1916b: 481.
- GERY & FLEEGER 1990: 294
- * BRAYDI, BODIN 1964: 145
- JAKOBI 1959: 279
- KUNZ 1984c: 37-
- KUNZ 1995: 93-

Phyllopodopsyllus briani Petkovski

1955 pp.126-130, figs.1-14.

--, COULL, 1973a: 15, 19; TAB. 2

--, VERVOORT, 1964: 277, 278, 286

--, Kunz, 1974, p. 266-271, figs. 73-79, 80-85, 86-88

-- MIELKE 1992b: 120-

-- MIELKE 1989b: 161

-- KITAZIMA 1981: 423-424

-- KUNZ 1984a: 19-, 31

-- KUNZ 1984b: 3; FIG. 1A, B, JUGOSLAVIA

-- KUNZ 1984c: 37-

Phyllopodopsyllus carinatus Mielke, 1992

-- Mielke 1992b: 101, 133-141, Figs. 22-27, n. sp.

female + male, Playa Manzanillo, Costa Rica

(Caribbean side)

Phyllopodopsyllus chavei Coull

- - Coull, 1970. p. 122

-- Coull, 1971, p. 208, 218

--, Coull & HERMAN, 1970: 394

--, Coull, 1973a: 15, 19; TAB. 2

-- Coull 1971: 221

-- REID 1990g: 202

-- KUNZ 1984c: 38-

Phyllopodopsyllus curtus Marcus, 1976

-- Kitazima 1981: 393-

-- Kunz 1984c: 38-

Phyllopodopsyllus danielae Bodin, 1964

- Bodin 1964: 146, 161-163, Pl. XVII, XIX, n. sp.
female + male, Gulf of Marseilles, Mediterranean
- Guille & Soyer 1966: 376
- Geddes 1968a: 22
- Coull 1971: 221
- Coull 1973a: 15, 19, tab. 2
- Mielke 1992b: 133, 141
- Kunz 1984c: 41-
- Kunz 1995: 97

Phyllopodopsyllus dissimilis

--, Vervoort, 1964, p. 277

Canthocamptidae-Harpacticoida

Phyllopodopsyllus gurgiger G.O. Sars, ①

— — Sars, 1908, p. 233: pl. 156 fig. 2. Norway
Farsund.

— — Sewell 1940, p. 304, fig. 70, South Arabian Coast

— — ♀ HANE, 1948, p. 886, fig. 356:3.

Phyllopsyllus^[?] —, Delamare Debutteville + Chappuis, 1957 (Bull. D.F.A.N.) p.

494. Luanda, Angola, Africa.

Habitat. Norway Coast.

— —, Por, 1964, p. 102, pl. 18 figs. 201-205

— —, COULL + HERMAN, 1970: TAB. 1

P. furcifer, Vervoort, 1964, p. 277

--, GEDDES, 1968a: 34

--, COULL, 1973a: 19; TAB. 2

PHYLLOPODOPSYLLYS FURCIFER, JAKOBI 1959: 279

P. FURCIFER, KRISHNASWAMY 1957d: 58, 59-60,
MADRAS, INDIA

-- COULL 1968: 72-, BERMUDA

-- REID: 1990g: 202

-- MIELKE 1989b: 144-152, FIGS. 31-35, GALAPAGOS

P. FURCIFER, BODIN 1964: 142-, PL. VIII FIG. 49, GULF
OF MARSEILLES

Phyllopodopsyllus furciger G.O. Sars, 1908

card 2

-- Kunz 1984c: 38--

-- KUNZ 1995: 83; FIGS. 1, 2A, TANZANIA

Phyllopodopsyllus aff. furciger Por

--- Kunz 1984c: 38-

Phyllopodopsyllus galapagoensis Mielke, 1989

- MIELKE 1989b: 153-162, FIGS. 36-41, n. sp. ♀♂,
GALAPAGOS; 161
- Mielke 1992b: 120, 125, 126

Phyllopodopsyllus geddesi Kunz, 1984

-- Kunz 1984a: 21-24, Figs. 14-19, n. sp. female + male,

Hawaii

-- Kunz 1984c: 41-

Phyllopodopsyllus gertrudi Kunz, 1984

- KUNZ 1984a: 24-27, FIG. 20-25, N.SP. ♀♂ HAWAII
- Fiers 1986f:110
- MIRLKE 1992b: 144, CF. P. GERTRUDI COSTARICENSIS
- KUNZ 1984b: 10
- KUNZ 1984c: 38-

Phyllopodopsyllus gertrudi costaricensis Mielke, 1992

--- Mielke 1992b: 101, 141-144, Figs. 28, 29, n. ssp.,

Playa Manzanillo, Playa Blanca^c/_^Osa, Costa Rica,

Caribbean & Pacific sides

PHYLOPODOPSYLLUS HARDINGI (ROE, 1935)

- COULL, 1973a: 19; TAB 2
- KUNZ 1984c: 41-

Phyllopodopsyllus hartmannorum Kunz, 1995

-- Kunz 1995: 83^e, Fig. 9B-D, 10-12, n. sp. female only,
Indian Ocean: Tanzania: Tanga

Phyllopodopsyllus hermani Coull

-- Coull B, 1969. p. 27

--, COULL, 1970: 345, 346, 347, 351, 352, 353; FIG. 3; TAB. 8

--, COULL & HERMAN, 1970: 394, 397

--, COULL, 1973 a: 15; TAB. 2

-- COULL 1968: 59-, BERMUDA

-- COULL 1971: 221

-- REID 1990 g: 202

-- KUNZ 1984 c: 38-

PHYLOPODOPSYLLUS HIBERNICUS (ROE, 1955)

—, COULL, 1973a: 20; TAB. 1

-- MIELKE 1992a: 98

-- MIELKE 1992b: 141

-- KUNZ 1984c: 41-

PHYLOPODOPSYLLUS INTERMEDIUS

--, = P. THIEBAUDI, WELLS, 1973: 137

--, WELLS + MCKENZIE, 1973: 137, POSSIBLY SYN. OF
P. THIEBAUDI.

--, FIERS 1986 F: 110

Phyllopodopsyllus kunzi Mielke, 1989

-- Mielke 1989b: 162-168, Figs. 42-44, n. sp. female +
male, Galapagos

-- Kunz 1995: 97

Phyllopodopsyllus langi Kunz, 1975

— Kunz, 1975 : 203-205, figs. 210-233

-- KITAZIMA 1981: 393-

.. KUNZ 1984c: 38-

PHYLLOPODOPSYLLUS LASPALMENSIS MARINOV, 1973

- MARINOV, 1973b:
- WELLS, 1979: 4
- MIELKE 1992a: 98
- MIELKE 1992b: 141
- KITAZIMA 1981: 393-

Phyllopodopsyllus laticauda Por

-- --, Por, 1964, p. 102, pl. 18 figs. 196-200

--, COLL, 1973a: 19; TAB. 2

-- DUSSART & DEFAYE 1990: 16, AS "PHYLLOGNATHOPUS
LATICAUDA" - PROBABLY A LAPSE.

Canthocamptidae - Harpacticoida

Phyllopodopsyllus longicaudatus A. Scott, ♂.

— — A. Scott, 1909, p. 223: pl. 60, f. 17-25.

— — Willey, 1935, p. 88: figs. 149-162.

— —, Hancock, 1948, p. 887.

— —, Wells, 1970, p. 264

Habitat. Paternoster Islands, Bermuda.

— —, Coull, 1973a: 15, 18; TAB. 2

— Wells, 1967: 193

--, Vervoort, 1964, p. 6, 277, 278, 279-284, figs. 112, 114a, d

-- KRISHNASWAMY 1957d: 58-59, FIG. 10, MADRAS, INDIA

-- SENSU VERVOORT, KUNZ 1984c: 37-

-- KUNZ 1995: 96-97

PHYLLOPODOPSYLLUS LONGIPALPATUS (CHAPPUIS, 1953)

- WELLS, 1979: 4
- COLL, 1973a: 19; TAB. 2
- KITAZIMA 1981: 412-
- KUNZ 1984a: 17-
- S. STR., KUNZ 1984b: 3, FIG. 1C-E, ITALY
- S. STR., KUNZ 1984c: 43-

Phyllopodopsyllus longipalpatus ^ahawiiensis Kunz, 1984

--- Kunz 1984^a: 14-17, Figs. 3-7, n. ssp. female + male,
Hawaii¹

--- Kunz 1984b: 5

--- Kunz 1984c: 43-

Phyllopodopsyllus longipalpatus madagascariensis Kunz, 1984

--- KUNZ 1984b: 3-5, FIG. 2, 3, N. S.S.P., AMBILO -
LEMAITSO, MADAGASCAR

--- Kunz 1984a: 17

--- KUNZ 1984c: 43 -

Phyllopodopsyllus medius Por, 1964

-- -- Por, 1964, pp. 99-101, pl. 18 figs. 190-195

--, COULL, 1973a: 19; TAB. 2

PHYLLOGNATHOMUS --, DUSSART + DEFAYE 1990: 16
(PROBABLY A HAMSE)

Phyllopodopsyllus mielkei mielkei Kunz, 1984

-- KUNZ 1984a: 27-30, FIG. 26-31, N.S.P. ♀♂, HAWAII

(also described ssp. californicus in same paper)

--- Mielke 1992b: 120

--- MIELKE 1989b: 161

P. MIELKE, GEE & FLEEGER 1990: 294

--- KUNZ 1984c: 37-

Phyllopodopsyllus mielkei californicus Kunz, 1984

-- KUNZ 1984a: 30-31, FIGS. 32-35, N. SSP.

♀ ONLY, SAN FRANCISCO, CALIFORNIA

--- Mielke 1992b: 120

-- KUNZ 1984c: 37-

Canthocamptidae - Harpacticoida

Phyllopodopsyllus minor (Thompson & Scott) ♂♀

Tetraگونiceps - Thompson & Scott, 1903, p. 268: pl. 8, f. 23-28

Phyllopodopsyllus - Willey, 1935, p. 91: figs. 163-172

--, Hancock, 1949, p. 586.

--, Vervoort, 1964, p. 278, 279, 284

--, Hicks, 1977a: TAB. 3, 4

Habitat. Bermuda, Ceylon.

-- KRISHNASWAMY 1957d: 58, MADRAS, INDIA

-- COULL, 1973a: 19; TAB 12

1971: 12 - ST. THOMAS

Phyllopodopsyllus minutus n. sp. Tetragonicepsidae-Harpacticoida

_____ Lang 1948, p. 887

--, Vervoort, 1964, p. 277, 278, 284

--, COULL, 1973a: 18; TAB. 2

- REID 1990g: 202

- KUNZ 1984c: 38-

~~Canthocamptidae - Harpacticoida~~

Phyllopodopsyllus mossmani T. Scott, ♂. ♀. ①

— — T. Scott, 1912, p. 563: pl. 5. figs. 1-14.

— —, LANG, 1945, p. 890.

--, Vervoort, 1964, p. 276

--, Lang, 1944, p. a syn. of Paraphyllopodopsyllus m.

--, WELLS, 1979: 4

Habitat. Falkland Islands.

--, COOK, 1973a: 20; TAB. 2

— Wells, 1967: 193

— MIELKE 1992 a: 98

Phyllopodopsyllus mossmani T. Scott, 1912

card 2

-- Mielke 1992b: 141

-- Kunz 1984c: 41-

Phyllopodopsyllus mossmani chiloensis Mielke, 1992

--- Mielke 1992a: 79, 92-98, Figs. 7-10, n. ssp.,

female + male, Quellon Voejo, Isla Chiloe, Chile

PHYLLOPODPSYLLUS OPISTHOCERATUS GEDDES,
1968

- GEDDES, 1968a: 25-31 ♀♂; FIGS. 5-7
- COULL, 1971: 221
- COULL, 1973a: 15, 18; TAB. 2
- REID 1990g: 202
- OPISTHOCERATUS, KUNZ 1984a: 24
- OPISTHOCERATUS, KUNZ 1984c: 41-

Phyllopodopsyllus pallaresae Kunz, 1995

-- Kunz 1995: 83-, Fig. 3C-E, Figs. 4-8, 9A, n. sp. female
+ male, Namibia

Phyllopodopsyllus paraborutzkyi Kunz

- Kunz, 1975: 205-207, figs. 242-258,

- KITAZIMA 1981: 393-

- KUNZ 1984c: 38-

PHYLOPODOPSYLLUS PARABRADYI LANG

— Lang, 1965: 389-395, fig. 212-216

--, COULL, 1973a: 18; TAB. 2

-- KUNZ 1984c: 37 -

PHYLLODOPSYLLUS PARAFURCIGER GEDDES, 1968

-- GEDDES, 1968a: 31-34 ♀♂; FIGS. 8-9

--, COULL, 1971: 201, 202, 223

--, COULL, 1973a: 15, 19; TAB. 2

-- REID 1990g: 202

- S. STR., KUNZ 1984c: 38-

PHYLOPODOPSYLLUS PARAFURCIGER CAROLINENSIS
COULL, 1971

- COULL, 1971: 221-223 ♀♂; FIG. XII
- COULL & HERMAN, 1970: 397
- COULL, 1973a: 15, 19; TAB. 2
- COULL, 1972: 49
- KUNZ 1984c: 38-

Canthocamptidae - Harpacticoida

Phyllopodopsyllus paramossmanni Lang, ♂♀.

— — Lang, 1935, p. 35: figs. 80-87.

— — Lang, 1948, p. 590

--, Wells, 1979: 4

--, Coull, 1973a: 20; TAB. 2

-- Kunz 1984c: 41-

Habitat, Campbell Island.

Phyllopodopsyllus paraxenus Coull, 1970

-- Coull 1970. p.119

- COULL & HERMAN, 1970: 394
- COULL, 1973a: 15, 19; TAB. 2
- COULL 1971: 221
- REID 1990g: 202
- KUNZ 1984c: 41-

Phyllopodopsyllus pauli Crisafi

①

Crisafi, P. ----1960, pp. 495-501, figs. 1-8.

-- --, Crisafi, 1961. pp. 263-267, fig. 1-3

--, Verveort, 1964, p. 277, 278, 286

- KUNZ 1984 a: 5, FIG. 4, TUNISIA

--, COULL, 1973 a: 19; TAB. 2

--, Kunz, 1974, p. 266, figs. 55-61, 62-67, 68-70

--, Crisafi, 19__ , p. 841

-- CECCHERELLI 1988: 155

-- MIELKE 1992 b: 133-

Phyllopodopsyllus pauli Crisafi, 1960

card 2

-- Kunz 1984c: 38-

-- KUNZ 1995: 97

Phyllopodopsyllus petkovskii Kunz, 1984

-- Kunz 1984a: 18-20, Figs. 8-13, n. sp. female + male,
Hawaii

-- Mielke 1992b: 120

-- MIELKE 1989a: 161

-- KUNZ 1984c: 37 -

PHYLOPODOPSYLLUS PIRGOS APOSTOLOV, 1969

-- COULL, 1973a: 15; TAB. 2

- KUNZ 1984c: 38, ? IMMATURE ♂ OF THIEBAUDI? -

SPECIES INCERTA

PHYLOPODOPSYLLUS PONTICUS APOSTOLOV, 1968

- APOSTOLOV, 1968 : 209-213
- , COULL, 1971: 221
- , COULL, 1973a: 15, 20; TAB. 2

Phyllopodopsyllus punctatus Kitazima, 1981

-- Kitazima 1981: 403-413, Figs. 8-14, n. sp. female +
male, Japan, Inland Sea, Mukaishima Island, Pl. I Figs. 3, 3

- KUNZ 1984 a: 17-

- KUNZ 1984 b: 5

- KUNZ 1984 c: 43-

Phyllopodopsyllus setouchiensis Kitazima, 1981

- Kitazima 1981: 413-424, Figs. 15-21, Pl. I Fig. 4,
n. sp. female + male, Japan, Inland Sea
- Mielke 1992b: 101, 120-126, Figs. 13-17, Punta
Morales, Playa Manzanillo, Playa Jimenez/Osa, and
Playa Blanca/Osa, Costa Rica (both Pacific and
Caribbean sides)
- MIELKE 1989b: 161
- KUNZ 1984a: 30, 31
- KUNZ 1984c: 37-

Phyllopodopsyllus simplex Kitazima, 1981

-- Kitazima 1981: 394-403, Figs. 1-7, n. sp. female +
male, Japan, Inland Sea, Mukaishima Island; Pl. I Fig. 1

- Kunz 1984c: 91-

- Kunz 1995: 97

Phyllopodopsyllus tenuis Wells & Rao, 1987

-- Kunz 1995: 97

Phyllopoppsyllus thiebaudi Petkovski, 1955

①

1955 pp. 130-134. figs. 15-20, *n. sp.*

--, Vervoort, 1964, p. 277, 278, 286

--, COULL, 1973 a: 15, 20; TAB. 2

--, WELLS, 1973: 133, 134, 137; FIG. 25 SYN:

P. INTERMEDIUS

-- WELLS + MCKENZIE 1973: 137, POSSIBLY P.

INTERMEDIUS IS SYN. OF P.T.

-- FIERS 1986 F: 106, 110

-- MIELKE 1992 a: 98

-- MIELKE 1989 b: 141, 144

Phyllopodopsyllus thiebaudi Petkovski, 1955

card 2

-- Kunz 1984c: 38-

PHYLLOPODOPSYLLUS CF. THIEBAUDI PETKOUSKI, 1955

--, WELLS + MCKENZIE 1973: 133, 134, 137, Fig. 2e-h;

P. INTERMEDIUS POSSIBLY SYN. OF P. T.; ALDABRA.

Phyllopodopsyllus thiebaudai santacruzensis Mielke 1989

-- Mielke 1989b: 135-144, Figs. 25-30, n. ssp., Galapagos

PHYLLOPODOPSYLLUS TRISTANENSIS (WIBORG, 1964)

--, GEDDES, 1968a: 22

--, COULL, 1971: 221

--, COULL, 1973a: 9, 15, 18

PHYLOPODOPSYLLUS XENUS (KUNZ, 1951)

-- COULL, 1973a: 20; TAB. 2

- KUNZ 1951: 41-

- KUNZ 1995: 83-, FIGS. 2B-F, 3A-B, NAMIBIA

PHYLLPODOPSYLLUS sp.

- COULL 1968: 59-, BERMUDA
- COTTARELLI 1977b: 98, CAMPECHE, MEXICO
- HARTZ BAND 1971: 12-, ST. THOMAS
- COTTARELLI 1971c: 29, ITALY
- ALVAREZ, M. 1987: 5, SÃO PAULO, BRAZIL

PHYLLIPODOPSYLLUS SP. ? OF BODIN, 1964

- - BODIN 1964: 145-146, PL. X FIGS. 57-61, IX FIGS. 50-52,
♂♂, GULF OF MARSEILLES
- - MIELKE 1992b: 117, PROBABLY = LAOPHONTELLA
HORRIDA (POR, 1964)