

in) Siphonostomatoida Monostelloida — Choniostomatidae.

Stenothocheres - valid

Hansen, 1897 - S. egregius.

Syn. Stenothocheres. Hansen, 1897, p. 88,

Stenothoë, amphipod genus, and cheres -
seeking or liking -

♀. Body longer than broad - naked | head not distinct; 1° ant! long - without distinct joints - 2° ant! medium - mouth with very short hairs - 1° mx. without branch: 2° mx. stout, without hairs. Myids small, 2° + 3° joints fused & all joints without hairs or spines - Trunk legs far from lateral margin - large - biramous - ramus 1-jointed - Basal part of abdomen stout - distal part narrowed - Gen. area wanting: gen. apertures partly on lateral surface -
♂ Head half & body; trunk legs close to margin only - narrower distal abdomen can be distinguished from trunk - Body naked, but covered dorsally with knobs & ventrally with keels & knots.

Choniostomatidae - Monstrilloida.

Stenothocheres egregius Hansen, ♀, ♂

— — Hansen, 1897, p. 89: pl. 1, fig. 1, a-l.

— — T. Scott, 1904, b, p. 250: pl. 15, fig. 5-10.

— — Stephensen, 1913, p. 350.

— — Wilson, 1920, p. 12 L.

Host. Marsupium of Metopa borealis, &
M. bruzelii (Goës). (Crustacean).

Greenland.

stenotocheres Hansen, 1923, p. 65

Stenothocheres sarsii, Hansen ♂, ♀.
^{Chonistomatidae}

— —, Hansen, 1897, p. 91; pl. I, figs. 2a-c.

Host. Stenothoe marina Sp. Bate.
(a crustacean)
Norway.