

in)

Caligoida
Lernaeidae.

Therodamas, valid.

Krøyer, 1863,

T. serrani.

Syn. Therodamas, Krøyer, 1863, p. 316.
Therodamas, Wilson, 1917, p. 41.

Θηροδάμας = A Scythian king who fed
lions with human flesh.

Head rounded, without horn-like appendages;
Mouth-opening at the base - long, slender neck.

Thorax much elongated, indistinctly segmented,
Gen. seg't. short - abdomen

small with minute anal laminae.

Post. antennae strongly uncinatae.

All four pairs of thorax legs present, biseriate; rami 3-jointed.

Inerodamas Kroyer, 1883

(2)

-- Cressy 1972:267

--, Jones & Hine, 1978, p. 218

- FRYER 1978a:345

- ABDELHALIM ET AL. 1993:45-, AMAZONICOPEUS ASYN.

Therodamas dawsoni Cressey

--, Cressey, R.F., 1972, pp. 265, 267, 268, 269, figs.
1-10

Therodamas fluviatilis Paggi, 1977

-- Fryer 1978a:345

Therodamas serrani, Kröyer, ♀.

— —, Kröyer, 1863, p. 316; pl. XV, fig. 4 a-f.

— —, B-Smith, 1899, p. 482.

— —, Wilson, 1913, p. 257; 1917, a, p. 41.

— —, Wilson, 1917, pp. 41-42.

— —, Thomsen, 1949, pp. 30-35, pl. XI, fig. 1-16 ^{gills of} Tachyurus barbus

Host. Gills of Serranus, sp

West Indies.

--, Cressey, R.F., 1972, pp. 267

Therodamas sphyricephalus Thomsen, 1949, pp. 35-36,
pl. 12, fig. 1-6. Host: Tachyurus barbus (Lac.),
on gills.

--, Cressey, R.F., 1972, pp. 267, 269

THERODAMAS spp.

-- KNOFF, LUQUE & TAKEMOTO 1995:45