

Tucca, valid, Poecilostomatoida Cyclopoida
Ergasilidae.
Kröyer, 1837 *T. impressa*.

Has no synonyms.

Tucca, a friend of Horace & Virgil.

Head small, rounded, with a lamellar appendage on either side: neck distinct.

No dorsal plates on Thorax seg'ts—
Gen. seg't large, oval. Abdomen small,
2-jointed. First antennae many-jointed,
setaceous: second pair small, uncinate.

Tucca Kroyer

-, Izawa, 1971, p. 182

-, DOJIRI + Cresssey 1987: 241

- VERVOORT 1962: 93, DIAGNOSIS, KEY

Tucca corpulentus, n.sp. Wilson. ♀.
Ergasilidae.

—. Wilson, 1911, p. 358: pls. 49,
figs. 116 & 117: pl. 50.

—. Wilson, 1932, p. 380: fig. 235

--, Ho, 1967, p. 286, 7, 8 = T. impressus

-- VERVOORT 1962: 95-96

Host. Pectoral fin of Sphoeroides maculatus.

Woods Hole.

Ergasilidae.

Tucca impressus. Kröyer. ♀. ♂, 1837

— —. Kröger, 1837, p. 479; pl. V, fig. 2, a-h.

— —, M. Edwards, 1840, p. 496.

— —, Nordmann, 1864, p. 491; pl. VI, figs. 7-10.

— —, B. Smith, 1899, p. 469.

Host. Diodon hystrix, Chilomycterus
Schoepfi. Atlantic.

- —, Wilson, 1911, p. 354, pls. 48 + 49.
— —, Wilson, 1913, p. 200
— —, Wilson, 1932, p. 379: fig. 234.
— —, Causey, 1955, pp. 2-19.. Gulf of Mexico
— —, Ho, 1967, pp. 285, 287-297, figs. 1-27
— —, Kabata, 1970, fig. 11
— PERE 1936: 582, FLORIDA-GULF, HOST
CHILOMYCTERUS SPINOSUS

Tucca impressus

card 2

--, HO, 1978, p. 5, 10, fig. 4

-- DOJIRI + CRESSY 1987: 240

-- PEARSE 1952: 191, HOST CHILONYCTERUS SCHOEPFI,
ALLIGATOR HARBOR, FLORIDA

-- VERVOORT 1962: 93 - 95

-- KAZACHENKO 1994: 34, FIG. 7B

Ergasilidae.

Tucca verrucosus, n. sp. Wilson. ♀.

— . Wilson, 1911, p. 359.

— impressus Nordmann, 1864, p. 491; pl. 6, fig. 7-10,

— , Ho, 1967, p. 286, 7

-- VERVOORT 1962: 96

Host. Diodon, sp.

Africa.

Tucca sp. Pearse, 1952

- Pearse 1952:12-13, figs. 23-27, Texas, hosts Lagocephalus laevigatus and Prionotus ophyras
- Ho 1967:287, possibly a chondracanthid?

Bomolochidae - Poecilostomatoida

Tuccinae Vervoort, 1962

- Vervoort 1962: 92-93, n. subfam.