

PROCEEDINGS

OF THE

ZOOLOGICAL SOCIETY

OF LONDON,

WITH ILLUSTRATIONS.



1848.

PRINTED FOR THE SOCIETY;
SOLD AT THEIR HOUSE IN HANOVER SQUARE,
AND BY MESSRS. LONGMAN, BROWN, GREEN, AND LONGMANS,
PATERNOSTER ROW.

turnal Lepidoptera are often almost impossible to describe, and it is only by the most accurate figures, or by comparison of specimens, that we can arrive at the determination of species.

I am indebted to Mr. Alfred Lambert of Sydney for the specimens figured and for the drawing of the larva. The following note accompanies the specimen:—

"The larva is figured in drawing No. 2. When I first found it I concluded that it was a *Cerura*, as in its habits it resembles the larva of that genus. It forms a strong cocoon, which is slightly attached to the trunk of the tree just below the surface of the ground. In form this cocoon is much like that of our common *Saturnia*, only exteriorly it is covered with points of sticks, grass, &c. The larva feeds on the *Eucalypti*, is found in January; the imago appears in July."

From this it will be seen that it is a winter insect.

3. DESCRIPTION OF TWENTY-NINE NEW SPECIES OF HELICINA, FROM THE COLLECTION OF H. CUMING, ESQ. BY DR. L. PFEIFFER.

1. *HELICINA ACUTA*, Pfr. *Hel. testū depresso-conicā, soliduld, obliquē confertim striatā et subgranulatā, opacā, luteā, supernē rubro-unifasciatā; spirā conoideā, acutā, mucronatā; anfractibus ferè 6 planiusculis, acutē carinatis, ultimo basi planiusculo; aperturā perobliquā, subtriangulari; columellā subverticali, brevissimā, basi angulatā, supernē in callum basalem tenuissimum abiente; peristomate simplice, aurantiaco, margine supero subrecto, basali subin-crassato.*

Diam. 15, altit. $7\frac{1}{2}$ mill.

From Sibonga, isle of Zebu; collected by Mr. Cuming.

2. *HELICINA ADAMSIANA*, Pfr. *Hel. testū depressā, tenuiusculd, sub lente seriebus confertis concentricis pustularum exiguarum subasperatā, nitiduld, diaphand, rubellā; spirā brevissimē conoidē; anfractibus $5\frac{1}{2}$ planiusculis, ultimo depresso, peripherid rotundato, anticē non scrobiculato; aperturā obliquā, subtriangulari; columellā verticali, brevissimā, basi subangulatā, supernē in callum tenuem, circumscriptum dilatatā; peristomate angulatim expanso, reflexiusculo, margine supero breviter soluto, stricto, basali prope columellam subdentato.*

Diam. 8, altit. $4\frac{1}{3}$ mill.

From Jamaïca.

3. *HELICINA AMENA*, Pfr. *Hel. testū subsemiglobosā, solidiusculd, obliquē striatulā lineisque impressis concentricis distantibus sculpta, nitiduld, roseo et luteo vel albo variegatā; spirā convexa, mucronulatā; anfractibus $5\frac{1}{2}$ vix convexiusculis, ultimo infra medium carinato, basi subplano; aperturā obliquā, subtriangulari, intus flavid; columellā brevi, verticaliter subrimata, basi angulosā, retrorsum in callum tenuem, diffusum abiente; peristomate simplice, margine supero late expanso, basali reflexo.* ? 123 1953

Diam. 15, altit. $9\frac{1}{2}$ mill.

From Honduras.

4. **HELICINA BESCKEI**, Pfr. *Hel. testā subsemiglobosā, solidā, subtilissimè striatulā, sublævigatā, opacā, citrind unicolorē vel fasciā 1 sanguinē juxta suturam ornatā, vel omnino rubicundā; spirā brevi, convexo-conoideā, submucronatā; anfractibus 5 subplanis, ultimo ad peripheriam carinā 1 acutā, pluribusque obtusioribus munito; aperturā obliquā, subtriangulari; columellā breviter recedente, basi obsoletē angulatā; peristome expando, subincrassato, margine supero strictiusculo, basali subarcuato; callo basali tenuissimo.*

Diam. 17, altit. 10 mill.

From Brazil (Bescke).

5. **HELICINA CAMPANULA**, Pfr. *Hel. testā campanulato-conicā, solidulā, lævigatā, nitidā, citrind; spirā elevatā, convexā, acuminatā; anfractibus 6 planiusculis, ultimo pone aperturam subconstricto, basi planulato; apertura obliquā, semilunari-subtriangulari; columellā breviter recedente, basi subtruncatā, callum nitidum, semi-circularem emittente; peristome simplice, tenui, breviter expando, margine basali strictiusculo.*

Diam. $8\frac{1}{3}$, altit. 7 mill.

From the island of Cuba.

6. **HELICINA CONCENTRICA**, Pfr. *Hel. testā depresso-trochiformi, tenuiusculd, striis longitudinalibus et obliquis sub lente subtilissimè sculptā, lineis concentricis elevatis utrinque munita, acutē carinatā, nitidulā, carneo-fusca, albido variegatā; spirā conoidea, subpapillatā; anfractibus $4\frac{1}{2}$ vix convexiusculis, ultimo utrinque convexiore; apertura obliquā, subsecuriformi, latiore quam alta; columellā subrimatā, breviter arcuatā, basi incrussata in callum album subcircumscripum retrorsum dilatatā; peristome simplice, breviter expando, margine basali immediate in columellam continuato.*

Diam. 10, altit. vix 6 mill.

From Venezuela and New Granada (De Lattre); a larger variety from Mirador, Mexico (Galeotti).

- (33.193)* 7. **HELICINA CONSTRICTA**, Pfr. *Hel. testā parvā, lenticulari, crassā, sublævigatā, non nitidā, opacā, albida, lineis undulatis rubris pictā; spirā vix elevatā, obtusa; anfractibus $4\frac{1}{2}$ planulatis, ultimo angulato, basi subturgido, pone aperturam constricto; apertura obliquā, subtriangulari, intus rubrā; columellā simplice, callum crassiusculum albidi vel igneum retrorsum emittente; peristome simplice, obtuso, latere dextro rotundato.*

Diam. 6, altit. $3\frac{1}{2}$ mill.

From Otaheite and the Sandwich Islands.

8. **HELICINA CONVEXA**, Pfr. *Hel. testā convexo-orbiculatā, solidā, lævigatā, nitidā, albā; spirā fornicate, nucronulatā; anfractibus $4\frac{1}{2}$, ultimis 2 convexiusculis, ultimo obsoletissimè angulato; apertura integrā, obliquā, semilunari; columellā breviter arcuatā, retrorsum in callum crassum, concolorem abiente; peristome incrassato, breviter expando, margine basali à callo columellari incisurā levissimā separato.*

Diam. $6\frac{1}{2}$, altit. $4\frac{1}{2}$ mill.
Locality unknown.

9. HELICINA CUMINGIANA, Pfr. *Hel. testa subglobosa, tenuiuscula, longitudinaliter et confertim plicatula, carneâ, rubro punctata et variegata; spirâ brevi, conoidea, obtusiuscula; anfractibus $5\frac{1}{2}$ planiusculis, ultimo inflato, obsoletè angulato; aperturâ subverticale, semiovali; columella basi dilatata, antrorsum arcuata, subtruncata, retrorsum in callum basi crassum, supernè diffusum abiente; peristomate subincrassato, breviter expanso, albo.*

Diam. 21, altit. $16\frac{1}{2}$ mill.
Locality unknown.

10. HELICINA DYSONI, Pfr. *Hel. testa orbiculato-conoidea, solidiuscula, striatula, nitidula, carnea, supernè fasciis 2 angustis, saturationibus ornata; spirâ elata, obtusiuscula; anfractibus 5 convexiusculis, lente accrescentibus, ultimo basi subplanato; aperturâ obliqua, semiovali, altiore quam lata; columella brevi, basi subtruncata, callum albido, linea subimpressa circumscriptum emitente; peristomate simplice, brevissimè reflexusculo, margine utroque leviter arcuato.*

Diam. 8, altit. $5\frac{1}{2}$ mill.

β . *Minor, testa saturatè carnea, supernè fasciis 2 angustis rubris et ad peripheriam 1 alba.*

γ . *Testa fulvo-rubella, supernè fasciis 2 angustis saturationibus.*

δ . *Minor, testa flavâ, supernè fasciis 2 angustis fulvis, ad peripheriam 1 albida ornata.*

From Cumana, Honduras (Dyson).

11. HELICINA EXIGUA, Pfr. *Hel. testa minutissima, conica, tenui, subtilissimè punctato-striatula, pellucidâ, pallide cornuta; spirâ conica, obtusiuscula; anfractibus 5 perconverxis, ultimo obsoletè angulata, basi planiusculo; aperturâ obliquâ, lunari; columella breviter recedente, callum exiguum emittente; peristomate simplice, tenui.*

Diam. $2\frac{1}{2}$, altit. 2 mill.

From Honduras (Dyson).

12. HELICINA FUNCKI, Pfr. *Hel. testa conico-subglobosa, tenuiuscula, sub lente tenuissimè obliquè striatula, vix nitidula, flavidâ, roseo-nebulosa; spirâ conoidea, obtusiuscula; anfractibus $5\frac{1}{2}$ planiusculis, ultimo utrinque convexiore, obsoletè angulata; aperturâ obliqua, semiovali; columella subarcuata, linea impressa verticali notata, basi subnodosa, in callum sensim tenuiorem retrorsum abiente; peristomate late expanso, margine supero subrepando.*

Diam. $13\frac{1}{2}$, altit. 9 mill.

From San Yago, New Granada (Funck).

13. HELICINA GONOCHEILA, Pfr. *Hel. testa conoideo-subglobosa, tenuiuscula, supernè striis spiralibus obsoletis sculpta et punctata, nitidula, fulvo-carnea; spirâ brevi, conoidea, subacuta; anfractibus $5\frac{1}{2}$ planiusculis, ultimo utrinque convexiore, obsoletè angulata; aperturâ obliqua, semiovali; columella subarcuata, linea impressa verticali notata, basi subnodosa, in callum sensim tenuiorem retrorsum abiente; peristomate late expanso, margine supero subrepando.*

No. CLXXXVIII.—PROCEEDINGS OF THE ZOOLOGICAL SOCIETY.

tibus 4½ vix convexiusculis, ultimo medio subcarinato, luteo-cingulato, basi convexiore, distinctius concentricè striato; apertura subobliqua, triangulari-semiovali, altiore quam latâ; columella subprecedente, supernè lineam impressam, brevem, curvatam emitente, basi acutè dentata; peristomate albo, rectangulè latè patente, margine basali substricto, cum columellâ angulum acutiusculum formante.

Diam. 10, altit. 6½ mill.
From Venezuela.

14. *HELICINA GOSSEI*, Pfr. *Hel. testâ depresso-globosâ, solida, pustulis seriatis subasperata, nitidula, fuscidulo-rubra; spirâ convexâ; anfractibus 4½ parum convexis, ultimo rotundato; apertura triangulare-semiovali, intus carnâ, nitidâ; columellâ basi truncata, retrorsum in callum crassum carneum abiente; peristomate subincassato, expanso, juxta columellam vix emarginato.*

Diam. 19, altit. 13½ mill.
From Jamaica (Gosse).

15. *HELICINA GULDINGIANA*, Pfr. *Hel. testâ depressâ, tenuiusculâ, sub lente subtilissimè granulatâ, diaphanâ, stramineâ vel albida, infra suturam fulvo-unifasciatâ; spirâ brevi, convexâ; anfractibus 4 vix convexiusculis, ultimo subdepresso, basi vix convexiore; apertura obliquâ, subtriangulari-semiovali; columellâ brevi, excavata, antrorsum in denticulum desinente, retrorsum in callum tenuem, semicircularem, flavescentem expansâ; peristomate tenui, breviter reflexo, margine supero repando, basali incisurâ levi à columellâ separato.*

Diam. 8, altit. 4½ mill.
Locality unknown.

16. *HELICINA HANLEYANA*, Pfr. *Hel. testâ globoso-conicâ, solida, lineis concentricis, impressis, subdistantibus sculptâ, vix diaphana, nitida, fulvo-carnâ; spirâ breviter conoidea, obtusiusculâ; anfractibus 5 vix convexiusculis, ultimo rotundato, anticè subdescendente; apertura parum obliquâ, subsemicirculari; columella brevissimâ, extrorsum denticulata, callum tenuem, albidum, diffusum emittente; peristomate albo, vix expansiusculo, intus subincassato, basi in denticulum columellæ abiente.*

Diam. 7½, altit. 5½ mill.
From New Orleans (Mr. Salle).

17. *HELICINA KIENERI*, Pfr. *Hel. testâ conoidea, tenuiusculâ, obliquè striata, lineis concentricis confertis subtilissimè decussata, albida, fusco-violaceo marmorata; spirâ convexo-conoidea, acuta; anfractibus 5½ vix convexiusculis, ultimo compressè carinato, basi convexiore; columellâ recedente, planata, supernè impressâ, basi incassata in callum basalem tenuem abiente; apertura obliquâ, integrâ, semiovali, altiore quam latâ; peristomate simplice, tenui, latè expanso.*

Diam. 16, altit. 11½ mill.
Locality unknown.

18. *HELICINA LINDENI*, Pfr. *Hel. testā globoso-conicā, tenuiusculd, subtilissimē striolatā et punctatā, subdiaphanā, pallidē stramineā vel carnea; spird conicā, acutiusculd; anfractibus 6 vix convexiusculis, ultimo inflato, obsoletē angulato; aperturā integrā, parum obliquā, semiovali, altiore quam latā; columellā leviter arcuatā, extrorsum in denticulum desinente, callum emittente exiguum, tenuem; peristomate breviter expanso, reflexiusculo.*

Diam. $11\frac{1}{2}$, altit. $8\frac{1}{3}$ mill.

From Tapinaba, Mexico (Linden).

19. *HELICINA ORBIGNYI*, Pfr. *Hel. testā depressā, sublenticulari, solidā, striatulā, vix nitidā, fusco-carnē; spird vix elatā; anfractibus $4\frac{1}{2}$ planiusculis, ultimo depresso, subangulato; aperturā obliquā, semiovali, altiore quam latā; columellā brevi, basi antrorsum dentatā, callum albidum, semicircularem retrorsum emitte; peristomate recto, subincrassato, juxta dentem columellæ non emarginato.*

Diam. $7\frac{1}{3}$, altit. 4 mill.

From the island of Cuba.

20. *HELICINA OWENIANA*, Pfr. *Hel. testā conicā, tenui, levigatā, sub lente lineolis impressis, antrorsum obliquis subtilissimē sculptā, nitidā, pellucidā, stramineā, sursum saturatiore; spird conicā, vertice obtusiusculo, castaneo; suturā linearī, albo-marginatā; anfractibus 6 planis, ultimo basi planiusculo; aperturā subobliquā, semiovali; columellā brevi, verticaliter rimatā, callum tenuissimum retrorsum emittente; peristomate aurantiaco, angulatim patente, reflexiusculo, margine utroque levissimē curvato.*

Diam. 9, altit. $7\frac{1}{2}$ mill.

From Chiapas, Mexico (Ghiesbreght).

21. *HELICINA PLICATULA*, Pfr. *Hel. testā depressē conoidea, solidulā, obliquē regulariter et elegantissimē plicatā, nitidā, cornē; spird brevi, conoidea, acutiusculd; anfractibus ferē 5 convexiusculis, ultimo supernē impresso, peripheriā obsoletissimē angulato; aperturā obliquā, semilunari; columellā brevissimā, simple, in callum tenuissimum diffusā; peristomate subincrassato, carneo, margine supero sinuato, basali juxta columellam subdentato.*

Diam. 5, altit. 3 mill.

From the island of Martinique.

22. *HELICINA REEVEANA*, Pfr. *Hel. testā conicā, solidulā, striis incrementi distinctis et lineolis obliquis impressis confertissimis sub lente clathratulā, nitidulā, albida, rufo nebulosā et tæniatā; spird elevatā, acutiusculd; suturā impressā; anfractibus 6 convexiusculis, ultimo angulato, basi vix convexiore; aperturā subsemiovali, intus castanē; columellā brevissimā, horizontaliter in callum parvulum, album, expansā; peristomate albo, angulatim patente, margine basali leviter arcuato, cum columellā extus subangulatim juncto.*

Diam. $8\frac{1}{2}$, altit. 6 mill.

From Cuba.

23. HELICINA ROHRI, Pfr. *Hel. testa conoidea, crassata, striatula et submalleata, opaca, vix nitidula, stramineo-albidata vel purpurea, albo-fasciata; spirae conoidea, acutiuscula; anfractibus 4½—5 planiusculis, ultimo supernè turgido, ad peripheriam carinata acutata, compressa, prominente munita, anticè deflexo, basi vix convexa; apertura obliqua, parvula, semiovali, altiore quam latata; columella subsimplice, basi obsoletè tuberculata, callum semicircularem album retrorsum emittente; peristome recto, acuto, intus crassè albolabiato, margine supero emarginato.*

Diam. 10, altit. 7 mill.

From the Marquesas Islands (Rohr).

24. HELICINA SANGUINEA, Pfr. *Hel. testa conoideo-orbiculata, crassa, punctato-striatula, opaca, sanguinea; spirae brevi, conoidea, acutiuscula; anfractibus 4½ planis, ultimo utrinque convexiusculo, medio subangulato; apertura obliqua, subtriangulari, altiore quam latata; columella basi antrorum dentata, callum tenuem, semicircularem retrorsum emittente; peristome recto, intus sublabiato, margine basali strictiusculo.*

Diam. 10½, altit. 6 mill.

Locality unknown.

25. HELICINA (TROCHATELLA) SEMILIRATA, Pfr. *Hel. testa conico-globosa, solidata, opaca, flavidata, supernè confertim albo-lirata; spirae conicae, acutiuscula; anfractibus 5½ planiusculis, ultimo convexiusculo, carinato, basi subtilissime concentricè striato; apertura perobliqua, subtriangulari; columella simplice, immediatè in marginem basalem abiente; peristome incrassato, angulatim expanso, marginibus callo tenuissimo junctis, supero sinuato.*

Diam. 10½, altit. 7½ mill.

From Venezuela (Linden).

26. HELICINA SOWERBIANA, Pfr. *Hel. testa depresso-trochiformis, tenuiuscula, lineis impressis spiraliter sulcatata, albata; spirae conicae, acutiuscula; anfractibus 6 planiusculis, ultimo subcarinato, basi convexiusculo; apertura parum obliqua, subtriangulari; columella tenui, basi nodiferata; peristome simplice, angulatim expanso, margine supero sinuato; callo basali tenuissimo.*

Diam. 21, altit. 14 mill.

From Guatimala (De Lattre).

27. HELICINA TENUILABRIS, Pfr. *Hel. testa subglobosa, solidata, sublævigata, alba et cinnamomeo variegata et subfasciata; spirae breviter conoidea, acutiuscula; anfractibus ferè 5 planiusculis, ultimo utrinque convexa, anticè vix descendente; apertura obliqua, semiovali, intus cerasinata, pallido-fasciata; columella recessente, angusta, retrorsum in callum tenuem dilatata, basi immediatè in peristoma tenue, expansiusculum, abiente.*

Diam. 10, altit. 7 mill.

Locality unknown.

28. HELICINA TENUIS, Pfr. *Hel. testa turbinata, tenuissima, vix*

striatula, pellucida, corneo-albidá, rubro obsoletè trifasciatâ; spirâ conicâ, acutâ; anfractibus 6 vix convexiusculis, ultimo basi planiusculo; aperturâ ferè verticali, triangulare-semiovali; columellâ brevi, basi retrorsum subdentata, superne in callum nitidum, circumscripsum, dilatata; peristomate tenui, angulatim expanso, margine basali cum columellâ basi angulum formante.

Diam. 11, altit. $8\frac{1}{2}$ mill.

From Yucatan.

29. *HELICINA UNIDENTATA*, Pfr. *Hel. testâ depressâ, tenuiusculâ, liris concentricis alternatim validis, obtusis et minoribus cinctâ, diaphanâ, nitidulâ, rubella; spirâ vix elevatâ; anfractibus $4\frac{1}{2}$ depresso, ultimo anticè descendente, basi medio profundè excavato; aperturâ perobliquâ, latè lunari; columellâ simplice, retrorsum in callum albidum circumscripsum dilatata; peristomate expanso, intus albo-labiato, margine basali prope columellam dente magno, prominente, instructo.*

Diam. 5, altit. $2\frac{1}{2}$ mill.

From Honduras (Dyson).

4. DESCRIPTION OF A NEW SPECIES OF PARROT.

BY G. R. GRAY, Esq., F.L.S. ETC.

(Aves, pl. 5.)

PSITTACUS RÜPPELLII.

Uniform dark bronze colour, with the lesser and under wing-coverts bright yellow; the feathers of the thighs orange-yellow.

Total length 9 inches 6 lines; bill, from gape, 10 lines; wings 5 inches 6 lines; tail 3 inches 3 lines; tarsi 6 lines.

The greater uniformity of colour at once distinguishes it from the allied species, *Psittacus Meyeri* and *P. rufiventris* of Dr. Rüppell.

The specimen from which this description is taken lived for upwards of twelve months in the Society's collection, and is believed to have been brought to this country from the river Nunez. I have named it in honour of my distinguished friend, whose labours have contributed so largely to our knowledge of African zoology.

The Meeting was adjourned by the Chairman until Tuesday, November 14th.