

# PROCEEDINGS

OF THE

## ZOOLOGICAL SOCIETY

OF LONDON.



---

PART XXIV.

1856.

---

PRINTED FOR THE SOCIETY ;  
SOLD AT THEIR HOUSE IN HANOVER-SQUARE,  
AND BY MESSRS. LONGMAN, BROWN, GREEN AND LONGMANS,  
PATERNOSTER-ROW.

*horizontaliter patente, concentrice striato, castaneo maculato, superne producto, ad anfr. penultimum subexciso.—Operc. albidum.*

Long. 13–15½, diam. 6–7½ mill.

Hab. Cordova.

#### DESCRIPTION OF PLATE XXXV.

- |      |             |   |
|------|-------------|---|
| Fig. | 1.          | Cylindrella Boucardi, <i>Pfr.</i>           |
|      | 2, 3.       | — polygyra, <i>Pfr.</i>                     |
|      | 4, 5.       | — apistoma, <i>Pfr.</i>                     |
|      | 6.          | Achatina (Varicella) Orizabæ, <i>Pfr.</i>   |
|      | 7.          | (Varicella) speciosa, <i>Pfr.</i>           |
|      | 8.          | Spiraxis Shuttleworthi, <i>Pfr.</i>         |
|      | 9.          | — turgidula, <i>Pfr.</i>                    |
|      | 10.         | Bulimus aurifluus, <i>Pfr.</i>              |
|      | 11.         | — sulphureus, <i>Pfr.</i>                   |
|      | 12.         | — Drouëti, <i>Pfr.</i>                      |
|      | 13, 14.     | Helix caduca and var., <i>Pfr.</i>          |
|      | 15, 16.     | Simpulopsis Salleana, <i>Pfr.</i>           |
|      | 17.         | Helicina Heloisæ, <i>Sallé.</i>             |
|      | 18, 19, 20. | — notata and var., <i>Pfr.</i>              |
|      | 21, 22.     | Proserpina (Ceres) Salleana, <i>Cuming.</i> |
|      | 23, 24.     | — eolina, <i>Ductos</i> (for comparison).   |
|      | 25.         | Cyclophorus Boucardi, <i>Sallé.</i>         |

#### 5. DESCRIPTIONS OF FIFTY-EIGHT NEW SPECIES OF HELICEA FROM THE COLLECTION OF H. CUMING, ESQ. BY DR. L. PFEIFFER.

1. VITRINA FLEMINGI, *Pfr.* *V. testa subdepressa, peripheria auriformis; solidula, superne plicato-striata striisque spiralibus sub lente notata, cæneo-micante, olivaceo-fulva; spira parum elata; sutura anguste albumarginata; anfr. 4½ convexiusculis, ultimo magno, infra medium obsoletissime angulato, basi levigato, nitidiore; apertura diagonali, lunato-ovali, intus margaritacea; perist. simplice, margine dextro subrepando, columellari arcuato, superne triangulatim reflexo, adnato.*

Diam. maj. 33, min. 24, alt. 17–18 mill.

Hab. Scinde, India (*Dr. Alex. Fleming*).

2. VITRINA PLANTI, *Pfr.* *V. testa deppressa, ambitu subauriformi, tenuissima, levigata, nitidissima, pellucida, albido-hyalina; spira parvula, plana; sutura vix impressa, submarginata; anfr. vix 3 planiusculis, ultimo magno, superne vix convexiore, ad suturam striatulo, subtus inflato, membranaceo-marginato; apertura magna, diagonali, lunato-ovali; perist. simplice, margine dextro antrorsum arcuato, columellari tenuissima.*

Diam. maj. 12, min. 8, alt. 6 mill.

Hab. Natal (*Mr. Plant*).

3. VITRINA BORNEENSIS, *Pfr.* *V. testa depresso-globosa, tenuissima, striatula, striis spiralibus nonnullis impressis notata, pellu-*

*cida, pallide cornea; spira brevissime conoidea, albida; sutura submarginata; anfr. 4 celeriter accrescentibus, superis vix convexis, ultimo subdepresso-rotundato, basi oblique ruguloso; apertura obliqua, lunato-rotundata; perist. simplice, levissime inflexo, margine columellari brevi, strictiusculo, filari.*

Diam. maj. 15, min.  $11\frac{2}{3}$ , alt. 9 mill.

Hab. Borneo.

4. **VITRINA CELEBENSIS**, Pfr. *V. testa helicoidea, solidula, striatula, nitida, coffeacea; spira breviter conoidea, obtusula; sutura pallide marginata; anfr. 4 convexiusculis, sensim accrescentibus, ultimo basi parum convexo, pallido; apertura diagonali, lunato-rotundata; perist. simplice, recto, marginibus subconvergentibus, columellari arcuato, subcalloso.*

Diam. maj.  $11\frac{1}{2}$ , min. 9, alt. 7 mill.

Hab. Celebes (Mrs. Ida Pfeiffer).

5. **VITRINA IDÆ**, Pfr. *V. testa depresso-globosa, ambitu subovali, tenui, arcuato-striatula, nitida, pellucida, lutescenti-cornea; spira vix elevata; sutura levi; anfr. 4 convexiusculis, ultimo magno, inflato, basi non impresso; apertura diagonali, rotundato-lunari; perist. simplice, marginibus conniventibus, regulariter arcuatis, columellari brevissime recedente, subcalloso.*

Diam. maj. 14, min. 11, alt.  $7\frac{2}{3}$  mill.

Hab. Celebes (Mrs. Ida Pfeiffer).

6. **VITRINA ? COMORENSIS**, Pfr. *V. testa subglobosa, tenui, rugoso-striata, vix nitidula, diaphana, virenti-cornea, punctis et strigis luteis variegata; spira convessa, obtusa; anfr. 3 convexis, rapide accrescentibus, ultimo inflato; apertura fere diagonali, lunato-ovali, intus margaritacea, albo variegata; perist. simplice, recto, marginibus convergentibus, regulariter arcuatis, columellari filari.*

Diam. maj.  $21\frac{1}{2}$ , min.  $16\frac{1}{2}$ , alt. 13–14 mill.

Hab. Mayotte, Comoro Islands (Mr. Clouë).

Shape very similar to that of *Helix aperta*, Born.

7. **VITRINA DARNAUDI**, Pfr. *V. testa semiovata, pertenui, sublente irregulariter striatula punctisque impressis notata, oleosomicante, pallide virenti-cornea; spira convessa, obtusa; sutura anguste marginata; anfr. vix 3 convexis, rapide accrescentibus, ultimo magno; apertura subdiagonali, sublunato-ovali, intus leviter margaritacea; perist. simplice, brevissime inflexo, marginibus approximatis, dextro antrorum dilatato, columellari arcuato, anguste membranaceo-marginato.*

Diam. maj. 16, min. 12, alt.  $7\frac{1}{2}$  mill.

Hab. Sennaar, Interior of Africa (Mr. Darnaud).

8. **VITRINA SENNAARIENSIS**, Pfr. *V. testa depressa, tenui, oblique rugosa, nitidula, pellucida, corneo-virente, plerumque luteo*

*nigro obducta ; spira subturbinata, mucronata ; anfr. 3 convexis, sensim accrescentibus, ultimo depresso, peripheria obsolete ungu-lato ; apertura perobliqua, lunato-circuluri ; perist. tenui, subin-flexo, marginibus convergentibus, columellari vix dilatato.*

Diam. maj.  $6\frac{2}{3}$ , min.  $5\frac{1}{3}$ , alt. 3 mill.

Hab. Sennaar, Interior of Africa (Mr. Darnaud).

9. **VITRINA LEUCOSPIRA**, Pfr. *V. testa depresse subglobosa, am-bitu ovali, tenuissima, striatula, pellucida, nitida, lutescenti-hy-a-lina ; spira parum convexa, alba ; sutura vix submarginata ; anfr. 4 convexiusculis, penultimo sensim pellucido-radiato, ultimo ro-tunduto ; apertura obliqua, lunato-subcirculari ; perist. tenui, marginibus conniventibus, subinflexis, dextro antrorum valde ar-cuato, columellari substricto, brevi, filari.*

Diam. maj.  $13\frac{1}{2}$ , min.  $11\frac{1}{2}$ , alt.  $8\frac{1}{2}$  mill.

Hab. Australia.

10. **SUCCINEA PLANTI**, Pfr. *S. testa conico-ovata, tenuiuscula, rugulosa, diaphana, sordide cornea ; spira brevi, subpapillata ; anfr. vix  $2\frac{1}{2}$ , penultimo conveexo, ultimo  $\frac{2}{3}$  longitudinis superante, medio turgido ; columella substricte recedente, superne levissime calloso-plicata ; apertura fere diagonali, ubique incumbente, acu-minato-ovali ; perist. simplice, margine dextro regulariter arcuato.*

Long. 5, diam.  $3\frac{1}{2}$ , alt.  $2\frac{2}{3}$  mill.

Hab. Cape Natal (Mr. Plant).

11. **SUCCINEA ASPERULA**, Pfr. *S. testa ovato-conica, tenuiuscula, striis rugosis subasperata, diaphana, corneo-rubella ; spira turbi-nata, acuta ; anfr.  $3\frac{1}{2}$  convexis, ultimo ventroso,  $\frac{2}{3}$  longitudinis formante ; columella leviter arcuata, compressa, albida ; apertura oblique, undique incumbente, subregulariter ovali, intus marg..ri-tacea ; perist. simplice, marginibus subsymmetricis.*

Long.  $10\frac{2}{3}$ , diam.  $6\frac{1}{3}$ , alt. 5 mill.

Hab. Flagstaff Hill, St. Helena.

12. **HELIX KERMANDECI**, Pfr. *H. testa imperforata, turbinata, tenui, superne subdistanter striata, hyalina ; spiru convexiusculo-turbinata, subacuminata ; anfr.  $5\frac{1}{2}$ -6 convexis, lente accrescenti-bus, ultimo subangulato, antice rotundato ; apertura vix obliqua, lunari ; perist. simplice, recto, marginibus distantibus, columellari superne subreflexo, adnato.*

Diam. maj.  $3\frac{2}{3}$ , min.  $3\frac{1}{3}$ , alt.  $2\frac{1}{2}$  mill.

Hab. Sunday Island, Kermandee Group (Lieut. Chimmo, R.N.).

13. **HELIX OPHIRIA**, Pfr. *H. testa subperforata, depresso turbinato-globosa, tenuissima, membranacea, radiato-rugata et lineis spirali-bus minutissimis sub lente sculpta, pellucida, virenti-cornea ; spira subconoidea, superne alba ; anfr. 5 convexiusculis, sensim accrescentibus, ultimo peripheria subangulato, antice rotundato, busi inflato ; apertura fere diagonali, rotundato-lunuri ; perist.*

*simplice, marginibus dextro et basali subinflexis, columellari fere verticali, superne subreflexo.*

Diam. maj. 20, min. 17, alt.  $11\frac{1}{2}$  mill.

Hab. Mount Ophir, Malacca (Dr. Traill).

*Vitrina heliciformis*, Pfr. 1854, is an imperfect form of this species.

14. **HELIX SHIPLAYI**, Pfr. *H. testa perforata, subturbanata, solida; superne arcuato-plicata striisque spiralibus eleganter granulata, isabellina; spira convexiusculo-conica, obtusula; sutura subcanaliculata; anfr. 6 convexis, lente accrescentibus, ultimo peripheria carina acuta, compressa, antice evanescente munito, basi convexo, levicre; apertura diagonali, subangulato-lunari, intus margaritacea; perist. simplice, obtusulo, margine columellari fere verticali, superne triangulatim reflexo.*

Diam. maj. 20, min. 18, alt.  $11\frac{1}{2}$  mill.

Hab. Neilgherries, India (Mr. Conway Shipplay).

15. **HELIX ACALLES**, Pfr. *H. testa vix perforata, turbinata, tenui, oblique rugosa, diaphana, pallide cornea; spira conoidea, acutiuscula; anfr.  $5\frac{1}{2}$  convexiusculis, sensim accrescentibus, ultimo peripheria carinato, antice non descendente, basi convexiore; apertura obliqua, subangulato-lunari; perist. simplice, recto, margine columellari superne anguste reflexo.*

Diam. maj. 13, min. 11, alt.  $7\frac{1}{2}$  mill.

Hab. Neilgherries, India (Mr. Conway Shipplay).

16. **HELIX CALABARICA**, Pfr. *H. testa perforata, turbinato-lentiformi, tenui, levigata, superne liris 6 argutis, filiformibus cincta, diaphana, cornea; spira conoidea, acutiuscula; anfr. 6 convexiusculis, sensim accrescentibus, ultimo non descendente, acute carinato, basi convexo; apertura parum obliqua, angulato-lunari; perist. simplice, recto, marginibus vix convergentibus, columellari arcuato, superne vix reflexo.*

Diam. maj.  $9\frac{1}{2}$ , min.  $8\frac{1}{3}$ , alt.  $4\frac{1}{2}$  mill.

Hab. Old Calabar, Guinea.

17. **HELIX DARNAUDI**, Pfr. *H. testa perforata, depresso conoideo-globosa, tenui, rugoso-striata, cornea, fasciis opacis albidis notata; spira conoidea; anfr. fere 5 vix convexiusculis, sensim accrescentibus, ultimo non descendente, rotundato; apertura vix obliqua, rotundato-lunari; perist. simplice, recto, marginibus subconvergentibus, columellari superne dilatato, patente.*

Diam. maj. 8, min. 7, alt. 5 mill.

Hab. Sennaar, Interior of Africa (Mr. Darnaud).

18. **HELIX ARGUTA**, Pfr. *H. testa subclause perforata, depressa, tenui, superne argente et confertim arcuato-striata, corneo-fusca; spira parum elevata, obtusa; sutura subcrenulata; anfr.  $4\frac{1}{2}$  celester accrescentibus, superis vix convexiusculis, ultimo peripheria carina fusiformi cincto, antice non descendente, basi levior, luteo-virente, medio excavato; apertura diagonali, ampla, lunato-ro-*

*tundata, intus margaritacea; perist. simplice, recto, marginibus conniventibus, columellari superne in laminam brevem callosam reflexo.*

Diam. maj. 36, min. 29, alt. 18 mill.

*Hab. Tenga Hills, Java (C. Shipley, Esq.).*

19. **HELIX CHIMMOI**, Pfr. *H. testa umbilicata, convexo-depressa, tenuiuscula, confertim plicato-striata, cornea, rufo irregulariter maculata; spira parum elevata; anfr. 5 convexis, sensim accrescentibus, ultimo non descendentे, subdepresso-rotundato; umbilico conico,  $\frac{1}{3}$  diametri subaequante; apertura parum obliqua, rotundato-lunari; perist. simplice, recto, marginibus convergentibus.*

Diam. maj.  $3\frac{2}{3}$ , min. 3, alt.  $1\frac{1}{2}$  mill.

*Hab. Sunday Island, Kermandee Group (Lieut. Chimmo, R.N.).*

20. **HELIX CONFERTA**, Pfr. *H. testa umbilicata, conoidea, solida, striis incrementi irregularibus et confertissimis spiralibus decussatula, sericea, fulvida; spira conoidea, obtusa; sutura pallide marginata; anfr. vix 5 convexiusculis, sensim accrescentibus, ultimo peripheria angulato, ad suturam turgidulo, basi convexo, juxta umbilicum angustum compresso; apertura perobliqua, subtetragono-lunuri; perist. obtuso, margine supero recto, basali incrassato, reflexiusculo, columellari declivi, subdentato, superne triangulatum reflexo.*

Diam. maj.  $34\frac{1}{2}$ , min. 30, alt. 20 mill.

*Hab. — ?*

21. **HELIX DAMAHOYI**, Pfr. *H. testa imperforata, globoso-depressa, tenuiuscula, oblique striata, parum nitente, saturate fusca, fasciis latis nigris, unica lutea mediana et superpositis non-nullis hydrophanis albidis ornata; spira vix elevata, obtusa; anfr. 4 convexiusculis, rapide accrescentibus, ultimo inflato, oblique malleato-plicato; columella declivi, compressa, dilatata, alba; apertura obliqua, truncato-ovali, intus albida; perist. albo, late expanso et reflexiusculo.*

Diam. maj. 47, min. 37, alt. 28 mill.

*Hab. Philippine Islands (Mr. Damahoy).*

22. **HELIX MEOBAMBENSIS**, Pfr. *H. testa obtecte umbilicata, conoideo-depressa, solida, subdistanter rugoso-striata, undique minute granulata, cinnamomea; spira conoideo-convexa, obtusa; anfr. 5 planiusculis, lente accrescentibus, ultimo antice descendente, peripheria subcarinato, utrinque convextiore; apertura perobliqua, tetragono-lunari; perist. albo, reflexo, marginibus callo junctis, columellari dilatato, adnato, stricto, ad dextram unidentato.*

Diam. maj. 32, min. 26, alt. 15 mill.

*Hab. Meobamba, Peru (Mr. Porte).*

23. **HELIX AMMIRALIS**, Pfr. *H. testa umbilicata, turbinato-de-*

*pressa, solidula, undique subruditer striata, fulva vel lutescente, plerunque fascia unica peripherica nigro-castanea ornata; spira regulariter conoidea; anfr. 6½ vix convexiusculis, lente accrescentibus, ultimo antice vix descendente, peripheria carinato, basi convexo, circa umbilicum angustum, castaneum subcompresso; apertura obliqua, lunari; perist. hepatico, breviter expanso, margine columellari superne fornicatim reflexo.*

Diam. maj. 36, min. 33, alt. 20 mill.

Hab. China (Admiral Cecille).

24. **HELIX MUCIDA**, Pfr. *H. testa umbilicata, turbinato-depressa, tenuiuscula, striatula, saturate rufa, quasi mucore obducta; spira conoidea, obtusula; anfr. 5 convexis, sensim accrescentibus, ultimo rotundato, antice descendente, circa umbilicum infundibuliformem subangulata; apertura fere diagonali, rotundato-lunari, intus nitida, carnea; perist. breviter expanso, marginibus vix convergentibus, columellari superne triangulatum dilatato, patente.*

Diam. maj. 20, min. 16½, alt. 11 mill.

Hab. Percy's Island (Lieut. Chimmo, R.N.).

25. **HELIX GUEINZII**, Pfr. *H. testa umbilicata, depressa, carinata, tenui, plicis subdistantibus retrorsum descendantibus sculpta, pellucida, pallide cornea, seriebus macularum rufarum ornata; spira vix elevata; anfr. 5 sensim accrescentibus, prope suturam turgidis, ultimo superne subacute carinato, antice non descendente, subtus per inflato, circa umbilicum ( $\frac{1}{6}$  diametri occupantem) conicum levigato, subangulata; apertura vix obliqua, subangulato-lunari; perist. tenui, undique breviter expanso, margine columellari superne dilatato, patente.*

Diam. maj. 20, min. 16, alt. 9 mill.

Hab. Meobamba, Peru (Mr. Gueinzius).

26. **HELIX BASIDENTATA**, Pfr. *H. testa umbilicata, conoideo-subsemiglobosa, solida, oblique striata, alba, fasciis 2 castaneis et 1 angustiore aurantiaca prope suturam ornata; spira convexo-conoidea, obtusa; anfr. 5 convexis, sensim accrescentibus, ultimo carinato, antice perdeflexo, basi planiusculo; umbilico angusto, pervio; apertura horizontali, elliptica; perist. continuo, undique reflexo, margine basali medio dente 1 valido, obtuso armato.*

Diam. maj. 26, min. 20, alt. 11 mill.

Hab. Philippine Islands.

27. **BULIMUS PHÆOSTYLOS**, Pfr. *B. testa imperforata, oblongo-ovata, tenui, oblique confertim striata, pallide fulvescenti-carnea, fasciis 3 nigricantibus (1 suturali, 1 infraperipherica, tertia columellari) ornata; spira convexo-conica, apice obtusula, nigro-violacea; anfr. 5 modice convexis, sensim accrescentibus, ultimo spira paulo breviore; columella compressa, plana, fusca, basi*

*subtruncata ; apertura obliqua, truncato-oblonga ; perist. castaneo, anguste reflexo.*

Long. 37, diam. 24 mill.

Hab. Philippine Islands.

28. **BULIMUS INÆQUALIS**, Pfr. *B. testa anguste umbilicata, oblique fusiformi, tenuiuscula, lœvigate, nitida, alba unicolo re vel strigis latis ramosis violaceo-nigricantibus picta ; spira gracili, conica, acuta ; anfr. 6 vix convexiusculis, ultimo spira longiore, antice subascendente, latere aperturali planulato, basi subattenuato ; apertura subverticali, elongato-auriformi, intus violaceo-limbata ; columella violacea, superne oblique plicata ; perist. albo, late expanso, margine columellari plano, patente, superne flexuoso.*

Long. 43, diam. 13 mill.

Hab. Banks of the Maranhon.

29. **BULIMUS GUEINZII**, Pfr. *B. testa profunde rimata, oblongo-conica, tenui, sublœvigate, nitida, albida, strigis angustis fuscis, basin versus undulatis, ornata ; spira conica, acutiuscula ; anfr. 6 convexiusculis, ultimo spira vix breviore, basi subattenuato, ad rimam violaceo ; columella compressa, subtorta, violacea ; apertura vix obliqua, truncato-oblonga, intus lilacina ; perist. tenui, margine dextro superne valde curvato, late expanso, columellari patente.*

Long. 23, diam. 10 mill.

Hab. Meobamba, Peru (*Mr. Gueinzius*).

30. **BULIMUS CLARUS**, Pfr. *B. testa rimato-umbilicata, ovato-oblonga, tenui, lœvigate, nitida, diaphana, sub epidermide fugace fulvescente-albida ; spira elongata, convexo-conica, obtusula ; anfr. 5½ modice convexis, ultimo ⅔ longitudinis subæquante, basi rotundato ; columella leviter arcuata ; apertura obliqua, truncato-ovali ; perist. tenui, marginibus conniventibus, dextro breviter expanso, columellari dilatato, patente.*

Long. 16, diam. 7½ mill.

Hab. Meobamba, Peru (*Mr. Gueinzius*).

31. **BULIMUS LACTIFLUUS**, Pfr. *B. testa breviter rimato-perforata, fusiformi-turrita, tenuiuscula, sublœvigate, cornea, strigis lacteis subserratis notata ; spira turrita, apice acutiuscula, sœpe nigro-cornæa ; anfr. 10-11 convexiusculis, ultimo ⅔ longitudinis subæquante, basi attenuato, compresso ; columella leviter arcuata ; apertura vix obliqua, ovali-oblonga ; perist. tenui, margine dextro anguste expanso, columellari a basi dilatato, fornicatim reflexo.*

Long. 16½-17, diam. 4½ mill.

Hab. Chili.

32. **BULIMUS FLORIDANUS**, Pfr. *B. testa anguste perforata, ovato-turrita, sublœvigate, griseo-hyalina, strigis et maculis opacis, albis notata ; spira elongato-conica, acutiuscula ; anfr. 6½ convexiusculis, superis interrupte fusco-fasciatis, ultimo ⅔ longi-*

*tudinis subæquante, infra medium subangulato, basi attenuato; columella subtorta, recedente; apertura parum obliqua, ovali; perist. tenui, margine dextro anguste expanso, columellari dilatato, reflexo, fere adnato.*

Long.  $15\frac{2}{3}$ –17, diam.  $7\frac{1}{2}$  mill.

Hab. Florida.

33. **BULIMUS DISTANS**, Pfr. *B. testa compresse umbilicata, ovato-conica, tenuiuscula, costis subarcuatis, chordæformibus subdistantibus sculpta, subdiaphana, albida; spira elevato-conica, obtusula; anfr. 7 convexis, ultimo  $\frac{3}{7}$  longitudinis subæquante, basi rotundato; columella profunde subtorta; apertura parum obliqua, acuminato-subovali; perist. tenui, marginibus convergentibus, dextro breviter expanso, columellari dilatato, patente.*

Long.  $22\frac{1}{2}$ , diam.  $10\frac{1}{2}$  mill.

Hab. Isle of Karah, Gulf of Persia.

34. **BULIMUS ÆSTIVUS**, Pfr. *B. testa umbilicata, oblongo-turrita, solidula, irregulariter striata, albida; spira elongata, convexiusculo-conica, acutiuscula; anfr. 6 modice convexis, ultimo  $\frac{3}{7}$  longitudinis subæquante, basi vix attenuato; columella subrecedente; apertura parum obliqua, acuminato-ovali; perist. tenui, margine dextro anguste expanso, columellari dilatato, fornicatim reflexo.*

Long. 17, diam.  $7\frac{1}{3}$  mill.

Hab. Meobamba, Peru (Mr. Gueinzius).

35. **BULIMUS CHARBONNIERI**, Pfr. *B. testa profunde rimata, ovato-conica, tenuiuscula, confertim costulato-striata, diaphana, corneo-albida; spira elevato-conica, obtusa; anfr. 6 convexiusculis, ultimo spira paulo breviore, juxta umbilicum rimæformem subcompresso; columella substicta; apertura parum obliqua, elliptico-ovali; perist. tenui, marginibus approximatis, dextro perarcuato, breviter expanso, columellari sursum dilatato, patente.*

Long. 15, diam. 8 mill.

Hab. Isle of Karah, Gulf of Persia.

36. **BULIMUS TIBETANUS**, Pfr. *B. testa profunde rimata, ovato-oblonga, pupæformi, solidula, striatula, albida, strigis et maculis corneis irregulariter notata; spira subcylindrica, sensim in conum obtusulum attenuata; anfr. 8– $8\frac{1}{2}$  vix convexiusculis, ad suturam plicatulis, ultimo  $\frac{1}{3}$  longitudinis subæquante, antice ascendentem, busi compresso; columella subuplicata; apertura verticali, sinuato-ovali; perist. albolabiato, expanso, marginibus callo subjunctis, columellari reflexiusculo.*

Long. 32, diam. 11 mill.

Hab. Tibet.

37. **BULIMUS COCHINCHINENSIS**, Pfr. *B. testa imperforata, fusiformi-ovata, solida, lœvigata, nitida, pallide sulphurea vel albida; spira convexiusculo-turrita, apice obtusula; sutura levi, pallida; anfr. 6–7, superis planiusculis, sequentibus convexiori-*

*bus, ultimo  $\frac{2}{5}$  longitudinis subæquante, basi attenuato, subcompresso; columella uniformi, leviter torta; apertura parum obliqua, elliptico-ovali; perist. subincrassato, marginc dextro breviter expanso, columellari dilatato, adnato.*

Long. 39, diam. 17 mill.

Hab. Cochin China.

38. **BULIMUS CHION**, Pfr. *B. testa perforata, oblonga, solida, striatula, alba; spira elongata, in conum acutiusculum terminata; anfr. 7-8 modice convexis, ultimo  $\frac{1}{3}$  longitudinis paulo superante, antice subascendente, basi rotundato; apertura verticali, ovato-lunari; perist. calloso, marginibus callo junctis, columellari brevi, substricto, dilatato, patente.*

Long. 12, diam.  $5\frac{1}{2}$  mill.

Hab. India; Punjab, Kurrachee, mouth of the Indus (Mr. Conway Shipley).

39. **BULIMUS KANAIENSIS**, Pfr. *B. testa subperforata, conico-ovata, solidula, striatula et irregulariter malleato-impressa, alba; spira conica, sursum interdum grisea, apice obtusa; anfr. 5 convexis, ultimo spiram vix superante, oblique descendente, basi rotundata; columella vix arcuata; apertura obliqua, truncato-ovali; perist. simplice, recto, margine columellari dilatato, reflexo, subappresso.*

Long. 14, diam. 8 mill.

Hab. Kanai, Sandwich Islands.

40. **BULIMUS LORRAINI**, Pfr. *B. testa subperforata, ovato-conica, tenui, rugata, subepidermide fulvida, glutinosa, alba; spira convexiusculo-conica, obtusula; anfr. 4 convexis, ultimo spiram paulo superante, basi parum attenuato; columella compressa, leviter arcuato; apertura obliqua, acuminato-ovali, intus margaritaceo-albida; perist. simplice, recto, margine columellari anguste reflexo, subadnato.*

Long. 20, diam. 9 mill.

Hab. Isle of Penang (Mr. Lorrain).

41. **BULIMUS WOODWARDI**, Pfr. *B. testa perforata, ovato-turrita, solidula, striata et submalleata, nitidula, fulvido-alba; spira elongato-conica, acutiuscula; anfr. 8 convexiusculis, ultimo spira paulo breviore, basi subattenuato; columella recedente; apertura obliqua, ovali-oblonga; perist. simplice, recto, margine dextro leviter arcuato, columellari sursum dilatato, fornicatim reflexo, perforationem angustum semitegente.*

Long. 31, diam.  $13\frac{1}{2}$  mill.

Hab. Andes of Peru.

42. **BULIMUS CASTELNEAUI**, Pfr. *B. testa subperforata, fusiformi-turrita, tenuiuscula, striatula (sub lente decussatula), nitida, griseo-albida, punctis rarissimis pellucidis conspersa; spira elongato-conica, apice acuta, cornea; anfr.  $7\frac{1}{2}$  convexiusculis, ultimo  $\frac{2}{5}$  lon-*

*gitudinis subæquante, basi attenuato; columella substricta; apertura obliqua, oblonga, intus carnea; perist. simplice, recto, margine columellari sursum dilatato, fornicato-reflexo.*

Long. 20, diam.  $8\frac{1}{2}$  mill.

Hab. Rio Pampas, Bolivia (Mr. Castelneau).

43. **BULIMUS NIGROAPICATUS**, Pfr. *B. testa perforata, ovato-conica, tenuiuscula, striata, nitida, albida, fasciis nigricantibus crebris, superioribus moniliformibus, ornata; spira conica, apice acutiuscula, nigra; anfr.  $5\frac{1}{2}$  parum convexis, ultimo spira paulo langiore, ventroso; columella levissime arcuata; apertura obliqua, elliptico-ovali; perist. simplice, recto, murgine columellari superne late reflexo, subappresso.*

Long. 22, diam.  $11\frac{1}{2}$  mill.

Hab. Rio Pampas, Bolivia (Mr. Castelneau).

44. **BULIMUS STENACME**, Pfr. *B. testa umbilicata, ovato-turrita, solidula, ruguloso-striata, albida, strigis angustis pallide corneis variegata; spira elongata, apicem versus acutiusculum attenuata; anfr. 7 convexiusculis, ultimo  $\frac{2}{5}$  longitudinis subæquante, juxta umbilicum angustum subcompresso; apertura obliqua, oblongo-ovali, intus fusco-carnea; perist. simplice, recto, margine columellari superne dilatato, fornicatum reflexo.*

Long.  $20\frac{1}{2}$ , diam. 9 mill.

Hab. Bolivia.

45. **BULIMUS MONACHUS**, Pfr. *B. testa anguste perforata, oblongo-turrita, tenui, striatula, diaphana, sordide cornea; spira convexo-turrita, apice acutiuscula; anfr.  $7\frac{1}{2}$  vix convexiusculis, ultimo  $\frac{3}{4}$  longitudinis subæquante, basi parum attenuato, rotundato; columella recedente; apertura obliqua; oblongo-ovali; perist. simplice, recto, margine columellari sursum dilatato, reflexo.*

Long. 31, diam.  $11\frac{1}{2}$  mill.

Hab. Meobamba, Peru (Mr. Gueinzii).

46. **PARTULA PURPURASCENS**, Pfr. *P. testa perforata, ovato-conica, solidu, oblique striata striisque spiralibus confertis distinete decussata, nitida, purpurascenti-fusca; spira convexo-conica, acuta; sutura levi; anfr.  $5\frac{1}{2}$  vix convexiusculis, ultimo spiram subæquante, peripheria subangulato, basi rotundato; columella leviter arcuata; apertura parum obliqua, truncato-oblonga; perist. fusco-violaceo, undique patente et reflexiusculo.*

Long. 22, diam. 14 mill.

Hab. —?

47. **PARTULA CALLIFERA**, Pfr. *P. testa umbilicata, ovato-conica, solida, sublævigata (sub lente punctulato-striata), albida; spira convexiusculo-conica, obtusula; sutura levi; anfr. 5 convexiusculis, ultimo spira paulo longiore, inflato; columella subverticali, superne tuberculifera, intus plicata; apertura vix obliqua, sinuato-oblonga, dente profundo parietali coarctata; perist. incrassato,*

*dilatato, marginibus callo junctis, dextro supra medium callo oblongo intus munito.*

Long. 19, diam. 11 mill.

Hab. — ?

48. PARTULA LÆVIGATA, Pfr. *P. testa profunde rimata, ovato-conica, solida, lævigata (sub lente vix striatula), nitida, lutescente; spira conica, obtusula; sutura mediocri; anfr. 5 convexiusculis, ultimo spiram subæquante, prope sutaram tumidiore, basi rotundato; columella supra medium introrsum nodoso-plicata; apertura vix obliqua, oblonga, tuberculo profundo anfractus penultimi coarctato; perist. crasso, albo, undique patente, margine columellari superne dilatato, adnato.*

Long. 20, diam. 10 mill.

Hab. — ?

49. PARTULA LILACINA, Pfr. *P. testa subobtecte perforata, ovato-conica, solidu, sub lente spiraliter striata, nitidula, lilacina; spira conica, acuta; sutura levi; unfr.  $5\frac{1}{2}$  vix convexiusculis, ultimo spiram subæquante, rotundato; columella superne tuberculo subcircumscripto munita; apertura obliqua, truncato-oblonga; perist. incrassato, albido, breviter expanso, margine columellari fere adnato.*

Long.  $17\frac{1}{2}$ , diam. 10 mill.

Hab. Marquesas Islands.

50. ACHATINELLA (ACHATINELLASTRUM) OVUM, Pfr. *A. testa sinistrorsu, imperforata, globoso-conica, solida, rugulosa-striata, nitidala, albida; spira concaviusculo-conica, apice acuta; sutura profunde marginata; anfr.  $5\frac{1}{2}$ , superis planis, sequentibus convexis, ultimo inflato, spira vix breviore; plica columellari crassa, tuberculiformi, pallide lilacea; apertura diagonali, sinuato-semicirculari; perist. recto, nigro-fusco, limbato, intus crenato-labiato.*

Long.  $19\frac{1}{2}$ , diam. 13 mill.

Hab. Oahu, Sandwich Islands (Dr. Newcomb).

51. ACHATINELLA (LAMINELLA) FARCIMEN, Pfr. *A. testa sinistrorsa, subperforata, oblongo-conica, solidula, rugulosa, sub epidermide nigricante albida; spira superne in conum acutum nudum terminata; anfr. 7, superis planis, 2 ultimis perconvexis, ultimo  $\frac{1}{3}$  longitudinis paulo superante, medio subangulato; plica columellari compressa, obliqua; apertura obliqua, semiovuli, intus alba; perist. simplice, recto, margine columellari subpatente.*

Long. 19, diam. 9 mill.

Hab. Mani, Sandwich Islands (Dr. Newcomb).

52. ACHATINELLA (LABIELLA) CALLOSA, Pfr. *A. testa imperforata, dextrorsa, fusiformi-oblonga, solida, leviter striatula, sub epidermide tenui fulvida alba; spiru elongata, ventroso-conica,*

13m

*apice obtusula; sutura sublacerata; anfr. 8 planiusculis, ultimo  $\frac{1}{3}$  longitudinis paulo superante, basi attenuato; plica columellari acute dentiformi, alba; apertura vix obliqua, acuminato-elliptica; perist. calloso, obtuso, margine dextro intus obsolete dentato.*

Long. 16, diam. 6 mill.

Hab. Oahu, Sandwich Islands (Dr. Newcomb).

53. *ACHATINA (VARICELLA) GUADELOUPENSIS*, Pfr. *A. testa oblongo-turrata, texui, laevigata, nitidissima, pellucida, virenticornea, varicibus arcuatis, vix prominulis, castaneo-marginatis passim munita; spira regulariter attenuata, apice obtusa; anfr. 7 convexiusculis, ultimo  $\frac{1}{3}$  longitudinis subæquante, basi attenuato; columella perarcuata, basi anguste truncata; apertura vix obliqua, acuminato-ovali; perist. tenui, margine dextro antrorum dilatato, rufo-limbato.*

Long. 14, diam.  $4\frac{1}{3}$  mill.

Hab. In insula Guadeloupe (Mr. Caillet).

54. *SPIRAXIS SANDWICHENSIS*, Pfr. *Sp. testa subperforata, oblongo-turrata, solidula, cerea; spira turrita, obtusula; anfr. 7  $\frac{1}{2}$  planiusculis, infra suturam plicatis, ultimo  $\frac{1}{3}$  longitudinis paulo superante; columella compressa, torta; apertura vix obliqua, ovali; perist. simplice, marginibus callo tenui junctis, dextro antrorum subdilatato, columellari subreflexo.*

Long. 9, diam. 3 mill.

Hab. Sandwich Islands (Dr. Newcomb).

55. *SPIRAXIS OBSOLETA*, Pfr. *Sp. testa subperforata, oblongo-turrata, tenui, levissime striatula, pellucida, nitida, pallide cornea; spira convexiusculo-turrata, obtusa; sutura marginata; anfr. 6  $\frac{1}{2}$  convexiusculis, ultimo  $\frac{2}{5}$  longitudinis formante; columella acuta, medio leviter torta; apertura vix obliqua, elliptico-ovali; perist. simplice, margine columellari brevissime reflexo, subadnato.*

Long. 8, diam. 3 mill.

Hab. Sandwich Islands (Dr. Newcomb),

56. *TORNATELLINA GOULDII*, Pfr. *T. testa ovato-conica, tenui, sublaevigata, pellucida, cornea; spira elongato-conica, apice obtusula; anfr. 5  $\frac{1}{2}$  convexis, ultimo  $\frac{2}{5}$  longitudinis subæquante, rotundata; apertura obliqua, erecto-lunari, lamella valida intrante parietali et plica crassa triangulari columellæ coarctata; perist. recto, tenui.*

Long. 4, diam.  $2\frac{1}{2}$  mill.

Hab. —?

57. *TORNATELLINA NEWCOMBI*, Pfr. *T. testa perforata, ovato-turrata, tenui, sublaevigata, parum nitida, diaphana, pallide cornea; spira subrectilineari, conica, obtusula; anfr. 7 subplanis, ultimo  $\frac{1}{3}$  longitudinis vix formante, basi conveexo; apertura obliqua, ro-*

*tundato-lunari, lamella mediocri parietali et plicis 2 parallelis columellæ coarctata; perist. simplice, recto.*

Long. 4, diam.  $2\frac{1}{3}$  mill.

Hab. Sandwich Islands (Dr. Newcomb).

58. **TORNATELLINA PERFORATA**, Pfr. *T. testa perforata, ovato-turrita, arcuato-striata, tenui, albido-hyalina; spira elongata, apice acuta; anfr. 7 convexis, ad suturam distanter plicatulis, ultimo  $\frac{1}{3}$  longitudinis paulo superunte; lamella parietali minuta; apertura vix obliqua, sinuato-ovali, acuminata; columella lamina torta munita; perist. tenui, margine dextro recto, superne antrorsum arcuato, basali expanso, columellari fornicatim reflexo, patente.*

Long. 18, diam.  $7\frac{1}{2}$  mill.

Hab. Venezuela.

#### 6. DESCRIPTIONS OF SIXTEEN NEW SPECIES OF PNEUMONOPOMA, FROM THE COLLECTION OF H. CUMING, ESQ.

BY DR. L. PFEIFFER.

##### A. OPISOPHTHALMA.

1. **DIPLOMMATINA CANTORI**, Pfr. *D. testa subrimata, sinistrorsa, ovato-oblonga, tenuiuscula, confertim et oblique plicata, albida; spira ovata, in conum acutiusculum terminata; anfr. 6 convexiusculis, ultimo ascendentem,  $\frac{1}{3}$  longitudinis vix superante; apertura subverticali, circulari; perist. subsimplice, continuo, superne adnato, cæterum breviter expanso.*

Long. 2, diam. 1 mill.

Hab. Lord Howe's Island, New Hebrides (Mr. Macgillivray).

2. **TRUNCATELLA CEYLANICA**, Pfr. *Tr. testa subrimata, attenuato-cylindracea, striatula, pellucida, nitida, corneo-rufa vel flavescente; anfr. superst. 4, supremis 2 rotundatis, reliquis planioribus, omnibus ad suturam plicato-crenatis, ultimo basi non compresso; apertura verticali, ampla, angulato-subovali; perist. continuo, margine dextro tenui, expansiusculo, columellari leviter arcuato, apppresso.*

Long. 6, diam. 2 mill.

Hab. Ceylon.

3. **TRUNCATELLA TERES**, Pfr. *Tr. testa vix subrimata, cylindrica, tenuiuscula, longitudinaliter magis minusve distincte costulata, pellucida, nitida, rufo-cornea; sutura marginata, valide plicato-crenata; anfr. superst. 4 subæqualibus, convexiusculis, ultimo basi breviter cristato, calloso, albido; apertura verticali, late orali, superne angulata, ad dextram dilatata; perist. simplice, continuo, margine dextro expansiusculo, columellari adnato.*

Long. 6, diam. 2 mill.

Hab. Isle of Mauritius, and Trinity Bay, Australia.