



















REGIO

CHIRURGORUM COLLEGIO

LONDINENSI

*DECIMUM QUINTUM*

HUNC

NATURÆ VIVARII

*FASCICULUM,*

D. D. D.

GEORGIUS SHAW.

E. NODDER.

TO THE  
ROYAL COLLEGE OF SURGEONS  
OF  
LONDON

*THIS FIFTEENTH VOLUME*  
OF THE  
NATURALIST'S MISCELLANY

RESPECTFULLY INSCRIBED

BY

GEORGE SHAW.

E. NODDER.



## TETRAO ARENARIA.

\*\*\*\*\*

### *CHARACTER GENERICUS.*

*Rostrum convexum, brevem.*

*Macula prope oculos nuda, aut papillosa, aut plumis rarius tecta.*

### *CHARACTER SPECIFICUS, &c.*

TETRAO griseo-flavescens, torque abdomine crisoque (maris) atris.

TETRAO ARENARIA. T. torque, abdomine et criso atris, rectricibus fusco et griseo fasciatis apice albis, intermediis duabus fulvescentibus.

*Lin. Syst. Nat. Gmcl. p. 755.*

*Pall. nov. comm. Petrop. p. 418.*

---

Primus accurate descripsit hanc speciem celeberimus Pallas in Academiæ Petropolitanae Commentariis.

“ Magnitudine hæc avis Perdicem superat, habitu Alchatam refert. Rostrum quam in perdice tenuius, prorsus ut in Alchata, cinereo cœrulescens, apice nigricante. Palpebræ nudæ, pallidæ, marginibus subpapillosis, flavescentibus. Supercilia plumosa, tecta. Caput albido cinereum in masculis, vertice usque

usque in cervicem grisco-flavescenti nebuloso. Gula ferrugineo-fulva, colore versus latera colli diffuso diluto, trianguloque atro submedio collo terminata. Collum jugulumque totum, in hoc sexu, cano albida, plumis vestita singularibus truncatis, densioribus, elasticis, columbarum similibus. Cervix inferior, dorsum totum adusque caudam, alarumque bases plumis testaceo-albidis, annulo singulis fusco terminali, maculam ovatam lutescentem cingente, notatis variantur. Inter pectus et jugulum circulus ater, hinc pectus albido; sed abdomen, femora, crissum atra. Subcaudales albæ, strigis aliquot transversis nigris. Tectrices primariæ remigibus concolores, canescentes; secundariæ extus luteofulvescentes, efficientes quasi speculum alare hujus coloris. Femina paulo major mare differt colore per totum corpus pallide flavescente, in capite, collo, juguloque nigro guttato, in dorso fasciolis transversis sagittatis, crebris variegato. Loca incolit deserta circa mare Caspium."





THE  
SAND PARTRIDGE.



*GENERIC CHARACTER.*

*Bill convex, short.*

*Spot near the eyes, either naked, granulated,  
or but sparingly covered with feathers.*

*SPECIFIC CHARACTER, &c.*

Yellowish-grey PARTRIDGE with the abdomen, vent, and collar (of the male) black.

Sand Grouse.

*Lath. syn. 2. p. 751.*

---

This species seems to have been first accurately described by the celebrated Dr. Pallas, in the Petersburg Transactions. Its size exceeds that of a common partridge, and its habit resembles that of the Alchata: the bill, which is more slender than in the partridge, is of a blueish ash-colour, with a black tip: the eye-lids naked, pale, and beset with slight yellowish papillæ round the edges: the brows covered with feathers: the head, in the male, of a whitish ash-colour, the crown being clouded as far as the neck with yellowish grey; the throat

throat orange-ferruginous, growing paler towards the sides of the neck, and terminated below by a black triangle: the neck is whitish-grey, and covered by feathers of a truncated shape, strong, and elastic like those of a pigeon: the remainder of the neck, and the back with pale testaceous feathers, each marked at the tip with a black ring including a yellowish spot: the abdomen, thighs, and vent are black: the smaller wing-coverts, and the quill-feathers are pale grey; but the larger coverts are of an orange-yellow, forming a speculum or oblong spot of that colour on the wings. The female, which is rather larger than the male, is almost entirely of a pale yellowish colour, marked on the head and neck with black specks, and variegated on the back by transverse streaks of the same colour. This bird is an inhabitant of the sandy deserts in the neighbourhood of the Caspian Sea.

# PAPILIO PROTENOR.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatæ.

*Alæ* (sedentis) erectæ, sursumque conniventes,  
(volatu diurno.)

*Lin. Syst. Nat.* p. 744.

## CHARACTER SPECIFICUS, &c.

**PAPILIO** alis dentatis nigris, anticis stria humerali rubra, posticis annulo rubro ad angulum ani.

**PAPILIO PROTENOR.** P. alis dentatis nigris, posticis subtus macula difformi rubra anguli ani.

*Fabr. spec. ins. 2. p. 7.*

*Eg. Troj.*

---

Indiam? incolit Papilio Protenor. Pulchrum specimen unde desumpta est figura hæc nostra in Museo celeberrimi Gulielmi Hunter asservatur.

## PROTENOR.



### *GENERIC CHARACTER.*

*Antennæ* gradually thickening towards the tip,  
and generally terminating in a knob.

*Wings* (when sitting) erect and meeting up-  
wards : (Flight diurnal.)

### *SPECIFIC CHARACTER.*

Black Butterfly with denticulated wings, the up-  
per pair marked by a red shoulder-streak,  
the lower by a red ring at the inner angle.

---

The Papilio Protenor is a native of India? The elegant specimen from which the present figure was taken, is preserved in the Museum of Dr. William Hunter.



2



# CANCER SEBANUS.



## *CHARACTER GENERICUS.*

*Pedes* octo (rarius sex aut decem :) insuper manus duæ chelatæ.

*Oculi* duo, distantes, plurimis pedunculati, elongati, mobiles.

*Cauda* articulata, inermis.

## *CHARACTER SPECIFICUS, &c.*

CANCER miniatus, thorace orbiculato, pedibus hirsutis, chela dextra maxima.

CANCER fluviatilis, sive Gammarus Americanus.

*Seb. mus.* 3. p. 43. t. 18. f. 4.

---

Americam testante Seba, incolit Cancer hic depictus, “cujus dextri lateris pes insolitæ est crassitiei atque magnitudinis, ut vel reliquis omnibus pedibus, totique simul corpori fere præponderet.”

THE  
SEBAN CRAB.



*GENERIC CHARACTER.*

*Legs* generally eight; (in some species six or ten,) besides two claspers or chelated arms.  
*Eyes* two, commonly distant; footstalked, moveable.

*Tail* articulated.

*SPECIFIC CHARACTER, &c.*

Red CRAB with orbicular thorax, hairy feet, and right claw excessively large.

Great-clawed American CRAB.

---

This Crab, according to Seba, is a native of America, and is remarkable for the excessive size of its right claw, which is said to exceed the weight of all the rest of the animal.





## PHYTOLITHUS FILICIS.



In Natura investiganda vix aliud jucundius quam vegetabilium et animalium certissimas reliquias per totum fere orbem sparsas, interdum etiam ipsis animalibus plantisque quæ in iisdem regionibus nunc dierum inveniuntur prorsus dissimiles attentius considerare. In re difficillima etiamnum satis obscura diu inter philosophos pugnatum est variis et sibi invicem contrariis argumentis. At inanca fortasse vel adhuc est telluris cognitio, et concedendum certe est hanc nostram a primæva terra longe esse diversam, quam magna sæpe vi convulsam fuisse satis constat et igni aquaque vastatam; adeo ut hodie super rudera quasi et ruinas ambuletur.

Hujusmodi permutationum quamvis causa lateat, vis est notissima. In summis montibus a mari longe remotis, in ipsis etiam Andibus Peruvianis reperiuntur plurima substantiarum marinorum tum animalium tum vegetabilium vestigia, quarum multas cum recentes nunquam viderimus, vel species deperditas putemus necesse est, vel in altissimo oceani profundo latere, unde forsitan nec insana tempestatum vi attolluntur.

In intimis etiam carbonis et schisti stratis manifesta insunt indicia piscium, crustatorum, planarum, &c.; sæpius autem filicum variarum, quarum aliæ ad species genuinas facile reduci possunt, aliæ vero ad illas referendæ sunt quæ aut diu perierunt, aut adhuc ignorantur. Harum quæ in

Museo

Museo Leveriano spectantur perfectior nulla est quam quæ in tabula cernitur, filix scilicet mas vulgaris seu *Polypodium filix mas Linnaei*, quo specimine vix bellius et integrius vel in ipso *horto sicco* videris, ipsæ enim in foliis fructificationes clare inspici possunt.

Dc planta vulgatissima speciatim tractare supervacancium foret; per totam fere Angliam in locis incultis et juxta sepes generatur.

Repertum est hoc specimen in fragmento lapidis schistosi immersum.





RPN

A

## FERN PHYTOLITHUS.



The indisputable remains of animal and vegetable bodies, so plentifully dispersed throughout almost all parts of the globe, and of which a great many are of a widely different appearance from the present natives of the regions in which they are found, form one of the most interesting points of speculation in the history of nature.

Various and even contradictory hypotheses have occasionally been proposed as elucidations of this subject, which yet remains in a considerable degree of obscurity ; the true theory of the earth being perhaps still but very imperfectly understood. It must be allowed, however, that vast and violent changes have been wrought in the body of the primeval earth ; that we walk as it were on the ruins of the original structure ; and that both fire and water have united their forces to derange the first formation.

Of these great convulsions of nature, though the causes yet remain unexplored by all the endeavours of improved philosophy, yet the effects are every where visible.

On the tops of mountains, remotely distant from the sea, even on those of the Andes in Peru, are found the remains of innumerable marine productions, both animal and vegetable ; many of which are still unknown in their recent state, and are therefore

fore either lost species, or else such as inhabit the deepest recesses of the ocean, from whence they are scarce ever raised by the utmost violence of storms and tempests.

In deep-laid strata of coal, slate, &c. are found plentiful specimens of fish, crustacea, plants, &c. The most frequent, however, seem to be those of Ferns, of which a great many different kinds are observed ; some easily reducible to their archetypes or proper species, while others appear to belong to species either lost or still unknown.

Among the most complete and curious specimens in the Leverian Museum, is that figured on the annexed plate, which represents in the most perfect manner the well-known European plant called the common male fern, (*Polypodium filix mas.* Lin.) So extremely perfect is this fossil vegetable, that the fructifications are distinctly seen on the leaves nearly in the same state as in a dried specimen.

To particularize the history of so well known a vegetable would be entirely unnecessary : it is sufficient to say that it is principally found in waste places, and by hedge sides, and is produced in most parts of this kingdom. The specimen is imbedded in a fragment of slaty stone.

## PIPRA RUPICOLA.

\*\*\*\*\*

### CHARACTER GENERICUS.

*Rostrum capite brevius, basi subtrigonum, integerrimum, apice incurvum.*

*Pedes gressorii.*

*Lin. Syst. Nat. p. 338.*

### CHARACTER SPECIFICUS, &c.

PIPRA RUPICOLA. *P. crista erecta margine purpurea, corpore croceo, tectricibus rectricum truncatis.*

*Lin. Syst. Nat. p. 338.*

Upupa Americana lutea.

*Ger. orn. 2. p. 64. t. 206.*

Coq de roche.

*Pl. Enl. 39. & 747.*

---

Avem hanc raram et perpulchram, in Peruvia et Surinamia generatam, insignit mira cristæ et tectricum conformatio. Incolit, ut plurimum, rupes et loca cavernosa, magnitudine columbæ minori par. Differt a mari temina penitus infuscata.

B







THE  
ROCK MANAKIN.



*GENERIC CHARACTER.*

*Bill* shorter than the head, somewhat triangular at the base, and bent at the tip.

*Feet* formed for walking.

*SPECIFIC CHARACTER, &c.*

Orange-coloured MANAKIN, with upright compressed crest, and truncated tail-coverts.

Hoopoe Hen.

*Edw. Glean.* t. 264.

Coq de roche du Perou.

*Buff. ois.* 4. p. 437.

*Museum Leverianum* 1. p. 15. pl. 4.

---

This rare and beautiful bird is a native of Peru and Surinam, and is remarkable for the singular conformation of its crest and tail-coverts. It is generally seen in the neighbourhood of rocks and caverns, and is about the size of a small pigeon. The female differs from the male in being entirely of a brown colour.



## LACERTA AURITA.



### CHARACTER GENERICUS.

*Corpus* tetrapodum, elongatum, caudatum, nudum.

### CHARACTER SPECIFICUS, &c.

LACERTA collo utrinque lobo semiorbiculato denticulato.

LACERTA AURITA. L. cauda tereti mediocri utrinque ad latus callosis punctis aspera, plica gulæ transversa subgemella, oris angulis utrinque in cristam semiorbiculatam mollem scabram dentatam dilatatis.

*Lin. Syst. Nat. Gmel.*

LACERTA Mystacea.

*Pall. ih. 3. p. 702.*

---

Luculente adeo et accurate explicit hanc speciem celeberrimus Pallas ut ipsissima auctoris verba sine ulla immutatione transcribere non dubitaverim.

“ Magnitudo adultis fere supra Geckonem. Caput retusum. Anguli oris dilatati utrinque in cristam semiorbiculatam mollem, extus punctis scabram, margine dentatam, in vivo animale sanguine turgescentem.

centem. Parotides utrinque muricatae; plica gulæ transversa subgemella. Corpus ventricosum, depresso, cum cauda totum punctis acute prominulis scabrum, quæ majora in pedibus. Caudæ latera (in tractu utrinque longitudinali) callulis muricata. Digihi pedum unguiculati, intermedii tres serrati, duo bifariam, interior uno versu. Color supra cinereo et lutescente nebulosus, atomis creberrimis fuscis; subtus sordide albus, litura sterni apiceque caudæ subtus atris. In collibus arenosis Naryn, ut et in deserti Comani sabuletis non infrequens.”





THE  
LOBE-CHEEKED LIZARD.



*GENERIC CHARACTER.*

*Body* four-footed, elongated, tailed; without any secondary integument.

*SPECIFIC CHARACTER, &c.*

LIZARD with a semiorbicular denticulated lobe on each side the neck.

LOBE-CHEEKED LIZARD.

*General Zoology*, vol. 3. p. 244.

---

This remarkable Lizard is thus accurately described by Dr. Pallas:—

The size of the full-grown animal is somewhat superior to that of the *Lacerta Gecko*: the head is obtuse, and rounded in front; and the corners of the mouth are dilated on each side into a soft, semiorbicular crest, which is roughened externally with small points, toothed on the edges, and, in the living animal is turgid with blood: the parotids are muricated, and the throat is furnished with a double transverse pleat: the body is ventricose, depressed, and together with the tail is roughened by small sharply prominent points, which are largest on the feet:

feet : the sides of the tail are muricated in a longitudinal direction with small longitudinal points : the toes are furnished with claws, and the three intermediate toes are serrated : two on both sides, and the other only on one : the colour above is a clouded variegation of yellowish and cinereous, with thickly-scattered dusky specks ; beneath dull white, with a black streak on the breast and beneath the tip of the tail. It inhabits the sandy and hilly deserts of Siberia.

## SPHINX CLUENTIUS.



### CHARACTER GENERICUS.

*A*ttennæ subprismaticæ, utroque fine attenuatæ.

*Lingua* exserta (plerisque.)

*Palpi* duo reflexi.

*Alæ* deflexæ.

### CHARACTER SPECIFICUS, &c.

SPHINX alis fuscis subferrugineo-variis, inferioribus basi fasciaque flavis, corpore utrinque luteo-maeulato.

## SPHINX CLUENTIUS.

*Cram. pap. t. 78. f. b. & 126. f. a.*

---

Surinamiam incolit Sphinx Cluentius, magnitudine vera in tabula depictus.







## CLUENTIUS.



### *GENERIC CHARACTER.*

*Antennæ* subprismatic, attenuated at each extremity.

*Tongue* generally exserted.

*Feelers* two, reflex.

*Wings* deflected.

### *SPECIFIC CHARACTER, &c.*

SPHINX with brown wings, with subferruginous variegations; the lower wings yellow at the base and in the middle, and the body marked on each side by deep-yellow spots.

### SPHINX CLUENTIUS.

*Cramer*, t. 78.

---

The superb Sphinx here exhibited is a native of Surinam, and is represented on the plate in its natural size.



# BUCCINUM RUMPHII.

\*\*\*\*\*

## *CHARACTER GENERICUS.*

*Animal* Limax.

*Testa* univalvis, spiralis, gibbosa.

*Apertura* ovata, desineus in canaliculum (s. retusam lacunam) dextrum, cauda rectusum.

*Labium* interius explanatum,

## *CHARACTER SPECIFICUS, &c.*

BUCCINUM carneo-flavescens fasciis longitudinalibus undulatis fuscis.

B: tuberosum. var.  $\gamma$ .

*Lin.* Gmel. p. 3473.

Le Papier marbré.

*Knorr.* 6. t. 18. f. 1.

*Rumph.* t. 23. f. c.

---

E mariibus Indicis extrahitur rarissima hæc cochlea, cuius magnitudinem veram cernere est in tabula.





596



RPN

THE  
MARLED BUCCINUM.



*GENERIC CHARACTER.*

*Animal* allied to a *Slug*.

*Shell* univalve, spiral, gibbous.

*Aperture* ovate, ending in a channel pointing towards the right.

*Interior Lip* expanded.

*SPECIFIC CHARACTER, &c.*

Yellowish flesh-coloured *BUCCINUM*, with longitudinal undulated brown bands.

Le Papier marbré.

*Rumph. Knorr. &c.*

---

This very rare shell is a native of the Indian seas, and is exhibited on the plate in its natural size.



## EMBERIZA CIRIS.



### CHARACTER GENERICUS.

*Rostrum conicum.*

*Mandibulae* basi deorsum a se invicem discedentes : inferiore lateribus inflexo-coaretata, superiore angustiore.

*Lin. Syst. Nat.* p. 308.

### CHARACTER SPECIFICUS, &c.

EMBERIZA capite cœruleo, abdomine fulvo, dorso viridi, pennis viridi-fuscis.

*Lin. Syst. Nat.* p. 313.

*Chloris ludoviciana* vulgo Papa dicta.

*Briss. av.* 3. p. 200.

*Fringilla Mariposa.*

*Scopoli ann.* 1. No. 222.

---

Sive formæ sive colorum habeatur ratio, merito numerari debet Americana haec avis in pulcherrimis sui generis. Magnitudinem naturalem ostendit tabula. Femina a mare differt corpore supra viridi, subtus flavescente.







THE  
PAINTED EMBERIZA.



*GENERIC CHARACTER.*

*Bill* conical : Mandibles separating a little from the base downwards : the sides of each mandible bending rather inwards.

*SPECIFIC CHARACTER, &c.*

Green EMBERIZA, with blue head, fulvous abdomen, and brownish-green wing-feathers.

The painted Finch.

*Edw. pl. 130.*

Le Pape.

*Buff. ois. 4. p. 176.*

*Pl. Enl. 159. f. 1.*

---

This species, which is a native of many parts of America, may be considered as one of the most elegant and beautiful of its tribe : it is represented on the plate in its natural size : the female differs from the male in being green above, and yellowish beneath.



# ANARHICHAS PANTHERINUS.



## CHARACTER GENERICUS.

*Caput* obtusiusculum.

*Dentes primores* supra infraque conici, divergentes, sex pluresve.

*Molares inferiores* palatique rotundati.

*Membr. branch.* radiis sex.

*Corpus* teretusculum, pinna caudæ distincta.

## CHARACTER SPECIFICUS, &c.

ANARHICHAS luteus, maculis rotundatis fuscis.

ANARHICHAS PANTHERINUS. A. maculis per totum corpus rotundis fuscis.

*Zouiew act. Petrop.* 1781.

*Lin. Syst. Nat. Gmel.* p. 1144.

---

In Actis Petropolitanis primum descripta fuit hæc species, cuius longitudo tres circiter pedes æquat.

“ *Corpus* teretusculum, ventricosum, lanceolatum, a medio ventre, ubi crassities ejus maxima est, ad caudam, sensim decrescens, nudum, lubricum, squamarum loco punctis minoribus concoloribus adspersum; cute firma, densa, coloris luteoflavescens, maculis per totum corpus et pinnam

dorsalem, excepta parte prona, rotundis, fuscis,  
majoribus notatum. Caput subglobosum; labia  
duplicata; rictus amplius; palato inserti dentes sep-  
tem; oculi majusculi; dissiti; branchiarum apertura  
lunata, coarctata, opercula ex laminis duabus con-  
flata; pinnæ pectorales amplissimæ, rotundatæ;  
anus post medium corporis ad exordium pinnæ ani:  
linea lateralis nulla: cauda lanceolata, ad pinnam  
subcaudalem torosior quam ad dorsalem.

“ Habitat in Oceano Septentrionali et Mari Albo.  
A Russis in esum non trahitur etiamsi caro ejus  
laudetur.”



१६५



१६५

THE  
PANTHER ANARHICHAS.



*GENERIC CHARACTER.*

*Head* rather obtuse.

*Fore-Teeth* both above and below conical, divergent, strong, six or more in number.

*Grinders* in the lower jaw and palate rounded.

*Gill Membrane* six-rayed.

*Body* roundish: tail-fin distinct.

*SPECIFIC CHARACTER, &c.*

Yellow Wolf-Fish, with rounded brown spots.

The Panther Wolf-Fish.

---

This species, which measures about three feet in length, seems to have been first described in the Petersburg Transactions. The body is subcylindric, ventricose, and lanceolate, gradually decreasing from the middle of the abdomen, where it is thickest, to the tail. It is naked, slippery, and beset in place of scales with small points of the same colour with the skin, which is thick and strong, and of a deep yellow, marked throughout the whole body except the under part, as well as on

the dorsal fin, with round, large, dusky spots : the head is subglobose ; the lips doubled, and the gape wide : in the palate are inserted seven teeth : the eyes are rather large and distant : the branchial apertures lunated and straitened, and the gill-covers consist of two plates on each side : the pectoral fins are very large and rounded : the vent is situated beyond the middle of the body, at the beginning of the anal fin : the lateral line is wanting : the tail is lanceolate, and is thicker towards the subcaudal fin than towards the dorsal one. It is a native of the Northern and White Seas, and though the flesh is said to be agreeable, is not eaten by the Russians.

## MUREX FEMORALE.



### CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, exasperata suturis membranaceis.

*Apertura* desinens in canalem integrum, rectum seu subadscendentem.

*Lin. Lyst. Nat.* p. 1213.

### CHARACTER SPECIFICUS.

MUREX FEMORALE. M. testa varicibus decussatis trigona rugosa anterius nodulosa, apertura edentula anterius transversa.

*Lin. Syst. Nat.*

*Gualt. test* t. 50. f. c.

*Seb. Mus.* 3. t. 63. f. 7.—10.

*Regenf. Conch* 1. t. 2. f. 21.

---

Maria Indica præcipue incolit elegans conchyliuni magnitudine vera in tabula depictum.





599



RN

THE  
FEMORAL MUREX.



*GENERIC CHARACTER.*

*Animal* allied to a *Slug*.

*Shell* univalve, spiral, commonly roughened by membranaceous sutures.

*Aperture*, terminating in a strait or subascending channel.

*SPECIFIC CHARACTER, &c.*

Yellow-brown roughened MUREX, with subtrigonal ventricose body, and toothless aperture.

*Argenv.* t. 10. f. b. ?

*Grew.* *Mus. Reg. Soc.* t. 10. f. 4. ?

---

This elegant shell is chiefly found in the Indian Seas, and is represented on the plate in its natural size.



# PAPILIO AMPHITRYON.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatæ.

*Alæ* (sedentis) erectæ sursumque conniventæ  
(volatu diurno.)

## CHARACTER SPECIFICUS, &c.

PAPILIO alis anticis apice flavo-maculatis, posticis fascia flavescente, subtus lunulis cæruleis.

PAPILIO AMPHITRION. P. alis nigris fascia inæquali flava, posterioribus subtus striga e punctis flavis lunulisque cæruleis.

*Lin. Syst. Nat. Gmel.* p. 2247.

# PAPILIO AMPHITRYON.

*Cram.* t. 7. f. A. B.

*Eq. Archiv.*

---

Americæ Australis partes calidiores incolit Papilio Amphitryon, coloribus interdum leviter varians. Magnitudinem veram ostendit tabula.





009  
A



114

## AMPHITRYON.



### *GENERIC CHARACTER, &c.*

*Antennæ* thickening towards the upper part, and generally terminating in a clavated tip.

*Wings* (when sitting) erect, and meeting upwards. (Flight diurnal.)

### *SPECIFIC CHARACTER.*

Blackish Butterfly, with the upper wings spotted with yellow at the tips, the lower wings marked by a yellowish band, and beneath by a row of blue crescents.

---

This insect is a native of several parts of South America, and is subject to some slight variation as to colour: the plate represents it in its natural size.



# MYCTERIA AUSTRALIS.



## CHARACTER GENERICUS.

*Rostrum* subascendens, acutum, mandibula superiore triquetra.

*Frons* calva.

*Nares* lineares.

*Pedes* tetradactyli.

*Lath. ind. orn.* p. 670.

## CHARACTER SPECIFICUS, &c.

MYCTERIA alba, capite colloque viridi-nigris, tectricibus, pennis scapularibus, caudaque nigris, rostro nigro, pedibus rubris.

# MYCTERIA AUSTRALIS.

*Lin. Trans.* Vol. 5. p. 34.

---

Generis Mycteriæ species Americana seu communis, physicis jamdiu cognita, varias Americæ Australis partes, incolit. Species autem quam depinximus, quamque generat Australasia, distat insigniter ab avi Americana, quod caput et collum non denu-dentur, sed plumis vestiantur saturatim viridibus, qualibus fere collucere solet Anas Boschas mas ; splendore insuper versicolori vel in cœruleum vel purpureum transeunte. Rostrum nigrat ; et a basi

D

mandibulæ inferioris dependet extensa aliquatenus in gulam cutis quædam nuda, seu sacculus membranaceus ruberrimus. Cætera avis nivei est candoris, alarum tectricibus omnibus exceptis nigerrimis, nigraque cauda. Crura pedesque ruberrima. Icones accuratissime delineatæ sunt apud Dominum White, coloniæ Britannicæ in Australasia chirurgum præcipuum, unde constat hanc speciem, cum plene adleveret, Mycteriæ ipsi Americanæ magnitudine esse parem ; sex enim vel septem pedes alta depingitur. Specimen autem nostrum quod ostendit tabula, quodque in Museo Leveriano asservatur, vix quinque pedes superat.





THE  
AUSTRALASIAN JABIRU.



*GENERIC CHARACTER.*

*Bill* subascendant, sharp; with the upper mandible triquetrous.

*Front* bare.

*Nostrils* linear. *Feet* tetradactylous.

*SPECIFIC CHARACTER, &c.*

White JABIRU, with the head and neck green-black; the coverts, scapulars, and tail black; the bill black, and the legs red.

New Holland JABIRU.

*Lin. Trans.* Vol. 5. p. 34.

---

The common or American Jabiru has long since been known to naturalists, and is found in many parts of South America. The present species is a native of Australasia, or New Holland, and differs very strikingly from the American bird in the appearance of the head and neck, which, instead of being bare, are entirely covered with feathers of a most elegant dark green, similar to those of a mallard, and accompanied by an elegant blue or purple

changeable lustre. The bill is black, and beneath the lower mandible, to some little distance down the throat, is an extensile naked skin or sacculus of a bright red colour: the rest of the bird is of a snowy whiteness, the coverts of the wings, and the tail excepted, which are jet-black: the legs and feet are of a bright red. According to some very accurate drawings, in the collection of Mr. White, Chief Surgeon to the Settlement at Botany Bay, this species, when full-grown, must be equal in size to the common Jabiru, being not less than six or seven feet high. The present specimen, however, which is preserved in the Leverian Museum, falls short of the height just mentioned, scarcely exceeding that of four and a half, or five feet.

# ARANEA TARANTULA.



## CHARACTER GENERICUS.

*Pedes* octo.

*Oculi* octo.

*Os* unguibus seu retinaculis duobus.

*Palpi* duo articulati; masculis genitalibus cappingtati.

*Anus* papillis textoriis.

*Lin. Syst. Nat.* p. 1030.

## CHARACTER SPECIFICUS, &c.

ARANEA fusca, abdome supra maculis triangonis nigris margine albidis, pedibus nigro variatis.

ARANEA TARANTULA. A. subtus atra, pedibus atro-fasciatis.

*Lin. Syst. Nat.* p. 1035.

---

Araneam Tarantulum, de qua abunde fabularum nugarumque laboriose doctarum andivimus, quæque ad hunc usque diem vulgi fidem et ignorantiam exercet, generant, calidiores Italiæ regiones, aliæque nonnullæ partes Europeæ, arva amantem, ut plurimum,

mum, sicca et soli exposita. Magnitudo et colores  
accurate in tabula exprimuntur. In ævo, cui plen-  
num scientiæ lumen affulsit, satis sit notare ficta et  
inania haberri quæ de symptomatibus morsum Taran-  
tulæ sequentibus, non nisi musica tollendis decan-  
tari solita sunt, et apud sanos omnes totam fabulam  
exolevisse.



602



✓

THE  
TARANTULA.



*GENERIC CHARACTER.*

*Legs* eight.

*Eyes* eight.

*Mouth* furnished with two hooks or holders.

*Palpi* or *Feelers* two, the tips of which (in the males) distinguish the sex.

*Abdomen* terminated by papillæ or *teats*, thro' which the insect draws its thread.

*SPECIFIC CHARACTER, &c.*

Brown Spider, with the back of the abdomen marked by trigonal black spots with whitish edges, and the legs variegated with black.

The TARANTULA Spider.

---

The Tarantula, of which so many idle recitals have been detailed in the works of the learned, and which even to this day continues to exercise the faith and ignorance of the vulgar, is a native of the warmer parts of Italy, and some other European regions, and is generally found in dry and sunny plains.

plains. Its size and colours are accurately represented on the annexed plate. In the present illuminated period, it may be sufficient merely to add, that the extraordinary symptoms supposed to ensue from its bite, as well as their supposed cure by the power of musick alone, are entirely fabulous, and are now sufficiently exploded among all rational philosophers.

# VESPA CAMPANARIA.



## CHARACTER GENERICUS.

Os corneum; maxilla compressa. Palpi quatnōr  
inæquales, filiformes.

*Antennæ* filiformes; articulo primo longiore cy-  
lindrico.

*Oculi* lunares.

*Corpus* glabrum.

*Aculeus* punctorius reconditus. *Alæ* superiores  
plicatæ.

## CHARACTER SPECIFICUS, &c.

VESPA Holsatica. V. nigra, linea utrinque ad  
humeros, maculisque scutellaribus luteis,  
abdomine luteo segmentis basi transverse  
punctisque contiguis nigris.

Latreille. *Ann. du Museum d'Histoire  
Naturelle*, p. 287.

---

Insectorum in nidificando æque ac avium ingeni-  
osa est solertia, et fortasse magis varia, illorum  
præcipue quæ *Hymenoptera* et *Neuroptera* dicuntur;  
aliorum rarior longe, at non minus miranda. Ingens  
species araneæ Americanæ nidum struit calido arti-  
ficio

ficio cylindraceum et tubulatum, cuius introitum, ut minus suspecta lateat, valvis mobilibus occludit, prædam si quam appropinquantem viderit, exiliendo arreptura. Nota est Myrmeleonis Formicaleonis Linnæi astutia, qui dum adhuc larva est, in arena cubans suo ipsius labore profunde et orbiculatim excavata, insectis quæ forte juxta repentina deciderint, vigilantissime insidiatur. Termitum quos parit Asia et Africa mirandum est ingenium, qui nidos ædificant in altitudinem multorum pedum, concameratos, et in magnam cellarum varietatem divisos. At longe præ aliis omnibus dædalisch apum vesparumque laboribus maximus semper habitus est honos; e quibus nulla elegantius nidificat quam species reliquis minus nota, quod cubile sibi faciat in locis solitariis, in summis scilicet stabulis, aliisque id generis. Struit haec nidum insignem festucæ, ut plurimum, seu frustulo ligni, aut tali alicui affixum, pendenteque de culmine horrei, aut septi cujuslibet, ore deorsum spectante, constantemque e globis concentricis tribus, quatuor, et interdum pluribus, substantiae fere ejusdem ac papyri rudioris, imo modice aperto rotundoque. In medio globi interioris construitur cellularum congeries, duodecim scilicet seu quindecim, seu etiam viginti, quasi papyriarum, more ipsius nidi, et circa columnam centralem dispositarum. Cæteris paulo humilior est circulus cellularum exterior. Reperitur nidos fere exente æstate. Quod impossibile sit internam nidi conformatiōnem plene repræsentare, ni exacta sit pars aliqua ut inspiçiantur cellulæ, ideo in tabula duæ proponuntur figuræ, quarum una nidum externum ostendit; altera

per

per globos omnes concentricos nidum totum aperit  
sectione recta longitudinali.

Notandum porro est Vespam quandam America-  
nam nidum formare eodem omnino modo quo haec  
nostra ; longe autem majorem, et e numerosioribus  
globis concentricis constantem, duodecim scilicet  
seu pluribus ; suspensum, ut plurimum, ramulis ar-  
borum, pinorum nempe et aliarum ; seu potius iis  
affixum.

THE  
CAMPANULAR WASP.



*GENERIC CHARACTER.*

*Mouth* horny ; with a compressed jaw : *Feelers* four, unequal, filiform.

*Antennæ* filiform ; with the first joint longer than the rest and cylindrical.

*Eyes* lunated.

*Body* smooth. *Sting* concealed. *Wings* pleated.

*SPECIFIC CHARACTER, &c.*

Black WASP, with a luteous line on each side the thorax, two luteous spots on the scutellum, and luteous abdomen with transverse annular black bands accompanied by black spots at their ends.

La Guepe de Holstein.

*Latreille. Ann. d'Hist. Nat.* p. 288.

---

The nidification of insects is scarcely less curious than that of birds, and the variety of structure is even greater. It is principally in the classes *Hymenoptera* and





and *Neuroptera* that this practice prevails; in the other classes it is less general, though not less singular. Thus a large species of American spider forms a very curious tubular or cylindrical nest in which it generally resides, and, in order to remove all suspicion of the dangerous inhabitant, it constantly forms a moveable valve or door, with which it closes the entrance, and when it perceives the approach of any other insect, springs out and seizes the prey.

The curious policy of the *Myrmeleon Formicaleo* in its larva state is well known. It lies in ambush at the bottom of a deep conical cavity formed in dry sand, into which other insects accidentally falling are sure of being seized by the watchful inhabitant.

The Termites of Africa and Asia afford wonderful examples of this nidificating power, and form stupendous structures of many feet in length, and divided into a vast variety of cells and passages. Of all insects however those whose operations are most conspicuous are the different kinds of Bees and Wasps, whose labours have been admired in all ages, and celebrated from the most remote antiquity. Amongst the most elegant of these structures may be numbered that of a species of Wasp, which, on account of its frequenting retired places, and fixing its nest in the upper parts of buildings, is less attended to than many others of its genus. The structure of the nest is singularly curious. It is commonly fastened or suspended by a straw, or slight fragment of wood, or other small projecting substance, from the upper part of some barn or outhouse, and consists of three or four, and sometimes more concentric globes,

globes, of an appearance resembling that of coarse paper, with a rather small round opening at the bottom. In the middle of the interior or central globe is placed the congeries of cells, to the number of twelve, fifteen, or even twenty : these are arranged round a kind of central column, and are composed of the same paper-like substance with rest of the nest ; the exterior circle of cells being somewhat lower or shorter than the rest. These curious nests are generally found about the latter end of summer. It being impossible to give a clear idea of the internal structure of the nest without laying it open, it is represented in two views on the annexed plate ; one of the figures shewing its complete exterior appearance ; the other being cut perpendicularly downwards in order to display the central cells.

It may not be improper to observe that there is an exotic species of Wasp, a native of America, which forms a nest on a plan exactly similar to our present species, but differing in being much larger, and having the concentric globes far more numerous, viz. from ten to twelve, or more. These American Wasp-nests are commonly suspended from, or rather affixed to the small branches of trees, as Firs, &c.

# PAPILIO MEROPE.



## *CHARACTER GENERICUS.*

*Antennæ* a piccm versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes,  
(volatu-diurno.)

## *CHARACTER SPECIFICUS, &c.*

PAPILIO alis flavis limbo nigro, posterioribus  
subtus fascia fusca.

PAPILIO Brutus.

*Lin. Syst. Nat. Gmel.* p. 2237.

PAPILIO MEROPE.

*Cram. t. 378. f. d. e.*

---

Africæ varias partes incolit Papilio Merope, cuius  
veram magnitudinem cernere est in tabula: colori-  
bus interdum paulo variat.

## MEROPE.



### GENERIC CHARACTER.

*Antennæ* thickening towards the upper part, and generally terminating in a clavated tip.

*Wings* (when sitting) erect, and meeting upwards. (Flight diurnal.)

### SPECIFIC CHARACTER, &c.

Butterfly with yellow wings bordered with black; the lower pair marked beneath by a transverse brown band.

### P. MEROPE.

*Cram.* t. 378. f. D. E.

---

This species is a native of several parts of Africa, and is represented on the plate in its natural size. In colour it sometimes varies a little.

N.Y.





## ANAS ÆGYPTIACA.



### CHARACTER GENERICUS.

*Rostrum* lainellosso-dentatum, convexum, obtusum.

*Lingua* ciliata, obtusa.

*Lin. Syst. Nat.* p. 194.

### CHARACTER SPECIFICUS, &c.

ANAS flavo-ferruginea, fusco-undulata, speculo alari candido, fascia nigra.

ANAS ÆGYPTIACA. A. rostro subcylindrico, corpore undulato, vertice albo, speculo alari albo fascia nigra.

*Lin. Syst. Nat.* p. 197.

---

Elegantem hanc speciem, in variis Africanis partibus generatam, abundantius alere dicitur Ægyptus. In Angliam jam olim illata est; commendatque eam tam venustas quam cicur et mansuetum ingenium. In avium numero est quarum imaginibus adornare soliti sunt antiqui Ægyptii sacras sculpturas; in obeliscis nonnullis præcipue notabilis, quos, inter alia vetera monumenta in Museo Britannico jam asservatos, a Gallis nuperrime eripuerunt celeberrimi nostri bellici ductores qui in Ægypto militarunt.

Magnitudo avis eadem fere est atque anseris communis.







THE  
EGYPTIAN GOOSE.

*GENERIC CHARACTER.*

*Bill* broad and flattened; the edges marked with sharp lamellæ.

*Tongue* broad and ciliated at the edges.

*SPECIFIC CHARACTER, &c.*

Yellowish-ferruginous GOOSE, with brown undulations, and white wing-coverts crossed by a black bar.

EGYPTIAN GOOSE.

*Lath. syn. 3. p. 453.*

L'Oye d'Egypte.

*Buff. 9. p. 79.*

*Pl. Enl. 379. 982. 983.*

---

This elegant species is found in several parts of Africa, and is said to be particularly plentiful in Egypt. In our own country it is often seen in a domestic state, and is highly esteemed on account of its agreeable appearance. It is one of those birds

which were frequently represented among the sacred sculptures of the ancient Egyptians, and is particularly observable on some obelisks lately brought from Egypt by our celebrated countrymen, and now deposited in the British Museum.

## PATELLA FUSCA.

\*\*\*\*\*

### CHARACTER GENERICUS.

*Animal* Limax.

*Testa univalvis, subconica, absque spira.*

*Lin. Syst. Nat. p. 1257.*

### CHARACTER SPECIFICUS, &c.

PATELLA testa subintegra ovata obtusa fusca,  
elevato-striata, intus alba.

PATELLA FUSCA. P. testa integerrima ovata,  
obtusa, striis elevatis.

*Lin. Syst. Nat.*


---

Maria Indica et Americana incolit Patella fusca,  
magnitudine vera in tabula depicta.





605



R.R.C.

THE  
BROWN PATELLA.



*GENERIC CHARACTER.*

*Animal* allied to a Limax or Slug.

*Shell* univalve, subconic, without spire.

*SPECIFIC CHARACTER, &c.*

PATELLA with nearly entire, ovate, brown,  
ribbed Shell; white within.

The brown Indian Limpet.

---

The Brown Patella is a native of the Indian and American seas, and is represented on the plate in its natural size.



# MADREPORA PEZIZOIDES.



## CHARACTER GENERICUS.

*Animal* Medusa.

*Corallium* cavitatibus lamelloso-stellatis.

## CHARACTER SPECIFICUS, &c.

MADREPORA subcærulea subfoliacca concava,  
supra stellis muricata, subtus aceroso-scabra.

MADREPORA cinerascens. M. subfoliacca  
explanata aggregata, subtus aceroso-sca-  
brosa, stellis remotiusculis elevatis, ambu-  
lacris scabrosis.

*Soland. et Ellis Zooph.* p. 157. t. 43.

---

E maribus Indicis extrahitur rara hæc Madrepora,  
interdum multo major quam in tabula depingitur.







28

THE  
PEZIZOID MADREPORE.



*GENERIC CHARACTER.*

*Animal* allied to a Medusa.

*Coral* marked by lamellar star-shaped cavities.

*SPECIFIC CHARACTER.*

Blueish subfoliaceous concave MADREPORE,  
muricated above with star-shaped pores,  
and of a chaffy-rough surface beneath.

---

This rare species of Madrepore is found in the Indian seas, and is sometimes much larger than the specimen figured on the present plate.



# PHALANGIUM DUBIUM.



## CHARACTER GENERICUS.

*Pedes* octo.

*Oculi* verticis duo contigui, duo laterales.

*Frons* antennis pediformibus.

*Abdomen* rotundatum.

*Lin. Syst. Nat.* p. 1028.

## CHARACTER SPECIFICUS.


PHALANGIUM fuscum, chelis minutis dorsa-  
libus.

---

In litoribus Americanis? generatur Phalangium dubium. A Donino Lathamo nacti sumus specimen quod depinximus. Ignotum credimus fuisse ipsum insectum, nec ab aliquo antea depictum.







THE  
DOUBTFUL PHALANGIUM.



*GENERIC CHARACTER.*

*Legs* eight.

*Eyes* two vertical and two lateral.

*Antennæ* resembling legs.

*Abdomen* rounded.

*SPECIFIC CHARACTER.*

Brown PHALANGIUM, with minute dorsal claspers.

---

This species is supposed to be a native of some of the American coasts. The specimen figured on the annexed plate was communicated by Mr. Latham. It is an insect which seems not to have been hitherto described or figured.



# PSITTACUS AUGUSTUS.



## CHARACTER GENERICUS.

*Rostrum* aduncum; mandibula superiore mobili; cera instructum.

*Nares* in rostri basi.

*Lingua* carnosa, obtusa, integra.

*Pedes* scandorii.

*Lin. Syst. Nat.* p. 130.

## CHARACTER SPECIFICUS, &c.

PSITTACUS macrourus cyaneus, rostro pedibusque nigris; orbitis basique mandibulæ inferioris luteis.

*Museum Leverianum.* No. 2. t. 2.

PSITTACUS hyacinthinus.

*Lath. ind. orn.* p. 84.

---

In Museo Leveriano asservatur avis hæc rarissima, quæ reliquis fere omniibus psittacini generis mole et magnificentia antecellere videtur. Tota avis est eximie cyanea, nisi quod super frontem et margines remigum levissima sit coloris thalassini tintura. Superficies inferior alarum caudæque ni-

gra est. Rostrum præter solitum magnum validumque omnino nigerrimum. Nigrant quoque crura et pedes, quorum ingens est robur. Orbitæ seu spatia nuda circum oculos coloris sunt lutei: lutea est etiam cutis nuda qua mandibulæ inferioris basis cingitur. Partium corporis proportio eadem fere est cum Psittaco Ararauna. In iisdem quoque Americæ Australis partibus fortasse generatur. Præter specimen quod descripsimus vix aliud in tota Europa inveniri dicitur.





THE  
HYACINTHINE MACAW.



*GENERIC CHARACTER.*

*Bill* hooked : upper mandible moveable, and furnished with a cere.

*Nostrils* round, placed at the base of the bill.

*Tongue* fleshy, broad, blunt at the end.

*Feet* scansorial.

*SPECIFIC CHARACTER, &c.*

Long-tailed deep-blue MACAW, with the bill and legs black ; the orbits and base of the lower mandible yellow.

HYACINTHINE MACAW.

*Museum Leverianum.* No. 2. pl. 2.

*Lath. ind. orn.* p. 84.

---

This extremely rare bird, which surpasses in the magnificence of its appearance almost all the rest of the Parrot tribe, is preserved in the Leverian Museum. The whole bird is of a very fine deep blue, except that the forehead and edges of the wing-feathers have a slight cast of sea-green. The under

surface of the wings and tail is black. The beak, which is uncommonly large and strong, is of a deep black, as are also the legs and feet, which are extremely stout. The orbits or naked spaces round the eyes are of a deep yellow, as is also the skin round the base of the lower mandible. The general proportions of the bird are nearly the same as those of the *Psittacus Ararauna*, and it is probably a native of the same parts of South America. The specimen now described is perhaps the only one known to exist at present in Europe.

# OVIS CANADENSIS.



## CHARACTER GENERICUS.

*Cornua* concava, retrorsum versa, intorta, rugosa.

*Dentes* primores inferiores octo.

*Laniarii* nulli.

## CHARACTER SPECIFICUS, &c.

OVIS pilosa fusco-ferruginea, fronte uropygio quoque albis, cauda brevissima, cornibus compressis lunatis.

Belier de Montagne.

*Geoffr. Ann. Mus. Nat.* No. 11. p. 360.

---


Physicis Europaeis non nisi imperrime innotuisse videtur Ovis quam describere pergitus, in Canada interiori generata, nec lana vestita, sed pilo denso, valido, et quasi cervino. Crura, si cum corpore comparentur, longa et gracilia. Cornua iis simillima quæ gerit aries nostras vulgaris: at foeminae minoræ longe quam mari. Color generalis fusco-ferrugineo-pallidus, idem fero ac cernere est in multis cervini generis. Genæ obscuriores. Rostrum et uropyginum alba. Cauda brevissima. Idem fere dicitur esse huic vivendi modus qui Ibici; scandenti nempe montium cacumina, et mira celeritate de rupe

rupe in rupem salienti. Ovium hujusmodi congregantur, ut plurimum, viginti seu triginta ; vocantque eas *Canadenses oves rupestres*. Juniorum caro in cibis lautissimis habetur. Ovi Canadensi eadem fere est magnitudo atque Ibici. Per pulchrum rari hujus quadrupedis specimen in Museo Britannico conspicitur.





THE  
CANADIAN SHEEP.



*GENERIC CHARACTER.*

*Horns* hollow, wrinkled, turning backwards,  
and spirally intorted.

*Front-Teeth* eight in the lower jaw.

*Canine-Teeth* none.

*SPECIFIC CHARACTER, &c.*

Ferruginous-brown hairy SHEEP, with white  
front and rump, very short tail, and com-  
pressed lunate horns.

Belier de Montagne.

*Geoffr. ann. mus. nat.* No. 11. p. 360.

---


The species of sheep here represented, and which appears to have been, till very lately, unknown to the naturalists of Europe, is a native of the interior parts of Canada. It is remarkable for being covered, instead of wool, with very thick and strong hair, greatly resembling that of a Deer. The legs are long in proportion to the body. The horns very much resemble

seimble those of the common ram, and those of the female are said to be much smaller than those of the male. The general colour is a pale ferruginous brown, similar to that of many of the Deer tribe : the cheeks are of a darker cast than the other parts, and the muzzle and rump are white : the tail is very short. The general habits of the animal are said to resemble those of the Ibex, frequenting chiefly the highest and most inaccessible parts of the mountainous regions, occasionally skipping from rock to rock with incredible swiftness. It is generally observed in small flocks of twenty or thirty together, and is known to the Canadians by the name of *Mountain Sheep*. The young are considered as the most delicate meat which that extensive country can afford. A very fine specimen of this rare quadruped may be seen in the British Museum.





# PHALÆNA REGIA.



## CHARACTER GENERICUS.

*Antennæ* setaceæ, a basi ad apicem sensim attenuatæ.

*Alæ* sedentis sæpius deflexæ. (*Volatu nocturno.*)

*Lin. Syst. Nat.* p. 808.

## CHARACTER SPECIFICUS, &c.

PHALÆNA (Bombyx) elinguis, alis deflexis griseo-fuscis flavo maculatis fulvoque nervosis; posticis corporeaque fulvis.

*Smith. Abbot. Ins. Amer.* p. 121. t. 61.

Bombyx regalis.

*Fabr. ins. emend.* vol. 4. p. 436.

PHALÆNA Laocoön.

*Cram. pap.* t. 117. B. C. ?

---


Quamvis crucam seu larvam pulcherrimi hujus insecti satis accurate delineaverunt Catesbeius aliqui, ipsam tamen phalænam primo depictam esse arbitror in Crameri pagina. Quod autem animalculi omnes permutationes subeuntis plena ad nos notitia pervenerit, debetur laudabili industriæ Domini Abbotti, qui opus nuperime edidit insoliti splendoris

de

de insectis Transatlantieis, doctis præterea ditatum  
observationibus Domini Smithii, M. D. Societatis  
Linnæanæ Londinensis præsidis. Docet Dominus  
Abbotus, erucam, (nomine *diaboli cornuti* vulgo  
dictam) folia Diospyri Virginianæ et Juglandis  
albæ præcipue depascere, et tela tenui involutam in  
chrysalidem converti mense Junio, mense autem Ju-  
lio erumpere phalænam.

Notavit recte omnino Dominus Smithius phalæ-  
nam hanc phalænæ nostræ Europææ *Ph: Humuli*  
Linnæi (Ghost-Moth) habitu generali evidenter esse  
affinem.

Variæ insecti vices ostenduntur in tabula juxta  
naturalem magnitudinem.





THE  
PERSIMON MOTH.



*GENERIC CHARACTER.*

*Antennæ* setaceous, decreasing in size from base to point.

*Wings* (when at rest) generally deflected. *Flight* chiefly nocturnal.

*SPECIFIC CHARACTER, &c.*

MOTH with deflected grey-brown wings, with yellow spots and fulvous nerves : the lower wings and body fulvous.

The PERSIMON MOTH.

*Abbot's American Insects.* p. 121. pl. 61.

---

The caterpillar of this highly-elegant insect has long ago been figured with sufficient accuracy by Catesby and others, but the Moth seems to have been first figured in the work of Cramer. For the complete knowledge of the insect in all its states we are indebted to the laudable industry of Mr. Abbot, whose researches on the Transatlantic Insects, assisted by the learned and accurate annotations of Dr. Smith, president of the Linnæan Society of

of London, have lately appeared with such superior splendour. Mr. Abbot informs us that the caterpillar (which is popularly known by the title of the Horned Devil) feeds principally on the leaves of the Persimon and the Hiccor, and that it changes into a chrysalis in the month of June, (first enveloping itself in an oval web,) the moth appearing in the month of July. Dr. Smith, in his annotations, has very judiciously added, that in point of general habit this moth is evidently allied to the European species called the Ghost-Moth (*Phalaena Humuli* Lin.) The plate represents the animal in its several changes, and in its natural size.

# MEROPS CÆRULOCEPHALUS.



## CHARACTER GENERICUS.

*Rostrum compressum, curvatum, carinatum.*

*Lingua apice laciniata.*

*Pedes gressorii.*

*Lin. Syst. Nat. p. 182.*

## CHARACTER SPECIFICUS, &c.

MEROPS RUBER, capite gula uropygioque  
cæruleo-viridibus.

MEROPS NUBICUS. M. ex cæruleo viridis  
subtus ruber, dorso, alis, candaque furcata  
lateritiis.

*Lin. Syst. Nat. Gmel. p. 464.*

---

Africæ partes interiores incolit Merops Cærulocephalus: magnitudo avis eadem fere est atque Meropis communis sive Europææ. Variat cauda sub-integra vel forcipata.

G







THE  
BLUE-HEADED BEE-EATER.



*GENERIC CHARACTER.*

*Bill* compressed, curvated, and carinated.  
*Tongue* generally laciniated at the tip.  
*Feet* formed for walking.

*SPECIFIC CHARACTER, &c.*

RED BEE-EATER, with the head, throat, and rump blue-green.

BLUE-HEADED BEE-EATER.

*Lath. Syn.* 1. p. 680.

Le Gnèpier rouge à tête bleue.

*Buf. ois.* 6. p. 506.

Gnèpier de Nubie.

*Pl. Ent.* 649.

---

This bird is a native of the interior parts of Africa, and is nearly similar in size to the common or European Bee-Eater. It varies in having the tail either even or slightly forked.



# ARANEA VENATORIA.



## CHARACTER GENERICUS.

*Pedes* octo.

*Oculi* octo.

*Os unguibus seu retinaculis* duobus.

*Palpi* duo articulati; masculis genitalibus capitati.

*Anus* papillis textoriis.

*Lin. Syst. Nat.* p. 1030.

## CHARACTER SPECIFICUS, &c.

ARANEA CASTANEA, thorace glabro, abdomine pubescente fusco.

ARANEA VENATORIA. A. thorace orbiculato glabro atro, abdomine ovato pubescente fusco.

*Lin. Syst. Nat. Gmel.* p. 2960.

*Fabr. sp. ins.* 1. p. 546.

---

Jamaican insulasque alias Americanas incolit Aranea venatoria, in locis siccis et arenosis nidum struens cylindraceum, quinque vel sex uncias longum, et operculo seu valvula instructum, quæ ad libitum resilit et clauditur. In hoc latibulo prædæ insidiatur, insectis nempe prope repentibus aut volantibus, quæ saltu subito arrepta et in nidum subducta, clauso operculo, devorat. Ostenditur in tabula animalis vera magnitudo; nidus autem magnitudine diminuta conspicitur.





614



PLATE 101

THE  
AMERICAN TARANTULA.



*GENERIC CHARACTER.*

*Legs* eight. -

*Eyes* eight.

*Mouth* furnished with two hooks or holders.

*Palpi* or *Feelers* two, the tips of which (in the males) distinguish the sex.

*Abdomen* terminated by papillæ or teats thro' which the insect draws its thread.

*SPECIFIC CHARACTER, &c.*

CHESNUT SPIDER, with smooth thorax, and brown pubescent body.

BLACK TARANTULA. *Brown's Jamaica.*

AMERICAN TARANTULA.

---

The Aranea venatoria inhabits Jamaica, and other American islands: it constructs in dry and sandy situations a cylindrical nest of five or six inches in length, furnished with an elastic closing valve. In this nest the animal waits the arrival of its prey, viz. any insect flying or creeping near; and suddenly rushing out, secures its victim, which it drags into the nest, and devours at leisure, the valve immediately closing behind it. In the annexed plate the spider itself is represented in its natural size; the nest of half its real dimensions.



# PHALÆNA CECROPIA.



## CHARACTER GENERICUS.

*Antennæ* setaceæ, a basi ad apicem sensim attenuatae.

*Alæ* sedentis saepius deflexæ. (Volatu nocturno).

*Lin. Syst. Nat.* p. 808.

## CHARACTER SPECIFICUS, &c.

PHALÆNA alis subfalcato-rotundatis griseofuscis, fascia ferruginea maculaque media fenestrata.

PHALÆNA CECROPIA. P. alis subfalcatis griseis, fascia fulva, primoribus ocello subfenestrato ferrugineo.

*Lin. Syst. Gmel.* p. 2401.

*Catesb. Carol.* 2. t. 86.

*Cram. pap.* 4. t. 42. f. A. B.

---


In variis Americæ borealis partibus conspicitur pulcherrima hæc Phalæna. Operi splendidissimo Domini Abboti de insectis Americanis effigiem debemus larvæ et pupæ.



7. 1653







R. P. Nodder

THE  
CECROPIAN MOTH.



*GENERIC CHARACTER.*

*Antennæ* setaceous, decreasing in size from base to point.

*Wings* (when at rest) generally deflected.  
Flight nocturnal.

*SPECIFIC CHARACTER, &c.*

GREY-BROWN MOTH, with rounded subfalcated wings, marked by a ferruginous band and semitransparent middle spot.

*Seb. Mus.* 4, *t.* 58, *f.* 4, 5?

*Drury.* 1, *pl.* 18, *f.* 2.

---

This highly beautiful insect is found in many parts of North America. It is to the splendid publication of Mr. Abbot on the American insects that we are indebted for the figure of the larva and pupa.



# TURDUS CYANURUS.



## CHARACTER GENERICUS.

*Rostrum* tereti-cultratum, mandibula superiore apice deflexo, emarginato.

*Nares* nudæ, superne membranula semitectæ.

*Fauæ* ciliata.

*Lingua* lacero-emarginata.

*Lin. Syst. Nat.* p. 291.

## CHARACTER SPECIFICUS, &c.

TURDUS SUBFERRUGINEUS subtus flavus, abdomine cœruleo lineato, cauda cyanea.

TURDUS CYANURUS. T. ruso-fuscus, subtus flavus, abdomine cœruleo fasciato, pileo lateribusque colli striga longitudinali nigra, fascia pectorali caudaque cœruleis.

*Lath. ind. orn.* p. 361.

---

In avibus quas plurimas alit America australis, haud facile elegantiorem reperies Turdo cyanuro. Caput fasciis nigris flavisque lateralibus decoratur :

H

dorsum

dorsum humerique nitidissime castanea : alarum remiges nigri ; tectricum apicibus albis fasciam concolorem per medium quasi alæ ducentibus. Tota avis inferior a gula pulchre flavet, lineis plurimis transversis lâte cernulis variata. Cauda brevis et acutula splendidissime cyanea est, pennis lateralibus a mediis duabus gradatim utrinque decrescentibus. Rostrum nigrum, pedes fusi. Avem hanc raritate mirabilem jactant per pauca musea. In Cayenna præcipue generari dicitur *Turdus cyanurus*.





NATIONAL MUSEUM OF VICTORIA

THE

## BLUE-TAILED THRUSH.



### GENERIC CHARACTER.

*Bill* strait, obtusely carinated at the top, bending a little at the point, and slightly notched near the end of the upper mandible.

*Nostrils* oval and naked.

*Tongue* slightly jagged at the end.

*Middle-toe* slightly connected to the outer as far as the first joint.

### SPECIFIC CHARACTER, &c.

SUBFERRUGINOUS THRUSH, yellow beneath, with deep blue tail, and abdomen marked by numerous blue lines.

L'Azurin. *Buff. ois.* 3. p. 410.

Merle de la Guiane. *Pl. Enl.* 355.

---

The Blue-Tailed Thrush may be considered as one of the most elegant of the South-American birds.

birds. The head is ornamented on each side by black and yellow stripes: the back and shoulders are of a bright chesnut colour: the wing-feathers black: the coverts tipped with white, forming a band of that colour across the middle of each wing: the whole under part of the bird, from the throat, is of a bright yellow, marked with numerous bright-blue transverse lines: the tail is short, and of a slightly sharpened shape, the feathers gradually shortening on each side from the middle ones: the bill is black and the legs brown. This highly rare and beautiful bird is but rarely seen in European museums, and is principally found in Cayenne.

## PHALÆNA STRIX.



### CHARACTER GENERICUS.

*Antennæ* setaceæ, a basi ad apicem sensim attenuatæ.

*Alæ* (sedentis) sæpius deflexæ. (Volatu nocturno.)

*Lin. Syst. Nat.* p. 808.

### CHARACTER SPECIFICUS, &c.

PHALÆNA ALBIDA, alis concoloribus fusco nigroque reticulatis nebulosisque

PHALÆNA STRIX? P. alis concoloribus albo nigroque reticulatis nebulosisque? *Lin. Syst. Nat. Gmel.* p. 2529. *Cram. pap.* vol. 2. t. 145. f. A.

---

Insulas Indieus, præcipue Javam et Amboynam incolit Phalæna Strix, magnitudine vera in tabula depicta.







## STRIX.



### *GENERIC CHARACTER.*

*Antennæ* setaceous, decreasing in size from base to point.

*Wings* (when at rest) generally deflected.

*Flight* nocturnal.

### *SPECIFIC CHARACTER, &c.*

WHITISH MOTH, with the wings clouded and reticulated by black and brown variegations.

INDIAN COSSUS or Owl-Moth.

---

The Phalæna Strix is a native of the Indian islands, and particularly of Java and Amboina. The plate represents it in its natural size.



# CYPRÆA ARGUS

ET

## CYPRÆA LEUCOPIS.



### CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, involuta, subovata, obtusa,  
lævis.

*Apertura* utrinque effusa, linearis, utrinque  
dentata, longitudinalis.

*Lin. Syst. Nat.* p. 1172.

### CHARACTER SPECIFICUS, &c.

CYPRÆA ARGUS. C. subferruginea, ocellis  
subconvexis fuscis, subtus maculis quatuor  
fuscis.

CYPRÆA ARGUS. C. testa subturbinata sub-  
cylindrica, adpersa ocellis, subtus maculis  
quatuor fuscis.

*Lin. Syst. Nat.* p. 1173.

Argus magnus. *Regenf.* t. 5. f. 57.

Porcellana major, Argus. *Rumph.* t. 38. d.

Argus mas. *Martini.* 1. p. 363. t. 28. f. 285. 286.

*CHARACTER SPECIFICUS, &c.*

CYPRÆA LEUCOPIS. C. subferruginea ocel-  
lis albis.

CYPRÆA EXANTHEMA? C. testa subturbi-  
nata ferruginea, maculis pallidis rotundis  
adspersa, linea longitudinalis subramosa.

• *Lin. Syst. Nat. p. 1172?*

ARGUS pupilla fusca in iride alba. *Martini* 1.  
p. 368. *t. 28. f. 289. & t. 29. f. 298. 299.*

---

E maribus Indicis præcipue extrahuntur rara hæc  
et pulchra conchylia, interdum majora quam in  
tabula ostenduntur.



THE  
ARGUS COWRY  
AND  
WHITE-EYED COWRY.

\*\*\*\*\*

GENERAL DESCRIPTION.

Animal resembling a Linnet or Spar.  
Shell univalve, smooth, opaque, orange.  
Aperture linear, longitudinal, rounded on both  
sides.

SPECIFIC DESCRIPTION. No.

ARGUS COWRY. Subtriangular Cowry, with  
subcircular brown, eye-shaped spots, and  
four brown patches beneath.

Le Grand Argus. Argus. pl. 15. 2.

Le Double Argus. Double Cowry. pl. 15. 3.

SPECIFIC DESCRIPTION. No.

WHITE-EYED COWRY. Subtriangular  
Cowry, with white, eye-shaped spots.

Le Petit Argus. Double.

Le Variable Argus. Double Cowry. pl. 15. 4.

---

The two last named shells are rarely found  
in the Indian seas and are sometimes seen at a  
large distance separated from the present place.

619





# ASTERIAS OLIGACTES.



## CHARACTER GENERICUS.

*Corpus* *depressum*; *crusta* *coriacea*, *tentaculis* *muricata*.

*Os centrale*, *quinquevalve*.

*Lyn. Syst. Nat.* p. 1098.

## CHARACTER SPECIFICUS, &c.

ASTERIAS ALBO-FLAVESCENS, *radiis longissimis simplicibus*.

ASTERIAS OLIGACTES. A. *radiis longissimis simplicibus*; *articulis singulis basi stylis binis mobilibus acentis*.

*Lin. Syst. Nat. GMEL.* p. 3167.

*Pall. nov. act. Petrop.* 2. p. 239.


---

A celeberrimo Pallas in Novis Actis Academæ Petropolitanae ita describitur Asterias oligactes.

" *Corpus* *calum duriusculum*, *exignum*, *magnitudine ea quam figura exprimit*, *pentagonum*, *angulis truncatis*; *superiore facie medio impressa*, *stellataque costis rotundatis denis*, *per paria versus angulos truncatos subparallelis*, *extrorsum crassescientibus*;  
inferiorc

inferiore facie plana, ore in medio ramis linearibus  
disceisso, stellato et ad ortum radiorum fissura utrin-  
que obliqua incisa. Radii proportione corporis  
enormi longitudine ad quindecim pollices et ultra  
explere visi, quantum mensurare filo potuere, tereti-  
filiformes, lentissime adtenuati, eompositi artieulis  
creberrimis, crustaceis, osseis, consistentia et colore  
ut in Asteria Medusæ. Singuli artieuli ad latus ori  
respondens instructi pedunculis seu stylis binis mobi-  
libus, ipso artieulo vix longioribus, approximatis.  
Color totius albo-flavescens, consistentia dura,  
crustacea."





THE  
SLENDER-LIMBED STAR-FISH.



*GENERIC CHARACTER.*

*Body* depressed; covered with a coriaceous crust, muricated with tentacula.

*Mouth* central, five valved.

*SPECIFIC CHARACTER, &c.*

**YELLOWISH WHITE STAR-FISH**, with very long, simple rays; the joints furnished at the base with two moveable styles.

**SLENDER-RAYED ASTERIAS.**

---

This curious species of Star-Fish is described by the celebrated Dr. Pallas in the New Petersburg Transactions. It is of the size represented on the plate, and consists of a small, hardish, pentagonal body with truncated angles: the upper surface is marked in a stellated manner by ten ribs or elevations disposed by pairs towards the corners: these ribs are somewhat thicker towards the corners than towards the middle of the body: the mouth is central,

tral, star-shaped and divided by linear processes: the rays or limbs are of enormous length in proportion to the body, measuring more than fifteen inches, and are of a rounded, thread-like appearance, tapering very gradually to their extremities, and consist of very numerous, crustaceous or bony joints, of similar consistence to those of the *Asterias Medusa*: each joint, on the side or surface answering to the mouth, is furnished with two moveable styles or processes, scarce longer than the joint itself: the whole animal is of a yellowish white colour, and of a hard, crustaceous substance.

## ARDEA TIGRINA.



### CHARACTER GENERICUS.

*Rostrum rectum, longum, acutum.*

*Nares lineares.*

*Lingua acuminata.*

*Pedes tetradactyli.*

### CHARACTER SPECIFICUS, &c.

ARDEA ferruginea, fasciis transversis nigris.

ARDEA tigrina. A. capitis levis vertice, caudaque albo-fasciata nigris, corpore nigromaculato supra rufa subtus ochrolenca, mento et crisso albis.

*Lin. Syst. Nat. Gmel. p. 638.*

ARDEA TIGRINA. A. maculis nigris difformibus notata, supra rufa subtus albida, vertice candaque nigris, rectricibus fasciis quatuor albis.

*Lath. ind. orn. p. 682.*

---

Partes Americæ Australis calidiores incolit Ardea  
tigrina,

tigrina, loca paludosa amans, moreque congenerum  
pisciculis, ranis, aliisque ejusmodi victitans: mag-  
nitudine Ardeam stellarem fere æquat.





THE  
TIGER BITTERN.



*GENERIC CHARACTER.*

*Bill* strait, long, sharp-pointed.

*Nostrils* linear.

*Tongue* pointed.

*Feet* tetractylous.

*SPECIFIC CHARACTER, &c.*

Ferruginous BITTERN, with transverse black bands.

TIGER BITTERN.

*Lath. syn.* 3, p. 63.

Heron tigré.

*Fermiu Surin.* 2, p. 151.

L'Onoré.

*Buff. ois.* 7, p. 431.

*Pl. Ent.* 790.

---

The Tiger Bittern is a native of the warmer regions of South-America, frequenting marshy places, and

preying, like the rest of this genus, on small fishes,  
frogs, &c. In size it nearly equals the common  
Bittern.

# SOLPUGA VENENOSA.



## CHARACTER GENERICUS.

*Mandibulæ maximæ, porrectæ, forcipatæ.*

*Labium inferius porrectum, subulatum.*

*Palpi elongati, pedibus anterioribus basi connecti.*

## CHARACTER SPECIFICUS, &c.

SOLPUGA tomentosa fusca, corpore ovato.

SOLPUGA arachnoides.

*Herbst. apt. p. 37. t. 1. f. 2.*

SOLPUGA araneoides.

*Fabr. suppl. entom. p. 294.*

Phalangium araneoides.

*Pall. sp. zool. 9. p. 37. t. 3. f. 8. g.*

---

In genere *Solpuga*, cui cum Phalangio et Aranea multa videtur esse affinitas, continentur plures species. Harum fortasse princeps haberi possit, cuius veram magnitudinem cernere est in tabula. Generant eam Europæ Australis Africæque loca calidiora.

In

In agris versatur, morsuque cutem irritando, tumores excitare solet acerbe molestos, interdum lethales. A celeberrimo Herbstio in opere suo præstanti de insectis apteris primo institutum est genus *Solpuga*.





THE  
POISONOUS SOLPUGA.

---

*GENERIC CHARACTER.*

*Mandibles* very large, stretched forwards, for-  
cipated at the tips.

*Lower lip* stretched forwards, subulate.

*Feelers* elongated, connected at their base with  
the fore-legs.

*SPECIFIC CHARACTER, &c.*

Brown, downy SOLPUGA, with ovate abdo-  
men.

Galeode araneoide.

*Olivier Nouveau Dict. D'Histoire Naturelle.*

Sonnini Voy. en Grece. 2. p. 115.

---

The genus Solpuga, much allied to those of Pha-  
langium and Aranea, contains several species, of  
which that represented in its natural size on the  
plate may perhaps be considered as the principal. It  
is a native of the warmer parts of the South of  
Europe, and of Africa, inhabiting fields, and by its  
bite producing most painful swellings on the skin,  
and

and even, as it is said, sometimes proving fatal. The genus *Solpuga* seems to have been first instituted by Mr. Herbst, in his excellent work on the Apteroous Insects.

# MADREPORA FILEUS.



## CHARACTER GENERICUS.

*Animal* Medusa.

*Corallium* cavitatibus lamelloso-stellatis.

*Lin. Syst. Nat.*

## CHARACTER SPECIFICUS, &c.

MADREPORA ovato-elongata albida, sulco longitudinali medio.

MADREPORA PILEUS. M. simplex acanlis oblonga, stella convexa conglomerato-lamellosa, lamellis abbreviatis, subtus concava.

*Lyn. Syst. Nat.* p. 1273.

Mitra Neptuni.

*Tournef. act. Paris.* 1700. p. 30.

Pileus Neptuni.


*Seb. mus.* 3. t. 111. f. 3. 5.

---

In oceano Indico reperta, quam descriptimus, Madrepora, major saepe est iconē quæ ostenditur in tabula. Color albo-flavescit.







## OBLONG MADREPORE.



### GENERIC CHARACTER.

*Animal* allied to a Medusa.

Coral marked by radiating lamellar cavities.

### SPECIFIC CHARACTER, &c.

Oblong whitish MADREPORE, with a longitudinal central furrow.

Mitra Polonica.

*Rumph. Amb.* 6. p. 248.

Oblong white MADREPORE.

---

The Madrepore figured on the present plate is a native of the Indian seas, and is frequently found much larger than here represented: its colour is yellowish white.



# ERPETON TENTACULATUS.



## CHARACTER GENERICUS.

*Corpus caudaque squamosa.*

*Abdomen scutatum.*

## CHARACTER SPECIFICUS, &c.

ERPETON rostro utrinque tentaculato.

ERPETON TENTACULATUS. *Cepede.*



Ab illustri Cepede institutum genus Erpeton ex unica constat specie quae insulas Indicas creditur incolere, quamque insigniunt tentacula duo, sed quasi cornua, super labium superius sita, parvulis squamis vestita, nec unquam retracta. Bipedali quasi longitudine est Erpeton tentaculatus, venuci expers, dentes habens minimos, eodem modo dispositos quo dentes serpentum innocuorum. Cumam viveudi modo se praecipue assuescat, non satis est comprehendim.

Specimen quod descripsit Cepede in museo Principis Aransiaci asservatur.







# TENTACULATED ERPETON.



## GENERIC CHARACTER.

*Body and tail scaly.*

*Abdomen lamellated.*

## SPECIFIC CHARACTER, &c.

ERPETON with the upper lip tentaculated on each side.

ERPETON tentaculé.


*Cepede Ann. Mus. National D'Hist. Nat.*

No. 10. p. 280. pl. 1.

---

The genus Erpeton, instituted by the Count de Cepede, consists at present of a single species, which is supposed to be a native of the Indian islands, and is remarkable for a pair of tentacula or elongated processes situated above the upper lip: these tentacula are covered with very minute scales, and are not retractile. The animal measures about two feet in length, and is not of a poisonous nature, the teeth, which are very small, resembling those of innoxions serpents. Of its particular habits or history nothing seems to be known.

The specimen described by Cepede belonged to the Museum of the Prince of Orange.



# PARADISEA MAGNIFICA.



## CHARACTER GENERICUS.

*Rostrum* capistri plumis tomentosis tectum.

*Pennæ* hypochondriorum longiores (plerisque.)

*Rectrices* duæ superiores singulares denudatæ.

## CHARACTER SPECIFICUS, &c.

PARADISEA rufo-castanea, subtus viridi-nitens, collo utrinque pennis flavis fasciulato.

PARADISEA MAGNIFICA. P. supra spadicea, gula viridi lunulis aurcis, cervicis fascieulo pennarum flavarum.

*Lin. Syst. Nat. Gmel.* p. 401.

---

Paradiseam magnificam commendat formæ elegantia et versicolor plumarum varietas. In insulis Moluccanis præcipue generatur, magitudine Paradiseæ apodæ paulo inferior.







THE  
MAGNIFICENT PARADISEA.



*GENERIC CHARACTER.*

*Bill* surrounded at the base by velvet-like feathers.

*Hypochondrial feathers* (in most species) long and loose.

*Tail-feathers* two upper naked or unwebbed.

*SPECIFIC CHARACTER, &c.*

RUFOUS-CHESNUT PARADISE-BIRD, Iridescent green beneath, with fasciculated yellow plumes on each side the neck.

MAGNIFICENT BIRD of PARADISE.

*Lath. Syn.* 1, p. 477.

Oiseau de Paradis, de la Nouvelle Guinée dit le Magnifique.

*Pl. Ent.* 631.

---

This elegant species, so remarkable for the splendor and variety of its colours, is principally found in the Molucca islands, and is of somewhat smaller size than the *Paradisea apoda* or common Paradise-Bird.



# SIMIA ASCANIUS.



## CHARACTER GENERICUS.

*Dentes primores* utrinque quatuor, approximati.

*Laniarii* solitarii, longiores, hinc remoti.

*Molares* obtusi.

## CHARACTER SPECIFICUS, &c.

SIMIA caudata, barbata, olivacea, subtus grisea, facie violacea, naso albo.

SIMIA ASCANIUS. S. caudata barbata, facie subcærulea, naso albo.

*Audeb. Singes. fam. 4. sect. 2. f. 13.*

---

Simiam Ascanium primus descriptsse videtur Dominus Audebertius, in eleganti opere, cui titulus “*Histoire des Singes, &c.*” Eadem fere est magnitudine atque Simia Sabæa Linnæi, et indole esse dicitur alacri, sed mansueta. De patria ambigitur; Africanam tamen crediderim.







# ASCANIUS.



## GENERIC CHARACTER.

*Front-teeth* in each jaw four, placed near together.

*Canine-teeth* solitary, longer than the others, distant from the grinders.

*Grinders* obtuse.

## SPECIFIC CHARACTER, &c.

Long-tailed, bearded, olive-coloured Monkey, grey beneath, with violaceous face, and white nose.

L'Ascagne.

*Audeb. Singes.*

---

The animal represented on the present plate seems to have been first described by Monsr. Audebert, in his elegant publication entitled "*Histoire des Singes*," &c. In size it nearly equals the Simia Sabæa or green monkey, and is said to be of a very lively, but gentle disposition. Its native country is not certainly known, but it is probably an African species.



# ONISCUS ENTOMON.



## CHARACTER GENERICUS.

*Antennæ* setaceæ.

*Palpi* inæquales, posteriores longiores.

*Corpus* ovale.

*Pedes* quatuordecim.

## CHARACTER SPECIFICUS, &c.

**ONISCUS** antennis quaternis, cauda oblonga acuta.

*Fabr. spec. ins.* 1. p. 375.

**ONISCUS ENTOMON.** O. abdomine subtus nudo, cauda subulata.


*Lin. Syst. Nat.* p. 1060.

---

Marina hæc species, Oniscorum Britannicorum maxima, adhæret plerumque piscibus majoribus. Color viventi livido-fusca, exsiccatæ obscuro-albidus.







THE  
SHARP-TAILED ONISCUS.



*GENERIC CHARACTER.*

*Antennæ* setaceous.

*Feelers* unequal, the hinder longest.

*Body* oval.

*Legs* fourteen.

*SPECIFIC CHARACTER, &c.*

Pale brown ONISCUS, with four antennæ, and  
oblong, sharpened tail.

SQUILLE ENTOMON.

*Degeer.* 7. p. 514.

---

This insect, the largest of the British Onisci, is a marine species, and is commonly found attached to the bodies of the larger fishes: its colour, when living, is pale brown, which fades in the dried animal into dull white.



## CANCER LINEARIS.



### CHARACTER GENERICUS.

*Pedes* octo, (rarius sex aut decem;) insuper  
manus duæ chelatae.

*Oculi* duo, distantes, plurimis pedunculati, elongati, mobiles.

*Cauda* articulata, inermis.

### CHARACTER SPECIFICUS, &c.

CANCER corpore linearis, manibus monodactylis.

CANCER LINEARIS. Cancer manibus quatuor monodactylis, pedibus decem.

*Fabr. sp. ins.* t. p. 517.

CANCER LINEARIS. C. macrourus articulatis, manibus quatuor monodactylis, pedibus decem.

*Linn. Syst. Nat.* p. 1056.

---

Circa littora varia Europaꝝ invenitur perpusilla  
haec Canceris species. Magnitudine vera nec non  
microscopio aucta in tabula annexa exprimitur.







22. *Leptoceras*

THE  
LINEAR CRAB.



GENERIC CHARACTER.

*Legs* generally eight, (in some species six or ten,) besides two claspers or chelated arms.

*Eyes* two, commonly distant, footstalked, moveable.

*Tail* articulated.

SPECIFIC CHARACTER, &c.

LINEAR-BODIED CANCER, with monodactylous feet.

ONISCUS SCOLOPENDROIDES?

*Pall. sp. zool. g. t. 4. f. 15.*

CHEVROLLE LINEAIRE,

*Latreille.*

---

This very small species of Crab is found about several of the European coasts. It is represented on the annexed plate both in its natural size, and as it appears when viewed by a microscope.



# MUSCICAPA MELANOLEUCA.



## CHARACTER GENERICUS.

*Rostrum* subtrigonum, utrinque emarginatum,  
apice incurvo: *vibrissæ* patentes versus  
fauces.

*Nares* subrotundæ.

*Lin. Syst. Nat.* p. 324.

## CHARACTER SPECIFICUS, &c.

MUSCICAPA NIGRA, dorso albo, femoribus  
albo nigroque annulatis, rectricibus albis  
apice nigris.

*Lath. ind. orn.* p. 469.

MUSCICAPA MELANOLEUCA.

*Guldenst. nov. comm. Petrop.* t. 19. p. 468.

---

Primus accurate descripsit hanc speciem Dominus  
Guldenstadt in Academiae Petropolitanæ Commen-  
tariis.

“ Longitudo avis extensæ ab apice rostri ad  
caudæ extremum sex pollicum, trium linearum :  
capistrum, caput infra usque ad medium collum et

L

ad

ad latera usque ad supercilia atrum; sed caput supra et corpus totum niveum, pectore levissime flavescente; basi obtecta pennarum omnium fusca; alæ totæ supra infraque atræ, apicis margine remigum secundariarum obsolete albido; complicatæ ultra mediam caudam procedentes: cauda integra, duos pollices et sex lineas longa; rectrices duodecim, albæ, apice nigræ, eoque magis quo interiores, attamen nec in medio pari nigredo ad medietatem accedit: femora usque ad genu plumosa, fusco et albido annulata; tibiæ decem lineas longæ, atræ-læves; digitæ pariter atri, quarum tres antici, quartus posticus, onines nigri, medio reliquis subæquilibus aliquantum longiore; ungues incurvi, acuti, nigri, subæquales. In femina ea quæ in mare nigra, fusca; et quæ in mare alba, sordide cinerea sunt. Habitat per æstatem in Georgia campestri ad fluviorum ripas fruticetis obsitas.





THE  
GEORGIAN FLYCATCHER.



*GENERIC CHARACTER.*

*Bill* somewhat triangular, flattened at the base, notched at the end of the upper mandible and beset with bristles.

*Toes* generally divided as far as their origin.

*SPECIFIC CHARACTER, &c.*

Black FLYCATCHER with white back, thighs annulated with black and white, and tail-feathers tipped with black.

Black-and-white WARBLER.

*Lath. Syn.* 2. p. 457.

---

The present species of *Muscicapa* seems to have been first accurately described in the Petersburg Transactions, by Mr. Guldenstadt, who informs us that it measures six inches and three lines from the tip of the bill to the end of the tail: the frontlet and the head beneath, as far as the middle of the

neck, and on each side as far as the eyebrows, is black; but the upper part of the head, as well as the whole body, is white, with a very slight cast of yellow on the breast; the covered parts or bases of all the feathers being brown: the whole wings, both above and beneath, are black, the edges of the tips of the secondary feathers whitish: the wings, when closed, reach beyond the middle of the tail, which is entire, and about two inches and six lines in length: the twelve tail-feathers are white: with black tips, that colour increasing gradually on the interior feathers; but even in the two middle feathers not reaching half way up the tail: the thighs are feathered as far as the knees, and annulated with brown and whitish: the legs are ten lines long, smooth, and black, as are likewise the toes, of which the middlemost is rather longer than the rest; the claws are curved, black, and of equal length. In the female all the parts which are black in the male are brown, and all the white parts are of a dull ash-colour. It is an inhabitant of Georgia during the summer, frequenting the shrubby banks of rivers.

# SQUALUS TENTACULATUS.



## CHARACTER GENERICUS.

Os in anteriore et inferiore capitis parte, dentibus numerosis seriatis.

*Spiracula* utrinque ad latera colli, plerisque quinque.

*Corpus* oblongum, teretinsculum.

## CHARACTER SPECIFICUS, &c.

**SQUALUS** rostro utrinque tentaculato, spinoso, spinis longioribus brevioribusque intermediis.

**PRISTIS** cirratus. P. rostro cirrato, spinis longioribus; brevioribusque intermediis.

*Lath. Lin. Trans.* 2, p. 281.

---

In mariibus australibus reperitur rara haec Squali species, longa forsitan, cum plene adoleverit, multos pedes. Quae tamen ab Australasia litoribus in Angliam delata fuerint specimenia vix tres quatinus pedes exuperant. Color totius animalis leviter fuscus, abdomine pallidiore.





633



THE  
TENTACULATED SHARK.



*GENERIC CHARACTER.*

*Mouth* situated beneath the anterior part of the head, with numerous teeth disposed in rows.

*Spiracles* on each side the neck, in most species five in number.

*Body* oblong, somewhat cylindric.

*SPECIFIC CHARACTER.*

SHARK with serrated snout tentaculated on each side, with short teeth interposed between the longer ones.

---

The rare species of Shark here represented is a native of the Southern Seas, and probably measures several feet in length, when full grown; tho' the specimens which have been hitherto observed have scarcely exceeded three or four feet: the colour of the whole animal is pale brown, with the abdomen of a still paler cast.



## MUREX BRANDARIS.



### CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, exasperata suturis membranaceis.

*Apertura* desinens in canalem integrum rectum seu subadscendentem.

### CHARACTER SPECIFICUS, &c.

MUREX testa subovata spinis rectis ciuita, canda medioeri subulata recta spinisque oblique circumdata.

*Lin. Syst. Nat.* p. 1214.

MUREX Purpura.



*Rondel, test.* 64.

---

E mari Mediterraneo præcipue extrahitur Murex Brandaris, magnitudine, spinis, coloribusque multum varians.







THE  
SNIPE MUREX.



GENERIC CHARACTER.

*Animal* allied to a *Limax* or *Slug*.

*Shell* univalve, spiral, roughened by membranaceous sutures.

*Aperture* ending in a strait subascending channel.

SPECIFIC CHARACTER, &c.

Subovate cinereous MUREX, with strait spines,  
and subulate, obliquely-spined beak.

Snipe's-Bill MUREX.

---

This shell is principally found in the Mediterranean sea, and varies very much in size, spines, and colours.



# PAPILIO JAÏRUS.



## *CHARACTER GENERICUS.*

*Antennæ apicem versus crassiores, sæpius clavato-capitatae.*

*Alæ (sedentis) erectæ sursumque conniventes volatu diurno.)*

*Lin. Syst. Nat. p. 744.*

## *CHARACTER SPECIFICUS, &c.*

PAPILIO alis integerrimis fuscis; posterioribus disco albo, supra ocello, subtus duobus.

*Lin. Syst. Nat. Gmel. p. 2277.*

PAPILIO alis integerrimis fuscis, posticis disco albo, supra ocello, subtus duobus.

*Fab. sp. ins. 2. p. 63.*

DANAI FESTIVI.

# PAPILIO JAÏRUS.

*Cramer t. 16. A. B.*

---

Indiam incolit Papilio Jaïrus, magnitudine vera in tabula depictus.







## JAÏRUS.



### *GENERIC CHARACTER.*

*Antennæ* generally thickening towards the extremities, and terminating in a clavated tip.

*Wings* (when sitting) erect, and meeting upwards. Flight diurnal.

### *SPECIFIC CHARACTER, &c.*

Brown BUTTERFLY with entire wings, a white cloud on each, an eye-shaped spot on each lower wing and two beneath.

## JAÏRUS.

*Cramer. t. 6.*

*Clerk ic. t. 29. f. 3.*

---

The Papilio Jaïrus is a native of India, and is represented on the plate in its natural size.



## PSITTACUS LORY.



### CHARACTER GENERICUS.

*Rostrum* aduncum: mandibula superiore mobili:

*Nares* in rostri basi.

*Lingua* carnosa, obtusa, integra.

*Pedes* scansorii.

*Lin. Syst. Nat.* p. 139.

### CHARACTER SPECIFICUS, &c.

PSITTACUS brachyurus coccineus, subtus cæruleus; pileo nigro-violacco, alis viridibus.

PSITTACUS brachyurus purpureus, pileo violaceo, alis viridibus, pectore gcnibus caudaque cæruleis, orbitis incarnatis.

*Lin. Syst. Nat.* p. 145.

---

In insulis Philippensibus generata pulcherrima  
haec avis indole est docili admodum et mansueta.  
Minor paululum turture, longa plerumque est cir-  
citer decem uncias.

M







RPN - Delft Sculpt.

THE  
PHILIPPINE LORY.



*GENERIC CHARACTER.*

*Bill hooked : upper mandible moveable.  
Nostrils round, placed in the base of the bill.  
Tongue fleshy, broad, blunt at the end.  
Feet scensorial.*

*SPECIFIC CHARACTER, &c.*

SHORT-TAILED SCARLET PARROT, blue beneath, with violet-black crown, and green wings.

BLACK-CAPPED LORY.

*Edw. pl. 170.*

LE LORI D'AMBOINE.

*Pl. Ent. 518.*

---

This most beautiful species is a native of the Philippine islands, and is remarkable for the gentleness of its manners in a state of captivity. It is somewhat smaller than a turtle-dove, usually measuring about ten inches in length.

M 2



# DELPHINUS GANGETICUS.



## CHARACTER GENERICUS.

*Dentes* in maxilla utraque.  
*Fistula* in capite.

## CHARACTER SPECIFICUS, &c.

DELPHINUS cinereus, rostro elongato.  
DELPHINUS Gangeticus.

*Roxb. Ind. Res. vol. 7.*

---

Quem depiuximus Delphinum norunt Indi nomine *Soosoo*. Horum in Gangi flumine congregantur interdum plurimi. Longus est piscis, ut plurimum, pedes circiter duodecim. Color cinereus, subtus pallidior. Maxillæ elongatae admodum et graciles continent utræque sexaginta dentes. Delphina Gangeticum primus accurate descriptissime videtur Dominus Roxburgh in volumine septimo Disquisitionum Asiaticarum.





Horn—Solefin



634

THE  
**GANGETIC DOLPHIN.**

---

*GENERIC CHARACTER.*

*Teeth* in both jaws.  
*Spiracle* on the head.

*SPECIFIC CHARACTER, &c.*

**CINEREOUS DOLPHIN**, with lengthened snout.

**GANGETIC DOLPHIN.**

**Narrow-Snouted INDIAN DOLPHIN.**

---

The species of Dolphin represented on the present plate is occasionally observed in great multitudes in the river Ganges, and is known in India by the name of Soosoo. Its general length is about twelve feet, and its colour cinereous, paler beneath : the jaws are remarkably long and slender, and in each jaw are sixty teeth. This animal seems to have been first distinctly described by Dr. Roxburgh in the seventh volume of the Asiatic Researches.

## 2000-2001

Year	Number of new cases	Number of new deaths	Number of new survivors	Number of new cases per 100,000 population	Number of new deaths per 100,000 population	Number of new survivors per 100,000 population
2000	1,000	200	800	2.0	0.4	1.6
2001	1,000	200	800	2.0	0.4	1.6
Total	2,000	400	1,600	4.0	0.8	3.2
Average	1,000	200	800	2.0	0.4	1.6

## CHAMA COR.



### CHARACTER GENERICUS.

*Animal* Tethys.

*Testa bivalvis, grossior.*

*Cardo callo gibbo, oblique inserto fossulæ obliquæ.*

### CHARACTER SPECIFICUS, &c.

CHAMA testa subrotunda lævi, natibus recurvatis, rima hiante.

*Lin. Syst. Nat. p. 1137.*

COR BOVIS.

*Argenv. t. 26. f. κ.*

CUCULLA fatui dichoncha.


*Seb. 3. p. 177. t. 86. f. 1.*

---

Maribus Europæis innascitur concha hic depicta,  
interdum major iconæ quæ in tabula ostenditur.







RPMC

THE  
**HEART CHAMA.**



**GENERIC CHARACTER.**

*Animal* allied to a *Tethys*.

*Shell* bivalve, strong.

*Hinge* with a gibbose callus obliquely inserted into an oblique fossule.

**SPECIFIC CHARACTER, &c.**

**CHAMA** with smooth roundish shell, with re-curved tips, and gaping orifice.

**The Smooth OX-HEART CHAMA.**

---

The present shell is a native of the European seas, and is sometimes found of a larger size than represented on the annexed plate.



# PAPILIO ANDROMACHUS.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque connientes ;  
(volatu-diurno.)

*Lin. Syst. Nat.* p. 744.

## CHARACTER SPECIFICUS, &c.

PAPILIO alis fuscis basi fulvis, primoribus sub-falcatis fascia flava, subtus albo-maculatis.

PAPILIO Andromachus.

*Cram. pap. t. 56. f. A. B.*

*Eq. Archiv.*

---

Americæ Australis partes calidiores incolit Papilio Andromachus, magnitudine vera in tabula depictus.

## ANDROMACHUS.



### GENERIC CHARACTER.

*Antennæ* generally thickening towards the extremities, and terminating in a clavated tip.

*Wings* (when sitting) erect, and meeting upwards.

Flight diurnal.

### SPECIFIC CHARACTER

BUTTERFLY with brown wings, fulvous at the base, with the upper wings subfalcated, marked with a yellow band, and spotted beneath with white.

---

This rare insect is an inhabitant of the warmer parts of South America, and is represented on the plate in its natural size.





## INDEX.

### PL.

- 505 *Anas Ægyptiaca.*  
 598 *Anarhichas pantherinus.*  
 614 *Aranea venatoria.*  
 602 —— *Tarantula.*  
 621 *Ardea tigrina.*  
 620 *Asterias oligactes.*  
 596 *Buccinum Rumphii.*  
 591 *Cancer Sebanus.*  
 627 —— *linearis.*  
 635 *Chama Cor.*  
 619 { *Cypræa Argus.*  
     —— *leucopis.*  
 634 *Delphinus Gangeticus.*  
 597 *Emberiza Ciris*  
 624 *Erpeton tentaculatus.*  
 594 *Lacerta aurita.*  
 623 *Madrepore Pileus.*  
 607 —— *pezizoides.*  
 599 *Murex Femorale.*  
 631 —— *Brandaris.*  
 618 *Metops cœruleocephalus.*  
 629 *Muscicapa melanoleuca.*  
 601 *Mycteria Australis.*  
 628 *Oniscus Entomon.*  
 610 *Ovis Canadensis.*  
 590 *Papilio Protenor.*  
 604 —— *Merope.*  
 600 597 —— *Amphytrion.*  
 632 —— *Jairus.*  
 635 —— *Andromachus.*  
 606 *Patella fusca.*  
 625 *Paradisea magnifica.*  
 608 *Phalangium dubium.*  
 618 *Phalæna Strix.*

## INDEX.

### PL.

- 598 *Anarhichas panther.*  
 613 *Bee-Eater blue-headed.*  
 621 *Bittern Tiger.*  
 596 *Buccinum marbled.*  
 635 *Butterfly Andromachus.*  
 632 —— *Jairus.*  
 604 —— *Merope.*  
 590 —— *Protenor.*  
 597 —— *Amphytrion.* 600  
 636 *Chama Heart.*  
 619 { *Cowry White-Eyed,*  
     —— *Argus.*  
 591 *Crab Seban.*  
 627 —— *linear.*  
 634 *Dolphin Gangetic.*  
 597 *Emberiza painted.*  
 624 *Erpeton tentaculatus.*  
 629 *Flycatcher Georgian.*  
 605 *Goose Egyptian.*  
 601 *Jabiru Australasian.*  
 594 *Lizard lobe-cheeked.*  
 607 *Madrepore pezizoid.*  
 623 —— *oblong.*  
 609 *Macaw hyacinthine.*  
 593 *Manakin rock.*  
 611 612 *Moth Persimon.*  
 615 616 —— *Cecropian.*  
 618 —— *Strix.*  
 626 *Monkey Ascanius.*  
 599 *Murex femoral.*  
 631 —— *Snipe.*  
 628 *Oniscus sharp-tailed.*  
 606 *Patella brown.*  
 625 *Paradisea magnificent.*

- |     |                       |             |     |                           |
|-----|-----------------------|-------------|-----|---------------------------|
| 611 | 612                   | — regia.    |     |                           |
| 615 | 616                   | — Cecropia. | 633 | Parrakeet Lory.           |
| 592 | Phytolithus Filicis.  |             | 589 | Partridge sand.           |
| 593 | Pipra rupicola.       |             | 608 | Phalangium doubtful.      |
| 609 | Psittacus Angustus.   |             | 592 | Phytolithus Fern.         |
| 633 | — Lory.               |             | 630 | Shark tentaculated.       |
| 626 | Simia Ascanius.       |             | 610 | Sheep Canadian.           |
| 595 | Sphinx Cluentias.     |             | 595 | Sphinx Cluentius.         |
| 622 | Solpuga venenosa.     |             | 622 | Solpuga poisonous.        |
| 630 | Squalus tentaculatus. |             | 620 | Star-Fish slender-limbed. |
| 589 | Tetrao arenaria.      |             | 602 | Tarantula.                |
| 617 | Turdus cyanurus.      |             | 614 | — American.               |
| 608 | Vespa campanaria.     |             | 617 | Thrush blue-tailed.       |
|     |                       |             | 603 | Wasp campanular.          |











