


MUSÉUM NATIONAL D'HISTOIRE NATURELLE
INSTITUT D'ÉCOLOGIE ET DE GESTION DE LA BIODIVERSITÉ
SERVICE DU PATRIMOINE NATUREL



European Register of Marine Species
A check-list of the marine species in Europe
and a bibliography of guides to their identification

Mark J. COSTELLO, Chris EMBLOW and Richard WHITE (editors)

European Register of Marine Species

A check-list of the marine species in Europe and a bibliography of guides to their identification

Mark J. COSTELLO⁽¹⁾, Chris EMBLOW⁽²⁾ and Richard WHITE⁽³⁾ (editors)

(1) Ecological Consultancy Services Ltd (EcoServe) - 17 Rathfarnham Road - Terenure - Dublin 6W - Ireland
E-mail: mcostello@ecoserve.ie

(2) Ecological Consultancy Services Ltd (EcoServe) - 17 Rathfarnham Road - Terenure - Dublin 6W - Ireland

(3) School of Biological Sciences - University of Southampton - Bassett Crescent East- Southampton SO16 7PX - UK

PUBLICATIONS SCIENTIFIQUES
DU MUSÉUM NATIONAL D'HISTOIRE NATURELLE

INSTITUT D'ÉCOLOGIE ET DE GESTION DE LA BIODIVERSITÉ (I.E.G.B.)
SERVICE DU PATRIMOINE NATUREL (S.P.N.)

associé par convention au
MINISTÈRE DE L'AMÉNAGEMENT DU TERRITOIRE ET DE L'ENVIRONNEMENT,
DIRECTION DE LA NATURE ET DES PAYSAGES ET DIRECTION DE L'EAU

Cette publication constitue le volume **50** de la collection Patrimoines Naturels

Rédacteur en chef : Philippe Keith
Assistante de rédaction : Gwénaëlle Chavassieu

Photo 1^e de couverture : *Simnia patula*, Mollusca, Gastropoda (photo : Claude Huyghens)

Photos 4^e de couverture : a b
c d

- a - *Himanthalia elongata* (Phaeophyceae) buttons on exposed lower eulittoral rock
(photo : Joint Nature Conservation Committee)
- b - *Prostheceraeus* sp., Plathyhelminthes, Turbellaria (photo : Claude Huyghens)
- c - *Botryllus schlosseri*, Tunicata (photo : Claude Huyghens)
- d - *Eunicella verrucosa* with *Alcyonium digitatum* (Cnidaria, Anthozoa),
erect sponges and faunal turf on moderately exposed rock (photo : Joint Nature Conservation Committee)

ISSN - 1281-6213

ISBN - 2-85653-538-0

© Publications Scientifiques du M.N.H.N. - Paris, 2001, for the present edition
Society for the Management of European Biodiversity Data Limited, Ireland, for all other uses

Recommended citation :

Costello M.J., Emblow C.S. and White R. (editors), 2001. European Register of Marine Species. A check-list of the marine species in Europe and a bibliography of guides to their identification. *Patrimoines naturels*, **50** : 463 p.

Sections may be cited as :

Clipson N, Landy E. and Otte M., 2001. Fungi. In : Costello M.J., Emblow C.S. and White R. (editors), European Register of Marine Species. A check-list of the marine species in Europe and a bibliography of guides to their identification. *Patrimoines naturels*, **50** : 463 p.

This document is a product of the project "A register of marine species in Europe to facilitate marine biodiversity research and management", a concerted action project funded under the European Union MAST research programme (MAS3-CT97-0146). Its short titles is "European Register of Marine Species (ERMS)"

All rights for the use and reuse of contents of this volume rest with the Society for the Management of European Biodiversity Data Limited. Any requests for electronic versions of the contents, or permission to use this data are welcomed, and must be addressed to the Secretary of the society : Dr M.J. Costello, c/o EcoServe, 17.Rathfarnham Road, Terenure, Dublin 6W, Ireland. E-mail : mcostello@ecoserve.ie
Project web site : <http://erms.biol.soton.ac.uk>

CONTENTS

CONTENTS	3
EXTENDED SUMMARY	5
Cadre géographique	5
Contenu taxonomique	5
Présentation des espèces dans le catalogue	5
INTRODUCTION	7
Geographic scope	7
Species included	7
Listing of species in the check-list	8
Layout of each section	9
Future development	9
Acknowledgements	9
SPECIES LISTS	15
FUNGI	15
PLANTS	20
PROTISTA	40
Alveolates	40
Apicomplexa (free-living species)	40
Ciliata (Phylum Ciliophora)	40
Dinoflagellata	47
Naked amoebae	53
Testate amoebae	54
Apusomonads	57
Choanoflagellata	57
Cryptophytes	59
Euglenozoa	59
Euglenoids (heterotrophic species)	59
Kinetoplastids	59
Foraminifera	60
Kathablepharids	75
Haptophytes	76
Prasinophytes	76
Stramenopiles	77
Bicosoecids	77
Labyrinthulids	77
Thaustrochyrids	77
Stramenopiles <i>incertae sedis</i>	78
Thaumatomonads	78
Xenophyophora	78
Protista <i>incertae sedis</i>	79
MYXOZOA	80
PORIFERA	85
PLACOZOA	103
MESOZOA	104
CNIDARIA	104
Anthozoa	104
Octocorallia	104
Hexacorallia	106
Scyphozoa	111
Hydrozoa	112
Hydroida	112
Siphonophora	120
CTENOPHORA	122
PLATYHELMINTHES	123
Turbellaria	123
Aspidogastrea	136
Digenea	136
Monogenea	142
Cestoda	146
ROTIFERA	149
GNATHOSTOMULIDA	151
NEMERTINI	152
GASTROTRICHA	156
CEPHALORHYNCHA	159
Kinorhyncha	159
Loricifera	160
Nematomorpha and Priapulida	160
NEMATODA	161
Nematoda (free-living)	161
Nematoda (parasitic)	174
ACANTHOCEPHALA	176
CYCLIOPHORA	177
ECHIURA	178
SIPUNCULA	178
ENTOPROCTA (Kamptozoa)	179
MOLLUSCA	180
ANNELIDA	214
Polychaeta	214
Oligochaeta	231
Hirudinea	234
Pogonophora	235
TARDIGRADA	236
CHELICERATA	237
Arachnida	237
Acarina	237
Pycnogonida	241
CRUSTACEA	243
Remipedia	243
Branchiura	243
Cladocera (Branchiopoda)	243
Ostracoda	244
Pentastomida	251
Mystacocarida	252
Copepoda	252
Tantulocarida	280
Cirripedia	280
Free-living	280
Ascothoracida	283
Rhizocephala	283
Stomatopoda	284
Euphausiacea	284
Decapoda	284
Thermosbaenacea	293
Mysidacea	293

Amphipoda	295
Cumacea	308
Tanaidacea	310
Isopoda	315
CHILOPODA	323
DIPLOPODA	323
INSECTA	323
PHORONIDA	324
BRYOZOA (Ectoprocta)	325
BRACHIOPODA	334
CHAETOGNATHA	335
HEMICORDATA	336
ECHINODERMATA	336
TUNICATA	352
Thaliacea	355
Appendicularia	356
CEPHALOCHORDATA	357
PISCES	357
TETRAPODA.....	375
BIBLIOGRAPHY OF IDENTIFICATION GUIDES	
TO MARINE SPECIES IN EUROPE	377
GENERAL	377
FUNGI	377
PLANTS	377
General and other algal taxa.....	377
Seaweeds	378
Bangiophyceae.....	378
Chlorophyta	378
Phaeophyta	378
Rhodophyta	378
PROTISTA	379
Foraminifera	379
FAUNA	379
GENERAL	379
PORIFERA	380
CNIDARIA	380
General	380
Anthozoa	381
Scyphozoa	381
Hydrozoa.....	381
CTENOPHORA	381
PLATYHELMINTHES	382
ROTIFERA	382
NEMERTINI	382
GASTROTRICHA	383
CEPHALORHYNCHA.....	383
Kinorhyncha	383
Priapulida (see under Sipunculida)	383
Nematomorpha	383
NEMATODA	383
ACANTHOCEPHALA	383
ECHIURA (see under sipunculida)	383
SIPUNCULIDA, ECHIURA & PRIAPULIDA	383
ENTOPOCTA	383
MOLLUSCA	383
General	383
Polyplacophora, Scaphopoda, Aplacophora ..	384
Gastropoda	384
Cephalopoda	385
Bivalvia	386
ANNELIDA	386
General	386
Polychaeta	386
Oligochaeta	387
Hirudinea	387
Pogonophora	387
TARDIGRADA	387
CHELICERATA.....	387
General	387
Pycnogonida	387
CRUSTACEA	388
General and other miscellaneous groups	388
Cladocera (Branchiopoda)	388
Ostracoda	388
Copepoda	388
Cirripedia	389
Stomatopoda	389
Decapoda.....	389
Mysidacea	390
Amphipoda.....	390
Cumacea	391
Tanaidacea	391
Isopoda	391
INSECTA	391
PHORONIDA	391
BRYOZOA	391
BRACHIOPODA	392
CHAETOGNATHA.....	392
HEMICORDATA	392
ECHINODERMATA	392
TUNICATA	393
General and other than Ascidiacea	393
Asciidiacea	393
CEPHALOCHORDATA	393
VERTEBRATA	393
General	393
Pisces	393
Tetrapoda.....	395
Reptilia	395
Aves	395
Mammalia	396
APPENDIX 1-Contact details of the persons and organisations contributing to ERMS	397
INDEX TO ORDERS, FAMILIES & GENERA	403

EXTENDED SUMMARY

European Register of Marine Species / *Répertoire des Espèces Marines d'Europe.*

Les noms scientifiques forment le véhicule de communication de tous les résultats de recherches et textes réglementaires sur la biodiversité, mais usages et habitudes d'emploi des noms varient encore d'un pays à l'autre. Pour la première fois, le présent volume présente un répertoire complet et coordonné de toutes les espèces –plantes, algues, invertébrés, vertébrés– à l'échelle des mers d'Europe. Il est le fruit du travail collectif de près de 170 scientifiques dans le cadre du projet *European Register of Marine Species* (ERMS), soutenu par la Commission Européenne dans le cadre de son programme Marine Science and Technology (MAST).

Cadre géographique

Le cœur géographique du répertoire est centré sur le plateau continental européen, les Canaries et les Açores, jusqu'à l'Islande et le nord de la Norvège, y compris bien entendu la Méditerranée et la mer Baltique. Dans la présente liste, le terme "espèces marines" s'applique également aux espèces intertidales (littorales) et à celles des eaux saumâtres. Dans la mesure du possible, les données concernant les eaux du large et les grandes profondeurs ont été prises en compte, de manière à couvrir la totalité des mers d'Europe (Figure 1). De nombreuses lacunes subsistent sans aucun doute aux frontières géographiques (Océan Arctique, Mer Noire, sud des Açores et des Canaries) et bathymétriques (faunes bathyales et abyssales), et l'objectif d'une prochaine édition du Répertoire devrait être précisément de combler ces lacunes. Une grande partie de la mer Baltique est occupée par de l'eau douce, et il a été laissé à l'appréciation des différents coordinateurs d'inclure, ou non, les espèces de ce secteur dans le Répertoire.

Contenu taxonomique

Le Répertoire catalogue environ 30 000 espèces dans l'ensemble des groupes taxonomiques (Tableau 1). Seuls les noms publiés et nominalement disponibles ont été pris en considération. A quelques très rares exceptions près, ne sont incluses que les espèces dont l'occurrence à l'intérieur du cadre géographique du projet avait déjà été reconnue dans la littérature. Certains petits groupes taxonomiques sont incomplètement couverts : ainsi, chez les Acariens, seule la famille Halacaridae a été traitée, et le catalogue des Rotifères et des Brachiopodes ne couvre que l'Atlantique nord-est. Les lichens, les diatomées et les cyanobactéries ne faisaient pas partie des objectifs du projet, de même que les bactéries (Eubacteria and Archaea) et les virus. L'opportunité de traiter plusieurs groupes de protistes et les champignons a été saisie, bien que non inclus à l'origine dans les objectifs. Les angiospermes des sols salés sont déjà couverts dans les catalogues de végétaux terrestres, et il n'a pas été jugé utile de les prendre en compte dans ce répertoire.

Présentation des espèces dans le catalogue

Pour faciliter la consultation, les groupes taxonomiques sont présentés selon une séquence qui va des champignons et des plantes aux organismes unicellulaires (protistes), puis aux animaux pluricellulaires, pour terminer avec les oiseaux et les mammifères. Chaque partie du Répertoire a été compilée par un coordinateur (dont les noms figurent à l'Annexe 1), dont c'est la responsabilité de choisir la classification suivie ici, jusqu'au niveau des familles. A l'intérieur de chaque famille, les genres sont présentés alphabétiquement, et à l'intérieur de chaque genre, les espèces le sont également.

Dans les listes d'espèces, les synonymes, ou tout autre nom employé pour désigner une même espèce, sont précédés du signe 'égal' (=). Les codes A et M après un nom d'espèce indiquent que l'espèce est présente dans, respectivement, l'Atlantique (y compris éventuellement l'Arctique, la Baltique, ou les grandes profondeurs) et la Méditerranée (y compris éventuellement

la Mer Noire). De telles indications synonymiques ou géographiques ne sont pas présentes uniformément pour l'ensemble des groupes traités.

Un index bibliographique de 700 titres oriente vers les principaux guides et outils de détermination de la flore et de la faune marines d'Europe.

Le catalogue *European Register of Marine Species* est destiné à devenir un outil de standardisation et de référence pour tous ceux qui sont impliqués dans la recherche, la formation et la gestion de la biodiversité marine en Europe. La présente version papier est publiée parallèlement à la liste électronique consultable sur le site Web du projet à l'adresse <http://erms.biol.soton.ac.uk>.

INTRODUCTION

The foundation of biodiversity research and management is correctly identifying and naming species. Despite the existence of a system for naming species, different names are used for the same species, and the same names for different species, in different parts of Europe. This leads to considerable confusion, and may cause regulatory problems where a species is listed as a priority for protection under an incorrect name. There will always be debate amongst taxonomic specialists about which name is more correct for some species, but a standard working list of names is essential for non-specialists to use. This volume is the first attempt to list all marine species in Europe. It has been produced as a product of the European Register of Marine Species (ERMS), a project part-funded by the European Commission (EC) Marine Science and Technology (MAST) research programme. All the information in this volume, and additional information, is available from the project web site <http://erms.biol.soton.ac.uk>.

Lists of species, such as provided here, are only the starting point for marine biodiversity management and research. The value of producing the lists is

- to provide a single nomenclature for European marine species, which will generate further research to clarify anomalies, and
- to form the basis for more elaborate species databases, with more synonyms, and data on species distribution, ecology, conservation importance, economic importance, and so on.

An added benefit has been that the co-operation amongst scientists in producing the register has increased communication and interest in the management and use of taxonomic data throughout Europe. It is anticipated that the Register will become a standard reference and technological tool for marine biodiversity training, research and management in Europe. The species register, with its accompanying bibliography of marine species identification guides, can be used to:

- Check the spelling or find the correct name of a species and the authority;
- For a given species, check what other (or how many) species exist in the same Genus, Family or higher taxa. These species may not be included in the local identification guides;
- Find information on the distribution of species among higher taxa;
- Indicate the level of knowledge of a group of species by analysing the rate of discovery of species;
- Find literature on the identification and other knowledge of marine species in Europe.

Geographic scope

The geographic scope of the register was initially the continental shelf seas of Europe, the Canaries and Azores to Iceland and northern Norway and includes the Mediterranean shelf and Baltic Seas. However, although not part of the EC contract, records from deeper waters were included where possible such that the ERMS area covers all European seas (Figure 1). Future work should aim to complete all lists for the deep sea. The EC contract restricted the project to western European organisations. Thus, the deep sea, Arctic Ocean, Black Sea, and southern limits of the Atlantic within the ERMS area (i.e. south of Azores and Canary Islands) may not be adequately covered. The northern parts of the Baltic Sea are freshwater and it was left to the discretion of list compilers whether to include species there within this check-list or not. Future work should involve the expertise of scientists wherever they are located to fill these gaps.

Species included

The check-list includes about 30,000 species representing all marine taxa in Europe (Table 1). Only published and taxonomically available species names are included. With very few exceptions, only species whose occurrence in the ERMS area (Figure 1) has been previously published are included. Synonyms and other names for a species have been included in some instances. The marine environment was broadly defined to include intertidal (littoral) and brackish water habitats. Parts of a few small groups have not been fully covered, namely other

families of Acarina apart from Halacaridae, and the Rotifera (Rotatoria) and Brachiopoda lists are limited to north-east Atlantic species. Although not part of the EC contract species lists of several protist groups and fungi have also been compiled. Lichens, diatoms and cyanobacteria were beyond the scope of the contract. Saltmarsh angiosperm plants were also excluded as these are generally included in terrestrial plant classifications. It was not possible to use the species concept in a similar manner for bacteria (Eubacteria and Archaea) and viruses, so these groups were also excluded from the project. Future work should aim to compile lists for all taxa where the species concept is applicable.

Listing of species in the check-list

Different phyla may have a few to several thousand species, and different experts have compiled different parts of some groups (e.g. parasitic and non-parasitic nematodes). To facilitate readers, the taxonomic groups have been listed starting from the fungi and plants, to single-celled organisms (protists), to lower animals, and to higher animals (birds, mammals). This project does not deal with systematics or classification of marine taxa. The suprageneric taxa named in the contents of this volume were selected to link with particular lists. The selection of groups so listed vary in their taxonomic rank, most phyla are listed, some classes, and some orders and a few families. Within a particular species list, the classification of families and higher taxa has been decided by the compilers of that list. Within a family, genera are listed alphabetically, and within a genus species are listed alphabetically. Where only some sub-groups are listed in the Contents, this is because these groups were compiled by different persons than the remainder of the list. They thus have a unique Preface and list of source references.




Figure 1. Map showing the geographic scope of the ERMS project. A - Atlantic Ocean and M - Mediterranean Sea.

Layout of each section

Species lists were provided to the project electronically, as spreadsheets and text files. These were converted into web pages and 'rich text format' (rtf) files in the format of this volume. Some lists had additional information on species distribution and other comments that were not included in this volume to maintain as much uniformity between lists as possible. It is anticipated that future editions of this volume will include such additional information.

A strict format for species lists, Preface text, and references, was not insisted upon so as not to delay or inhibit the timely arrival of contributions. Thus the content, style of writing, and format of references, varies between species groups.

In the species lists, synonyms, or other names erroneously used for certain species (i.e. chresonyms) follow an equals (=) sign. The letters A and M after a species name indicate it occurs in the Atlantic (may include Arctic, Baltic and/or deep sea) and Mediterranean (may include Black Sea) respectively. The name of the person who described a species, called the 'taxonomic authority', is provided for almost all species names. The manner of citation of the authority follows normal practice in general zoology (only first authority given), some worm and parasite taxa (person who changed genus name also indicated) and botany; see the Preface to Plants for a fuller description of the citation of authorities.

Some species are not provided with synonyms or chresonyms, and/or distribution or other data. This was generally because the time available did not allow this information to be assembled. However, most list compilers have additional information on the species and would be interested to know what information would be of greatest interest to readers. If you cannot find a species name in this volume there are four possibilities:

1. the name you have is not available (check the spelling)
2. it has been treated as a synonym by the list compiler but not listed as such
3. it is a senior synonym and the list has used the wrong name
4. it has been omitted from the list due to lack of knowledge, uncertainty, or error.

The 'Prefaces' to each species list cite key publications where you will find more detailed information on the species. In addition a bibliography of identification guides to the marine species of Europe is provided at the end of this volume. Having consulted these texts, if you feel the list may be in error or should indicate the synonym for other readers, please contact the list compiler and/or the Secretary of the Society for the management of European biodiversity data. They will note your comments so that future editions may address it. The contact details of compilers are provided in Appendix 1.

Future development

This Register will continue to be updated as new knowledge becomes available. The present lists will vary in quality because of the availability of recent reviews of taxa, existing databases, and time available for people to produce and check them. To facilitate the updating of the Register its copyright has been transferred to a new organisation, namely the 'Society for management of European biodiversity data'. Contributors to ERMS are members of the society and elect a governing council that authorises updates, elaborations, and publications arising from the species lists. Comments on this volume, and offers to revise and expand certain groups, should be addressed to the Secretary of the Society for the management of European biodiversity data: Dr M. J. Costello, EcoServe, 17 Rathfarnham Road, Terenure, Dublin 6W, Ireland; E-mail: mcostello@ecoserve.ie

Acknowledgements

The production of this volume would not have been possible without the good will and co-operation of over 170 scientists. Many of these are named within the volume as compilers of lists or who assisted compilation of lists (Table 1). Other people whose role is less obvious must also be thanked. Their contact details are provided in Appendix 1. Frank Bisby (University of Reading, UK) and Wouter Los (University of Amsterdam, The Netherlands) provided invaluable advice in how to design, execute and continue this project. Dusan Zavodnik (Centre for Marine

Research, Rovinj, Croatia) provided details of species recorded in the Adriatic Sea, including endemic, and other helpful comments, and Mateo Garrido (Universidad of Las Palmas de Gran Canaria) provided lists of species recorded in the Canary Islands. Guenther Radach (Institut für Meereskunde, Kiel) and Herman Hummel (Netherlands Institute of Ecology, Yerseke) participated in discussions and provided advice in relation to data management. David Connor, Frank Bisby, Carlo Heip, Salvatore Aricò, Pamela Harding, Colleen Skule Adam, Pierre Lasserre, Ioannis Karakassis, Jan Marcin Weslawski and others helped the project interact with external (potential end-user) organisations, including the European Environment Agency, International Council for the Exploration of the Sea (ICES), Oslo and Paris Convention for the protection of the marine environment of the north-east Atlantic (OSPAR), Species 2000, Jakarta Marine Mandate of the Convention on Biological Diversity, Diversitas, CIESM and other organisations. John W. Patching (National University of Ireland, Galway) explained how the species concept as used here cannot be similarly applied to bacteria. Ulf Scheller (Järvpås, Sweden) confirmed that the Pauropoda are a strictly terrestrial group, although one species *Amphipauropus rhenanus* (Hüther, 1971) lives in coastal sand dunes. Many other people helped the project directly and through assisting others more directly involved. The compilers named in this volume also contributed to the project in its design, communication with other experts, and implementation. This project has demonstrated a co-operative and collegial spirit amongst the scientific community in Europe.

Table 1. The species lists included in this volume, the persons who compiled them and assisted in their compilation, the number of species per group, and an indicator of how complete a list is of the described species. The contact details of the list compilers are in Appendix 1. The status of a list is scored as follows: + = preliminary list, known to be or likely to be incomplete; ++ = compiled from recent authoritative literature; +++ = compiled by expert in the group; ++++ = checked by additional expert in the group; +++++ = checked by several experts in the group; C = confident of reasonable coverage of all European seas, including Arctic, deep sea and Black Sea.

Species group	Compiler(s)	Assisted by	Number species	Status
Fungi	N. Clipson, E. Landy, M. Otte	<i>G. Bremer, G. Jones</i>	318	++++
Macroalgae of Rhodophycota, Phaeophycota, Chlorophycota, and two genera of Xanthophycota	M. D. Guiry	<i>G. Furnari, F. Rindi, E. Nic Dhonchha, S. Lawson</i>	1702	+++++ C
Seagrass	M. D. Guiry		5	+++++ C
Apicomplexa (free-living species)	S. Brandt		3	++
Ciliates - Chonotricha	A. W. Jankowski		37	+++
Ciliates - folliculinids	M. Mulisch		30	+++
Ciliates - aloricate oligotrichs	S. Agatha		82	+++
Ciliates - Rhynchodida	A. W. Jankowski		42	+++
Dinoflagellates	S. Brandt	<i>M. Elbrächter</i>	718	++
Amoebae - naked	A. Rogerson, A. Goodkov		74	++++
Amoebae - testate	R. Meisterfeld		97	+++
Apusomonads	S. Brandt		3	+++
Choanoflagellates	S. Brandt		98	+++
Cryptophytes	S. Brandt		14	+
Euglenoids - heterotrophic euglenoids	S. Brandt		26	+
Euglenoids - kinetoplastids	S. Brandt		13	+++
Foraminifera	O. Gross		1167	+++++ C
Kathablepharids	S. Brandt		2	++
Haptophytes	S. Brandt		36	+
Prasinophytes	S. Brandt		24	+
Bicosoecids	S. Brandt		17	+++
Labyrinthulids	M. Dick, S. Brandt		10	+++
Thaustrochytrids	M. Dick, S. Brandt		15	+++
Stramenopiles <i>incertae sedis</i>	S. Brandt		4	+++
Thaumatomonads	S. Brandt		17	+++
Xenophyophora	O. Tendal, J. van der Land		20	++++
Protista <i>incertae sedis</i> (heterotrophic species)	S. Brandt		40	+++
Myxozoa	E. Karlsbakk		230	+++++ C
Porifera	R. W. M. van Soest	<i>N. Boury-Esnault</i>	1640	+++
Placozoa	J. van der Land		2	++
Mesozoa	J. van der Land, J. Hallan		36	+++
Octocorallia (excl. Pennatulacea)	L. van Ofwegen, M. Grasshoff, J. van der Land		92	+++++ C
Octocorallia - Pennatulacea	G. C. Williams, J. van der Land		37	+++++ C
Actiniaria	J. van der Land, J. H. den Hartog	<i>J. Ryland, K. Riemann-Zürneck</i>	243	+++++ C
Antipatharia	J. van der Land, D. M. Opresco		28	+++++ C
Scleractinia	S. D. Cairns, B. W. Hoeksema, J. van der Land	<i>H. Zibrowius</i>	86	+++++ C
Scyphozoa	P. Cornelius, G. Jarms, Y. M. Hirano, J. van der Land,		53	+++++ C
Cubozoa	P. Cornelius		1	+++++ C
Hydrozoa	J. van der Land, W. Vervoort, S. D. Cairns, P. Schuchert		684	+++++ C
Siphonophora	G. M. Mapstone	<i>P. R. Pugh</i>	105	++++
Ctenophora	J. van der Land		38	++
Turbellaria	A. Faubel, C. Noreña		1137	+++++ C
Aspidogastrea	D. Gibson		4	+++++ C
Digenea	D. Gibson	<i>M. Køie, P. Bartoli</i>	592	+++++ C
Monogenea	R. Bray	<i>L. Euzet, G. Kearn</i>	353	+++++ C
Cestoda	R. Bray	<i>B. B. Georgiev, L. Euzet</i>	312	+++++ C
Rotifera	M. O'Reilly		139	++
Gnathostomulida	J. van der Land		25	+++

Species group	Compiler(s)	Assisted by	Number species	Status
Nemertini (Nemertea)	R. Gibson		478	+++++ C
Gastrotrichia	J. L. d'Hondt, J. van der Land		240	++++ C
Cephalorhyncha (= Loricifera, Priapulida, Kinorhyncha, Nematomorpha)	B. Neuhaus, J. van der Land		52	++++ C
Nematoda - free living	G. De Smet, M. Vincx, A. Vanreusel, S. Vanhove, J. Vanaverbeke, M. Steyaert	F. Riemann	1625	++++ C
Nematoda - parasitic	D. Gibson	F. Moravec, H.-P. Fagerholm	212	++++ C
Acanthocephala	D. Gibson	C. R. Kennedy, Z. M. Dimitrova	67	+++++ C
Cyclophora	C. S. Emblow		1	+++++ C
Echiura	J. van der Land	J. I. Saiz-Salinas	19	+++ C
Sipuncula	J. van der Land	J. I. Saiz-Salinas	44	+++ C
Entoprocta	P. J. Hayward		45	+++++ C
Mollusca	S. Gofas, J. Le Renard, P. Bouchet,	R. Giannuzzi-Savelli, A. Guerra, D. Heppell, T. Hoisaeter, E. Platts, S. Smith, J.-A. Sneli, A. Warén	3353	+++++ C
Polychaeta	G. Bellan	K. Arvanitidis, J. C. Dauvin, F. Gentil, G. Bachelet, H. Hansson, R. Barnich, D. Fiege, M. E. Petersen, T. Brattegard, T. Holthe	1848	++++ C
Oligochaeta	C. Erséus, B. Healy		190	+++++ C
Hirudinea	J. van der Land		36	++
Pogonophora	J. van der Land, E. Southward	T. Brattegard	23	+++++ C
Tardigrada	J. van der Land		76	++++ C
Chelicerata	A. Legakis		1	+
Acarina - Halacaridae	I. Bartsch		214	+++++ C
Pycnogonida	J. van der Land, F. Krapp	J. Stock, Child, R. Bamber	146	++++ C
Remipedia	G. Boxshall		1	++++ C
Branchiura	G. Boxshall		2	++++ C
Cladocera - Branchiopoda	G. Boxshall		9	++++ C
Ostracoda	D. Horne, A. Bruce, J. Whittaker		769	+++++ C
Pentastomida	J. van der Land		2	+++ C
Mystacocarida	G. Boxshall		2	++++ C
Copepoda - Calanoida	G. Boxshall		649	++++ C
Copepoda - Cyclopoida	G. Boxshall		177	++++ C
Copepoda - Harpacticoida	R. Huys		1357	++++ C
Copepoda - Misophrioida	G. Boxshall		16	++++ C
Copepoda - Monstrilloida	G. Boxshall		33	++++ C
Copepoda - Mormonilloida	G. Boxshall		2	++++ C
Copepoda - Platycopioida	G. Boxshall		3	++++ C
Copepoda - Poecilostomatoida	G. Boxshall	M. O'Reilly, D. Zavadnik	353	++++ C
Copepoda - Siphonostomatoida	G. Boxshall		354	++++ C
Tantulocarida	G. Boxshall		13	++++ C
Cirripedia - non-parasitic	A. Southward		107	++++ C
Thoracica				
Cirripedia - parasitic	G. Boxshall		10	++++ C
Ascothoracida				
Cirripedia - parasitic Rhizocephala	G. Boxshall		28	++++ C
Stomatopoda	J. van der Land	P. Noël	22	+++ C
Euphausiacea	J. van der Land		41	+++
Decapoda	M. Türkay		672	++++ C
Thermosbaenacea	J. van der Land		7	++
Mysidacea	J. van der Land, T. Brattegard		198	++++ C
Amphipoda	D. Bellan-Santini, M. J. Costello	S. Ruffo, J. C. Dauvin, L. Collier	1183	+++++ C
Cumacea	L. Watling	T. Brattegard	188	+++++ C
Tanaidacea	G. Bird	M. Gutu	280	+++++ C
Isopoda - excluding Epicaridea	J. van der Land		605	++
Isopoda, Epicaridea, Bopyridae	J. C. Markham		54	++++ C
Chilopoda	A. Minelli		6	++++
Diplopoda	A. Minelli		2	++++
Insecta - Chironomidae	D. Murray		15	++++ C

Species group	Compiler(s)	Assisted by	Number species	Status
Insecta	A. Legakis		19	++++
Phoronida	C. Emig, C. Roldán, J. M. Viéitez		9	+++++
Bryozoa	P. J. Hayward	J. Harmelin	724	+++++ C
Brachiopoda	C. Howson		18	++
Chaetognatha	J. van der Land, H. Kapp		42	++++ C
Hemichordata	J. van der Land		17	+++
Echinodermata	H. G. Hansson	S. Stöhr, C. Massin, A. Gebruk, A. Mironov, A. Smirnov, D. Zavodnik, M. Garrido	648	+++++ C
Asciacea & Sorberacea	C. Monniot, D. Connor, P. Lozouet, J. Marmayou		393	+++++ C
Thaliacea	J. van der Land, R. van Soest		35	++++ C
Appendicularia	J. van der Land		53	++
Cephalochordata	J. van der Land		2	++
Pisces - Agnatha	J. van der Land, M. J. Costello, N. Bailly, W. N. Eschmeyer, R. Froese	L. Collier	5	+++++ C
Pisces - Chondrichthyes	J. van der Land, M. J. Costello, D. Zavodnik, R. Serrão Santos, F. Mora Porteiro, N. Bailly, W. N. Eschmeyer, R. Froese	L. Collier	145	+++++ C
Pisces - Osteichthyes	J. van der Land, M. J. Costello, D. Zavodnik, R. Serrão Santos, F. Mora Porteiro, N. Bailly, W. N. Eschmeyer, R. Froese	L. Collier	1199	+++++ C
Tetrapoda - Aves	J. van der Land	M. Ramos, J. Templado	74	+++++ C
Tetrapoda - Reptilia	J. van der Land	M. Ramos, J. Templado	5	+++++ C
Tetrapoda - Mammalia	J. van der Land	M. Ramos, J. Templado	50	+++++ C
TOTAL			29,714	

SPECIES LISTS**FUNGI**

Compiled by Nicholas Clipson, Eleanor Landy and Marinus Otte

Fungi from coastal and marine ecosystems are a neglected but significant part of marine biodiversity. Fungi in general are able to degrade a wide range of recalcitrant biological molecules and particularly in coastal ecosystems fungal activity may be critical in the early stages of biodegradative pathways. Fungi in the marine environment have only been fully recognised since about 1960, and within the group, marine fungi have shown high decadal indices (% increases in species number over a 10 year period). Between 1981 and 1991, Hawksworth (1991, *Mycological Research* 95, 641-655) calculated that marine fungi had the highest decadal index (49%) for any fungal group. Furthermore, Hawksworth put forward the opinion that less than 10% of fungal biodiversity had yet been discovered. Recent collections of marine fungi have largely been restricted to the Far East, with only slight extension of the European list over the last 15 years. Although no precise figure presently exists for a total worldwide number of marine fungal species, this is probably in excess of 1000. Our provisional list for European marine fungi extends to 318 species, largely of ascomycetes and deuteromycetes, but also several basidiomycetes (see accompanying list).

Two problems have arisen during the production of this list. Firstly, maximal fungal activity in marine systems is found adjacent to terrestrial areas. No precise definition of a marine fungus exists, although they are generally defined as fungi that can complete lifecycles in marine environments. In coastal regions where fungal isolations are made many terrestrial species are found. The sporulating nature of fungi means that it is unclear whether such species exist actively in the marine environment and, except where there is clear evidence that they fulfil our definition, they have been excluded from this list. Secondly, the taxonomy of microeukaryotes is very confused and molecular evidence now restricts the Fungi to a monophyletic group consisting of chytridiomycetes, zygomycetes, ascomycetes and basidiomycetes. Only members of these have been included.

The list at present is preliminary and it is expected that further species will be added. Nevertheless, we believe that the list at present includes in excess of 90% of identified European species. We continue to identify and approach experts in this area to validate this list. We are grateful to Mr G. Bremer and the University of Portsmouth Marine Fungal Collection who have allowed us to use information from this collection and Dr. E.B.G. Jones of the University of Hong Kong for his expertise and editorial input.

References

Kohlmeyer, J. & Kohlmeyer, E. 1979 *Marine Mycology – The Higher Fungi*. Academic Press, New York.

KINGDOM FUNGI**PHYLUM ZYgomycota***incertae sedis*

Rhizopus stolonifer var. *stolonifer*

Class Zygomycetes**Order Mucorales****Family Mortierellaceae**

Mortierella
alpina Peyronel, 1913
gamsii Milko, 1974
minutissima Tiegh, 1878
parvispora Linnem., 1941
polycephala Coem., 1863
turficola Y. Ling, 1930

Family Mucoraceae

Absidia
spinosa Lendl., 1905
Actinomucor
repens Schostak, 1898
Mucor
hiemalis Wehmer, 1903
plumbeus Bonard, 1864
racemosus Tiegh, 1878

PHYLUM ASCOMYCOTA*incertae sedis***Order Diaporthales****Family Valsaceae**

Gnomonia
salina E.B.G. Jones, 1962

Order Dothideales**Family Arthopyreniaceae**

Leiophloea
pelvetia (G.K. Sutherl.) Kohlm. & E. Kohlm, 1979

Family Didymosphaeriaceae

Lautitia
danica (Berl.) Schatz, 1984

Family Herpotrichiellaceae

Capronia
ciliomaris (Kohlm.) E. Mull., Petrini & P.J. Fischer.
 Samuels & Rossman, 1987

incertae sedis

Didymella
fucicola (G.K. Sutherl.) Kohlm., 1968

Passeriniella
obiones (AM Crouan & P.L. Crouan) K.D. Hyde &
 Mouzouras, 1988

Pontogeneia

calospora (Pat.) Kohlm.
cnormis (Pat. & Hariot) Kohlm.
Thalassoascus
tregoubovii Ollivier
cystoseirae (Ollivier) Kohlmeyer

Family Leptosphaeriaceae

Leptosphaeria
arundinacea (Sowerby) Sacc., 1873
chondrii Rosenv., 1906
culmifraga Sacc., 1863
fuckelii Niessl., 1882
luctuosa Niessl ex Sacc., 1875
marina Rostr.
michotii (Westend.) Sacc., 1878
microscopica P. Karst, 1872
microspora Ellis & Everh.
neomaritima R. Gessner & Kohlm., 1976

- nigrans* (Desm.) Ces & De Not., 1863
orae-maris Linder, 1944
pelagica E.B.G. Jones, 1962
pontiformis (Fuckel) Sacc., 1883
spartinae Ellis & Everh., 1885
Ophiobolus
culturum
littoralis (P. Crouan & H. Crouan) Sacc.
- Family Lophiostomataceae**
Lophiotrema
littorale Speg.
Massarina
thalassiae Kohlm. & Volk.-Kohlm., 1987
- Family Microthyriaccae**
Microthyrium
graminum E. Bommer, M. Rousseau & Sacc., 1891
nigroannulatum J. Webster, 1952
- Family Monoblastiaceae**
Microthelia
linderi Kohlm., 1971
verruculosa Anastasiou
- Family Mycosphaerellaceae**
Micronectriella
agropyri Apinis & Chesters, 1964
Mycosphaerella
ascophylli Cotton, 1909
elymifoliae Munk, 1952
lineolata (Roberge ex Desm.) J. Schrot., 1894
pelvetiae Sutherland, 1915
salicorniae (Auerswald) Petrak, 1903
saticicola (Pat.) S. Dias
tassiana (De Not.) Johanson, 1884
- Pharcidia*
balani (Winter) Bauch, 1936
laminaricola Kohlm.
rhachiana Kohlm.
- Sphaerulina*
albispiculata Tubaki
orae-maris Linder, 1944
- Family Phaeosphaeriaceae**
Lautitia
danica (Berlesse) Schatz, 1984
Phaeosphaeria
albopunctata (Westend.) Shoemaker & C.E. Babcock, 1989
ammophilae (Lasch) Kohlm. & E. Kohlm., 1979
macrosporidium (E.B.G. Jones) Shoemaker & Babcock, 1989
typharum (Desm.) L. Holm, 1957
- Family Pleomassariaceae**
Kirschsteinothelia
maritima (Linder) D.L. Hawks, 1985
- Family Pleosporaceae**
Pleospora
gaudefroyi Pat., 1886
herbarum P. Karst, 1863
insectoria Fuckel, 1870
lignicola J. Webster & Lucas, 1961
pelvetiae G.K. Sutherl., 1915
rubelloides (Plowr. ex Cooke) J. Webster, 1957
rubicunda Niessl., 1876
spartinae (J. Webster & Lucas) Apinis & Chesters, 1964
straminis Sacc. & Speg., 1879
trichostoma (Fr.) Fuckel
triglochinicoila J. Webster, 1969
- Trematia*
halophila Kohlm. & Volk.-Kohlm. & O.E. Erikss., 1995
- Family Sporomiaceae**
Preussia
funiculata Fuckel., 1870
terricola Cain, 1961
- vulgaris*
- Family Zopfiaceae**
Halothia
posidoniae (Dur. & Mont.) Kohlm., 1963
Pontoporeia
biturbinata (Dur. & Mont.) Kohlm.
- Order Eurotiales**
- Family Monascaceae**
Monascus
purpureus Went, 1895
- Family Trichocomaceae**
Byssochlamys
nivea Westling, 1909
Emericella
nidulans (Eidam) Vuill., 1927
Emericellopsis
stolkiae Davidson & Christensen.
terricola J.F.H.. Beyma, 1939
Talaromyces
dupontii (Griffon & Maubl.) Apinis, 1963
- Order Halosphaeraiales**
- Family Halosphaeriaceae**
Aniptodera
chesapeakensis C. Shearer & J. Miller
Anisostagma
rotundatum K.R.L. Petersen & J. Koch
Arenariomyces
trifurcatus Hohnk., 1954
Appendichordella
amicta (Kohlm.) Johnson, Jones & Moss, 1987
Bovicornua
intricata Jorgen Koch & E.B.G. Jones, 1993
Carbosphaerella
leptosphaerooides I. Schmidt
pleosporoides I. Schmidt
Ceriosporopsis
caduca E.B.G. Jones & Zainal, 1988
cambrensis I.M. Wilson, 1954
halima Linder, 1944
tubulifera (Kohlm.) P.W. Kirk ex Kohlm., 1972
Chaedosaudia
corallinarum (P. Crouan & H. Crouan) E. Mull. & Arx, 1973
Corollospora
armoricana Kohlm. & Volk.-Kohlm., 1997
intermedia I. Schmidt
lacera (Linder) Kohlm.
luteola Nakagiri and Tubaki, 1982
maritima Werderm., 1922
pulchella (Kohlm.) I. Schmidt & Nair
Haligena
elaterophora Kohlm., 1961
salina C.A. Farrant & E.B.G. Jones, 1986
Halosphaeria
appendiculata Linder, 1944
Halosarpheia
hamata
retoquens Shearer & J.L. Crane, 1981
spartinae (E.B.G. Jones) C. Shearer & J.L. Crane, 1981
trullifera (Kohlm.) E.B.G. Jones, S.T. Moss & Cuomo, 1983
unicaudata (E.B.G. Jones & LeChampion-Alsumard) R.G. Johnson, E.B.G. Jones & S.T. Moss, 1987
viscosa (I Schmidt) Shearer & J.L. Crane ex Kohlm. & Volk.-Kohlm., 1991
Halosphaeriopsis
mediosetigera (Cribb & J.W. Cribb) T.W. Johnson, 1958
Iwilsoniella
rotunda E.B.G. Jones, 1991
Kohlmeyeriella
tubulata (Kohlm.) E.B.G. Jones, R.G. Johnson & S.T.

- Moss, 1983
Lautosporopsis
circumvestita (Kohlm.) E.B.G. Jones, Yusoff & S.T.
Moss
Ligninicola
laevis Hohnk, 1955
Limacospora
sundica Jorgen Koch & E.B.G. Jones, 1995
Lindra
inflata I.M. Wilson, 1956
marinera Meyers
Lulworthia
crassa Nagakiri, 1984
floridana Meyers, 1957
fucicola G.K. Sutherl., 1916
kneippii Kohlm.
lignoarenaria J. Koch & E.B.G. Jones, 1984
medusa (Ellis & Everh.) Cribb & J.W. Cribb, 1955
purpurea (I.M. Wilson) T.W. Johnson, 1958
rufa (I.M. Wilson) T.W. Johnson, 1958
uniseptata Nagakiri, 1984
Luttrellia
estuarina Shearer
Marinospora
longissima (Kohlm.) A.R. Caval.
calyprata (Kohlm.) A.R. Caval, 1960
Mycophycophila
corallinarum
Nais
inornata Kohlm., 1962
Naupagella
spinibarbata (J. Koch) Kohlm. & Volkm.-Kohlm., 1998
Nautosphaeria
cristaminuta E.B.G. Jones, 1964
Nereiospora
comata (Kohlm.) E.B.G. Jones, Johnson, R.G. &
Moss, S.T., 1983
cristata (Kohlm.) E.B.G. Jones, Johnson, R.G. &
Moss, S.T., 1983
Ocostaspora
apilongissima E.B.G. Jones, R.G. Johnson & Moss,
S.T., 1983
Ondiniella
torquata (Kohlm.) Jones, Johnson & Moss, 1984
Remispora
crispa Kohlm., 1981
galerita Tubaki
maritima Linder, 1944
pilleata Kohlm.
quadriremis (Hohnk) Kohlm.
stellata Kohlm.
Tirispora
unicaudata E.B.G. Jones & Vrijmoed
Trailia
ascophylli G.K. Sutherl., 1915
- Order Hypocreales**
- Family Clavicipitaceae**
Barya
aurantiaca Plowr. & A.S. Wilson, 1884
Claviceps
purea (Fr.: Fr.) Tul., 1883
- Family Hypocreaceae**
Halonectria
milfordensis E.B.G. Jones, 1965
Nectria
gliocladioides Smalley & H.N. Hansen, 1957
inventa Pethybr., 1919
Nectriella
laminariae O. Erikss
- incertae sedis*
- Family Dactylosporaceae**
Dactylospora
- canariensis* Kohl. & Volkm.-Kohlm., 1995
incertae sedis
Amylocarpus
encephaloides Currey, 1986
Botrylloides
physasca Jorgen Koch & E.B.G. Jones, 1989
Botrytis
cinerea Pers. ex Nocca & Balb., 1821
Chaetomastix
typhicola (P. Karst) M.E. Barr, 1989
Crinigera
maritima I. Schmidt, 1989
Cryptovalsa
suaedicola Spooner, 1981
Drysosphaera
navigans J. Koch & E.B.G. Jones, 1989
Eiona
tunicata Kohlm.
Marisolaris
ansata J. Koch & E.B.G. Jones, 1989
Orcadia
ascophylli G.K. Sutherl., 1915
Peziza
paulula
Phomatospora
dinemasperium J. Webster, 1955
Pseudoeurotium
zonatum Van Beyma, 1937
Sphaerooides
fimicola (E.C. Hansen) P.F. Cannon & D.L. Hawks., 1982
Torpedospora
ambispinosa Kohlm.
radiata Meyers, 1957
Tunicatispora
australiensis K.D. Hyde, 1990
- Family Magnaportheaceae**
Gaeumannomyces
graminis var. *tritici* Walker
- Order Leotiales**
- Family Dermateaceae**
Mollisia
arenevaga (Desm.) W. Phillips, 1887
arundinacea (D.C.) W. Phillips, 1887
palustris (Robege ex Desm.) P. Karst, 1871
- Pyrenopezizidae**
graminis Sacc., 1883
- Family Hyaloscyphaceae**
Dasyphyllum
acutipilus (P. Karst) Sacc., 1889
carneolus (Sacc.) Sacc., 1889
controversus Cooke, 1881
palearum (Desm.) Massee, 1887
rhodoleucus (Sacc.) Sacc., 1889
- Order Microascales**
- Family Microascaceae**
Microascus
cirrosus Curzi, 1930
- Order Onygenales**
- Family Gymnoascaceae**
Arachniotus
albicans Apinis, 1964
ruber (Teigh) J. Schrot., 1893
- Gymnoascus**
reesi Baran.
roseus (Raillo) Apinis, 1964
- Family Myxotrichaceae**
Myxotrichum
ochraceum Berk. & Broome, 1875
- Family Onygenaceae**
Auxoarthron

umbrinum (Boud.) G.F. Orr & Plunkett, 1963

Order Pezizales

Family Ascobolaceae

Ascobolus

crenulatus P. Karst., 1868

microsporus Berk & Broome, 1865

Ascophanus

carneus (Pers: Fr.) Boudier, 1869

Family Otideaceae

Humaria

rutilans (Fr.) Sacc., 1889

Order Phyllochoraceae

Family Phyllochorales

Haloguignardia

cystoseira Kolhm. & Demoulin, 1981

Marinosphaera

mangrovei K.D. Hyde, 1989

Order Pyrenolales

Family Pyrenolaceae

Pyrenocolla

pelvetica (G.K. Sutherl.) D.L. Hawks., 1988

Order Saccharomycetales

Family Metschnikowiaceae

Metschnikowia

bicuspidata (Metschnikoff) Kamienski, 1899

Order Sordariales

Family Chaetomiaceae

Chaetomium

aterrimum Ellis & Everh., 1910

bostrychodes Zopf., 1877

cochlioides Palliser, 1910

elatum Kunze & Schmidt, 1824

erectum Skolko & J.W. Groves, 1948

funicola Cooke, 1873

globosum Kunze: Fr., 1824

indicum Corda, 1840

Thielavia

sepedonium C.W. Emmons, 1932

terricola (J.C. Gilman & E.V. Abbott) C.W. Emmons, 1930

Family Chaetosphaeriaceae

Chaetosphaeria

chaetosa Kohlmeyer

Family Coniochaetaceae

Coniochaeta

lignaria (Grev.) Massee, 1887

incertae sedis

Savoryella

lignicola E.B.G. Jones & Eaton, 1969

Family Lasiosphaeriaceae

Apiospora

montagnei Sacc., 1875

Biconiosporella

corniculata Schaumann

Phaeonectriella

lignicola Eaton & E.B.G. Jones, 1971

Pleurage

curvula (de Bary ex G. Winter) Kuntze, 1898

decipiens (G. Winter ex Fuckel) Kuntze, 1898

minuta (Fuckel) Kuntze, 1898

Family Nitschkiaceae

Groenhiella

bivestia J. Koch, E.B.G. Jones & S.T. Moss, 1983

Family Sordariaceae

Gelaninospora

calospora

Gelasinospora

retispora Cain., 1950

Sordaria

fimicola (Roberge ex Desm.) Ces. & De Not., 1863

tetraspora G. Winter, 1871

Order Xylariales

Family Amphisphaeriaceae

Amphisphaeria

culmicola Sacc., 1873

Family Xylariaceae

Anthostomella

lugubris (Roberge & Desm.) Sacc., 1882

phaeosticta (Berk.) Sacc., 1878

PHYLUM BASIDIOMYCOTA

Class Basidiomycetes

Order Agaricales

Family Agaricaceae

Psalliota

litoralis A. Pearson & Wakef.

Family Tricholomataceae

Helotium

robustum P. Karst.

Laccaria

trullisata (Ellis) Peck.

Order Cortinariales

Family Cortinariaceae

Inocybe

decipiens Bres.

Order Melanogastrales

Family Niaceae

Nia

epidermoidea Rossello and Descals, 1993

globospora Barrata & Basilio, 1997

vibrissa R.T. Moore & Meyers, 1961

Order Stereales

Family Amylocorticaceae

Digitatispora

marina Douget, 1962

Order Tulostomatales

Family Tulostomataceae

Tulostoma

macrocephalum Long.

Class Tiliaceae

Order Ustilaginales

Family Usticomycetes

Melanotaenium

ruppiiæ G. Feldmann

MITOSPORIC FUNGI

Acremonium

strictum

Allescheria

terrestris Apinis, 1963

Alternaria

alternata (Fr.: Fr.) Keissl., 1912

maritima G.K. Sutherl., 1916

tenuis Nees., 1822

Arthropsis

hispanica Gene, Ulfig & Guarro, 1995

Ascochyta

obiones (Jaap) P.K. Buchanan, 1987

Aspergillus

fumigatus Fresen., 1863

glaucus (L.: Fr.) Link, 1863

nidulans (Eidam) G. Winter, 1884

niger Tiegh

terreus Thom., 1918

ustus (Bainier) Thom. & Church

versicolor (Vuill.) Tirab.

Asteromyces

cruciatus Moreau & M. Moreau, 1941

<i>Camarosporium</i>	<i>Scopulariopsis</i>
<i>obiones</i> Jaap, 1905	<i>brevicaulis</i> (Sacc.) Bainier, 1907
<i>metableticum</i> Trail	<i>fusca</i> Zach, 1934
<i>roumequerii</i> Sacc.	<i>Sigmoidea</i>
<i>Cephalosporium</i>	<i>marina</i> Haythorn & E.B.G. Jones, 1980
<i>curtipes</i> Sacc., 1881	<i>Sporidesmium</i>
<i>Cephalotrichum</i>	<i>salinum</i> E.B.G. Jones, 1963
<i>microsporum</i> (Sacc.) Kirk, 1984	<i>Stemphylium</i>
<i>nanum</i> (Ehrenb.)	<i>maritimum</i> Johnson, 1957
<i>stemonitis</i> (Pers.) Nees, 1809	<i>triglochincola</i> Sutton & Pirozynski, 1963
<i>Chrysosporium</i>	<i>Trichocladium</i>
<i>panorum</i> (Link) S. Hughes, 1958	<i>achrasporum</i> Meyers & R.T. Moore, 1971
<i>Cirrenalia</i>	<i>alopallonellum</i> (Meyers & R.T. Moore) Kohlm. &
<i>fusca</i> I. Schmidt, 1985	Volkm.-Kohlm., 1995
<i>macrocephala</i> (Kohlm.) Meyers & R.T. Moore, 1960	<i>asperum</i> Harz., 1871
<i>Cladosporium</i>	<i>constrictum</i> I. Schmidt, 1974
<i>alagrum</i> Cooke & Massee in Cooke	<i>lignicola</i> I. Schmidt, 1974
<i>herbarum</i> (Pers.: Fr.) Link, 1821	<i>Trichoderma</i>
<i>macrocarpum</i> Preuss, 1848	<i>koningii</i> Oudem., 1902
<i>Clavariopsis</i>	<i>viride</i> Pers.: Fr., 1829
<i>aquatica</i> De Wild.	<i>Tolypocladium</i>
<i>Colletotrichum</i>	<i>geodes</i> W. Gams, 1971
<i>typhae</i> H.C. Greene, 1955	<i>Tetracladium</i>
<i>Coniothyrium</i>	<i>marchalianum</i> De Wild.
<i>obiones</i> Jaap, 1907	<i>Trichothecium</i>
<i>Cumulospora</i>	<i>roseum</i> (Pers.: Fr.) Link
<i>marina</i> I. Schmidt, 1985	<i>Tubicularia</i>
<i>Cylindrocarpon</i>	<i>pulverulenta</i> Spegazzini
<i>destructans</i> (Zinssm.) Scholten, 1964	<i>Verticillium</i>
<i>Dendryphiella</i>	<i>bulbillosum</i> W. Gams & Malla, 1971
<i>arenaria</i> Nicot, 1958	<i>Volutella</i>
<i>salina</i> (G.K. Sutherl.) Pugh & Nicot., 1976	<i>ciliata</i> (Alb. & Schwein.: Fr.) Fr., 1832
<i>Dictyosporium</i>	<i>Zalerion</i>
<i>pelagica</i> (Corda) Gueg., 1905	<i>varium</i> Anastasiou
<i>toruloides</i> (Corda) Gueg., 1905	
<i>Fusarium</i>	
<i>culmorum</i> (W.G. Sm.) Sacc., 1895	
<i>oxysporum</i> Schiltl.: Fr., 1824	
<i>solani</i> (Mart.) Sacc., 1881	
<i>Gliocladium</i>	
<i>roseum</i> Bainier, 1907	
<i>Gliomastix</i>	
<i>guttuliformis</i> Brown & Kendrick, 1958	
<i>murorum</i> (Corda) S. Hughes, 1958	
<i>Macrophoma</i>	
<i>gymnogongri</i>	
<i>Monodictys</i>	
<i>pelagica</i> (T.W. Johnson) E.B.G. Jones, 1963	
<i>putredinis</i> (Wallr.) S. Hughes, 1958	
<i>Oidiodendron</i>	
<i>maius</i> Barron, 1962	
<i>tenuissimum</i> (Peck) S. Hughes, 1958	
<i>Orbimyces</i>	
<i>spectabilis</i> Linder, 1944	
<i>Ostracoblabe</i>	
<i>implexa</i> Born & Flahault	
<i>Paecilomyces</i>	
<i>aegyptiacus</i> S. Ueda & Udagawa, 1983	
<i>Papularia</i>	
<i>sphaerosperma</i> (Per.: Fr.) Hohnk, 1916	
<i>Penicillium</i>	
<i>chrysogenum</i> Thom., 1910	
<i>digitatum</i> (Pers. ex St.-Am.) Sacc., 1882	
<i>frequentans</i> Westling, 1911	
<i>spinulosum</i> Thom., 1910	
<i>variabile</i> Sapp., 1912	
<i>Phialophorophoma</i>	
<i>litoralis</i> Linder	
<i>Phoma</i>	
<i>laminariae</i> Cooke & Massee in Cooke	
<i>suaedae</i> Jaap, 1907	
<i>Pullularia</i>	
<i>pullulans</i> (de Bary) Berkhoult	
<i>Scolecobasidium</i>	
<i>salinum</i> (G.K. Sutherl.) M.B. Ellis	

PLANTS

Compiled by Michael Guiry

This list includes all benthic (attached) marine algae belonging to the phyla (divisions) Chlorophycota (Chlorophyta), Phaeophycota (Phaeophyta), Rhodophycota (Rhodophyta) and the genera *Vaucheria* and *Tribonema* (Xanthophycota) that are generally known as seaweeds. It also includes the seagrasses (Magnoliophyta). This list is extracted from a worldwide database of algae and seagrasses being prepared as part of the Species 2000 project. The database is searchable at:

<http://www.algaebase.org>

where it incorporates detailed references, notes, some synonyms, common names, some pictures, and much of the distributional data that forms part of the present list. As this database is "live" it may not include exactly the same data in the list below; it does, however, explain much of the sources for the records included here.

The present list was compiled and checked by Michael Guiry. Its initial foundation was the species listed in the *Species Directory of the Marine Fauna and Flora of the British Isles and Surrounding Seas* (Guiry, 1997), which was then systematically built up from the most recent key works and check-lists for each geographical area as follows:

North Eastern Atlantic: East Greenland (Pedersen, 1976), Iceland (Caram & Jónsson, 1972; Munda, 1979), Spitsbergen (Vinogradova, 1995), Faroes (Irvine, 1982), Norway (Rueness, 1997), Baltic Sea (Nielsen *et al.*, 1995), Netherlands (Stegenga *et al.*, 1997), Belgium and France (Coppejans, 1995; Feldmann, 1954; Feldmann & Magne, 1964), Northern Spain (Veiga, Cremades & Bárbara, 1998, and others), Portugal (Ardé, 1970, and others), Southern Spain (Seoane-Camba, 1965; Conde *et al.*, 1996, and others), Morocco (Dangeard, 1949), Madeira and the Salvage Islands (Levrin, 1974; Audiffred & Weisscher, 1984), and the Azores (Neto, 1994; Tittley & Neto, 1994). Unpublished lists from Helgoland and the Canary Islands provided by Drs Bartsch and Kuhlenkamp and Drs Haroun, Gil-Rodríguez and Díaz de Castro, respectively, were invaluable. This list is not exhaustive but is included to give an indication of the sources.

Mediterranean: The excellent brown (Ribera *et al.*, 1992) and green (Gallardo *et al.*, 1993) check-lists were used as the basis for an initial list; however, no overall summary exists for the red algae. The Mediterranean list was completed from a wide range of sources which are far too numerous to cite here but which are listed in the database version cited above.

The overall arrangement of phyla (divisions) classes, orders and families is largely that of Silva, Basson & Moe (1996) with some modifications resulting from later work. Individual taxonomic decisions are too numerous to detail here but are also listed in the web version of the database. Categories below the level of species (subspecies, varieties and formae) are not included here for reasons of space. Nomenclatural authorities are given in full and, as far as possible, fit in with the full names as listed in Brummitt & Powell (1992). Non-botanists should note that it is not the normal practice to cite the date of publication or date of combination for plants; however, I have, for convenience, included the date

when known. Those familiar with the *International Code of Zoological Nomenclature* should also note that it is the practice under the *Botanical Code* to cite the original author(s) in parentheses and the author(s) making a new combination outside such parentheses. So, there are only two possibilities: there may be the name of an authority or authorities with no parentheses or there will be the names of an authority or authorities in parentheses with another name or names outside it. Botanical authors should note the legitimate use of diacritical marks in certain instances (e.g., *Hydrolithon samoënsis*; *Schottera nicaëensis*) to facilitate correct pronunciation. It should also be noted that whilst "ex" is part of nomenclatural authority citation, "in" is part of bibliographical citations and should not be used except in this context. The following recently altered spellings should also be noted: *Acanthophora nayadiformis* for *A. najadiformis*; *Chondracanthus teedii* for *C. teedii*; and *Halopithys* for *Halopitys*. A number of nomenclaturally illegitimate names are still included (but not indicated) such as *Sargassum vulgare*. The binomial *Scytoniphon lomentaria* is included on the assumption that its proposed conservation will be accepted. A number of doubtful records, such as *Ecklonia buruncinata* from the Canary Islands, are included pending clarification. The genus *Pilinia* is included in the Phaeophycota even though there is doubt as to whether *P. rimosa*, the type, is a green or a brown alga.

The seagrass lists were compiled from a variety of sources and the nomenclature is that of Phillips & Meñez (1988).

I am grateful to all those who helped me in the compilation of this list, in particular to Professor G. Furnari, Dr Fabio Rindi, Ms Eilís Nic Dhonncha, and Ms Sandy Lawson.

References

- Ardé, F. 1970. Contribution à l'étude des algues marines du Portugal. I. La flore. *Portugalia Acta Biologica* sér. B, 10: 137-555.
- Audiffred, P. A. J. and Weisscher, F. L. M. 1984. Marine algae of Selvagem Grande (Salvage Islands, Macaronesia). *Boletim do Museu Municipal do Funchal* 36: 5-37.
- Brummitt, R. K. and Powell, C. E., Eds. 1992. *Authors of Plant Names. A list of authors of scientific names of plants, with recommended standard forms of these names including abbreviations*. Key, Royal Botanic Gardens.
- Caram, B. and Jónsson, S. 1972. Nouvel inventaire des algues marines de l'Islande. *Acta Botanica Islandica* 1: 5-31.
- Conde, F., Flores-Moya, A., Soto, J., Altamirano, M. and Sánchez, A. 1996. Check-list of Andalusia (S. Spain) seaweeds. III. Rhodophyceae. *Acta Botanica Malacitana* 21: 7-33.
- Coppejans, E. 1995. *Flore algologique des côtes du Nord de la France et de la Belgique*. Meise, Jardin Botanique National de la Belgique.
- Dangeard, P. 1949. Les algues marines de la côte occidentale du Maroc. *Botaniste* 34: 89-189.
- Feldmann, J. 1954. Inventaire de la flore marine de Roscoff. Algues, champignons, lichens et spermatophytes. *Travaux Station Biologique de Roscoff* Ser. 2, Suppl. 6: 152.
- Feldmann, J. and Magne, M. F. 1964. Additions à l'inventaire de la flore marine de Roscoff algues, champignons, lichens. *Travaux Station Biologique de Roscoff* 15 (New supplement): 1-23 [+ 5].
- Gallardo, T., Gómez Garreta, A., Ribera, M. A., Cormaci, M., Furnari, G., Giaccone, G. and Boudouresque, C. F. 1993. Check-list of Mediterranean Seaweeds, II. Chlorophyceae Wille s.l. *Botanica Marina* 36: 399-421.
- Guiry, M. D. 1997. Benthic red, brown and green algae. *The Species Directory of the Marine Fauna and Flora of the British Isles and Surrounding Seas*.

- British Isles and Surrounding Seas.* C. M. Howson and B. E. Picton. Belfast & Ross-on-Wye, Ulster Museum & Marine Conservation Society: 341-367.
- Irvine, D. E. G. 1982. Seaweeds of the Faroes I: The flora. *Bulletin British Museum Natural History, Botany* 10: 109-131.
- Levrang, T. 1974. The marine algae of the Archipelago of Madeira. *Boletim Museu Municipal do Funchal* 28: 5-111.
- Munda, I. M. 1979. Addition to the check-list of benthic marine algae from Iceland. *Botanica marina* 22: 459-463.
- Neto, A. I. 1994. Checklist of the benthic marine macroalgae of the Azores. *Arquipélago. Ciências Biológicas e Marinhas* 12A: 15-34.
- Nielsen, R., Kristiansen, A., Mathiesen, L. and Mathiesen, H. 1995. Distributional index of the benthic marine macroalgae of the Baltic Sea area. *Acta Botanica Fennica* 155: 1-70.
- Pedersen, P. M. 1976. Marine benthic algae from southernmost Greenland. *Meddelelser om Grönland* 199(3): 1-80.
- Phillips, R. C. and Meñez, E. G. 1988. *Seagrasses*. Washington, D.C., Smithsonian Institution Press.
- Ribera, M. A., Gómez-Garreta, A., Gallardo, T., Cormaci, M., Furnari, G. and Giaccone, G. 1992. Check-list of Mediterranean Seaweeds. I. *Fucophyceae* (Warming 1884). *Botanica marina* 35: 109-130.
- Rueness, J. 1997. Algae. *Distribution of marine, benthic macro-organisms in Norway. A tabulated catalogue, preliminary Edition. Research Report No. 1997-1.* T. Brattegard and T. Holthe. Trondheim, Direktoratet for Naturforvaltninhg.
- Seoane-Camba, J. 1965. Estudios sobre las algas bentónicas en la costa sur de la Península Ibérica (litoral de Cádiz). *Investigacion Pesquera* 29: 3-216.
- Silva, P. C., Basson, P. W. and Moe, R. L. 1996. Catalogue of the benthic marine algae of the Indian Ocean. *University of California Publications in Botany* 79: 1-1259.
- Stegenga, H., Mol, I., Prud'homme van Reine, W. F. and Lokhorst, G. M. 1997. Checklist of the marine algae of the Netherlands. *Gorteria Supplement* 4: 3-57.
- Tittley, I. and Neto, A. I. 1994. "Expedition Azores 1989": Benthic marine algae (seaweeds) recorded from Faial and Pico. *Arquipélago, Life and Marine Sciences* 12: 1-13.
- Veiga, A. J., Cremades, J. and Bárbara, I. 1998. A catalogue of the marine benthic algae of the Sisargas Islands (N.W. Iberian Peninsula, Spain). *Boletim do Museu Municipal do Funchal Suppl. no. 5:* 481-493.
- Vinogradova, K. L. 1995. The checklist of the marine algae from Spitsbergen. *Botanicheskii Zhurnal SSR* 80: 50-61.

KINGDOM ALGAE

PHYLUM CHLOROPHYCOTA

Class Chlorophyceae

Order Acrosiphoniales

Family Acrosiphonaceae

Acrosiphonia

<i>arcata</i> (Dillwyn) Gain, 1912	A
<i>arctiuscula</i> (Kützing) Jónsson	A
<i>flagellata</i> Kjellman	A
<i>grandis</i> Kjellmann, 1893	A
<i>incurva</i> Kjellman	A
<i>sonderi</i> (Kützing) Kornmann, 1962	A
<i>spinescens</i> (Kützing) Kjellmann, 1883	A

Spongomorpha

<i>aeruginosa</i> (Linnaeus) van den Hoek, 1963	A M
<i>centralis</i> (Lyngbye) Kützing	A
<i>vernalis</i> (Kjellman) Jónsson, 1903	A

Urospora

<i>elongata</i> (Rosenvinge) Hagem	A
<i>hartzii</i> Rosenvinge	A
<i>laeta</i> (Thuret ex Bornet) Børgesen	A M
<i>neglecta</i> (Kornmann) Lokhorst & Trask, 1981	A
<i>penicilliformis</i> (Roth) Areschoug, 1874	A M
<i>wormskioldii</i> (Mertens ex Hornemann) Rosenvinge, 1893	A

Order Bryopsidales

Family Bryopsidaceae

Bryopsisella

<i>neglecta</i> (Berthold) H. Rietema, 1975	A M
<i>ostreobiformis</i> Calderón-Sáenz & Schnetter	M

Bryopsis

<i>adriatica</i> (J. Agardh) Frauenfeld, 1854	A M
<i>balbisiana</i> J.V. Lamouroux ex Harvey	A M
<i>corymbosa</i> J. Agardh, 1842	A M
<i>cupressina</i> J.V. Lamouroux	A M
<i>dichotoma</i> De Notaris	M
<i>duplex</i> De Notaris	A M
<i>feldmannii</i> Gallardo & G. Furnari	M
<i>hypnoides</i> J.V. Lamouroux, 1809	A M
<i>implexa</i> De Notaris, 1849	M
<i>lyngbyei</i> Hornemann, 1818	A
<i>muscosa</i> J.V. Lamouroux, 1809	M
<i>penicillum</i> Meneghini	M
<i>pennata</i> J.V. Lamouroux, 1809	A M
<i>plumosa</i> (Hudson) C. Agardh, 1823	A M
<i>secunda</i> J. Agardh	M
<i>Trichosolen</i>	
<i>myura</i> (J. Agardh) W.R. Taylor, 1962	A M

Family Derbesiaceae

Derbesia

<i>boergesenii</i> (Iyengar & Ramanathan) Mayhoub, 1976	M
<i>corallicola</i> Funk	M
<i>marina</i> (Lyngbye) Solier, 1846	A M
<i>minima</i> Funk	M
<i>tenuissima</i> (Moris & De Notaris) P.L. Crouan & H.M. Crouan, 1867	A M

Pedobesia

<i>lamourouxi</i> (J. Agardh) Feldmann, Loreau, Codomier & Couté, 1975	A M
<i>solieri</i> Feldmann ex Abéard & Knoepffler	A M

Order Caulerpales

Family Caulerpaceae

Caulerpa

<i>cupressoides</i> (Vahl) C. Agardh, 1817	A
<i>mexicana</i> Sonder ex Kützing, 1849	A M
<i>ollivieri</i> Dostál	M
<i>peltata</i> J.V. Lamouroux, 1809	A
<i>prolifera</i> (Forsskål) J.V. Lamouroux, 1809	A M
<i>racemosa</i> (Forsskål) J. Agardh, 1873	A M
<i>seal pelliformis</i> (R. Brown ex Turner) C. Agardh, 1817	A M
<i>sertularioides</i> (S.G. Gmelin) M. Howe, 1905	A M
<i>subserrata</i> Okamura, 1897	M
<i>taxifolia</i> (M. Vahl) C. Agardh, 1817	A M
<i>webbiana</i> Montagne, 1837	A

Family Chaetosiphonaceae

Blastophysa

<i>rhizopus</i> Reinke, 1889	A M
------------------------------	-----

Chaetosiphon

<i>moniliformis</i> Huber, 1892	M
---------------------------------	---

Family Udoteaceae

Avrainvillea

<i>canariensis</i> A. Gepp & E.S. Gepp	A
--	---

Halimeda

<i>discoidea</i> Decaisne, 1842	A
<i>opuntia</i> (Linnaeus) J.V. Lamouroux, 1812	M
<i>tuna</i> (J. Ellis & Solander) J.V. Lamouroux, 1816	A M

Penicillus

<i>capitatus</i> Lamarck, 1813	A M
--------------------------------	-----

Pseudochlorodesmis

<i>furcellata</i> (Zanardini) Børgesen, 1925	A M
<i>tenuis</i> Ercegovic	M

Rhipilia

<i>tenaculosa</i> A. Gepp & E.S. Gepp	A
---------------------------------------	---

Order Chaetophorales

Family Chaetophoraceae

Arthrochaete

<i>penetrans</i> Rosenvinge, 1898	A
-----------------------------------	---

Bolbocoleon

<i>piliferum</i> Pringsheim, 1862	A M
-----------------------------------	-----

<i>Didymosporangium</i>			
<i>repens</i> Lambert	M	<i>stellata</i> (Wulfen) C. Agardh, 1822	A M
<i>Entocladia</i>		<i>Microdictyon</i>	
<i>endolithica</i> (Ercegovic) R. Nielsen	M	<i>boergesenii</i> Setchell, 1925	A
<i>leptocheete</i> (Huber) Burrows, 1991	A M	<i>calodictyon</i> (Montagne) Duchesne	A
<i>major</i> (Feldmann) R. Nielsen	M	<i>laxereticulatum</i> Setchell	M
<i>majus</i> (Feldmann) R. Nielsen	M	<i>tenius</i> Decaisne ex J.E. Gray, 1866	A M
<i>pennata</i> (Feldmann) R. Nielsen	M	<i>umbilicatum</i> (Velley) Zanardini	M
<i>reticulata</i> (Printz) Levring	A		
<i>Entodictyon</i>		Family Boodleaceae	
<i>schilleri</i> Schiffner, 1931	M	<i>Phyllodictyon</i>	
<i>Periplegmatium</i>	M	<i>anastomosans</i> (Harvey) Kraft & M.J. Wynne	A
<i>ceramii</i> Kützing, 1843	M	<i>pulcherrimum</i> J.E. Gray	A
<i>Pringsheimiella</i>			
<i>conchyliophila</i> Feldmann, 1937	A M	Family Cladophoraceae	
<i>sanctae-luciae</i> R. Nielsen & McLachlan	A	<i>Bryobesia</i>	
<i>scutata</i> (Reinke) Höhnel ex Marchewianka, 1924	A M	<i>johannae</i> Weber-van Bosse	A
<i>Pseudendoconium</i>		<i>Chaetomorpha</i>	
<i>conchiliophila</i> Feldmann	M	<i>adriani</i> Feldmann, 1937	M
<i>marinum</i> (Reinke) Aleem & E. Schulz, 1952	A M	<i>antennina</i> (Bory) Kützing, 1849	A
<i>submarinum</i> Wille, 1901	A M	<i>crassa</i> (C. Agardh) Kützing, 1845	A M
<i>Pseudopringsheimia</i>		<i>fibrosa</i> Kützing	A
<i>confluens</i> (Rosenvinge) Wille, 1909	A M	<i>gracilis</i> Kützing, 1845	A M
<i>Stromatella</i>		<i>implexa</i> (Harvey) Feldmann	A
<i>monostromatica</i> (P.J.L. Dangeard) Kornmann &	A M	<i>ligustica</i> (Kützing) Kützing, 1849	A M
Sahling		<i>linum</i> (O.F. Müller) Kützing, 1845	A M
<i>papillosa</i> (P.J.L. Dangeard) Kornmann & Sahling	A M	<i>litoraea</i> Harvey, 1858	A M
<i>Syncyrene</i>		<i>melagonium</i> (F. Weber & D. Mohr) Kützing, 1845	A
<i>reinkei</i> R. Nielsen & P.M. Pedersen, 1977	A	<i>pachynema</i> (Montagne) Kützing, 1847	A M
<i>Tellamia</i>		<i>zernovii</i> Woronichin	M
<i>contorta</i> Batters, 1895	A M		
<i>Ulvella</i>		Cladophora	
<i>acervus</i> P.J.L. Dangeard	A M	<i>aegagropila</i> (Linnaeus) Trevisan, 1845	A M
<i>lens</i> P.L. Crouan & H.M. Crouan, 1859	A M	<i>albida</i> (Nees) Kützing, 1843	A M
<i>nadsonii</i> (Rochlina) Gallardo <i>et al.</i>	M	<i>battersii</i> van den Hoek	A M
<i>setchellii</i> P.J.L. Dangeard, 1931	A M	<i>booleoides</i> Børgesen	A
<i>stellata</i> P.J.L. Dangeard, 1965	M	<i>catenata</i> (Linnaeus) Kützing, 1843	M
<i>Uronema</i>		<i>coelothrix</i> Kützing	A M
<i>curvata</i> Printz	A	<i>conferta</i> P.L. Crouan & H.M. Crouan, 1865	A
<i>Zygomitrus</i>		<i>crispata</i> (Roth) J. Agardh	A
<i>reticulatus</i> Bornet & Flahault	A	<i>crystallina</i> (Roth) Kützing, 1843	A M
Order Chlorococcales		<i>cymopoliae</i> Børgesen, 1925	A
Family Chlorococcaceae		<i>dalmatica</i> Kützing, 1843	A M
<i>Characium</i>		<i>echinus</i> (Biasoletto) Kützing, 1849	A M
<i>marinum</i> Kjellman, 1883	A	<i>feredai</i> Harvey	A M
<i>Chlorococcum</i>		<i>flexuosa</i> (O.F. Müller) Kützing, 1843	A M
<i>submarinum</i> Alvik, 1934	A	<i>fracta</i> (O.F. Müller ex Vahl) Kützing	A M
<i>Coccomyxa</i>		<i>globulina</i> (Kützing) Kützing	A M
<i>astericola</i> Rosenvinge	A	<i>glomerata</i> (Linnaeus) Kützing	A M
<i>litoralis</i> (Hansgirg) Wille	A	<i>hutchinsiae</i> (Dillwyn) Kützing, 1845	A M
<i>Sykidion</i>		<i>inclusa</i> Børgesen	A
<i>droebakense</i> Wille	A	<i>laetevirens</i> (Dillwyn) Kützing, 1843	A M
<i>dyeri</i> E.P. Wright, 1881	A	<i>lehmannaiana</i> (Lindenberg) Kützing, 1843	A M
Family Endosphaeraceae		<i>liebetrichii</i> Grunow, 1884	A M
<i>Chlorochytrium</i>		<i>liniformis</i> Kützing, 1849	A M
<i>cohnii</i> E.P. Wright, 1877	A M	<i>michaelsenii</i> O.C. Schmidt	A
<i>crassum</i> (Naccari) Rabenhorst	M	<i>nigrescens</i> Zanardini ex Frauenfeld	A M
<i>dermatocolax</i> Reinke, 1889	A	<i>oblitterata</i> J. Söderström	A
<i>facciolae</i> (Borzi) Bristol, 1920	A	<i>parriaudii</i> van den Hoek	A M
<i>schmitzii</i> Rosenvinge, 1893	A	<i>pellucida</i> (Hudson) Kützing, 1843	A M
<i>Gomontia</i>		<i>prolifera</i> (Roth) Kützing, 1843	A M
<i>polyrhiza</i> (Lagerheim) Bornet & Flahault, 1888	A M	<i>pygmaea</i> Reinke, 1888	A
Order Chlorocystidales		<i>retroflexa</i> (Bonnemaison ex P.L. Crouan & H.M. Crouan) van den Hoek, 1963	A M
Family Chlorocystidaceae		<i>ruchingeri</i> (C. Agardh) Kützing, 1845	A M
<i>Halochlorococcum</i>		<i>rupestris</i> (Linnaeus) Kützing, 1843	A M
<i>dilatum</i> Kornmann & Sahling	A	<i>sericea</i> (Hudson) Kützing, 1843	A M
<i>marinum</i> P.J.L. Dangeard	A	<i>sivashensis</i> C. Meyer	M
<i>moorei</i> (N.L. Gardner) Kornmann & Sahling, 1917	A M	<i>socialis</i> Kützing, 1849	A M
<i>operculatum</i> Kornmann & Sahling, 1983	A	<i>theotonii</i> O.C. Schmidt	A
Order Cladophorales		<i>vadorum</i> (Areschoug) Kützing, 1849	A M
Family Anadyomenaceae		<i>vagabunda</i> (Linnaeus) van den Hoek, 1963	A M
<i>Anadyomene</i>		<i>weizenbaurii</i> O.C. Schmidt	A
<i>saldanhae</i> A.B. Joly & E.C. Oliveira, 1969	A	Rhizoclonium	
		<i>africanum</i> Kützing, 1853	A
		<i>kochianum</i> Kützing, 1845	A M
		<i>pachydermum</i> Kjellman	A
		<i>riparium</i> (Roth) Kützing ex Harvey, 1849	A M
		<i>tortuosum</i> (Dillwyn) Kützing, 1845	A M

Family Witrockiellaceae		Family Boodleaceae	
<i>Witrockiella</i>		<i>Boodlea</i>	
<i>paradoxa</i> Wille, 1909	A	<i>struveoides</i> M. Howe, 1918	A
Order Codiales		<i>Struvea</i>	
Family Codiaceae		<i>anastomosans</i> (Harvey) Piccone & Grunow ex Piccone, 1884	A
<i>Codium</i>		<i>ramosa</i> Dickie, 1874	A
<i>adhaerens</i> C. Agardh	A M		
<i>bursa</i> (Linnaeus) C. Agardh, 1822	A M		
<i>carolinanum</i> Searles	A		
<i>cattaniae</i> Vouk	M		
<i>coralloides</i> (Kützing) P.C. Silva	M		
<i>decoratum</i> (Woodward) M. Howe, 1911	A M		
<i>effusum</i> (Rafinesque) Delle Chiaje	A M		
<i>elisabethae</i> O.C. Schmidt	A		
<i>fragile</i> (Suringar) Hariot, 1889	A M		
<i>guineense</i> P.C. Silva	A		
<i>intertextum</i> Collins & Hervey, 1917	A		
<i>lucasii</i> Setchell, 1935	A		
<i>repens</i> P.L. Crouan & H.M. Crouan ex Vickers, 1905	A		
<i>taylorii</i> P.C. Silva, 1960	A M		
<i>tomentosum</i> Stackhouse, 1797	A M		
<i>vermilara</i> (Oliv) Delle Chiaje	A M		
<i>Flabellia</i>			
<i>petiolata</i> (Turra) Nizamuddin	A M		
<i>Ostreobium</i>			
<i>quekettii</i> Bornet & Flahault, 1889	A M		
Order Dasycladales			
Family Dasycladaceae			
<i>Batophora</i>			
<i>occidentalis</i> (Harvey) S. Berger & Kever ex M.J. Wynne	A		
<i>oersledii</i> J. Agardh	A		
<i>Cymoplia</i>			
<i>barbata</i> (Linnaeus) J.V. Lamouroux	A		
<i>Dasycladus</i>			
<i>vermicularis</i> (Scopoli) Krasser, 1898	A M		
Family Polyphysaceae			
<i>Acetabularia</i>			
<i>acetabulum</i> (Linnaeus) P.C. Silva, 1952	A M		
<i>calyculus</i> J.V. Lamouroux, 1824	A M		
<i>parvula</i> Solms-Laubach, 1895	A M		
<i>polyphysoides</i> P.L. Crouan & H.M. Crouan	A		
Order Klebshormidiales			
Family Stichococcaceae			
<i>Stichococcus</i>			
<i>bacillaris</i> Nägeli, 1849	A M		
Order Microsporales			
Family Microsporaceae			
<i>Microspora</i>			
<i>ficulinae</i> P.A. Dangeard	A		
Order Phaeophilales			
Family Phaeophilaceae			
<i>Phaeophila</i>			
<i>dendroides</i> (P.L. Crouan & H.M. Crouan) Batters, 1902	A M		
<i>endophytum</i> (M. Möbius) R. Nielsen	A		
<i>hirsuta</i> (Ercegovic) R. Nielsen	M		
<i>tenuis</i> (Kylin) R. Nielsen	A		
Order Prasiolales			
Family Prasiolaceae			
<i>Prasiola</i>			
<i>calophylla</i> (Carmichael ex Greville) Kützing, 1845	A		
<i>crispa</i> (Lightfoot) Kützing, 1843	A M		
<i>fluvialis</i> (Sommerfelt) Areschoug	A		
<i>furfuracea</i> (Mertens ex Hornemann) Kützing, 1843	A		
<i>stipitata</i> Suhr ex Jessen, 1848	A M		
<i>Rosenvingiella</i>			
<i>constricta</i> (Setchell & N.L. Gardner) P.C. Silva	A		
<i>polyrhiza</i> (Rosenvinge) P.C. Silva, 1957	A M		
Order Siphonocladales			
Family Boodleaceae			
<i>Boodlea</i>			
<i>struveoides</i> M. Howe, 1918			
Family Siphonocladaceae			
<i>Cladophoropsis</i>			
<i>gerloffii</i> Nizamuddin, 1998			M
<i>macromeres</i> W.R. Taylor, 1928			A
<i>membranacea</i> (Hofman Bang ex C. Agardh)			
<i>Børgesen, 1905</i>			
<i>modonensis</i> (Kützing) Reinbold, 1905			M
<i>Siphonocladus</i>			
<i>pussillus</i> (C. Agardh ex Kützing) Hauck			M
<i>tropicus</i> (P.L. Crouan & H.M. Crouan) J. Agardh, 1887			A
Family Valoniaceae			
<i>Dictyosphaeria</i>			
<i>ocellata</i> (M. Howe) J.L. Olsen			A
<i>Ernadesmis</i>			
<i>verticillata</i> (Kützing) Børgesen, 1912			A M
<i>Valonia</i>			
<i>aegagropila</i> C. Agardh, 1823			A M
<i>caespitosa</i> P.L. Crouan & H.M. Crouan			M
<i>macrophysa</i> Kützing, 1843			A M
<i>utricularis</i> (Roth) C. Agardh, 1823			A M
<i>Ventricaria</i>			
<i>ventricosa</i> (J. Agardh) J.L. Olsen & J.A. West, 1988			M
Order Tetrasporales			
Family Palmellopsidaceae			
<i>Palmophyllum</i>			
<i>crassum</i> (Naccari) Rabenhorst			M
Order Ulotrichales			
<i>Aphanochaete</i>			
<i>repens</i> A. Braun			A
Family Borodinellaceae			
<i>Planophila</i>			
<i>microcystis</i> (P.J.L. Dangeard) Kornmann & Sahling			A M
Family Ulotrichaceae			
<i>Eugomontia</i>			
<i>sacculata</i> Kornmann, 1960			A M
<i>Ulothrix</i>			
<i>brunnihaleri</i> Schüssnig			M
<i>discifera</i> Kjellman			A
<i>flacca</i> (Dillwyn) Thuret, 1863			A M
<i>flexuosa</i> Schüssnig			M
<i>implexa</i> (Kützing) Kützing, 1849			A M
<i>longicauda</i> Schüssnig			M
<i>palusalsa</i> Lohhorst, 1978			A
<i>speciosa</i> (Carmichael) Kützing, 1849			A
<i>subflaccida</i> Wille, 1901			A M
<i>terriformis</i> Kützing			A
<i>tenuissima</i> Kützing, 1833			M
<i>zonata</i> (F. Weber & D. Mohr) Kützing, 1833			A M
Order Ulvales			
Family Capsosiphonaceae			
<i>Capsosiphon</i>			
<i>fulvescens</i> (C. Agardh) Setchell & N.L. Gardner, 1920			A M
Family Kornmanniaceae			
<i>Kornmannia</i>			
<i>leptoderma</i> (Kjellman) Bliding, 1969			A
Family Monostromataceae			
<i>Monostroma</i>			
<i>arcticum</i> Wittrock, 1866			A
<i>balticum</i> (Areschoug) Wittrock			A
<i>bullosum</i> (Roth) Thuret			M
<i>grevillei</i> (Thuret) Wittrock, 1866			A M
<i>lubricum</i> Kjellman			A
<i>obscurum</i> (Kützing) J. Agardh			A M

<i>oxyccum</i> (Kützing) Thuret	A	<i>flustrae</i> Reinke, 1889	A M
<i>oxyspermum</i> (Kützing) Doty, 1947	A	<i>halimedae</i> (Hansgirg) Reinke, 1889	M
<i>saccodemeum</i> Kjellman	A	<i>heterotricha</i> (Yarish) R. Nielsen	A M
<i>Protomonostroma</i>		<i>perforans</i> (Huber) R. Nielsen	A M
<i>undulatum</i> (Wittrock) K.L. Vinogradova, 1969	A	<i>phillipsii</i> (Batters) R. Nielsen	A
<i>Ulvaria</i>		<i>pontica</i> Rochlina	M
<i>fusca</i> Ruprecht, 1850	A	<i>testarum</i> (Kylin) R. Nielsen	A
Family Ulvaceae		Ochlochaete	
<i>Blidingia</i>		<i>hystrix</i> Thwaites ex Harvey, 1849	A M
<i>chadefaudii</i> (Feldmann) Bliding, 1963	A M	Protoderma	
<i>marginata</i> (J. Agardh) P.J.L. Dangeard ex Bliding, 1958	A M	<i>concharum</i> Hansgirg	M
<i>minima</i> (Nägeli ex Kützing) Kylin, 1947	A M	<i>Pseudodendroclonium</i>	
<i>ramifera</i> (Bliding) Garbary & Barkhouse	A M	<i>dynamenae</i> R. Nielsen	A
<i>subsalsa</i> (Kjellman) Kornmann & Sablina ex Scagel et al.	A M	<i>fucicola</i> (Rosenvinge) R. Nielsen, 1980	A
Enteromorpha		Order Volvocales	
<i>adriatica</i> Bliding	M	Family Volvocaceae	
<i>aragoensis</i> Bliding, 1960	M	<i>Chaetopeltis</i>	
<i>chaetomorphoides</i> Børgesen, 1911	A	<i>orbicularis</i> Berthold	A
<i>clathrata</i> (Roth) Greville, 1830	A M	PHYLUM HETEROKONTOPHYTA	
<i>compressa</i> (Linnaeus) Nees, 1820	A M	Class Xanthophyceae	
<i>fasciculata</i> P.J.L. Dangeard, 1949	A	Order Vaucherales	
<i>flexuosa</i> (Wulfen) J. Agardh, 1883	A M	Family Triboneemataceae	
<i>gelatinosae</i> Kützing	A	<i>Tribonema</i>	
<i>hendayensis</i> P.J.L. Dangeard & Parriaud	A M	<i>marinum</i> Feldmann	M
<i>intestinalis</i> (Linnaeus) Nees, 1820	A M	Family Vaucheriaceae	
<i>intestinaloides</i> Koeman & van den Hoek, 1982	A	<i>Vaucheria</i>	
<i>jugoslavica</i> Bliding	A M	<i>arcassoniensis</i> P.J.L. Dangeard, 1939	A
<i>kylinii</i> Bliding, 1948	A M	<i>bermudensis</i> W.R. Taylor & Bernatowicz	A
<i>lingulata</i> J. Agardb, 1883	A M	<i>compacta</i> (Collins) Collins ex W.R. Taylor, 1937	A
<i>linza</i> (Linnaeus) J. Agardh, 1883	A M	<i>coronata</i> Nordstedt, 1879	A M
<i>maeotica</i> Proshkina-Lavrenko	M	<i>dichotoma</i> (Linnaeus) Martius, 1817	A M
<i>multiramosa</i> Bliding	A M	<i>erythrospora</i> T.A. Christensen, 1956	A
<i>muscoidea</i> (Clemente) Cremades, 1990	A M	<i>intermedia</i> Nordstedt, 1879	A
<i>procera</i> Ahlner	M	<i>litorea</i> C. Agardh, 1823	A
<i>prolifera</i> (O.F. Müller) J. Agardh, 1883	A M	<i>medusa</i> T.A. Christensen, 1952	A
<i>pseudolinza</i> Koeman & van den Hoek, 1982	A M	<i>minuta</i> Blum & Conover, 1953	A
<i>radiata</i> J. Agardh, 1883	A M	<i>piloboloides</i> Thuret, 1954	A M
<i>ralfsii</i> Harvey, 1850	A M	<i>schleicheri</i> De Wildeman	A
<i>robertii-lamii</i> Parriaud, 1961	A	<i>sescupilaria</i> T.A. Christensen, 1952	A M
<i>simplex</i> (K.L. Vinogradova) Koeman & van den Hoek	A M	<i>subsimplex</i> P.L. Crouan & H.M. Crouan, 1878	A
<i>stipitata</i> P.J.L. Dangeard	A M	<i>synandra</i> Woronin, 1869	A
<i>torta</i> (Mertens) Reinbold, 1893	A M	<i>velutina</i> C. Agardh, 1824	A M
<i>Ppercursaria</i>		<i>vipera</i> Blum, 1960	A
<i>percursa</i> (C. Agardh) Rosenvinge, 1893	A M	PHYLUM PHAEOPHYCOTA	
<i>Ulva</i>		Class Phaeophyceae	
<i>bifrons</i> Ardré, 1967	A M	Order Chordariales	
<i>curvata</i> (Kützing) De Toni, 1889	A M	Family Acrotrichaceae	
<i>fasciata</i> Delile, 1813	A M	<i>Acrotrix</i>	
<i>gigantea</i> (Kützing) Bliding	A M	<i>gracilis</i> Kylin, 1907	A
<i>lactuca</i> Linnaeus, 1753	A M	Family Chordariaceae	
<i>linearis</i> P.J.L. Dangeard, 1957	A M	<i>Adriogloia</i>	
<i>neapolitana</i> Bliding	M	<i>adriatica</i> Ercegovic	M
<i>olivascens</i> P.A. Dangeard	A M	<i>Chordaria</i>	
<i>pseudocurvata</i> Koeman & van den Hoek, 1981	A	<i>flagelliformis</i> (O.F. Müller) C. Agardh, 1817	A
<i>rhacodes</i> (Holmes) Papenfuss, 1960	A	<i>Cladosiphon</i>	
<i>rigida</i> C. Agardh, 1823	A M	<i>contortus</i> (Thuret) Kylin, 1940	A M
<i>rotundata</i> Bliding	A M	<i>cylindricus</i> (Sauvageau) Kylin	M
<i>scandinavica</i> Bliding, 1968	A M	<i>irregularis</i> (Sauvageau) Kylin	M
<i>uvoides</i> Bory	A	<i>mediterraneus</i> Kützing	M
Family Ulvaceae		<i>occidentalis</i> Kylin	A
<i>Acrochaete</i>		<i>zosterae</i> (J. Agardh) Kylin, 1940	A M
<i>cladophorae</i> (Hornby) R. Nielsen, 1995	A	<i>Dalmatogloia</i>	
<i>geniculata</i> (N.L. Gardner) O'Kelly, 1982	A M	<i>bryozoi</i> Ercegovic, 1948	M
<i>heteroclada</i> J.A. Correa & R. Nielsen	A	<i>Eudesme</i>	
<i>inflata</i> (Ercegovic) Gallardo et al.	A M	<i>virescens</i> (Carmichael ex Harvey) J. Agardh, 1882	A M
<i>leptochaete</i> (Huber) R. Nielsen	A M	<i>Gontrania</i>	
<i>operculata</i> J.A. Correa & R. Nielsen	A	<i>lubrica</i> Sauvageau	M
<i>polymorpha</i> (L. Moewus) R. Nielsen	A	<i>Halonema</i>	
<i>repens</i> Pringsheim, 1863	A M	<i>subsimplex</i> Jaasund, 1951	A
<i>viridis</i> (Reinke) R. Nielsen, 1979	A M	<i>Liebmannia</i>	
<i>wittrockii</i> (Wille) R. Nielsen, 1983	A		
<i>Epicladia</i>			
<i>cyclostoma</i> (Aaern) R. Nielsen, 1995	A		

<i>leveillei</i> J. Agardh, 1842	A M	
<i>Mesogloia</i>		
<i>lanosa</i> P.L. Crouan & H.M. Crouan, 1867	A M	
<i>vermiculata</i> (J.E. Smith) S.F. Gray, 1821	A M	
<i>Myriocladia</i>		
<i>ekmani</i> (Areschoug) Kylin, 1907	A	
<i>loveni</i> J. Agardh, 1841	A	
<i>tomentosa</i> P.L. Crouan & H.M. Crouan, 1867	A	
<i>Myriogloea</i>		
<i>sciurus</i> (Harvey) Kuckuck ex Oltmanns, 1922	M	
<i>Papenfussiella</i>		
<i>callitrichia</i> (Rosenvinge) Kylin, 1940	A	
<i>kuromae</i> (Yendo) Inagaki, 1958	A	
<i>Saundersella</i>		
<i>simplex</i> (Saunders) Kylin, 1940	A	
<i>Sauvageaugloia</i>		
<i>griffithsiana</i> (A.W. Griffiths ex Harvey) G. Hamel ex Kylin, 1940	A M	
<i>Sphaerotrichia</i>		
<i>divaricata</i> (C. Agardh) Kylin, 1940	A M	
<i>Strepisithalia</i>		
<i>bifhamiana</i> (Batters) Batters, 1902	A	
<i>curvata</i> Sauvageau, 1896	A	
<i>liagorae</i> Sauvageau, 1896	A M	
<i>liebmanniae</i> Miranda, 1928	A M	
Family Elachistaceae		
<i>Elachista</i>		
<i>chondrii</i> Areschoug, 1875	A	
<i>flaccida</i> (Dillwyn) Fries, 1835	A M	
<i>fucicola</i> (Vellay) Areschoug, 1842	A M	
<i>globulosa</i> (C. Agardh) J. Agardh, 1848	A	
<i>intermedia</i> P.L. Crouan & H.M. Crouan, 1867	A M	
<i>jabukae</i> Ercegovic	M	
<i>neglecta</i> Kuckuck, 1929	M	
<i>scutulata</i> (J.E. Smith) Duby, 1830	A M	
<i>stellaris</i> Areschoug, 1842	A M	
<i>Halothrix</i>		
<i>lumbricalis</i> (Kützing) Reinke, 1888	A M	
<i>Leptonematella</i>		
<i>fasciculata</i> (Reinke) P.C. Silva, 1959	A M	
<i>neapolitana</i> (Schüssnig) Cormaci & G. Furnari	M	
Family Leathesiaceae		
<i>Corynophlaea</i>		
<i>crispa</i> (Harvey) Kuckuck, 1929	A	
<i>cystophorae</i> J. Agardh, 1882	A	
<i>flaccida</i> (C. Agardh) Kützing	M	
<i>hamelii</i> Feldmann, 1937	M	
<i>umbellata</i> (C. Agardh) Kützing	M	
<i>Cylindrocarpus</i>		
<i>microscopicus</i> P.L. Crouan & H.M. Crouan, 1851	A M	
<i>Leathesia</i>		
<i>diformis</i> (Linnaeus) J.E. Areschoug, 1847	A M	
<i>mucosa</i> Feldmann, 1937	M	
<i>verruculiformis</i> Y.P. Lee & I.K. Lee, 1988	A	
<i>Microcoryne</i>		
<i>ocellata</i> Strömfelt, 1888	A	
<i>Myriactula</i>		
<i>arabica</i> (Kützing) Feldmann, 1937	A M	
<i>areschougei</i> (P.L. Crouan & H.M. Crouan) G. Hamel, 1935	A	
<i>chordae</i> (Areschoug) Levring, 1937	A	
<i>clandestina</i> (P.L. Crouan & H.M. Crouan) Feldmann, 1943	A	
<i>elongata</i> (Sauvageau) G. Hamel	M	
<i>sucorum</i> (Rosenvinge) Kylin, 1947	A	
<i>gracilis</i> van der Ben	M	
<i>haydenii</i> (Gatty) Levring, 1973	A	
<i>microscopica</i> (Ercegovic) Ercegovic	M	
<i>rigida</i> (Sauvageau) G. Hamel	M	
<i>rivulariae</i> (Suhr) J. Feldmann, 1937	A M	
<i>stellulata</i> (Harvey) Levring, 1937	A M	
<i>vlastae</i> J. Feldmann, 1954	A	
<i>Petrospongium</i>		
<i>berkeleyi</i> (Greville) Nügeli ex Kützing, 1858	A M	
Family Myrionomataceae		
<i>Microspongium</i>		
<i>gelatinosum</i> Reinke, 1888	A M	
<i>globosum</i> Reinke, 1888	A	
<i>immersum</i> (Levrings) P.M. Pedersen, 1984	A	
<i>Microsporangium</i>		
<i>kuckuckianum</i> Schiffner	M	
<i>Myrionema</i>		
<i>balticum</i> (Reinke) Foslie, 1894	A M	
<i>conchicola</i> (Feldmann) Boudouresque	M	
<i>corunnae</i> Sauvageau, 1897	A	
<i>feldmannii</i> Loiseaux, 1964	A	
<i>furcatum</i> Jaasdund, 1951	A	
<i>liechtensternii</i> Hauck, 1877	A M	
<i>orbiculare</i> J. Agardh, 1848	M	
<i>papillosum</i> Sauvageau, 1897	A	
<i>seriatum</i> (Reinke) Kylin, 1947	A	
<i>strangulans</i> Carmichael ex Greville, 1827	A M	
<i>Protectocarpus</i>		
<i>speciosus</i> (Børgesen) Kuckuck ex Kornmann, 1955	A M	
<i>Ulonema</i>		
<i>rhizophorum</i> Foslie, 1894	A	
Family Ralfsiaceae		
<i>Hapalospongion</i>		
<i>macrocarpum</i> (Feldmann) León Álvarez & González, 1993	A M	
<i>Lithoderma</i>		
<i>fatiscens</i> Areschoug, 1875	A	
<i>Porterinema</i>		
<i>fluviatile</i> (A. Porter) Waern, 1952	A	
<i>Pseudolithoderma</i>		
<i>adriaticum</i> (Hauck) Verlaque	A M	
<i>extensem</i> (P.L. Crouan & H.M. Crouan) S. Lund, 1959	A M	
<i>roscoffense</i> Loiseaux	A M	
<i>rosenvingei</i> (Waern) S. Lund, 1959	A	
<i>subextensem</i> (Waern) S. Lund, 1959	A	
<i>Ralfsia</i>		
<i>bornetii</i> Kuckuck, 1894	A	
<i>disciformis</i> P.L. Crouan & H.M. Crouan, 1867	A M	
<i>expansa</i> (J. Agardh) J. Agardh, 1848	M	
<i>fungiformis</i> (Gunnerus) Setchell & N.L. Gardner, 1924	A	
<i>lucida</i> S. Lund, 1967	A	
<i>ovata</i> Rosenvinge, 1893	A	
<i>pusilla</i> (Strömfelt) Batters, 1892	A	
<i>verrucosa</i> (Areschoug) Areschoug, 1845	A M	
<i>Stragularia</i>		
<i>clavata</i> (Harvey) G. Hamel, 1931	A M	
<i>spongiocarpa</i> (Batters) G. Hamel	A	
Family Spermatophchnaceae		
<i>Halorhiza</i>		
<i>vaga</i> Kützing	A	
<i>Nemacystus</i>		
<i>erythraeus</i> (J. Agardh) Sauvageau, 1897	A	
<i>flexuosus</i> (C. Agardh) Kylin	A M	
<i>hispanicus</i> (Sauvageau) Kylin, 1940	A M	
<i>Spermatochnus</i>		
<i>paradoxus</i> (Roth) Kützing, 1843	A M	
<i>Stilophora</i>		
<i>nodulosa</i> (C. Agardh) P.C. Silva, 1996	A	
<i>tenella</i> (Esper) P.C. Silva, 1996	A M	
<i>Stilosiphis</i>		
<i>lejolisi</i> (Thuret) Kuckuck & Nienburg ex G. Hamel, 1937	A M	
Order Cutleriales		
Family Cutleriaceae		
<i>Cutteria</i>		
<i>adspersa</i> (Mertens ex Roth) De Notaris, 1842	A M	
<i>chilosa</i> (Falkenberg) P.C. Silva, 1957	A M	
<i>multifida</i> (J.E. Smith) Greville, 1830	A M	
<i>Zanardinia</i>		
<i>typus</i> (Nardo) G. Furnari, 1999	A M	

Order Desmarestiales			
Family Arthrocladiaceae			
<i>Arthrocladia</i>			A
<i>villosa</i> (Hudson) Duby, 1830	A M		
Family Desmarestiaceae			
<i>Desmarestia</i>			
<i>aculeata</i> (Linnaeus) J.V. Lamouroux, 1813	A M		
<i>dresnayi</i> J.V. Lamouroux ex Leman, 1819	A M		
<i>ligulata</i> (Lightfoot) J.V. Lamouroux, 1813	A M		
<i>viridis</i> (O.F. Müller) J.V. Lamouroux, 1813	A M		
Order Dictyosiphonales			
Family Asperococcaceae			
<i>Asperococcus</i>			
<i>bullosus</i> J.V. Lamouroux, 1813	A M		
<i>compressus</i> A.W. Griffiths ex W.J. Hooker, 1833	A M		
<i>fistulosus</i> (Hudson) W.J. Hooker, 1833	A M		
<i>scaber</i> Kuckuck, 1899	A M		
Family Buffhamiaceae			
<i>Buffhamia</i>			
<i>speciosa</i> Batters, 1895	A		
Family Coelodesmaceae			
<i>Coelodesme</i>			
<i>bulligera</i> Strömfelt, 1886	A		
Family Delamareaceae			
<i>Delamarea</i>			
<i>attenuata</i> (Kjellman) Rosenvinge, 1893	A		
Family Dictyosiphonaceae			
<i>Acrocytis</i>			
<i>groenlandicus</i> Rosenvinge, 1933	A		
<i>Dictyosiphon</i>			
<i>chordarius</i> Areschoug, 1847	A M		
<i>ekmani</i> Areschoug, 1875	A		
<i>filiformis</i> (Foslie) De Toni, 1895	A		
<i>foeniculaceus</i> (Hudson) Greville, 1830	A		
Family Giraudiaceae			
<i>Giraudia</i>			
<i>sphaelarioides</i> Derbès & Solier, 1851	A M		
Family Myriotrichiaceae			
<i>Desmotrichum</i>			
<i>punctarioides</i> A.D. Zinova & Dimitrova-Konaklieva	M		
<i>Leblondiella</i>			
<i>densa</i> (Batters) G. Hamel, 1939	A		
<i>Litosiphon</i>			
<i>laminariae</i> (Lyngbye) Harvey, 1849	A		
<i>tenuis</i> Levring, 1937	A		
<i>Myriotrichia</i>			
<i>adriatica</i> Hauck	M		
<i>canariensis</i> Kützing	A		
<i>clavaeformis</i> Harvey, 1833	A M		
<i>repens</i> Hauck, 1879	A		
<i>Protasperococcus</i>			
<i>myriotrichiiformis</i> Sauvageau	M		
Family Pogotrichiaceae			
<i>Omphalophyllum</i>			
<i>ulvaceum</i> Rosenvinge, 1893	A		
<i>Pogotrichium</i>			
<i>filiforme</i> Reinke, 1892	A		
<i>setiforme</i> (Rosenvinge) P.M. Pedersen, 1978	A		
Family Punctariaceae			
<i>Chilionema</i>			
<i>foecundum</i> (Strömfelt) R.L. Fletcher, 1987	A		
<i>hispanicum</i> (Sauvageau) R.L. Fletcher, 1987	A M		
<i>ocellatum</i> (Kützing) Kornmann, 1953	A		
<i>Clathrodiscus</i>			
<i>mandouli</i> (Sauvageau) G. Hamel, 1935	A		
<i>Hecatonema</i>			
<i>liagorae</i> (Feldmann) G. Hamel	M		
<i>terminale</i> (Kützing) Kylin, 1937	M		
<i>Punctaria</i>			
<i>crispata</i> (Kützing) Batters, 1900	A		
<i>crouanii</i> (Thuret) Bornet	A		
<i>glacialis</i> Rosenvinge, 1910			A
<i>latifolia</i> Greville, 1830			A M
<i>plantaginea</i> (Roth) Greville, 1830			A M
<i>tenuissima</i> (C. Agardh) Greville, 1830			A M
<i>Trachynema</i>			
<i>mortensenii</i> (Lund) Pedersen, 1985			A
Family Striariaceae			
<i>Coelocladia</i>			
<i>arctica</i> Rosenvinge, 1893			A
<i>Dermatocelis</i>			
<i>laminariae</i> Rosenvinge, 1898			A
<i>Foslia</i>			
<i>curta</i> (Foslie) Reinke, 1891			A
<i>Isthmoplea</i>			
<i>sphaerophora</i> (Carmichael ex Harvey) Kjellman, 1879			A M
<i>Stictyosiphon</i>			
<i>adriaticus</i> Kützing, 1843			A M
<i>arcticus</i> Pedersen & Kristiansen, 1992			A
<i>griffithianus</i> (Le Jolis) Holmes & Batters, 1891			A
<i>laxus</i> (J. Agardh) Athanasiadis, 1996			A
<i>lofotensis</i> (Foslie) Jaasund, 1965			A
<i>soriferus</i> (Reinke) Rosenvinge, 1935			A M
<i>tortilis</i> (Ruprecht) Reinke, 1889			A M
<i>Striaria</i>			
<i>attenuata</i> (C. Agardh) Greville, 1828			A M
Order Dictyotales			
Family Dictyotaceae			
<i>Dictyopteris</i>			
<i>delicatula</i> J.V. Lamouroux, 1809			A
<i>divaricata</i> (Okamura) Okamura, 1932			M
<i>plagiogramma</i> (Montagne) Vickers, 1905			A
<i>polypodioides</i> (De Candolle) J.V. Lamouroux, 1809			A M
<i>tripolitana</i> Nizamuddin, 1981			M
<i>Dictyota</i>			
<i>adnata</i> Zanardini			A
<i>bartayresiana</i> J.V. Lamouroux, 1809			A
<i>cervicornis</i> Kützing, 1859			A
<i>ciliolata</i> Sonder ex Kützing, 1859			A
<i>crenulata</i> J. Agardh, 1848			A
<i>dichotoma</i> (Hudson) J.V. Lamouroux, 1809			A M
<i>divaricata</i> J.V. Lamouroux, 1809			A
<i>fasciola</i> (Roth) J.V. Lamouroux, 1809			A M
<i>humifusa</i> Hörning, Schnetter & Coppejans, 1992			A
<i>implexa</i> J.V. Lamouroux			A
<i>kohlmeyeri</i> (Nizamuddin & Gerloff) Hörning, Schnetter & Coppejans			A
<i>liturata</i> J. Agardh, 1848			A
<i>mediterranea</i> (Schiffner) G. Furnari, 1987			M
<i>menstrualis</i> (Hoyt) Schnetter, Hörning & Weber-Peukert, 1987			A
<i>mertensii</i> (Martius) Kützing, 1859			A
<i>naevosa</i> (Suhr) J. Agardh, 1848			A
<i>pinnatifida</i> Kützing, 1859			A
<i>spiralis</i> Montagne, 1846			A M
<i>volubilis</i> Kützing, 1849			A
<i>Lobophora</i>			
<i>variegata</i> (J.V. Lamouroux) Womersley ex E.C. Oliveira, 1977			A M
<i>Padina</i>			
<i>boergesenii</i> Allender & Kraft, 1983			M
<i>boryana</i> Thivy, 1966			M
<i>gymnospora</i> (Kützing) Sonder, 1871			A
<i>pavonica</i> (Linnaeus) Thivy, 1960			A M
<i>tenuis</i> Bory, 1827			M
<i>Padinopsis</i>			
<i>adriatica</i> Ercegovic			M
<i>Spatoglossum</i>			
<i>schroederi</i> (C. Agardh) Kützing, 1859			A M
<i>soli</i> (Chauvin ex Montagne) Kützing, 1843			A M
<i>variabile</i> Figari & De Notaris, 1853			M
<i>Stylopodium</i>			
<i>schimperi</i> (Buchinger ex Kützing) Verlaque & Boudouresque			M

<i>zonale</i> (J.V. Lamouroux) Papenfuss, 1940	A	<i>holmesii</i> (Batters) Kornmann, 1956	A
<i>Taonia</i>		<i>Laminariocolax</i>	
<i>atomaria</i> (Woodward) J. Agardh, 1848	A M	<i>tomentosoides</i> (Farlow) Kylin, 1947	A
<i>lacheana</i> Cormaci, G. Furnari and Pizzuto, 1994	M	<i>Laminarinema</i>	
<i>Zonaria</i>		<i>elisabethiae</i> Kawai & Tokuyama, 1995	A
<i>inarticulata</i> (J.V. Lamouroux) Papenfuss	A	<i>Mikrosyphar</i>	
<i>subarticulata</i> (J.V. Lamouroux) Papenfuss, 1944	A	<i>polysiphoniae</i> Kuckuck, 1897	A
<i>tournefortii</i> (J.V. Lamouroux) Montagne, 1846	A M	<i>porphyrae</i> Kuckuck, 1897	A
Order Ectocarpales		<i>sphaelariae</i> Levring, 1974	A
Family Ectocarpaceae		<i>zosterae</i> Kuckuck, 1895	A
<i>Acinetospora</i>		<i>Phaeostroma</i>	
<i>crinita</i> (Carmichael ex Harvey) Kornmann, 1953	A M	<i>bertholdii</i> Kuckuck	M
<i>Acrospongium</i>		<i>endophyticum</i> S. Lund, 1959	A
<i>ralfsioides</i> Schiffner	M	<i>parasiticum</i> Børgesen, 1902	A
<i>Asteronema</i>		<i>pustulosum</i> Kuckuck, 1893	A
<i>rhodochortonoides</i> (Børgesen) D.G. Müller & E. Parodi, 1994	A M	<i>Pilinia</i>	
<i>Climacosorus</i>		<i>minor</i> (Hansgrig) Kützing	A M
<i>mediterraneus</i> Sauvageau, 1933	A M	<i>rimosa</i> Kützing, 1843	A M
<i>Dichosporangium</i>		<i>Pleurocladia</i>	
<i>chordariae</i> Wollny	A	<i>lacustris</i> A. Braun, 1855	A
<i>Ectocarpus</i>		<i>lucifuga</i> (Kuckuck) Wilce, 1966	A
<i>borealis</i> (Kjellman) Kjellman, 1890	A	<i>Spongonema</i>	
<i>caspicus</i> Henckel	M	<i>tomentosum</i> (Hudson) Kützing, 1849	A M
<i>fasciculatus</i> Harvey, 1849	A M	<i>Streblonema</i>	
<i>flagelliformis</i> Kützing, 1843	M	<i>breve</i> (Sauvageau) De Toni, 1895	A
<i>fulvescens</i> Schousboe ex Thuret	A	<i>deformans</i> (P.A. Dangeard) G. Hamel, 1931	A
<i>parvulus</i> Kützing, 1849	M	<i>fasciculatum</i> Thuret, 1863	A
<i>parvus</i> (Saunders) Hollenberg, 1971	M	<i>helophorum</i> (Rosenvinge) Batters, 1902	A
<i>rufulus</i> Kützing, 1843	M	<i>inclusum</i> Kylin, 1910	A
<i>siliculosus</i> (Dillwyn) Lyngbye, 1819	A M	<i>infestans</i> (H. Gran) Batters, 1902	A M
<i>similiusculus</i> C. Agardh, 1827	M	<i>integratum</i> P.J.L. Dangeard, 1970	A
<i>tomentosoides</i>	A	<i>intestinum</i> (Reinsch) Batters, 1892	A
<i>variabilis</i> Vickers, 1905	A	<i>oligosporum</i> Strömfelt, 1884	A M
<i>virescens</i> Thuret ex Sauvageau	A M	<i>parasiticum</i> (Sauvageau) Levring, 1937	A M
<i>Feldmannia</i>		<i>stilophorae</i> (P.L. Crouan & H.M. Crouan) Kylin, 1908	AM
<i>battersiidea</i> (Ercegovic) Cormaci & G. Furnari	M	<i>tenuissimum</i> Hauck, 1884	A M
<i>irregularis</i> (Kützing) G. Hamel, 1939	A M	<i>tuberculi</i> Jaasdun, 1965	A
<i>kjellmanii</i> Kylin, 1947	A	<i>turnerellae</i> (Foslie) Printz, 1926	A
<i>lebelii</i> (Areschoug ex P.L. Crouan & H.M. Crouan) G. Hamel, 1939	A M	<i>zanardinii</i> (P.L. Crouan & H.M. Crouan) De Toni, 1895	A
<i>Gononema</i>		<i>Streblonemopsis</i>	
<i>alariae</i> P.M. Pedersen, 1981	A	<i>irritans</i> Valiante, 1883	M
<i>Herponema</i>		<i>Zosterocarpus</i>	
<i>desmarestiae</i> (Gran) Cardinal, 1964	A	<i>oedogonium</i> (Meneghini) Bornet	M
<i>graniferum</i> Kuckuck	M	Family Lithodermataceae	
<i>minutum</i> Levring, 1974	A	<i>Jonsonia</i>	
<i>solitarium</i> (Sauvageau) G. Hamel, 1939	A M	<i>pulvinata</i> S. Lund, 1959	A
<i>valiantei</i> (Bornet ex Sauvageau) G. Hamel, 1939	A M	<i>Petroderma</i>	
<i>velutinum</i> (Greville) J. Agardh, 1890	A M	<i>maculiforme</i> (Wollny) Kuckuck, 1897	A
<i>Hincksia</i>		<i>Sorapion</i>	
<i>dalmatica</i> (Ercegovic) Cormaci & G. Furnari	M	<i>kjellmanii</i> (Wille) Rosenvinge, 1898	A
<i>fenestrata</i> (Berkeley ex Harvey) P.C. Silva, 1987	A	<i>simulans</i> Kuckuck, 1894	A
<i>fuscata</i> (Zanardini) P.C. Silva, 1987	A M	Family Pilayellaceae	
<i>geniculata</i> (Ercegovic) Cormaci & G. Furnari	M	<i>Bachelotia</i>	
<i>granulosa</i> (J.E. Smith) P.C. Silva, 1987	A M	<i>antillarum</i> (Grunow) Gerloff, 1959	A
<i>hauckii</i> (Ercegovic) Cormaci & G. Furnari	M	<i>Pilayella</i>	
<i>hincksiae</i> (Harvey) P.C. Silva, 1987	A M	<i>littoralis</i> (Linnaeus) Kjellman, 1872	A M
<i>mitchelliae</i> (Harvey) P.C. Silva, 1987	A M	<i>varia</i> Kjellman, 1883	A
<i>onslowensis</i> (Amsler & Kapraun) P.C. Silva, 1987	A	Family Sorocarpaceae	
<i>ovata</i> (Kjellman) P.C. Silva, 1987	A M	<i>Botrytella</i>	
<i>rallisiae</i> (Vickers) P.C. Silva, 1987	A	<i>micromora</i> Bory, 1822	A
<i>sandriana</i> (Zanardini) P.C. Silva, 1987	A M	<i>reinboldii</i> (Reinke) Kornmann & Sahling, 1988	A
<i>sandriana</i>	A	Order Fucales	
<i>secunda</i> (Kützing) P.C. Silva, 1987	A M	Family Cystoseiraceae	
<i>Kuckuckia</i>		<i>Bifurcaria</i>	
<i>kylinii</i> Cardinal, 1964	A M	<i>bifurcata</i> R. Ross, 1958	A
<i>spinosa</i> (Kützing) Kornmann, 1958	A M	<i>Cystoseira</i>	
<i>Kuetzingiella</i>		<i>abies-marina</i> (S.G. Gmelin) C. Agardh	A M
<i>battersii</i> (Bornet) Kornmann, 1956	A M	<i>algeriensis</i> Feldmann	M
		<i>amentacea</i> Bory	M
		<i>baccata</i> (S.G. Gmelin) P.C. Silva, 1952	A M
		<i>barbata</i> (Stackhouse) C. Agardh, 1842	A M
		<i>barbatula</i> Kützing	M
		<i>brachycarpa</i> J. Agardh	M

<i>compressa</i> (Esper) Gerloff & Nizamuddin, 1975	A M	<i>pinnatifida</i> (Harvey) Suringar, 1873	A M
<i>concatenata</i> (Linnaeus) C. Agardh	A M		
<i>corniculata</i> (Turner) Zanardini, 1841	M		
<i>crinita</i> Duby	A M	Family Chordaceae	
<i>crinitophylla</i> Ercegovic	M	<i>Chorda</i>	
<i>dubia</i> Valiante	M	<i>filum</i> (Linnaeus) Stackhouse, 1797	A M
<i>elegans</i> Sauvageau	M	<i>Haloisiphon</i>	
<i>foeniculacea</i> (Linnaeus) Greville, 1830	A M	<i>tomentosus</i> (Lyngbye) Jaasund, 1957	A
<i>gibraltarica</i> (Sauvageau) P.J.L. Dangeard, 1949	A		
<i>humilis</i> Schousboe ex Kützing, 1860	A M	Family Laminariaceae	
<i>hyblaea</i> Giaccone	M	<i>Laminaria</i>	
<i>jabukae</i> Ercegovic	M	<i>agardhii</i> Kjellman, 1877	A
<i>mauritanica</i> Sauvageau	A M	<i>digitata</i> (Hudson) J.V. Lamouroux, 1813	A
<i>mediterranea</i> Sauvageau, 1912	A M	<i>hyperborea</i> (Gunnerus) Foslie, 1884	A
<i>montagnei</i> J. Agardh	M	<i>japonica</i> Areschoug, 1851	M
<i>nodiculis</i> (Withering) M. Roberts, 1967	A M	<i>nigripes</i> J. Agardh, 1868	A
<i>pelagoae</i> Ercegovic	M	<i>ochroleuca</i> De La Pylaie, 1824	A M
<i>planiramea</i> Schiffner	M	<i>rodriguezii</i> Bornet	M
<i>platyclada</i> Sauvageau, 1912	A M	<i>saccarina</i> (Linnaeus) J.V. Lamouroux, 1813	A
<i>sauvageana</i> G. Hamel, 1939	M	<i>solidungula</i> J. Agardh, 1868	A
<i>schiffneri</i> G. Hamel, 1939	M		
<i>sedoides</i> (Desfontaines) C. Agardh, 1820	M	Family Phyllophoraceae	
<i>spinosa</i> Sauvageau, 1912	A M	<i>Phyllophora</i>	
<i>squarrosa</i> De Notaris	M	<i>brevipes</i> (C. Agardh) E.C. Henry & South, 1987	M
<i>susanensis</i> Nizamuddin, 1985	M	<i>purascens</i> (C. Agardh) E.C. Henry & South, 1987	A M
<i>tamariscifolia</i> (Hudson) Papenfuss, 1950	A M	Saccorhiza	
<i>usneoides</i> (Linnaeus) M. Roberts, 1968	A M	<i>dermatodea</i> (De La Pylaie) J. Agardh, 1868	A
<i>wildprettii</i> Nizamuddin, 1995	A	<i>polyschides</i> (Lightfoot) Batters, 1902	A M
<i>zosteroides</i> C. Agardh	M		
<i>Halidrys</i>		Order Scytoniphonales	
<i>siliquosa</i> (Linnaeus) Lyngbye, 1819	A	Family Lithodermataceae	
Family Fucaceae		<i>Sympylocarpus</i>	
<i>Ascophyllum</i>		<i>strangulans</i> Rosenvinge, 1893	A M
<i>nodosum</i> (Linnaeus) Le Jolis, 1863			
<i>Fucus</i>		Family Scytoniphonaceae	
<i>ceranoides</i> Linnaeus, 1753	A	<i>Colpomenia</i>	
<i>cottoni</i> M.J. Wynne & Magne, 1991	A	<i>peregrina</i> (Sauvageau) Hamel, 1937	A M
<i>distichus</i> Linnaeus, 1767	A	<i>sinuosa</i> (Mertens ex Roth) Derbès & Solier, 1851	A M
<i>evanescens</i> C. Agardh, 1820	A	Compsoneura	
<i>serratus</i> Linnaeus, 1753	A	<i>gracile</i> Kuckuck, 1899	M
<i>spiralis</i> Linnaeus, 1753	A M	<i>microspodium</i> (Batters) Kornmann, 1953	A
<i>vesiculosus</i> Linnaeus, 1753	A M	<i>minutum</i> (C. Agardh) Kornmann, 1953	A M
<i>virsoidea</i> J. Agardh	M	<i>saxicolum</i> (Kuckuck) Kornmann, 1953	A
<i>Pelvetia</i>		Endorachne	
<i>canaliculata</i> (Linnaeus) Decaisne & Thuret, 1845	A	<i>binghamiae</i> J. Agardh, 1896	A
Family Himanthaliaceae		Hydroclathrus	
<i>Himanthalia</i>		<i>clathratus</i> (Bory ex C. Agardh) M. Howe, 1920	A M
<i>elongata</i> (Linnaeus) S.F. Gray, 1821	A		
Family Sargassaceae		Nemodictya	
<i>Sargassum</i>		<i>tingitanum</i> Schousboe ex Bornet	A M
<i>acinarium</i> (Linnaeus) Setchell, 1933	A M	Petalonia	
<i>cymosum</i> C. Agardh, 1820	A	<i>fascia</i> (O.F. Müller) Kuntze, 1898	A M
<i>desfontainesii</i> (Turner) J. Agardh	A M	<i>filiformis</i> (Batters) Kuntze	A
<i>filipendula</i> C. Agardh, 1824	A	<i>zosterifolia</i> (Reinke) Kuntze, 1898	A M
<i>flavifolium</i> Kützing, 1849	A M	Rosenvingea	
<i>furcatum</i> Kützing, 1843	A	<i>intricata</i> (J. Agardh) Børgesen, 1914	A M
<i>hornschtchii</i> C. Agardh	M	<i>sanctae-crucis</i> Børgesen, 1914	A
<i>muticum</i> (Yendo) Fencholt, 1955	A M	Scytoniphon	
<i>natans</i> (Linnaeus) Gaillon, 1828	A	<i>complanatus</i> (Rosenvinge) Doty, 1947	A
<i>polycystum</i> C. Agardh, 1824	A	<i>dotyi</i> M.J. Wynne, 1969	A M
<i>ramentaceum</i> Zarmouth & Nizamuddin, 1991	M	<i>lomentaria</i> (Lyngbye) Link, 1833	A M
<i>thunbergii</i> (Mertens ex Roth) Kuntze, 1893	A		
<i>trichocarpum</i> J. Agardh	M	Order Sphaeriales	
<i>vulgare</i> C. Agardh, 1820	A M	Family Choristocarpaceae	
Order Laminariales		<i>Discosporangium</i>	
Family Alariaceae		<i>mesarthrocarpum</i> (Meneghini) Hauck, 1885	A M
<i>Alaria</i>			
<i>esculenta</i> (Linnaeus) Greville, 1830	A	Family Sphaerulariaceae	
<i>grandifolia</i> J. Agardh, 1872	A	<i>Choristocarpus</i>	
<i>pylai</i> (Bory) Greville, 1830	A	<i>tenellus</i> (Kützing) Zanardini, 1860	A M
<i>Ecklonia</i>		<i>Cladostephus</i>	
<i>biruncinata</i> (Bory) Papenfuss	A	<i>spongiosus</i> (Hudson) C. Agardh, 1817	A M
<i>muratii</i> Feldmann	A		
<i>Undaria</i>		<i>Sphaeraria</i>	
		<i>arctica</i> Harvey, 1858	A
		<i>brachygonia</i> Montagne, 1843	A
		<i>caespitula</i> Lyngbye, 1819	A
		<i>cirrosa</i> (Roth) C. Agardh, 1824	A M
		<i>fusca</i> (Hudson) S.F. Gray, 1821	A M
		<i>mirabilis</i> (Reinke ex Batters) Prud'homme van Reine, 1982	A
		<i>nana</i> Nägeli ex Kützing, 1855	A M

<i>plumigera</i> Holmes, 1883	A	<i>gracile</i> Børgesen, 1915	A M
<i>plumosa</i> Lyngbye, 1819	A	<i>gynandrum</i> (Rosenvinge) G. Hamel, 1927	A
<i>plumula</i> Zanardini, 1865	A M	<i>hallanicum</i> (Kylin) G. Hamel, 1927	A M
<i>racemosa</i> Greville, 1824	A	<i>hauckii</i> Schiffner, 1916	M
<i>radicans</i> (Dillwyn) C. Agardh, 1824	A M	<i>immersum</i> (Rosenvinge) G. Hamel, 1927	A
<i>reticulata</i> Lyngbye, 1818	A	<i>inclusum</i> (Levring) Papenfuss, 1945	A
<i>rigidula</i> Kützing, 1843	A M	<i>incrassatum</i> Ercegovic	M
<i>solitaria</i> Kylin, 1947	A	<i>infestans</i> M. Howe & Hoyt	A
<i>tribuloides</i> Meneghini, 1840	A M	<i>kylini</i> G. Hamel, 1927	A
<i>Sphaecella</i>		<i>lenormandii</i> (Suhr ex Kützing) Nägeli	M
<i>subtilissima</i> Reinke, 1890	A M	<i>leptonema</i> (Rosenvinge) Børgesen, 1915	A M
Family Stylocaulaceae		<i>liagorae</i> Børgesen, 1915	A
<i>Halopteris</i>		<i>lorrain-smithiae</i> (Lyle) L. Newton, 1931	A
<i>silicina</i> (Grateloup) Kützing, 1843	A M	<i>macula</i> (Rosenvinge) G. Hamel, 1927	A
<i>Stylocaulon</i>		<i>mediterraneum</i> (Levring) Boudouresque, 1970	M
<i>scoparium</i> (Linnaeus) Kützing, 1843	A M	<i>microfilum</i> Jao, 1936	A
Order Sporochiales		<i>microscopicum</i> (Nägeli ex Kützing) Nägeli, 1862	A M
Family Sporochnaceae		<i>minutum</i> (Suhr) G. Hamel, 1927	A
<i>Carpomitra</i>		<i>mirabilis</i> (Suhr) Nägeli, 1962	A
<i>costata</i> (Stackhouse) Batters, 1902	A M	<i>molinieri</i> Coppejans & Boudouresque, 1976	M
<i>Nereia</i>		<i>moniliforme</i> (Rosenvinge) Børgesen, 1915	A M
<i>filiformis</i> (J. Agardh) Zanardini	A M	<i>naumannii</i> (Askenasy) Askenasy	A
<i>tropica</i> (W.R. Taylor) W.R. Taylor	A	<i>nemalii</i> (De Notaris ex L. Dufour) Bornet	A M
<i>Sporochnus</i>		<i>nemalionis</i> (De Notaris ex L. Dufour) Bornet, 1904	M
<i>bolleanus</i> Montagne	A	<i>nitophylli</i> (Levring) Papenfuss, 1945	A
<i>gaertneri</i> (S.G. Gmelin) C. Agardh, 1820	A M	<i>occidentale</i> Børgesen, 1915	A
<i>pedunculatus</i> (Hudson) C. Agardh, 1817	A M	<i>pallens</i> (Zanardini) Nägeli, 1862	M
Order Syringodermatales		<i>parvulum</i> (Kylin) Hoyt, 1920	A M
Family Syringodermataceae		<i>pulchellum</i> Børgesen	A
<i>Syringoderma</i>		<i>radiatum</i> Jao, 1936	M
<i>floridana</i> E.C. Henry, 1984	A	<i>reductum</i> (Rosenvinge) G. Hamel, 1927	A M
Order Tilopteridales		<i>robustum</i> Børgesen, 1915	A
Family Tilopteridaceae		<i>rosulatum</i> (Rosenvinge) Papenfuss	A
<i>Haplospora</i>		<i>sanctae-mariae</i> (Darbshire) G. Hamel	A
<i>globosa</i> Kjellman, 1872	A	<i>savianum</i> (Meneghini) Nägeli, 1862	A M
<i>Tilopteris</i>		<i>secundatum</i> (Lyngbye) Nägeli, 1862	A M
<i>mertensi</i> (Turner) Kützing, 1849	A	<i>seiriolanum</i> (Gibson) Rosenvinge, 1924	A
PHYLUM RHODOPHYCOTA			
Class Rhodophyceae		<i>sparsum</i> (Carmichael ex Harvey) Nägeli, 1862	A
Order Acrochaetales		<i>stilophorae</i> (Levring) Kylin, 1944	A
Family Acrochaetiaceae		<i>subpinnatum</i> Bornet	M
<i>Acrochaetium</i>		<i>subseriatum</i> Børgesen	M
<i>alariae</i> (Jónsson) Bornet, 1904	A	<i>subtilissimum</i> (Kützing) G. Hamel, 1927	A M
<i>asparagopsis</i> (Chemin) Papenfuss, 1945	A	<i>trifilum</i> (Buffham) Batters	A M
<i>balticum</i> (Rosenvinge) Aleem & Schulz, 1952	A	<i>unifilum</i> Jao, 1936	A
<i>barbadensis</i> (Vickers) Børgesen, 1915	A	<i>virgatum</i> (Harvey) Bornet, 1904	A M
<i>battersianum</i> G. Hamel, 1927	A		
<i>boergesenii</i> Schiffner, 1931	M	<i>Audouinella</i>	
<i>bonnemaisoniae</i> (Batters) J. Feldmann & G. Feldmann, 1939	A M	<i>boryana</i> Abdel-Rahman & Magne, 1984	A
<i>brebneri</i> (Batters) G. Hamel, 1828	A	<i>conchicola</i> (Lami) South & Tittley, 1986	A
<i>caespitosum</i> (J. Agardh) Nägeli, 1862	A M	<i>Rhodochorton</i>	
<i>canariense</i> Børgesen, 1927	A	<i>atlanticum</i> Miranda, 1936	A
<i>catenulatum</i> M. Howe, 1914	A	<i>concrescens</i> K.M. Drew, 1928	A
<i>cheminii</i> Feldmann, 1954	A	<i>membranaceum</i> (Magnus) Hauck, 1883	A M
<i>chylocladiae</i> (Batters) Batters, 1902	A	<i>purpureum</i> (Lightfoot) Rosenvinge, 1900	A M
<i>codicolum</i> Børgesen, 1927	A M	<i>velutinum</i> (Hauck) G. Hamel, 1927	A M
<i>codiij</i> (P.L. Crouan & H.M. Crouan) G. Hamel, 1927	A M	<i>Schmitziella</i>	
<i>collopodum</i> (Rosenvinge) G. Hamel, 1927	A	<i>endophloea</i> Bornet & Batters, 1892	A M
<i>corymbiferum</i> (Thuret) Batters	A M	<i>Order Ahnfeltiales</i>	
<i>crassisipes</i> (Børgesen) Børgesen	A M	Family Ahnfeltiaceae	
<i>cymopoliae</i> Børgesen, 1927	A	<i>Ahnfeltia</i>	
<i>cystophagum</i> (Rosenvinge) G. Hamel, 1927	A	<i>plicata</i> (Hudson) Fries, 1836	A
<i>daviesii</i> (Dillwyn) Nägeli, 1862	A M	Order Bangiales	
<i>duboscqii</i> Feldmann, 1939	M	Family Bangiaceae	
<i>dumontiae</i> (Rosenvinge) G. Hamel, 1927	A	<i>Bangia</i>	
<i>efflorescens</i> (J. Agardh) Nägeli	A M	<i>atropurpurea</i> (Roth) C. Agardh, 1824	A M
<i>emergens</i> (Rosenvinge) Weber-van Bosse, 1921	A	<i>Porphyra</i>	
<i>endophyticum</i> Batters, 1896	A	<i>abyssicola</i> Kjellman, 1883	A
<i>endozoicum</i> (Darbshire) Batters	A	<i>amethystea</i> Kützing, 1843	A
<i>extensum</i> Ercegovic	M	<i>amplissima</i> (Kjellman) Setchell & Hus, 1900	A
		<i>carolinensis</i> Coll & J. Cox, 1977	A
		<i>dioica</i> Brodie & L.M. Irvine, 1997	A
		<i>drachii</i> J. Feldmann, 1978	A
		<i>heleneae</i> A.D. Zinova, 1948	A
		<i>insolita</i> Kornmann & Sahlung, 1991	A
		<i>leucosticta</i> Thuret, 1863	A M

<i>linearis</i> Greville, 1830	A M	<i>1851</i>	A
<i>miniata</i> (C. Agardh) C. Agardh, 1824	A	<i>granulatum</i> (Ducluzeau) C. Agardh, 1828	A M
<i>ochotensis</i> Nagai, 1941	A	<i>tetragonum</i> (Withering) S.F. Gray, 1821	A M
<i>purpurea</i> (Roth) C. Agardh, 1824	A M	<i>tetricum</i> (Dillwyn) S.F. Gray, 1821	A
<i>tenuissima</i> (Strömfelt) Setchell & Hus, 1900	A	<i>Centroceras</i>	
<i>thulae</i> Munda & P.M. Pedersen, 1978	A	<i>clavulatum</i> (C. Agardh) Montagne, 1846	A M
<i>umbilicalis</i> (Linnaeus) Kützing, 1843	A M	<i>Centroceroocolax</i>	
Order Bonnemaisoniales		<i>ubatubensis</i> A.B. Joly	A
Family Bonnemaisoniaceae		<i>Ceramium</i>	
<i>Asparagopsis</i>		<i>abyssale</i> Petersen, 1924	A
<i>armata</i> Harvey, 1855	A M	<i>arborescens</i> J. Agardh, 1894	A
<i>taxiformis</i> (Delile) Trevisan, 1845	A M	<i>arcticum</i> J. Agardh, 1894	A
<i>Bonnemaisonia</i>		<i>areschougi</i> Kylin, 1907	A M
<i>asparagooides</i> (Woodward) C. Agardh, 1822	A M	<i>atlanticum</i> Petersen, 1911	A
<i>clavata</i> G. Hamel, 1930	A M	<i>bertholdii</i> Funk, 1922	A M
<i>hamifera</i> Hariot, 1891	A M	<i>boergesenii</i> Petersen, 1911	A
Order Ceramiales		<i>botryocarpum</i> A.W. Griffiths ex Harvey, 1848	A
Family Ceramiaceae		<i>callipterum</i> Feldmann-Mazoyer, 1938	A M
<i>Acrothamnion</i>		<i>ciliatum</i> (J. Ellis) Ducluzeau, 1806	A M
<i>preissii</i> (Sonder) E.M. Wollaston, 1967	M	<i>cimbricum</i> H.E. Petersen, 1924	A M
<i>Aglaothamnion</i>		<i>cingulatum</i> Weber-van Bosse, 1923	A M
<i>bipinnatum</i> (P.L. Crouan & H.M. Crouan) Feldmann- Mazoyer, 1941	A M	<i>circinatum</i> (Kützing) J. Agardh, 1851	A M
<i>caudatum</i> (J. Agardh) Feldmann-Mazoyer, 1941	M	<i>codii</i> (H. Richards) Feldmann-Mazoyer, 1938	A M
<i>cordatum</i> (Børgesen) Feldmann-Mazoyer, 1941	A M	<i>comptum</i> Børgesen, 1924	A M
<i>diaphanum</i> L'Hardy-Halos & Maggs, 1991	A	<i>crouanianum</i> J. Agardh	A
<i>fasciculatum</i> (Harvey) Maggs & L'Hardy-Halos, 1993A		<i>danicum</i> Petersen, 1924	A
<i>feldmanniae</i> Halos, 1965	A M	<i>derbesii</i> Solier ex Kützing	A M
<i>gallicum</i> (Nägeli) Halos ex Ardré, 1970	A M	<i>deslongchampii</i> Chauvin ex Duby, 1830	A M
<i>hookeri</i> (Dillwyn) Maggs & Hommersand, 1993	A M	<i>diaphanum</i> (Lightfoot) Roth, 1806	A M
<i>priceanum</i> Maggs, Guiry & Rueness, 1991	A	<i>echionotum</i> J. Agardh, 1844	A M
<i>pseudobryssoides</i> (P.L. Crouan & H.M. Crouan) Halos, 1965	A	<i>flaccidum</i> (Kützing) Ardisson, 1871	A M
<i>roseum</i> (Roth) Maggs & L'Hardy-Halos, 1993	A	<i>fosliei</i> (Petersen) Petersen, 1935	A
<i>sepositum</i> (Gunnerus) Maggs & Hommersand, 1983	A	<i>fruticulosum</i> (Kützing) J. Agardh, 1894	A M
<i>tenuissimum</i> (Bonnemaison) Feldmann-Mazoyer, 1941	A M	<i>furcatum</i> (Petersen) Petersen, 1925	A
<i>tripinnatum</i> (C. Agardh) Feldmann-Mazoyer, 1941	A M	<i>gaditanum</i> (Clemente) Cremades, 1990	A M
<i>westbrookiae</i> Rueness & L'Hardy-Halos, 1991	A	<i>giacconei</i> Cormaci & G. Furnari, 1991	M
<i>Anotrichium</i>		<i>graeicum</i> Lazaridou & Boudouresque, 1992	M
<i>barbatum</i> (C. Agardh) Nägeli, 1862	A M	<i>hellenicum</i> Giaccone	M
<i>furcellatum</i> (J. Agardh) Baldock, 1976	A M	<i>incospicuum</i> Zanardini	M
<i>tenue</i> (C. Agardh) Nägeli, 1862	A M	<i>lepto cladum</i> Schiffner, 1933	M
<i>Antithamnion</i>		<i>pallidum</i> (Nägeli ex Kützing) Maggs & Hommersand, 1993	A
<i>amphigeneum</i> A.J.K. Millar, 1990	M	<i>pedicellatum</i> (Duby) J. Agardh	A M
<i>antillanum</i> Børgesen, 1917	A M	<i>penicillatum</i> Areschoug, 1849	A
<i>compactum</i> (Grunow) Schiffner	M	<i>petiti</i> Feldmann-Mazoyer	M
<i>cruciatum</i> (C. Agardh) Nägeli, 1847	A M	<i>pleurosporum</i> Schiffner, 1938	M
<i>decipiens</i> (J. Agardh) Athanasiadis, 1996	A M	<i>pseudostrictum</i> Schiffner, 1938	M
<i>densum</i> (Suhr) M. Howe, 1914	A	<i>recissum</i> Kylin	A
<i>diminutum</i> Wollaston, 1968	A	<i>rescissum</i> Kylin, 1907	A
<i>heterocladum</i> Funk	A M	<i>rosenvingei</i> Petersen, 1908	A
<i>pectinatum</i> (Montagne) Brauner ex Athanasiadis & Tittley	A M	<i>rubrum</i> (Hudson) C. Agardh, 1811	A M
<i>piliferum</i> Cormaci et G. Furnari	M	<i>secundatum</i> Lyngbye, 1819	A M
<i>tenuissimum</i> (Hauck) Schiffner, 1916	A M	<i>septentrionale</i> Petersen, 1911	A
<i>villosum</i> (Kützing) Athanasiadis, 1993	A	<i>shuttleworthianum</i> (Kützing) Rabenhorst, 1846	A
<i>Antithamnionella</i>		<i>siliquosum</i> (Kützing) Maggs & Hommersand, 1993	A M
<i>boergesenii</i> (Cormaci & G. Furnari) Athanasiadis, 1996	M	<i>siliquosum</i>	M
<i>elegans</i> (Berthold) J.H. Price & D.M. John, 1986	A M	<i>spinoso-pilosum</i> Kützing	A
<i>floccosa</i> (O.F. Müller) Whittick, 1980	A	<i>strobiliforme</i> G.W. Lawson & D.M. John	M
<i>spiropigraphidis</i> (Schiffner) E.M. Wollaston, 1968	A M	<i>terrimum</i> (G. Martens) Okamura, 1921	A M
<i>ternifolia</i> (J.D. Hooker & Harvey) Lyle, 1922	A M	<i>tenue</i> J. Agardh	A
<i>Balliella</i>		<i>tenuicornis</i> (Kützing) Waern, 1949	A
<i>cladoderma</i> (Zanardini) Athanasiadis, 1987	A M	<i>valvata</i> Schiffner	M
<i>Bornetia</i>		<i>Compsothamnion</i>	
<i>secundiflora</i> (J. Agardh) Thuret, 1855	A M	<i>decompositum</i> (J. Agardh) Maggs & L'Hardy-Halos, 1993	A
<i>Callithamniella</i>		<i>gracillimum</i> De Toni, 1903	A M
<i>tingitana</i> (Schousboe ex Bornet) Feldmann-Mazoyer	A M	<i>thywoides</i> (J.E. Smith) Nägeli, 1862	A M
<i>Callithamnion</i>		<i>Corallophila</i>	
<i>corymbosum</i> (J.E. Smith) Lyngbye, 1819	A M	<i>cinnabarina</i> (Grateloup ex Bory) R.E. Norris, 1993	M
<i>ellipticum</i> Montagne	A	<i>Crouania</i>	
<i>gailloni</i> (P.L. Crouan & H.M. Crouan) J. Agardh,		<i>attenuata</i> (C. Agardh) J. Agardh, 1842	A M
		<i>francescoi</i> Cormaci, G. Furnari & Scammacca, 1978	M
		<i>ischiana</i> (Funk) Boudouresque & M. Perret	M
		<i>Diplothamnion</i>	
		<i>jolyi</i> van den Hoek, 1978	A

<i>Dohniella</i>			
<i>nana</i> Mayhoub, 1975	M	<i>fernandeziana</i> Reiss, 1977	A
<i>neapolitana</i> Funk	M	<i>ferulacea</i> Suhr ex J. Agardh, 1863	A M
<i>Grallatoria</i>		<i>fibrata</i> (Dillwyn) Harvey, 1833	A
<i>reptans</i> M. Howe, 1920	A	<i>fibrillosa</i> (Dillwyn) Sprengel, 1827	A M
<i>Griffithsia</i>		<i>flexella</i> (C. Agardh) J. Agardh	A M
<i>capitata</i> Børgesen, 1930	A	<i>flocculosa</i> (C. Agardh) Kützing	A M
<i>corallinoides</i> (Linnaeus) Trevisan, 1845	A M	<i>foeniculacea</i> (C. Agardh) Sprengel	M
<i>devoniensis</i> Harvey, 1846	A	<i>foetidissima</i> Cocks ex Bornet, 1892	A M
<i>genoveseae</i> Feldmann	M	<i>fucoides</i> (Hudson) Greville, 1824	A M
<i>globulifera</i> (Harvey) J. Agardh	A	<i>funebris</i> De Notaris ex J. Agardh, 1851	A M
<i>opuntioides</i> J. Agardh, 1842	A M	<i>furcellata</i> (C. Agardh) Harvey, 1833	A M
<i>phyllamphora</i> J. Agardh, 1842	A M	<i>gonatophora</i> Kützing	A
<i>radicans</i> Kützing	A	<i>harveyi</i> J. Bailey, 1848	A M
<i>schousboei</i> Montagne, 1853	A M	<i>havanensis</i> Montagne, 1837	A
<i>Gulsonia</i>		<i>hemisphaerica</i> Areschoug, 1875	A
<i>nodulosa</i> (Ercegovic) J. Feldmann & G. Feldmann-		<i>hirta</i> J. Agardh	A
Mazoyer	M	<i>hochstetteriana</i> O.C. Schmidt	A
<i>Gymnophycus</i>		<i>kampsaxii</i> Børgesen, 1939	M
<i>hapsiphorus</i> Huisman & Kraft	A	<i>kieliana</i> Kaminski & Nizamuddin, 1975	A
<i>Gymnothamnion</i>		<i>lanosa</i> (Linnaeus) Tandy, 1931	A M
<i>elegans</i> (Schousboe ex C. Agardh) J. Agardh, 1892	A M	<i>mottei</i> Lauret, 1967	M
<i>pteroton</i> (Schousboe ex Bornet) Athanasiadis, 1996	A	<i>myriococca</i> Montagne	A
<i>Halosia</i>		<i>neglecta</i> Bornet	M
<i>elisae</i> Cormaci & G. Furnari, 1994	M	<i>nigra</i> (Hudson) Batters, 1902	A M
<i>Halurus</i>		<i>nutans</i> Montagne, 1841	A
<i>equisetifolius</i> (Lightfoot) Kützing, 1843	A M	<i>opaca</i> (C. Agardh) Moris & De Notaris, 1839	A M
<i>flosculosus</i> (J. Ellis) Maggs & Hommersand, 1993	A M	<i>ornata</i> J. Agardh, 1842	M
<i>flosculosus</i>	M	<i>orthocarpa</i> Rosenvinge, 1924	A
<i>Lejolisia</i>		<i>paniculata</i> Montagne, 1842	A M
<i>mediterranea</i> Bornet	M	<i>parvula</i> Suhr ex Kützing, 1864	M
<i>Microcladia</i>		<i>perforans</i> Cormaci, G. Furnari, Pizzuto & Serio, 1998	M
<i>glandulosa</i> (Solander ex Turner) Greville, 1830	A M	<i>polyspora</i> (C. Agardh) Montagne, 1841	A M
<i>Microgeliopsis</i>		<i>pulvinata</i> (Roth) Sprengel, 1827	A M
<i>horrida</i> Ercegovic	M	<i>requienii</i> Montagne	M
<i>Monosporus</i>		<i>rhuensis</i> Thuret	A
<i>pedicellatus</i> (J.E. Smith) Solier, 1845	A M	<i>ruchingeri</i> J. Agardh	M
<i>Neosiphonia</i>		<i>sanguinea</i> (C. Agardh) Zanardini	M
<i>sphaerocarpa</i> (Børgesen) M.S. Kim & I.K. Lee	A M	<i>scopulorum</i> Harvey, 1855	A M
<i>Pleonosporium</i>		<i>senticulosa</i> Harvey, 1862	A
<i>borreri</i> (J.E. Smith & Sowerby) Nägeli, 1862	A M	<i>sertularioides</i> (Grateloup) J. Agardh, 1863	A M
<i>caribaeum</i> (Børgesen) R.E. Norris, 1985	A M	<i>setigera</i> Kützing, 1849	M
<i>flexuosum</i> (C. Agardh) Bornet	A	<i>simulans</i> Harvey, 1849	A
<i>Plumaria</i>		<i>spinosa</i> (C. Agardh) J. Agardh	M
<i>plumosa</i> (Hudson) Kuntze, 1891	A	<i>stricta</i> (Dillwyn) Greville, 1824	A M
<i>Polysiphonia</i>		<i>stuposa</i> Zanardini ex Kützing	M
<i>adriatica</i> Schiffner, 1933	M	<i>subtilissima</i> Montagne, 1840	M
<i>affinis</i> Moore	A	<i>subulata</i> (Ducluzeau) P.L. Crouan & H.M. Crouan	A M
<i>allochroa</i> (Roth) Fries, 1845	A	<i>subulifera</i> (C. Agardh) Harvey, 1834	A M
<i>arachnoidea</i> (C. Agardh) Zanardini	M	<i>tenerrima</i> Kützing, 1843	M
<i>arctica</i> J. Agardh, 1863	A	<i>tepida</i> Hollenberg, 1958	A
<i>atlantica</i> Kapraun & J.N. Norris, 1982	A M	<i>tripinnata</i> J. Agardh	A M
<i>atra</i> Zanardini	M	<i>Pseudospora</i>	
<i>azorica</i> O.C. Schmidt	A	<i>adriatica</i> Schiffner, 1931	M
<i>banyulensis</i> Coppejans	M	<i>Pterothamnion</i>	
<i>barbatula</i> Kützing, 1849	M	<i>crispum</i> (Ducluzeau) Nägeli, 1862	A M
<i>beguinottii</i> Schiffner, 1938	M	<i>plumula</i> (J. Ellis) Nägeli, 1855	A M
<i>biformis</i> Zanardini, 1860	M	<i>polyacanthum</i> (Kützing) Nägeli, 1862	A M
<i>breviarticulata</i> (C. Agardh) Zanardini	A M	<i>spinescens</i> (Strömfelt) Athanasiadis, 1996	A
<i>brodiei</i> (Dillwyn) Sprengel, 1827	A M	<i>Ptilocladia</i>	
<i>carettia</i> Hollenberg, 1971	A	<i>horrida</i> Berthold, 1882	M
<i>caspica</i> Kützing	M	<i>Ptilota</i>	
<i>castagneyi</i> Kützing, 1863	M	<i>gunneri</i> P.C. Silva, Maggs & L.M. Irvine, 1993	A
<i>castelliana</i> De Notaris ex L. Dufour	M	<i>serrata</i> Kützing, 1847	A
<i>ceramiaeformis</i> P.L. Crouan & H.M. Crouan, 1867	A M	<i>Ptilothamnion</i>	
<i>cladorhiza</i> Ardisson, 1864	M	<i>codicolum</i> (E.Y. Dawson) I.A. Abbott, 1971	A
<i>codicola</i> Zanardini ex Frauenfeld, 1855	M	<i>pluma</i> (Dillwyn) Thuret, 1863	A M
<i>denudata</i> (Dillwyn) Greville ex Harvey, 1833	A M	<i>sphaericum</i> (P.L. Crouan & H.M. Crouan ex J.	
<i>derbesii</i> Solier ex Kützing, 1849	M	Agardh) Maggs & Hommersand, 1993	A
<i>deusta</i> (Roth) Sprengel	M	<i>Scagelia</i>	
<i>devoniensis</i> Maggs & Hommersand, 1993	A	<i>americana</i> (Harvey) Athanasiadis, 1996	A
<i>dichotoma</i> Kützing	A M	<i>pylaisaei</i> (Montagne) M.J. Wynne, 1985	A
<i>elongata</i> (Hudson) Sprengel, 1827	A M	<i>Scageliopsis</i>	
<i>elongella</i> Harvey, 1833	A M	<i>patens</i> Wollaston	A
<i>erythraea</i> Schousboe ex J. Agardh	A	<i>Scagelothamnion</i>	
		<i>pusillum</i> (Ruprecht) Athanasiadis, 1996	A

<i>Seirospora</i>		Hommersand, 1993	A
<i>apiculata</i> (Meneghini) G. Feldmann-Mazoyer	M	<i>minuta</i> (Kylin) Maggs & Hommersand, 1993	A
<i>giraudyi</i> (Kützing) De Toni, 1903	M	<i>spectabilis</i> J. Ernst & J. Feldmann, 1957	A
<i>humilis</i> Kützing, 1849	M	<i>Erythroglossum</i>	
<i>interrupta</i> (J.E. Smith) F. Schmitz, 1893	A M	<i>baleanicum</i> J. Agardh ex Kylin	M
<i>sphaerospora</i> J. Feldmann, 1942	M	<i>laciniatum</i> (Lightfoot) Maggs & Hommersand, 1993	A M
<i>Spermothamnion</i>		<i>lusitanicum</i> Ardré, 1970	A
<i>flabellatum</i> Bornet, 1876	A M	<i>sandrianum</i> (Kützing) Kylin, 1924	A M
<i>irregularare</i> (J. Agardh) Ardisson, 1883	A M	<i>Gonimocolax</i>	
<i>johannis</i> G. Feldmann	M	<i>roscoffensis</i> J. Feldmann & G. Feldmann, 1961	A
<i>repens</i> (Dillwyn) Rosenvinge, 1924	A M	<i>Gonimophyllum</i>	
<i> speluncarum</i> (Collins & Hervey) M. Howe	A	<i>buffhamii</i> Batters, 1892	A
<i>strictum</i> (C. Agardh) Ardisson, 1883	A M	<i>Haraldia</i>	
<i>Sphondylothamnion</i>		<i>lenormandii</i> (Derbès & Solier) Feldmann, 1939	A M
<i>multiplicifidum</i> (Hudson) Nägeli, 1862	A M	<i>Haraldiophyllum</i>	
<i>Spyridia</i>		<i>bonnemaisonii</i> (Kylin) A.D. Zinova, 1981	A M
<i>filamentosa</i> (Wulfen) Harvey, 1833	A M	<i>Hypoglossum</i>	
<i>hypnoides</i> (Bory) Papenfuss	A M	<i>hypoglossoides</i> (Stackhouse) Collins & Hervey, 1917	A M
<i>Tiffaniella</i>		<i>tenuifolium</i> (Harvey) J. Agardh	A
<i>capitata</i> (Schousboe ex Bornet) Doty & Mefez	A M	<i>Membranoptera</i>	
<i>feldmanniae</i> (P. Huve) Gillis & Coppejans	M	<i>alata</i> (Hudson) Stackhouse, 1809	A
<i>gorgonea</i> (Montagne) Doty & Mefez	A	<i>Myriogramme</i>	
<i>Vickersia</i>		<i>carnnea</i> (J.J. Rodríguez) Kylin	A M
<i>baccata</i> (J. Agardh) Karsakoff, 1896	A M	<i>costata</i> P.J.L. Dangeard, 1949	A
<i>Wrangelia</i>		<i>distromatica</i> Boudouresque, 1971	M
<i>argus</i> (Montagne) Montagne, 1856	A	<i>repens</i> Hollenberg, 1945	M
<i>penicillata</i> (C. Agardh) C. Agardh, 1828	A M	<i>tristromatica</i> (J.J. Rodríguez ex Mazza) Boudouresque	M
Family Dasyaceae		<i>unistromaticum</i> Coppejans	M
<i>Dasya</i>		<i>Nitophyllum</i>	
<i>baillouviana</i> (S.G. Gmelin) Montagne, 1841	A M	<i>albidum</i> Ardisson	M
<i>caraibica</i> Børgesen	A	<i>flabellatum</i> Ercegovic	M
<i>corymbifera</i> J. Agardh, 1841	A M	<i>marmoratum</i> J.J. Rodríguez, 1889	M
<i>crouaniana</i> J. Agardh, 1890	A	<i>micropunctatum</i> Funk, 1955	M
<i>elongata</i> Sonder, 1845	M	<i>punctatum</i> (Stackhouse) Greville, 1830	A M
<i>hutchinsiae</i> Harvey, 1833	A M	<i>rotundum</i> Funk, 1955	M
<i>ocellata</i> (Grateloup) Harvey, 1833	A M	<i>Pantoneura</i>	
<i>punicea</i> (Zanardini) Meneghini ex Zanardini, 1841	A M	<i>angustissima</i> (Turner) Kylin, 1924	A
<i>rigidula</i> (Kützing) Ardisson, 1878	A M	<i>Phycodrys</i>	
<i>villosa</i> Harvey	M	<i>rubens</i> (Linnaeus) Batters, 1902	A
<i>Dasyella</i>		<i>Platysiphonia</i>	
<i>gracilis</i> Falkenberg, 1901	M	<i>caribaea</i> D.L. Ballantine & M.J. Wynne	A
<i>Eupogodon</i>		<i>delicata</i> (Clemente) Cremades, 1990	A
<i>apiculatus</i> (C. Agardh) P.C. Silva	M	<i>miniata</i> (C. Agardh) Børgesen, 1931	A
<i>penicillata</i> (Zanardini) P.C. Silva	M	<i>Polyneura</i>	
<i>planus</i> (C. Agardh) Kützing, 1845	A M	<i>bonnemaisonii</i> (C. Agardh) Maggs & Hommersand, 1993	A
<i>spinellus</i> (C. Agardh) Kützing	A M	<i>Radicilingua</i>	
<i>Halydictyon</i>		<i>adriatica</i> (Kylin) Papenfuss	M
<i>mirabile</i> Zanardini	A M	<i>reptans</i> (Kylin) Papenfuss	M
<i>Heterosiphonia</i>		<i>thyisanorhizans</i> (Holmes) Papenfuss, 1956	A M
<i>crispella</i> (C. Agardh) M.J. Wynne, 1985	A M	<i>Taenioma</i>	
<i>plumosa</i> (J. Ellis) Batters, 1902	A M	<i>nanum</i> (Kützing) Papenfuss, 1952	A M
Family Delesseriaceae		<i>perpusillum</i> (J. Agardh) J. Agardh, 1863	A M
<i>Acrosorium</i>		Family Rhodomelaceae	
<i>reptans</i> (P.L. Crouan & H.M. Crouan) Kylin	A M	<i>Acanthophora</i>	
<i>venulosum</i> (Zanardini) Kylin, 1924	A M	<i>muscoidea</i> (Linnaeus) Bory, 1828	M
<i>Apoglossocolax</i>		<i>nayadiformis</i> (Delile) Papenfuss, 1968	M
<i>pusilla</i> Maggs & Hommersand, 1993	A	<i>Aiolocolax</i>	
<i>Apoglossum</i>		<i>pulchellus</i> Pocock, 1956	A
<i>gregarium</i> (E.Y. Dawson) M.J. Wynne, 1985	M	<i>Alsidium</i>	
<i>ruscifolium</i> (Turner) J. Agardh, 1898	A M	<i>corallinum</i> C. Agardh, 1827	A M
<i>Arachnophyllum</i>		<i>helminthochorton</i> (Latourrette) Kützing	A M
<i>conservaceum</i> (Meneghini) Zanardini	M	<i>Aphanocladia</i>	
<i>Asterocolax</i>		<i>stichidiosa</i> (Funk) Ardré, 1970	A M
<i>erythroglossi</i> J. Feldmann & G. Feldmann, 1951	A	<i>Boergesenella</i>	
<i>Branchioglossum</i>		<i>deludens</i> (Falkenberg) Kylin	M
<i>prostratum</i> C.W. Schneider, 1974	A	<i>fruticulosa</i> (Wulfen) Kylin, 1956	A M
<i>Cottoniella</i>		<i>martensiana</i> (Kützing) Ardré, 1970	A
<i>filamentosa</i> (M. Howe) Børgesen, 1920	A M	<i>thuyoides</i> (Harvey) Kylin, 1956	A M
<i>libyensis</i> Nizamuddin & Godeh	M	<i>Bostrychia</i>	
<i>Cryptopleura</i>		<i>scorpioides</i> (Hudson) Montagne ex Kützing, 1849	A M
<i>ramosa</i> (Hudson) Kylin ex L. Newton, 1931	A M	<i>Brongniartella</i>	
<i>Delesseria</i>		<i>byssoides</i> (Goodenough & Woodward) F. Schmitz, 1893	A M
<i>sanguinea</i> (Hudson) J.V. Lamouroux, 1813	A M	<i>Chondria</i>	
<i>Drachiella</i>			
<i>heterocarpa</i> (Chauvin ex Duby) Maggs &			

<i>arcuata</i> Hollenberg, 1945	A	<i>pelagosae</i> (Schiffner) K.W. Nam	M
<i>boryana</i> (J. Agardh) De Toni	M	<i>pinnatifida</i> (Hudson) Stackhouse, 1809	A M
<i>capillaris</i> (Hudson) M.J. Wynne, 1991	A M	<i>ramosissima</i> (Oeder) Athanasiadis, 1996	A M
<i>coerulescens</i> (J. Agardh) Falkenberg, 1901	A M	<i>verlaquei</i> G. Furnari	M
<i>collinsiana</i> M. Howe	M		
<i>curvilineata</i> Collins & Hervey, 1917	A M	<i>Pterosiphonia</i>	
<i>dasyphylla</i> (Woodward) C. Agardh, 1817	A M	<i>ardeana</i> Maggs & Hommersand, 1993	A M
<i>densa</i> P.J.L. Dangeard, 1951	A	<i>complanata</i> (Clemente) Falkenberg, 1901	A M
<i>mairei</i> Feldmann-Mazoyer, 1949	A M	<i>parasitica</i> (Hudson) Falkenberg, 1901	A M
<i>oppositiclada</i> E.Y. Dawson, 1945	M	<i>pennata</i> (C. Agardh) Sauvageau, 1897	A M
<i>polyrhiza</i> Collins & Hervey, 1917	M	<i>pinnulata</i> (Kützing) Maggs & Hommersand, 1993	A M
<i>pygmaea</i> Garbary & Vandermeulen	M	<i>spinifera</i> (Kützing) Ardré, 1967	M
<i>scintillans</i> G. Feldmann, 1964	A M		
<i>Chondrophycus</i>		<i>Rhodomela</i>	
<i>corallopsis</i> (Montagne) Garbary & Harper	A	<i>confervoides</i> (Hudson) P.C. Silva, 1952	A
<i>paniculatus</i> (C. Agardh) G. Furnari	A M	<i>lycopodioides</i> (Linnaeus) C. Agardh, 1822	A
<i>papillosum</i> (C. Agardh) Garbary & Harper	A M		
<i>patentirameus</i> (Montagne) K.W. Nam	A M	<i>Rodriguezella</i>	
<i>perforatus</i> (Bory) K.W. Nam	A	<i>bornetii</i> (J.J. Rodríguez) F. Schmitz ex J.J. Rodríguez, 1896	M
<i>poiteaui</i> (J.V. Lamouroux) K.W. Nam	A	<i>ligulata</i> Feldmann, 1942	M
<i>Ctenosiphonia</i>		<i>pinnata</i> (Kützing) F. Schmitz ex Falkenberg	M
<i>hypnoidea</i> (Welwitsch ex J. Agardh) Falkenberg, 1897A		<i>strafforelooi</i> F. Schmitz ex J.J. Rodríguez	M
<i>Digenea</i>			
<i>simplex</i> (Wulfen) C. Agardh, 1822	A M	<i>Rytiophloea</i>	
<i>Dipterosiphonia</i>		<i>tinctoria</i> (Clemente) C. Agardh, 1824	A M
<i>dendritica</i> (C. Agardh) F. Schmitz, 1897	A M		
<i>rigens</i> (Schousboe ex C. Agardh) Falkenberg	A M	<i>Stictothamnion</i>	
<i>Erythrocystis</i>		<i>cymatophilum</i> Børgesen	A M
<i>montagnei</i> (Derbès & Solier) P.C. Silva	A M	<i>Streblocladia</i>	
<i>Halophysis</i>		<i>collabens</i> (C. Agardh) Falkenberg, 1901	A M
<i>incura</i> (Hudson) Batters, 1902	A M	<i>Sympyocladia</i>	
<i>Janczewskia</i>		<i>merchantioides</i> (Harvey) Falkenberg, 1897	A
<i>verrucaformis</i> Solms-Laubach, 1877	A M	<i>Womersleyella</i>	
<i>Laurencia</i>		<i>setacea</i> (Hollenberg) R.E. Norris, 1992	A M
<i>brongniartii</i> J. Agardh, 1841	A		
<i>caspica</i> A.D. Zinova & Zaberzhinskav	M	Order Compsopogonales	
<i>chondrioides</i> Børgesen, 1918	A M	Family Compsopogonaceae	
<i>corallopsis</i> (Montagne) M. Howe, 1918	A	<i>Compsopogon</i>	
<i>coronopus</i> J. Agardh	M	<i>aegyptiacus</i> Aleem, 1981	M
<i>decumbens</i> Kützing, 1865	A		
<i>epiphylla</i> Boisset & Lino	M	Order Corallinales	
<i>fastigiata</i> Montagne, 1846	M	Family Corallinaceae	
<i>flexilis</i> Setchell, 1926	A	<i>Amphiroa</i>	
<i>glandulifera</i> (Kützing) Kützing, 1849	A M	<i>beauvoisii</i> J.V. Lamouroux, 1816	A M
<i>intricata</i> J.V. Lamouroux, 1813	A M	<i>cryptarthrodia</i> Zanardini, 1843	A M
<i>majuscula</i> (Harvey) A.H.S. Lucas, 1947	A M	<i>fragilissima</i> (Linnaeus) J.V. Lamouroux, 1816	A M
<i>microcladia</i> Kützing, 1865	A M	<i>rigida</i> J.V. Lamouroux, 1816	A M
<i>minuta</i> Vandermeulen, Garbary & Guiry, 1990	A M	<i>vanbosseae</i> Lemoine	A
<i>obtusa</i> (Hudson) J.V. Lamouroux, 1813	A M		
<i>oophora</i> Kützing, 1865	M	<i>Boreolithon</i>	
<i>pyramidalis</i> Bory ex Kützing, 1849	A M	<i>van-heurckii</i> (Heydrich) A.S. Harvey & Woelkerling, 1995	A M
<i>radicans</i> (Kützing) Kützing, 1849	M		
<i>tenera</i> C.K. Tseng, 1946	A	<i>Choreonema</i>	
<i>viridis</i> Gil-Rodríguez & Haroun	A	<i>thurettii</i> (Bornet) F. Schmitz, 1889	A M
<i>Leptosiphonia</i>		<i>Clathromorphum</i>	
<i>schousboei</i> (Thuret) Kylin, 1956	A	<i>circumspectum</i> (Strömfelt) Foslie, 1898	A
<i>Lophocladia</i>		<i>compactum</i> (Kjellman) Foslie, 1898	A
<i>lallemandii</i> (Montagne) F. Schmitz, 1893	M	<i>Corallina</i>	
<i>trichoclados</i> (Mertens ex C. Agardh) F. Schmitz, 1893	A M	<i>elegans</i> J.V. Lamouroux	A M
<i>Lophosiphonia</i>		<i>elongata</i> J. Ellis & Solander, 1786	A M
<i>cristata</i> Falkenberg, 1901	A M	<i>millegrana</i> Lamarck	A
<i>obscura</i> (C. Agardh) Falkenberg, 1897	A M	<i>officinalis</i> Linnaeus, 1758	A M
<i>reptabunda</i> (Suhr) Kylin, 1956	A M		
<i>Odonthalia</i>		<i>Dermatolithon</i>	
<i>dentata</i> (Linnaeus) Lyngbye, 1819	A	<i>geometricum</i> M. Lemoine	A
<i>Ophidocladus</i>		<i>papillosum</i> (Zanardini) Foslie, 1909	A
<i>simpliciusculus</i> (P.L. Crohan & H.M. Crohan)	A M		
Falkenberg, 1897		<i>Exilicrusta</i>	
<i>Osmundaria</i>		<i>parva</i> Y.M. Chamberlain, 1992	A
<i>volubilis</i> (Linnaeus) R.E. Norris, 1991	A M		
<i>Osmundea</i>		<i>Fosliella</i>	
<i>hybrida</i> (A.P. Candolle) K.W. Nam, 1994	A	<i>paschalis</i> (M. Lemoine) Setchell & N.L. Gardner, 1930	A
<i>maggsiana</i> Serio, Cormaci & G. Furnari, 1999	M	<i>zonalis</i> (P.L. Crohan & H.M. Crohan) Feldmann	A M
<i>osmundula</i> (S.G. Gmelin) K.W. Nam & Maggs, 1994	A		
<i>pelagiensis</i> G. Furnari	M	<i>Goniolithon</i>	
		<i>byssoides</i> (Lamarck) Foslie, 1898	M
		<i>Haliptilon</i>	
		<i>attenuata</i> (Kützing) Garbary & Johansen, 1982	M
		<i>cubense</i> (Montagne ex Kützing) Garbary & H.W. Johansen, 1982	A
		<i>squamatum</i> (Linnaeus) H.W. Johansen, L.M. Irvine & A. Webster, 1973	A M
		<i>subulatum</i> (Ellis & Solander) Johansen, 1970	M
		<i>virgatum</i> (Zanardini) Garbary & H.W. Johansen	A M

<i>Hydrolithon</i>			
<i>boreale</i> (Foslie) Y.M. Chamberlain	A M	<i>sonderi</i> Hauck, 1883	A M
<i>cruciatum</i> (Bressan) Y.M. Chamberlain, 1994	A M	<i>trophiforme</i> Unger, 1858	A
<i>farinosum</i> (J.V. Lamouroux) Penrose & Y.M. Chamberlain, 1993	A M	<i>tusserense</i> Foslie, 1905	A
<i>oligocarpum</i> (Foslie) Foslie	A	<i>ungeri</i> Kjellman, 1883	
<i>onkodes</i> (Heydrich) Penrose & Woelkerling, 1992	A	<i>valens</i> Foslie, 1909	M
<i>samoënsse</i> (Foslie) Keats & Y.M. Chamberlain, 1994	A	<i>Melobesia</i>	
<i>sargassi</i> (Foslie) Y.M. Chamberlain, 1994	A	<i>conferoides</i> Funk	M
<i>Jania</i>		<i>membranacea</i> (Esper) J.V. Lamouroux, 1812	A M
<i>adhaerens</i> J.V. Lamouroux, 1816	A M	<i>Mesophyllum</i>	
<i>capillacea</i> Harvey, 1853	A	<i>alternans</i> (Foslie) Cabioch & Mendoza, 1998	A M
<i>crassa</i> J.V. Lamouroux, 1821	A	<i>arcticum</i> (Kjellman) Athanasiadis, 1996	A
<i>longiarthra</i> E.Y. Dawson, 1953	M	<i>brachycladum</i> (Foslie) W.H. Adey	A
<i>longifurca</i> Zanardini	A M	<i>canariense</i> (Foslie) Lemoine	A
<i>micrarthrodia</i> J.V. Lamouroux, 1816	A	<i>caraiensis</i> (Foslie) M. Lemoine	A
<i>pumila</i> J.V. Lamouroux, 1816	A	<i>ectocarpum</i> (Foslie) W.H. Adey	A
<i>rubens</i> (Linnaeus) J.V. Lamouroux, 1812	A M	<i>lichenoides</i> (J. Ellis) M. Lemoine, 1928	A M
<i>Kvaleya</i>		<i>macroblastum</i> (Foslie) W.H. Adey, 1970	M
<i>epilaeve</i> W.H. Adey & Sperapani, 1971	A	<i>Neogoniolithon</i>	
<i>Leptophytum</i>		<i>accretum</i> (Foslie & M. Howe) Setchell & Mason	A
<i>bisporum</i> (Foslie) W.H. Adey	A	<i>brassica-florida</i> (Harvey) Setchell & Mason, 1943	M
<i>elatum</i> Y.M. Chamberlain, 1990	A	<i>caribaeum</i> (Foslie) W.H. Adey	A
<i>Litholepis</i>		<i>hirtum</i> (M. Lemoine) Afonso Corillo	A
<i>mediterranea</i> Foslie, 1905	A M	<i>illitum</i> (M. Lemoine) Afonso Corillo	A
<i>Lithophyllum</i>		<i>mamillosum</i> (Hauck) Setchell & Mason	M
<i>applicatum</i> M. Lemoine	A	<i>orotavicum</i> (Foslie) M. Lemoine ex Afonso Corillo	A
<i>azorum</i> M. Lemoine	A	<i>Phymatolithon</i>	
<i>bipartitum</i> M. Lemoine	A	<i>bisporum</i> (Foslie) Afonso-Carillo, 1984	A
<i>byssoides</i> (J.V. Lamouroux) Foslie, 1900	M	<i>bornetii</i> (Foslie) Foslie	A M
<i>capense</i> Rozanov	A	<i>brunneum</i> Y.M. Chamberlain, 1994	A
<i>crodelioides</i> Boudouresque	M	<i>calcareum</i> (Pallas) W.H. Adey & D.L. McKibbin, 1970	A M
<i>crouanii</i> Foslie, 1899	A	<i>foecundum</i> (Kjellman) L. Düwel & S. Wegeberg, 1996	A
<i>cystoseirae</i> (Hauck) Heydrich, 1897	A	1996	A
<i>decussatum</i> (Ellis & Solander) Philippi	A M	<i>investiens</i> (Foslie) Foslie, 1905	A
<i>dentatum</i> (Kützing) Foslie, 1900	A M	<i>laevigatum</i> (Foslie) Foslie, 1898	A
<i>duckeri</i> Woelkerling, 1983	A M	<i>lamii</i> (M. Lemoine) Y.M. Chamberlain, 1991	A
<i>fasciculatum</i> (Lamarck) Foslie, 1900	A M	<i>lenormandii</i> (J.E. Areschoug) W.H. Adey, 1966	A M
<i>frondosum</i> (Dufour) G. Furnari, Cormaci & Alongi, 1996	A M	<i>purpureum</i> (P.L. Crouan & H.M. Crouan)	
<i>hibernicum</i> Foslie, 1906	A	Woelkerling & L.M. Irvine, 1986	A M
<i>incrustans</i> Philippi, 1837	A	<i>tenue</i> (Rosenvinge) L. Düwel & S. Wegeberg, 1996	A
<i>irregularis</i> (Foslie) Huvé ex Steentoft	A M	<i>tenuissimum</i> (Foslie) W.H. Adey	A M
<i>lichenoides</i> Philipp, 1837	A	<i>Phaeophyllum</i>	
<i>lobatum</i> M. Lemoine	A M	<i>caudatae</i> (Foslie) P.L. Jones & Woelkerling, 1984	A M
<i>nitorum</i> W.H. Adey & P.J. Adey, 1973	A	<i>confervicolum</i> (Kützing) Y.M. Chamberlain, 1983	A M
<i>orbiculatum</i> (Foslie) Foslie, 1900	A M	<i>coronatum</i> (Rosanoff) Penrose	M
<i>papillosum</i> (Zanardini ex Hauck) Foslie	A M	<i>fragile</i> Kützing, 1843	A M
<i>polycephalum</i> Foslie	A	<i>limitatum</i> (Foslie) Y.M. Chamberlain, 1983	A
<i>racemus</i> (Lamarck) Foslie	A M	<i>lobescens</i> Y.M. Chamberlain, 1983	A
<i>tortuosum</i> (Esper) Foslie, 1900	A M	<i>myriocarpum</i> (P.L. Crouan & H.M. Crouan) Y.M.	
<i>trochanter</i> (Bory) H. Huvé	M	Chamberlain, 1983	A
<i>vickersiae</i> M. Lemoine, 1929	A	<i>subplanum</i> (Rosenvinge) Y.M. Chamberlain ex South & Tittley, 1986	A
<i>Lithoporella</i>		<i>Pseudolithophyllum</i>	
<i>sauvageaui</i> (Foslie) W.H. Adey	A	<i>esperi</i> M. Lemoine	A
<i>Lithothamnion</i>		<i>Spongites</i>	
<i>breviaxe</i> Foslie, 1895	A	<i>fruticulosus</i> Kützing, 1841	A M
<i>coralloides</i> (P.L. Crouan & H.M. Crouan) P.L. Crouan & H.M. Crouan, 1867	A M	<i>Tenarea</i>	
<i>crispatum</i> Hauck	M	<i>adhaerens</i> M. Lemoine	A
<i>ectocarpum</i> Foslie	A	<i>Titanoderma</i>	
<i>esperi</i> Heydrich	A	<i>byssoides</i> (Lamarck) Y.M. Chamberlain & Woelkerling, 1988	M
<i>flavescens</i> Kjellman, 1883	A	<i>corallinae</i> (P.L. Crouan & H.M. Crouan) Woelkerling, Y.M. Chamberlain & P.C. Silva, 1985	A M
<i>fornicatum</i> Foslie, 1891	A	<i>laminariae</i> (P.L. Crouan & H.M. Crouan) Y.M.	
<i>glaciele</i> Kjellman, 1883	A	Chamberlain, 1991	A
<i>grande</i> Foslie, 1905	A	<i>pustulatum</i> (J.V. Lamouroux) Nägeli, 1858	A M
<i>hamelii</i> M. Lemoine, 1931	A	<i>verrucatum</i> (J.V. Lamouroux) Y.M. Chamberlain, 1986	A M
<i>intermedium</i> Kjellman, 1883	A	<i>Family Sporolithaceae</i>	
<i>lemoineae</i> W.H. Adey, 1970	A	<i>Sporolithon</i>	
<i>minervae</i> Basso, 1995	M	<i>africanum</i> (Foslie) Afonso-Carrillo, Chacaba & Sansón, 1984	
<i>murmanicum</i> Elenkin, 1905	A	<i>ptychoides</i> Heydrich, 1897	M
<i>nodulosum</i> Foslie, 1895	A	<i>Order Cryptonemiales</i>	
<i>norvegicum</i> (Areschoug) Kjellman, 1883	A M		
<i>philippii</i> Foslie, 1897	M		
<i>propontidii</i> Foslie			

Family Acrosymphtaceae			
<i>Acrosymphton</i>			
<i>purpuriferum</i> (J. Agardh) Sjöstedt	A M		M
<i>taylorii</i> I.A. Abbott, 1962	M		A
Family Corynomorphaceae			
<i>Corynomorpha</i>			
<i>prismatica</i> (J. Agardh) J. Agardh	A		A M
Family Dumontiaceae			
<i>Dilsea</i>			
<i>carnosa</i> (Schmidel) Kuntze, 1898	A		
<i>integra</i> (Kjellman) Rosenvinge, 1898	A		
<i>Dudresnaya</i>			
<i>canariensis</i> Tabares, Afonso-Carrillo, Sansón & Reyes, 1997	A		A M
<i>crassa</i> M. Howe	A		
<i>verticillata</i> (Withering) Le Jolis, 1863	A M		
<i>Dumontia</i>			
<i>canariensis</i> Montagne	A		
<i>contorta</i> (S.G. Gmelin) Ruprecht, 1850	A		
<i>Pikea</i>			
<i>californica</i> Harvey, 1853	A		
Family Gloiosiphoniaceae			
<i>Cruoriopsis</i>			
<i>danica</i> Rosenvinge, 1918	A		
<i>Gloiosiphonia</i>			
<i>capillaris</i> (Hudson) Carmichael ex Berkeley, 1833	A M		
<i>Plagiospora</i>			
<i>gracilis</i> Kuckuck, 1897	A		
<i>Schimmelmannia</i>			
<i>schousboei</i> (J. Agardh) J. Agardh, 1851	M		
<i>Thuretella</i>			
<i>schousboei</i> (Thuret) F. Schmitz, 1897	A M		
Family Halymeniacae			
<i>Acrodiscus</i>			
<i>vidovichii</i> (Meneghini) Zanardini	M		
<i>Aedes</i>			
<i>marginata</i> (Roussel) F. Schmitz	M		
<i>Cryptonemia</i>			
<i>crenulata</i> (J. Agardh) J. Agardh, 1851	A		
<i>hibernica</i> Guiry & L.M. Irvine, 1974	A		
<i>lomatia</i> (Bertoloni) J. Agardh, 1851	A M		
<i>seminervis</i> (C. Agardh) J. Agardh, 1851	A M		
<i>tunaeformis</i> (A. Bertolini) Zanardini	M		
<i>Dermocorynus</i>			
<i>montagnae</i> P.L. Crouan & H.M. Crouan, 1858	A		
<i>Grateloupia</i>			
<i>cosentinii</i> Kützing	M		
<i>cuneifolia</i> J. Agardh	M		
<i>dichotoma</i> J. Agardh, 1842	A M		
<i>doryphora</i> (Montagne) M. Howe, 1914	A M		
<i>filiicina</i> (J.V. Lamouroux) C. Agardh, 1822	A M		
<i>proteus</i> Kützing, 1843	A M		
<i>Halymenia</i>			
<i>asymmetrica</i> Gargiulo, de Masi & Tripodi, 1986	M		
<i>dichotoma</i> J. Agardh	A M		
<i>elongata</i> C. Agardh, 1822	A M		
<i>imbriata</i> Zanardini	M		
<i>floresia</i> (Clemente) C. Agardh, 1822	A M		
<i>floridana</i> J. Agardh	A		
<i>latifolia</i> P.L. Crouan & H.M. Crouan ex Kützing, 1866	A M		
<i>pluribola</i> Ercegovic	M		
<i>rosea</i> M. Howe & W.R. Taylor	A		
<i>vinacea</i> M. Howe & W.R. Taylor	A		
Family Kallymeniaceae			
<i>Callocolax</i>			
<i>neglectus</i> F. Schmitz ex Batters, 1895	A		
<i>Callophyllis</i>			
<i>cristata</i> (C. Agardh) Kützing, 1849	A		
<i>fastigiata</i> (J. Agardh) J. Agardh, 1855	A		
<i>laciniosa</i> (Hudson) Kützing, 1843	A M		
<i>Kallymenia</i>			
<i>seldmannii</i> Codomier, 1971	A M		
<i>lacerata</i> Feldmann	M		
<i>palens</i> (J. Agardh) Parkinson			M
<i>perforata</i> J. Agardh			A
<i>reniformis</i> (Turner) J. Agardh, 1842			A M
<i>requienii</i> J. Agardh			A M
<i>schizophylla</i> J. Agardh, 1847			A
<i>spathulata</i> (J. Agardh) P.G. Parkinson			M
<i>Meredithia</i>			
<i>microphylla</i> (J. Agardh) J. Agardh, 1892			A M
Family Peyssonneliaceae			
<i>Metapeyssonnelia</i>			
<i>feldmannii</i> Boudouresque, Coppejans & Marcot			M
<i>Peyssonnelia</i>			
<i>armorica</i> (P.L. Crouan & H.M. Crouan) Weber-van Bosse, 1916			A M
<i>atropurpurea</i> P.L. Crouan & H.M. Crouan, 1867			A M
<i>bornetii</i> Boudouresque & Denizot			M
<i>coriacea</i> J. Feldmann, 1941			A M
<i>crispata</i> Boudouresque & Denizot			M
<i>dubyi</i> P.L. Crouan & H.M. Crouan, 1844			A M
<i>harveyana</i> P.L. Crouan & H.M. Crouan ex J. Agardh, 1852			A M
<i>immersa</i> Maggs & L.M. Irvine, 1983			A M
<i>inamoena</i> Pilger, 1911			A M
<i>magna</i> Ercegovic, 1949			M
<i>orientalis</i> (Weber-van Bosse) Cormaci & G. Furnari			M
<i>polymorpha</i> (Zanardini) F. Schmitz, 1879			A M
<i>rosa-marina</i> Boudouresque & Denizot, 1973			A M
<i>rosenvingei</i> Schmitz, 1893			A
<i>rubra</i> (Greville) J. Agardh, 1851			A M
<i>squamaria</i> (S.G. Gmelin) Decaisne, 1841			A M
<i>stoechas</i> Boudouresque & Denizot			M
<i>Polystrata</i>			
<i>compacta</i> (Foslie) Denizot, 1968			M
<i>dura</i> Heydrich, 1905			A
<i>fosliei</i> (Weber-van Bosse) Denizot, 1968			M
Family Rhizophyllidaceae			
<i>Contarinia</i>			
<i>peyssonneliaeformis</i> Zanardini, 1843			M
<i>squamariae</i> (Meneghini) Denizot, 1968			M
Order Erythropeltidales			
Family Erythropeltidaceae			
<i>Erythrocladia</i>			
<i>irregularis</i> Rosenvinge, 1909			A M
<i>polystromatica</i> P.A. Dangeard, 1932			A
<i>violacea</i> P.J.L. Dangeard, 1968			A
<i>Erythrotrichia</i>			
<i>carnea</i> (Dillwyn) J. Agardh, 1883			A M
<i>investiens</i> (Zanardini) Bornet, 1892			A M
<i>reflexa</i> (P.L. Crouan & H.M. Crouan) Thuret ex De Toni, 1897			A M
<i>rosea</i> P.J.L. Dangeard			M
<i>simplex</i> P.J.L. Dangeard, 1968			A
<i>welwitschii</i> (Ruprecht) Batters, 1902			A
<i>Porphyropsis</i>			
<i>coccinea</i> (J. Agardh ex Areschoug) Rosenvinge, 1909			A M
<i>imperfecta</i> Kornmann & Sahling, 1985			A
<i>Porphyrostromium</i>			
<i>boryanum</i> (Montagne) Trevisan, 1848			A M
<i>ciliare</i> (Carmichael ex Harvey) M.J. Wynne, 1986			A M
<i>Sahlingia</i>			
<i>subintegra</i> (Rosenvinge) Kornmann, 1989			A M
Order Gelidiales			
Family Gelidiaceae			
<i>Beckerella</i>			
<i>mediterranea</i> H. Huvé			M
<i>Gelidiella</i>			
<i>acerosa</i> (Forsskål) Feldmann & G. Hamel, 1934			A
<i>antipai</i> Celan, 1938			A M
<i>calcicola</i> Maggs & Guiry, 1987			A
<i>lubrica</i> (Kützing) Feldmann & G. Hamel, 1934			M
<i>machrisiana</i> Dawson, 1957			M
<i>nigrescens</i> (Feldmann) Feldmann & G. Hamel			A M
<i>pannosa</i> (J. Feldmann) J. Feldmann & G. Hamel,			

1934		A M	<i>strafforellii</i> Ardissonne	M
<i>ramellosa</i> (Kützing) Feldmann & G. Hamel, 1934		A M		
<i>sanctarum</i> Feldmann & G. Hamel, 1934		A		
<i>tinerfensis</i> Seoane-Camba, 1977		A		
<i>Gelidium</i>			Family Furcellariaceae	
<i>affine</i> Schiffner		M	<i>Furcellaria</i>	
<i>arbuscula</i> Bory ex Børgesen		A	<i>lumbricalis</i> (Hudson) J.V. Lamouroux, 1813	A M
<i>bipectinatum</i> Furnari, 1999		M	<i>Halarachnion</i>	
<i>canariensis</i> (Grunow) Seoane Camba ex Haroun, Gil-Rodríguez & Díaz de Castro, in		A	<i>ligulatum</i> (Woodward) Kützing, 1843	A M
<i>corneum</i> (Hudson) J.V. Lamouroux, 1813		A M	<i>Neurocaulon</i>	
<i>crinale</i> (Turner) Gaillón, 1828		A M	<i>foliosum</i> (Meneghini) Zanardini	A M
<i>divaricatum</i> G. Martens, 1868		M		
<i>hypnosum</i> Zanardini		M		
<i>intertextum</i> P.J.L. Dangeard, 1949		A	Family Gigartinaceae	
<i>maggiae</i> Rico & Guiry, 1997		A	<i>Chondracanthus</i>	
<i>microdon</i> Kützing, 1849		A M	<i>acicularis</i> (Roth) Fredericq, 1993	A M
<i>micropterum</i> Kützing, 1868		M	<i>teedei</i> (Mertens ex Roth) Kützing, 1843	A M
<i>minusculum</i> (Weber-van Bosse) R.E. Norris, 1992		M	<i>Chondrus</i>	
<i>pulchellum</i> (Turner) Kützing, 1868		A M	<i>crispus</i> Stackhouse, 1797	A
<i>pulvinatum</i> (Kützing) Thuret ex Bornet		A	<i>Gigartina</i>	
<i>pusillum</i> (Stackhouse) Le Jolis, 1863		A M	<i>pistillata</i> (S.G. Gmelin) Stackhouse, 1809	A M
<i>replans</i> (Suhr) Kylin, 1938		A		
<i>secundatum</i> Zanardini		M		
<i>spathulatum</i> (Kützing) Bornet		A M	Family Haemeschariaceae	
<i>spinosum</i> (S.G. Gmelin) P.C. Silva, 1996		A M	<i>Haemeschia</i>	
<i>torulosum</i> Kützing		A	<i>hennedyi</i> (Harvey) K.L. Vinogradova & T. Yacovleva, 1989	A
<i>venetum</i> Schiffner		M		
<i>versicolor</i> (S.G. Gmelin) J.V. Lamouroux		A	Family Hypnaceae	
<i>Pterocladia</i>			<i>Hypnea</i>	
<i>tripolitana</i> De Toni & Levi		M	<i>arbuscula</i> P.J.L. Dangeard	A
<i>Pterocladiella</i>			<i>coccinea</i> (Clemente) Cremades, 1990	A
<i>capillacea</i> (S.G. Gmelin) Santelices & Hommersand, 1997		A M	<i>cornuta</i> (Kützing) J. Agardh, 1851	M
<i>melanoidea</i> (Schousboe ex Bornet) Santelices & Hommersand, 1997		A M	<i>esperi</i> Bory, 1828	M
Order Gigartinales			<i>flagelliformis</i> Greville ex J. Agardh	A
Family Calosiphonaceae			<i>furnariana</i> Cormaci, Alongi and Dinaro, 1993	M
<i>Calosiphonia</i>			<i>musciformis</i> (Wulfen) J.V. Lamouroux, 1813	A M
<i>dalmatica</i> (Kützing) De Toni		M	<i>spinella</i> (C. Agardh) Kützing, 1847	A M
<i>vermicularis</i> (J. Agardh) F. Schmitz, 1889		A M	<i>valentiae</i> (Turner) Montagne, 1841	A M
<i>Schmitzia</i>			<i>Hypnocolax</i>	
<i>hiscockiana</i> Maggs & Guiry, 1985		A	<i>stellaris</i> Børgesen	A
<i>neapolitana</i> (Berthold) Lagerheim ex P.C. Silva, 1959	A M			
Family Caulacanthaceae			Family Nemastomataceae	
<i>Catenella</i>			<i>Itonoa</i>	
<i>caespitosa</i> (Withering) L.M. Irvine, 1976		A M	<i>marginifera</i> (J. Agardh) Masuda & Guiry, 1995	A M
<i>Caulacanthus</i>			<i>Nemastoma</i>	
<i>ustulatus</i> (Mertens ex Turner) Kützing, 1843		A M	<i>canariense</i> (Kützing) J. Agardh	A
<i>Feldmannophycus</i>			<i>confusum</i> Kraft & D.M. John	A
<i>rayssiae</i> (Feldmann & G. Feldmann) Augier & Boudouresque		A M	<i>constrictum</i> Ercegovic, 1949	M
Family Choreocolacaceae			<i>dichotomum</i> J. Agardh, 1842	A M
<i>Choreocolax</i>			<i>gelatinosum</i> M. Howe	A
<i>polysiphoniae</i> Reinsch, 1875		A	<i>Predaea</i>	
<i>Harveyella</i>			<i>feldmannii</i> Børgesen	A
<i>mirabilis</i> (Reinsch) F. Schmitz & Reinke, 1889		A	<i>huismanii</i> Kraft, 1984	A
Family Cruoriaceae			<i>masonii</i> (Setchell & N.L. Gardner) De Toni fil., 1936	A
<i>Cruoria</i>			<i>olivieri</i> Feldmann	A M
<i>cruoriaeformis</i> (P.L. Crouan & H.M. Crouan)		A M	<i>pusilla</i> (Berthold) Feldmann	A M
Denizot, 1968		A	<i>Tsengia</i>	
<i>pellita</i> (Lyngbye) Fries, 1835			<i>bardii</i> (Farlow) K.C. Fan & Y.P. Fan, 1962	A
Family Cystocloniaceae			Family Petrocelldaceae	
<i>Calliblepharis</i>			<i>Mastocarpus</i>	
<i>ciliata</i> (Hudson) Kützing, 1843		A M	<i>stellatus</i> (Stackhouse) Guiry, 1984	A
<i>jubata</i> (Goodenough & Woodward) Kützing, 1843		A M	Family Phyllophoraceae	
<i>Cystoclonium</i>			<i>Ahnfeltiopsis</i>	
<i>purpureum</i> (Hudson) Batters, 1902		A M	<i>devoniensis</i> (Greville) P.C. Silva & DeCew, 1992	A M
<i>Fimbrifolium</i>			<i>pusilla</i> (Montagne) P.C. Silva & De Cew, 1992	M
<i>dichotomum</i> (Lepechin) G. Hansen, 1980		A	<i>Ceratocolax</i>	
<i>Rhodophyllis</i>			<i>hartzii</i> Rosenvinge, 1898	A
<i>divaricata</i> (Stackhouse) Papenfuss, 1950		A M	<i>Coccotylus</i>	
<i>madeirensis</i> Levring, 1974		A	<i>truncatus</i> (Pallas) M.J. Wynne & J.N. Heine, 1992	A M
			<i>Erythrodermis</i>	
			<i>trailii</i> (Holmes ex Batters) Guiry & Garbary, 1990	A
			<i>Gymnogongrus</i>	
			<i>crenulatus</i> (Turner) J. Agardh, 1851	A M
			<i>griffithsiae</i> (Turner) Martius, 1828	A M
			<i>patens</i> (Goodenough & Woodward) J. Agardh, 1851	A M
			<i>tenuis</i> (Turner) J. Agardh	A
			<i>Phyllophora</i>	
			<i>crispa</i> (Hudson) P.S. Dixon, 1964	A M
			<i>fimbriata</i> Ercegovic, 1957	M
			<i>gelidiooides</i> P.L. Crouan & H.M. Crouan ex Karsakoff	A
			<i>heredia</i> (Clemente) J. Agardh	A M

<i>pseudoceranoïdes</i> (S.G. Gmelin) Newroth & A.R.A.			
<i>Taylor, 1971</i>	A M		
<i>sicula</i> (Kützing) Guiry & L.M. Irvine, 1976	A M		
<i>Schottera</i>			
<i>nicaeensis</i> (J.V. Lamouroux ex Duby) Guiry &			
<i>Hollenberg, 1975</i>	A M		
<i>Stenogramme</i>			
<i>interrupta</i> (C. Agardh) Montagne ex Harvey, 1848	A M		
Family Polyidaceae			
<i>Polyides</i>			
<i>rotundus</i> (Hudson) Greville, 1830	A M		
Family Rissoellaceae			
<i>Rissoella</i>			
<i>verruculosa</i> (Bertoloni) J. Agardh, 1848	M		
Family Sarcodiaceae			
<i>Chondrymenia</i>			
<i>lobata</i> (Meneghini) Zanardini	M		
Family Schizymeniaceae			
<i>Platoma</i>			
<i>cyclocarpa</i> (Montagne) F. Schmitz	A M		
<i>incrassata</i> Schousboe ex Bornet	A		
<i>Schizymenia</i>			
<i>dubyi</i> (Chauvin ex Duby) J. Agardh, 1851	A M		
Family Sebdeniaceae			
<i>Sebdenia</i>			
<i>dichotoma</i> Berthold	M		
<i>macaronesica</i> Soler-Onis, Haroun & Prud'homme van			
Reine	A		
<i>monardiana</i> (Montagne) Berthold	M		
<i>rodrigueziana</i> (Feldmann) P.G. Parkinson	M		
Family Solieriaceae			
<i>Agardhiella</i>			
<i>subulata</i> (C. Agardh) Kraft & M.J. Wynne, 1979	A		
<i>Meristotheca</i>			
<i>decumbens</i> Grunow	A		
<i>Rhabdonia</i>			
<i>decumbens</i> Grunow, 1888	A		
<i>Sarcodiotheca</i>			
<i>divaricata</i> W.R. Taylor	A		
<i>Solieria</i>			
<i>chordalis</i> (C. Agardh) J. Agardh, 1842	A M		
<i>dura</i> (Zanardini) F. Schmitz, 1895	M		
<i>Turnerella</i>			
<i>pennyi</i> (Harvey) F. Schmitz, 1893	A		
Family Wurdemanniaceae			
<i>Wurdemannia</i>			
<i>miniatia</i> (Sprengel) Feldmann & G. Hamel, 1934	A M		
Order Goniotrichales			
Family Goniotricaceae			
<i>Chroodactylon</i>			
<i>ornatum</i> (C. Agardh) Basson, 1979	A M		
<i>wolleanum</i> Hansgrig, 1885	A		
<i>Colacodium</i>			
<i>reticulatum</i> (Batters) Feldmann, 1955	A		
<i>Styloclonia</i>			
<i>alsidii</i> (Zanardini) K.M. Drew, 1956	A M		
<i>cornu-cervi</i> Reinsch, 1875	A M		
<i>subcoeruleum</i> (P.A. Dangeard) M.J. Wynne, 1985	A		
Order Gracilariales			
Family Gracilariaeae			
<i>Gracilaria</i>			
<i>arcuata</i> Zanardini, 1858	M		
<i>armata</i> (C. Agardh) Greville, 1830	A M		
<i>bursa-pastoris</i> (S.G. Gmelin) P.C. Silva, 1952	A M		
<i>cervicornis</i> (Turner) J. Agardh, 1852	A M		
<i>conferta</i> (Schousboe ex Montagne) Montagne, 1846	A M		
<i>corallicola</i> Zanardini, 1865	M		
<i>dendroides</i> Gargiulo, De Masi & Tripodi, 1985	M		
<i>dura</i> (C. Agardh) J. Agardh, 1842	A M		
<i>foliifera</i> (Forsskål) Børgesen, 1932	A M		
<i>gracilis</i> (Stackhouse) Steentoft, L.M. Irvine &			
Farnham, 1995	A M		
<i>heteroclada</i> (Montagne) Feldmann & G. Feldmann,			
1943			
<i>longa</i> Gargiulo, De Masi & Tripodi, 1987		M	
<i>multipartita</i> (Clemente) Harvey, 1846		A M	
<i>rubra</i> (C. Agardh) J. Agardh		A	
<i>vermiculata</i> P.J.L. Dangeard, 1949		A	
<i>Gracilariaopsis</i>			
<i>longissima</i> (S.G. Gmelin) Steentoft, L.M. Irvine &			
Farnham, 1995		A	
Family Pterocladiobiaceae			
<i>Gelidiocolax</i>			
<i>christianae</i> Feldmann & G. Feldmann, 1963	A M		
<i>deformans</i> Secoane, 1982	A		
<i>margaritoides</i> (Martin & Pocock) Fan & Papenfuss	A		
<i>microsphaericus</i> N.L. Gardner	A		
<i>Holmsella</i>			
<i>pachyderma</i> (Reinsch) Sturck, 1926	A		
Order Hildenbrandiales			
Family Hildenbrandiaceae			
<i>Hildenbrandia</i>			
<i>crouanii</i> J. Agardh, 1852	A M		
<i>occidentalis</i> Setchell, 1917	A		
<i>rivularis</i> (Liebmamn) J. Agardh, 1863	A		
<i>rubra</i> (Sommerfelt) Meneghini, 1841	A M		
Order Nemaliales			
Family Galaxauraceae			
<i>Galaxaura</i>			
<i>cyclindrica</i> (Solander) J.V. Lamouroux	A		
<i>obtusata</i> (J. Ellis & Solander) J.V. Lamouroux, 1816	A		
<i>rugosa</i> (J. Ellis & Solander) J.V. Lamouroux, 1816	A		
<i>Scinaia</i>			
<i>australis</i> (Setchell) Huisman, 1985	A		
<i>caribaea</i> (W.R. Taylor) Huisman	A		
<i>complanata</i> (Collins) A.D. Cotton, 1907	A M		
<i>furcellata</i> (Turner) J. Agardh, 1852	A M		
<i>interrupta</i> (De Candolle) M.J. Wynne, 1989	M		
<i>Tricleocarpa</i>			
<i>cylindrica</i> (J. Ellis & Solander) Huisman &			
Borowitzka, 1990	A		
<i>fragilis</i> (Linnaeus) Huisman & Townsend, 1993	M		
Family Liagoraceae			
<i>Cylindraxis</i>			
<i>rotundatus</i> Kraft		M	
<i>Dermoneema</i>			
<i>abbottiae</i> Afaf-Husain, Nizamuddin & Shameel	M		
<i>virens</i> (J. Agardh) Pedroche & Ávila Ortiz	M		
<i>Garonema</i>			
<i>farinosum</i> (J.V. Lamouroux) K.C. Fan & Y.C. Wang,			
1974	A M		
<i>lubrica</i> Afonso-Carrillo, Sansón & Reyes, 1998	A		
<i>Helminthocladia</i>			
<i>agardhiana</i> P.S. Dixon, 1962	A M		
<i>calvadosii</i> (J.V. Lamouroux ex Duby) Setchell, 1925	A		
<i>Helminthora</i>			
<i>divaricata</i> (C. Agardh) J. Agardh, 1852	A M		
<i>Liagora</i>			
<i>albicans</i> J.V. Lamouroux, 1816	A		
<i>canariensis</i> Børgesen, 1927	A		
<i>ceranoides</i> J.V. Lamouroux, 1816	A M		
<i>distenta</i> (Mertens ex Roth) J.V. Lamouroux, 1816	A M		
<i>divaricata</i> C.K. Tseng, 1941	A		
<i>fragilis</i> Zanardini, 1851	A		
<i>gymnarthon</i> Børgesen	A		
<i>maderensis</i> Kützing	A		
<i>tetrasporifera</i> Børgesen	A M		
<i>valida</i> Harvey, 1853	A		
<i>viscidia</i> (Forsskål) C. Agardh, 1822	A M		
<i>Nemalion</i>			
<i>helminthoides</i> (Velley) Batters, 1902	A M		
<i>Trichogloeopsis</i>			
<i>pedicellata</i> (Howe) I.A. Abbott & Doty, 1960	A		
Family Nacciariaceae			
<i>Atractophora</i>			

<i>hypnoides</i> P.L. Crouan & H.M. Crouan, 1848	A	<i>repens</i> (C. Agardh) Montagne & Bory	A M
<i>Naccaria</i>		<i>Gloiocladia</i>	
<i>corymbosa</i> J. Agardh, 1899	A	<i>atlantica</i> (Searles) R.E. Norris	A
<i>wiggii</i> (Turner) Endlicher, 1836	A M	<i>blomquistii</i> (Searles) R.E. Norris	A
Order Palmariales		<i>furcata</i> (C. Agardh) J. Agardh	M
Family Palmariaceae		<i>Halicrysis</i>	
<i>Devaleraea</i>		<i>depressa</i> (J. Agardh) F. Schmitz, 1889	A
<i>ramentacea</i> (Linnaeus) Guiry, 1982	A	<i>Leptofaucha</i>	
<i>Palmaria</i>	A	<i>brasiliensis</i> A.B. Joly	A
<i>palmata</i> (Linnaeus) Kuntze, 1891		Family Lomentariaceae	
Family Rhodophysemataceae		<i>Gelidiopsis</i>	
<i>Halosaccioncolax</i>		<i>intricata</i> (C. Agardh) Vickers, 1905	A
<i>kjellmanii</i> S. Lund, 1959	A	<i>planicalis</i> (W.R. Taylor) W.R. Taylor, 1960	A
<i>Meiodiscus</i>		<i>variabilis</i> (Greville ex J. Agardh) F. Schmitz, 1895	A
<i>spetsbergensis</i> (Kjellman) Saunders & McLachlan, 1991	A	<i>Lomentaria</i>	
Rhodophysema		<i>articulata</i> (Hudson) Lyngbye, 1819	A M
<i>elegans</i> (P.L. Crouan & H.M. Crouan ex J. Agardh)		<i>baileyan</i> (Harvey) Farlow, 1876	A
P.S. Dixon, 1964	A M	<i>chyoctadiella</i> Funk, 1955	M
<i>feldmannii</i> Cabioch, 1975	A	<i>claviformis</i> Ercegovic, 1956	M
<i>georgii</i> Batters, 1900	A M	<i>clavellosa</i> (Turner) Gaillon, 1828	A M
Family Rhodothamniellaceae		<i>compressa</i> (Kützing) Kylin, 1931	M
<i>Rhodothamniella</i>		<i>ercegovicia</i> Verlaque, Boudouresque, Meinesz, Giraud & Marcot Coqueugniot	M
<i>floridula</i> (Dillwyn) Feldmann, 1978	A	<i>firma</i> (J. Agardh) Falkenberg	M
Order Plocamiales		<i>jabukae</i> Ercegovic	M
Family Plocamiaceae		<i>orcadensis</i> (Harvey) Collins ex W.R. Taylor, 1937	A
<i>Plocamium</i>		<i>subdichotoma</i> Ercegovic, 1956	A M
<i>cartilagineum</i> (Linnaeus) P.S. Dixon, 1967	A M	<i>uncinata</i> Meneghini ex Zanardini	M
<i>raphelisianum</i> P.J.L. Dangeard, 1949	A M	<i>verticillata</i> Funk, 1944	M
Order Porphyridiales		Family Rhodymeniaceae	
Family Phragmonemataceae		<i>Asteromenia</i>	
<i>Goniotrichiopsis</i>		<i>peltata</i> (W.R. Taylor) Huisman & A.J.K. Millar	A
<i>sublittoralis</i> G.M. Smith, 1943	A	<i>Botryocladia</i>	
<i>Kyliniella</i>		<i>boergesenii</i> Feldmann, 1942	A M
<i>latvica</i> Skuja, 1926	A	<i>botryoides</i> (Wulff) Feldmann, 1941	A M
<i>Neevea</i>		<i>chiajeana</i> (Meneghini) Kylin	A M
<i>repens</i> Batters, 1900	A	<i>madagascariensis</i> G. Feldmann	M
Family Porphyridiaceae		<i>microphysa</i> (Hauck) Kylin, 1931	A M
<i>Chrootheca</i>		<i>occidentalis</i> (Børgesen) Kylin	A
<i>rupestris</i> Hansgirg	M	<i>pyriformis</i> (Børgesen) Kylin, 1931	A
<i>Porphyridium</i>		<i>wynnei</i> D.L. Ballantine	A
<i>aeruginosum</i> Geitler, 1923	A	<i>Chrysomenia</i>	
<i>purpureum</i> (Bory) Ross, 1965	A M	<i>bullosa</i> Levring, 1974	A
<i>Rhodella</i>		<i>enteromorpha</i> Harvey	A
<i>maculata</i> L. Evans, 1970	A	<i>ventricosa</i> (J.V. Lamouroux) J. Agardh, 1876	A M
<i>violacea</i> (Kornmann) Wehrmeyer, 1971	A	<i>Coelothrix</i>	
<i>Rhodosorus</i>		<i>irregularis</i> (Harvey) Børgesen, 1920	A
<i>marinus</i> Geitler, 1930	A	<i>Cordylecladia</i>	
<i>Rhodospora</i>		<i>erecta</i> (Greville) J. Agardh, 1852	A M
<i>sordida</i> Geitler, 1927	M	<i>guiryi</i> Gargiulo, G. Furnari & Cormaci, 1990	M
Order Rhodochaetales		<i>Irvinea</i>	
Family Rhodochaetaceae		<i>ardreana</i> (Brodie & Guiry) Guiry, 1999	A
<i>Rhodochaete</i>		<i>Rhodymenia</i>	
<i>parvula</i> Thuret	M	<i>ardissonei</i> J. Feldmann, 1941	A M
Order Rhodymeniales		<i>caespitosa</i> P.J.L. Dangeard, 1949	A
Family Champiaceae		<i>delicatula</i> P.J.L. Dangeard, 1949	A M
<i>Champia</i>		<i>holmesii</i> Ardisson, 1893	A
<i>parvula</i> (C. Agardh) Harvey, 1853	A M	<i>ligulata</i> Zanardini, 1843	M
<i>salicornoides</i> Harvey, 1853	A	<i>pertusa</i> (Postels & Ruprecht) J. Agardh, 1852	A
<i>vieillardii</i> Kützing, 1866	A	<i>pseudopalmaria</i> (J.V. Lamouroux) P.C. Silva, 1952	A M
<i>Chylocladia</i>		Order Sphaerococcales	
<i>pelagoae</i> Ercegovic	M	Family Sphaerococcaceae	
<i>verticillata</i> (Lightfoot) Bliding, 1928	A M	<i>Sphaerococcus</i>	
<i>Gastroclonium</i>		<i>coronopifolius</i> Stackhouse, 1797	A M
<i>clavatum</i> (Roth) Ardisson, 1883	A M	<i>rhizophylloides</i> J.J. Rodriguez	M
<i>ovatum</i> (Hudson) Papenfuss, 1944	A M	KINGDOM PLANTAE	
<i>reflexum</i> (Chauvin) Kützing, 1849	A M	PHYLUM MAGNOLIOPHYTA	
Family Faucheaceae		Class Liliopsida	
<i>Fauchea</i>		Order Hydrocharitales	
<i>microspora</i> Bornet	A M	Family Hydrocharitaceae	
		<i>Halophila</i>	
		<i>decipiens</i> Ostenfeld	A

<i>stipulacea</i> (Forsskål) Ascherson	M
Order Potamogetonales	
Family Cymodoceaceae	
<i>Cymodocea</i>	
<i>nodosa</i> (Ucria) Ascherson	A M
Family Posidoniaceae	
<i>Posidonia</i>	
<i>oceanica</i> (Linnaeus) Delile	A M
Family Zosteraceae	
<i>Zostera</i>	
<i>marina</i> Linnaeus	A M
<i>noltii</i> Hornemann	A M

PROTISTA

Compiled by Sue Brandt

Individual lists contributed by Sue Brandt, Anatol Wiktor Jankowski, Maria Mullisch, Sabine Agatha, Andrew Rogerson & Andrew Zubkov, Ralf Meisterfeld, Onno Gross, Michael Dick, Ole Tendal and Jacob van der Land.

Protists are ubiquitous in marine waters, freshwater and soil. They gave rise to the plants, animals and fungi, and contain all eukaryotes not included within those kingdoms. Protists (previously artificially separated into "algae", "protozoa" and certain fungal-like groups) are typically single-celled and between 1 and 500 µm in size, although the largest are up to 10⁵ µm (Andersen 1998). Many species appear to have a global distribution (Finlay *et al.* 1999) and some marine species also occur in fresh water or soil.

Nutritional modes vary from photosynthesis, to the ingestion of prey, to the absorption of organic molecules, and some protists are mixotrophic, carrying out both photosynthesis and ingestion or absorption. Almost 100% of the primary production in marine systems is carried out by protists and prokaryotic microorganisms (cyanobacteria), and these groups account for some 40% of global primary production (Andersen 1998). Protists are also important in biogeochemical cycles. By forming "marine snow" they contribute to the effect of the ocean as a "carbon sink", reducing the rate at which atmospheric carbon-dioxide levels are rising. Fossilised species of Foraminifera are used in oil prospecting, and these and other shelled species are the constituents of rocks such as limestone and chalk.

During the last 40 years, the use of electron microscopy and molecular biology has led to a demolition of the traditional view of the classification of protists. During the phase of reconstruction, many overlapping or contradictory schemes have been proposed, leading to confusion. Despite this, there is confidence that many species have now been assigned to monophyletic assemblages and the lists included in this Register reflect some of those groups. Those interested in the current state of knowledge are referred to Patterson (1999) who lists 71 protist lineages without identifiable sister taxa. As a complete modern classification of the protists does not exist, taxa included in the Register have been given rankless colloquial names – the terms being familiar to those currently working with protists.

Protists were outside the original scope of the ERMS project, and it has only been possible to include a small proportion of taxa in the time-scale available. However, it is hoped that the number of taxa included in the Register will be increased in the future, in recognition of their importance in marine ecosystems. The global nature of many taxa means that these "European" lists also have an application in other regions.

References

- Andersen, R.A. 1998. What to do with Protists? *Australian Systematic Botany*, 11: 185-201.
- Finlay, B.J., Esteban, G.F., Olmo, J.L. & Tyler, P.A. 1999. Global distribution of free-living microbial species. *Ecography* 22: 138-144.
- Patterson, D.J. 1999. The Diversity of Eukaryotes. The *American Naturalist* 154 (supplement), pS97-S124.

Alveolates

Apicomplexa (free-living species)

Compiled by Sue Brandt

Comprehensive list for free-living taxa only. There are many parasitic taxa.

References

- Droop, M.R. 1953 On the ecology of flagellates from some brackish water and fresh water rockpools of Finland. *Acta Bot. Fennica* 51: 1-52
- Simpson, A.B.G & Patterson, D.J. 1996 Ultrastructure and identification of the predatory flagellate *Colpodella pugnaxi* Cienkowski (Apicomplexa) with a description of *Colpodella turpis* n. sp. and a review of the genus *Systematic Parasitology*, 33: 187-198
- Tong, S.M. 1997b Heterotrophic flagellates and other protists from Southampton Water, U.K. *Ophelia* 47: 71-131.

Apicomplexa, free-living

- Colpodella perforans* (Hollande, 1938) Patterson & Zöllfel, 1991
= *Bodo perforans* Hollande, 1938
= *Spiromonas perforans* (Hollande, 1938) Brugerolle & Mignot, 1979
- edax* (Klebs, 1892) Simpson & Patterson, 1996
= *Bodo edax* Klebs, 1892
= *Colpodella pugnax* sensu Dangeard 1900 non Cienkowski 1865
= *Nephromonas hyalina* Droop, 1953
- Perkinsus atlanticus* Azevedo, 1989

A

Ciliata (Phylum Ciliophora)

Chonotrichia

Compiled by Anatol Wiktor Jankowski

A comprehensive list of European species.

References

- Batisse, A. 1994 Sous-classe des Chonotrichia. Puytorac P. De, ed. *Traité de Zoologie*, t. 2, fasc. 2, Masson, Paris, 433-473
- Batisse, A. & Crumeyrolle, J.-B. 1988 Sur deux chonotriches nouveaux de l'Archipel Kerguelen (Ciliophora, Chonotrichida) *C. R. Acad. Sci. Paris*, série III, 307: 635-640
- Canu, E. 1886 Note sur le genre *Spirochona* (Stein) *Bull. Scient. Dept. du Nord et des Pays Voisins*, 17 (=Bull. Biol. France Belg., ser. 2, t. 9): 21-31
- Doflein, F. J. Th. 1897a Studien zur Naturgeschichte der Protozoen. I. *Kentrochona nebuliae* Rompel *Zool. Jahrb., Abth. Anat. Ontog. Thiere*, 10: 619-641
- Doflein, F. 1897b Studien zur Naturgeschichte der Protozoen. II. *Kentrochonopsis multipara* n. g., n. sp., ein Ciliat mit multipler Knospung *Zool. Jahrb., Abth. Anat. Ontog. Thiere*, 10: 642-646
- Dons, C. 1940 Zweie neue Ciliaten. Det Kongel. Norske Videnskabers Selskab. *Forhandl.*, 13: 115-128
- Dovgal, I. V. 1983 *Heliochona pontica* (Ciliophora, Chonotrichida) in the sea of Azov. *Vestnik Zoologii* (Kiev), yr. 1983: 72-74
- Dovgal, I. V. 1999 The distribution and variability of chonotrichs (Ciliophora, Chonotrichia) of the Ukrainian fauna. Report 2. *Heliochona pontica* Jankowski, 1973. *Vestnik Zoologii* (Kiev), yr. 1999, in press
- Fenchel, T. 1965 On the ciliate fauna associated with the marine species of the amphipod genus *Gammarus* J. G. Fabricius. *Ophelia*, 2: 281-303
- Giard, A. 1894 Contribution à la faune du Pas de Calais et de la Manche. *C. R. Soc. Biol. Paris*, ser. X, 1: 245-247
- Jankowski, A. W. 1971 Commensological notes. I. Chonotrichs (Ciliata) on epibionts of Cetacea. *Zool. Zh.*, 50: 1445-1456

- Jankowski, A. W. 1972 Commensological sketches. 3. New genera of Chonotricha, Exogemmina, symbiotic with Isopoda, Amphipoda and Decapoda. *Zool. Zh.*, 51: 1617-1524
- Jankowski, A. W. 1973 Commensological sketches. 4. New genera of Chonotricha Endogemmina symbiotic with Leptostraca. *Zool. Zh.*, 52: 15-24
- Jankowski, A. W. 1973 Subclass Chonotricha. In: *Fauna of the USSR, Infusoria*, 2(1) (=New series, No. 103), Nauka, Leningrad: 1-356
- Jankowski, A. W. 1989 *Segmentochona* gen. n. (Chonotricha), symbiont of chelurids (Amphipoda). Sveshnikov V. A., ed. *Symbiosis in marine animals*. IEMEZ, Moscow: 7-13
- Kahl, A. 1933 Ciliata libera et ectocommensalia. Grimpe G. & Wagler E., *Die Tierwelt der Nord- und Ostsee*. Lief. 23, Teil II-C-3, Leipzig: 29-146
- Kahl, A. 1935 Urtiere oder Protozoa. I. Wimpertiere oder Ciliata (Infusoria). Dahl, F., ed. *Die Tierwelt Deutschlands*. Jena: 3: 799-805
- Kent W. S. 1881 *A manual of the Infusoria: including description of all known flagellate, ciliate and tentaculiferous Protozoa, British and foreign*. D. Bogue, London. Vol. 2: 660-664. Vol. 3, Atlas, pl. 33
- Matsudo, H. & Mohr, J. L. 1965 A new species of *Heliochona*, *H. psychra* n. sp. (Ciliata: Chonotrichida) from Nuuvuk Lake, Alaska. *J. Protozool.*, 12: 522-527
- Mohr, J. L., Le Veque, J. A. & Matsudo, H. 1963 On a new collar ciliate of a gribble: *Lobochona prorates* n. sp. on *Limnoria tripunctata*. *J. Protozool.*, 10: 226-233
- Mohr, J. L., Matsudo, H. & Leung, Y. M. 1963 A ciliate protozoan of whale lice. *Proc. XVI Intern. Congr. Zool.*, Washington, 1: 121
- Mohr, J. L., Matsudo, H. & Leung, Y. M. 1970 The ciliate taxon Chonotricha. Barnes H., ed., *Oceanography and marine biology*. G. Allen & Unwin, L.: 8: 415-456
- Plate, L. H. 1888 Protozoen-Studien. Habilitationsschrift. Über die verschiedenen Formen der Fortpflanzung im Thierreiche. V. *Heliochona sessilis* mihi, eine neue Vorticelline. *Zool. Jahrb., Abth. Anat. Ontog. Thiere*, 3: 135-198
- Precht, H. 1935 Epizoen der Kieler Bucht. *Nova Acta Leopoldina Carol.* = Abhandl. Kaiserl. Leopoldinisch-Carol. Deutsche Akad. der Naturforsch., N. F., 3: 405-474
- Rompel, J. 1894 *Kentrochona nebaliae* n. g., n. sp. Ein neues Infusor aus der Familie der Spirochoninen, zugleich ein Beitrag zur Lehre von der Kerntheilung und dem Centrosoma. *Z. Wiss. Zool.*, 58: 618-635
- Small, E. B. & Lynn, D. H. 1985 Phylum Ciliophora Doflein, 1901. Lee, J. J., Hutner, S. H. & Bovee, E. C., ed., *An illustrated guide to the Protozoa. Society of Protozoologists*, Lawrence, Kansas: 393-575
- Wallengren, H. 1895 Studier of ver Ciliata Infusorier. II. Slaget *Heliochona* Plate, slaget *Chilodochona* n. g., slaget *Hemispeira* Fabre-Domergue. *Lunds Universitets Arsskrift* (=Acta Univ. Lund, 1; = Acta Reg. Soc. Physiogr. Lunds, 6), 31: 1-77
- Chonotrichia**
- Order Exogemmida**
- Suborder Lobochonina**
- Family Loboconidae**
- Lobochna limnoriae* Dons, 1940
prorates Mohr, Le Veque & Matsudo, 1963
= *Spirochona crystallina* Canu, 1886
- Family Filichonidae**
- Aurichona armata* Jankowski, 1973
- Family Phyllochonidae**
- Phyllochona vexillata* Jankowski, 1972
- Family Segmentochonidae**
- Segmentochona spiralis* (Bonnier in Giard, 1894) Jankowski, 1989
= *Stylochona spiralis* Bonnier in Giard, 1894
- Family Heliochonidae**
- Heliochona alternans* Jankowski, 1972
elegans Jankowski, 1972
pontica Jankowski, 1973
sessilis Plate, 1888
- Family Heterochonidae**
- coronata* (Kent, 1881) Jankowski, 1972
= *Stylochona spiralis* subsp. *coronata* Kent, 1881
crinita Jankowski, 1972
exilis Jankowski, 1973
peregra Jankowski, 1973
psychra (Matsudo & Mohr, 1965) Jankowski, 1972
= *Heliochona psychra* Matsudo & Mohr, 1965
scheutennii (Stein, 1854) Jankowski, 1972
= *Spirochona scheutennii* Stein, 1854
trifaria Jankowski, 1972
- Suborder Chilodochonina**
- Family Chilodochonidae**
- Chilodochona aranei* Jankowski, 1973
carcini Jankowski, 1973
palinuri Jankowski, 1973
quvennerstedti Wallengren, 1895
= *Chilodochona microchilus* Wallengren, 1895
- Vasichona paguri* Jankowski, 1972
- Order Cryptogemmida**
- Family Isochonidae**
- Cyamichona megapterae* Jankowski, 1971
mysticeti Jankowski, 1971
- Isochona crassa* Jankowski, 1973
tisbica Jankowski, in press
- Family Actinichonidae**
- Cristichona tessellata* Jankowski, 1973
Kentrochonopsis multipara Doflein, 1897
- Family Stylochonidae**
- Armichona hexadenta* Jankowski, 1973
insecta Jankowski, 1973
- Eriochona horrida* Jankowski, 1973
- Flectichona crenata* Jankowski, 1973
- Stylochona spiralis* subsp. *glabra* Jankowski, 1973
spiralis subsp. *nebalina* Kent, 1881
- Family Inversochonidae**
- Ceratochona scandens* Jankowski, 1973
- Chonosaurus galeatus* Jankowski, 1973
- Inversochona peregra* Jankowski, 1973
- Kentrochona nebaliae* Rompel, 1894
- Folliculinidae**
- Compiled by Maria Mulisch
- A partial list of European species.
- Folliculinidae**
- Aulosfolliculina labyrinthica* Hadzi, 1951
Diagfolliculina rotunda Hadzi, 1951

M

M

<i>similis</i> Hadzi, 1951	M	(Ciliophora, Oligotrichidea) from coastal brackish water basins of Germany. - <i>Europ. J. Protistol.</i> 34: 10-17.
<i>Donsia</i>	M	
<i>mirabilis</i> (Dons, 1917) Hadzi, 1951	M	Agatha, S. 1998 Morphology and infraciliature of some planktonic ciliates (Ciliophora) from coastal waters of Schleswig-Holstein (Northern Germany). - <i>J. Euk. Microbiol.</i> 45, Abstract 46, 9A.
<i>Epifolliculina</i>	M	
<i>diaphana</i> Hadzi, 1951	M	
<i>Eufolliculina</i>	M	Auf dem Venne, H. 1994 Zur Verbreitung und ökologischen Bedeutung planktischer Ciliaten in zwei verschiedenen Meeresgebieten: Grönlandsee und Ostsee. <i>Ber. Inst. Meereskd.</i> 262: 1-160.
<i>ampullacea</i> (Laakmann, 1910) Hadzi, 1951	M	
<i>baltica</i> Sjögren, 1964	A	Biemacka, I. 1963 Die Protozoenfauna in Danziger Bucht II. Die Charakteristik der Protozoen in Untersuchten Biotopen der Seekuste. - <i>Polskie Archiwum Hydrobiol.</i> 11: 17-75.
<i>latemarginata</i> Sahrhage, 1916	A	Busch, W. 1921 Studien über Ciliaten des Nordatlantischen Ozeans und Schwarzen Meeres. - <i>Arch. Protistenk.</i> 42: 364-379.
<i>moebiusi</i> (Möbius, 1887) Kahl, 1932	A	Dragesco, J. 1960 Ciliés mésopsammiques littoraux, systématique, morphologie, écologie. - <i>Trav. Stn biol. Roscoff</i> 12: 1-356.
<i>uhligi</i> Mulisch & Patterson, 1983	A	Entz, G. 1884 Über Infusorien des Golfs von Neapel. - <i>Mitt. zool. Stn Neapel</i> 5: 289-444.
<i>Folliculinopsis</i>	A M	Fauré-Fremiet, E. & Ganier, M.-Cl. 1970 Structure fine du <i>Strombidium sulcatum</i> Cl. et L. (Ciliata Oligotrichida). - <i>Protistologica</i> 6: 207-223.
<i>annulata</i> Faure-Fremiet, 1936	A M	Fauré-Fremiet, E. 1924 Contribution à la connaissance des infusoires planctoniques. - <i>Bull. biol. Fr. Belg.</i> , Suppl. 6: 1-171.
<i>Lagotia</i>	M	Fauré-Fremiet, E. 1932 <i>Strombidium calkinsi</i> , a new thigmotactic species. - <i>Biol. Bull.</i> 62: 201-204.
<i>donsi</i> Hadzi, 1951	M	Fauré-Fremiet, E. 1969 Remarques sur la systématique des ciliés Oligotrichida. - <i>Protistologica</i> 5: 345-352.
<i>expansa</i> (Levinson, 1893) Hadzi, 1951	A M	Gruber, A. 1884 Die Protozoen des Hafens von Genua. - <i>Nova Acta Leop. Carol.</i> 46: 289-444.
<i>faurefremieti</i> Hadzi, 1951	A	Hartwig, E. 1974 Verzeichnis der im Bereich der deutschen Meeresküste angetroffenen interstitiellen Ciliaten. - <i>Mitt. zool. Mus. Hamb.</i> 71: 7-21.
<i>lutea</i> (Faure-Fremiet, 1936) Hadzi, 1951	A	Hovasse, R. 1932 Trois infusoires planctoniques du Bosphore. - <i>Arch. Zool. Exp. Gén. Paris</i> , Notes and Revue 73: 1-8.
<i>minima</i> Hadzi, 1951	M	Kahl, A. 1932 Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 3. Spirotricha. - <i>Tierwelt Dtl.</i> 25: 399-650.
<i>minor</i> (Dons, 1948) Hadzi, 1951	A	Leegaard, C. 1915 Untersuchungen über einige Planktonciliaten des Meeres. - <i>Nytt Mag. Naturvid.</i> 53: 1-37.
<i>obstretrica</i> (Wright, 1862) Hadzi, 1951	A	Lohmann, H. 1908 Untersuchungen zur Feststellung des vollständigen Gehaltes des Meeres an Plankton. - <i>Wiss. Meeresunters.</i> , Abt. Kiel 10: 129-370.
<i>spirorbis</i> (Dons, 1912) Hadzi, 1951	A	Lynn, D.H. & Gilron, G.L. 1993 Strombidiid ciliates from coastal waters near Kingston Harbour, Jamaica (Ciliophora, Oligotrichia, Strombidiidae). - <i>J. mar. biol. Ass. U.K.</i> 73: 47-65.
<i>viridis</i> Wright, 1858	A	Lynn, D.H. & Montagnes, D.J.S. 1988 Taxonomic descriptions of some conspicuous species of strobiliid ciliates (Ciliophora: Choretrichida) from the Isles of Shoals, Gulf of Maine. - <i>J. mar. biol. Ass. U.K.</i> 68: 639-658.
<i>Metafolliculina</i>	A	Lynn, D.H., Montagnes, D.J.S. & Small, E.B. 1988 Taxonomic descriptions of some conspicuous species in the family Strombidiidae (Ciliophora: Oligotrichida) from the Isles of Shoals, Gulf of Maine. - <i>J. mar. biol. Ass. U.K.</i> 68: 259-276.
<i>andrewsi</i> Hadzi, 1938	A	Lynn, D.H., Montagnes, D.J.S., Dale, T., Gilron, G.L. & Strom, S.L. 1991 A reassessment of the genus <i>Strombidinopsis</i> (Ciliophora, Choretrichida) with description of four new planktonic species and remarks on its taxonomy and phylogeny. - <i>J. mar. biol. Ass. U.K.</i> 71: 597-612.
<i>ballerina</i> Mulisch <i>et al.</i> , 1986	A	Maeda, M. & Carey, P.G. 1985 An illustrated guide to the species of the family Strombidiidae (Oligotrichida, Ciliophora), free swimming protozoa common in the aquatic environment. - <i>Bull. Ocean Res. Inst., Univ. Tokyo</i> 19: 1-68.
<i>nordgardi</i> Dons, 1924	A	Maeda, M. 1986 An illustrated guide to the species of the families Halteriidae and Strobiliidae (Oligotrichida, Ciliophora), free swimming protozoa common in the aquatic environment. - <i>Bull. Ocean Res. Inst., Univ. Tokyo</i> 21: 1-67.
<i>Parafolliculina</i>	M	
<i>glutinata</i> Hadzi, 1951	M	
<i>violacea</i> (Giard, 1888) Hadzi, 1951	A M	
<i>Planifolliculina</i>	M	
<i>cumbens</i> Hadzi, 1951	M	
<i>Platyfolliculina</i>	M	
<i>sahrhageana</i> (Sahrhage, 1917) Hadzi, 1938	M	
<i>Splitofolliculina</i>	M	
<i>adherens</i> Hadzi, 1951	M	
<i>Splitoiculina</i>	M	
<i>longicollis</i> Hadzi, 1951	M	
<i>Stentofolliculina</i>	M	
<i>tubicola</i> Hadzi, 1938	M	

Oligotrichaea (alorate species)

Compiled by Sabine Agatha

The alorate oligotrichs are a highly abundant, widely distributed and ecologically important group of ciliates (Protozoa, Ciliophora) in the marine environment. However, identification is difficult due to their fragility, small body size, similar body shape and the minute differences in the ciliary pattern only recognisable after time-consuming silver preparations. The determination of live specimens also needs very experienced ciliate taxonomists, whilst reliable identifications from Lugol-preserved material are almost impossible. Therefore, the following list of alorate oligotrich ciliates is comprised only of the species which were originally described or re-described from the area under consideration, or were mentioned by a reliable reference, namely, based on silver-impregnated material investigated by an experienced ciliate taxonomist. Bare species lists based on Lugol-preserved or live material were not included due to the rather doubtful species identifications. The classification of the alorate oligotrichs follows, for the sake of simplicity, Petz & Foissner (1992).

References

- Agatha, S. & Riedel-Lorjé, J.C. 1997 Morphology, infraciliature, and ecology of halteriids and strombidiids (Ciliophora, Oligotrichida) from coastal brackish water basins. - *Arch. Protistenk.* 148: 445-459.
- Agatha, S. & Riedel-Lorjé, J.C. 1998 Morphology, infraciliature, and ecology of some strobiliid ciliates

- Martin, A.J. & Montagnes, D.J.S. 1993 Winter ciliates in a British Columbian fjord: six new species and an analysis of ciliate putative prey. - *J. Euk. Microbiol.* 40: 535-549.
- Meunier, A. 1910 *Campagne arctique de 1907. Microplancton des Mers de Barents et de Kara*. - Bulens, Bruxelles.
- Montagnes, D.J.S. & Taylor, F.J.R. 1994 The salient features of five marine ciliates in the class Spirotrichea (Oligotrichia), with notes on their culturing and behaviour. - *J. Euk. Microbiol.* 41: 569-586.
- Montagnes, D.J.S., Lynn, D.H., Stoecker, D.K. & Small, E.B. 1988 Taxonomic descriptions of one new species and redescription of four species in the family Strombidiidae (Ciliophora, Oligotrichida). - *J. Protozool.* 35: 189-197.
- Montagnes, D.J.S., Taylor, F.J.R. & Lynn, D.H. 1990 *Strombidium inclinatum* n. sp. and a reassessment of *Strombidium sulcatum* Claparède and Lachmann (Ciliophora). - *J. Protozool.* 37: 318-323.
- Petz, W. & Foissner, W. 1992 Morphology and morphogenesis of *Strobilidium caudatum* (Fromentel), *Meseres corlissi* n. sp., *Halteria grandinella* (Müller), and *Strombidium rehwaldi* n. sp., and a proposed phylogenetic system for oligotrich ciliates (Protozoa, Ciliophora). *J. Protozoology* 39 (1): 159-176.
- Petz, W., Song, W. & Wilbert, N. 1995 Taxonomy and ecology of the ciliate fauna (Protozoa, Ciliophora) in the endopagial and pelagic of the Weddell Sea, Antarctica. - *Stapfia* (Linz) 40: 1-223.
- Rassoulzadegan, F. 1977 Évolution annuelle des ciliés pélagiques en Méditerranée nord-occidentale ciliés oligotriches "non tintinnides" (Oligotrichina). - *Annls Inst. océanogr.*, Paris 53: 125-134.
- Song, W. & Bradbury, P.C. 1998 Studies on some new and rare reported marine planktonic ciliates (Ciliophora: Oligotrichia) from coastal waters in North China. - *J. mar. biol. Ass. U.K.* 78: 767-794.
- Song, W. & Packhoff, G. 1996/97 Taxonomische Untersuchungen an marinen Ciliaten aus China mit Beschreibung von zwei neuen Arten, *Strombidium globosaneum* nov. spec. und *S. platum* nov. spec. (Protozoa, Ciliophora). - *Arch. Protistenk.* 147: 331-360.
- TRIPOS (ed.) 1997 Tempelmann, D. & Agatha, S.: Biomonitoring van microzooplankton in de Nederlandse zoute wateren. - *Rijkswaterstaat, Rijksinstituut voor Kust en Zee* (RIKZ), No. 97.T0017-1b.
- Tuolesco, J. 1962 Études protozoologiques sur les eaux roumaines I. Espèces nouvelles d'infusoires de la Mer Noire et des bassins salés paramarins. - *Arch. Protistenk.* 106: 1-36.
- Wulff, A. 1919 Ueber das Kleinplankton der Barentssee. - *Wiss. Meeresunters., Abt. Helgoland* 13: 95-124.
- Oligotrichida, aloricate**
- Family Halteriidae**
- Halteria*
- chlorelligera* Kahl, 1932
 - = *H. grandinella* (Müller, 1773) Dujardin, 1841 var. *chlorelligera* Kahl
- Pelagoalteria*
- cirrifera* (Kahl, 1932) Foissner, Skogstad & Pratt, 1988
 - = *Halteria cirrifera* Kahl, 1932
 - = *Halteria grandinella* (Müller, 1773) Dujardin, 1841 var. *cirrifera* Kahl
- Family Strombidiidae**
- Laboea*
- strobila* Lohmann, 1908 A M
 - = *Conocylis helix* Meunier, 1910
 - = *Strombidium strobilum* (Lohmann, 1908) Wulff, 1919
 - = *Strombidium strobilus* (Lohmann, 1908) Wulff, 1919
- Metastrombidium*
- sonnifer* Jankowski, 1980 A
- Novistrombidium*
- testaceum* (Anigstein, 1914) Song & Bradbury, 1998 M
 - = *Strombidium testaceum* Anigstein, 1914
- Spirostrombidium*
- cinctum* (Kahl, 1932) Petz *et al.*, 1995
- = *Strombidium cinctum* Kahl, 1932
- elegans* (Florentin, 1901) Petz *et al.*, 1995
- = *Strombidium elegans* Florentin, 1901
- pulchrum* (Leegaard, 1915) Agatha, 1998
- = *Laboea pulchra* Leegaard, 1915
- = *Strombidium pulchrum* (Leegaard, 1915) Kahl, 1932
- sauerbreyae* (Kahl, 1932) Petz *et al.*, 1995
- = *Strombidium coronatum* Sauerbrey, 1928
- = *Strombidium sauerbreyae* Kahl, 1932
- oblongum* (Entz, 1884) Petz *et al.*, 1995 M
- = *Clypeolum corsicum* Gourret & Roeser, 1888
- = *Strombidium oblongum* (Entz, 1884) Kahl, 1932
- = *Strombidium sulcatum* Claparède & Lachmann, 1859 var. *oblongum* Entz
- Strombidium*
- acutum* Leegaard, 1915 A
 - arenicola* Dragesco, 1960 A
 - atlanticum* (Busch, 1921) Maeda & Carey, 1985 A
 - = *Buehringa atlantica* Busch, 1921
 - = *S. atlantica* (Busch, 1921) Maeda & Carey, 1985
 - bilobum* Lynn & Gilron, 1993
 - buehringae* Busch, 1921 A M
 - capitatum* (Leegaard, 1915) Kahl, 1932
 - = *Laboea capitata* Leegaard, 1915
 - clavellinae* Buddenbrock, 1922
 - conicoides* (Leegaard, 1915) Kahl, 1932 A M
 - = *Woodania conicoides* Leegaard, 1915
 - constrictum* (Meunier, 1910) Wulff, 1919 A
 - = *Conocylis constricta* Meunier, 1910
 - = *Laboea constricta* (Meunier, 1910) Leegaard, 1915
 - cornutum* (Leegaard, 1915) Kahl, 1932 A M
 - = *Laboea cornuta* Leegaard, 1915
 - costatum* Tucolesco, 1962
 - cylindromorphum* Perejaslawzewa, 1886
 - datum* Lynn *et al.*, 1988
 - elongatum* Leegaard, 1915
 - = *Laboea conica* Lohmann, 1908 var. *elongatum* Leegaard A
 - faurei* Dragesco, 1960
 - filiforme* Kahl, 1932 A
 - fourneti* Dragesco, 1960
 - = *S. sauerbreyae* Kahl, 1932 var. *fourneti* Dragesco
 - grande* Levander, 1894
 - kahli* Bock, 1952
 - kielum* Maeda & Carey, 1985
 - lagenula* Fauré-Fremiet, 1924 A
 - latum* Kahl, 1932
 - macronucleatum* Dragesco, 1960
 - minor* (Kahl, 1935) Maeda & Carey, 1985
 - = *S. calkinsi* Kahl, 1932
 - = *S. caudatum* Calkins, 1901
 - = *S. stylifer* Levander, 1894 var. *minor* Kahl
 - oblongum* Leegaard, 1915
 - opisthostomum* Tucolesco, 1962
 - purpureum* Kahl, 1932
 - reticulatum* (Leegaard, 1915) Busch, 1921 A
 - = *Laboea reticulata* Leegaard, 1915
 - schizostomum* Kahl, 1932
 - spirale* Busch, 1950 A
 - stylifer* Levander, 1894
 - sulcatum* Claparède & Lachmann, 1859 A M
 - = *S. minutum* Wulff, 1919
 - tintinnodes* Entz, 1884 M
 - = *Strobilidium elegans* Hada, 1970
 - = *Strobilidium oculatum* (Gruber, 1884) Kahl, 1932
 - = *S. hadai* (Hada, 1970) Maeda & Carey, 1985
 - = *S. oculatum* Gruber, 1884
 - = *S. typicum* (Lankester, 1874) Bütschli, 1889
 - triquetrum* Agatha & Riedel-Lorjé, 1997
 - turcicum* Hovasse, 1932
 - urceolare* Stein, 1867
 - vestitum* (Leegaard, 1915) Kahl, 1932 A
 - = *Laboea delicatissima* Leegaard, 1915
 - = *Laboea vestita* Leegaard, 1915

= <i>S. delicatissimum</i> (Leegaard, 1915) Busch, 1921			
<i>virgatum</i> Wulff, 1919			
<i>wulffii</i> Kahl, 1932			
= <i>S. striatum</i> Wulff, 1919			
<i>compressum</i> (Leegaard, 1915) Kahl, 1932	A M		
= <i>Laboea compressa</i> Leegaard, 1915			
<i>conicum</i> (Lohmann, 1908) Wulff, 1919			
= <i>Laboea acuminata</i> Leegaard, 1915	A		
= <i>Laboea conica</i> Lohmann, 1908			
= <i>S. acuminatum</i> (Leegaard, 1915) Kahl, 1932	A M		
<i>coronatum</i> (Leegaard, 1915) Kahl, 1932			
= <i>Laboea coronata</i> Leegaard, 1915	A		
<i>crassulum</i> (Leegaard, 1915) Kahl, 1932			
= <i>Laboea crassula</i> Leegaard, 1915			
<i>emergens</i> (Leegaard, 1915) Kahl, 1932	A M		
= <i>Laboea emergens</i> Leegaard, 1915			
<i>obliquum</i> Kahl, 1932			
<i>ovale</i> (Leegaard, 1915) Kahl, 1932			
= <i>Laboea ovalis</i> Leegaard, 1915			
<i>Tontonia</i>			
<i>appendiculariformis</i> Fauré-Fremiet, 1924	A		
<i>caudata</i> (Lohmann, 1908) Kahl, 1932			
= <i>Strombidium caudatum</i> Lohmann, 1908			
<i>gracillima</i> Fauré-Fremiet, 1924	A		
= <i>Paratontonia gracillima</i> (Fauré-Fremiet, 1924)			
Jankowski, 1978			
= <i>Strombidium gracillimum</i> (Fauré-Fremiet, 1924)			
Alekperov & Mamajeva, 1992			
<i>simplicidens</i> Lynn & Gilron, 1993			
Family Strobilidiidae			
<i>Leegaardiella</i>			
<i>ovalis</i> Lynn & Montagnes, 1988			
<i>sol</i> Lynn & Montagnes, 1988			
<i>Lohmanniella</i>	M		
<i>oviformis</i> Leegaard, 1915			
<i>Parastrombidium</i>			
<i>faurei</i> (Kahl, 1932) Maeda, 1986			
= <i>Strobilidium faurei</i> Kahl, 1932			
<i>Pelagostrobilidium</i>			
<i>neptuni</i> (Montagnes & Taylor, 1994) Petz et al., 1995			
= <i>Strobilidium neptuni</i> Montagnes & Taylor, 1994			
<i>spiralis</i> (Leegaard, 1915) Petz et al., 1995	M		
= <i>Lohmanniella spiralis</i> Leegaard, 1915			
= <i>Strobilidium spiralis</i> (Leegaard, 1915) Lynn & Montagnes, 1988			
<i>Rimostrombidium</i>			
<i>caudatum</i> (Kahl, 1932) Agatha & Riedel-Lorjé, 1998			
= <i>R. kahli</i> (Petz & Foissner, 1992) Petz & Foissner, 1992			
= <i>Strobilidium caudatum</i> Kahl, 1932			
= <i>Strobilidium kahli</i> Petz & Foissner, 1992			
<i>conicum</i> (Kahl, 1932) Petz & Foissner, 1992			
= <i>Strobilidium conicum</i> Kahl, 1932			
<i>multinucleatum</i> (Lynn & Montagnes, 1988) Petz et al., 1995			
= <i>Strobilidium multinucleatum</i> Lynn & Montagnes, 1988			
<i>undinum</i> (Martin & Montagnes, 1993) Petz et al., 1995			
= <i>Strobilidium undinum</i> Martin & Montagnes, 1993			
<i>veniliae</i> (Montagnes & Taylor, 1994) Petz et al., 1995			
= <i>Strobilidium veniliae</i> Montagnes & Taylor, 1994			
<i>Strobilidium</i>			
<i>elegans</i> (Wulff, 1919) Maeda, 1986			
= <i>Lohmanniella elegans</i> (Wulff, 1919) Kahl, 1932			
= <i>Sphaerotrichium elegans</i> Wulff, 1919			
<i>marinum</i> (Fauré-Fremiet, 1910) Fauré-Fremiet, 1924	A		
= <i>Strombidium marinum</i> Fauré-Fremiet, 1910			
<i>pelagicum</i> Fauré-Fremiet, 1924	A		
= <i>Torquatella pelagica</i> (Fauré-Fremiet, 1924) Fauré-Fremiet, 1969			
<i>proboscidiiformum</i> (Milne, 1886) Kahl, 1932			
= <i>Strombidinopsis proboscidiiformum</i> Milne, 1886			
<i>striatum</i> (Meunier, 1910) Kahl, 1933			
= <i>Conocylis striata</i> Meunier, 1910			
= <i>Strombidium striatum</i> (Meunier, 1910) Kahl, 1932			
<i>sulcatum</i> Tucolesto, 1962			
<i>tonsuratum</i> (Meunier, 1910) Maeda, 1986			
= <i>Cephalotrichum tonsuratum</i> Meunier, 1910		A	
<i>typicum</i> (Lankester, 1874) Fauré-Fremiet, 1924			
= <i>Torquatella typica</i> Lankester, 1874			
Family Strombidinopsidae			
<i>Strombidinopsis</i>			
<i>acuminatum</i> Fauré-Fremiet, 1924	A		
= <i>Strobilidium acuminatum</i> (Fauré-Fremiet, 1924) Kahl, 1932			
<i>spiniferum</i> (Leegaard, 1915) Lynn et al., 1991			
= <i>Strobilidium spiniferum</i> (Leegaard, 1915) Kahl, 1932			
= <i>Strombidium spiniferum</i> Leegaard, 1915			
<i>minimum</i> (Gruber, 1884) Lynn et al., 1991			
= <i>Arachnidium becheri</i> Buddenbrook, 1920			
= <i>Strobilidium minimum</i> (Gruber, 1884) Kahl, 1932			
= <i>Strombidium minimum</i> Gruber, 1884			
Rhynchodida			
Compiled by Anatol Wiktor Jankowski			
A comprehensive list of European species.			
References			
Banning, P. van 1979 Protistan parasites observed in the European flat oyster (<i>Ostrea edulis</i>) and the cockle (<i>Cerastoderma edule</i>) from some coastal areas of the Netherlands <i>Haliotis</i> , 8: 33-37			
Bower, S.M. & Meyer, G.R. 1993 <i>Stegotricha enterikos</i> gen. nov., sp. nov. (class Phyllopharyngea, order Rhynchodida), a parasitic ciliate in the digestive gland of Pacific oysters (<i>Crassostrea gigas</i>), and its distribution in British Columbia <i>Canad. J. Zool.</i> , 71: 2005-2017			
Burreson, E.M. 1973 Symbiotic ciliates from solitary ascidians in the Pacific Northwest, with a description of <i>Parahypocoma rhamphiskarya</i> n. Sp. <i>Trans. Amer. Micr. Soc.</i> , 92: 517-522			
Butschli, O. 1887-1889 Protozoa. Abtheilung I II. Infusoria und System der Radiolaria. In: Bronn, H.G., hrsg., <i>Klassen und Ordnungen des Thier-Reichs</i> , Bd. I. C.F. Winter Verlag, Leipzig: 1098-2035			
Certes, A. 1882 Sur les parasites intestinaux de l'huître <i>Bull. Soc. Zool. France</i> , 7: 347-353			
Chatton, E. & Lwoff, A. 1921 Sur une famille nouvelle d'acinetiens, les sphenophryides, adaptés aux branchies des mollusques acéphales <i>C. R. Hebdom. Séances Acad. Sci. Paris</i> , 173: 1495-1497			
Chatton, E. & Lwoff, A. 1922 Sur l'évolution des infusoires des lamellibranches. Relation des hypocomides avec les ancistrides. Le genre nouveau <i>Hypocomides</i> n. Gen. <i>C. R. Hebdom. Séances Acad. Aci. Paris</i> , 175: 787-790			
Chatton, E. & Lwoff, A. 1922 Sur l'évolution des infusoires des lamellibranches. Le genre <i>Pelecyophryxa</i> , intermédiaire entre les hypocomides et les sphenophryides. Bourgeonnement et conjugaison <i>C. R. Hebdom. Séances Acad. Sci. Paris</i> , 175: 915-918			
Chatton, E. & Lwoff, A. 1922 Sur l'évolution des infusoires des lamellibranches. Relation des sphenophryides avec les hypocomides <i>C. R. Hebdom. Séances Acad. Sci. Paris</i> , 175: 1444-1447			
Chatton, E. & Lwoff, A. 1924 Sur l'évolution des infusoires des lamellibranches. Morphologie comparée des hypocomides. Les nouveaux genres <i>Hypocomina</i> et <i>Hypocomella</i> <i>C. R. Hebdom. Séances Acad. Sci. Paris</i> , 178: 1928-1931			
Chatton, E. & Lwoff, A. 1926 Diagnoses de ciliés thigmotriches nouveaux <i>Bull. Soc. Zool. France</i> , 51: 345-352			
Chatton, E. & Lwoff, A. 1934 Sur un cilié thigmotriches nouveau: <i>Gargarius gargaricus</i> n. Gen., n. Sp., de <i>Mytilus edulis</i> <i>Bull. Soc. Zool. France</i> , 59: 375-376			
Chatton, E. & Lwoff, A. 1939 Sur le sucoir des infusoires thigmotriches rhynchoïdes (Hypocomidae et Sphenophryidae) et sa genèse <i>C. R. Hebdom. Séances Acad. Sci. Paris</i> , 209: 333-336			
Chatton, E. & Lwoff, A. 1939 Sur la systématique de la tribu des thigmotriches rhynchoïdes. Les deux familles des Hypocomidae Butschli et des Ancistrocomidae n. Fam. Les			

- deux nouveaux genres *Heterocoma* et *Parhypocoma* C. R. *Hebdom. Séances Acad. Sci. Paris*, 209: 429-433
- Chatton, E. & Lwoff, A. 1949 Recherches sur les ciliés thigmotriches. I. *Arch. Zool. Exp. Gen.*, 86: 169-253
- Chatton, E. & Lwoff, A. 1950 Recherches sur les ciliés thigmotriches. II. *Arch. Zool. Exp. Gen.*, 86: 393-485
- Chatton, E., Lwoff, A. & Lwoff, M. 1931 L'infraciliature de l'infusoire thigmotrichie *Sphenophrya dosinia* Ch. Et Lw. Topographie, structure et développement. *C. R. Soc. Biol. Paris*, 107: 532-536
- Chatton, E. & Seguela, J. 1936 Un hypotrope de la branchie de *Ciona intestinalis* L., intermédiaire entre les Euplotidae et les Aspidiscidae: *Euplotaspis cionaecola* n. G., n. Sp. *Bull. Soc. Zool. France*, 61: 332-340
- Cheng, T.C. 1967 Marine molluscs as hosts for symbioses, with a review of known parasites of commercially important species *Adv. Mar. Biol.*, 5: XIII + 424 pp.
- Collin, B. 1907 Note préliminaire sur quelques acinétiens: I. *Ephelota gemmipara* (Hertwig); II. *Hypocoma acinetarum* n. Sp. *Arch. Zool. Exp. Gen.*, ser. 4, Notes et Revue, 7: 93-103
- Collin, B. 1911 Etude monographique sur les acinétiens. I. Recherches expérimentales sur l'étendue des variations et les facteurs térotogènes *Arch. Zool. Exp. Gen.*, 48 (=ser. 5, t. 8): 421-497
- Collin, B. 1912 Etude monographique sur les acinétiens. II. Morphologie, physiologie, systématique *Arch. Zool. Exp. Gen.*, 51: 1-457
- Collin, B. 1914 Notes protistologiques. Sur deux infusoires ciliés parasites d'hétéropodes, et sur un flagelle vivant chez ces mêmes hôtes *Arch. Zool. Exp. Gen.*, 54, Notes et Revue, 85-97
- Comps, M., Razet, D. & Quiros-Ramos, H. 1973 Protiste parasite de l'huître du Pacifique, *Crassostrea gigas* (Thunberg) C. R. *Hebdom. Séances Acad. Sci. Paris*, ser. D, 277: 2797-2799
- Corliss, J.O. 1979 *The ciliated Protozoa. Characterization, classification, and guide to the literature* Pergamon Press. Oxford, New York, Toronto, XVI + 455 pp.
- Delphy, J. 1938 Etudes de morphologie et de physiologie sur la faune d' Arcachon *Bull. Soc. Sci. Arcachon*, 35: 49-75
- Deroux, G. 1974 Les dispositifs adhésifs ciliaires chez les Cyrtophorida et la famille des Hypocomidae *Protistologica*, 10: 379-396
- Deroux, G. 1976 Le plan cortical des Cyrtophorida. Unité d'expression et marges de variabilité. II. Cyrtophorida à thigmotactisme ventral généralisé *Protistologica*, 12: 483-500
- Deroux, G. 1978 *Les Cyrtophorida. Ciliés thigmotactiques de parois immergées* Thèse, Université de Clermont-Ferrand, ser. E, no. 260. Tome I. Texte, p. 1-262. Tome II. Figures, p. 1-111.
- Ergens, R. & Lom, J. 1970 *Puvodci parazitarnich nemoci ryb* Praha, Academia, Naklad. CzSAV: 1-383
- Fenchel, T. 1964 On *Ancistrum caudatum* sp. nov. and *Hypocomides modiolariae* Chatton et Lwoff (Ciliata, Thigmotrichida) from the lamellibranch *Musculus niger* (Gray) *Ophelia*, 1: 113-120
- Fenchel, T. 1965 Ciliates from Scandinavian molluscs *Ophelia*, 2: 71-174
- Fenchel, T. 1965 On the ciliate fauna associated with the marine species of the amphipod genus *Gammarus* J. G. Fabricius *Ophelia*, 2: 281-303
- Gajewskaja, A.W. ed. 1990 *Parasites, commensals and diseases of Black sea mussel* Kiev, Naukova Dumka: 1-132
- Grell, K.G. & Meister, A. 1982 Über die taxonomische Stellung der Hypocomidae (Ciliata) *Zeitschr. für Naturforsch.*, ser. C, 37: 1050-1052
- Grell, K.G. & Meister, A. 1983 Die Ultrastruktur von *Hypocoma acinetarum* Collin *Protistologica*, 19: 51-72
- Gruber, A. 1884 Die Protozoen des Hafens von Genua *Nova Acta Acad. Caesar. Leop. Carol. Natur. Cur.* (=Nova Acta Leopoldina), 46: 1-67 (Pages of Separatum, Halle, 1884)
- Gutierrez, M. 1977 Datos sobre parasitología e histopatología del ostión, *Crassostrea angulata* Lmk. de la costa sudatlántica de España *Invest. Pesquera*, 41: 543-567
- Harant, H. 1931 Les ascidies et leurs parasites. II partie. Les parasites des ascidies *Ann. Inst. Océanogr. Paris*, 8: 347-365
- Herrbach, B. 1971 Sur une affection parasitaire de la glande digestive de l'huître plate, *Ostrea edulis* Linne *Revue des Travaux Inst. Peches Marit.*, 35: 79-87
- Holodkovskaya, E.V. 1989 Fauna of parasites and commensals of the mussel *Mytilus galloprovincialis* Lam. in north-western part of the Black sea. (Systematics, ecology, practical significance) Moscow State Univ., Biol., Dept., Thesis, Abstract: 1-18
- Jankowski, A.W. 1967 Ciliates from the mantle cavity of *Balanus* in Barents sea *Parasitology* (Moscow), 1: 82-93
- Jankowski, A.W. 1975 Conspectus of a new system of subphylum Ciliophora Doflein, 1901 *Scient. Session, Zool. Inst. Acad. Sci. USSR*, Leningrad: 26-27
- Jankowski, A.W. 1976 Revision of a system of cyrtophorines Materials of the II All-Union Conference of Protozoologists. Part I. General Protozoology. Kiev, ed. Naukova Dumka: 167-168
- Jankowski, A.W. 1980 Conspectus of a new system of the phylum Ciliophora *Proc. Zool. Inst. Acad. Sci. USSR*, Leningrad, 94: 102-121
- Jankowski, A.W. 1986 *Ciliophora - commensals and parasites of freshwater molluscs Parasites and diseases of water invertebrates*. Ed, Popova, T.I. & Bauer, O.N. Moscow: 163-164
- Kahl, A. 1934 Ciliata ectocommensalia et parasitica Die *Tierwelt der Nord- und Ostsee*, akad. Verlagsges., Leipzig, Lief. 26-II-C-4: 147-183, 224-225
- Khan, M.A. 1969 Fine structure of *Ancistrocoma pelseneeri* (Chatton et Lwoff), a rhynchodine thigmotrichid ciliate *Acta Protozool.*, 7: 29-47
- Khan, M.A. 1970 Ultrastructural evidence of heavy protein synthesis in *Ancistrocoma pelseneeri* (Thigmotrichida: Ciliata) *Current Science*, 39: 438-439
- Kirby, H. 1941 Relationships between certain Protozoa and other animals Calkins, G.N. & Summers, F.M., eds, *Protozoa in Biological Research*, chapter XIX, Columbia Univ. Press, New York: 890-1008
- Kofoid, C.A. & Bush, M. 1936 The life cycle of *Parachaenia myae* gen. nov., sp. nov., a ciliate parasite in *Mya arenaria* Linn. from San Francisco Bay, California *Bull. Mus. Royal Hist. Nat. de Belgique*, 12: 1-15
- Kozloff, E.N. 1946 Studies on ciliates of the family *Ancistrocomidae* Chatton and Lwoff (order Holotrichia, suborder Thigmotrichia). I. *Hypocomina tegularum* sp. nov. and *Crebricoma* gen. nov. *Biol. Bull. Woods Hole*, 90: 1-7
- Kozloff, E.N. 1946 Studies... II. *Hypocomides mytili* Chatton and Lwoff, *Hypocomides botulae* sp. nov., *Hypocomides parva* sp. nov., *Hypocomides kelliæ* sp. nov., and *Insignicoma venusta* gen. nov., sp. nov. *Biol. Bull. Woods Hole*, 90: 200-212
- Kozloff, E.N. 1946 Studies... III. *Ancistrocoma pelseneeri* Chatton and Lwoff, *Ancistrocoma dissimilis* sp. nov., and *Hypocomagalma pholadidis* sp. nov. *Biol. Bull. Woods Hole*, 91: 189-199
- Kozloff, E.N. 1961 A new genus and two new species of ancistrocomid ciliates (Holotrichia: Thigmotricha) from sabellid polychaete and from a chiton *J. Protozool.*, 8: 60-63
- Kozloff, E.N. 1965 *Colligocineta furax* gen. nov., sp. nov., an ancistrocomid ciliate (Holotrichia: Thigmotricha) from the sabellid polychaete *Laonome kroyeri* Malmgren *J. Protozool.*, 12: 333-334
- Kozloff, E.N. 1976 New ancistrocomid ciliates from polychaete annelids *Trans. Amer. Micr. Soc.*, 95: 622-627
- Lauckner, G. 1983 Diseases of Mollusca: *Bivalvia* Kinne, O., ed., *Diseases of marine animals*. Vol. 2: *Introduction, Bivalvia to Scaphopoda*. Biol. Anstalt Helgoland, Hamburg: 477-961

- Lichtenstein, J.L. 1921 *Hypocoma patellarum* n. sp., acinétiens parasite de *Patella coerulea* C. R. *Hebdom. Séances Soc. Biol. Paris*, 85: 796-799
- Lom, J. & Kozloff, E.N. 1968 Observations on the ultrastructure of the suctorial tube of ancistrocomid ciliates *Folia Parasitol.*, Praha, 15: 291-308
- Lom, J. & Kozloff, E.N. 1969 Ultrastructure of the cortical regions of ancistrocomid ciliates *Protistologica*, 5: 173-192
- Mjassnikowa, M. 1930 Über einen neuen Vertreter der Familie Sphaenophryidae aus *Mya truncata* L. *Arch. Protistenk.*, 72: 377-389
- Plate, L.H. 1888 Studien über Protozoen. Die Gattung *Acinetoides* n. g., eine Intermediar-Form zwischen den Ciliaten Infusorien und den Acineten. *Zool. Jahrb., Abth. Anat. Syst. Ontog. der Thiere*, 3: 155-200
- Puytorac, P. de 1969 Remarques à propos de l'ultrastructure du sucoïd d'*Ancistrocoma myae* (K. et B.), cilié Rhynchodea C. R. *Hebdom. Séances Acad. Sci. Paris*, ser. D, 268: 820-822
- Puytorac, P. de 1994 Sous-classe des Rhynchodia Chatton et Lwoff, 1939 Grassé, P.-p., ed., *Traité de Zoologie. Anatomie, systématique, biologie*. - Puytorac, P. de, ed. Infusoires ciliés. T. II, fasc. 2. Masson, Paris, Milan, Barcelone: 475-492
- Raabe, Z. 1934 Über einige an den Kiemen von *Mytilus edulis* L. und *Macoma balthica* (L.) parasitierende Ciliaten-Arten *Annales Mus. Zool. Polon.*, 10: 289-303
- Raabe, Z. 1935 *Rhynchophrya cristallina* g. n., sp. n., nouvelle forme d'infusoire de la famille des Sphaenophryidae Chatten et Lwoff *Bull. Inst. Océanogr. Monaco*, 676: I-6
- Raabe, Z. 1938 Weitere Untersuchungen an parasitischen Ciliaten aus dem Polnischer Teil der Ostsee. II. Ciliata Thigmotricha aus den Familien: Hypocomidae Butschli und Sphaenophryidae Ch. et Lw. *Annales Mus. Zool. Polon.*, 13: 41-75
- Raabe, Z. 1949 Remarks on protozoan parasitocenose of some representatives of genus *Mytilus* *Annales Univ. Mariae Curie-Sklodowska, Lublin*, sect. C, Biol. Sci., 4: 1-16
- Raabe, Z. 1950 Recherches sur les ciliés thigmotriches (Thigmotricha Ch. Lw.). V. Ciliés thigmotriches du lac Balaton (Hongrie). *Annales Univ. Mariae Curie-Sklodowska, Lublin*, sect. C, Biol. Sci., 5: 197-215
- Raabe, Z. 1949 Recherches sur les ciliés thigmotriches (Thigmotricha Ch. Lw.). III. Développement non-parallèle de deux espèces du genre *Sphenophrya* Ch. Lw. *Annales Univ. Mariae Curie-Sklodowska, Lublin*, sect. C, Biol. Sci., 4: 119-135
- Raabe, Z. 1956 Badania nad parazytofauna miedzakow slodkowodnych w wodach slonawych. (In Polish) (Investigations on the parasitofauna of freshwater molluscs in the brackish waters) *Acta Parasitol. polon.*, 4: 375-406
- Raabe, Z. 1967 Ordo Thigmotricha (Ciliata Holotricha). I. *Acta Protozool.*, 5: 1-36
- Raabe, Z. 1970 Ordo Thigmotricha (Ciliata Holotricha). II. Familiae Ancistrocomidae et Sphenophryidae *Acta Protozool.*, 7: 385-463
- Sand, R. 1901 Etude monographique sur le groupe des infusoires tentaculifères *Annales Soc. Belge de Microscopie*, t. 24, 25, 26. 1899-1900. Separate edition as book: Bruxelles, A. Castaigne ed., 1901: 1-445
- Tunberg, B. 1984 Population ecology of *Dosinia exoleta* (L.) (Bivalvia) in Raunefjorden, western Norway *Sarsia*, 69: 159-168
- Rhynchodida**
- Family Hypocomidae**
- Crateristoma kindi* Jankowski, 1967
- Enigmocoma acinetarum* (Collin, 1907) Jankowski, 1976
= *Hypocoma acinetarum* Collin, 1907
- Harmocoma rostellum* (Deroux, 1976) Jankowski, 1980
= *Hypocoma rostellum* Deroux, 1976
- Hypocoma ascidiarum* Collin, 1912
- parasitica* Gruber, 1884
- = *Acinetoides greeffii* Plate, 1888
- = *Acinetoides zoothamni* Plate, 1888
- Parahypocoma collini* Chatton & Lwoff, 1939
- Rhynchocoma hyperparasitica* (Chatton & Lwoff, 1939) Jankowski, comb. n.
= *Heterocoma hyperparasitica* Chatton & Lwoff, 1939
- Family Ancistrocomidae**
- Ancistrocoma myae* (Kofoid & Bush, 1936) Chatton & Lwoff, 1950
= *Parachaenia myae* Kofoid & Bush, 1936
- pelseneeri* Chatton & Lwoff, 1926
- pholadis* Chatton & Lwoff, 1926
- thorsoni* Fenchel, 1965
- Anisocomides zyprhaeae* (Chatton & Lwoff, 1926) Chatton & Lwoff, 1950
= *Hypocomides zyprhaeae* Chatton & Lwoff, 1926
- Colligocineta affinis* Kozloff, 1976
- finleyi* Kozloff, 1976
- scolepidis* Kozloff, 1976
- Crebricoma carinata* (Raabe, 1934) Kozloff, 1946
= *Hypocomina carinata* Raabe, 1934
- Goniocoma macomae* (Chatton & Lwoff, 1926) Chatton & Lwoff, 1950
= *Hypocomella macomae* Chatton & Lwoff, 1924
- Holocoma primigenius* Chatton & Lwoff, 1950
- Hypocomagalma dreissenae* Jarocki & Raabe, 1932
- Hypocomatidium jarockii* Chatton & Lwoff, 1950
- Hypocomella cardii* (Chatton & Lwoff, 1922) Chatton & Lwoff, 1924
= *Hypocoma cardii* Chatton & Lwoff, 1922
- krzysiki* (Jarocki, 1934) Raabe, 1970
= *Encelys pisii* Delphy, 1938
= *Heterocineta krzysiki* Jarocki & Raabe, 1934
- raabei* Chatton & Lwoff, 1950
- roscoffensis* Chatton & Lwoff, 1950
- Hypocomides astartae* Fenchel, 1965
- elsinora* Fenchel, 1965
- hiatellae* Fenchel, 1965
- modiolariae* Chatton & Lwoff, 1922
- musculus* Fenchel, 1965
- Hypocomidium fabius* Raabe, 1938
- gramum* Raabe, 1938
- Hypocomina patellarum* (Lichtenstein, 1921) Chatton & Lwoff, 1924
= *Hypocoma patellarum* Lichtenstein, 1921
- Isocomidae**
- myili* (Chatton & Lwoff, 1922) Chatton & Lwoff, 1950
= *Hypocomides myili* Chatton & Lwoff, 1922
- Raabella helensis* Chatton & Lwoff 1950, 1950
- Stegotricha enterikos* Bower & Meyer, 1993
- Syringopharynx pterotrachaeae* Collin, 1914
- Family Sphenophryidae**
- Gargarius gargarius* Chatton & Lwoff, 1934
= *Rhynchophrya cristallina* Raabe, 1935
- Parasphenophrypa sphaerii* (Mjassnikowa, 1930) Jankowski, comb. n.
= *Sphenophrya sphaerii* Mjassnikowa, 1930

Pelecyophrya
tapetis Chatton & Lwoff, 1922
Sphenophrya
cardii Chatton & Lwoff, 1950
dosiniae Chatton & Lwoff 1950, 1921
myae Mjassnikowa, 1930

Dinoflagellata

Compiled by Sue Brandt

Preliminary list based on regional lists compiled by Dr. M. Elbrächter and a few other surveys. Dr Elbrächter is thanked for allowing the editor access to his lists. The list below is still incomplete in terms of geographical coverage, and, particularly, in terms of synonymy.

References

- Dodge, J.D. 1982 *Marine dinoflagellates of the British Isles*. HMSO, London
 Elbrächter, M.L. 2000 Personal communication; species from the Mediterranean, German Bight, and coastal Finland
 Griessmann, K. 1913 Über marine Flagellaten. *Archiv. Protistenk.* 32: 1-78
 Larsen, J. 1985 Algal studies of the Danish Wadden Sea II. A taxonomic study of psammobious dinoflagellates *Opera Botanica*, 79: 14-37
 Thomsen, H.A. (ed.) 1992 *Plankton i de indre danske farvande*. Miljøministeriet, Miljøstyrelsen, Copenhagen.

Dinoflagellata

Achradina
pulchra Lohmann
Actiniscus
pentasterias (Ehrenberg) Ehrenberg, 1854
Adenoides
eludens (Herdman) Balech
Adinomonas
oviforme Schiller
Alexandrium
affine (Inoue & Fukuyo) Balech, 1985
balechii (Steidinger) Balech
foedum Balech, 1990
leei Balech, 1985
minutum Halim, 1960
ostenfeldii (Paulsen) Balech & Tangen
 = *Gessnerium ostenfeldii* (Paulsen) Loeblich & Loeblich
 = *Goniiodoma ostenfeldii* Paulsen
 = *Gonyaulax globosa* (Braarud) Balech
 = *Gonyaulax tamarensis* Lebour var. *globosa* Braarud
 = *Heteraulacus ostenfeldii* (Paulsen) Loeblich III
 = *Triadinium ostenfeldii* (Paulsen) Dodge
pseudogonyaulax (Biecheler) Yuki & Fukuyo
tamarensis (Lebour) Balech
 = *Gonyaulax excavata* (Braarud) Balech
 = *Gonyaulax tamarensis* Lebour
 = *Gonyaulax tamarensis* Lebour var. *excavata* Braarud
taylori Balech, 1994
Amoeobaphya
ceratii (Koeppen) Cachon
Amphidiniopsis
hirsum (Balech, 1956) Dodge, 1982
kofoidii Woloszynska, 1928
swedmarkii (Balech, 1956) Dodge, 1982
Amphidinium
acutissimum Schiller, 1933
acutum Lohmann
bipes Herdman, 1924
britannicum (Lebour) Herdman, 1924
 = *Amphidinium asymmetricum* var. *britannicum* Herdman 1922
 = *Amphidinium asymmetricum* var. *compactum* Herdman 1922
carterae Hulbert, 1967
conus Schiller
corpulentum Kofoid & Swezy, 1921

crassum Lohmann, 1908
curvatum Schiller
flagellans Schiller
flexum Herdman, 1923
globosum Schröder, 1911
herdmani Kofoid & Swezy, 1921
 = *Amphidinium operculatum* auct.
hoeferi Schiller & Diskus
incoloratum Campbell, 1973
kesselitzii Schiller
lacustre Stein, 1883
lacustriforme Schiller
lanceolatum Schröder
latum Lebour, 1925
longum Lohmann, 1908
manannini Herdman, 1924
massartii Biecheler, 1952
operculatum Claparède & Lachmann, 1859
 = *Amphidinium klebsii* Kofoid & Swezy, 1921
ovoideum (Lemmermann) Lemmermann, 1900
ovum Herdman, 1924
pellucidum Herdman, 1922
poecilochroum Larsen, 1985
psittacus Larsen, 1985
schroederii Schiller
scissum Kofoid & Swezy, 1921
semilunatum Herdman, 1924
 = *Thecadinium semilunatum* (Herdman) Dodge, 1982
sphenoides Wulff, 1916
 = *Gymnodinium filum* Lebour, 1917
steinii (Lemmermann) Kofoid & Swezy, 1921
stigmatum Schiller
subsalum Biecheler, 1952
testudo Herdman, 1924
Amphidoma
caudata Halldal, 1953
Amphisolenia
bidentata Schröder
brevicaudata Kofoid
extensa Kofoid
globifera Stein, 1883
inflata Murray & Whitting
palmata Stein, 1883
rectangulata Kofoid
schroeteri Kofoid
spinulosa Kofoid
truncata Kofoid & Michener
Amylax
triacantha (Jørgensen) Sournia
Archaeosphaerodiniopsis
verrucosa Rampi, 1943
Blepharocysta
paulsenii Schiller, 1937
splendor-maris Ehrenberg, 1873
Boreadinium
pisiforme Dodge
Brachydinium
capitatum Taylor, 1963
Cachonina
niei Loeblich
Cachonodinium
caudatum (J. & M. Cachon) Loeblich
Calcidinellum
operosum De Flandre
Centrodinium
complanatum (Cleve) Kofoid
intermedium Pavillard
maximum Pavillard
Ceratium
arietinum Cleve, 1900
azoricum Cleve, 1900
belone Cleve
breve (Ostenfeld & Schmidt) Schröder
brunellii Rampi
buceros (Zacharias) Schiller

- candelabrum* (Ehrenberg) Stein, 1883
carriense Gourret
cephalotum (Lemmermann) Jörgensen
coarctatum Pavillard
compressum Gran, 1902
concilians Jörgensen
contortum (Gourret) Cleve
contrarium (Gourret) Pavillard
declinatum (Karsten) Jörgensen
deflexum (Kofoid) Jörgensen
digitatum Schütt
euarcatum Jörgensen
extensum (Gourret) Cleve
falciforme Jörgensen
falcatum (Kofoid) Jörgensen
furca (Ehrenberg) Claparède & Lachmann
fusus (Ehrenberg) Dujardin
geniculatum (Lemmermann) Cleve
gibberum Gourret
gravidum Gourret, 1883
hexacanthum Gourret, 1883
horridum (Cleve) Gran
= *Ceratium intermediate* (Cleve) Jörgensen, 1911
incisum (Karsten) Jörgensen
inflatum (Kofoid) Jörgensen
karstenii Pavillard
kofoidii Jörgensen
limulus (Gourret) Gourret
lineatum (Ehrenberg) Cleve
longipes (Bailey) Gran
longirostrum Gourret
longissimum (Schröder) Kofoid
macroceros (Ehrenberg) Vanhöffen
massiliense (Gourret) Jörgensen
minutum Jörgensen
paradoxides Cleve
pavillardii Jörgensen
pentagonum Gourret
praelongum (Lemmermann) Kofoid
pulchellum Schröder
ranipes Cleve
schoeteri Schröder
setaceum Jörgensen
strictum (Kofoid) Jörgensen
symmetricum Pavillard
teres Kofoid
trichoceros (Ehrenberg) Kofoid
tripos (O.F. Müller) Nitzsch, 1817
vultur Cleve
Ceratocorys
armata (Schütt) Kofoid
gourrei Paulsen
horrida Stein
magna Kofoid
Ceratoperidinium
yeye Margalef, 1969
Citharistes
regius Stein, 1883
Cladopyxis
brachiolata Stein
caryophyllum (Kofoid) Pavillard
spinosa (Kofoid) Schiller
Clipedinium
pileus Pascher
Cochlodinium
achromaticum Lebour, 1925
adriaticum Schiller
archimedes (Pouchet) Lemmermann
brandtii Wulff, 1916
cnidophorum Biecheler, 1939
constrictum (Schütt) Lemmermann
geminatum (Schütt) Schütt
helicoideas Lebour, 1925
= *Cochlodinium helix* Kofoid & Swezy
helix (Pouchet) Lemmermann, 1899
= *Gymnodinium helix* Pouchet, 1887
heterolobatum Silva
pellucidum Lohmann, 1908
platycorne Daday, 1888
pulchellum Lebour, 1917
pupa Lebour, 1925
schuetii Kofoid & Swezy, 1921
strangulatum Schütt
vincutum Kofoid & Swezy, 1921
Coolia
monotis Meunier
Corythodinium
diploconus (Stein, 1883) Taylor, 1976
frenguellii (Rampi) Balech
michaelsarsii (Gaarder, 1954) Taylor, 1976
reticulatum (Stein, 1883) Taylor, 1976
tesselatum (Stein) Loeblich & Loeblich
Craspedotella
pileolus Kofoid
Cryptocodinium
cohnii (Seligo) Chatton
setense Biecheler
Cymbodinium
elegans Cachon & Cachon
Desmocapsa
gelatinosa Pascher
Dinophysis
acuminata Claparède & Lachmann
acuta Ehrenberg
acutoides Balech
alata Jörgensen
amandula Sournia
apicata (Kofoid & Skogsberg) Abé
argus (Stein) Abé
baltica (Paulsen) Kofoid & Skogsberg
biceps Schiller
caudata Saville-Kent, 1881
circumsuta (Kofoid) Balech
contracta (Kofoid & Skogsberg) Balech, 1967
cuneus (Schütt) Abé
dens Pavillard, 1915
dentata Schiller
diegensis Kofoid
dolichopterygium (Murray & Whitting) Balech
doryphora (Stein) Abé
elongata (Jörgensen) Abé
expulsa Kofoid & Michener
fava (Kofoid & Michener) Abé
fortii Pavillard
gigantea (Kofoid & Michener) Balech
greenlandica (Schiller) Balech
hastata Stein, 1883
hindemarchi (Schütt) Abé
infundibulus Schiller
intermedia Pavillard
irregularis (Lebour) Balech
lachmanni Paulsen
lenticula Pavillard
limbata (Kofoid & Michener) Balech
mitra (Schütt) Abé
muconata (Kofoid & Skogsberg, 1928) Sournia, 1973
norvegica Claparède & Lachmann
odiosa (Pavillard) Tai & Skogsberg
operculata (Stein) Balech
operculoides (Schütt) Balech
ovata Claparède & Lachmann
ovum Schütt, 1895
parva Schiller
parvula (Schütt) Balech
paulseni (Schiller) Balech
planiceps (Schiller) Balech
porodictylum (Stein) Abé
pugiunculus (Jörgensen) Balech
pulchella (Lebour) Balech
pulchra (Kofoid & Michener) Balech

<i>punctata</i> Jørgensen, 1923	
<i>pusilla</i> Jørgensen	
<i>rapa</i> (Stein) Abé	
<i>recurva</i> Kofoid & Skogsberg, 1928	
<i>rotundata</i> Claparède & Lachmann	
<i>schroederii</i> Pavillard	
<i>schuettii</i> Murray & Whitting	
<i>similis</i> Kofoid & Skogsberg	
<i>skagii</i> Paulsen	
<i>sphaerica</i> Stein	
<i>striata</i> (Schütt) Abé	
<i>tripos</i> Gourret	
<i>umbosa</i> Schiller	
<i>uracantha</i> Stein	
<i>Diplopelta</i>	
bomba Stein ex. Jørgensen	
= <i>Diplopelta asymmetrica</i> Mangin	
= <i>Diplopelta asymmetricum</i> Lebour	
= <i>Diplopsalopsis asymmetricum</i> Abé, 1941	
= <i>Diplosalis lenticulata</i> Stein, 1883	
= <i>Dissodium asymmetricum</i> Loeblich, 1970	
= <i>Peridiniopsis asymmetrica</i> Lebour, 1922	
= <i>Peridiniopsis asymmetrica</i> Mangin, 1911	
<i>pusilla</i> Balech & Akselman	
<i>symmetrica</i> Pavillard, 1913	
<i>Diplopsalis</i>	
<i>excentrica</i> Nie, 1943	
<i>lenticula</i> Bergh	
<i>minima</i> Mangin	
<i>pilula</i> Ostenfeld	
<i>Diplopsalopsis</i>	
<i>latipeltata</i> Balech & Borgese, 1990	
<i>orbicularis</i> (Paulsen) Meunier	
<i>Dissodinium</i>	
<i>pseudocalani</i> (Gönnert) Drebess	
<i>pseudolunula</i> Swift	
<i>Endodinium</i>	
<i>chattonii</i> Hovasse, 1922	
<i>Erythropsodinium</i>	
<i>agile</i> (Hertwig) Silva	
<i>Filodinium</i>	
<i>hovassei</i> Cachon & Cachon	
<i>Fragilidium</i>	
<i>fissile</i> Balech, 1990	
<i>subglobosum</i> (von Stosch) Loeblich, 1980	
= <i>Helgolandium subglobosum</i> von Stosch, 1969	
<i>Glenodinium</i>	
<i>armatum</i> Levander	
<i>bacilliferum</i> Biecheler, 1952	
<i>berghii</i> Lemmermann	
<i>cinctum</i> Ehrenberg	
<i>cohnii</i> Seligo, 1885	
<i>danicum</i> Paulsen	
<i>gymnodinium</i> Pernard	
<i>monensis</i> Herdman, 1923	
<i>paululum</i> Lindemann	
<i>pulvisculus</i> Stein	
<i>Gleodinium</i>	
<i>marinum</i> Bouquaqueux	
<i>Goniadoma</i>	
<i>acuminata</i> (Ehrenberg, 1834) Stein, 1883	
<i>sphaericum</i> Murray & Whitting, 1899	
<i>Gonyaulax</i>	
<i>africana</i> Schiller	
<i>alaskensis</i> Kofoid	
<i>apiculata</i> (Pernard) Entz	
<i>birostis</i> Stein	
<i>buxus</i> Balech	
<i>diegensis</i> Kofoid	
<i>digitale</i> (Pouchet) Kofoid	
<i>dimorpha</i> Biecheler, 1952	
<i>elegans</i> Rambi	
<i>fragilis</i> (Schütt) Kofoid	
<i>fusiformis</i> Graham	
<i>gracilis</i> Schiller	
<i>kofoidii</i> Pavillard	
<i>ligustica</i> Rambi	
<i>milneri</i> (Murray & Whitting) Kofoid	
<i>minima</i> Matzenauer	
<i>mitra</i> (Schütt) Kofoid	
<i>monacantha</i> Pavillard	
<i>monospina</i> Rambi	
<i>ovalis</i> Schiller	
<i>pacifica</i> Kofoid	
<i>polyedra</i> Stein	
<i>polygramma</i> Stein	
<i>rostrata</i> Dangeard	
<i>rotundata</i> Rambi	
<i>rugosum</i> Wailes	
<i>scrippsae</i> Kofoid	
<i>sphaeroidea</i> Kofoid	
<i>spinifera</i> (Clap. & Lach.) Diesing	
<i>turbynei</i> Murray & Whitting	
<i>unicornis</i> Lebour	
<i>verior</i> Sournia, 1973	
= <i>Amylax diacantha</i> Meunier, 1919	
= <i>Gonyaulax diacantha</i> Schiller, 1937	
= <i>Gonyaulax longispina</i> Lebour, 1925	
<i>Greuetodinium</i>	
<i>cylindricum</i> Loeblich	
<i>Gymnodinium</i>	
<i>abbreviatum</i> Kofoid & Swezy, 1921	
<i>achromaticum</i> Lebour, 1917	
<i>adriaticum</i> (Schmarda) Kofoid & Swezy	
<i>agiliformis</i> Schiller	
<i>arenicola</i> Dragesco	
<i>areuginosum</i> Stein	
<i>caput</i> Schiller	
<i>catenatum</i> Davies	
<i>chlorophorum</i> Elbrächter & Schnepf	
<i>coeruleum</i> Dogiel	
<i>conicum</i> Kofoid & Swezy, 1921	
<i>corii</i> Schiller	
<i>cucumis</i> Schütt	
<i>diploconus</i> Schütt	
<i>elongatum</i> Hope, 1954	
<i>endofasciculum</i> Campbell, 1973	
<i>fuscum</i> (Ehrenberg) Stein	
<i>fusus</i> Schütt	
<i>galatheanum</i> Braarud, 1957	
<i>gibbera</i> Schiller	
<i>gleba</i> Schütt	
<i>gracile</i> Bergh, 1881	
<i>grammaticum</i> (Pouchet) Kofoid & Swezy	
<i>halophilum</i> Biecheler, 1952	
<i>helveticae</i> Pernard	
<i>herbaceum</i> Kofoid	
<i>incertum</i> Herdman, 1924	
<i>lantzschii</i> Utermöhl	
<i>lebourae</i> Pavillard, 1921	
<i>lohmannii</i> Paulsen, 1908	
<i>maguelonnense</i> Biecheler, 1952	
<i>marinum</i> Saville-Kent, 1881	
<i>micrum</i> (Leadbeater & Dodge) Loeblich, 1970	
<i>mikimotoi</i> Miyake & Kominami	
<i>minor</i> Lebour	
<i>multistriatum</i> Kofoid & Swezy	
<i>najadeum</i> Schiller	
<i>neapolitanum</i> Schiller	
<i>nollerii</i> Ellegaard & Moestrup, 1999	
<i>ostenfeldii</i> Schiller, 1928	
<i>paulseini</i> Schiller	
<i>pigmentosum</i> (Dodge, 1967) Loeblich, 1970	
<i>placidum</i> Herdman, 1922	
<i>pseudonociluca</i> Pouchet, 1885	
<i>pulchrum</i> Schiller	
<i>pygmaeum</i> Lebour, 1925	
<i>rhomboides</i> Schütt, 1895	
<i>rubrocinctum</i> Lebour, 1925	
<i>rubrum</i> Kofoid & Swezy, 1921	

- sangineum* Hirasaka, 1924
 = *Gymnodinium splendens* Lebour, 1925
semidivisum Schiller
simplex (Lohmann) Kofoid & Swezy, 1921
striassimum Hulbert
ubberimum Alm. In Kofoid & Swezy
variabile Herdman, 1924
venescum Ballantine, 1956
vestigii Schütt, 1895
vitiligo Ballantine, 1956
vorticella Stein
voukii Schiller
Gyrodinium
acutum (Schütt) Kofoid & Swezy
aureolum Hulbert, 1957
britannicum Kofoid & Swezy, 1921
calyptoglyphe Lebour, 1925
cochlea Lebour, 1925
conicum Schiller
contortum (Schütt) Kofoid & Swezy
corsicum Paulmier, Berland, Billard & Nezan, 1995
crassum (Pouchet, 1885) Kofoid & Swezy, 1921
cuneatum Kofoid & Swezy, 1921
dominans Hulbert, 1957
estuariale Hulbert, 1957
fissum (Levander) Kofoid & Swezy
flagellare Schiller
foliaceum Kofoid & Swezy
fusiforme Kofoid & Swezy, 1921
impedans Larsen, 1996
impudicum Fraga & Bravo, 1995
lacryma (Meunier) Kofoid & Swezy
linguliferum Lebour, 1925
longum (Lohmann) Kofoid & Swezy, 1921
mitra Kofoid & Swezy
obtusum (Schütt) Kofoid & Swezy, 1925
ochraceum Kofoid & Swezy
opimum (Schütt, 1895) Lebour, 1925
ovatum (Gourret) Kofoid & Swezy
ovum (Schütt) Kofoid & Swezy
parvulum (Schütt) Kofoid & Swezy, 1921
pavillardii Biecheler, 1934
pellucidum (Wulff) Schiller, 1933
pepo (Schütt) Kofoid & Swezy, 1921
pingue (Schütt) Kofoid & Swezy
prunus (Wulff) Lebour, 1925
resplendens Hulbert, 1957
schuetii (Lemmermann) Kofoid & Swezy
spirale (Bergh) Kofoid & Swezy, 1921
uncatenatum Hulbert, 1957
undulans Hulbert, 1957
varians (Wulff) Schiller
vorax Biecheler, 1952
wulffii Schiller
Hemidinium
nasutum Stein, 1883
ochraceum Levander
Herdmania
litoralis Dodge, 1981
Heterocapsa
rotundata (Lohmann) Hansen, 1995
triquetra (Ehrenberg) Stein
Heterodinium
agassizi Kofoid
balechii Rambi
detonii Rambi
dispar Kofoid & Adamson
doma (Murray & Whitting) Kofoid
dubium Rambi
globosum Kofoid
grahamii Rambi
inaequale (Kofoid) Schiller
laticeps Léger
leiorhynchum (Murray & Whitting) Kofoid
mediterraneum Pavillard
milneri (Murray & Whitting, 1899) Kofoid, 1906
minutum Kofoid & Michener
murrayi Kofoid, 1906
richardii Pavillard
scrippsi Kofoid
whittingae Kofoid
Histioneis
alata Rambi
bernardii Rambi
depressa Schiller
detonii Rambi
elegans Halim
expansa Rambi
faouzii Halim
guberans Schütt
imbricata Halim
inclinata Kofoid & Michener
joergensenii Schiller
ligustica Rambi
longicollis Kofoid
marchesonii Rambi
oxyptera Schiller
pavillardii Rambi
rampii Halim
robusta Rambi
speciosa Rambi
subcarinata Rambi
sub-longicollis Halim
variabilis Schiller
villafranca Halim
vouckii Schiller
Katodinium
asymmetricum (Massart) Loeblich, 1965
fungiforme (Annisimova) Loeblich III, 1965
glandulum (Herdman) Loeblich, 1970
= *Gymnodinium glandula* Herdman, 1924
= *Massartia glandula* (Herdman) Schiller, 1933
glaucum (Lebour) Loeblich, 1965
nieuportense (Conrad) Loeblich & Loeblich, 1966
rotundatum (Lohmann) Loeblich
Kofoidinium
pavillardii Cachon & Cachon
spendens Cachon & Cachon
velleloides Pavillard, 1928
Kolkwitziella
acuta (Apstein) Elbrächter
Kryptoperidinium
foliaceum (Stein) Lindemann
Lebouraia
minuta Abé
Leptodiscus
medusoides Hertwig
Leptophyllus
dasypus Cachon & Cachon
Mesoporus
perforatus (Gran) Lillick
= *Mesoporus adriaticus* (Schiller) Lillick
= *Mesoporus asymmetricum* (Schiller) Lillick
= *Mesoporus bisimpressus* (Schiller) Lillick
= *Mesoporus globulus* (Schiller) Lillick
Micracanthodinium
claytonii (Holmes, 1956) Dodge, 1982
setiferum (Lohmann, 1902) Deflandre, 1937
Micranthodinium
bacilliferum (Schiller) Deflandre
quadrispinum (Pavillard) Margalef
Myxodinium
pipiens Cachon, Cachon & Bouquaheux, 1970
Nematodinium
armatum (Dogiel) Kofoid & Swezy, 1921
lebourae (Kofoid & Swezy) Schiller
Noctiluca
scintillans (Macartney) Kofoid
Oblea
rotunda (Lebour) Balech ex. Sournia

<i>Ornithocercus</i>	<i>kornmannii</i> Drebes & Schnepf
<i>bilobatus</i> Rampi	<i>vonstoschii</i> Drebes & Schnepf
<i>carolinae</i> Kofoid	<i>Pavillardinium</i>
<i>geniculatus</i> Dangeard	<i>intermedium</i> (Pavillard) De Toni
<i>heteroporus</i> Kofoid	<i>punctatum</i> (Cleve) De Toni
<i>magnificus</i> Stein	<i>splendidum</i> Rampi
<i>quadratus</i> Schütt	<i>Pentapharsodinium</i>
<i>splendidus</i> Schütt	<i>dalei</i> Indelicato & Loblich
<i>steinii</i> Schütt	<i>tyrrhenicum</i> (Balech) Montressor, Zingone &
<i>thumi</i> (Schmidt) Kofoid & Skogsberg	Marino, 1993
<i>Ostreopsis</i>	<i>Peridiniella</i>
<i>ovata</i> Fukuyo, 1981	<i>catenata</i> (Levander) Balech
<i>Oxyphysis</i>	<i>Peridiniopsis</i>
<i>oxytoxoides</i> Kofoid	<i>borgei</i> Lemmermann
<i>Oxyrrhis</i>	<i>dinobryonis</i> (Woloszynska) Bourrelly
<i>marina</i> Dujardin, 1841	<i>elpatiewskyi</i> (Ostenfeld) Bourrelly
<i>Oxytoxum</i>	<i>pernardiiforme</i> (Lindemann) Bourrelly
<i>aceratum</i> Rampi	<i>pernardii</i> (Lemmermann) Lebour
<i>adriaticum</i> Schiller	<i>polonicum</i> (Woloszynska) Bourrelly
<i>areolatum</i> Rampi	<i>pygmeus</i> (Lindemann) Bourrelly
<i>brunellii</i> Rampi	<i>quadridens</i> (Stein) Bourrelly
<i>caudatum</i> Schiller	<i>Peridinium</i>
<i>challengeroides</i> Kofoid	<i>balticum</i> (Levander) Lemmermann
<i>compressum</i> Kofoid	<i>bipes</i> Stein
<i>constrictum</i> (Stein) Blütschli	<i>chattonii</i> Biecheler, 1952
<i>coronatum</i> Schiller	<i>cinctum</i> (O.F. Müller) Ehrenberg
<i>crassum</i> Schiller	<i>dalei</i> (Indel. & Loeb.) Balech
<i>cribosum</i> Stein	<i>gargantua</i> Biecheler, 1952
<i>cristatum</i> Kofoid	<i>goslavicense</i> Woloszynska
<i>curvatum</i> Kofoid	<i>hangoei</i> Schiller
<i>curvicaudatum</i> Kofoid	<i>inconspicuum</i> Lemmermann
<i>depressum</i> Schiller	<i>michaelis</i> Ehrenberg
<i>diploconus</i> Stein	<i>palatinum</i> Lauterborn
<i>elegans</i> Pavillard	<i>palustre</i> Lefèvre
<i>gigas</i> Kofoid	<i>pusillum</i> (Pernard) Lemmermann
<i>globosum</i> Schiller	<i>tabulatum</i> Ehrenberg
<i>laticeps</i> Schiller, 1937	<i>umbonatum</i> Stein
<i>ligisticum</i> Rampi	<i>volzii</i> Lemmermann
<i>longiceps</i> Schiller	<i>willei</i> Huitfeld-Kaas
<i>longum</i> Schiller	<i>Petalodinium</i>
<i>margalefii</i> Rampi	<i>porcelio</i> Cachon & Cachon
<i>mediterraneum</i> Schiller	<i>Pheopolykrikos</i>
<i>milneri</i> Murray & Whitting, 1899	<i>beauchampii</i> Chatton
<i>minutum</i> Rampi	<i>Planodinium</i>
<i>mitra</i> Stein	<i>striatum</i> Saunders & Dodge
<i>obesum</i> Rampi	<i>Podolampas</i>
<i>obliquum</i> Schiller	<i>bipes</i> Stein
<i>ovale</i> Schiller	<i>curvatus</i> Schiller
<i>parvum</i> Schiller	<i>elegans</i> Schütt
<i>punctulatum</i> Rampi	<i>spinifera</i> Okamura
<i>radiosum</i> Rampi	<i>Podolampus</i>
<i>rampii</i> Sournia	<i>palmipes</i> Stein, 1883
<i>reticulatum</i> (Stein) Schütt	<i>Polykrikos</i>
<i>sceptrum</i> (Stein) Schröder	<i>beauchampii</i> (Chatton) Dodge, 1982
<i>scolopax</i> Stein, 1883	<i>hartmannii</i> Zimmermann
<i>sphaeroideum</i> Stein, 1883	<i>kofoidii</i> Chatton
<i>spinosum</i> Rampi	<i>lebourae</i> Herdman
<i>subulatum</i> Kofoid	<i>schwarzii</i> Blütschli
<i>tenuistriatum</i> Rampi	<i>Pomatodinium</i>
<i>tonollii</i> Rampi	<i>impatiens</i> Cachon & Cachon-Enjumet
<i>variabilis</i> Schiller	<i>Preperidinium</i>
<i>viride</i> Schiller	<i>meunieri</i> (Pavillard) Elbrächter
<i>Pachydinium</i>	<i>pseudooblea</i> (Elbrächter) Elbrächter
<i>mediterraneum</i> Pavillard	<i>Pronostiluca</i>
<i>Palaeophalacroma</i>	<i>pelagica</i> Fabre-Domerque, 1889
<i>verrucosum</i> Schiller	<i>spinifera</i> (Lohmann) Schiller
<i>Palaephalacroma</i>	<i>Prorocentrum</i>
<i>unicinctum</i> Schiller, 1928	<i>adriaticum</i> Schiller
<i>Parahistioneis</i>	<i>aporum</i> (Schiller, 1918) Dodge, 1975
<i>acutiformis</i> Rampi	<i>arcuatum</i> Isse
<i>karstenii</i> (Kofoid & Michener) Kofoid & Skogsberg	<i>balticum</i> (Lohmann) Loeblich, 1970
<i>mediterranea</i> Schiller	<i>cassubicum</i> (Woloszynska) Dodge
<i>paraformis</i> Kofoid & Skogsberg	<i>compressum</i> (Bail.) Abé ex Dodge
<i>Paulsenella</i>	<i>cornutum</i> Schiller
<i>chaetoceratis</i> (Paulsen) Chatton	<i>dactylus</i> (Stein) Dodge

- dentatum* Stein, 1883
gibbosum (Schiller) Schiller
gracile Schütt, 1895
lima (Ehrenberg) Stein
maximum (Gourret) Schiller
micans Ehrenberg
minimum (Pavillard) Schiller, 1933
nanum Schiller, 1918
oblongum (Schiller) Abé
obtusidens Schiller
ovum (Schiller) Dodge
pyriforme (Schiller) Taylor
redfieldii Bursa
rostratum Stein
rotundatum Schiller
schillerii Böhm
scutellum Schröder, 1900
triangulatum Martin
triestinum Schiller, 1918
vaginulum (Ehrenberg) Dodge
- Protoceratum*
areolatum Kofoid
pepo Kofoid & Michener
reticulatum (Clap. & Lach.) Bütschli
= *Gonyaulax grindleyi* Reinecke, 1967
= *Peridinium reticulatum* Clap. & Lach., 1859
= *Protoceratum aceros* Bergh, 1881
spinulosum (Murray & Whitting) Schiller
- Protoperidinium*
achromaticum (Levander) Balech
avellana (Meunier) Balech
biconicum (Dangeard) Balech
bipes (Paulsen) Balech
= *Glenodinium bipes* Paulsen, 1904
= *Minuscula bipes* Lebour, 1925
= *Peridinium minisculum* Pavillard, 1905
breve Paulsen
brevipes (Paulsen) Balech
brochii (Kofoid & Swezy) Balech
conicooides (Paulsen) Balech
conicum (Gran) Balech
crassipes (Kofoid) Balech
curlipes (Jørgensen) Balech
denticulatum (Gran & Braarud) Balech
depressum (Bailey) Balech
diabolus (Karsten) Balech
divergens (Ehrenberg) Balech
excentricum (Paulsen) Balech
grande (Kofoid) Balech
grani (Ostenfeld) Balech
groenlandicum (Wołoszynska) Balech
heterocanthum (Dangeard) Balech
inflatum Okamura
islandicum (Paulsen, 1904) Balech, 1973
laticeps (Grøntved & Seidenfaden, 1938) Balech, 1974
leonis (Pavillard) Balech
ligusticum (Rampi) Balech
fusiformis Balech
mariaelebouriae (Paulsen) Balech
mediterraneum (Kofoid) Balech
minutum (Loeblich) Kofoid
mite (Pavillard) Balech
monospinum (Paulsen) Zonefeld & Dale, 1994
murray Kofoid
nudum (Meunier) Balech
oblongum (Aurivillius+H282) Parke & Dodge
obtusum (Karsten) Parke & Dodge
oceanicum (Vanhöffen) Balech
ovatum Pouchet
oviforme (Dangeard) Balech
ovum (Schiller) Balech
pallidum (Ostenfeld) Balech
pellucidum Bergh
pendunculatum (Schütt) Balech
- pentagonum* (Gran) Balech
punctulatum (Paulsen) Balech
pyriforme (Paulsen) Balech
quarnerense (Schröder) Balech
saltans (Meunier) Balech
scilleri (Paulsen) Balech
simulum (Paulsen) Balech
solidcorne (Mangin) Balech
sphaericum (Murray & Whitting) Balech
sphaeroides (Dangeard) Balech
steinii (Jørgensen) Balech
subcurvipes (Lebour, 1923) Balech, 1974
subinerme (Paulsen) Loeblich III
thorianum (Paulsen) Balech
thulsesense (Balech) Balech
tregouboffii Halim
tristylum (Stein) Balech
wiesneri (Schiller) Balech
- Pseliodinium*
vaubanii Sournia
- Ptychodiscus*
noctiluca Stein, 1883
- Pyrocystis*
elegans Pavillard
fusiformis (Haeckel) Blackman
gerbaultii Pavillard
lanceolata Schröder
lunata (Schütt) Schütt
noctiluca Murray
obtusa Pavillard
robusta Kofoid
- Pyrophacus*
horologicum Stein
steinii (Schiller) Wall & Dale
- Roscoffia*
capitata Balech, 1956
- Sabulodinium*
undulatum Saunders & Dodge
- Scaphodinium*
mirabile Margalef
- Scripsiella*
crystallina Lewis
lachrymosa Lewis
minima Gao & Dodge
rotundata Lewis
trifida Lewis
trochoidea (Stein) Loeblich
- Scriptiella*
precaria Montressor & Zingone, 1988
ramonii Montressor, 1995
spinifera Honsell & Cabrini, 1991
subsalsa (Ostenfeld) Steidinger & Balech, 1977
- Sinophysis*
ebriolum (Herdman) Balech
- Spatulodinium*
pseudonoctiluca (Pouchet) Cachon & Cachon
- Spiraulax*
jollifei (Murray & Whitting) Kofoid
kofoidii Graham
- Styloodium*
gastrophilum Cachon, Cachon & Bouquaheux, 1965
- Sylloidinium*
listii Drebes
- Symbiodinium*
microadriaticum Freudenthal, 1962
- Thecadinium*
dragescoi Balech, 1956
kofoidii (Herdman) Larsen, 1985
= *Amphidinium kofoidii* var. *petasorum* Herdman 1922
= *Phalacroma kofoidii* Herdman, 1924
= *Thecadinium kofoidii* Schiller, 1933
= *Thecadinium petasorum* (Herdman) Kofoid & Skogsberg, 1928
neopetasatum Saunders & Dodge, 1984
- Thoracosphaera*

- heimii* (Lohmann) Kamptner
Torodinium
robustum Kofoed & Swezy
teredo (Pouchet) Kofoed & Swezy
Triposolenia
bicornis Kofoed
depressa Kofoed
truncata Kofoed
Warnowia
fusus Lindemann
Woloszynskia
pascieri (Suchlandt) Stosch
teuissima (Lauterborn) Thompson
Zygapikodinium
lenticulatum (Pauls.) Loeb. & Loeb.
= *Diplopeltopsis minor* Pavillard, 1913

Naked amoebae

Compiled by Andrew Rogerson and Andrew Goodkov
A comprehensive list of European species.

References

- Anderson O.R., Rogerson A. & Hannah F. 1997 Three new limax amoebae isolated from marine surface sediments: *Vahlkampfia caledonica*n.sp., *Saccamoeba marina* n.sp., and *Hartmannella vacuolata* n.sp. *Journal of Eukaryotic Microbiology* 44: 33-42.
- Bovee E.C. & Sawyer T.K. 1979 *Marine flora and fauna of the north-eastern United States. Protozoa: amoebae*. Washington: National Oceanographic and Atmospheric Administration. Technical Report 419
- Butler H. 1994 *The ecological importance of marine benthic naked amoebae* PhD Thesis, University of London
- Goodkov A.V. & Burkyakov V. Yu. 1987 A new species of limax amoeba (Sarcodina, Lobosea) from the White Sea. *Zool. Zh.* 66 (6): 928-930.
- Goodkov A.V. & Burkyakov V. Yu. 1988 *Mayorella dactylifera* sp.n. (Gymnamoebia, Paramoebidae) and brief review of marine species of mayorellas. *Zool. Zh.* 67 (6): 927-931.
- Grell K.G. 1994 *Reticulamoeba gemmipara* n. gen., n. sp., an amoeboflagellate with reticulopodia and zoospores. *Archives für Protistenkunde* 144: 55-61.
- Grell K.G. 1994 *Thalassomyxa canariensis* n.sp. from a tide pool of Tenerife (Canary Islands). *Archives für Protistenkunde* 144: 319-324.
- Grell K.G. 1994 The feeding community of *Synamoeba arenaria* n.gen., n.sp. *Archives für Protistenkunde* 144: 143-146.
- Grell K.G. 1995 *Reticulamoeba minor* n.sp. and its reticulopodia. *Archives für Protistenkunde* 145: 309.
- Hollande A., Nicolas G. & Escaig J. 1981 Vêtue glycostylaire et ultrastructure d'une Amibe marine libre (*Mayorella pussardi* nov. sp.: Paramoebidae) observée après congélation ultrarapide suivie de cryosubstitution. *Protistologica* 17: 147-154.
- Janicki C. 1912 Paramoebenstudien (*P. pigmentifera* Grassi und *P. chaetognatha* Grassi). *Z. wiss. Zool.* 103: 449-518.
- Kadiri G.E., Joyon L. & Pussard M. 1992 *Pernina chaumonti*, n.g., n.sp., a new marine amoeba (Rhizopoda, Heterolobosea). *European Journal of Protistology* 28: 43-50.
- Kuhn S.F. 1997 *Rhizamoeba schneppii* sp. nov., a naked amoeba feeding on marine diatoms (North Sea, German Bight). *Archives für Protistenkunde* 147: 277-282.
- Page F.C. 1983 *Marine Gymnamoebae* Institute of Terrestrial Ecology, Lavenham Press Ltd, Suffolk, UK54 pp
- Rogerson A. 1991 On the abundance of marine naked amoebae on the surfaces of five species of macroalgae. *FEMS Microbiology Ecology* 85: 301-312.
- Rogerson A. 1993 *Parvamoeba rugata* n.g., n.sp., (Gymnamoebia, Thecamoebidae): an exceptionally small marine naked amoeba. *European Journal of Protistology* 29: 446-452.
- Rogerson, A., Hannah F. & Anderson O.R. 1998 A redescription of *Rhabdamoeba marina* Dunkerly 1921, an inconspicuous marine amoeba from benthic sediments. *Invertebrate Biology* 117 (4): 261-270.
- Rogerson, A., Hannah F. & Gothe G. 1995 The grazing potential of some unusual marine benthic amoebae feeding on bacteria. *European Journal of Protistology* 34 (2): 271-279.
- Sawyer T.K. 1975 *Clydonella* n.g. (Amoebida: Thecamoebidae), proposed to provide an appropriate generic home for Schaeffer's marine species of *Rugipes*, *C. vivax* (Schaeffer, 1926) n.comb. *Transactions of the American Microscopical Society* 94: 395-400
- Sawyer T.K. 1975 Marine amoebae from surface waters of Chincoteague Bay, Virginia: two new genera and nine new species within the families Mayorellidae, Flabellulidae and Stereomyxidae *Transactions of the American Microscopical Society* 94: 71-92.
- Schaeffer A.A. 1926 Taxonomy of the amebas. *Washington: Carnegie Institution of Washington* 24: 3-116.
- Smirnov A.V. & Fenchel T. 1996 *Vahlkampfia anaeronica* n.sp. and *Vannella peregrinia* n. sp. (Rhizopoda) - anaerobic amoebae from marine sediment. *Archives für Protistenkunde* 147: 189-198.
- Smirnov A.V. 1996 *Stygamoeba regulata* n.sp (Rhizopoda) - a new amoeba with an unusual combination of light microscopical and ultrastructural features *Archives für Protistenkunde* 146: 299-307
- Smirnov A.V. 1997 Two new species of marine amoebae: *Hartmannella lobiferan*.sp., and *Korotnevella nivon*.sp. (Lobosea, Gymnamoebida). *Archives für Protistenkunde* 147: 238-292.
- Smirnov A.V. 1999 An atlas of gymnamoebae - Euamoebida and Leptomyxida, isolated from anaerobic sediments of the Nivo Bay (Baltic Sea, The Sound). *Ophelia* (in press)

AMOEBAE, NAKED

Family Acanthamoebidae

- Acanthamoeba gigantea* Schmoller, 1964
Protacanthamoeba caledonica Page, 1981

Family Amoebidae

- Saccamoeba admirata* Goodkov & Buryakov, 1987

Family Flabellulidae

- Flabellula baltica* Smirnov, 1999
demetica Page, 1980
trinovantica Page, 1980
calkinsi (Hogue, 1914) Page, 1983
= *Vahlkampfia calkinsi* Hogue, 1914
citata Schaeffer, 1926
Parflabellula hoguae (Sawer, 1975) Page & Willumsen, 1983
= *Flabellula hoguae* Sawyer, 1975
reniformis (Schmoller, 1964) Page & Willumsen, 1983
= *Flabellula reniformis* Schmoller, 1964

Family Hartmannellidae

- Hartmannella abertlawensis* Page, 1980
lobifera Smirnov, 1997
vacuolata Anderson, Rogerson & Hannah, 1997
Saccamoeba marina Anderson, Rogerson & Hannah, 1997

Family Leptomyxidae

- Rhizamoeba polyura* Page, 1972
saxonica Page, 1974
schneppii Kuhn, 1997

Family Paramoebidae

- Korotnevella nivo* Smirnov, 1997

Mayorella
dactylifera Goodkov & Buryakov, 1988
pussardi Hollande, Nicolas & Escaig, 1981
kuwaitensis (Page, 1981) Page, 1982
= *Hollandella kuwaitensis* Page, 1981
gemmifera Schaeffer, 1926

Paramoeba
eilhardi Schaudinn, 1896

Family Reticulomyxidae

Reticulamoeba
geminipara Grell, 1994
minor Grell, 1995

Thalassomyxa
canariensis Grell, 1994

Family Stereomyxidae

Corallomyxa
chattoni Grell & Benwitz, 1978

Stereomyxa
angulosa Grell, 1996
ramosa Grell, 1996

Family Thecamoebidae

Paramoeba
rugata Rogerson, 1993

Thecamoeba

hilla Schaeffer, 1926

mundula Schaeffer, 1926

orbis Schaeffer, 1926

pulchra Page, 1977

= *Amoeba pulchra* Biernacka, 1963

Family Vahlkampfiidae

Heteramoeba
clara Droop, 1962

Tetramitus
salinus Entz, 1883

Vahlkampfia

baltica Schmoller, 1961

damariscottae Page, 1974

longicauda Schmoller, 1964

trilaminata Schmoller, Jonas & Ludvik, 1983

anaerobica Smirnov & Fenchel, 1996

dumnonica Page, 1983

caledonica Anderson, Rogerson & Hannah, 1997

Family Vannellidae

Clydonella
rosenfieldi Sawyer, 1975

Platyamoeba
langae Sawyer, 1975

plurinucleolus Page, 1974

murchelanoi Sawyer, 1975

calycinucleolus Page, 1974

bursella Page, 1974

flabellata Page, 1974

Vannella

aberdonica Page, 1980

anglica Page, 1980

caledonica Page, 1979

devonica Page, 1979

peregrinia Smirnov & Fenchel, 1996

septentrionalis Page, 1980

platypodia Glaser, 1912

simplex Bovee, 1965

= *Hyalodiscus simplex* Wohlfarth-Botterman, 1960

Family Vexilliferidae

Neoparamoeba

aestuarina (Page, 1970) Page, 1987

= *Paramoeba aestuarina* Page, 1970

pemaquidensis (Page, 1980) Page, 1987

= *Paramoeba pemaquidensis* Page, 1970

Pseudoparamoeba

pagei (Sawer, 1975) Page, 1979

= *Vexillifera pagei* Sawyer, 1975

Vexillifera

armata Page, 1979

minutissima Bovee & Sawyer, 1979

telmathalassa Bovee, 1956

Family incertae sedis

Gruberella

flavescens (Gruber, 1889) Page, 1984

= *Amoeba flavescens* Gruber, 1889

Janickina

chaetognathi Grassi, 1881

pigmentifera Grassi, 1881

Nolandella

hibernica (Page, 1980) Page, 1983

= *Hartmannella hibernica* Page, 1980

Pernina

chaumonti Kadiri, Joyon & Pussard, 1992

Rhabdamoeba

marina Dunkerly, 1921

Stygameba

polymorpha Sawyer, 1975

regulata Smirnov, 1996

Synamoeba

arenaria Grell, 1994

Trichosphaerium

sieboldi Schneider, 1878

Testate amoebae

Compiled by Ralf Meisterfeld

A comprehensive list of European species.

References

- Anderson, O.R., Rogerson, A. & Hannah, F. 1996 A description of the Testate Amoeba *Ovulina parva* gen.nov., sp.nov. from coastal marine sediments. *J. mar. biol. Ass. U.K.* 76, 851-865
- Biernacka, I. 1962 Die Protozoenfauna in der Danziger Bucht. I. Die Protozoen in einigen Biotopen der Seeküste. *Polsk. Arch. Hydrobiol.* 10, 39-109
- Cash, J.; Wailes, G.H. & Hopkinson, J. 1915 *British Freshwater Rhizopoda and Heliozoa III* Ray Soc, London
- Chardez, D. 1971 Etude sur les thécamoebiens des biotopes interstitiels, psammons littoraux et zones marginales souterraines des eaux douces. *Bull. Rech. Agr. Gembloux* 6, 3-4, 257-268.
- Chardez, D. 1972 *Voluta hemispiralis*. Thécamoebien psammophile nouveau. *Rev. Verv. Hist. Nat.* 29, 10-12,
- Chardez, D. 1977 Thécamoebiens du mesopsammon des Plages de la Mer du Nord. *Rev. Verv. Hist. Nat.* 34, 4-6, 18-34.
- Chardez, D. 1978 *Hyalosphenia savoiei* sp. nov. (Thécamoebien mesopsammique). *Rev. Verv. Hist. Nat.* 35.
- Chardez, D. 1984 Study on mesopsammal testate amoebae. *Acta Protozool.* 23, 4, 247-253.
- Chardez, D. 1986 Thécamoebiens des plages de la Mer du Nord en Angleterre. *Acta Protozool.* 25, 4, 375-378.
- Chardez, D. 1993 Thécamoebiens mésopsammiques du littoral de la Mer du Nord. *Notes fauniques de Gembloux* 26, 3-7.
- Chardez, D. & Thomas, R. 1980 Meso-Psammon Thecamoeba from beaches of Lacanau and Leporge Ocean (Gironde, France) - (Protozoa, Rhizopoda, Testacea). *Acta Protozool.* 19, 3, 277-285.
- Coûteaux, M.-M. & Golemansky, V. 1984 Premières observations sur les Thécamoebiens interstitiels du supralittoral marin des Antilles françaises. *Bull. Mus. natn. Hist. nat.*, Paris 6, Sect. A, 263-277
- De Saedeleer 1934 Beitrag zur Kenntnis der Rhizopoden morphologische und systematische Untersuchungen und ein Klassifikationsversuch *Verh. Koninkl. nat. Mus. Beige*, 60, 1-112
- Decloitre, L. 1977 Thécamoebiens des bords de mer zone littorale et supralittorale Six-Fours-les-Plages (Var) *Ann. Soc. Sci. Nat. Archéol. Toulon et Var.* 1977, 168-170
- Golemansky, V. 1967 Matériaux sur la systématique et l'écologie des Thécamoebiens (Protozoa, Rhizopoda) du lac d'Ohrid *Sect. Sci. Nat. Univ. Skopje* 14, 4, 3-26

- Golemansky, V. 1970 *Chardezia caudata* gen. n. sp.n. et *Rhumbleriella filosa* gen. n. sp. n. - deux Thécamoebiens nouveaux du Psammon littoral de la Mer Noire. *Bull. Inst. Zool. Mus.* 32, 121-125.
- Golemansky, V. 1970 Contribution à la connaissance des Thécamoebiens (Rhizopoda, Testacea) des eaux souterraines littorales du Golfe de Gdansk (Pologne). *Bull. Inst. Zool. Mus.* 31,32, 77-87.
- Golemansky, V. 1970 Rhizopodes nouveaux du psammon littoral de la Mer Noire. *Protistologica* 6, 4, 365-371.
- Golemansky, V. 1970 Thécamoebiens (Rhizopoda, Testacea) des eaux souterraines littorales de quelques Plages de Cuba. *Bull. Inst. Mus.Zool.* 32, 151-158.
- Golemansky, V. 1970 Thécamoebiens (Rhizopoda, Testacea) nouveaux des eaux souterraines littorales de la Mer Noire. *Acta Protozool.* 8,2, 41-46.
- Golemansky, V. 1971 Taxonomische und zoogeographische Notizen über die Thekamöbe Fauna (Rhizopoda, Testacea) der Küstengrundwässer der sowjetischen Fernostküste (Japanisches Meer) und der Westküste Kanadas (Stiller Ozean). *Arch. Protistenkd.* 113, 235-249.
- Golemansky, V. 1973 Deuxième contribution à la connaissance des Thécamoebiens (Rhizopoda, Testacea) du psammal littoral de la Mer Baltique. *Bull. Inst. Zool. Mus.* 38, 49-60.
- Golemansky, V. 1973 *Messemvriella filosa* n.gen. n.sp. - une nouvelle thécamoebienne psammobionte (Rhizopoda, Testacea) des eaux souterraines littorales de la Mer Noire. *Zool. Anz.* 190, 5-6, 302-304.
- Golemansky, V. 1974 *Lagenidiopsis valkanovi* gen. n., sp. n. - un nouveau thécamoebien (Rhizopoda, Testacea) du psammal supralittoral des mers. *Acta Protozool.* 8, 1,1-4.
- Golemansky, V. 1976 Contribution à l'étude des Rhizopodes et des Heliozoaires du psammal supralittoral de la Méditerranée. *Acta Protozool.* 15, 1, 35-45.
- Golemansky, V. 1979 *Cyphoderia compressa* sp. n. (Rhizopoda, Arcellinida) - un nouveau thécamoebien psammobionte de supralittoral des mers. *Acta Protozool.* 18, 3, 429-434.
- Golemansky, V. 1981 Description de trois thécamoebiens (Protozoa, Rhizopodae) nouveaux des eaux souterraines littorales des mers. *Acta Protozool.* 20, 1, 115-119.
- Golemansky, V. 1982 Thécamoebiens (Protozoa, Rhizopodea) interstitiels du supralittoral Grec de la Mer Egée. *Biol. Gallo-Hellenica* 9, 2, 271-276.
- Golemansky, V. 1974 Sur la composition et la distribution horizontale de l'association thécamoebienne (Rhizopoda, Testacea) des eaux souterraines littorales de la Mer Noire en Bulgarie. *Bull. Inst. Zool. Mus.* 40, 195-202.
- Golemansky, V. & Coûteaux, M-M. 1982 Etude en microscopie électronique à balayage de huit espèces de thécamoebiens interstitiels du supralittoral marin. *Protistologica* 18, 4, 473-480.
- Golemansky, V. & Ogden, C.G. 1980 Shell structure of three littoral species of testate amoebae from the Black Sea (Rhizopoda, Protozoa). *Bull. Br. Mus. Nat. Hist. (Zool.)* 38, 1, 1-6.
- Golemansky, V.G. 1982 Révision du genre *Ogdeniella* nom. n. (= *Amphorellopsis* Golemansky, 1970) (Rhizopoda, Gromida) avec considérations sur son origine et évolution dans le milieu interstitiel. *Acta Zool. Bulg.* 19, 3-12.
- Golemansky, V.G. 1983 Interstitial Thecamoebas (Rhizopoda, Arcellinida and Gromida) of the Soviet Finn Bay Coast. *Acta Zool. Bulg.* 23, 3-8.
- Golemansky, V.G. 1990 Interstitial testate amoebas (Rhizopoda, Testacea) from the Mediterranean basin. *Stylogolia* 5, 1, 49-54.
- Golemansky, V.G. 1991 Mesopsammal Testate Amoebas (Rhizopoda, Arcellinida, Gromida and Monothalamida) from the Atlantic Sublittoral Near Roscoff (France). *Arch. Protistenkd.* 140, 1, 35-43.
- Golemansky, V.G. 1992 Thécamoebiens interstitiels (Rhizopoda, Arcellinida, Gromida et Monothalamida) du supralittoral français de l'Atlantique dans la région du Roscoff (Bai de Morlaix). *Acta Zool. Bulg.* 45, 3-13.
- Golemansky, V.G. 1998 Interstitial Testate Amoebae (Rhizopoda: Arcellinida and Gromida) from the Finnish Coast of the Baltic Sea and Summary Check List of the Interstitial Testate Amoebae in the Baltic Sea. *Acta Protozoologica* 37, 133-137
- Golemansky, V.G. 1998 Interstitial Testate Amoebae (Rhizopoda: Testacea) from the Italian Coast of the Mediterranean Sea. *Acta Protozoologica* 37, 139-144
- Grospeitsch, T. 1965 Monographische Studie der Gattung *Hylosphenia* Stein (Rhizopoda, Testacea). *Hydrobiologia* 26, 1-2, 211-241
- Jepps, M. W. 1926 Contribution to the study of *Gromia oviformis* Dujardin. *Quart. J. microscop. Sci.* 70, 701-719
- Johnson, P.W., Hargraves, P.E. & Sieburth, J.MCN. 1988 Ultrastructure and ecology of *Calycomonas ovalis* Wulff, 1919, Chrysophyceae) and its redescription as a testate Rhizopod, *Paulinella ovalis* n. comb. (Filosea, Euglyphina). *J. Protozool.* 35, 4, 618-626
- Laminger, H. 1973 Notes on some Testacea (Protozoa, Rhizopoda) of the Yugoslavian coast of the Adria near Rab. *Hydrobiologia* 42, 1, 153-154
- Ogden, C.G. & Hedley, R.H. 1980 *An atlas of freshwater testate amoebae*. British Museum & Oxford University Press
- Ogden, C.G. & Coûteaux, M-M. 1986 The nature of the shell of *Alepiella tricornuta* a marine testate amoeba (Sarcodina, Rhizopoda). *Protistologica* 22, 2, 213-220.
- Ogden, C.G. & Coûteaux, M-M. 1989 Interstitial marine rhizopods (Protozoa) from littoral sands on the east coast of England. *Europ. J. Protistol.* 24, 281-290.
- Pankow, H. 1982 *Paulinella chromatophora* Lauterb., eine bisher nur im Süßwasser nachgewiesene Thekamöbe, in den Boddengewässern des Darß und des Zingst (südliche Ostsee). *Arch. Protistenkd.* 126, 261-263.
- Sheehan, R. ; Banner, F.T 1973 Trichosphaerium - an extraordinary testate rhizopod from coastal waters. *Estuar. Coast. Mar. Sci.* 1, 245-260
- Valkanov, A. 1936 Notizen über die Brackwässer Bulgariens. 2. Versuch einer hydrographischen und biologischen Erforschung derselben. *Annuaire Univ. Sofia. III. Fac. Physico-Math. Livre 3 (Sci. Nat.)*, 209-341.
- Valkanov, A. 1970 Beitrag zur Kenntnis der Protozoen des Schwarzen Meeres. *Zool. Anz.* 184, 3-4, 241-290
- Vørs, N 1992 Heterotrophic Amoebae, Flagellates and Heliozoa from the Twärminne Area, Gulf of Finland, in 1988-1990. *Ophelia* 36, 1-109.
- Vørs, N 1993 Marine heterotrophic amoebae, flagellates and heliozoa from Belize (Central America) and Tenerife Canary Islands, with descriptions of new species. *Luffisphaera bulbochaete* N.Sp., *L. longihastis* N.Sp., *L. turiformis* N.Sp. and *Paulinella intermedia* N.Sp. *J. Euk. Microbiol.*, 40, 3: 272-87

AMOEBAE, TESTATE

Order Trichosida

Family Trichosidae

Trichosphaerium sieboldi Schneider, 1878

Order Arcellinida

Suborder Arcellina

Family Microchlamyiidae

Microchlamys patella (Claparède & Lachmann, 1859) Cockerell, 1911
= *Pseudochlamys patella* Claparède & Lachmann, 1859

Family Arcellidae

Antarcella atava Collin, 1914
Arcella hemisphaerica Perty, 1852
vulgaris Ehrenberg, 1832
Pyxidicula cymbalum Penard, 1902

Suborder Diffugina

Family Centropyxidae

Centropyxis

- aculeata* (Ehrenberg, 1832) Stein, 1859
 = *Arcella aculeata* Ehrenberg, 1832
aerophila Deflandre, 1929
constricta (Ehrenberg, 1841) Deflandre, 1929
 = *Arcella constricta* Ehrenberg, 1841
minuta Deflandre, 1929
platystoma (Penard, 1890) Deflandre, 1929
 = *Difflugia platystoma* Penard, 1890
- Family Centropyxidae ?**
- Centropyxiella*
arenaria Valkanov, 1970
elegans Valkanov, 1970
gibbula Valkanov, 1970
gibbulina Chardez & Thomas, 1980
golemanskyi Chardez, 1977
lucida Golemansky, 1971
oopyxiformis Chardez, 1977
platystoma Golemansky, 1981
- Family Difflugiidae**
- Difflugia*
algaephila Decloitre, 1977
linearis (Penard, 1890) Gauthier & Lièvre, 1958
 = *Difflugia pyriformis* var. *linearis* Penard, 1890
lithophila (Penard, 1902) Gauthier-Lièvre & Thomas, 1958
 = *Difflugia hydrostatica* var. *lithophila* Penard, 1902
lucida Penard, 1890
submarina (Chardez, 1980) Chardez & Thomas, 1980
subterranea Golemansky, 1970
 = *Difflugia subterranea* f. *circularis* Chardez, 1971
- Family Hyalospheniidae**
- Hyalosphenia*
cuneata Stein, 1857
minuta Cash, 1892
savoiei Chardez, 1978
- Family Lesquereliidae**
- Pomoriella*
valkanovi Golemansky, 1970
- Family Trigonopyxidae**
- Cyclopaxis*
eurystoma Deflandre, 1929
- Suborder Phryganellina**
- Family Cryptodifflugiidae**
- Cryptodifflugia*
collum (Chardez, 1971)
 = *Difflugiella collum* Chardez, 1971
lamingerae (Chardez, 1977)
 = *Difflugiella lamingerae* Chardez, 1977
lanceolata Golemansky, 1970
paludosa Golemansky, 1981
psammophila (Golemansky, 1970)
 = *Difflugiella psammophila* Golemansky, 1970
- Wailesella*
eboracensis (Wailes & Penard, 1911) Deflandre, 1928
 = *Cryptodifflugia eboracensis* Wailes & Penard, 1911
- Family Phryganellidae**
- Phryganella*
marinus Chardez, 1971
- Order Gromida**
- Family Gromiidae**
- Gromia*
oviformis Dujardin, 1835
- Family Pseudodifflugiidae**
- Ovulina*
parva Anderson, Rogerson & Hannah, 1996
- Pseudodifflugia*
andreevi Golemansky, 1976
delamarei Coûteaux & Golemansky, 1984
- Family Psammonobiotidae**
- Alepiella*
tricornuta Golemansky, 1970
- Chardezia*
caudata Golemansky, 1970
- Micramphora*
amphoriformis Chardez & Thomas, 1980
atlantica Chardez & Thomas, 1980
pontica Valkanov, 1970
tokioensis Sudzuki, 1979
- Micropsammella*
infralitoralis Chardez, 1993
minima Chardez & Thomas, 1980
retorta Golemansky, 1970
- Ogeniella*
carinata (Chardez, 1977) Golemansky, 1982
 = *Amphorellopsis carinata* Chardez, 1977
elegans (Golemansky, 1970) Golemansky, 1982
 = *Amphorellopsis elegans* Golemansky, 1970
lucida (Golemansky, 1970) Golemansky, 1982
 = *Amphorellopsis lucida* Golemansky, 1970
 = *Micramphora hellebauti* Chardez, 1971
maxima (Golemansky, 1970) Golemansky, 1982
 = *Amphorellopsis maximus* Golemansky, 1970
pusilla (Chardez, 1977) Golemansky, 1982
 = *Amphorellopsis pusilla* Chardez, 1977
septentrionalis Golemansky, 1982
taschevi (Golemansky, 1970) Golemansky, 1982
 = *Amphorellopsis taschevi* Golemansky, 1970
- Psammonobiotus*
balticus Golemansky, 1973
communis Golemansky, 1967
golemanskyi Chardez, 1971
linearis Golemansky, 1970
minutus Golemansky, 1970
plana Chardez, 1971
septentrionalis Chardez, 1977
- Order Euglyphida**
- Family Cyphoderidae**
- Campascus*
interstitialis Golemansky, 1981
vulgaris Valkanov, 1936
- Corythionella*
acolla Golemansky, 1970
anteroplanata Chardez, 1977
minima Golemansky, 1970
pontica Golemansky, 1970
sudzukii Chardez, 1977
- Cyphoderia*
ampulla (Ehrenberg, 1840) Leidy, 1878
 = *Cyphoderia margaritacea* Schlumberger, 1845
 = *Difflugia ampulla* Ehrenberg, 1840
compressa Golemansky, 1979
littoralis Golemansky, 1973
- Messemvriella*
filosa Golemansky, 1973
- Pseudocorythion*
acutum (Wailes, 1927) Valkanov, 1970
 = *Corythion acutum* Wailes, 1927
mannei Chardez, 1971
undulacollis Chardez & Thomas, 1980
wailesi Golemansky, 1971
- Family Euglyphidae**
- Assulina*
muscorum Greeff, 1888
- Euglypha*
compressa Carter, 1864
laevis Perty, 1849
rotunda Wailes & Penard, 1911
tuberculata Dujardin, 1841
- Tracheuglypha*
dentata (Vejdovsky, 1882) Deflandre, 1928
 = *Euglypha beta* Vejdovsky, 1882
- Family Paulinellidae**
- Paulinella*
chromatophora Lauterborn, 1895
intermedia Vørs, 1993

ovalis Johnson, Hargraves & Sieburth, 1988

Family Trinematidae

Trinema

- complanatum* Penard, 1890
- enchelys* (Ehrenberg, 1938) Leidy, 1878
- = *Difflugia enhelys* Ehrenberg, 1838
- = *Trinema acinus* Dujardin, 1841
- lineare* Penard, 1890

Filose testate amoebae incertae sedis

Family Amphitremidae ?

- Paramphitrema*
- pontica* Valkanov, 1970

Family Volutellidae

- Volutella*
- hemispiralis* (Chardez, 1972) Chardez, 1974
- = *Voluta hemispiralis* Chardez, 1972

Family incertae sedis

- Euglyphinopsis*
- pontica* Valkanov, 1970
- Lesquerella*
- mesopsammophila* Chardez & Thomas, 1980
- Rhumblierella*
- filosa* Golemansky, 1970
- Sudzukiella*
- marina* Golemansky, 1991

Apusomonads

Compiled by Sue Brandt

Preliminary list covering most records to 1995.

References

- Mylnikov, A.P. 1989 The ultrathin structure of the flagellate
Amastigomonas caudata (in Russian) *Tsitologiya* 31: 489-491
- Thomsen, H.A. (ed.) 1992 *Plankton i de indre danske farvande*
Miljøministeriet, Miljøstyrelsen, Copenhagen.

APUSOMONADS

- Amastigomonas*
- debruynei* De Saedeleer, 1931
- = *Thecamonas trahens* Larsen & Patterson, 1990
- mutabilis* (Griessmann, 1913) Patterson & Zöllfel, 1991
- = *Rhynchomonas mutabilis* Griessmann, 1913
- = *Thecamonas mutabilis* Larsen & Patterson, 1990
- caudata* Mylnikov, 1989

Choanoflagellata

Compiled by Sue Brandt

The choanoflagellates are divided into three taxa on the basis of the composition of the covering surrounding the cell. The Acanthoecidae, with a siliceous lorica, are particularly conspicuous in marine waters. No members of this taxon have been reported from freshwater, although some species may occur in brackish water. Members of the Codosigidae ("naked" cells, although a thin investment is sometimes visible, particularly with electron microscopy) and the Salpingoecidae (with organic thecae) appear to be more diverse in freshwaters. However, the identity of many species within these two families is in great need of revision.

The list below includes a comprehensive list of the Acanthoecidae, produced from recent literature. The lists of the Codosigidae and Salpingoecidae are preliminary, covering most records to 1995.

References

Boucaud Camou, E. 1967 Les Choanoflagellés des Côtes de la Manche: I. Systématique *Bull. Soc. Linn. Normandie*, 10: 191-209

Ellis, W.N. 1930 Recent researches on the Choanoflagellata (Craspedomonadines) (freshwater and marine) with description of new genera and species *Annls. Soc. R. Zool. Belg.*, 60: 49-88

Espeland, G. & Thronsen, J. 1986 Flagellates from Kilsfjorden, Southern Norway, with description of two new species of Choanoflagellida *Sarsia* 71: 209-226

Leadbeater, B.S.C. 1973 External morphology of some marine choanoflagellates from the coast of Jugoslavia *Arch. Protistenk.*, 115: 234-252

Leadbeater, B.S.C. 1980 Four new species of loricate choanoflagellates from South Brittany, France *Cahiers de Biologie Marine* 21: 345-353

Parke, M. & Leadbeater, B.S.C. 1977 Check-list of British marine Choanoflagellida - second revision *J. mar. biol. Ass. U.K.* 57: 1-6

Thomsen, H.A. (ed.) 1992 *Plankton i de indre danske farvande* Miljøministeriet, Miljøstyrelsen, Copenhagen.

Thronsen, J. 1974 Planktonic choanoflagellates from North Atlantic waters *Sarsia* 56: 96-122

Tong, S.M. 1997 Choanoflagellates in Southampton Water, including the description of three new species. *J. mar. biol. Ass. U.K.* 77: 929-958.

Tong, S.M., Nygaard, K., Bernard, C., Vørs, N & Patterson, D.J. 1998 Heterotrophic flagellates from the water column in Port Jackson, Sydney, Australia *Europ. J. Protistol.*, 34: 162-194

Vørs, N. 1992 Heterotrophic amoebae, flagellates and heliozoa from the Tvarminne area, Gulf of Finland, in 1988-1990 *Ophelia* 36: 1-109

CHOANOFAGELLATA

Family Acanthoecidae

Acanthocoris

- apoda* (Leadbeater, 1972) Hara & Takahashi, 1984
- = *Acanthoecopsis apoda* Leadbeater, 1972
- asymmetrica* (Thomsen, 1979) Hara & Takahashi, 1984
- = *Acanthoecopsis asymmetrica* Thomsen, 1979
- = *Acanthoecopsis spiculifera* sensu Thomsen, 1977
- campanula* (Espeland, 1986) Thomsen in Thomsen et al., 1991
- = *Acanthoecopsis campanula* Espeland, 1986
- haurakiana* Thomsen in Thomsen et al., 1991
- unguiculata* (Thomsen, 1973) Hara & Takahashi, 1984
- = *Acanthoecopsis unguiculata* Thomsen, 1973
- = *Pseudomicrosportalora ornata* Scagel & Stein, 1961

Acanthoeca

- brevipoda* Ellis, 1930
- spectabilis* Ellis, 1930

Amoenoscopra

- caudata* Hara & Takahashi, 1987

Bicosta

- antennigera* Moestrup, 1979
- minor* (Reynolds, 1976) Leadbeater, 1978
- = *Salpingoeca minor* Reynolds, 1976
- = *Salpingoeca virgata* Parke & Leadbeater, 1977
- spinifera* (Thronsen, 1970) Leadbeater, 1978
- = *Salpingoeca spinifera* Thronsen, 1970

Calliantha

- longicaudata* (Leadbeater, 1975) Leadbeater, 1978
- = *Salpingoeca longicaudata* Leadbeater in Manton et al., 1975

- multispina* Manton & Oates, 1979

- natans* (Grøntved, 1956) Leadbeater, 1978

- = *Salpingoeca natans* Grøntved, 1956

- simplex* Manton & Oates, 1979

Campanoeca

- dilatata* Thronsen, 1974

Campyloacantha

- spinifera* (Leadbeater, 1973) Hara & Takahashi, 1987
- = *Parvicorbicula spinifera* Leadbeater, 1973

- Cosmoeca**
norvegica Thomsen in Thomsen & Boonruang., 1984
ventricosa Thomsen in Thomsen & Boonruang., 1984
= *Pleurasiga oculaeformis* Schiller, 1925
- Crinolina**
isefjordensis Thomsen, 1976
- Crucispina**
cruciformis (Leadbeater, 1974) Espeland in Espeland & Thronsen, 1986
= *Salpingoeca cruciformis* Leadbeater, 1974
- Diaphanoeca**
grandis Ellis, 1930
pedicellata Leadbeater, 1972
= *Campanoeca pedicellata* (Leadbeater, 1972) Thronsen, 1974
sphaerica Thomsen, 1982
undulata Thomsen, 1982
- Diplotheca**
costata Valkanov, 1970
- Monocosta**
fennica Thomsen, 1979
- Nannoeca**
minuta (Leadbeater, 1972) Thomsen, 1988
= *Pleurasiga minima* Thronsen, 1970 var. *minuta* Leadbeater 1972
- Parvicorbicula**
aculeatus Tong, 1997
circularis Thomsen, 1976
manubriata Tong, 1997
pedicellata Leadbeater, 1973
pedunculata Leadbeater, 1980
quadricostata Thronsen, 1970
socialis (Meunier, 1910) Deflandre, 1960
= *Corbicula socialis* Meunier, 1910
superpositus Booth, 1990
- Platyleura**
infundibuliformis (Leadbeater, 1974) Thomsen in Thomsen & Boonruang, 1983
= *Parvicorbicula infundibuliformis* Leadbeater, 1974
- Pleurasiga**
echinocostata Espeland in Espeland & Thronsen, 1986
= *Pleurasiga reynoldsi* Thronsen aff. Leadbeater, 1973
minima Thronsen, 1970
reynoldsi Thronsen, 1970
- Polyfibula**
sphyrelata (Thomsen, 1973) Manton in Manton & Bremer, 1981
= *Pleurasiga caudata* Leadbeater, 1975
= *Pleurasiga sphyrelata* Thomsen, 1973
= *Polyfibula caudata* Manton in Manton & Bremer, 1981
= *Polyfibula hexacostata* Manton in Manton & Bremer, 1981
= *Polyfibula stipitata* Manton in Manton & Bremer, 1981
- Polyoeca**
dichotoma Kent, 1881
= *Acanthoecopsis spiculifera* Norris, 1965
= *Polyoeca dumosa* Dunkerley, 1910
- Saepicula**
pulchra Leadbeater, 1980
- Saroea**
attenuata Thomsen, 1979
- Savillea**
micropora (Norris, 1965) Leadbeater, 1975
= *Pleurasiga micropora* Norris, 1965
parva (Ellis, 1930) Loeblich III, 1967
= *Diaphanoeca parva* Ellis, 1930
= *Ellisiella parva* Norris, 1965
- Stephanacantha**
campaniformis (Leadbeater, 1973) Thomsen in Thomsen & Boonruang, 1983
= *Parvicorbicula campaniformis* Leadbeater, 1973
- Stephanoeca**
ampulla (Kent, 1880) Ellis, 1930
= *Salpingoeca ampulla* Kent, 1880
- aphelis* Thomsen, Buck & Chavez, 1991
campanula (Kent, 1880) Boucard Camou, 1967
= *Salpingoeca campanula* Kent, 1880
cauliculata Leadheater, 1980
complexa (Norris, 1965) Thronsen, 1974
= *Pleurasiga complexa* Norris, 1965
constricta Ellis, 1930
cupula (Leadbeater, 1972) Thomsen, 1988
= *Pleurasiga cupula* Leadbeater, 1972
diplocostata Ellis, 1930
= *Stephanoeca pedicellata* Leadbeater, 1972
elegans (Norris, 1965) Thronsen, 1974
= *Pleurasiga elegans* Norris, 1965
kentii Ellis, 1930
norrisii Thomsen, 1973
= *Sportelloeca kentii* Norris, 1965
paucicostata Thronsen, 1969
= *Stephanoeca diplocostata* Ellis, 1930 subsp.
paucicostata Thronsen, 1969
pyxidoides Leadbeater, 1980
supracostata Hara, Lee Chen, Sheu & Takahashi, 1996
urnula Thomsen, 1973
- Family Codosigidae**
- Codonocladium**
cymosum (Kent, 1880) Boucard Camou, 1967
- Codosiga**
botrylis (Ehrenberg) Kent, 1880
gracilis (Kent, 1880) De Saedeleer, 1927
= *Monosiga gracilis* Kent, 1880
pyriformis Kent, 1880
- Desmarella**
moniformis Kent, 1880
- Dicraspedella**
stokesi Ellis, 1930
- Kentrosiga**
echina Tong, 1997
thienemanni Schiller, 1953
- Monosiga**
brevicollis Ruinen, 1938
lagenula Thronsen, 1969
marina Grøntved, 1952
micropelagica Thronsen, 1974
ovata Kent, 1880
singuosa Möbius
- Proterospongia**
dybosensis (Grøntved, 1956) Loeblich III
nana (Braarud, 1935) Thronsen, 1974
= *Protospongia nana* Braarud, 1935
- Family Salpingoecidae**
- Choanoeca**
perplexa Ellis, 1930
- Diploeca**
placita Ellis, 1930
- Pachysoeca**
depressa Ellis, 1930
longicollis Ellis, 1930
- Salpingoeca**
aggregata Valkanov, 1970
amphoridium James-Clark, 1867
= *Salpingoeca ampullacea* Stein, 1878
= *Salpingoeca ampulloides* Bicudo & Bicudo, 1983
collaris Stokes, 1894
cornuta Kent, 1880
huxleyi Ellis, 1930
infusionum Kent, 1880
= *Salpingoeca longipes* Kent, 1880
inquillata Kent, 1880
= *Salpingoeca curvipes* Kent, 1880
= *Salpingoeca ringens* Kent, 1880
marina James-Clark, 1867
megacheila Ellis, 1930
napiformis Kent, 1880
= *Salpingoeca desaedeleeri* Ellis, 1930
procera Möbius

stenotheaca Boucard Camou, 1967
teres Kent, 1880
tuba Kent, 1880
= *Salpingoeca petiolata* Kent, 1880
= *Salpingoeca vaginalis* sensu Dunkerly, 1910
undulata Ellis, 1930
urceolata Kent, 1880
= *Salpingoeca cardiformis* Ellis, 1930

Cryptophytes

Compiled by Sue Brandt

A preliminary list, based mainly on a survey of plankton in Danish waters.

References

- Kent, W.S. 1880 *A manual of the Infusoria*. (1880-1882) Bogue, London.
Larsen, J. & Patterson, D.J. 1990 Some flagellates (Protista) from tropical marine sediments. *Journal of Natural History*, 24: 801-937.
Thomsen, H.A. (ed.) 1992 *Plankton i de indre danske farvande* Miljøministeriet, Miljøstyrelsen, Copenhagen.

CRYPTOPHYTES

Chilomonas
amygdalum Kent, 1880
Chroomonas
baltica (Büttner) Carter
vectensis Carter
Goniomonas
amphinema (Fresenius) Larsen & Patterson, 1990
pacificica Larsen & Patterson, 1990
truncata (Fresenius) Stein, 1878
Hemiselmis
virescens Droop
Plagioselmis
prolonga Butcher
Platychilomonas
psammobia Larsen & Patterson, 1990
Rhodomonas
baltica Karsten
= *Cryptomonas baltica* (Karsten) Butcher
= *Cryptomonas pseudobaltica* Butcher
marina (Dangeard) Lemmermann
= *Chroomonas marina* Butcher
= *Cryptomonas marina* Büttner
= *Cryptomonas marina* Dangeard
= *Rhodomonas pelagica* Lohmann
salina (Wislouch) Hill et Wetherbee
= *Chroomonas salina* (Wislouch) Butcher
= *Cryptomonas salina* Wislouch
= *Pyrenomonas salina* (Wislouch) Santore
Teleaulax
acuta (Butcher) Hill
= *Cryptomonas acuta* Butcher
= *Cryptomonas prora* Conrad et Kuferath
amphioxiae (Conrad) Hill
= *Chroomonas amphioxiae* (Conrad) Butcher
= *Rhodomonas amphioxiae* Conrad

Euglenozoa

Euglenoids (heterotrophic species)

Compiled by Sue Brandt

A preliminary list of heterotrophic species.

References

- Griessmann, K. 1913 Über marine Flagellaten. *Archiv. Protistenk.* 32: 1-78
Larsen, J 1987 Algal studies of the Danish Wadden Sea IV. A taxonomic study of the interstitial euglenoid flagellates *Nordic Journal of Botany*, 7: 589-607

Larsen, J. & Patterson, D.J. 1990 Some flagellates (Protista) from tropical marine sediments. *Journal of Natural History*, 24: 801-937.
Leadbeater, B.S.C 1972 Identification, by means of electron microscopy, of flagellate nanoplankton from the coast of Norway *Sarsia* 49: 107-124
Tong, S.M. 1997 Heterotrophic flagellates and other protists from Southampton Water, U.K. *Ophelia* 47: 71-131.

Kinetoplastids

Compiled by Sue Brandt

Comprehensive list produced from recent literature.

References

- Bernard, C., Simpson, A.G.B. & Patterson, D.J. 2000 Some free-living flagellates (Protista) from anoxic habitats. *Ophelia*, in press.
Elbrächter, M.L., Schnepp, E. & Balzer, I. 1996 *Hemistasia phaeocysticola* (Scherffel) comb. Nov., redescription of a free-living, marine, phagotrophic kinetoplastid flagellate. *Arch. Protistenk.* 147: 125-136
Ellis, W.N. 1930 Recent researches on the Choanoflagellata (Craspedomonadines) (freshwater and marine) with description of new genera and species *Annls. Soc. R. Zool. Belg.*, 60: 49-88
Larsen, J. & Patterson, D.J. 1990 Some flagellates (Protista) from tropical marine sediments. *Journal of Natural History*, 24: 801-937.
Ruijten, J. 1938 Notizen über Salzflagellaten II Über die Verbereitung der Salzflagellaten *Archiv für Protistenkunde*, 90, 210-258.
Schnepp, E. 1994 Light and electron microscopical observations in *Rhynchopus coscinodiscivorus* spec. nov., a colourless phagotrophic Euglenozoon with concealed flagella. *Arch. Protistenk.*, 144: 63-74
Thomsen, H.A. (ed.) 1992 *Plankton i de indre danske farvande* Miljøministeriet, Miljøstyrelsen, Copenhagen.
Tong, S.M. 1997 Heterotrophic flagellates and other protists from Southampton Water, U.K. *Ophelia* 47: 71-131.
Tong, S.M., Nygaard, K., Bernard, C., Vørns, N & Patterson, D.J. 1998 Heterotrophic flagellates from the water column in Port Jackson, Sydney, Australia *Europ. J. Protistol.*, 34: 162-194
Vørns, N. 1992 Heterotrophic amoebae, flagellates and heliozoa from the Tvarminne area, Gulf of Finland, in 1988-1990 *Ophelia* 36: 1-109

EUGLENOZOA

Euglenoids, heterotrophic

Anisonema
acinus Dujardin, 1841
prosgeobium Skuja, 1939
trepidum Larsen, 1987
Dinematomonas
litorale (Skuja) Silva, 1960
Euglena
deses Ehrenberg, 1830
Eutreptiella
gymnastica Thronsen, 1969
Heteronema
ovale Kahl, 1928
Metanema
sedatum Larsen, 1987
strenuum (Skuja, 1948) Larsen, 1987
Notoselenus
canellatus Skuja, 1948
esulcis Larsen, 1987
rhombeicus Larsen, 1987
= *N. obliquus* sensu Skuja, 1939
Petalomonas
abscissa (Dujardin) Stein, 1859
mediocanellata Stein, 1878
minuta Holland, 1942
= *P. angusta* (Klebs) Lemmermann, 1913 var. *pusilla* (Klebs) Lemmermann, 1913

= *P. mediocanellata* Stein, 1878 var. *pusilla* Klebs, 1892

= *P. minutula* Christen, 1962

pusilla Skuja, 1948

= *P. poosilla* Larsen & Patterson, 1990

Ploeoetia

corrugata Larsen & Patterson, 1990

= *P. currugata* Larsen & Patterson, 1990

vitrea Dujardin, 1841

Pseudoperanema

dolichonema Larsen & Patterson, 1990

= *Peranema dolichonema* Larsen & Patterson, 1990

fusiforme Larsen, 1987

trichophorum (Ehrenberg) Larsen, 1987

= *Peranema trichophorum* sensu Stein p.p., 1878

Urecoelus

costatus Stein ex Lemmermann, 1910

= *Phialonema cyclostomum* Stein p.p., 1878

cyclostomus (Stein) Mereschkowsky, 1881

= *Phialonema cyclostomum* Stein p.p., 1878

pascheri Skvortzow, 1924

= *U. gobii* Skvortzow, 1924

= *U. ovatus* Roskin, 1931

sabulosus Stokes, 1888

= *Urecoelopsis sabulosa* (Stokes) Stokes, 1887

= *U. ovatus* Möbius, 1888

Kinetoplastids

Bodo

curvifilus Griessmann, 1913

designis Skuja, 1948

saliens Larsen & Patterson, 1990

= *B. angusta* sensu Hänel, 1979

saltans Ehrenberg, 1832

= *Pleuromonas jaculans* Perty, 1962

Rhynchomonas

nasuta (Stokes, 1888) Klebs, 1892

Bodo

platyrrhynchus Larsen & Patterson, 1990

Hemistasia

phaeocystidicola (Scherffel, 1900) Elbrächter & Schneppf, 1996

= *Cryptaulax marina* Thronsen, 1969

= *H. klebsii* Griessmann, 1913

= *Oxyrrhis phaeocysticola* Scherffel, 1900

= *Pronoctiluca phaeocysticola* (Scherffel, 1900)

Pavillard, 1922

Diplonema

breviciliata Griessmann, 1913

Bodo

caudatus (Dujardin) Stein, 1883

edax Klebs, 1893

Rhynchopus

amitus Skuja, 1948

= *R. coscinodiscivorus* Schneppf, 1994

Diplonema

ambulator Larsen & Patterson, 1990

Rhynchobodo

taeniata (Skuja, 1956) Vørs, 1992

Foraminifera

Compiled by Onno Gross

Study of recent benthic Foraminifera in European waters began in the middle of the 18th century (e.g. Carpenter's paper (1868) on benthic Foraminifera of the Faeroe-Shetland Channel). Since then, many other scientists have contributed papers on the taxonomy, distribution, and ecology of these organisms. However, to my knowledge this is the first time an attempt has been made to summarise a species list of European Foraminifera.

During the course of the ERMS project a literature survey, and databases from universities and museums were used to establish such a species list. The database from which this checklist was compiled now holds 1233 species and 364 genera and includes another 431 synonyms of Foraminifera. The species fall into the four main suborders: 3% are members of the membranous or tectinous suborder Allogromiina, 28% are agglutinated species of Textulariina, 20% porcelaneous taxa of the suborder Miliolina and the majority (49%) are members of the suborder of hyaline calcareous Rotaliina (combining the "calcareous" suborders Fusulinina, Lagenina, Robertinina, Globigerinina, and others).

However, our knowledge of the biodiversity patterns of benthic and planktonic Foraminifera is still far from complete. The general difficulties of the foraminiferal species concept, the taxonomic uncertainties due to frequent synonyms, the present lack of knowledge of the diversity of soft-walled Foraminifera (e.g. Allogromiina) especially in the deep-sea, and the unknown species richness of foraminiferal communities in many geographic regions (e.g., along the southern Mediterranean Sea and North-African coast) have to be kept in mind when looking at this checklist. The catalogue awaits further inputs, and revision and should be seen as a baseline for future biodiversity studies.

The catalogue of Foraminifera benefited greatly from databases and comments of Thomas Cedhagen (University of Aarhus), Samuel Bowser (NY State Department of Health, Albany), Wolfgang Kuhnt and Jutta Wollenburg (University of Kiel), Bob Jones (BP Exploration Operating Company), John Murray (Southampton Oceanography Centre), Ralf Schiebel and Gerhard Schmiedl (University of Tübingen), Igor Dovgal (University of Kiev) and Dusan Zavodnik (Centre for Marine Research Rovinji, Croatia). I would like to thank these colleagues for their help.

References

- Brönnimann, P. & Whittaker, J.E., 1980, A revision of *Reophax* and its type species, with remarks on several other recent hormosinid species (Protozoa: Foraminiferida) in the collections of the British Museum. *Bull. Br. Mus. Nat. Hist.*, 39, p. 259-272.
- Brönnimann, P. & Whittaker, J.E., 1988, *The Trochamminacea of the Discovery Reports*: British Museum, London, 10 volumes, 152 p.
- Caralp, M.H., 1984, Quaternary calcareous benthic foraminifers, Leg 80: in de Graciansky, P.C. & Poag, C.W. et al. (eds): *Initial Reports of the Deep Sea Drilling Project*, Leg 80, Washington, 725-755.
- Cedhagen, T., 1991, *Foraminifera i Nordostatlanten och Arkis*, Manuscript, 50 p.
- Cedhagen, T., 1993, Taxonomy and biology of *Pelosina arborescens* with comparative notes on *Astrorhiza limicola* (Foraminiferida): *Ophelia*, v. 37, p. 143-162.

- Cimerman, F. & Langer, M.R., 1991, *Mediterranean Foraminifera*: Slovenska Akademija Znanosti, Ljubljana, 118 p.
- Gabel, B., 1971, Die Foraminiferen der Nordsee: *Helgoländer wiss. Meeresunters.*, 22, 1-65.
- Gooday, A.J. & Haynes, J., 1983, Abyssal foraminifers, including two new genera, encrusting the interior of *Bathysiphon rusticus* tubes: *Deep Sea Res.*, v. 30, p. 591-614.
- Gooday, A.J. & Nott, J.A., 1982, Intracellular Barite Crystals in two Xenophyophores, *Aschemonella ramuliformis* and *Galatheammina* sp. (Protozoa: Rhizopoda) with comments on the Taxonomy of *A. ramuliformis*: *J. Mar. Biol. Assoc. U.K.*, v. 62, p. 595-605.
- Gooday, A.J., 1986, The genus *Rhabdammina* in the northeast Atlantic: A new species, a redescription of *R. major* de Folin, 1887, and some speculations on species relationships: *J. Foram. Res.*, v. 16, p. 150-160.
- Gooday, A.J., 1988, The genus *Bathysiphon* (Protista, Foraminiferida) in the north east Atlantic: a neotype for *B. filiformis* G.O. Sars, 1872 and the description of a new species: *J. Nat. Hist.*, v. 22, p. 95-105.
- Gooday, A.J., 1990, *Tinogummia riemannii* sp. nov. (Allogromiina: Foraminiferida), a new species associated with organic detritus in the deep sea: *Bull. Br. Mus. nat. Hist. (Zool.)* 56, 93-103.
- Groß, O., 1991, *Die Verbreitung der Meiofauna und der Foraminiferen am untermeerischen Hang der Färöer Inseln*. Diploma thesis, University of Hamburg, Inst. Hydrobiologie und Fischereiwissenschaften, 147 p., 9 plt.
- Groß, O., 1998, *Investigations on autecology, migration and bioturbation of living benthic deep-sea Foraminifera (Protozoa)*, Berichte aus dem Zentrum für Meeres- und Klimaforschung, No. 15, University of Hamburg, 225 p., 21 plt.
- Gudmundsson, G., 1998, Distribution limits of Pyrgo species at the biogeographic boundaries of the Arctic and the North-Atlantic boreal regions, *J. Foram. Res.*, 28, p. 240-256.
- Haynes, J., 1973, Cardigan Bay Recent Foraminifera (Cruises of the R.V. Antur, 1962-1964), *Bull. Br. Mus. nat. Hist. (Zool.) Suppl.* 4, London.
- Hayward, B.W., Grenfell, H.R., Reid, C.M. & Hayward, K.A., 1999, Recent New Zealand shallow-water benthic Foraminifera. *Institut of Geological & Nuclear Science monograph* 21, 264 p.
- Hemleben, Ch., Spindler, M. & Anderson, O.R., 1989, *Modern Planktonic Foraminifera*. Springer Verlag, New York.
- Hermelin, J.O.R. & Scott, D.B., 1985, Recent benthic Foraminifera from the central North Atlantic: *Micropaleontology*, v. 31, p. 199-220.
- Höglund, 1947, Foraminifera in the Gullmar Fjord and the Skagerak, *Zoologiska Bidrag från Uppsala*, 26, 1-328.
- Hottinger, L., Halicz, E. & Reiss, Z., 1993, Recent Foraminifera from the Gulf of Aquaba, Red Sea, in *Slovenska Akademija Znanosti in Umetnosti*, 33, Ljubljana, 230 p.
- Jones, R.W., 1994, *The Challenger Foraminifera*, Oxford Science Publications, 149 p.
- Jorissen, F.J., 1988, Benthic Foraminifera from the Adriatic Sea; principles of phenotypic variation: *Utrecht Micropal. Bull.*, 37.
- Loeblich, A.R. & Tappan, H., 1988, *Foraminiferal genera and their classification*: Van Nostrand Reinhold Company, New York, pp. 970.
- Mikhalevich, V.I., 1968, *The keys for identification of the Black and Asov Sea fauna*. In 3 volumes. Voll Free-living invertebrates: Protists, sponges, coelenterates, vermes, tentaculifers. Kiev. Naukova dumka. 437 p. (e-mail 17.10.99 by Igor V. Dovgal).
- Moodley, L., 1992, *Experimental Ecology of Benthic Foraminifera in Soft Sediments and its (Paleo) Environmental Significance*; Diss. Univ. Enschede, 140 p.
- Murray, J.W., 1991, *Ecology and palaeoecology of benthic Foraminifera*. Longman Scientific & Technical, Harlow, U.K., 397 p.
- Parker, F.L., 1958, Eastern Mediterranean Foraminifera, Sediment Cores from the Mediterranean Sea and the Red Sea: *Rept. Swed. Deep Sea Exped.* 1947 1948, 8, no.4, 219-283.
- Röttger, R., 1995, Benthische Foraminiferen (Nord und Ostsee) in Roettger, R. (eds), *Praktikum der Protozoologie*, 217 p., p. 100-110.
- Tendal, O.S., 1996, Synoptic checklist and bibliography of the Xenophyophorea (Protista), with a zoogeographical survey of the group, *Gathea Rep.*, v. 17, p. 79-101.
- Thalmann, H.E., 1960, *An index to the genera and species of the Foraminifera 1890-1950*, George Vanderbilt Foundation, Stanford University, California, USA.
- Timm, S., 1992, Rezente Tiefsee Benthosforaminiferen aus Oberflächensedimenten des Golfs von Guinea (Westafrika) Taxonomie, Verbreitung, Ökologie und Korngrößenfraktionen: *Ber. Geol. Paläontol. Inst. Univ. Kiel in Geol. Pal. Inst. und Museum Kiel* (eds), 19.
- Wollenburg, J. & A. Mackensen, 1998, Living benthic foraminifers from the central Arctic Ocean: faunal composition, standing stock and diversity, *Mar. Micropal.*, 34, 153-185.
- Wollenburg, J. & A. Mackensen, 1998, On the vertical distribution of living (Rose Bengal stained) benthic foraminifers in the Arctic Ocean, *J. Foram. Res.*, 28, 268-285.
- Wollenburg, J., 1995, *Benthische Foraminiferenfaunen als Wassermassen-, Produktions- und Eisdriftanzeiger im Arktischen Ozean: Berichte zur Polarforschung in AWI*, Bremerhaven (eds), Bremerhaven, v. 179, p. 1-227.
- Southern North Sea**
- Moodley, L., 1992, *Experimental Ecology of Benthic Foraminifera in Soft Sediments and its (Paleo) Environmental Significance*; Diss. Univ. Enschede, 140 p.
- Mediterranean Sea:**
- Cimerman, F. & Langer, M.R., 1991, *Mediterranean Foraminifera*: Slovenska Akademija Znanosti, Ljubljana, 118 p.
- Jorissen, F.J., 1988, Benthic Foraminifera from the Adriatic Sea; principles of phenotypic variation: *Utrecht Micropal. Bull.*, 37.
- Mullineaux, L.S. & Lohmann, G.P., 1981, Late Quaternary stagnations and recirculation of the eastern Mediterranean: Changes in the deep water recorded by fossil benthic Foraminifera: *J. Foram. Res.*, 11, 20-39.
- Parker, F.L., 1958, Eastern Mediterranean Foraminifera, Sediment Cores from the Mediterranean Sea and the Red Sea: *Rept. Swed. Deep Sea Exped.* 1947 1948, 8, no.4, 219-283.
- Wright, R., 1978, Neogene benthic foraminifers from DSDP Leg 42A, Mediterranean Sea; in Hsü, K. & Montadert, L. et al. (eds.), *Initial Repts. Deep Sea Drilling Project (Joides)*, 42, 709-726.
- North Atlantic**
- Barker, R.W., 1960, Taxonomic Notes on the Species Figured by H. B. Brady in his Report on the Foraminifera Dredged by H. M. S. Challenger During the years 1873-1876: *Soc. Econ. Paleontol. Mineral. Spec. Pub.* 9, 238 p.
- Caralp, M.H., 1984, Quaternary calcareous benthic foraminifers, Leg 80: in de Graciansky, P.C. & Poag, C.W. et al. (eds): *Initial Reports of the Deep Sea Drilling Project*, Leg 80, Washington, 725-755.
- Gooday, A.J., 1986, Soft shelled Foraminifera in meiofaunal samples from the bathyal northeast Atlantic: *Sarsia*, 71, 275-287.
- Hermelin, J.O.R. & Scott, D.B., 1985, Recent benthic Foraminifera from the central North Atlantic: *Micropaleontology*, v. 31, p. 199-220.
- Arctic**
- Gudmundsson, G., 1998, Distribution limits of Pyrgo species at the biogeographic boundaries of the Arctic and the North-Atlantic boreal regions, *J. Foram. Res.*, 28, p. 240-256.
- Wollenburg, J. & A. Mackensen, 1998, Living benthic foraminifers from the central Arctic Ocean: faunal

composition, standing stock and diversity, *Mar. Micropal.*, 34, 153-185.
Wollenburg, J., 1995, Benthische Foraminiferenfaunen als Wassermassen, Produktions und Eisdrifianzeiger im Arktischen Ozean: Berichte zur Polarforschung in AWI, Bremerhaven (eds), Bremerhaven, v. 179, p. 1-227

FORAMINIFERA

Order Foraminiferida

Suborder Allogromiina

Family Allogromiidae

<i>Allogromia</i>	
<i>laticollaris</i> Arnold, 1948	A M
<i>mollis</i> Gruber, 1881	M
<i>ovoidea</i> Rhumbler, 1904	A M
= <i>Gromia oviiformis</i> Rhumbler, 1904	
<i>Boderia</i>	
<i>turneri</i> Wright, 1867	A
<i>Cylindrogullmia</i>	
<i>alba</i> Nyholm, 1974	A
<i>Dactylosacus</i>	
<i>vermiformis</i> Rhumbler, 1894	
<i>Dendrotuba</i>	
<i>nodulosa</i> Rhumbler, 1894	
<i>Gloio(gull)mia</i>	
<i>eurystoma</i> Nyholm, 1974	
<i>Hospitella</i>	
<i>fulva</i> Rhumbler, 1911	
<i>Marsepulina</i>	
<i>schultzei</i> Rhumbler, 1904	A M
<i>Micatuba</i>	
<i>flexilis</i> (Höglund, 1947)	
<i>Micrometula</i>	
<i>hyalostriata</i> Nyholm, 1952	A
<i>Nemogullmia</i>	
<i>longevaribilis</i> Nyholm, 1956	A
<i>Nodellum</i>	
<i>membranaceum</i> (Rhumbler, 1913) Brady, 1913	A M
= <i>Reophax membranaceum</i> Brady, 1913	
<i>Peritygma</i>	
<i>lunothalamia</i> Arnold, 1978	A M
<i>Phainogullmia</i>	
<i>aurata</i> Nyholm, 1955	A
<i>Placopsisilina</i>	
<i>bradyi</i> Cushman & McCulloch, 1939	M
<i>Placopsilinella</i>	
<i>aurantiaca</i> Earland, 1910	A
<i>Rhynchogromia</i>	
<i>variabilis</i> Rhumbler, 1894	A
<i>Sheppardella</i>	
<i>taeniformis</i> Siddall, 1880	A M
<i>Sphairogullmia</i>	
<i>aurea</i> Nyholm, 1974	A
<i>Thalamophaga</i>	
<i>incerta</i> (Rhumbler, 1911)	A
<i>ramosa</i> Rhumbler, 1911	A
<i>Tinogullmia</i>	
<i>hyalina</i> Nyholm, 1954	A
<i>riemannii</i> Gooday, 1990	A

Family Lagynidae

<i>Lagynis</i>	
<i>baltica</i> Schultze, 1854	A
<i>parva</i> Schultze, 1854	A
<i>Myxotheca</i>	
<i>arenilega</i> Schaudin, 1893	A M
<i>Ophiotuba</i>	
<i>gelatinosa</i> Rhumbler, 1894	A
<i>Pilalla</i>	
<i>exigua</i> Rhumbler, 1938	A
<i>Rhumblierinella</i>	

bacillifera Schmidt, 1929

A

Suborder Textulariina

Family Ammodiscidae

<i>Ammodiscus</i>	
<i>anguillae</i> Höglund, 1947	A
<i>catinus</i> Hoeglund, 1947	M
<i>flavidus</i> Hoeglund, 1947	A
<i>gullmarenensis</i> Höglund, 1948	A
<i>incertus</i> (d'Orbigny, 1839)	A M
<i>intermedius</i> Höglund, 1947	A
<i>minimus</i> Hoeglund, 1947	A
<i>planorbis</i> Höglund, 1947	A M
<i>planus</i> Höglund, 1947	A M
<i>rhumbleri</i> Rottgardt, 1952	A
<i>slevicus</i> Rottgardt, 1952	A
<i>tenuis</i> Brady, 1881	A

Ammolagena

<i>clavata</i> (Jones & Parker, 1860)	A M
---------------------------------------	-----

Glomospira

<i>gordialis</i> (Jones & Parker, 1860)	A M
= <i>Trochammina squamata</i> Jones & Parker, 1860 var.	

charoides Jones & Parker, 1860

Repmanina

<i>charoides</i> (Jones & Parker, 1860)	A M
= <i>Glomospira charoides</i> (Jones & Parker, 1860)	

Spirillinoides

<i>circumcinctus</i> Rhumbler, 1938	A
-------------------------------------	---

Tolyppammina

<i>schaudinni</i> Rhumbler, 1904	A
<i>vagans</i> (Brady, 1879)	A

Turritellella

<i>shoneana</i> (Siddall, 1878)	A
<i>spectabilis</i> (Brady, 1881)	A

Family Ammosphaeroidinidae

<i>Adercotryma</i>	
<i>glomerata</i> (Brady, 1878)	A M
= <i>Glomospira glomerata</i> Höglund, 1947	
= <i>Haplophragmoides glomeratum</i> (Brady, 1878)	
= <i>Lituolea glomerata</i> d'Orbigny, 1826 (Brady, 1878)	
<i>wrighti</i> Brönnimann & Whittaker, 1987	

Family Astrorhizidae

<i>Armoredella</i>	
<i>sphaerica</i> Heron-Allen & Earland, 1932	A M
<i>Astrorhiza</i>	
<i>angulosa</i> Brady, 1881	A
<i>arenaria</i> Norman, 1876	A
<i>granulosa</i> (Brady, 1879) Parker, 1960	A
<i>limicola</i> Sandahl, 1857	A
<i>Bathysiphon</i>	
<i>acuta</i> (Höglund, 1947)	A
= <i>Hippocrepinella acuta</i> (Höglund, 1947)	
<i>alba</i> (Heron-Allen & Earland, 1932)	A
= <i>Hippocrepinella alba</i> Heron-Allen & Earland, 1932	
= <i>Hippocrepinella ampulacea</i> Rhumbler, 1935	
<i>ampulacea</i> (Rhumbler, 1935)	A
<i>argentea</i> Heron-Allen & Earland, 1913	A
<i>capillare</i> de Folin, 1886	A
<i>crassatina</i> (Brady, 1884)	A
= <i>Astrorhiza crassatina</i> Brady, 1884	
<i>filiformis</i> M. Sars, 1871	A
<i>flexibilis</i> (Wiesner, 1931)	A
= <i>Hippocrepinella flexibilis</i> (Wiesner, 1931)	
<i>flexilis</i> Höglund, 1947	A
<i>hirudinea</i> (Heron-Allen & Earland, 1932)	A
= <i>Hippocrepinella hirudinea</i> Heron-Allen & Earland, 1932	
<i>historix</i> Gooday, 1988	A
<i>minutus</i> Pearcey, 1900	A
<i>rufus</i> De Folin, 1886	A
<i>virgulata</i> (Rhumbler, 1935)	A
= <i>Hippocrepinella virgulata</i> Rhumbler, 1935	
<i>Bigenerina</i>	
<i>cylindrica</i> Cushman, 1922	M
= <i>B. digitata</i> (d'Orbigny, 1826)	

<i>laevigata</i> d'Orbigny, 1826	M	<i>pseudospiralis</i> (Williamson, 1858)	A M
<i>paupa</i> d'Orbigny, 1826		= <i>Proteonina pseudospiralis</i> Williamson, 1858	
<i>nodosaria</i> d'Orbigny, 1826	M	<i>runiana</i> (Heron-Allen & Earland, 1916)	A M
<i>Botellina</i>		= <i>Haplophragmium runianum</i> Heron-Allen & Earland,	
<i>labyrinthica</i> Brady, 1881	A	1916	
<i>Dendrophyra</i>	A	<i>tenuimargo</i> (Brady, 1882)	A M
<i>arborescens</i> (Norman, 1876)		Family Eggerellidae	
= <i>Psammatodendron arborescens</i> (Norman, 1881)	A	<i>Eggerella</i>	
<i>erecta</i> Wright, 1861	A	<i>affixa</i> (Cushman)	A
<i>tolli</i> Awerinzew, 1911	A	<i>europeum</i> (Christensen, 1958)	
<i>Halophysema</i>	A	<i>parkerae</i> Uchio, 1960	A
<i>ramulosa</i> Bowerbank, 1862	A	= <i>E. bradyi</i> (Cushman, 1911)	
<i>tumanowiczii</i> Bowerbank, 1862	A	<i>Eggerelloides</i>	
<i>Hippocrepina</i>	A	<i>advenus</i> (Cushman, 1922)	A M
<i>cylindrica</i> Höglund, 1947	A	<i>medius</i> (Höglund, 1947)	A
<i>indivisa</i> Parker, 1870	A M	= <i>Verneuilina media</i> Höglund, 1947	
<i>pusilla</i> Heron-Allen & Earland, 1930	A	<i>propinqua</i> (Brady, 1884)	A
<i>Jaculella</i>	A	<i>scaber</i> (Williamson, 1858)	A
<i>acuta</i> Brady, 1879	A	= <i>Bulimina scabra</i> Williamson, 1858	
<i>obtusa</i> Brady, 1882	A	= <i>Eggerella scabra</i> (Williamson, 1858)	
<i>Loeblichopsis</i>	A	= <i>E. scabrus</i> (Williamson, 1858)	
<i>cylindrica</i> (Brady, 1884)	A	= <i>Verneuilina polystropha</i> (Reuss, 1856)	
= <i>Reophax cylindrica</i> Brady, 1884		<i>Karreriella</i>	
<i>Marsipella</i>	A	<i>apicularis</i> (Cushman, 1911)	A
<i>arenaria</i> Christiansen, 1958	A	<i>bradyi</i> (Cushman, 1911)	A M
<i>dextrospiralis</i> Chapman & Parr	A	<i>novangliae</i> (Cushman, 1922)	A
<i>elongata</i> Norman, 1878	A	Family Haplophragmoididae	
<i>spiralis</i> Heron-Allen & Earland, 1912	A	<i>Ammotium</i>	
<i>Nubeculariella</i>	A	<i>fragile</i> Warren, 1957	A
<i>birulai</i> Awerinzew, 1911	A	= <i>Haplophragmoides fragile</i> Warren, 1957	
<i>Pelosina</i>	A	<i>Conglophragmium</i>	
<i>arborescens</i> Pearcey, 1914	A	<i>coronatum</i> (Brady, 1879)	A M
<i>arctica</i> Awerinzew, 1911	A	= <i>Haplophragmoides coronatum</i> (Brady, 1879)	
<i>cylindrica</i> Brady, 1884	A	<i>Haplophragmoides</i>	
<i>didera</i> (Loeblich & Tappan, 1953)	A	<i>canariensis</i> (d'Orbigny, 1839)	A M
<i>fusciformis</i> Earland, 1933	A	<i>membranaceum</i> Höglund, 1947	A
<i>rotundata</i> Brady, 1879	A	<i>sphaeriloculum</i> Cushman, 1910	A
<i>variabilis</i> Brady, 1879	A	<i>wilberti</i> Andersen, 1953	A
<i>Pelosphaera</i>	A	<i>Labrospira</i>	
<i>cornuta</i> Heron-Allen & Earland, 1932	A	<i>rotulata</i> (Brady, 1881)	A
<i>Rhabdaminella</i>	A	<i>Veleroninoides</i>	
<i>cylindrica</i> (Brady, 1882)	A	<i>wiesneri</i> (Parr, 1950)	A M
= <i>Marsipella cylindrica</i> Brady, 1882		= <i>Alveophragmium wiesneri</i> (Parr, 1950)	
<i>Rhabdammina</i>	A	= <i>Cribrostomoides wiesneri</i> (Parr, 1950)	
<i>abyssorum</i> M. Sars, 1868	A	= <i>Labrospira wiesneri</i> (Parr, 1950)	
= <i>Bathysiphon discreta</i> (Brady, 1884)		Family Hippocrepinidae	
<i>cornuta</i> (Brady, 1879)	A	<i>Hyperammina</i>	
<i>discreta</i> Brady, 1884	A	<i>cylindrica</i> Parr, 1950	A
<i>inaequalis</i> Le Calvez, 1935	A	<i>elongata</i> Brady, 1884	A
<i>linearis</i> Brady, 1879	A	<i>fragilis</i> Höglund, 1947	A
<i>major</i> (DeFolin, 1986)	A	<i>fribilis</i> Brady, 1884	A M
<i>neglecta</i> Gooday, 1986	A	<i>laevigata</i> Wright, 1891	A M
<i>scabra</i> Höglund, 1947	A	<i>subnodosa</i> Brady, 1884	A
<i>Rhizammina</i>	A	<i>Saccorhiza</i>	
<i>algaeformis</i> Brady, 1879	A	<i>ramosa</i> (Brady, 1879)	A M
<i>algaeformis</i> Brady, 1879 f. <i>nuda</i> Rhumbler, 1935	A	= <i>Hyperammina ramosa</i> Brady, 1879	
<i>Testulosphon</i>	A M	Family Hormosinidae	
<i>indivisus</i> (Brady, 1884)		<i>Cuneata</i>	
= <i>Rhizammina indivisa</i> (Brady, 1884)		<i>arctica</i> (Brady, 1881)	A
<i>Vanhoefenella</i>	A	= <i>Reophax arctica</i> (Brady, 1881)	
<i>gaussi</i> Rhumbler, 1905	A	<i>Hormosina</i>	
<i>gaussi</i> Rhumbler, 1905 f. <i>imperfecta</i> Rhumbler, 1935	A	<i>globulifera</i> Brady, 1879	A
Family Ataxophragmiidae		<i>normani</i> Brady, 1881	A
<i>Martinottiella</i>	M	<i>semiglobosa</i> Wieser	M
<i>communis</i> (d'Orbigny, 1846)	M	<i>bacillaris</i> (Brady, 1881)	A M
<i>gymnesica</i> (Colom, 1964)	M	= <i>Martinottiella bacillaris</i> (Brady, 1881)	
= <i>Listerella gymnesica</i> Colom, 1984		<i>Hormosinella</i>	
Family Baculellidae		<i>carpenteri</i> (Brady, 1881)	A
<i>Edgertonia</i>	A	= <i>Hormosina carpenteri</i> Brady, 1881	
<i>floccula</i> Tendal & Hessler, 1977		<i>distans</i> (Brady, 1881)	A
<i>tolerans</i> Tendal & Hessler, 1977		= <i>Hormosina distans</i> (Brady, 1881)	
Family Discamminidae		= <i>Reophax distans</i> Brady, 1881	
<i>Ammoscalaria</i>	A	<i>guttifera</i> (Brady, 1881)	
<i>foliacea</i> (Brady, 1881)	A	= <i>Reophax guttifer</i> Brady, 1881	A

<i>ovicula</i> (Brady, 1879)	A M	<i>triangularis</i> Saidova, 1961	A
= <i>Hormosina ovicula</i> Brady, 1879		= <i>Labrospira triangularis</i> (Saidova, 1961)	
<i>Leptohyalis</i>		Cyclammina	
<i>scottii</i> Chaster, 1892	A M	<i>cancellata</i> Brady, 1879	A
= <i>Reophax scottii</i> Chaster, 1892		<i>orbicularis</i> Brady, 1884	A
<i>Protoschista</i>		<i>pusilla</i> Brady, 1881	A
<i>findens</i> (Parker, 1870)	A	<i>senegalensis</i> Colom, 1956	M
<i>Reophax</i>		<i>trullissata</i> (Brady, 1879)	A
<i>agglutinans</i> Cushman, 1913	A	Discammina	
<i>bilocularis</i> Flint, 1897	A M	<i>compressa</i> (Goes, 1882)	M
<i>catella</i> Höglund, 1947	A	Recurvooides	
<i>catenata</i> Höglund, 1947	A	<i>contortus</i> Earland, 1934	A
<i>curtus</i> Cushman, 1920	A	<i>trochamminiforme</i> Höglund, 1947	A
<i>decorata</i> (Earland, 1933)	A	= <i>Cribrostomoides trochamminiforme</i> (Höglund, 1947)	
<i>dentaliniformis</i> Brady, 1881	A M	<i>turbinatus</i> (Brady, 1881)	A
= <i>Nodulina dentaliniformis</i> (Brady, 1884)		= <i>Haplophragmium turbinatum</i> Brady, 1881	
<i>dentaliniformis</i> subsp. <i>fragilis</i> Brady, 1884 n. subsp.	A	Family Lituotubidae	
<i>fusiformis</i> (Williamson, 1858)	A M	<i>Lituotuba</i>	
= <i>Lagenammina fusiformis</i> (Williamson, 1858)		<i>lituiformis</i> (Brady, 1879)	A M
= <i>Proteonina fusiformis</i> (Williamson, 1858)		Family Nouriidae	
= <i>Saccamina fusiformis</i> (Brady, 1879)		<i>Nouria</i>	
<i>gracilis</i> Kiär, ?	A	<i>johnsoni</i> Cushman, 1935	A
<i>helena</i> (Rhumbler, 1913)	A	<i>polymorphinoides</i> Heron-Allen & Earland, 1914	A M
<i>longicollis</i> Wiesner, 1931	A	Family Pseudogaudryinidae	
<i>micaceus</i> Earland, 1934	A M	<i>Pseudoclavulina</i>	
<i>nana</i> Rhumbler, 1913	A M	<i>crustata</i> Cushman, 1936	A M
<i>nodulosus</i> Brady, 1879	A M	Family Rzehakinidae	
<i>pilulifer</i> Brady, 1884	A M	<i>Miliammina</i>	
<i>regularis</i> Höglund, 1947	A	<i>arenacea</i> (Chapman, 1916)	A
<i>rostrata</i> Höglund, 1947	A	<i>fusca</i> (Brady, 1870)	A M
<i>scorpiurus</i> Montfort, 1808	A M	= <i>Quinqueloculina fusca</i> Brady, 1870	
<i>spiculifer</i> Brady, 1879	A	<i>fusca</i> subsp. <i>subterranea</i> Rhumbler, 1936	A
<i>subfusiformis</i> Earland, 1933	A M	<i>groenlandica</i> Cushman, 1933	M
<i>calcareus</i> (Cushman, 1947)	A	<i>rugosa</i> Mikhalevich, 1968	M
<i>Scherochorella</i>	A	Family Saccaminidae	
<i>moniliforme</i> (Siddall, 1886)		<i>Ammopempix</i>	
= <i>Reophax moniliformis</i> (Siddall, 1886)		<i>arctica</i> (Cushman, 1933)	A
<i>Subreophax</i>	A	<i>Amphifенestrella</i>	
<i>aduncus</i> (Brady, 1882)		<i>wiesneri</i> Rhumbler, 1935	A
= <i>Reophax aduncus</i> Brady, 1882		<i>Causia</i>	
<i>incertae sedis</i>		<i>injudicata</i> Rhumbler, 1938	A
<i>Ammotrochoides</i>		<i>Crithiona</i>	
<i>bignoti</i> Janin, 1984	A	<i>abyssorum</i> Kiaer, 1899	A
<i>Jullienella</i>	A	<i>granum</i> Goes, 1894	A
<i>foetida</i> Schlumberger, 1890	A	<i>Crithionina</i>	
<i>Schizammina</i>	A	<i>heincke</i> Rhumbler, 1928	A
<i>furcata</i> Heron-Allen & Earland, 1929	A	<i>hispida</i> Flint, 1899	A
Family Lituolidae		<i>mamilla</i> Goes, 1894	A
<i>Ammobaculites</i>		<i>pisum</i> Goes, 1896	A
<i>agglutinans</i> (d'Orbigny, 1846)	A M	<i>sphaerica</i> (Höglund, 1947)	A
<i>americanus</i> d'Orbigny, 1826 Cushman, 1910	A	= <i>Thorammina sphaerica</i> (Höglund, 1947)	
<i>filiformis</i> Earland, 1934	A	<i>Hemisphaeramma</i>	
<i>ponticus</i> Mikhalevich, 1968	M	<i>bradyi</i> Loeblich & Tappan, 1957	A M
<i>Ammotium</i>		= <i>Webbinella hemisphaerica</i> (Jones & Parker & Brady, 1866)	
<i>cassis</i> (Parker, 1870)	A	<i>marisalbi</i> (Schedrina, 1962)	A
= <i>Lituola cassis</i> Dawson, 1870		<i>Lagenammina</i>	
<i>salsum</i> (Cushman & Brönnimann, 1948)		<i>arenulata</i> (Skinner, 1961)	A
= <i>Ammobaculites salsum</i> Cushman & Brönnimann, 1948		<i>atlantica</i> (Cushman, 1944)	A M
<i>Cribrostomoides</i>		= <i>Proteonina atlantica</i> Cushman, 1944	
<i>crassimargo</i> (Norman, 1892)	A	= <i>Saccamina atlantica</i> Cushman, 1944	
= <i>Haplophragmium crassimargo</i> Norman, 1892		<i>diffugiformis</i> (Brady, 1879)	A M
= <i>Labrospira crassimargo</i> (Norman, 1892)		= <i>Proteonina diffugiformis</i> (Brady, 1879)	
<i>jeffreysi</i> (Williamson, 1858)	M	= <i>Reophax diffugiformis</i> Brady, 1879	
= <i>Labrospira jeffreysi</i> (Williamson, 1858)		<i>hancocki</i> (Cushman & Mc Culloch, 1948)	A
= <i>Nonionina jeffreysi</i> (Williamson, 1858)		<i>laguncula</i> Rhumbler, 1911	A
<i>kosterensis</i> (Höglund, 1947)	A M	<i>tubulata</i> Rhumbler, 1931	A M
= <i>Labrospira kosterensis</i> Höglund, 1947	A	= <i>Reophax tubulatum</i> (Rhumbler, 1931)	
<i>ringens</i> (Brady, 1879)	A	<i>Pilulina</i>	
= <i>Labrospira ringens</i> (Brady, 1879)		<i>argentea</i> Höglund, 1947	A
<i>scitulum</i> (Brady, 1881)	A	<i>jeffreysi</i> Carpenter, 1875	A
= <i>Recurvooides laevigatum</i> Höglund, 1947	A M	<i>Psammosphaera</i>	
<i>subglobosus</i> (Cushman, 1910)		<i>bowmanni</i> Heron-Allen & Earland, 1912	A
= <i>Labrospira subglobosa</i> (Cushman, 1910)		<i>exigua</i> (Rhumbler, 1935)	A
= <i>Lituola subglobosa</i> (Cushman, 1910)			
<i>subturbanata</i> (Cushman, 1920)	A		

<i>frankei</i> (Rhumbler, 1935)	A	<i>Globotextularia</i>	
<i>fusca</i> Schultze, 1875	A M	<i>anceps</i> (Brady, 1884)	M
<i>fusca</i> subsp. <i>adhaerens</i> Rhumbler, 1935	A	<i>Sahulina</i>	
<i>fusca</i> subsp. <i>asperrima</i> Rhumbler, 1935	A	<i>conica</i> (d'Orbigny, 1839)	M
<i>fusca</i> subsp. <i>tapetifera</i> Rhumbler, 1935	A	= <i>Siphonotextularia conica</i> (d'Orbigny, 1839)	
<i>parva</i> Flint, 1899	A	= <i>Textularia conica</i> (d'Orbigny, 1839)	
<i>testacea</i> Flint, 1899	A	<i>Siphonotextularia</i>	
<i>Pseudowebbinella</i>		<i>affinis</i> (Fornasini)	M
<i>goesi</i> (Högglund, 1947)	A	<i>concava</i> (Karrer, 1868)	A M
= <i>Criithionina goesi</i> (Högglund, 1947)		<i>curta</i> (Cushman, 1922)	A
= <i>Liebusella goesi</i> Högglund, 1947		<i>eastatlantica</i> Cushman, 1922	A
<i>Radicula</i>		<i>flintii</i> (Cushman, 1911)	A
<i>limosa</i> Christiansen, 1958	A	= <i>Gaudryina flintii</i> (Cushman, 1911)	
<i>Saccammina</i>		<i>heterostoma</i> (Forsanini, 1896)	A
<i>sphaerica</i> M. Sars, 1868	A M	<i>occidentalis</i> subsp. <i>bermudezi</i> Mikhalevich, 1978	A
<i>sphaerica</i> M. Sars, 1868 var. <i>catenulata</i> Cushman, 1917	A	<i>rolshauseni</i> Phleger & Parker, 1951	A
<i>Saccodendron</i>		<i>Spirorutilus</i>	
<i>heronalleni</i> Rhumbler, 1935	A	<i>carinatus</i> (d'Orbigny, 1846) Jones, 1994	M
<i>heronalleni</i> Rhumbler, 1935 var. <i>laticerium</i> Rhumbler, 1935	A	= <i>Textularia carinata</i> d'Orbigny, 1839	
<i>Sorosphaera</i>		<i>Textularia</i>	
<i>confusa</i> Brady, 1879	A	<i>aciculata</i> d'Orbigny, 1826	M
<i>perforata</i> Stschedrina, 1939	A	<i>agglutinans</i> d'Orbigny, 1839	M
<i>Storthosphaera</i>		<i>australis</i> Parr, 1950	A
<i>albida</i> Schulze, 1875	A	<i>bigeneroides</i> Lacroix, 1932	A M
<i>Technitella</i>		<i>bocki</i> Högglund, 1947	M
<i>bradyi</i> Earland, 1934	M	<i>calva</i> Lalicker, 1940	M
<i>erinacea</i> Hofker, 1972	A	<i>cochleata</i> Lacroix, 1932	A
<i>legumen</i> Norman, 1878	A M	<i>earlandi</i> (Parker, 1952)	A
<i>melo</i> Norman, 1878	A	= <i>Prolixoplecta earlandi</i> (Parker, 1952)	
<i>teivyense</i> Haynes, 1973	A	= <i>Spiroplectammina earlandi</i> (Parker, 1952)	
<i>Telammina</i>		<i>foeda</i> Reuss	M
<i>minuta</i> Gooday & Haynes, 1983		<i>gibbosa</i> d'Orbigny	M
<i>Tholosina</i>		<i>gracillima</i> Högglund, 1947	A
<i>bulla</i> (Brady, 1881)	A M	<i>kattegatensis</i> (Högglund, 1947)	
= <i>Placopsisilina bulla</i> (Brady, 1881)		<i>mexicana</i> Cushman, 1922	A
<i>bulla</i> subsp. <i>appressa</i> Rhumbler, 1935	A	<i>millettii</i> Cushman, 1911	A
<i>centroforata</i> Rhumbler, 1935	A	<i>panamensis</i> Cushman, 1918	A
<i>confusa</i> Cushman, 1920	A	<i>porrecta</i> (Brady, 1884)	M
= <i>Placopsisilina confusa</i> Cushman, 1920		<i>pseudogrammen</i> Chapman & Parr, 1937	A M
<i>laevis</i> Rhumbler, 1931	A	= <i>T. gramen</i> d'Orbigny	
<i>margineforata</i> Rhumbler, 1935	A	<i>pseudorugosa</i> Lacroix, 1932	M
<i>protea</i> Heron-Allen & Earland, 1932	A	<i>skagerakensis</i> Högglund, 1947	A
<i>vesicularis</i> (Brady, 1879)	A	<i>tenuissima</i> Earland, 1933	A
<i>Thurammina</i>		<i>torquata</i> Parker, 1952	A
<i>albicany</i> Brady, 1879	A	<i>trochus</i> d'Orbigny, 1839	M
<i>basispicula</i> Zheng	A	<i>truncata</i> Högglund, 1947	A M
<i>compressa</i> Brady, 1879	A	<i>Vulvulina</i>	
<i>faerleensis</i> Högglund, 1948	A	<i>pennatula</i> (Batsch, 1791)	A M
<i>papillata</i> Brady, 1879	A M	= <i>Nautilus pennatula</i> Batsch, 1791	
<i>Tumidotubus</i>		= <i>Orthocera pennatula</i> Batsch, 1791	
<i>albus</i> Gooday & Haynes, 1983		= <i>V. capreolus</i> (d'Orbigny, 1839)	
<i>Webbinella</i>		Family Trochamminidae	
<i>arctica</i> Cushman, 1848	A	<i>Atlantiella</i>	
<i>earlandi</i> Rhumbler, 1935	A	<i>atlantica</i> (Parker, 1952)	A
Family Spiroplectamminidae		= <i>Trochammina atlantica</i> Parker, 1952	
<i>Morulaeplecta</i>		<i>Deuterammina</i>	
<i>bulbosa</i> (Högglund, 1947)	A	<i>discorbis</i> Earland, 1934	
= <i>Textularia bulbosa</i> (Högglund, 1947)		= <i>Trochammina discorbis</i> Earland, 1934	A
<i>Spiroplectammina</i>		<i>grahami</i> Brönniman & Whittaker, 1988	A
<i>biformis</i> (Parker & Jones, 1865)	A	<i>grisea</i> Earland, 1934	A
= <i>Prolixoplecta biformis</i> (Parker & Jones, 1865)		= <i>Trochammina grisea</i> Earland, 1934	
= <i>Textularia agglutinans</i> Parker & Jones, 1865 var.		<i>montagui</i> Heron-Allen & Earland, 1911	
<i>biformis</i> Parker & Jones, 1865		= <i>Trochammina montagui</i> Brönnimann & Whittaker, 1988	
<i>sagittula</i> (Defrance, 1824)	A	<i>rotaliformis</i> (Heron-Allen & Earland, 1911)	A M
= <i>Prolixoplecta sagittula</i> (Defrance, 1824)		= <i>Trochammina rotaliformis</i> Heron-Allen & Earland, 1911	
= <i>Spiroplectinella sagittula</i> (d'Orbigny, 1839)		<i>Earlandammina</i>	
= <i>Textularia sagittula</i> Defrance, 1824		<i>inconspicua</i> Earland, 1934	A
<i>Spiroplectinella</i>		= <i>Trochammina inconspicua</i> Earland, 1934	
<i>wrightii</i> (Silvestri, 1903)	M	<i>Jadammina</i>	
= <i>Prolixoplecta wrightii</i> (Silvestri, 1903)		<i>macrescens</i> (Brady, 1870)	A M
= <i>Spiroplectammina wrightii</i> (Silvestri, 1903)		= <i>Trochammina macrescens</i> (Brady, 1870)	
Family Textulariidae		<i>Lepidodeuterammina</i>	
		<i>ochracea</i> (Williamson, 1858)	A

= <i>Rotalina ochracea</i> Williamson, 1858			
= <i>Trochammina ochracea</i> (Williamson, 1858)			
<i>Polystomammina</i>			
<i>nitida</i> (Brady, 1881)	A M		
= <i>Cribrostomoides nitida</i> (Goes, 1896)			M
= <i>Labrospira nitida</i> Goes, 1896			
= <i>Trochammina nitida</i> (Brady, 1881)			
<i>Portatrocchammina</i>			
<i>karica</i> (Shchedrina, 1946)	A		
= <i>P. bipolaris</i> Brönnimann & Whittaker, 1980			
= <i>Trochammina karica</i> Shchedrina, 1946			
<i>murrayi</i> Brönnimann & Zaninetti, 1984			
<i>Recurvoidella</i>			
<i>bradyi</i> (Robertson, 1891)	A		
= <i>Cribrostomoides bradyi</i> (Robertson, 1891)			
= <i>Haplophragmoides bradyi</i> (Robertson, 1891)			
= <i>Labrospira bradyi</i> (Robertson, 1891)			
= <i>Trochammina bradyi</i> (Robertson, 1891)			
<i>Tiphochrocha</i>			
<i>comprimata</i> (Cushman & Brönnimann, 1948)	A		
= <i>Trochammina comprimata</i> Cushman & Brönnimann, 1948			
<i>Tritaxis</i>			
<i>britannica</i> Brönnimann & Whittaker, 1990			
<i>challengeri</i> (Hedley, Hurdle & Burdett, 1964) Jones, 1994	A M		
<i>conica</i> (Parker & Jones, 1865)	A		
= <i>Valvulina conica</i> Parker & Jones, 1865			
<i>fusca</i> (Williamson, 1858)	A		
= <i>Valvulina fusca</i> (Williamson, 1858)			
<i>squamata</i> (Jones & Parker, 1860)	M		
= <i>Trochammina squamata</i> (Jones & Parker, 1860)			
<i>Trochammina</i>			
<i>adoptera</i> Rhumbler	A		
<i>astrifica</i> Rhumbler, 1938	A		
<i>bellingshausenii</i> (Brönnimann & Whittaker, 1988)	A		
<i>globigeriniformis</i> Parker & Jones, 1865	M		
= <i>Ammoglobigerina globigeriniformis</i> (Parker & Jones, 1865)			
= <i>Lituola nautiloides</i> Lamarck, 1804 var. <i>globigeriniformis</i> Parker & Jones, 1865			
<i>helgolandica</i> Rhumbler, 1938	A		
<i>inflata</i> (Montagu, 1808)	A M		
= <i>Nautilus inflatus</i> Montagu, 1808			
<i>intermedia</i> Rhumbler, 1938	A		
<i>japonica</i> Ishiwada, 1950	A		
<i>multiloculata</i> Höglund, 1947	A		
<i>shannoni</i> Brönnimann & Whittaker, 1988	A		
<i>Trochamminopsis</i>			
<i>pusilla</i> (Höglund, 1947)	A		
= <i>Haplophragmoides pusillum</i> (Höglund, 1947)			
= <i>Trochammina pusilla</i> (Höglund, 1947)			
= <i>T. quadriloba</i> (Höglund, 1947)			
Family Verneuilinidae			
<i>Dorothia</i>			
<i>bradyana</i> Cushman, 1936	A		
= <i>Martinottiella bradyana</i> Cushman, 1936			
<i>pseudoturris</i> (Cushman, 1922)	A		
= <i>Textularia pseudoturris</i> (Cushman, 1922)			
<i>scabra</i> (Brady, 1884)	A M		
<i>Gaudryina</i>			
<i>rudis</i> Wright, 1900	A		
= <i>Connemarella rudis</i> (Wright, 1900)			
= <i>Dorothia rudis</i> (Wright, 1900)			
<i>Verneuilina</i>			
<i>arctica</i> Höglund, 1947	A		
= <i>Cribrostomoides arctica</i> (Höglund, 1947)			
= <i>Eggerella arctica</i> Höglund, 1947			
= <i>Eggerelloides arctica</i> (Höglund, 1947)			
<i>Verneuilinella</i>			
<i>advena</i> (Cushman, 1922)	A		
= <i>Eggerelloides advena</i> Cushman, 1922			
= <i>Trochammina advena</i> Cushman, 1922			
= <i>Verneuilina advena</i> Cushman, 1922			
Suborder Miliolina			
		Family Cornuspiridae	
		<i>Cornuspira</i>	
		<i>carinata</i> (Costa, 1856)	A M
		<i>foliacea</i> (Philippi, 1844)	M
		= <i>Cornuspirodes foliaceus</i> (Philippi, 1844)	
		<i>involvens</i> (Reuss, 1850)	M
		<i>planorbis</i> Schultze, 1854	A
		<i>Cornuspirella</i>	
		<i>diffusa</i> (Heron-Allen & Earland, 1913)	M
		= <i>Cornuspirodes diffusa</i> Heron-Allen & Earland, 1916	
		<i>Cornuspiroides</i>	
		<i>striolatus</i> (Brady, 1882)	M
		Family Fischerinidae	
		<i>Trisegmentina</i>	
		<i>compressa</i> Wiesner, 1931	M
		Family Hauerinidae	
		<i>Affinetrina</i>	
		<i>planciana</i> (d'Orbigny, 1839)	M
		= <i>Lachlanella planciana</i> (d'Orbigny, 1839)	
		= <i>Quinqueloculina planciana</i> (d'Orbigny, 1839)	
		= <i>Triloculina planciana</i> d'Orbigny, 1839	
		<i>Articularia</i>	
		<i>sagra</i> (d'Orbigny, 1839)	M
		= <i>Articulina sagra</i> (d'Orbigny, 1839)	
		<i>Articulina</i>	
		<i>carinata</i> Wiesner, 1923	M
		<i>pacifica</i> Cushman, 1944	M
		<i>tubulosa</i> (Seguenza, 1862)	M
		= <i>Quinqueloculina tubulosa</i> Seguenza, 1862	
		<i>Biloculinella</i>	
		<i>cryptella</i> d'Orbigny, 1839	M
		= <i>Triloculina cryptella</i> d'Orbigny, 1839	
		<i>cylindrica</i> Todd, 1948	M
		<i>depressa</i> (d'Orbigny, 1826)	M
		= <i>Biloculina depressa</i> (d'Orbigny, 1826)	
		= <i>Pyrgo depressa</i> (d'Orbigny, 1826)	
		= <i>Pyrgo rotalaria</i> Loeblich & Tappan, 1953	
		<i>globula</i> (Bornemann, 1855)	M
		<i>inflata</i> (Wright, 1902)	A M
		<i>labiata</i> (Schlumberger, 1891)	M
		<i>subsphaerica</i> (Wiesner, 1923)	M
		<i>wiesneri</i> (Le Calvez, J. & Y., 1958)	M
		<i>Cruciloculina</i>	
		<i>triangularis</i> d'Orbigny, 1826	M
		<i>Cycloforina</i>	
		<i>colomi</i> (Le Calvez, J. & Y., 1958)	M
		<i>contorta</i> (d'Orbigny, 1846)	M
		= <i>Quinqueloculina contorta</i> d'Orbigny, 1839	
		<i>juleana</i> (d'Orbigny, 1846)	M
		<i>rugosa</i> (d'Orbigny, 1826)	M
		<i>tenuicollis</i> (Wiesner, 1923)	M
		<i>villafanca</i> (Le Calvez, J. & Y., 1958)	M
		<i>Lachlanella</i>	
		<i>undulata</i> (d'Orbigny, 1826)	M
		= <i>Quinqueloculina undulata</i> d'Orbigny, 1852	
		<i>variolata</i> (d'Orbigny, 1826)	M
		<i>Sigmoihauerina</i>	
		<i>bradyi</i> (Cushman, 1917)	M
		= <i>Hauerina compressa</i> d'Orbigny, 1839	
		<i>Wellmanellina</i>	
		<i>striata</i> (Sidebottom, 1904)	M
		Family Miliolidae	
		<i>Adelosina</i>	
		<i>bicornis</i> (Walker & Jacob, 1798)	A
		= <i>Massilina bicornis</i> (Walker & Jacob, 1798)	
		= <i>Quinqueloculina bicornis</i> (Walker & Jacob, emend. Haynes)	
		<i>carinata-striata</i> Wiesner, 1923	M
		= <i>Spiroloculina carinata</i> Wiesner, 1923	
		= <i>Spiroloculina nitida</i> d'Orbigny, 1839	
		<i>cliarensis</i> (Heron-Allen & Earland, 1930)	M
		= <i>Quinqueloculina cliarensis</i> Heron-Allen & Earland, 1930	
		<i>dubai</i> (d'Orbigny, 1826)	M

= <i>Triloculina dubia</i> (d'Orbigny, 1826)			
<i>duthiersi</i> Schlumberger, 1886	M		A
<i>elegans</i> (Williamson, 1858)	M		
<i>ferussaci</i> (d'Orbigny, 1826)			
<i>intricata</i> (Terquem, 1878)	M		
= <i>Quinqueloculina intricata</i> (Terquem, 1878)			
<i>mediterranensis</i> (Le Calvez, J. & Y., 1958)	M		
<i>partschii</i> (d'Orbigny, 1846)	M		
<i>pulchella</i> (d'Orbigny, 1826)	M		
= <i>Quinqueloculina pulchella</i> (d'Orbigny, 1826)			
= <i>Spiroloculina pulchella</i> (d'Orbigny, 1826)			
<i>Ammomassilina</i>			
<i>alveoliniformis</i> (Millett, 1891)	A		
<i>Flintina</i>			
<i>crassatina</i> (Brady, 1884)	M		
<i>Flintinoides</i>			
<i>labiosa</i> (d'Orbigny, 1839)	M		
= <i>Miliolinella labiosa</i> (d'Orbigny, 1839)			
<i>Massilina</i>			
<i>amygdalooides</i> (Brady, 1884) Jones, 1994	A		
= <i>Quinqueloculina amygdalooides</i> (Brady, 1884)			
<i>arenaria</i> (Brady)	M		
<i>granulocostata</i> (Germeraad, 1946)	M		
= <i>Triloculina lineiana</i> (Brady, 1884)			
<i>gualteriana</i> (d'Orbigny, 1839)	M		
= <i>Triloculina gualteriana</i> (d'Orbigny, 1839)			
<i>Miliolinella</i>			
<i>chukchiensis</i> Loeblich & Tappan, 1950	A		
<i>dilatata</i> (d'Orbigny, 1839)	M		
<i>elongata</i> Kruit, 1955	M		
<i>grata</i> (Terquem, 1878)	M		
<i>meridionalis</i> (d'Orbigny, 1839)	A		
<i>oblonga</i> (Montagu, 1803)	A M		
<i>pratiense</i> Meyer-Abich	A		
<i>semicostata</i> (Wiesner, 1923)	M		
<i>subrotunda</i> (Montagu, 1803)	A M		
= <i>Quinqueloculina subrotunda</i> Montagnu, 1803 f.			
<i>hauerinoides</i> Rhumbler, 1936			
= <i>Vermiculum subrotundum</i> Montagnu, 1803			
<i>valvularis</i> (Reuss)	M		
<i>webbiana</i> (d'Orbigny, 1839)	M		
<i>Parrina</i>			
<i>bradyi</i> (Millett, 1898)	M		
<i>Planispirinoides</i>			
<i>bucculentus</i> (Brady, 1884)	A		
<i>Pseudomassilina</i>			
<i>australis</i> (Cushman, 1932)	A M		
<i>macilenta</i> (Brady)	M		
<i>Pseudotriloculina</i>			
<i>cuneata</i> (Karrer, 1867)	M		
<i>laevigata</i> (d'Orbigny, 1826)	M		
<i>sidebottomi</i> (Martinotti, 1920)	M		
<i>Ptychomiliola</i>			
<i>seperans</i> (Brady)	M		
<i>Pyrgo</i>			
<i>anomala</i> (Schlumberger, 1891)	M		
<i>appendiculata</i> (Reuss)	A		
<i>bradyi</i> (Lamarck, 1804) Schlumberger, 1891	A M		
<i>bulloides</i> (d'Orbigny)	M		
<i>carinata</i> (d'Orbigny, ??)			
<i>comata</i> (Brady, 1884)	M		
<i>fornasinii</i> Chapman & Parr			
<i>inornata</i> (d'Orbigny, 1839)	M		
= <i>Triloculina inornata</i> d'Orbigny, 1839			
<i>laevis</i> Defrance	A		
<i>lucernula</i> (Schwager, 1891)	A M		
<i>murrhina</i> (Schwager, 1866)	A M		
= <i>Biloculina murrhina</i> Schwager, 1866			
<i>patagonica</i> (d'Orbigny, 1839)	A M		
= <i>Quinqueloculina patagonica</i> (d'Orbigny, 1839)			
<i>pisum</i> (Schlumberger, 1891)	M		
<i>ringens</i> (Lamarck, 1804)	A		
<i>simplex</i> (d'Orbigny, 1839)	M		
<i>subglobulus</i> Parr	A		
<i>subsphaerica</i> (d'Orbigny, 1839)	M		
<i>vespertilio</i> (Schlumberger, 1967)			
<i>williamsoni</i> (Silvestri, 1923)			
= <i>Biloculina williamsoni</i> Silvestri, 1923			
= <i>Biloculinella elongata</i> (Wiesner, 1923)			
= <i>P. elongata</i> (d'Orbigny, 1826)			
= <i>Quinqueloculina elongata</i> Natland, 1938			
= <i>Spiroloculina elongata</i> d'Orbigny, 1839			
<i>Pyrgoella</i>			
<i>irregularis</i> (d'Orbigny, 1839)			A
= <i>Biloculinella irregularia</i> (d'Orbigny, 1839)			
= <i>Nummoloculina irregularis</i> (d'Orbigny, 1839)			
= <i>Siphonaptera irregularis</i> (d'Orbigny, 1826)			
<i>sphaera</i> (d'Orbigny, 1839)			A M
<i>Pyrulina</i>			
<i>cylindroides</i> (Roemer, 1838)			
<i>gutta</i> d'Orbigny, 1826			
<i>Quinqueloculina</i>			
<i>akneriana</i> d'Orbigny, 1846			A
<i>anguina</i> Terquem, 1876			M
<i>angustidentata</i> Terquem, 1876			
<i>annectens</i> (Schlumberger, 1893)			A M
= <i>Massilina annectens</i> (Schlumberger, 1893)			
<i>arctica</i> Cushman, 1933			A
<i>auberiana</i> d'Orbigny, 1826			M
<i>berthelotiana</i> d'Orbigny, 1839			M
<i>bicarinata</i> d'Orbigny, 1826			M
<i>bidentata</i> d'Orbigny, 1839			A M
= <i>Spiroloculina bidentata</i> (Hadley, 1935)			
<i>bosciana</i> d'Orbigny, 1839			M
<i>bradyana</i> Cushman, 1917			
= <i>Q. undosa</i> Karrer, 1868			
<i>concava</i> Reuss			M
<i>cuvieriana</i> d'Orbigny, 1839			M
<i>denticulata</i> O.G. Costa			M
<i>depressa</i> d'Orbigny, 1839			M
<i>dimidiata</i> Terquem, 1876			A
<i>disciformis</i> Williamson			M
<i>disparilis</i> d'Orbigny, 1826			M
<i>dunkeriana</i> (Heron-Allen & Earland, 1930)			A
= <i>Miliolinella dunkeriana</i> Heron-Allen & Earland, 1930			
<i>dutemplei</i> d'Orbigny, 1846			M
ex gr. <i>agglutinans</i> d'Orbigny, 1839			A
<i>flexuosa</i> d'Orbigny, 1839			M
<i>goesi</i> (Wiesner, 1923)			M
<i>inconstans</i> Terquem, 1876			M
<i>jugosa</i> Cushman, 1944			M
<i>laevigata</i> d'Orbigny, 1839			M
<i>lamarckiana</i> d'Orbigny, 1839			M
<i>lata</i> Terquem, 1876			A
<i>limbata</i> d'Orbigny, 1826			M
<i>linnaeana</i> (d'Orbigny, 1839)			M
<i>longidentata</i> Terquem			M
<i>lucida</i> Karrer, 1868			M
<i>milletti</i> (Wiesner, 1923)			M
= <i>Massilina milletti</i> (Wiesner, 1923)			
<i>nodulosa</i> Wiesner, 1923			M
<i>nussdorfensis</i> d'Orbigny, 1839			M
<i>oblonga</i> (Montagu, 1893)			M
= <i>Pseudotriloculina oblonga</i> (Montagu, 1803)			
= <i>Pyrgo oblonga</i> (d'Orbigny, 1839)			
= <i>Triloculina oblonga</i> (Montagu, 1803)			
<i>osinclinatum</i> Le Calvez			M
<i>ovulcula</i> Terquem, 1876			M
<i>padana</i> Perconig, 1954			A M
<i>parvula</i> Schlumberger, 1894			M
<i>pauperata</i> d'Orbigny, 1839			M
<i>peregrina</i> d'Orbigny, 1839			M
<i>peruviana</i> d'Orbigny, 1839			M
<i>phoenica</i> Martinotti			M
<i>polygona</i> d'Orbigny, 1839			M
<i>pseudobuchiana</i> Luczkowska, 1974			M
<i>pseudoseminula</i> Mikhalevich, 1968			M
<i>pygmaea</i> Reuss, 1850			A M
<i>rugosa</i> d'Orbigny, 1839			A M

<i>sadkovi</i> (Shchedrina, 1946)	A	<i>Triloculina</i>	
= <i>Miliolina sadkovi</i> Shchedrina, 1946		<i>adriatica</i> Le Calvez, J. & Y., 1958	M
<i>schlumbergeri</i> (Wiesner, 1923)	A M	<i>affinis</i> d'Orbigny, 1826	M
= <i>Miliolinella schlumbergeri</i> Wiesner, 1923		<i>costifera</i> Terquem, 1876	M
<i>sclerotica</i> Karrer, 1868	A	<i>eburnea</i> d'Orbigny, 1839	M
<i>seminula</i> (Linné, 1758)	A M	<i>fischeri</i> Schlumberger	M
= <i>Serpula seminulum</i> Linné, 1758		<i>frigida</i> Lagoe, 1977	A M
<i>semistriata</i> d'Orbigny, 1839	M	<i>gibba</i> d'Orbigny, 1839	M
<i>sinuosa</i> Terquem, 1876	M	<i>hemisphaerica</i> Terquem, 1876	M
<i>stalkeri</i> Loeblich & Tappan, 1953	A M	<i>inflata</i> d'Orbigny, 1826	A M
<i>stelligera</i> Schlumberger, 1893	M	<i>lamellidens</i> Reuss	M
<i>subcarinata</i> d'Orbigny, 1839	M	<i>longirostra</i> (d'Orbigny, 1826)	A M
<i>tricarinella</i> (Wiesner)	M	= <i>Adelosina longirostra</i> (d'Orbigny, 1846)	
<i>ungeriana</i> d'Orbigny, 1846	A M	= <i>Quinqueloculina longirostra</i> d'Orbigny, 1826	
<i>variolata</i> d'Orbigny, 1826	M	<i>marioni</i> Schlumberger, 1893	M
<i>venusta</i> Karrer, 1868	A M	<i>ornata</i> Le Calvez, J. & Y., 1958	M
<i>viennensis</i> Le Calvez & Le Calvez, 1958	M	<i>plicata</i> Terquem, 1878	M
<i>vulgaris</i> d'Orbigny, 1826	A M	<i>rotunda</i> d'Orbigny, 1826	A
<i>Sigmoilina</i>		= <i>Pseudotriloculina rotunda</i> (d'Orbigny, 1826)	
<i>coelata</i> O.G. Costa, 1855	M	= <i>Spiroloculina rotunda</i> d'Orbigny, 1826	
<i>sigmoidea</i> (Brady, 1884)	M	= <i>Spiroloculina rotundata</i> d'Orbigny, 1826	
<i>distorta</i> Phleger & Parker, 1951	A M	<i>schreibersiana</i> d'Orbigny, 1839	M
<i>Sigmoilinella</i>		= <i>Quinqueloculina schreibersiana</i> (d'Orbigny, 1839)	
<i>costata</i> (Schlumberger, 1893)	M	<i>terquemiana</i> Brady, 1884	M
= <i>Nummoloculina contraria</i> (d'Orbigny, 1846)		<i>tricarinata</i> d'Orbigny, 1826	A M
= <i>Sigmoilina costata</i> Schlumberger		= <i>Spiroloculina tricarinata</i> d'Orbigny, 1839	
<i>grata</i> (Terquem, 1878)	M	<i>trigona</i> (Lamarck, 1804)	A M
= <i>Sigmoilina grata</i> (Terquem, 1878)		= <i>Miliolites trigonula</i> Lamarck, 1804	
<i>tenuis</i> (Czjek, 1848)	M	<i>Zoiaella</i>	
= <i>Sigmoilina tenuis</i> (Czjek, 1848)		<i>trochamminoides</i> (Goes, 1879)	
<i>Sigmoilopsis</i>		<i>Discospirina</i>	
<i>asperula</i> (Karrer, 1868)	A M	<i>italica</i> (Costa, 1856)	M
= <i>Ammomassilina asperula</i> (Karrer, 1868)		= <i>D. tenuissima</i> (Costa, 1856)	
<i>procericollis</i> Mikhalevich, 1977	A	<i>Pateoris</i>	
<i>schlumbergeri</i> (Silvestri, 1904)	A M	<i>hauerinoides</i> (Rhumbler, 1936)	A
= <i>Sigmoilina schlumbergeri</i> Silvestri, 1904		Family Nubeculariidae	
<i>Siphonaptera</i>		<i>Calcituba</i>	
<i>agglutinans</i> (d'Orbigny, 1839)	M	<i>polymorpha</i> v. Roboz, 1884	M
<i>aspera</i> (d'Orbigny, 1826)	M	<i>Fischerina</i>	
= <i>Quinqueloculina aspera</i> (d'Orbigny, 1826)		<i>communis</i> (Seguenza, 1880)	A
<i>dilatata</i> (Le Calvez, J. & Y., 1958)	M	<i>schlumbergi</i> (Sidebottom)	M
<i>hauerina</i> (Wiesner, 1923)	M	<i>Nodobacularia</i>	
<i>Spiroloculina</i>		<i>fusca</i> Wiesner, 1923	M
<i>angulifera</i> Terquem, 1876	M	<i>lapidea</i> Wiesner, 1923	M
<i>angulosa</i> Terquem, 1878	A M	<i>Nubecularia</i>	
<i>antillarium</i> d'Orbigny, 1839	M	<i>elongata</i> Terquem, 1878	M
<i>arenilega</i> Rhumbler, 1909	A	<i>lucifuga</i> Defrance, 1825	A M
<i>atlantica</i> Cushman, 1947	A	<i>Nubeculina</i>	
<i>barthelotiana</i> Wiesner, 1923	M	<i>divaricata</i> (Brady, 1879)	M
<i>californica</i> Cushman & Todd, 1944	A	<i>Planispirinella</i>	
<i>communis</i> Cushman & Todd, 1944	A	<i>exigua</i> (Brady, 1879)	M
= <i>Quinqueloculina canaliculata</i> Terquem, 1876		<i>Vertebralina</i>	
= <i>S. canaliculata</i> d'Orbigny, 1846		<i>striata</i> d'Orbigny, 1826	M
<i>cymbium</i> d'Orbigny, 1839	M	Family Ophtalmidiidae	
<i>depressa</i> d'Orbigny, 1826	M	<i>Cornuloculina</i>	
<i>dilatata</i> d'Orbigny, 1846	M	<i>balkwilli</i> (Macfayden, 1939)	
<i>elevata</i> Wiesner, 1923	M	<i>inconstans</i> (Brady, 1879) Loeblich & Tappan, 1964	A M
<i>excavata</i> d'Orbigny, 1846	M	<i>Nodophthalmidium</i>	
<i>grateloupi</i> (d'Orbigny)		<i>simplex</i> Cushman & Todd, 1947	M
<i>henbesti</i> Thalman, 1955	M	= <i>Nodobacularia tibia</i> (Jones & Parker, 1860)	
= <i>S. planulata</i> Lamarck	M	<i>Ophthalmina</i>	
<i>interrupta</i> Terquem, 1876	M	<i>kiliansensis</i> Rhumbler, 1936	
<i>krumbachi</i> Wiesner, 1912	M	<i>Spirophthalmidium</i>	
<i>lancea</i> Ehrenberg, 1861	A	<i>acutimargo</i> (Brady, 1884)	A M
<i>lyra</i> d'Orbigny, 1839	M	<i>longiscata</i> (Terquem & Berthelin, 1875)	A
<i>ornata</i> d'Orbigny, 1839	M	<i>Spirosigmoilina</i>	
<i>ornata</i> d'Orbigny, 1839 var. <i>tricarinata</i> Le Calvez, 1958	M	<i>pusilla</i> (Earland, 1934)	A
<i>perforata</i> d'Orbigny, 1839	M	= <i>Spirophthalmidium pusillum</i> (Earland, 1934)	
<i>planissima</i> Lamarck	A	Family Peneroplididae	
<i>regularis</i> Cushman & Todd, 1944	M	<i>Archaias</i>	
<i>rostrata</i> Reuss	M	<i>angulatus</i> (Fichtel & Moll, 1798)	M
<i>subangulosa</i> Terquem, 1876	M	<i>Coscinospira</i>	
<i>subimpressa</i> Parr	A	<i>arietina</i> (Batsch, 1791)	M
<i>tenuiseptata</i> Brady, 1884	M	= <i>Coscinopira hemprichii</i> Ehrenberg, 1839	

= <i>Spirolina arietina</i> (Batsch, 1791) Jones, 1994		
<i>Laevipeneroplis</i>		
<i>karreri</i> (Wiesner, 1923)	M	
<i>Peneroplis</i>		
<i>arbusculus</i> (d'Orbigny, 1839)	M	
<i>bradyi</i> Cushman, 1930	M	
= <i>P. laevigatus</i> (Karrer, 1868)		
<i>ellipticus</i> d'Orbigny, 1839	M	
<i>pertusus</i> (Forskal, 1775)	A M	
= <i>Nautilus pertusus</i> Forskal, 1775		
<i>planatus</i> (Fichtel & Moll, 1798)	A M	
= <i>Nautilus planatus</i> Fichtel & Moll, 1798		
<i>proteus</i> d'Orbigny, 1839	M	
<i>Spirolina</i>		
<i>cylindracea</i> (Lamarck, 1804) Jones, 1994	M	
= <i>Peneroplis cylindracus</i> (Lamarck, 1804)		
Family Soritidae		
<i>Amphisorus</i>		
<i>hemprichii</i> Ehrenberg, 1839	A M	
<i>Sorites</i>		
<i>orbiculus</i> (Forskal, 1775)	M	
= <i>Nautilus orbiculus</i> Forskal, 1775		
<i>orbitolitoides</i> (Hofker)	M	
Suborder Rotaliina		
Family Acervulinidae		
<i>Acervulina</i>		
<i>inhaerens</i> Schultze, 1854	A M	
<i>Gypsina</i>		
<i>vesicularis</i> (Parker & Jones)	M	
<i>Sphaerogypsina</i>		
<i>globula</i> (Reuss, 1848)	M	
Family Alabaminidae		
<i>Gyroidina</i>		
<i>laevigata</i> (d'Orbigny, 1826)	M	
<i>longispira</i> Tedeschi & Zanmatti, 1957	M	
<i>Osangularia</i>		
<i>cultur</i> (Parker & Jones, 1865)	A	
<i>Osangulariella</i>		
<i>umbonifera</i> (Cushman, 1933)	A	
= <i>Osangularia umbonifera</i> (Cushman, 1933)		
<i>Svartkina</i>		
<i>tuberculata</i> (Balkwill & Wright, 1885)	M	
Family Ammoniidae		
<i>Ammonia</i>		
<i>aberdoeyensis</i> Haynes, 1973	A	
<i>ammoniformis</i> (d'Orbigny, 1846)	M	
<i>batava</i> (Hofker, 1951)	A	
= <i>A. batavus</i> (Hofker, 1951)		
= <i>Streblius batavus</i> Hofker, 1956		
<i>beccarii</i> (Linné, 1758)	A M	
= <i>Nautilus beccarii</i> Linné, 1758	A	
<i>falsobeccarii</i> (Rouvilleo, 1974)	A	
<i>flevenensis</i> (Hofker)	A	
<i>inflata</i> (Seguenza, 1862)	M	
<i>limnetes</i> (Todd & Brörmimarm, 1957)	A	
<i>papillosa</i> (Brady)	M	
<i>parkinsoniana</i> (d'Orbigny, 1839)	M	
= <i>Rosalina parkinsoniana</i> d'Orbigny, 1839		
<i>perlucida</i> (Heron-Allen and Earland, 1913)	A M	
= <i>Aubignyna planidorso</i> (Atkinson, 1989)		
= <i>Rotalia perlucida</i> Heron-Allen and Earland, 1913	A M	
<i>tepida</i> (Cushman, 1926)		
Family Amphisteginidae		
<i>Amphistegina</i>		
<i>gibbosa</i> d'Orbigny, 1839	A	
<i>lessonii</i> d'Orbigny, 1826	A M	
<i>madagascariensis</i> d'Orbigny, 1903	A M	
Family Anomaliniidae		
<i>Anomalina</i>		
<i>ammonoides</i> (Reuss)	M	
<i>Melonis</i>		
<i>umbilicatulus</i> (Walker & Jacob, 1798)	M	
<i>Cibicides</i>		
		A
		A
		M
		A
		A
		A M
		A
		A
		Cibicidoides
		<i>cicatricosus</i> (Schwager, 1866)
		<i>floridanus</i> (Cushman, 1918)
		<i>globulosus</i> Chapman & Parr, 1937
		= <i>Anomalina globulosa</i> Chapman & Parr, 1937
		<i>kullenbergi</i> (Parker, 1953)
		= <i>Cibicides kullenbergi</i> Parker, 1953
		<i>pachyderma</i> (Rzehak, 1886)
		<i>phlegeri</i> (Uchio, 1960)
		<i>pseudoungerianus</i> (Cushman, 1922)
		= <i>Cibicides pseudoungerianus</i> (Cushman, 1931)
		= <i>Truncatulina pseudoungeriana</i> Cushman, 1922
		<i>robertsonianus</i> (Brady, 1881)
		<i>rugosus</i> (Phleger & Parker, 1951)
		<i>wuellerstorfi</i> (Schwager, 1866)
		= <i>Anomalina wuellerstorfi</i> Schwager, 1866
		= <i>Fontbotia wuellerstorfi</i> (Schwager, 1866)
		= <i>Planulina wuellerstorfi</i> (Schwager, 1866)
		<i>Discanomalina</i>
		<i>coronata</i> (Parker & Jones, 1857)
		= <i>Paromalina coronata</i> (Parker & Jones, 1865)
		<i>Discoanomalina</i>
		<i>semiunctata</i> (Bailey, 1851)
		= <i>Rotalina semiunctata</i> Bailey, 1851
		<i>Gyroidina</i>
		<i>altiformis</i> R.E. & K.C. Stewart, 1930
		<i>neosoldanii</i> Brotzen, 1936
		= <i>Gyroidinoides neosoldanii</i> (Brotzen, 1936)
		<i>orbicularis</i> d'Orbigny, 1826
		= <i>Gyroidinoides orbicularis</i> d'Orbigny, 1826
		<i>polia</i> (Phleger & Parker, 1951)
		<i>zeelandica</i> Finley, 1939
		<i>Nuttallides</i>
		<i>rugosus</i> (Phleger & Parker, 1951)
		= <i>Pseudoparella rugosa</i> Phleger & Parker, 1951
		<i>umboniferus</i> (Cushman, 1933)
		= <i>Pulvinulina umbonifera</i> Cushman, 1933
		<i>Planulina</i>
		<i>ariminensis</i> d'Orbigny, 1826
		<i>Family Asterigerinatidae</i>
		<i>Asterigerinata</i>
		<i>mamilla</i> (Williamson, 1858)
		= <i>Eponidella mamilla</i> (Williamson, 1858)
		= <i>Rotalia mamilla</i> Williamson, 1858
		<i>Family Bagginiidae</i>
		<i>Cancris</i>
		<i>auricula</i> (Fichtel & Moll, 1798)
		= <i>Nautilus auricula</i> Fichtel & Moll, 1798
		<i>congoensis</i> Margerel & Kouyoumotzakis, 1978
		<i>oblongus</i> (Williamson, 1858)
		<i>Valvularinia</i>
		<i>allomorphinoides</i> (Reuss, 1860)
		<i>araucana</i> (d'Orbigny, 1932)
		<i>arctica</i> Green, 1960
		<i>bradyana</i> (Fornasini, 1900)
		<i>complanata</i> (d'Orbigny, 1846)
		<i>laevigata</i> Phleger & Parker, 1951
		<i>obtusa</i> (d'Orbigny, 1839)
		<i>Family Bolivinidae</i>
		<i>Bolivina</i>
		<i>albatrossi</i> Cushman, 1922

<i>arctica</i> Herman, 1973	A	<i>elegans</i> d'Orbigny, 1826	A M
<i>britannica</i> MacFadyen, 1942	A	<i>elongata</i> d'Orbigny, 1826	A M
<i>cincta</i> Heron-Allen & Earland, 1932	A	<i>etnea</i> Seguenza	M
<i>compacta</i> Sidebottom, 1905	A	<i>fusiformis</i> subsp. <i>marginata</i> Fornasini, 1901	M
<i>compacta</i> Sidebottom, 1905	A	<i>gibba</i> Fornasini, 1902	A M
<i>difformis</i> (Williamson)	A	<i>inflata</i> Seguenza, 1862	A M
<i>inflata</i> Heron-Allen & Earland, 1913	A	<i>marginata</i> d'Orbigny, 1826	A M
<i>minima</i> Phleger & Parker, 1951	A	<i>striata</i> d'Orbigny, 1826	A
<i>ordinaria</i> Phleger & Parker, 1952	A	<i>striata</i> subsp. <i>mexicana</i> Cushman, 1922	A M
<i>pacifica</i> Cushman & McCulloch, 1942	A	<i>subornata</i> (Brady, 1884)	A
<i>persiensis</i> Lutze, 1974	A	<i>subulata</i> Cushman & Parker, 1947	A
<i>plicata</i> (d'Orbigny, 1848)	M	= <i>Laryngosigma subulata</i> (Chapman & Parr, 1937)	
<i>pseudoplicata</i> Heron-Allen & Earland, 1930	A M	<i>Furcenkoina</i>	
<i>quadrilatera</i> (Schwager)	A	<i>acuta</i> (d'Orbigny, 1846)	M
<i>superba</i> Emiliani, 1949	A	<i>bradyi</i> (Cushman, 1922)	A
<i>thalmanni</i> Renz, 1948	A	<i>davisi</i> (Chapman & Parr, 1931)	A
<i>tongi</i> Cushman, 1929	A	= <i>Virgulina davisi</i> Chapman & Parr, 1931	
<i>variabilis</i> (Williamson, 1858)	A M	<i>rotundata</i> (Parr, 1950)	A
= <i>Brizalina variabilis</i> (Williamson, 1858)		= <i>Virgulina rotundata</i> Parr, 1950	
<i>Bolivinella</i>		<i>schreibersiana</i> (Czjzek, 1848)	
<i>pseudopunctata</i> (Högglund, 1947)	A M	<i>squammosa</i> (d'Orbigny, 1839)	M
= <i>Bolivina pseudopunctata</i> Högglund, 1947		<i>subsquammosa</i> (d'Orbigny, 1839)	M
<i>Brizalina</i>		<i>Globobulimina</i>	
<i>alata</i> (Seguenza, 1862)	M	<i>affinis</i> (d'Orbigny, 1839)	A M
= <i>Bolivina alata</i> (Seguenza, 1862)		= <i>Bulimina affinis</i> d'Orbigny, 1839	
= <i>Vulvulina alata</i> Seguenza, 1862		<i>pseudospinescens</i> (Emiliani)	M
<i>cataensis</i> (Seguenza, 1862)	A M	<i>turgida</i> (Bailey, 1851)	M
= <i>Bolivina cataensis</i> Seguenza, 1862		= <i>Bulimina turgida</i> Bailey, 1851	
<i>difformis</i> (Williamson, 1858)	A M	<i>Praeglobobulimina</i>	
<i>dilatata</i> (Reuss, 1850)	M	<i>pupoides</i> (d'Orbigny, 1846)	M
<i>earlandi</i> (Parr, 1950)	M	= <i>Pseudoglobobulimina pupoides</i> (d'Orbigny, 1846)	
= <i>Bolivina punctata</i> (d'Orbigny, 1848)		<i>spinescens</i> (Brady, 1884)	A
<i>lowmani</i> (Phleger & Parker, 1951)	A	= <i>Bulimina pyrula</i> d'Orbigny, 1846 var. <i>spinescens</i>	
= <i>Bolivina lowmani</i> Phleger & Parker, 1951		Brady, 1888	
<i>pygmaea</i> (Brady, 1881)	A	<i>Pseudobolivina</i>	
= <i>Bolivina pygmaea</i> (Brady, 1881)		<i>antarctica</i> (Wiesner, 1931)	A
<i>skagerrakensis</i> (Qvale & Nigam, 1985)	A	<i>fusiformis</i> (Chaster, 1892)	A
= <i>Bolivina skagerrakensis</i> Qvale & Nigam, 1985		<i>Stainforthia</i>	
<i>spathulata</i> (Williamson, 1858)	A M	<i>complanata</i> (Egger, 1893)	A
= <i>Bolivina dilatata</i> Reuss, 1850		<i>concava</i> (Högglund, 1947)	A M
= <i>Bolivina spathulata</i> (Williamson, 1858)		<i>fusiformis</i> (Williamson, 1858)	A
<i>striatula</i> (Cushman, 1922)	A M	= <i>Bulimina pupoides</i> d'Orbigny, 1846 var. <i>fusiformis</i>	
= <i>Bolivina striatula</i> Cushman, 1922		Williamson, 1858	
<i>subaenariensis</i> (Cushman, 1922)	A M	= <i>Furcenkoina fusiformis</i> (Williamson, 1858)	
= <i>Bolivina subaenariensis</i> Cushman, 1922		<i>loeblichi</i> Feyling-Hanssen, 1954	A
<i>subspathulata</i> (Boomgart, 1949)	A	= <i>Furcenkoina loeblichi</i> (Feyling-Hanssen, 1954)	
= <i>Bolivina subspathulata</i> (Boomgart, 1949)	M	<i>Stetsonia</i>	
<i>subspinescens</i> (Cushman, 1922)		<i>arctica</i> Green, 1960	A
= <i>Bolivina subspinescens</i> Cushman, 1922		<i>horvathi</i> Green, 1960	A
= <i>Bolivina textileoides</i> Reuss, 1850		Family Cassidulinidae	
= <i>B. spinescens</i> (Cushman, 1911)		<i>Cassidulina</i>	
<i>Loxostomina</i>		<i>crassa</i> d'Orbigny, 1839	A M
<i>limbata</i> (Brady, 1881)	A	= <i>C. oblonga</i> Reuss, 1850	
= <i>Bolivina limbata</i> Brady, 1881		= <i>Globocassidulina oblonga</i> (Reuss, 1850)	
<i>Rectobolivina</i>		<i>laevigata</i> d'Orbigny, 1826	A M
<i>columellaris</i> (Brady, 1884)	M	<i>mexicana</i> (Cushman, 1922)	A
Family Bolivinitidae		= <i>Rutherfordoides mexicana</i> Cushman, 1922	
<i>Abditodentrix</i>		= <i>Virgulina mexicana</i> Cushman, 1922	
<i>rhomboidalis</i> (Millett, 1899)	M	<i>minuta</i> Cushman, 1933	A
<i>Orthomorphina</i>		<i>obtusa</i> Williamson, 1858	A
<i>challengeriana</i> (Thalmann, 1937)	A	<i>reniforme</i> (Norvang, 1945)	A
<i>radicula</i> (Linné, 1804)	M	= <i>C. crassa</i> d'Orbigny, 1839 var. <i>reinforme</i> Norvang, 1945	
<i>Stilosomella</i>		<i>teretis</i> Tappan, 1951	A M
<i>advena</i> (Cushman & Laiming, 1931)	A	= <i>Islandiella teretis</i> (Tappan, 1951)	
<i>antillea</i> (Cushman, 1923)	A	<i>Cassidulinoides</i>	
<i>consorbina</i> (d'Orbigny, 1839)	M	<i>bradyi</i> (Norman, 1881)	A M
<i>fistuca</i> (Schwager, 1866)	A	<i>Ehrenbergina</i>	
<i>longiscata</i> (d'Orbigny, 1839)	M	<i>bradyi</i> Cushman, 1922	A
<i>pauperata</i> (d'Orbigny, 1839)	M	<i>trigona</i> (Goes, 1896)	A
Family Buliminidae		<i>Globocassidulina</i>	
<i>Bulimina</i>		<i>subglobosa</i> (Brady, 1881)	A M
<i>aculeata</i> d'Orbigny, 1826	A M	= <i>Cassidulina subglobosa</i> (Brady, 1881)	
<i>alazanensis</i> Cushman, 1927	A M	<i>Islandiella</i>	
<i>buchiana</i> d'Orbigny	M	<i>australis</i> (Phleger & Parker, 1951)	A
<i>costata</i> d'Orbigny, 1852	A M		

<i>helenae</i> Feyling-Hanssen & Buzas, 1976	A	<i>pauperata</i> (Parker & Jones, 1865)	A
<i>islandica</i> (Norvang, 1945)	A	<i>Neoponides</i>	
= <i>Cassidulina islandica</i> Norvang, 1945		<i>auberii</i> (d'Orbigny, 1839)	M
<i>norcrossi</i> (Cushman, 1933)	A	= <i>Discorbis turbo</i> (d'Orbigny, 1839)	
= <i>Cassidulina norcrossi</i> Cushman, 1933		<i>Rosalina</i>	
<i>Paracassidulina</i>		<i>anomala</i> Terquem, 1875	A
<i>neocarinata</i> (Thalman, 1950)	A	<i>bradyi</i> (Cushman, 1915)	M
= <i>Cassidulina neocarinata</i> (Thalman, 1950)		<i>complanata</i> d'Orbigny, 1846	A
<i>Sphaeroidina</i>	A M	<i>floridana</i> (Cushman, 1922)	M
<i>bulloides</i> (d'Orbigny, 1826)		= <i>Discorbis floridana</i> Cushman, 1922	
Family Ceratobuliminidae		= <i>R. floridensis</i> (Cushman, 1922)	
<i>Ceratobulimina</i>		<i>globularis</i> d'Orbigny, 1826	A M
<i>arctica</i> Green, 1960	A	<i>irregularis</i> (Rhumbler, 1906)	A
<i>jonesiana</i> (Brady, 1881)	A	<i>macropora</i> (Hofker, 1951)	M
= <i>C. pacifica</i> Cushman & Harris, 1927		<i>neapolitana</i> (Hofker, 1951)	A
<i>Ceratocancris</i>	M	<i>posidonicola</i> (Colom, 1942)	M
<i>haueri</i> (d'Orbigny, 1839)		<i>valvulata</i> d'Orbigny	M
<i>Lamarcinia</i>	A	<i>williamsoni</i> (Chapman & Parr, 1932)	A
<i>halictidea</i> (Heron-Allen & Earland)		= <i>Discorbis williamsoni</i> Chapman & Parr, 1932	
Family Chilostomellidae		= <i>Neocorbinella williamsoni</i> (Chapman & Parr, 1932)	
<i>Pullenia</i>		<i>Rotaliella</i>	
<i>bulloides</i> (d'Orbigny, 1826)	A	<i>chasteri</i> (Heron-Allen & Earland, 1913)	
= <i>Nonionina bulloides</i> d'Orbigny, 1846		<i>Strebloides</i>	
= <i>Nonionina sphaerooides</i> d'Orbigny, 1826		<i>advena</i> (Cushman, 1922)	M
<i>osloensis</i> Feyling-Hanssen, 1954	A	= <i>Cibicidina advena</i> (d'Orbigny, 1839)	
<i>quinqueloba</i> (Reuss, 1851)	A M	= <i>Discorbis advena</i> (Cushman, 1922)	
= <i>Nonionina quinqueloba</i> Reuss, 1851		<i>Tretomphalus</i>	
<i>salisburyi</i> R. E. & K.C. Stewart, 1930	M	<i>bulloides</i> (d'Orbigny, 1839)	M
<i>sphaerooides</i> (d'Orbigny, 1839)	M	Family Discorbinellidae	
<i>subcarinata</i> d'Orbigny, 1839	A M	<i>Hanzawaia</i>	
Family Cibicididae		<i>boueana</i> (d'Orbigny)	M
<i>Cyclocibicides</i>		<i>concentrica</i> (Cushman, 1918)	A
<i>vermiculatus</i> (d'Orbigny, 1826)	M	= <i>Cibicides concentricus</i> Cushman, 1918	
= <i>Discoanomalina vermiculata</i> (d'Orbigny, 1839)		<i>nitidula</i> (Bandy)	M
Family Cymbaloporidae		<i>rhodiensis</i> (Terquem, 1878)	A
<i>Cymbalopora</i>	M	= <i>Discorbinella rhodiensis</i> (Terquem, 1878)	
<i>tabelaeformis</i> (Brady)		<i>solei</i> Colom, 1957	A
<i>Cymbaloporeta</i>	M	Family Ellipsolagenidae	
<i>squamosa</i> (d'Orbigny)		<i>Entosolenia</i>	
Family Discorbidae		<i>lineata</i> (Williamson, 1848)	M
<i>Buccella</i>		= <i>Favulina lineata</i> (Williamson, 1848)	
<i>florifomis</i> Voloshinova, 1960	A	= <i>Oolina lineata</i> (Williamson, 1848)	
<i>frigida</i> (Cushman, 1921)	A M	<i>Favulina</i>	
= <i>Pulvinulina frigida</i> Cushman, 1921		<i>hexagona</i> (Williamson, 1848)	M
<i>granulata</i> (di Napoli Alliata, 1952)	A M	= <i>Oolina hexagona</i> (Williamson, 1848)	
<i>karsteni</i> (Reuss)	M	<i>melo</i> (d'Orbigny, 1839)	A
<i>tenerrima</i> (Bandy, 1950)	A M	<i>scalariformis</i> (Williamson, 1848)	M
= <i>Rotalia tenerrima</i> Bandy, 1950		<i>variata</i> (Brady, 1884)	M
<i>Disconorbis</i>	M	<i>Fissurina</i>	
<i>bulbosus</i> (Parker, 1954)		<i>alveolata</i> (Brady, 1884)	A
= <i>Discorbis bulbosus</i> (Parker, 1954)		<i>biancae</i> Seguenza, 1862	A
<i>Discorbinella</i>	A M	<i>cucullata</i> Silvestri, 1902	A M
<i>bertheloti</i> (d'Orbigny, 1839)		<i>danica</i> (Madsen, 1895)	A
<i>Discorbs</i>		<i>elliptica</i> (Cushman, 1923)	A
<i>globularis</i> Parr, 1950	M	<i>fimbriata</i> (Brady, 1881)	A
<i>orbicularis</i> (Terquem, 1875)	M	<i>kerguelensis</i> Parr, 1950	A
<i>rosacea</i> (d'Orbigny, 1839)	M	<i>laevigata</i> Reuss, 1850	A M
<i>villardeboanus</i> (d'Orbigny, 1839)	A M	<i>lucida</i> (Williamson, 1848)	M
= <i>Rosalina villardeboana</i> d'Orbigny, 1839		<i>marginata</i> (Montagu, 1803) (Walker & Boys?)	A M
= <i>Rosalina villardeboanus</i> d'Orbigny, 1839		<i>neptuni</i> (Büchner, 1940)	A M
<i>wrightii</i> (Brady, 1881)	A M	<i>orbigniana</i> Seguenza, 1862	A M
<i>Discorbinoides</i>		= <i>Palliolatella orbigniana</i> (Seguenza, 1862)	
<i>milletti</i> (Wright, 1911)		<i>radiata</i> (Rhumbler, 1911)	A M
= <i>Rosalina milletti</i> (Wright, 1911)		<i>semimarginata</i> (Boomgart, 1949)	A
<i>Gavelinopsis</i>		= <i>Entosolenia semimarginata</i> Boomgart, 1949	
<i>isabellanus</i> (d'Orbigny, 1839)	M	<i>serrata</i> (Schlumberger)	A
<i>lobatula</i> (Parr, 1950)	M	<i>truncata</i> (Brady, 1884)	A
= <i>Discorbis lobatulus</i> (Parr, 1950)		<i>Lagenosolenia</i>	
<i>praegeri</i> (Heron-Allen & Earland, 1913)	A M	<i>seguenziana</i> (Fornasini, 1886)	M
= <i>Discorbina praegeri</i> Heron-Allen & Earland, 1913		<i>Oolina</i>	
= <i>Rosalina praegeri</i> (Heron-Allen & Earland)		<i>ampulladistoma</i> (Rymer-Jones, 1874)	M
<i>translucens</i> (Phleger & Parker, 1951)	A M	= <i>Lagena ampulladistoma</i> (Rymer-Jones, 1874)	
<i>Laticarinina</i>		<i>borealis</i> Loeblich & Tappan, 1954	A
<i>halophora</i> (Stache, 1864)	A	= <i>O. apiopleura</i> (Loeblich & Tappan, 1953)	
		<i>globosa</i> (Montagu, 1803)	A

= <i>Favulina globosa</i> (Montagu, 1803)		
<i>heronalleni</i> Haynes, 1973	A	
<i>laevigata</i> d'Orbigny, 1839	A	
<i>longispina</i> (Brady, 1881)	A	
<i>melo</i> d'Orbigny, 1839	A	
<i>ovum</i> Ehrenberg, 1843 var. <i>pontica</i> Mikhalevich, 1968	M	
= <i>Lagena ovum</i> Ehrenberg, 1843 var. <i>pontica</i> Mikhalevich, 1968		
<i>squamosa</i> (Montagu, 1803)	A	
<i>williamsoni</i> (Alcock, 1865)	A	
Parafissurina		
<i>arctica</i> Green, 1960	A	
<i>fusiformis</i> (Wiesner, 1931)	A	
<i>fusuliformis</i> Loeblich & Tappan, 1953	A	
<i>groenlandica</i> Shchedrina, 1946	A	
<i>lateralis</i> (Cushman, 1913)	A M	
= <i>Lagena lateralis</i> (Cushman, 1913) var. <i>pontica</i> Mikhalevich, 1968		
<i>tectulostoma</i> Loeblich & Tappan, 1953	A	
Family Elphidiidae		
Criboelphidium		
<i>poeyanum</i> (d'Orbigny, 1826)	M	
= <i>Elphidium poeyanum</i> (d'Orbigny, 1826)		
= <i>Polystomella poeyanum</i> d'Orbigny, 1826		
<i>albiumbilicatum</i> (Weiss, 1954)	A	
= <i>Elphidium albiumbilicatum</i> (Weiss, 1954)		
= <i>Haynesina albiumbilicatula</i> (Weiss, 1954)		
= <i>Nonion pauciloculum</i> Cushman, 1954 var. <i>albiumbilicatum</i> Weiss, 1954		
<i>depressum</i> (Walker & Jacob, 1784)	M	
<i>hallandense</i> (Brotzen, 1943)	A	
<i>incertum</i> Williamson, 1858	A	
= <i>Elphidium incertum</i> (Williamson, 1858)		
= <i>Polystomella umbilicatula</i> Williamson, 1858 var. <i>incerta</i> Williamson, 1858		
<i>magellanicum</i> Heron-Allen & Earland, 1932	A	
<i>maricobi</i> (Bogdanovich, 1947)	M	
Cribrononion		
<i>excavatum</i> (Terquem, 1875)	M	
<i>incertum</i> (Williamson, 1858)	M	
<i>translucens</i> (Natland, 1938)	M	
Elphidiella		
<i>arctica</i> (Parker & Jones, 1864)	A	
= <i>Elphidium arcticum</i> (Parker & Jones, 1864)		
<i>groenlandica</i> (Cushman, 1936)	A	
<i>nitida</i> Cushman, 1941	A	
Elphidium		
<i>aculeatum</i> (d'Orbigny, 1846)	M	
<i>advenum</i> (Cushman, 1922)	A M	
= <i>Cribrononion advenum</i> (Cushman, 1922)		
= <i>Polystomella advena</i> Cushman, 1922		
<i>articulatum</i> (d'Orbigny, 1839)	A	
<i>asklundi</i> Brotzen, 1943	A	
<i>bartletti</i> Cushman, 1933	A	
= <i>Criboelphidium bartletti</i> (Cushman, 1933)		
<i>clavatum</i> Cushman, 1930	A	
= <i>E. incertum</i> (Williamson, 1858) var. <i>clavatum</i> Cushman, 1930		
<i>complanatum</i> (d'Orbigny, 1839)	A	
<i>crispum</i> (Linné, 1758)	A M	
= <i>Nautilus crispus</i> Linné, 1758		
<i>cuvilieri</i> Levy, 1966	A M	
<i>decipiens</i> (O.G. Costa, 1856)	M	
<i>depressulum</i> Cushman, 1933	M	
<i>earlandi</i> Cushman, 1936	A	
<i>excavatum</i> (Terquem, 1875)	A	
= <i>E. selseyense</i> (Terquem, 1875)		
= <i>Polystomella excavata</i> Terquem, 1875		
<i>gerthi</i> Van Voorthuysen, 1957	A M	
<i>granosum</i> (d'Orbigny, 1826)	A M	
= <i>Nonionina granosa</i> d'Orbigny, 1826		
<i>gunteri</i> Cole, 1931	A	
<i>iranicum</i> (Yassini & Ghahreman, 1977)	A M	
= <i>Criboelphidium iranicum</i> Yassini & Ghahreman, 1977		
<i>jensenii</i> (Cushman, 1924)	M	
<i>lessoni</i> (d'Orbigny, 1826)		M
<i>lidoense</i> (Cushman, 1936)		A M
= <i>Cribrononion lidoense</i> (Cushman, 1936)		
<i>macellum</i> (Fichtel & Moll, 1798)		A M
= <i>Nautilus macellum</i> Fichtel & Moll, 1798		
<i>magellanicum</i> Heron-Allen & Earland, 1932	A	
<i>maiorensis</i> Colom, 1942		M
<i>margaritaceum</i> (Cushman, 1930)		M
<i>ponticum</i> (Dolgopolskaja & Pauli, 1931)		M
<i>semistriatum</i> (d'Orbigny)		M
<i>striatopunctatum</i> (Fichtel & Moll, 1798)		M
<i>subarcticum</i> Cushman, 1944		A
<i>subnodosum</i> (Münster)		M
<i>translucens</i> Natland, 1938		A M
<i>verruculatum</i> (Brady)		M
<i>waddense</i> Van Voorthuysen, 1975		A
<i>oceanensis</i> (d'Orbigny, 1826)		A
= <i>Polystomella oceanensis</i> d'Orbigny, 1826		
<i>williamsoni</i> Haynes, 1973		A
Haynesina		
<i>depressula</i> (Walker & Jacob, 1798)		A M
<i>germanica</i> (Ehrenberg, 1840)		A M
= <i>Nonion germanica</i> Ehrenberg, 1840		
<i>orbiculare</i> (Brady, 1881)		A
= <i>H. orbicularis</i> (Brady, 1881)		
= <i>Nonionina orbiculare</i> Brady, 1881		
Family Epistominidae		
Eiloehedra		
<i>levicula</i> (Resig, 1958)		A
= <i>Epistominella levicula</i> Resig, 1958		
<i>nipponica</i> Kuwano, 1962		A
Epistominella		
<i>arctica</i> Green, 1960		A
<i>exigua</i> (Brady, 1884)		A M
= <i>Pulvinulina exiguia</i> Brady, 1884		
<i>laevicula</i> Resig, 1958		A
<i>naraensis</i> (Kuwano, 1962)		A
<i>pulchra</i> (Cushman, 1933)		A
<i>rugosa</i> (Phleger & Parker, 1951)		M
<i>vitrea</i> Parker, 1953		A
Hoeglundina		
<i>elegans</i> (d'Orbigny, 1826)		A M
= <i>Rotalina elegans</i> d'Orbigny		
Family Eponididae		
Alabaminella		
<i>weddellensis</i> (Earland, 1936)		A
= <i>Eponides weddelensis</i> Earland, 1936		
Eponides		
<i>concameratus</i> (Williamson, 1858)		M
<i>punctulatus</i> (d'Orbigny, 1839)		M
<i>pusillus</i> Parr, 1950		M
<i>repandus</i> (Fichtel & Moll, 1798)		A
= <i>Nautilus repandus</i> Fichtel & Moll, 1798		
<i>repandus</i> subsp. <i>concameratus</i> (Montagu, 1808)		A
<i>tenera</i> (Brady, 1884)		M
<i>tuberculata</i> (Blackwill & Wright, 1885)		A
<i>turgidus</i> Phleger & Parker, 1951		A
<i>umbonatus</i> (Reuss, 1860)		M
Ioanella		
<i>tumidula</i> (Brady, 1884)		A
= <i>Eponides tumidulus</i> (Brady, 1884)		
= <i>Truncatulina tumidula</i> Brady, 1884		
Neoeponides		
<i>bradyi</i> (Le Calvez, 1974)		
= <i>Eponides bradyi</i> Earland, 1934		
<i>schreibersi</i> (d'Orbigny)		M
Oridorsalis		
<i>tener</i> (Brady, 1884)		A
= <i>O. umbonatus</i> subsp. <i>tenerus</i> (Brady, 1884)		
= <i>O. umbonatus</i> subsp. <i>umbonatus</i> (Reuss, 1851)		
= <i>Rotalina umbonatus</i> Reuss, 1851		
<i>umbonata</i> (Reuss, 1851)		
Planopulvinulina		
<i>dispansa</i> (Brady, 1884)		M
Poroeponides		

<i>lateralis</i> (Terquem, 1878)	A M	<i>Turborotalita</i>
= <i>Eponides lateralis</i> (Terquem, 1878)		<i>humilis</i> (Brady, 1884)
<i>Pseudoeponides</i>	M	<i>quinqueloba</i> Natland, 1938
<i>falsobeccharii</i> Rouvillois, 1974		
Family Fursenkoinidae		Family Globorotaliidae
<i>Sigmavirgulina</i>	A M	<i>Dentigloborotalia</i>
<i>tortuosa</i> (Brady, 1881)		<i>anfracta</i> (Parker, 1967)
= <i>Bolivina tortuosa</i> Brady, 1881		= <i>Globorotalia anfracta</i> Parker, 1967
<i>Virgulina</i>	A	<i>Globigerinata</i>
<i>exilis</i> (Cushman & Ellisor, 1939)	A	<i>glutinata</i> (Egger, 1893)
<i>squamosa</i> d'Orbigny, 1826	A	<i>uvula</i> (Ehrenberg, 1861)
<i>Virgulopsis</i>	A	<i>Neogloboquadrina</i>
<i>pustulata</i> Finlay, 1939		<i>dutertrei</i> (d'Orbigny, 1839)
		<i>incompta</i> Cifelli, 1961
Family Gavelinellidae		Family Hastigerinidae
<i>Anomalinoidea</i>	A M	<i>Hastigerina</i>
<i>minimus</i> (Forster, 1892)		<i>pelagica</i> (d'Orbigny, 1839)
= <i>Truncatulina minima</i> Forster, 1892		
<i>Gyroidinoides</i>	M	Family Homotrema
<i>lamarckiana</i> (d'Orbigny, 1839)		<i>Homotrema</i>
= <i>Gyroidina lamarckiana</i> (d'Orbigny, 1839)		<i>rubra</i> (Lamarck, 1816)
<i>soldanii</i> (d'Orbigny, 1826)	A M	<i>Miniacina</i>
= <i>Gyroidina soldanii</i> (d'Orbigny, 1826)		<i>miniacea</i> Pallas, 1766
= <i>Hansenisca soldanii</i> d'Orbigny, 1826		
<i>umbonatus</i> (Silvestri, 1898)	A M	<i>Rupertina</i>
= <i>Gyroidina umbonata</i> (Silvestri, 1898)		<i>stabilis</i> (Wallich, 1877)
= <i>Gyroidina umbonata</i> (Silvestri, 1898)		= <i>Rupertia stabilis</i> Wallich, 1877
Family Glabratellidae		Family Lagenidae
<i>Conorbella</i>	M	<i>Lagena</i>
<i>erecta</i> (Sidebottom, 1908)	M	<i>acuticosta</i> (Reuss, 1862)
<i>imperatoria</i> (d'Orbigny, 1846)	M	<i>apiopleura</i> Loeblich & Tappan, 1953
<i>patelliformis</i> (Brady, 1884)	M	<i>aspera</i> Reuss, 1861
<i>pulvinata</i> (Brady, 1884)	M	<i>crenata</i> Parker & Jones, 1865
<i>Glabrarella</i>	A	<i>doveyenensis</i> Haynes, 1973
<i>chasteri</i> (Heron-Allen & Earland, 1913) var. <i>bispinosa</i>		<i>gracilis</i> Williamson, 1848
(Heron-Allen & Earlan, 1913)		= <i>L. distoma</i> Parker & Jones, 1864
		<i>hexagona</i> (Williamson)
Family Glandulinidae	M	<i>hibernica</i> Haynes, 1973
<i>Glandulina</i>	M	<i>hispida</i> Reuss, 1863
<i>laevigata</i> (d'Orbigny, 1839)		<i>hispidula</i> Cushman, 1913
<i>rorundata</i> (Reuss, 1860)		<i>interrupta</i> Williamson, 1848
		<i>laevis</i> (Montagu, 1803)
Family Globigerinidae	M	<i>lagenoides</i> (Williamson)
<i>Beella</i>	M	<i>nebulosa</i> (Cushman, 1923)
<i>digitata</i> (Brady, 1879)		<i>pacifica</i> (Sidebottom, 1912)
= <i>Globigerina digitata</i> Brady, 1879		<i>perlucida</i> (Montagu, 1803)
<i>Globigerina</i>	A M	<i>perlucida</i> (Montagu, 1803) var. <i>basispiculata</i>
<i>bulloides</i> d'Orbigny, 1826	M	Mikhalevich, 1968
<i>eggeri</i> Rhumbler, 1900	M	<i>semistriata</i> (Williamson, 1848)
<i>inflata</i> (d'Orbigny)	M	<i>spicata</i> (Cushman & McCulloch, 1950)
<i>Globigerinella</i>	A	<i>staphyllearia</i> (Schwager) var. <i>basispiculata</i>
<i>siphonifera</i> (d'Orbigny, 1839)		Mikhalevich, 1968
= <i>G. aequilateralis</i> (Brady, 1879)		= <i>Fissurina staphyllearia</i> (Schwager, 1866)
= <i>Hastigerina aequilateralis</i> (Brady, 1879)		<i>striata</i> (d'Orbigny, 1839)
<i>Globigerinoides</i>	M	<i>substrata</i> Williamson, 1848
<i>conglobatus</i> (Brady, 1884)	M	<i>sulcata</i> (Walker & Jacob, 1789)
<i>elongatus</i> (d'Orbigny, 1839)	M	
<i>gomitulus</i> (Seguenza)	M	
<i>ruber</i> (d'Orbigny, 1839)	A M	Family Mississippinidae
<i>sacculifer</i> (Brady, 1884)	A M	<i>Stomatorbina</i>
<i>trilobus</i> (Reuss)	M	<i>concentrica</i> (Parker & Jones, 1864)
<i>Globorotalia</i>		= <i>Mississippiina concentrica</i> (Parker & Jones, 1864)
<i>hirsuta</i> (d'Orbigny, 1839)	M	<i>torrei</i> (Cushman & Mermudez, 1949)
<i>inflata</i> d'Orbigny, 1839		
<i>menardii</i> (d'Orbigny, 1826)	M	Family Nodosariidae
<i>patagonica</i> (d'Orbigny, 1839)		<i>Dentalina</i>
<i>scitula</i> (Brady, 1884)	M	<i>advena</i> (Cushman, 1923)
<i>truncatulinoides</i> (d'Orbigny, 1839)		<i>baggi</i> Galloway & Wissler, 1927
<i>Globoturborotalia</i>	M	<i>communis</i> (d'Orbigny, 1826)
<i>rubescens</i> (Hofker, 1956)		<i>consobrina</i> d'Orbigny
<i>tenella</i> (Parker, 1958)		<i>flintii</i> (Cushman, 1923)
<i>Neogloboquadrina</i>	M	= <i>Nodosaria flintii</i> Cushman, 1923
<i>pachyderma</i> (Ehrenberg, 1861)		<i>frobisherensis</i> Loeblich & Tappan, 1953
= <i>Globigerina pachyderma</i> (Ehrenberg, 1861)		<i>intorta</i> (Dervieux, 1894)
<i>Orbulina</i>	M	<i>leguminiformis</i> (Batsch, 1791)
<i>biloba</i> (d'Orbigny, 1839)		<i>pauperata</i> (d'Orbigny, 1846)
<i>universa</i> d'Orbigny, 1839		<i>subarcuata</i> (Montagu, 1803)
		<i>subsoluta</i> Cushman, 1923
		<i>throndheimensis</i> Feyling-Hansen, 1964

<i>Grigelis</i>			
<i>orectus</i> Loeblich & Tappan, 1994	A	<i>iridea</i> Heron-Allen & Earland, 1932	A
= <i>Dentalina guttifera</i> (d'Orbigny, 1846)		<i>opima</i> Cushman, 1947	A M
<i>Laevidentalina</i>		<i>stella</i> Cushman & Moyer, 1930	A
<i>filiformis</i> (d'Orbigny, 1826)	A	<i>turgida</i> (Williamson, 1858)	A M
= <i>Dentalina filiformis</i> (d'Orbigny, 1826)		= <i>Rotalina turgida</i> Williamson, 1858	
<i>flexuosa</i> (Reuss, 1866)	A M	<i>Nonionellina</i>	
<i>inflexa</i> (Reuss, 1866)	M	<i>labradorica</i> (Dawson, 1860)	A
= <i>Dentalina inflexa</i> (Reuss, 1866)		= <i>Nonionina labradorica</i> Dawson, 1860	
<i>Lingulina</i>		<i>Nonionoides</i>	
<i>carinata</i> d'Orbigny, 1826	M	<i>boueanum</i> (d'Orbigny, 1846)	A
<i>seminuda</i> Hantken	M	= <i>Nonionina boueanum</i> d'Orbigny, 1846	
<i>Marginulina</i>		Family Nummulitidae	
<i>silicostata</i> Fornasini	M	<i>Heterostegina</i>	
<i>glabra</i> d'Orbigny, 1826	M	<i>depressa</i> d'Orbigny, 1826	M
<i>hirsuta</i> d'Orbigny, 1826	M	Family Pararotaliidae	
<i>laevigata</i> d'Orbigny, 1826	M	<i>Neorotalia</i>	
<i>Marginulinopsis</i>		<i>calcar</i> (d'Orbigny, 1839)	A
<i>costata</i> (Batsch, 1791)	M	= <i>Calcarina calcar</i> d'Orbigny, 1839	
<i>Nodosaria</i>		Family Patellinidae	
<i>abbreviata</i> Neugeboren	M	<i>Patellina</i>	
<i>calymorpha</i> Reuss, 1865	A	<i>corrugata</i> Williamson, 1858	A M
<i>obliqua</i> (Linné, 1804)	M	<i>nitida</i> Terquem, 1878	M
<i>ovicula</i> d'Orbigny, 1839	M	Family Planorbulinidae	
<i>raphanistrum</i> (Linné, 1804)	M	<i>Gypsina</i>	
<i>raphanus</i> (Linné, 1804)	M	<i>plana</i> (Carter, 1876)	A M
<i>substriata</i> d'Orbigny, 1839	M	<i>Planogypsina</i>	
<i>Procerolagena</i>		<i>acervalis</i> (Brady, 1884)	M
<i>clavata</i> (d'Orbigny, 1846)	A	<i>Planorbulina</i>	
= <i>Lagena clavata</i> (d'Orbigny, 1846)	M	<i>distoma</i> Terquem, 1876	A
<i>gracillima</i> (Seguenza, 1862)	A M	<i>mediterranensis</i> d'Orbigny, 1826	M
= <i>Hyalinonetrion gracilis</i> (Costa, 1856)		<i>variabilis</i> (d'Orbigny, 1826)	A M
= <i>Lagena gracillima</i> (Costa, 1856)		= <i>Cibicidella variabilis</i> (d'Orbigny, 1839)	
<i>Siphonagena</i>		= <i>Truncatulina variabilis</i> d'Orbigny, 1826	
<i>benevestita</i> (Büchner, 1940)	M	Family Planulinidae	
Family Nonionidae		<i>Hyalinea</i>	
<i>Astrononion</i>		<i>balthica</i> (Schröter, 1783)	A M
<i>echolisi</i> Kennett, 1967	A	= <i>Nautilus balthicus</i> Schröter, 1783	
<i>gallowayi</i> Loeblich & Tappan, 1953	A	Family Plectofrondiculariidae	
= <i>A. stelligerum</i> (d'Orbigny, 1839)		<i>Frondicularia</i>	
<i>sidebottomi</i> Cushman & Edwards, 1937	A M	<i>tetragona</i> O.G. Costa	M
<i>Chilostomella</i>		Family Pleurostomellidae	
<i>oolina</i> Schwager, 1878	A M	<i>Pleurostomella</i>	
<i>ovoidea</i> Reuss, 1850	A M	<i>alternans</i> Schwager, 1866	A
<i>Melonis</i>		<i>brevis</i> Schwager, 1886	A
<i>barleeanus</i> (Williamson, 1858)	A M	Family Polymorphinidae	
= <i>M. barleeanum</i> (Williamson, 1858)		<i>Globulina</i>	
= <i>Nonion barleeanum</i> (Williamson, 1858)		<i>amygdalooides</i> (Reuss)	M
= <i>Nonionina barleeanana</i> Williamson, 1858		<i>gibba</i> d'Orbigny, 1826	M
<i>pompilioides</i> (Fichtel & Moll, 1798)	A	<i>inaequalis</i> Reuss, 1850	A
= <i>Nonion pompilioides</i> (Fichtel & Moll, 1798)		<i>myristiformis</i> (Williamson, 1858)	M
<i>zaandami</i> (Van Voorthuysen, 1952)	A M	<i>punctata</i> d'Orbigny, 1846	M
= <i>M. zaandamae</i> (Van Voorthuysen, 1952)		<i>serroria</i> (Reuss)	M
<i>Nonion</i>		<i>tuberculata</i> d'Orbigny	M
<i>asterizans</i> (Fichtel & Moll, 1798)		<i>Guttilina</i>	
= <i>Florilus asterizans</i> (Fichtel & Moll, 1798)		<i>communis</i> (d'Orbigny, 1826)	M
<i>boueanum</i> (d'Orbigny, 1846)	M	= <i>G. problema</i> (d'Orbigny)	
<i>commune</i> (d'Orbigny, 1846)	A M	<i>concava</i> (Williamson)	M
= <i>Florilus communis</i> (d'Orbigny, 1839)		<i>glacialis</i> Cushman & Ozawa	A
= <i>Nonionina communis</i> d'Orbigny, 1846		<i>harrisi</i> Haynes, 1973	A
<i>depressulus</i> (Walker & Jacob, 1798)	A M	<i>lactea</i> (Walker & Jacob)	M
= <i>Nautilus depressulus</i> Walker & Jacob, 1798		<i>oblonga</i> (d'Orbigny)	M
<i>depressum</i> (Walker & Jacob, 1798)	M	<i>regina</i> (Brady, Parker & Jones)	M
<i>granosum</i> (d'Orbigny, 1846)	M	<i>Laryngosigma</i>	
<i>labradoricum</i> (Dawson)	A	<i>hyalascidia</i> Loeblich & Tappan, 1953	A
<i>pacificum</i> Cushman, 1924	A	<i>lactea</i> (Walker & Jacob, 1798)	A
<i>padanum</i> Perconig, 1954	M	<i>williamsoni</i> (Terquem, 1878)	A M
<i>pauperatum</i> (Balkwill & Wright, 1885)	A	= <i>Fissuripolymorpha williamsoni</i> (Terquem, 1878)	
<i>umbilicatum</i> (Walker & Jacob, 1798)	M	<i>Lingulopyrulinoides</i>	
<i>Nonionella</i>		<i>concava</i> (Williamson, 1858)	M
<i>atlantica</i> Cushman, 1936	A M	<i>Polymorphina</i>	
<i>auricula</i> Heron-Allen & Earland, 1930	A	<i>angusta</i> Egger, 1857	A
<i>cordiformis</i> Cushman, 1947	M	<i>hispida</i> Terquem, 1878	M
<i>digitata</i> Norvang, 1945	A	<i>Pseudopolymorpha</i>	
<i>grateloupi</i> (d'Orbigny, 1836)	A		

<i>compressa</i> (d'Orbigny, 1846)	M	<i>arquatensis</i> Papp, 1963	A
<i>novangliae</i> (Cushman, 1923)	A	<i>elongatastriata</i> (Colom, 1952)	M
<i>suboblonga</i> Cushman & Ozawa, 1930	A	<i>nicoli</i> Mathews, 1945	A
<i>Sigmoidella</i>		<i>phlegeri</i> Le Calvez, 1958	AM
<i>elegantissima</i> (Parker & Jones)	M	<i>Trifarina</i>	
Family Reussellidae		<i>angulosa</i> (Williamson, 1858)	AM
<i>Reussella</i>		= <i>Angulogerina angulosa</i> (Williamson, 1858)	
<i>spinulosa</i> (Reuss, 1850)	AM	= <i>Uvigerina angulosa</i> Williamson, 1858	
= <i>Verneuilina spinulosa</i> Reuss, 1850		<i>bradyi</i> Cushman, 1923	A
Family Robertinidae		<i>carinata</i> (Cushman, 1923)	A
<i>Robertina</i>		= <i>Angulogerina carinata</i> Cushman, 1927	
<i>arctica</i> (d'Orbigny, 1839)	A	<i>fluens</i> (Todd, 1948)	A
<i>subcylindrica</i> (Brady, 1881)		<i>fornasinii</i> (Sell, 1948)	A
<i>tasmanica</i> Parr, 1950	M	= <i>Angulogerina fornasinii</i> Sell, 1948	
<i>translucens</i> Cushman & Parker, 1936	AM	<i>pauperata</i> (Heron-Allen & Earland, 1932)	A
= <i>Robertinoides translucens</i> Cushman & Parker, 1936		<i>Uvigerina</i>	
<i>Robertinoides</i>		<i>auberiana</i> d'Orbigny, 1839	AM
<i>bradyi</i> (Cushman & Parker, 1936)	M	<i>auberiana</i> subsp. <i>bellula</i> Bandy	A
= <i>Robertina bradyi</i> (Cushman & Parker, 1936)		<i>bifurcata</i> d'Orbigny, 1839, <i>sensu</i> Lutze, 1986	A
<i>charlottensis</i> (Cushman, 1925)	A	<i>bononiensis</i> Fornasini, 1888, <i>sensu</i> Lutze, 1986	A
<i>normani</i> (Goes, 1894)	A	<i>canariensis</i> d'Orbigny, 1839	AM
= <i>Bulimina normani</i> Goes, 1894		<i>cylindrica</i> subsp. <i>cylindrica</i> (d'Orbigny, 1826), <i>sensu</i>	
<i>pumillum</i> Höglund, 1947	A	Lutze, 1986	A
<i>sueicum</i> Höglund, 1947	A	<i>elongatastriata</i> (Colom, 1952), <i>sensu</i> Lutze, 1986	A
Family Rosalinidae		<i>finisterensis</i> Colom, 1952	A
<i>Neoconorbina</i>		<i>hispidia</i> Schwager, 1866	A
<i>terquemi</i> (Rzehak, 1888)	AM	<i>hollicki</i> Thalmann, 1950, <i>sensu</i> Lutze, 1986	A
= <i>Discorbina terquemi</i> Rzehak, 1888		<i>mediterranea</i> Hofker, 1932, <i>sensu</i> Lutze, 1986	AM
Family Rotaliidae		<i>peregrina</i> Cushman, 1923, <i>sensu</i> Lutze, 1986	A M
<i>Asterorotalia</i>		<i>peregrina</i> subsp. <i>dirupta</i> Todd, 1948	A
<i>gaimardi</i> (d'Orbigny, 1826)		<i>proboscidea</i> Schwager, 1866	AM
= <i>Rotalia gaimardi</i> d'Orbigny, 1826		<i>pygmaea</i> d'Orbigny, 1826, <i>sensu</i> Lutze, 1986	AM
= <i>Rotalinoides gaimardii</i> (d'Orbigny, 1826)		<i>tenuistrata</i> Reuss	M
<i>Parafrondicularia</i>		Family Vaginulinidae	
<i>advena</i> (Cushman)	A	<i>Amphicoryna</i>	
<i>Placopsisina</i>		<i>hispida</i> d'Orbigny	M
<i>confusa</i> Cushman, 1920	A	<i>papillosa</i> (Silverstr, 1872)	M
<i>Remaneica</i>		<i>scalaris</i> (Batsch, 1791)	AM
<i>helgolandica</i> Rhumbler, 1938	A	= <i>Nautilus scalaris</i> Batsch, 1791	
<i>plicata</i> (Terquem, 1876)	A	<i>Astacolus</i>	
= <i>Patellina plicata</i> Terquem, 1876		<i>crepidulus</i> (Fichtel & Moll, 1798)	M
Family Siphogenerinoididae		<i>planulatus</i> Galloway & Wissler, 1927	A
<i>Hopkinsina</i>		<i>Lenticulina</i>	
<i>pacifica</i> Cushman, 1933	AM	<i>atlantica</i> (Barker, 1960)	A
<i>Hopkinsinella</i>		<i>calcar</i> (Linné, 1758)	M
<i>glabra</i> (Millett, 1903)	A	<i>culturata</i> (Montfort, 1808)	M
Family Siphoninidae		<i>curvisepia</i> (Seguenza)	M
<i>Siphonina</i>		<i>echinata</i> (d'Orbigny)	M
<i>georgiana</i> Haynes, 1973	A	<i>gibba</i> (d'Orbigny, 1826)	M
<i>reticulata</i> (Czjzek, 1884)	M	<i>obtusata</i> subsp. <i>subalata</i> Brady, 1884	M
= <i>Discorbis reticulata</i> (Czjzek, 1848)		<i>orbicularis</i> (d'Orbigny, 1826)	M
<i>tenuicarinata</i> Cushman, 1927	M	<i>rotundata</i> (Lamarck, 1804)	M
Family Spirillinidae		<i>Neolenticulina</i>	
<i>Spirillina</i>		<i>peregrina</i> (Schwager, 1866)	M
<i>decorata</i> Brady, 1884	A	= <i>Lenticulina peregrina</i> (Schwager, 1866)	
<i>limbata</i> Brady, 1884	M	<i>Saracenaria</i>	
<i>obconica</i> Brady, 1879	A	<i>italica</i> (Defrance, 1824)	AM
<i>perforata</i> (Schultze, 1854)	A	<i>Vaginulina</i>	
<i>vivipara</i> Ehrenberg, 1843	AM	<i>costata</i> (Cornuel)	M
Family Trichohyalidae		<i>legumen</i> (Linné, 1804)	M
<i>Aubignya</i>		<i>margaritifera</i> (Batsch, 1791)	M
<i>hamblensis</i> Murray, Whittaker & Alve, 2000		<i>striata</i> d'Orbigny, 1839	M
Family Turrilinidae		<i>vertebralis</i> Parr, 1932	M
<i>Buliminella</i>		= <i>Nodosaria vertebralis</i> d'Orbigny, 1839	
<i>borealis</i> Haynes, 1973	A	Kathablepharids	
<i>elegantissima</i> (d'Orbigny, 1839) var. <i>hensonii</i> Lagoe, 1977	A	Compiled by Sue Brandt	
<i>elegantissima</i> (d'Orbigny, 1839)	AM	These species were previously thought to be cryptophytes. A comprehensive list (to 1995).	
= <i>Bulimina elegantissima</i> d'Orbigny, 1839		References	
<i>mississippiensis</i> Seiglie, 1969	A	Clay, B.L. & Krugens, P. 1999 Description and ultrastructure of <i>Kathablepharis tenuis</i> sp. Nov. and <i>K. obesa</i> sp. Nov. -	
Family Uvigerinidae			
<i>Rectuvigerina</i>			

- Two new freshwater kathablepharids from Colorado and Wyoming. *Europ. J. Protistol.*, 35: 435-447
- Tong, S.M. 1997 Heterotrophic flagellates and other protists from Southampton Water, U.K. *Ophelia* 47: 71-131
- Vørs, N. 1992 Ultrastructure and autecology of the marine heterotrophic flagellate *Leucocryptos marina* (Braarud) Butcher 1967 (Katablepharidaceae/Kathablepharidae), with a discussion of the genera *Leucocryptos* and *Kathablepharis*. *Kathablepharis*. *Europ. J. Protistol.*, 28: 369-389

KATHABLEPHARIDS

- Leucocryptos*
marina (Braarud, 1935) Butcher, 1967
= *Bodo marina* Braarud, 1935
= *Chilomonas marina* (Braarud, 1935) Halldal, 1953
- Kathablepharis*
remigera (Vørs, 1992) Clay & Krugens, 1999
= *Leucocryptos remigera* Vørs, 1992

Haptophytes

Compiled by Sue Brandt

A preliminary list based on a few surveys.

References

- Espeland, G. & Thronsdene, J. 1986 Flagellates from Kilsfjorden, Southern Norway, with description of two new species of Choanoflagellida. *Sarsia* 71: 209-226
- Ikävalko, J & Thomsen, H.A. 1997 The Baltic Sea Ice Biota (March 1994): a study of the protistan community. *Europ. J. Protistol.*, 33: 229-243
- Leadbeater, B.S.C 1972 Identification, by means of electron microscopy, of flagellate nanoplankton from the coast of Norway. *Sarsia* 49: 107-124
- Thomsen, H.A., Bjørn, P.P., Højlund, L., Olesen, J. & Pedersen, J.B. 1995 *Ericiolus* gen. Nov. (Prymnesiophyceae), a new cocolithophorid genus from polar and temperate regions. *Eur. J. Phycol.*, 30: 29-34

HAPTOPHYTES

- Acanthoica*
acanthifera Lohm, 1902
- Anthosphaera*
quadricornu (Schiller) Halldal & Markali, 1955
= *Syracosphaera quadricornu* Schiller, 1925
- Balaniger*
balticus Thomsen & Oates, 1978
- Chrysochromulina*
acantha Leadbeater & Manton, 1971
alifera Parke & Manton, 1956
apheles Moestrup & Thomsen
bergenensis Leadbeater, 1972
brevifilum Parke & Manton, 1955
chiton Parke & Manton, 1958
cymbium Leadbeater & Manton, 1969
ephippium Parke & Manton, 1956
ericina Parke & Manton, 1956
fragilis Leadbeater, 1972
herdagensis Leadbeater, 1972
hirta Manton, 1978
kappa Parke & Manton, 1955
macra Manton, 1972
mantoniae Leadbeater, 1972
megacylindra Leadbeater, 1972
microcylindra Leadbeater, 1972
minor Parke & Manton, 1955
parkeae Green & Leadbeater, 1972
polylepis Manton & Parke, 1962
pringsheimii Parke & Manton, 1962
spinifera (Fournier) Pienaar & Norris, 1979
= *Chrysocampanula spinifera* Fournier, 1971
strobilus Parke & Manton, 1959
- Dicrateria*
inornata Parke, 1949

- Emiliania*
huxleyi (Lohmann) Hay & Mohler, 1967
= *Pontosphaera huxleyi* Lohmann, 1902
- Ericiolus*
spiculiger Thomsen, 1995
- Hymenomonas*
carterae (Braarud & Fagerland) Manton & Peterfi, 1969
- Ophiaster*
hydroideus (Lohmann) Lohmann, 1913
= *Meringosphaera hydroidea* Lohmann, 1903
- Pappomonas*
flabellifera Manton & Oates, 1975
virgulosa Manton & Sutherland, 1975
- Phaeocystis*
pouchetii (Hariot) Legerheim, 1896
- Trigonaspis*
minutissima Thomsen, 1980
- Turrisphaera*
borealis Manton, Sutherland & Oates, 1976

Prasinophytes

Compiled by Sue Brandt

A preliminary list based on a survey of plankton in Danish Waters.

References

- Thomsen, H.A. (ed.) 1992 *Plankton i de indre danske farvande* Miljøministeriet, Miljøstyrelsen, Copenhagen.

PRASINOPHYTES

- Cymbomonas*
tetramitiformis Schiller
- Dolichomastix*
lepidota Manton
nummiliifera Manton
- Halosphaera*
viridis Schmitz
- Mamiella*
gilva (Parke & Rayns) Moestrup
= *Nephroselmis gilva* Parke & Rayns
- Mantoniella*
squamata (Manton & Parke) Desikachary
= *Micromonas squamata* Manton & Parke
- Micromonas*
pusilla (Butcher) Manton & Parke
- Nephroselmis*
minuta (N. Carter) Butcher
= *Heteromastix minuta* Butcher
- pyriformis* (Butcher) Ettl
= *Heteromastix longifilis* Rayns
- rotunda* (N. Carter) Fott
= *Heteromastix rotunda* Manton
- Pseudoscourfieldia*
marina (Thronsdene) Manton
= *Scourfieldia marina* Thronsdene
- Pyramimonas*
circulanae Pennick
cordata McFadden, Hill & Wetherbee
- disomata* McFadden, Hill & Wetherbee
- grossii* Parke
longicaudavan Meel
= *Pyramimonas ostendensis* van Meel
- mantoniae* Moestrup & Hill
- nansenii* Braarud
- norrisii* Sym & Pienaar
- octopus* Moestrup & Aa. Kristiansen
- orientalis* McFadden, Hill & Wetherbee
- propulsa* Moestrup & Hill
- virginica* Pennick
- Tetraselmis*
contracta (N. Carter) Butcher
= *Platymonas contracta* N. Carter

Stramenopiles

Bicosoecids

Compiled by Sue Brandt

Comprehensive list produced from recent literature.

References

- Griessmann, K. 1913 Über marine Flagellaten. *Archiv. Protistenk.* 32: 1-78
 Hallfors, G. & Niemi, Å. 1984 A preliminary checklist of the phytoplankton of the Baltic Sea. *Acta Botanica Fennica*, 128: 1-26
 Ikävalko, J & Thomsen, H.A. 1997 The Baltic Sea Ice Biota (March 1994): a study of the protistan community. *Europ. J. Protistol.*, 33: 229-243
 Järnfelt, H 1956 Zur Limnologie einiger Gewässer Finlands *XVI Ann. Zool. Soc. Zool.-Bot. Fenn "Vanamo"*, 17: 1-201
 Kent, W.S. 1880 *A manual of the Infusoria*. (1880-1882) Bogue, London.
 Larsen, J. & Patterson, D.J. 1990 Some flagellates (Protista) from tropical marine sediments. *Journal of Natural History*, 24: 801-937.
 Moestrup, Ø. & Thomsen, H.A. 1976 Fine structural studies on the flagellate genus *Bicoeca* l. *Bicoeca maris* with apicular emphasis on the flagellar apparatus *Protistologica*, 12: 101-120
 Preisig, H.R., Vørs N. & Hälfors, G. 1991 Diversity of heterotrophic heterokont flagellates In: Patterson, D.J. & Larsen, J. (eds): *The biology of free-living heterotrophic flagellates*. Systematics Association Special Volume, 45, pp361-99, Clarendon Press, Oxford
 Tong, S.M. 1997b Heterotrophic flagellates and other protists from Southampton Water, U.K. *Ophelia* 47: 71-131.
 Valkanov, A. 1970 Beitrag zur Kenntnis der Protozoen des Schwarzen Meeres. *Zool. Anz.*, 184: 241-290
 Vørs, N. 1992 Heterotrophic amoebae, flagellates and heliozoa from the Tvarminne area, Gulf of Finland, in 1988-1990 *Ophelia* 36: 1-109
 Vørs, N. 1992 Ultrastructure and autecology of the marine heterotrophic flagellate *Leucocryptos marina* (Braarud) Butcher 1967 (Katablepharidae/Cathablepharidae), with a discussion of the genera *Leucocryptos* and *Katablepharis/Kathablepharis*. *Europ. J. Protistol.*, 28: 369-389

Bicosoecids

Bicosoeca

- ainikkaiae* Järnfelt, 1956
 = *B. tubiformis* Skuja, 1956
eiphytica Hilliard, 1971
 = *B. pulchra* Hilliard, 1971
gracilipes James-Clark, 1867
lacustris James-Clark, 1867
maris Picken, 1941
 = *B. pocillum* sensu Griessmann, 1913
mediterranea Pavillard, 1916
mignotii Moestrup, Thomsen & Hibberd in Vørs, 1992
planctoronica Kissilew, 1931
 = *B. annulata* (Lackey) Bourely, 1951
 = *B. multiannulata* Skuja, 1948
pontica Valkanov, 1970
Caecitellus
parvulus (Griessmann, 1913) Patterson, Nygaard, Steinberg & Turley, 1993
 = *Bodo parvulus* Griessmann, 1913
Cafeteria
marsupialis Larsen & Patterson, 1990
minuta (Griessmann, 1913) Larsen & Patterson, 1990
 = *Pseudobodo minuta* Griessmann, 1913
roenbergensis Fenchel & Patterson, 1988
Pseudobodo
minima Griessmann, 1913
tremulans Griessmann, 1913

Labyrinthulids

Compiled by Michael Dick and Sue Brandt

Comprehensive list of European species derived from a global checklist.

References

- Chadefaud, M. 1956 *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences, Paris*, Série D, 243:1796.
 Cienkowsky, L. 1867 *Archiv für Mikroskopische Anatomie und Entwickelungsmechanik* 3: 275.
 Dangeard, P.-A. 1932 *Le Botaniste* 24: 223.
 Hollande, A. & Enjumet, M. 1955 *Annales des Sciences Naturelles, Zoologie et Biologie Animale*, Série 11, 17: 357.
 Karling, J.S. 1944 *American Journal of Botany* 31: 50.
 Schmoller, H. 1960 *Archiv für Mikrobiologie* 36: 371.
 Schmoller, H. 1966 *Archiv für Protistenkunde* 109:237.
 Valkanov, A. 1972 *Progress in Protozoology*. Abstracts 3rd International Congress on Protozoology: 373.
 Zopf, W. 1892 *Beiträge zur Physiologie und Morphologie* 2: 61.

Labyrinthulids

Labyrinthomyxa

- pohlia* H. Schmoller, 1966
Labyrinthula
algeriensis A. Holland & M. Enjumet, 1955
chattonii P.-A. Dangeard, 1932
 = *Phagomyxa chattonii* (P.-A. Dangeard, 1932) J.S. Karling, 1944
cienkowskii W. Zopf, 1892
 = *Labyrinthorhiza cienkowskii* (W. Zopf, 1892) M. Chadefaud, 1956
coenocystis H. Schmoller, 1960
macrocystis L. Cienkowski, 1867
 = *Labyrinthula macrocystis* L. Cienkowski, 1867 var. *atlantica* H.S. Vishniac & S.W. Watson 1953
magnifica (A. Valkanov, 1969) L.S. Olive, 1975
 = *Labyrinthodictyon magnificum* A. Valkanov, 1969
roscoffensis M. Chadefaud, 1956
valkonovii J.S. Karling, 1944
 = *Labyrinthula zoppii* A. Valkanov, 1929
vitellina L. Cienkowski, 1867
 = *Chlamydomyx labyrinthuloides* W. Archer, 1875
 = *Labyrinthula macrocystis* L. Cienkowski, 1867 var. *vitellina* E.L. Young 1943
 = *Labyrinthula vitellina* L. Cienkowski, 1867 var. *pacifica* S.W. Watson 1951
 = *Pseudoplasmodium aurantiacum* H. Molisch, 1926

Thaustrochytrids

Compiled by Michael Dick and Sue Brandt

Comprehensive list of European species derived from a global checklist.

References

- Artemchuk, N. Ya. 1972 *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 13: 233.
 Gaertner, A. 1967 *Helgoländer wissenschaftliche Meeresuntersuchungen* 15: 186.
 Gaertner, A. 1972 *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 19: 69.
 Gaertner, A. 1977 *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 16: 150.
 Raghu Kumar, S. 1977 *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 16: 160.
 Raghu Kumar, S. 1980 *Transactions of the British Mycological Society* 74: 612.
 Raghu Kumar, S. 1988 *Transactions of the British Mycological Society* 90: 627.
 Schlotz, E. 1958 *Archiv für Mikrobiologie* 29: 361.
 Schneider, J. 1967 *Kieler Meeresforschung* 23: 16.
 Ulken, A. 1965 *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 9: 290, 291.

Thraustochytrids

Elina

- marisalba* N. Ya. Artemchuk, 1972
- sinorifica* N. Ya. Artemchuk, 1972

Schizochytrium

- minutum* A. Gaertner, 1972
- octosporum* S. Raghu Kumar, 1988
- Thraustochytrium*
- aggregatum* A. Ulken, 1965
- arudimentale* N. Ya. Artemchuk, 1972
- benthicola* S. Raghu Kumar, 1980
- kinnei* A. Gaertner, 1967
- pachydermum* E. Scholz, 1958
- striatum* J. Schneider, 1967

Ulkenia

- minuta* S. Raghu Kumar, 1977
- profunda* A. Gaertner, 1977
- radiata* A. Gaertner, 1977
- sarkariana* A. Gaertner, 1977
- visurgensis* (A. Ulken, 1965) A. Gaertner, 1977
- = *Thraustochytrium visurgense* A. Ulken, 1965

Stramenopiles incertae sedis

Compiled by Sue Brandt

Stramenopile taxa which do not fit into any known taxa above the level of genus.

References

- Tong, S.M. 1997 Heterotrophic flagellates and other protists from Southampton Water, U.K. *Ophelia* 47: 71-131.
- Vørs, N. 1992 Heterotrophic amoebae, flagellates and heliozoa from the Tvarminne area, Gulf of Finland, in 1988-1990 *Ophelia* 36: 1-109

Stramenopiles incertae sedis

Bordnamonas

- tropicana* Larsen & Patterson, 1990

Diplophysrys

- archeri* Barker, 1868
- = *Diplophysrys marina* Dykstra & Porter, 1984

Developayella

- elegans* Tong, 1996

Pendulomonas

- adriperis* Tong, 1997

Thaumatomonads

Compiled by Sue Brandt

Comprehensive list produced from recent literature.

References

- Beech, P.L. & Moestrup, Ø 1986 Light and electron microscopical observations on the heterotrophic protist *Thaumatomastix salina* comb. Nov. (syn. *Chrysosphaerella salina*) and its allies *Nordic Journal of Botany*, 6: 865-877
- Larsen, J. & Patterson, D.J. 1990 Some flagellates (Protista) from tropical marine sediments. *Journal of Natural History*, 24: 801-937.
- Larsen, J. 1985 Algal studies of the Danish Wadden Sea II. A taxonomic study of psammobious dinoflagellates *Opera Botanica*, 79: 14-37
- Thomsen, H.A., Hälfors, G., Hälfors, S. & Ikävalko, J. 1993 New observations on the heterotrophic protist genus *Thaumatomastix* (Thaumatomastigaceae, Protista incertae sedis), with particular emphasis on material from the Baltic Sea *Ann. Bot. Fennici* 30: 87-108
- Thomsen, H.A. (ed.) 1992 *Plankton i de indre danske farvande* Miljøministeriet, Miljøstyrelsen, Copenhagen.
- Tong, S.M. 1997 Heterotrophic flagellates and other protists from Southampton Water, U.K. *Ophelia* 47: 71-131.

THAUMATOMONADS

Protaspis

gemmifera

- Larsen & Patterson, 1990*
- glans* Skuja, 1939
- major* Skuja, 1939
- metarhiza* Skuja, 1939
- obliqua* Larsen & Patterson, 1990
- simplex* Vørs, 1992
- verrucosa* Larsen & Patterson, 1990

Thaumatomastix

- bipartita* Beech & Moestrup, 1986
- dybsoeana* Thomsen, Hälfors, Hälfors & Ikävalko, 1993
- formosa* Thomsen, Hälfors, Hälfors & Ikävalko, 1993
- patelliformis* (Takahashi & Hara) Beech & Moestrup, 1986
- = *Chrysosphaerella patelliformis* Takahashi & Hara, 1984
- = *Spinifermonas patelliformis* (Takahashi & Hara) Nicholls, 1984
- sagittifera* (Conrad) Beech & Moestrup, 1986
- = *Reckertia sagittifera* Conrad, 1920
- salina* (Birch-Andersen) Beech & Moestrup, 1986
- = *Chrysosphaerella salina* (Birch-Andersen) Birch-Andersen, 1973
- = *Spinifermonas salina* (Birch-Andersen) Nicholls, 1984
- spinososa* Thomsen, Hälfors, Hälfors & Ikävalko, 1993
- thomsoni* Tong, 1997
- triangulata* (Balonov) Beech & Moestrup, 1986
- = *Chrysosphaerella triangulata* Balonov, 1980
- tripus* (Takahashi & Hara) Beech & Moestrup, 1986
- = *Chrysosphaerella tripus* Takahashi & Hara, 1984
- = *Spinifermonas tripus* (Takahashi & Hara) Nicholls, 1984

Xenophyophora

Compiled by Ole Tendal and Jacob van der Land

The European list derived from URMO which is based on Tendal (1995). Ole Tendal checked the ERMS list in 1999.

References

- Tendal, O.S. 1995. Synoptic checklist and bibliography of the Xenophyophora (Protista), with a zoogeographical survey of the group. *Galathea Report* 17 1995-1996: 79-101, illustr.

XENOPHYOPHORA

Family Cerelasmidae

- Cerelasma*
- massa* Tendal, 1972

Family Psammettidae

- Homogammina*
- crassa* Gooday, 1991
- lamina* Gooday & Tendal, 1988
- maculosa* Gooday & Tendal, 1988

Family Psamminidae

- Galatheammina*
- calcarea* (Haeckel, 1889)
- discoveryi* Gooday & Tendal, 1988
- erecta* Gooday, 1991
- irregularis* Gooday, 1991
- microconcha* Gooday & Tendal, 1988
- Psammina*
- delicata* Gooday & Tendal, 1988
- sabulosa* Gooday & Tendal, 1988
- fusca* Gooday & Tendal, 1988

Family Syringamminidae

- Aschemonella*
- catenata* (Norman, 1876)

- ramuliformis* Brady, 1884
scabra Brady, 1879
Syringammina
fragillissima Brady, 1883
reticulata Gooday, 1996

Protista incertae sedis

Compiled by Sue Brandt

The species in this list are from genera where the nearest relative is not known. The individual genera are sometimes placed within their own families, orders, classes etc. This list is comprehensive for heterotrophic species, and produced from recent literature.

References

- Drebes, G., Kühn, S.F., Gmelch, A. & Schnepf, E. 1996 *Cryothecononas aestivalis* sp. Nov., a colourless nanoflagellate feeding on the marine centric diatom *Guinardia delicatula* (Cleve) Hasle *Helgoländer Meeresuntersuchungen*, 50: 497-515
- Entz, G. 1904 Fauna der Kontinentalen Kochsalzwässer Mathematische und Naturwissenschaftliche Berichte aus Ungarn, 19, 89-124.
- Fenchel, T., Bernard, C., Esteban, G., Finlay, B.J., Hansen, P.J. & Iversen N. 1995 Microbial diversity and activity in a Danish fjord with anoxic deep water *Ophelia* 43: 45-100
- Griessmann, K. 1913 Über marine Flagellaten. *Archiv. Protistenk.* 32: 1-78
- Kent, W.S. 1880 *A manual of the Infusoria*. (1880-1882) Bogue, London.
- Kühn, S.F., Drebes, G. & Schnepf, E. 1996 Five new species of the nanoflagellate *Pirsonia* in the German Bight, North Sea, feeding on planktonic diatoms. *Helgoländer Meeresuntersuchungen*, 50: 205-222
- Patterson, D.J. & Zöllfel, M. 1991 Heterotrophic flagellates of uncertain taxonomic position In: Patterson, D.J. & Larsen, J. (eds): *The biology of free-living heterotrophic flagellates*. Systematics Association Special Volume, 45, pp 427-475, Clarendon Press, Oxford
- Patterson, D.J., Nygaard, K., Steinberg, G. & Turley, C.M. 1993 Heterotrophic flagellates and other protists associated with oceanic detritus throughout the water column in the mid north Atlantic *J. mar. biol. Ass. U.K.* 73: 67-95
- Pavillard, M.J. 1916 Flagellés nouveaux, épiphytes des Diatomées pélagiques. *C. r. hebdo. Séanc. Acad. Sci., Paris*, 163: 65-68
- Ruinen, J. 1938 Notizen über Salzflagellaten II Über die Verbereitung der Salzflagellaten *Archiv für Protistenkunde*, 90, 210-258.
- Thomsen, H.A. (ed.) 1992 *Plankton i de indre danske farvande* Miljøministeriet, Miljøstyrelsen, Copenhagen.
- Thomsen, H.A., Buck, K.R., Bolt, P.A. & Garrison, D.L. 1991 Fine structure and biology of *Cryothecononas* gen. Nov. (*Protista incertae sedis*) from the ice biota *Can. J. Zool.*, 69: 1048-1070
- Tong, S.M. 1997 Heterotrophic flagellates and other protists from Southampton Water, U.K. *Ophelia* 47: 71-131.
- Valkanov, A. 1970 Beitrag zur Kenntnis der Protozoen des Schwarzen Meeres. *Zool. Anz.*, 184: 241-290
- Vørs, N. 1992 Heterotrophic amoebae, flagellates and heliozoa from the Tvärminne area, Gulf of Finland, in 1988-1990 *Ophelia* 36: 1-109
- Vørs, N. 1992 Ultrastructure and autecology of the marine heterotrophic flagellate *Leucocryptos marina* (Braarud) Butcher 1967 (Katablepharidaceae/Kathablepharidae), with a discussion of the genera *Leucocryptos* and *Katablepharis*/Kathablepharis. *Europ. J. Protistol.*, 28: 369-389
- tachyploon* Sandon, 1924
- Amphimonas*
cuneatus Namyslovski, 1913
divaricans Kent, 1880
- Ancyromonas*
sigmoides Kent, 1880
= *Bodo cephaloporus* Larsen & Patterson, 1990
- Bordnamonas*
tropicana Larsen & Patterson, 1990
- Chloraster*
agilis Kent, 1880
- Cryothecononas*
inermis Thomsen, Buck, Bolt & Garrison, 1991
scybalophora Thomsen, Buck, Bolt & Garrison, 1991
vesiculata Thomsen, Buck, Bolt & Garrison, 1991
aestivalis Drebes, Kühn & Schnepf, 1996
- Discocelis*
saleuta Vørs, 1988
- Ebria*
tripartita (Shumann) Lemmermann, 1899
- Jakoba*
libera (Ruinen, 1938) Patterson, 1990
= *Cryptobia libera* Ruinen, 1938
- Kiioksia*
ystava Vørs, 1992
- Luffisphaera*
hamatus Tong, 1997
turriformis Vørs, 1992
cucumiformis Belcher & Swale, 1975
myosurus Belcher & Swale, 1975
- Metopion*
fluens Larsen & Patterson, 1990
- Metromonas*
simplex (Griessmann, 1913) Larsen & Patterson, 1990
- Ministeria*
marisola Patterson, Nygaard, Steinberg & Turley, 1993
vibrans Tong, 1997
- Phylloclitus*
granulatus Larsen & Patterson, 1990
- Pirsonia*
guinardiae Schnepf, Drebes & Elbrächter, 1990
verrucosa Kühn, 1996
formosa Kühn, 1996
diadema Kühn, 1996
eucampiae Kühn, 1996
mucosa Kühn, 1996
- Postgaardi*
mariagerensis Fenchel, Bernard, Esteban, Finlay, Hansen & Iversen, 1995
- Quadricilia*
rotundata (Skuja, 1948) Vørs, 1992
= *Collodictyon sphaericum* Norris, 1964
= *Tetramitus rotundata* Skuja, 1948
- Rhizomonas*
setigera (Pavillard, 1916) Patterson, Nygaard, Steinberg & Turley, 1993
= *Solenicola setigera* Pavillard, 1916
- Rigidomastix*
devoratum Tong, 1997
- Telonema*
antarctica Thomsen in Vørs, 1992
subtile Griessmann, 1913
= *T. subtilis* Griessmann, 1913
- Tetracilia*
paradoxa Valkanov, 1970
- Trichomastix*
salina Entz, 1904
= *T. salinus* Entz, 1904
- Triflagellum*
salinum Namyslovski, 1913
- Trimastix*
marina Kent, 1880

PROTISTA INCERTAE SEDIS

- Abollifer*
probabens Vørs, 1992
Allantion

MYXOZOA

Compiled by Egil Karlsbakk

Myxozoa are multicellular but microscopic parasites of vertebrates (mainly fishes, but some amphibian and chelonian hosts are known) and invertebrates (mainly annelids, and mainly oligochaeta). Known life cycles describe sporogony in a vertebrate host producing myxospores. These infect an oligochaete or polychaete in which a different type of sporogony produces actinospores. A deviating myxozoan group also produce spores in freshwater bryozoans.

The list is based on an unpublished list, compiled by E. Karlsbakk, based on literature search and the Aquatic Sciences and Fisheries Abstracts. Almost exclusively original species descriptions have been used as the basis for species information. The preliminary list was then checked against other lists or sources: Gurley (1894), Thélohan (1895), Labbé (1899), Auerbach (1910), Kudo (1920), Shulman (1962), Shulman (1966), Naidenova *et al.* (1975), Donetz and Shulman (1984), Lom and Dyková (1992) and others. A total of 230 species are listed as valid, in addition, some important synonyms are included. It is likely that the actual number of species is much higher, since many fish species have never been examined. Also, many old myxozoan species described from several unrelated hosts will likely turn out to be complexes of sibling species (e.g. *Myxidium oviiforme*).

The classification used is based on Lom and Dyková (1992). However, it differs in that; (a) the recent genus *Polysporoplasma* Sitjá-Bobadilla & Alvarez-Pellitero, 1995 is placed in family Sphaerosporidae and seems closely related to *Sphaerospora* Thélohan, 1892, (b) Genus *Paramyxoproteus* Wierzbicka, 1986 is considered as a synonym of *Schulmania* Kovaleva, Zubchenko & Krasin, 1983, (c) *P. reinhardtii* Wierzbicka, 1986 is a synonym of *S. quadriolobata* Kovaleva, Zubchenko & Krasin, 1983. The valid binomen for *Sphaerospora caudata* Parisi, 1910 is considered to be *Hosferellus caudatus* (Parisi, 1910) (large polysporic plasmodia, spore shape, posterior filaments).

Hosts are listed as valid genus or family of type host, and then other families or simply "other fishes" when the hosts are many and unrelated. When type hosts were not designated in the original description, the first listed is considered the type-host. Only the geographic distribution in the ERMS area is considered. However, several of the myxozoan species in the area are recorded also from, for instance, Australia and New Zealand.

References

- Auerbach, M. 1910. *Die Cnidosporidien (Myxosporidien, Actinomyxidien, Microsporidien). Eine Monographische Studie.* Leipzig: Werner Klinkhardt. 261 pp.
- Donetz, E.S. & Shulman, S.S. 1984. 1984. Phylum cnidosporidians - Cnidosporidia Doflein, 1901, emend. Shulman & Podlipaev, 1980. In: *Keys to the parasites of freshwater fishes in the USSR fauna. Vol. 1. Parasitic protozoa*, pp. 88-251. S.S. Shulman, editor. Leningrad: Nauka. (in Russian)
- Gurley, R.R. 1894. Myxosporidia, or psorosperms of fishes, and the epidemics produced by them. Report to the U.S. Commissioner of Fish and Fisheries for 1892. 1892: 65-304.
- Kudo, R. 1920. Studies on Myxosporidia. A synopsis of genera and species of Myxosporidia. *Illinois Biological Monographs* 5: 239-503.
- Labbé, A. 1899. *Sporozoa. Das Tierreich. Vol. 5.* Berlin: Deutsche Zool. Gesellsch. 180 pp.
- Lom, J. & Dyková, I. 1992. *Protozoan parasites of fishes. Developments in Aquaculture and Fisheries Science*, Vol. 26. Amsterdam: Elsevier. 315 pp.
- Naidenova, N.N., S.S. Shulman and Z.S. Donetz. 1975. Class cnidosporidians - Cnidosporidia Doflein, 1901, emend. Shulman, 1967. In: *Identification of the parasites of vertebrates in the Black- and Azov Seas*, p. 20-50. A.V. Gaevskaya *et al.*, editors. : Kiev: Akademiya Nauk Ukrainskoj SSR, "Naukova Dumka". (in Russian)
- Shulman, S.S. 1962. Order Myxosporidia Bütschli, 1881. Slime sporozoans. In: *Key to parasites of freshwater fish of the U.S.S.R.*, p. 48-130. I.E. Bykhovskaya-Pavlovskaya *et al.*, editors. Moscow-Leningrad: Izdatel'stvo Akademii Nauk SSSR. (in Russian)
- Shulman, S.S. 1966. *Myxosporidia of the USSR*. Moscow: Akademiya Nauk SSSR, Zoologicheskii Institut. Nauka. 504 pp. (in Russian)
- Thélohan, P. 1895. Recherches sur les Myxosporidies. *Bulletin scientifique de la France & de la Belgique* 5: 100-394.
- Bibliography**
1. Auerbach, M. 1907. Weitere Mitteilungen über *Myxobolus aeglefini* Auerb. *Zool. Anz.* 31: 115-119.
 2. Auerbach, M. 1909. Bemerkungen über Myxosporidien. *Zool. Anz.* 34: 65-82.
 3. Auerbach, M. 1909. Biologische und morphologische Bemerkungen über Myxosporidien. *Zool. Anz.* 35: 57-63.
 4. Auerbach, M. 1910. Cnidosporidenstudien. *Zool. Anz.* 35: 767-777.
 5. Averintzev, S. 1908. Studien über parasitische Protozoen. I-VII. *Trav. Soc. Imp. Nat. St.-Petersburg* 28: 1-139. (in Russian)
 6. Bahri, S. and A. Marques. 1996. Myxosporean parasites of the genus *Myxobolus* from *Mugil cephalus* in Ichkeul lagoon, Tunisia: description of two new species. *Dis. Aquat. Org.* 27: 115-122.
 7. Bakay, Y.I. and M.A. Grudnev. 1998. New species of Myxosporidia (Cnidospora: Myxosporea) in redfishes of the North Atlantic. *Parazitologiya* 32: 372-375. (in Russian)
 8. Basikalova, A.Y. 1932. Contribution to the parasitology of Murman fishes. *Sbornik nauchno-promyshlonykh rabot na Murmane.* Snabtekhizdat, Moscow. p.136-153. (in Russian)
 9. Canning, E.U., A. Curry, C.L. Anderson and B. Okamura. 1999. Ultrastructure of *Myxidium trachinorum* sp. nov. from the gallbladder of the lesser weever fish *Echiichthys vipera*. *Parasitol. Res.* 85: 910-919.
 10. Davies, A.J. 1985. *Zschokkella russelli* Tripathi (Myxozoa: Myxosporea) from five-bearded rockling, *Ciliata mustela* L., (Teleostei: Gadidae) in Wales. *J. Fish Dis.* 8: 299-308.
 11. Diamant, A. and I. Paperna. 1992. *Zschokkella icterica* sp. nov. (Myxozoa, Myxosporea), a pathogen of wild rabbitfish *Siganus luridus* (Ruppell, 1829) from the Red Sea. *Europ. J. Protistol.* 28: 71-78.
 12. Diamant, A., J. Lom and I. Dyková. 1994. *Myxidium leei* n. sp., a pathogenic myxosporean of cultured sea bream *Sparus aurata*. *Dis. Aquat. Org.* 20: 137-141.
 13. Doflein, F. 1898. Studien zur Naturgeschichte der Protozoen III. *Zool. Jahrb. Abt. Anatomie* 11: 281-350.
 14. Dunkerly, J.S. 1921. Fish Myxosporidia from Plymouth. *Parasitology* 12: 328-333.
 15. Gaevskaya, A.V. and A.A. Kovaleva 1984. Materials on the myxosporidian fauna of the Celtic Sea fishes. *Vestn. Zool.* 1984: 3-7. (in Russian)
 16. Gaevskaya, A.V. and A.A. Kovaleva. 1979. New and rarely occurring Myxosporidia species from the Celtic Sea fishes. *Parazitologiya* 13: 159-165. (in Russian)
 17. Gaevskaya, A.V. and A.A. Kovaleva. 1980. New species of giant myxosporidians from the Atlantic bathypelagic fishes. *Zool. Zh.* 59: 942-944. (in Russian)
 18. Gaevskaya, A.V. and A.A. Kovaleva. 1984. Addition to the Myxosporidia fauna (Protozoa: Myxosporidia) of

- fishes of the Northeast Atlantic. *Gidrobiol. Zh.* 1984. 20(3): 49-53 (in Russian) (Transl. in: *Hydrobiol. J.* 20 (4): 49-53.)
19. Gaevskaya, A.V., A.A. Kovaleva and B.A. Umnova. 1980. *Davisia amoena* sp. n. (Myxosporidia, Sinuolineidae), a parasite of fishes of the family pleuronectidae from north-western Atlantic. *Parazitologiya* 14: 276-279. (in Russian)
 20. Georgévitch, J. 1916. Note sur les myxosporidies des poissons de la baie de Villefranche & de Monaco. *Bull. Inst. Oceanogr.* (322): 1-7.
 21. Georgévitch, J. 1916. Note sur les Myxosporidies recueillies à Roscoff. *Bull. Soc. Zool. France.* 41: 86-95.
 22. Georgévitch, J. 1917. Esquisses protistologiques. *Bull. Inst. Oceanogr.* (328): 1-12.
 23. Georgévitch, J. 1917. Recherches sur le développement de *Ceratomyxa herouardi* Georgév. *Arch. Zool. Exp. Gen.* 56: 375-399.
 24. Georgévitch, J. 1926. Protistologica III. Sur la *Cocomyxa* de la sardine. *Arch. Zool. Exp. Gen.* 65(3): 57-63.
 25. Georgévitch, J. 1929. Recherches sur *Ceratomyxa maenae* nov. spec. *Arch. Protistenkd.* 65: 106-123.
 26. Gurley, R.R. 1894. Myxosporidia, or psorosperms of fishes, and the epidemics produced by them. *Rep. U.S. Comm. Fish Fisheries* for 1892. 1892: 65-304.
 27. Holliman, A. 1994. Sea trout, *Salmo trutta*, a new host for the myxosporean *Kudoa thyrsites* (Gilchrist)? *Vet. Rec.* 134: 524-525.
 28. Iskov, M.P. and A.K. Karataev. 1984. New species of myxosporidian from Black Sea scad. *Vestn. Zool.* 1984(4): 77-78. (in Russian)
 29. Kabata, Z. 1960. On two myxosporidian parasites of marine fishes, including one new species (*Ceratomyxa tenuispora*). *Ann. Mag. Nat. Hist.* 29: 305-308.
 30. Kabata, Z. 1962. Five new species of Myxosporidia from marine fishes. *Parasitology* 52: 177-186.
 31. Karataev, A.K. 1983. A new Myxosporidia species from the Black Sea broad-nosed pipefish. *Vestn. Zool.* 1983(4): 80-81. (in Russian)
 32. Karataev, A.K. and M.P. Iskov. 1984. New Myxosporidia species from Black Sea sand smelt *Atherina mochon pontica*. *Vestn. Zool.* 1984(1): 59-60. (in Russian)
 33. Kovaleva, A.A. 1988. Myxosporidia of the genus *Chloromyxum* (Cnidospora, Myxosporea) of cartilaginous fishes from the Atlantic coast of Africa. *Parazitologiya* 22: 384-388. (in Russian)
 34. Kovaleva, A.A. and A.V. Gaevskaya. 1979. Two new species of Myxosporidia of the genus *Myxoproteus* from the Celtic Sea fishes. *Parazitologiya* 13: 437-439. (in Russian)
 35. Kovaleva, A.A. and A.V. Gaevskaya. 1981. On new findings of myxosporidia of the genus *Parvicapsula* in the Atlantic fishes. *Zool. Zh.* 60: 771-773. (in Russian)
 36. Kovaleva, A.A. and A.V. Gaevskaya. 1988. Myxosporidia (Protozoa, Myxosporea) of *Beryx splendens* from the Atlantic ocean. *Parazitologiya* 22: 250-254. (in Russian)
 37. Kovaleva, A.A. and A.V. Zubchenko. 1984. New information on the North Atlantic myxosporidiae fauna. *Parazitologiya* 18: 484-487. (in Russian)
 38. Kovaleva, A.A., A.V. Zubchenko and V.K. Krasin. 1983. Foundation of a new myxosporidean family (Protozoa, Myxosporidia) with a description of two new genera. *Parazitologiya* 17: 195-202. (in Russian)
 39. Kovaleva, A.A., S.S. Shulman and V.N. Jakovlev. 1979. Myxosporidians of the genus *Kudoa* (Myxosporidia, Multivalvulae) from the Atlantic Ocean basin. *Trudy Zool. Inst. Akad. Nauk. SSSR* 78: 42-64. (in Russian)
 40. Kovaleva, A.A., Z.S. Donetz and M.G. Kolesnikova. 1989. New Myxosporea species (Cnidospora) from the Black Sea fishes. *Vestn. Zool.* 1989(5): 75-79. (in Russian)
 41. Labb  , A. 1899. Sporozoa. In: *Das Tierreich*. Vol. 5., Berlin: Deutsche Zool. Gesellsch. 180 p..
 42. Levsen, A. 1990. *Some parasites of flatfish (Pleuronectiformes) from the west coast of Norway*. Cand. Scient. thesis, University of Bergen. 102 p.
 43. Leydig, F. 1851. Über Psorospermien und Gregarinien. *Arch. Anat. Physiol.* 1851: 221-234.
 44. Lom, J., E.R. Noble and M. Laird. 1975. Myxosporidia from the deep - sea fish, *Macrourus berglax*, off Newfoundland and Iceland. *Folia Parasitol.* 22: 105-109.
 45. Lom, J., I. Dykov   and S. Lhot  kov  . 1983. *Kudoa lunata* n. sp. (Myxozoa, Myxosporea) and notes on the nature of muscular "cysts" of the genus *Kudoa*. *Arch. Protistenkd.* 127: 387-397.
 46. Lubat, V., B.M. Radujkovic, A. Marques and G. Bouix. 1989. Parasites des poissons marins du Montenegro: Myxosporidies. *Acta Adriatica* 30: 31-50.
 47. MacKenzie, K. 1983. Parasites as biological tags in fish population studies. *Adv. Appl. Biol.* 7: 251-331.
 48. Moser, M. and E.R. Noble. 1977. Three genera of myxosporidia (Protozoa) in macrourid fishes. *Int. J. Parasitol.* 7: 93-96.
 49. Myklebust, C.F. 1999. *Parasites of European hake (Merluccius merluccius (L.))*. Potential pathogens in future hake culture? Cand. Scient. thesis, University of Bergen. 81 p. (in Norwegian).
 50. Naidenova, N.N. 1968. *Ortholinea gobiusi* sp. n. (Myxosporidia, Bivalvulae) from *Gobius ophiocephalus* in the Black Sea. *Biologiya Morya*, Kiev 14: 15-18. (in Russian)
 51. Naidenova, N.N. 1970. Parasite fauna of fishes in the family Gobiidae in the Azov Sea. *Biologiya Morya*, Kiev 20: 84-113. (in Russian)
 52. Naidenova, N.N. and W.E. Zaika. 1969. Two new species of Protozoa from the fishes of the Black Sea. *Parazitologiya* 3: 97-101. (in Russian)
 53. Naidenova, N.N., S.S. Shulman and Z.S. Donetz. 1975. Class cnidosporidians - Cnidosporidia Doflein, 1901, emend. Shulman, 1967. In: *Identification of the parasites of vertebrates in the Black- and Azov Seas*, V.N. Gaevskaya et al., editors. Akademiya Nauk Ukrainskoj SSR, "Naukova Dumka": Kiev. p. 20-50. (in Russian)
 54. Naville, A. 1930. Recherches sur la sexualit   chez les myxosporidies. *Arch. Protistenkd.* 69: 327-400.
 55. Nemeczek, A. 1922.   ber *Zschokkella rovignensis* spec. nov. *Arch. Protistenkd.* 45: 390-400.
 56. Palmer, R. 1995. *Kudoa* in Ireland. Abstracts book, European Association of Fish Pathologists, Seventh international conference "Diseases of Fish and Shellfish". 1995. p. 132. Palma, Mallorca, Spain: EAAP.
 57. Pampoulie, C., A. Marques, E. Rosecchi, A.J. Crivelli and J.-L. Bouchereau. 1999. A new myxosporean parasite, *Kudoa camarguensis* n. sp., recorded on two goby species (Teleostei: Pisces) in the Rh  ne delta (Mediterranean Sea, France). *J. Euk. Microbiol.* 46: 304-310.
 58. Parenzan, P. 1966. *Myxobolus mugilis* e *Myxobolus branchialis*, due nuovi Missosporidi parassiti di *Mugil chelo* dello Jonio. *Boll. Soc. Nat. Napoli.* 75: 3-8.
 59. Parisi, B. 1912. Primo contributo alla distribuzione geografica dei Missosporidi in Italia. *Atti Soc. Ital. Sci. Nat. Mus. Civ. Milano* 50: 283-299.
 60. Perard, C. 1928. Sur une maladie du maquereau (*Scomber scomber* L.) due 脳 une Myxosporidie: *Chloromyxum histolyticum* n. sp. *C.R. Acad. Sci. Paris* 186: 108-110.
 61. Petrushevsky, G.K. 1932. Zur Systematik und Cytologie der Myxosporidia aus einigen Fischen des Wei  en Meeres. *Arch. Protistenkd.* 78: 543-556.
 62. Pogoreltseva, T.P. 1964. Data for the study of parasitic protozoans of fish in the Black Sea. *Problemy Parazitologii. Trudy Ukr. Resp. Obshch. Parazitol.* 3: 16-29. (in Russian)
 63. Polyanskii, Yu.I. 1955. Parasites of the fish of the Barents Sea. *Izdatelstvo Akademii Nauk SSSR*, Moskva 19 (English translation by Israel program for scientific translations, Jerusalem 1966, 158 pp).
 64. Priebe, K. 1987. Disseminated degeneration of somatic musculature of a blue ling (*Molva dypterygia*) caused by a *Trilospora* species. *Bull. Eur. Ass. Fish Pathol.* 7: 86-89.
 65. Pulsford, A. and R.A. Matthews. 1982. An ultrastructural study of *Myxobolus exiguis* Thelohan, 1895

- (Myxosporea) from the grey mullet, *Crenimugil labrosus* (Risso). *J. Fish Dis.* 5: 509-526.
66. Schröder, O. 1907. Beiträge zur Entwicklungsgeschichte der Myxosporidien. *Sphaeromyxa sabrazei* (Laveran & Mesnil). *Arch. Protistenkd.* 9: 359-381.
67. Schubert, G., V. Sprague and R. Reinboth. 1975. Observations on a new species of *Unicapsula* (Myxosporida) in the fish *Maena smaris* (L.) by conventional and electronmicroscopy. *Z. Parasitenkd.* 46: 245-252.
68. Shulman, S.S. 1953. New and poorly studied myxosporidia (Myxosporidia, Bivalvulae) from Fishes in the White Sea. *Zool. Zh.* 32: 384-393.
69. Shulman, S.S. 1966. *Myxosporidia of the USSR*. Moskva: Akademiya Nauk SSSR, Zoologicheskii Institut. Nauka.
70. Shulman, S.S. and R.E. Shulman-Albova. 1953. *Parasites of Fishes of the White Sea*. Izdatelstvo Akademii Nauk SSSR, Moskva: 198 pp. (in Russian)
71. Shulman, S.S., A.A. Kovaleva and V.R. Dubina. 1979. New myxosporidians from fishes of the Atlantic coast of Africa. *Parazitologiya* 13: 71-79. (in Russian)
72. Sitjà-Bobadilla, A. and P. Alvarez-Pellitero. 1990. *Sphaerospora testicularis* sp. nov. (Myxosporea: Sphaerosporidae) in wild and cultured sea bass, *Dicentrarchus labrax* (L.), from the Spanish Mediterranean area. *J. Fish Dis.* 13: 193-203.
73. Sitjà-Bobadilla, A. and P. Alvarez-Pellitero. 1992. Light and electron microscopic description of *Sphaerospora dicentrarchi* n. sp. (Myxosporea: Sphaerosporidae) from wild and cultured sea bass, *Dicentrarchus labrax* L. *J. Protozool.* 39(2): 273-281.
74. Sitjà-Bobadilla, A. and P. Alvarez-Pellitero. 1993. Light and electron microscopical description of *Ceratomyxa labracis* n. sp. and a redescription of *C. diploidae* (Myxosporea: Bivalvulida) from wild and cultured Mediterranean sea bass *Dicentrarchus labrax* (L.) (Teleostei: Serranidae). *Syst. Parasitol.* 26: 215-223.
75. Sitjà-Bobadilla, A. and P. Alvarez-Pellitero. 1993. *Zschokkella mugilis* n. sp. (Myxosporea: Bivalvulida) from mullets (Teleostei: Mugilidae) of Mediterranean waters: light and electron microscopic description. *J. Euk. Microbiol.* 40: 755-764.
76. Sitjà-Bobadilla, A. and P. Alvarez-Pellitero. 1995. Light and electron microscopic description of *Polysporoplasma* n. g. (Myxosporea: Bivalvulida), *Polysporoplasma sparis* n. sp. from *Sparus aurata* (L.), and *Polysporoplasma mugilis* n. sp. from *Liza aurata*. *Europ. J. Protistol.* 31: 77-89.
77. Sitjà-Bobadilla, A., O. Palenzuela and P. Alvarez-Pellitero. 1995. *Ceratomyxa sparosaurati* n. sp. (Myxosporea: Bivalvulida), a new parasite from cultured gilthead seabream (*Sparus aurata* L.) (Teleostei: Sparidae); light and electron microscopic description. *J. Euk. Microbiol.* 42: 529-539.
78. Stempell, W. 1919. Untersuchungen über *Leptotheca coris* n. sp. und das in dieser schmarotzende *Nosema marionis* Thél. *Arch. Protistenkd.* 40: 113-157.
79. Thélohan, P. 1895. Recherches sur les Myxosporidies. *Bull. Sci. Fr. Belg.* 5: 100-394.
80. Théodorides, J. 1955. Myxosporidies & microsporidies parasites de Poissons marins de Banyuls. *Vie Milieu* 6: 60-66.
81. Tripathi, Y.R. 1948. Some new myxosporidia from Plymouth with a proposed new classification of the order. *Parasitology* 39: 110-118.
82. Tuzet, O. and R. Ormières. 1957. *Ceratomyxa anguillae* n. sp. myxosporidie parasite de la vesicule biliaire des anguilles de l'Etang de Thau. *Ann. Parasitol. Hum. Comp.* 32: 189-196.
83. Vismanis, K., Z. Petrina, R. Eglite, A. Volkova and B. Sable. 1980. Materialen über Parasitenfauna einiger Objekte des Fischfangs im Rigaer Meerbusen. *Zoologijas Muz. Rak. P. Stuckas Latvij. Valsts Univ.* 1980: 1-12. (in Russian)
84. Wierzbicka, J. 1986. *Paramyxoproteus reinhardtii* gen. n. & sp. n. (Bivalvulida, Myxospora), a parasite of *Reinhardtius hippoglossoides* Walbaum, 1792. *Acta Protozool.* 25: 227-234.
85. Woodcock, H.M., 1903. On myxosporidia in flat-fish. *Rep. Lancashire Sea Fish. Lab.* 12/13: 46-62.
86. Yoshino, T.P. and E.R. Noble. 1973. Myxosporidia in macrourid fishes of the North Atlantic. *Can. J. Zool.* 51: 745-752.
87. Yurakhno, M.V. 1993. New data on the fauna of Myxosporidians from fishes of the Black Sea. *Parazitologiya* 27: 320-326. (in Russian)
88. Yurakhno, V.M. 1988. New data on fish Myxosporea of the Black Sea. *Parazitologiya* 22: 521-523. (in Russian)
89. Yurakhno, V.M. 1991. New species of Myxosporidia from fishes of the Black Sea. *Parazitologiya* 25: 104-109. (in Russian)
90. Zaika, V.E. 1966. Fauna of protozoans, parasites of fish in the Black Sea. In: *The helminth fauna of animals in the southern seas*. Naukova Dumka: Kiev. p. 13-31. (in Russian)
91. Zaika, V.E. 1968. Parasite Protozoa of the Black Sea fishes. *Biol. Morya*, Kiev 14: 41-46. (in Russian)
92. Zubchenko, A.V. 1981. *Myxidium profundum* Zubtschenko, 1981 (Myxosporidia: Myxidiidae), a new name for *M. noblei* Zubtschenko, 1980. *Parazitologiya* 15: 368. (in Russian)
93. Zubchenko, A.V. and V.K. Krasin. 1980. Mixosporidia of the genus *Myxidium* in some macrourids from northern Atlantic and Pacific. *Parazitologiya* 14: 168-176. (in Russian)

PHYLUM MYXOZOA

Class Myxospora

Order Bivalvulida

Suborder Sphaeromyxina

Family Sphaeromyxidae

<i>Sphaeromyxa</i>		
<i>atherinae</i> Karataev & Iskov, 1984	M	
<i>balbianii</i> Thélohan, 1892	A	M
<i>gasterostei</i> Georgévitch, 1916	A	
<i>heltandi</i> Auerbach, 1909	A	
<i>incurvata</i> Doflein, 1898		M
<i>longa</i> Dunkerley, 1921	A	
<i>minuta</i> Polyanskii, 1955	A	
<i>ovata</i> Dunkerley, 1921	A	
<i>sabrazesi</i> Laveran & Mesnil, 1900	A	M
<i>sebastopoli</i> Naidenova, 1970	M	

Suborder Variisporina

Family Myxidiidae

<i>Myxidium</i>		
<i>adriaticum</i> Lubat, Radujkovic, Marques & Bouix, 1989	M	
<i>benthophili</i> Naidenova, 1970	M	
<i>bergense</i> Auerbach, 1909	A	
<i>cochleatum</i> Yurakhno, 1991	M	
<i>coryphaenoidium</i> Noble, 1966	A	
<i>depressum</i> Parisi, 1912	M	
<i>euzeti</i> Lubat, Radujkovic, Marques & Bouix, 1989	M	
<i>gadi</i> Georgévitch, 1916	A	M
<i>giganteum</i> Doflein, 1898	M	
<i>incervatum</i> Thélohan, 1892	A	M
<i>inflatum</i> Auerbach, 1909	A	
<i>intermedium</i> Dunkerley, 1921	A	
<i>laticurvum</i> Kabata, 1962	A	
= <i>M. trachinorum</i> Canning, Curry, Anderson & Okamura, 1999		
<i>leei</i> Diamant, Lom & Dykova, 1994	M	
<i>macrocapsulare</i> Auerbach, 1910	A	
<i>melanocetum</i> Noble, 1966	A	
<i>melanostigmum</i> Noble, 1966	A	
<i>melanostomi</i> Naidenova, 1970	M	
<i>obliquelineolatum</i> Gaevskaya & Kovaleva, 1984	A	

<i>oviforme</i> Parisi, 1912	A	<i>hepseti</i> Thélohan, 1895	M
<i>parvum</i> Yurakhno, 1991	M	<i>informis</i> Auerbach, 1910	A M
<i>procerum</i> Auerbach, 1909	A	= <i>Ceratomyxa informis</i> (Auerbach, 1910) of authors	
<i>profundum</i> Zubchenko & Krasin, 1981	A	<i>kovaljovae</i> Bakay & Grudnev, 1998	A
= <i>M. noblei</i> Zubchenko & Krasin, 1980		<i>longipes</i> Auerbach, 1910	A
<i>pulchrum</i> Yurakhno, 1991	M	<i>macroformis</i> Gaevskaya & Kovaleva, 1984	A
<i>scorpii</i> Shulman-Albova, 1950	A	<i>polymorpha</i> Labb��, 1899	M
<i>sphaericum</i> Th��lohan, 1895	A M	= <i>L. elongata</i> Th��lohan, 1895 part	
<i>virens</i> Gaevskaya & Kovaleva, 1984	A	<i>quadritaenia</i> Kovaleva & Zubchenko, 1984	A
<i>Zschokkella</i>		<i>renicola</i> Th��lohan, 1895	A M
<i>admiranda</i> Yurakhno, 1993	M	<i>scombri</i> Pogoreltzeva, 1964	M
<i>dogieli</i> Pogoreltseva, 1964	M	<i>simplex</i> Kovaleva & Zubchenko, 1984	
<i>hildae</i> Auerbach, 1909	A	<i>vikrami</i> Tripathi, 1948	A
<i>icterica</i> Diamant & Paperna, 1992	M		
<i>iskovi</i> Kovaleva, Donetz & Kolesnikova, 1989	M	<i>Ceratomyxa</i>	
<i>kudoi</i> Moser & Noble, 1977	A	<i>acanthuri</i> Kpatcha, Diebakate, Faye & Toguebaye, 1996	
<i>mugilis</i> Sitj��-Bobadilla & Alvarez-Pellitero, 1993	M	<i>allantoidea</i> Gaevskaya & Kovaleva, 1984	A
<i>nova</i> Klokaewa, 1914	A M	<i>anguillae</i> Tuzet & Ormieres, 1957	M
<i>rovignensis</i> Nemeczek, 1922	M	<i>appendiculata</i> Th��lohan, 1892	A M
<i>russelli</i> Tripathi, 1948	A	<i>arcuata</i> Th��lohan, 1892	A M
<i>sturionis</i> Tripathi, 1948	A	<i>artedielli</i> Polyanskii, 1955	A
<i>Coccoomyxa</i>		<i>auerbachi</i> Kabata, 1962	A
<i>morovi</i> L��ger & Hesse, 1907	A M	<i>beloneae</i> Lubat, Radujkovic, Marques & Bouix, 1989	M
Family Ortholineidae		<i>chromis</i> Lubat, Radujkovic, Marques & Bouix, 1989	M
<i>Ortholinea</i>		<i>coris</i> Georg��vitch, 1916	M
<i>divergens</i> (Th��lohan, 1895)	A M	= <i>Leptotheca coris</i> Stempell, 1918	
= <i>Sphaerospora divergens</i> Th��lohan, 1895		= <i>Leptotheca macrospora</i> of Kalenscher, 1927	
<i>gobiusi</i> Naidenova, 1968	M	<i>diploidae</i> Lubat, Radujkovic, Marques & Bouix, 1989	A M
<i>orientalis</i> (Shulman & Shulman-Albova, 1953)	A	<i>drepanopsetae</i> Averintzev, 1908	A
<i>sakinachanumae</i> Ibragimov, 1988		<i>dubia</i> Dunkerley, 1921	A
<i>Sinuolineidae</i>		<i>elegans</i> Jameson, 1929	M
<i>Sinuolinaea</i>		<i>globulifera</i> Th��lohan, 1895	M
<i>cycloptera</i> Bazikalova, 1932	A	<i>herouardi</i> Georg��vitch, 1916	M
<i>murmanica</i> Bazikalova, 1932	A	<i>hopkinsi</i> Jameson, 1929	A
<i>rebae</i> Tripathi, 1948	A M	<i>inaequalis</i> Doflein, 1898	M
<i>schulmani</i> Gaevskaya & Kovaleva, 1979	A	<i>labracis</i> Sitj��-Bobadilla & Alvarez-Pellitero, 1993	A M
<i>schulmani</i> Karataev & Iskov, 1984	M	<i>lata</i> Dunkerley, 1921	A
<i>sinuosa</i> Shulman, 1953	A	<i>linospora</i> Doflein, 1898	M
<i>Davisia</i>		<i>longispina</i> Petruschewsk, 1932	A
<i>amoena</i> Gaevskaya, Kovaleva & Umnova, 1980	A	<i>macrospora</i> (Auerbach, 1909)	A
= <i>D. longibrachia</i> of Kabata, 1962 part		= <i>Leptotheca macrospora</i> Auerbach, 1909	
<i>longibrachia</i> Kabata, 1962	A	<i>maenae</i> Georg��vitch, 1929	M
<i>newfoundlandia</i> Yoshino & Noble, 1973	A	<i>markewitchi</i> Iskov & Karataev, 1984	M
<i>ophidioni</i> Zaika, 1966	M	<i>maxima</i> Gaevskaya & Kovaleva, 1980	
<i>Myxoproteus</i>		<i>merlangi</i> Zaika, 1966	M
<i>ambiguus</i> (Th��lohan, 1895)	A M	<i>orientalis</i> (Dogiel, 1948)	A
<i>caudatus</i> Shulman, 1953	A	<i>pallida</i> Th��lohan, 1895	M
<i>elongatus</i> Shulman, 1953	A	<i>parva</i> (Th��lohan, 1895)	A M
<i>formosus</i> Kovaleva & Gaevskaya, 1979	A	<i>peculiaria</i> Yurakhno, 1991	M
<i>irregularis</i> (Kabata, 1962)	A	<i>platichthys</i> (Fujita, 1923)	A
= <i>Sphaerospora irregularis</i> Kabata, 1962		<i>ramosa</i> Averintzev, 1908	A
<i>meridianalis</i> Evdokimova, 1977	A	= <i>C. ramosa</i> Averintzev, 1907 (<i>nomen nudum</i>)	
= <i>Conispora renalis</i> Sankurathri, 1977		<i>reticulata</i> Th��lohan, 1895	M
= <i>M. meridianalis</i> of authors		<i>scorpaenarum</i> Labb��, 1899	M
= <i>M. renalis</i> (Sankurathri, 1977)		<i>sparusaurati</i> Sitj��-Bobadilla, Palenzuela & Alvarez-Pellitero, 1995	A M
<i>scoleciformis</i> Kovaleva & Gaevskaya, 1979	A	<i>sphaerulosa</i> Th��lohan, 1892	A M
<i>Bipteria</i>		<i>tenuispora</i> Kabata, 1960	A
<i>admiranda</i> Kovaleva, Zubchenko & Krasin, 1983		= <i>C. magna</i> Gaevskaya & Kovaleva, 1980	
<i>Schulmania</i>		<i>trichiuri</i> Kpatcha, Diebakate, Faye & Toguebaye, 1996	
<i>aenigmatica</i> Kovaleva, Zubchenko & Krasin, 1983		<i>truncata</i> Th��lohan, 1895	A M
<i>ovale</i> Kovaleva, Zubchenko & Krasin, 1983			
<i>quadriolobata</i> Kovaleva, Zubchenko & Krasin, 1983	A		
= <i>Paramyxoproteus reinhardti</i> Wierzbicka, 1986			
<i>Fabespora</i>			
<i>nana</i> Naidenova & Zaika, 1969	M		
Family Ceratomyxidae			
<i>Leptotheca</i>			
<i>adeli</i> Bakay & Grudnev, 1998	A		
<i>agilis</i> (Th��lohan, 1892)	A M	<i>Sphaerospora</i>	
= <i>Ceratomyxa agilis</i> Th��lohan, 1892		<i>brevis</i> (Polyanskii, 1955)	A
<i>chromis</i> Lubat, Radujkovic, Marques & Bouix, 1989	M	= <i>Leptotheca brevis</i> (Polyanskii, 1955)	
<i>coelorhyncha</i> Yoshino & Noble, 1973	A	= <i>Myxoproteus brevis</i> Polyanskii, 1955	
= <i>L. coelorhynchus</i> of authors		<i>caspialosae</i> (Dogiel, 1939)	M
<i>elongata</i> Th��lohan, 1895	A M	<i>dicentrarchi</i> Sitj��-Bobadilla & Alvarez-Pellitero, 1992	A M

<i>Hoferebellus</i>			
<i>caudatus</i> (Parisi, 1910)	A M	<i>improvisus</i> Izyumova in Shulman, 1966	M
= <i>Mitraspora caudata</i> (Parisi, 1910)		<i>merlucii</i> (Perugia, 1891)	M
= <i>Sphaerospora caudata</i> Parisi, 1910		<i>muellicheri</i> Bütschli, 1882	A M
<i>Myxobolatus</i>		= <i>M. milleri</i>	
<i>convexum</i> Yurakhno, 1991	M	<i>mugauratus</i> Landsberg & Lom, 1991	M
<i>platessae</i> (Bazikalova, 1932)	A M	= <i>M. mugilis</i> Pogoreltseva, 1964	
<i>Palliatius</i>		<i>mugchelo</i> Landsberg & Lom, 1991	M
<i>indecorus</i> Shulman, Kovaleva & Dubina, 1979	A	= <i>M. mugilis</i> Parenzan, 1966	
<i>mirabilis</i> Shulman, Kovaleva & Dubina, 1979	A	<i>mugilis</i> (Perugia, 1891)	M
Family Chloromyxidae		<i>parenzani</i> Landsberg & Lom, 1991	M
<i>Chloromyxum</i>		= <i>M. branchialis</i> Parenzan, 1966	
<i>dogieli</i> Kovaleva, 1988	A	<i>parvus</i> Shulman, 1962	M
<i>kotorensis</i> Lubat, Radujkovic, Marques & Bouix, 1989	M	<i>petruschewskii</i> Zhukov, 1964	A
<i>leydigi</i> Mingazzini, 1890	A M	<i>raibauti</i> Fall, Kpatcha, Diebakate, Faye & Toguebaye, 1997	
<i>osmanovi</i> Karataev, 1983	M	<i>spinacurvatura</i> Maeno, Sorimachi, Ogawa & Egusa, 1990	M
<i>ovatum</i> Jameson, 1929	M	<i>Thelohanellus</i>	
<i>partistriatus</i> Kovaleva, Donetsk & Kolesnikova, 1989	M	<i>pyriformis</i> (Thélohan, 1892)	
<i>psetti</i> Kovaleva, Donetsk & Kolesnikova, 1989	M	<i>Henneguya</i>	
<i>schulmani</i> Kovaleva, 1988	A M	<i>brachideuteri</i> Kpatcha, Faye, Diebakate, Fall & Toguebaye, 1997	
<i>striatellus</i> Kovaleva, 1988	A	<i>kayarensis</i> Kpatcha, Faye, Diebakate, Fall & Toguebaye, 1997	
<i>trachuri</i> Iskov & Karataev, 1984	M	<i>mbourensis</i> Kpatcha, Faye, Diebakate, Fall & Toguebaye, 1997	
Family Auerbachiidae		<i>neapolitana</i> Parisi, 1912	M
<i>Auerbachia</i>		<i>ouakamensis</i> Kpatcha, Faye, Diebakate, Fall & Toguebaye, 1997	
<i>pulchra</i> Lom, Noble & Laird, 1975	A	<i>priacanthi</i> Kpatcha, Faye, Diebakate, Fall & Toguebaye, 1997	
Family Alatasporidae		<i>sympodae</i> Lubat, Radujkovic, Marques & Bouix, 1989	M
<i>Alataspora</i>		<i>yoffensis</i> Kpatcha, Faye, Diebakate, Fall & Toguebaye, 1997	
<i>africana</i> Shulman, Kovaleva & Dubina, 1979	A	Order Multivalvulida	
<i>caproi</i> Gaevskaya & Kovaleva, 1984	A	Family Trilosporidae	
<i>contrariocapsularia</i> Shulman, Kovaleva & Dubina, 1979	A	<i>Trilospora</i>	
<i>lepidum</i> Gaevskaya & Kovaleva, 1979	A	<i>minuta</i> Kovaleva & Zubchenko, 1984	A
<i>serenum</i> Gaevskaya & Kovaleva, 1979	A	<i>muscularis</i> Priebe, 1987	A
<i>solomonii</i> Yurakhno, 1988	M	<i>Unicapsula</i>	
<i>tetricum</i> Gaevskaya & Kovaleva, 1979	A	<i>pflugfelderi</i> Schuberg, Sprague & Reinboth, 1975	M
<i>Pseudalataspora</i>		Family Kudoidae	
<i>beryxi</i> Kovaleva & Gaevskaya, 1988	A	<i>Kudoa</i>	
<i>pontica</i> Kovaleva, Donetsk & Kolesnikova, 1989	M	<i>camarguensis</i> Pampoulie, Marques, Crivelli & Bouchereau, 1999	A M
<i>sebastei</i> Bakay & Grudnev, 1998	A	<i>clupeidae</i> (Hahn, 1917)	A
<i>umbraculiformis</i> Gaevskaya & Kovaleva, 1984	A	= <i>Chloromyxum clupeidae</i> Hahn, 1917	
Family Parvicapsulidae		= <i>Chloromyxum</i> sp. Auerbach, 1910	
<i>Parvicapsula</i>		<i>histolytica</i> (Pérard, 1928)	A
<i>asymmetrica</i> Shulman, 1953	A	= <i>Chloromyxum histolyticum</i> Pérard, 1928	
<i>schulmani</i> Kovaleva & Gaevskaya, 1981	A	<i>insolita</i> Shulman & Kovaleva in Kovaleva, Shulman & Yakovlev, 1979	A
<i>unicornis</i> Kabata, 1962	A	<i>kabatai</i> Shulman & Kovaleva in Kovaleva, Shulman & Yakovlev, 1979	A
<i>Neoparvicapsula</i>		<i>lunata</i> Lom, Dyková & Lhotákova, 1983	M
<i>ovalis</i> Gaevskaya, Kovaleva & Shulman, 1982	A	<i>nova</i> Naidenova, 1975	A M
Suborder Platysporina		= <i>K. quadratum</i> (Thélohan, 1895) of Naidenova, 1970	
Family Myxobolidae		<i>quadratum</i> (Thélohan, 1895)	A M
<i>Myxobolus</i>		<i>stellula</i> Yurakhno, 1991	M
<i>aeglefini</i> Auerbach, 1906	A	<i>thyrsites</i> (Gilchrist, 1924)	A
= <i>M. esmarkii</i> Woodcock in Johnstone & Woodcock, 1906			
= <i>M. lairdi</i> Moser & Noble, 1977			
<i>asymmetricus</i> (Parisi, 1912)	M		
<i>bizerti</i> Bahri & Marques, 1996	M		
<i>branchialis</i> (Markewitsch, 1932)	A M		
= <i>Lentospora branchialis</i> Markewitsch, 1932			
= <i>Myxosoma branchialis</i> (Markewitsch, 1932)			
<i>episquamalis</i> Egusa, Maeno & Sorimachi, 1990	A M		
= <i>M. exiguum</i> Théloha, 1895 of Bahri & Marques, 1995			
= <i>M. sp.</i> Menezes, 1984			
= <i>M. sp.</i> Overstreet & Howse, 1977			
<i>exiguus</i> Thélohan, 1895	A M		
= <i>Myxosporidium mugilis</i> Perugia, 1891			
<i>gobii</i> Naidenova, 1975	M		
= <i>M. naidenovae</i> Shulman, 1984			
<i>goreensis</i> Fall, Kpatcha, Diebakate, Faye & Toguebaye, 1997			
<i>hannensis</i> Fall, Kpatcha, Diebakate, Faye & Toguebaye, 1997			
<i>ichkeulensis</i> Bahri & Marques, 1996	M		

PORIFERA

Compiled by R.W.M. van Soest

Data consist of a species name considered valid or status unknown, followed by author and year of erection. Obvious synonyms (based on general consensus of past authors) are omitted, but all doubtful or contested cases are included. No subsequent records or distribution data are given. The entries are in most cases not recently validated, so are based on opinions of previous authors or experience of the present list author. Care should be taken when using these entries for other purposes than simple reference data. There is no guarantee for completeness of the species list. The system followed here is that of the most recent insights in the higher taxa classification. A project in process of near completion is the Systema Porifera initiative (editors J.N.A. Hooper & R.W.M. van Soest). This will result in a 25 author collective revision of all Porifera taxa above the species level, including keys to their identification. Until this volume appears, the present list remains tentative as far as classification and the use of genus and family names is concerned.

N. Boury-Esnault assisted in the compilation of the Calcarea.

References

- Ackers, R.G.A., Moss, D. & Picton, B.E. 1992. *Sponges of the British Isles ('Sponge V'), a colour guide and working document*. Marine Conservation Society: 1-175.
- Arndt, W. 1935. Porifera (systematischer Teil). IIIa. In: Grimpe, & G. & E. Wagler, (eds.). *Die Tierwelt der Nord- und Ostsee*: 1-140.
- Bergquist, P.R. 1978. *Sponges*. Hutchinson University Press. London: 1-268, pls. 1-12.
- Borojévic, C., Cabioch, L. & C.Lévi, 1968. *Inventaire de la faune marine de Roscoff. Spongaires*: 1-44 (Roscoff, Station Biologique).
- Boury-Esnault, N. & Lopes, T. 1985. Les Démospanges littoraux de l'archipel des Açores. *Annales de l'Institut Océanographique de Monaco* 61(2): 149-225.
- Boury-Esnault, N., Pansini, M. & Uriz, M.J. 1994. Spongaires bathyaux de la mer d'Alboran et du golfe ibéro-marocain. *Mémoires du Muséum national d'Histoire naturelle* 160 (Zoologie): 1-174.
- Bowerbank, J.S. 1864, 1866, 1874, 1882. *A monograph of the British Spongidae I, II, III, and IV*. Ray Society, London.
- Burton, M. 1930. Norwegian sponges from the Norman collection. *Proceedings of the Zoological Society of London* 1930(2): 487-546, pls. 1-2.
- Cabioch, L. 1968. Contribution à la connaissance de la faune des spongaires de la Manche occidentale. Démospanges de la région de Roscoff. *Cahiers de Biologie marine* 9: 211-246.
- Koltun, V.M. 1959. *[The cornacuspongida of the northern and fareastern seas of the USSR]. [Keys to the fauna of the USSR]*, Moscow, Leningrad, Akademia Nauk SSSR: 1-236, pls 1-43. (in Russian).
- Koltun, V.M. 1966. *[Tetraxonid sponges of the northern and far eastern seas of the USSR. (Order Tetraxonida)]. [Keys to the fauna of the USSR]*, 90: 1-111, 38 pls. (in Russian).
- Lévi, C. 1973. Systématique de la classe de Demospongia (Démospanges). *Traité de Zoologie* 3(1): 577-631.
- Lundbeck, W. 1902, 1905, 1910. . Homorrhaphidae, Heterorrhaphidae, Desmacidonidae. *Danish Ingolf Expedition* 6(1): 1-108, 6(2): 1-219, 6(3): 1-124.
- Pulitzer-Finali, G. 1983. A collection of Mediterranean Demospongiae (Porifera) with, in appendix, a list of the Demospongiae hitherto recorded from the Mediterranean sea. *Annali del Museo Civico di Storia Naturale di Genova* 84: 445-621.
- Sarà, M. 1972. Guida ai Poriferi della Fauna Italiana. *Quaderni della Civica Stazione Idrobiologica di Milano* 3-4: 53-97.
- Soest, R.W.M. Van, Picton, B.E. & Morrow, C. 1999. *Sponges of the North East Atlantic*. CD-ROM ETI World Biodiversity Series, Amsterdam.
- Topsent, E. 1894, 1895, 1900. Étude monographique des Spongaires de France. I. Tetractinellida. II. Carnosa. III. Hadromerida. *Arch. Zool. exp. gén.*, (3) 2: 259-400, (3) 3: 493-590, (3) 8: 1-331.
- Topsent, E. 1892, 1904, 1928. Contribution à l'étude des Spongaires de l'Atlantique Nord. *Spongaires des Açores. Spongaires de l'Atlantique et de la Méditerranée provenant des croisières du Prince Albert Ier de Monaco. Résultats des Campagnes scientifiques du Prince de Monaco* 2: 1-165, 25: 1-280, 74: 1-376.
- Vacelet, J. 1969. Eponges de la roche du large et de l'étage bathyal de Méditerranée (récoltées de la soucoupe plongeante cousteau et dragages). *Mémoires du Muséum national d'Histoire naturelle* (A, Zoologie) 59(2): 146-220.
- Vosmaer, G.C.J. 1933-35. The sponges of the Bay of Naples. *Porifera Incalcarea*. *Capita Zoologica* 5: 457-848.

PHYLUM PORIFERA

Class Demospongiae

Subclass Homoscleromorpha

Order Homosclerophorida

Family Plakinidae

Corticium

- bowerbanki* Sara, 1960
- candelabrum* Schmidt, 1862
- reductum* Pulitzer-Finali, 1983
- ropsenti* Pouliquen, 1972

Octavella

- galangai* Tuzet & Paris, 1963

Oscarella

- lobularis* (Schmidt, 1862)
- = *Halisarca lobularis*

Placinocephala

- moncharmonti* (Sarà, 1960)
- = *Diactinolopha moncharmonti*

Plakina

- brachylopha* Topsent, 1928
- dilopha* Schulze, 1880
- monolopha* Schulze, 1880
- trilopha* Schulze, 1880

Plakinastrella

- copiosa* Schulze, 1880
- mixta* Maldonado, 1992
- oxeata* Topsent, 1904
- trunculifera* Topsent, 1928

Plakortis

- simplex* Schulze, 1880
- Pseudocorticium*

- jarrei* Boury-Esnault *et al.*, 1995

Subclass Tetractinomorpha

Order Astrophorida

Family Ancorinidae

Ancorina

- cerebrum* Schmidt, 1862
- radix* (Marenzeller, 1889)
- verruca* Schmidt, 1862
- wageneri* (Schmidt, 1862)
- = *Stelletta wageneri*

Calthropella

- geodioides* (Carter, 1876)
- = *Pachastrella geodioides*
- inopinata* Pulitzer-Finali, 1983
- pathologica* (Schmidt, 1868)
- recondita* Pulitzer-Finali, 1972
- simplex* Sollas, 1888
- simplex* subsp. *durissima* Topsent, 1892
- stelligera* (Schmidt, 1868)
- = *Corticium stelligera*

- Dercitancorina*
lesinensis (Lendenfeld, 1894)
- Dragmastra*
normani (Sollas, 1880)
 = *Stelletta normani*
normani subsp. *rhiphiophora* (Hentschel, 1929)
- Holoxea*
furtiva Topsent, 1892
- Penares*
candidata (Schmidt, 1868)
helleri (Schmidt, 1864)
 = *Stelletta helleri*
sclerobesa Topsent, 1904
- Stelletta*
addita (Topsent, 1938)
aspera Carter, 1871
boglicii Schmidt, 1862
carbonaria Schmidt, 1880
collingsi (Bowerbank, 1866)
defensa Pulitzer-Finali, 1983
dichoclada Pulitzer-Finali, 1983
dorsigera Schmidt, 1862
grubii Schmidt, 1862
hispida (Buccich, 1886)
 = *Ancorina hispida*
(Pilocrota) inermis Topsent, 1904
lactea Carter, 1871
mediterranea (Topsent, 1893)
pachastrelloides Carter, 1876
pumex (Nardo, 1847)
rhiphiophora Hentschel, 1929
simplicissima (Schmidt, 1868)
 = *Ancorina simplicissima*
stellata Topsent, 1893
stellata subsp. *adriatica* Sara, 1960
tuberosa (Topsent, 1892)
 = *Astrella tuberosa*
(Pilocrota) ventricosa Topsent, 1904
- Stryphnus*
fortis (Vosmaer, 1885)
 = *Stelletta fortis*
mucronatus (Schmidt, 1868)
 = *Stelletta mucronatus*
niger (Carter, 1871)
 = *Dercitus niger*
ponderosus (Bowerbank, 1866)
 = *Ecionemia ponderosus*
ponderosus subsp. *rudis* Sollas, 1888
- Family Pachastrellidae**
- Characella*
tripodaria (Schmidt, 1868)
- Dercitus*
abyssi (Carter, 1876)
 = *Corticium abyssi*
plicatus (Schmidt, 1868)
bucklandi (Bowerbank, 1866)
 = *Hymeniacidon bucklandi*
- Nethea*
dissimilis Sara, 1959
- Pachastrella*
abyssi Schmidt, 1870
debilis Topsent, 1892
echinorhabda Pulitzer-Finali, 1972
intexta Carter, 1876
monilifera Schmidt, 1868
ovisternata Von Lendenfeld, 1897
stylifera Von Lendenfeld, 1897
- Poecillastra*
compressa (Bowerbank, 1866)
 = *Ecionemia compressa*
amygdaloidea (Carter, 1876)
 = *Pachastrella amygdaloidea*
compressa (Bowerbank, 1866)
crassiuscula Sollas, 1888
cumana Vosmaer, 1894
- fragilis* Vosmaer, 1894
pachastrelloides (Carter, 1874)
rudiastra Pulitzer-Finali, 1983
saxicola (Topsent, 1892)
 = *Characella saxicola*
scabra (Schmidt, 1868)
sollasi (Topsent, 1892)
symbiotica Topsent, 1904
- Sphinctrella* (= *Vulcanella*)
aberrans Maldonado & Uriz, 1996
annulata Schmidt, 1870
armata (Hanitsch, 1895)
 = *Poecillastra armata*
cribrifera Sollas, 1888
gracilis Sollas, 1888
horrida Schmidt, 1870
linaresi Ferrer-Hernandez, 1914
ornata Sollas, 1888
verruculosa Pulitzer-Finali, 1983
- Triptolemus*
simplex Sara, 1959
- Family Geodiidae**
- Caminella*
loricata Lendenfeld, 1894
- Caminus*
vulcani Schmidt, 1862
- Erylus*
cantabricus (Ferrer-Hernandez, 1912, 1918)
 = *Scutastraea cantabricus*
corsicus Pulitzer-Finali, 1983
deficiens Topsent, 1927
discophorus (Schmidt, 1862)
 = *Stelletta discophorus*
discophorus subsp. *deficiens* Babio & Gondar, 1978
euastrum (Schmidt, 1868)
 = *Stellettinopsis euastrum*
euastrum subsp. *aspidotiscus* Topsent, 1928
expletus Topsent, 1928
granularis Topsent, 1904
mamillaris (Schmidt, 1862)
 = *Stelletta mamillaris*
nummulifer Topsent, 1892, 1904
oblongus Topsent, 1928
papillatus Topsent, 1928
papulifer Pulitzer-Finali, 1983
stellifer Topsent?
topsenti Von Lendenfeld, 1903
- Geodia*
atlantica (Stephens, 1914)
 = *Sidonops atlantica*
barretti Bowerbank, 1858
conchilega Schmidt, 1862
cydonium (Jameson, 1811)
 = *Alcyonium cydonium*
cydonium subsp. *aegagropila* Corriero, 1989
depressa Bowerbank, 1873
echinastrella Topsent, 1904
eosaster (Sollas, 1888)
geodina (Schmidt, 1868)
gigas Schmidt, 1862
glariosa Sollas, 1862
globus Schmidt, 1870
macandrewi Bowerbank, 1872
megastrella Carter, 1876
mesotriaena (Hentschel, 1929)
 = *Sidonops mesotriaena*
micropunctata Row, 1911
muelleri (Fleming, 1828)
nodastrella Carter, 1876
normani (Sollas, 1888)
 = *Cydonium normani*
pergamentacea Schmidt, 1870
placenta Schmidt, 1862
pyriformis (Vosmaer, 1885)
 = *Synops pyriformis*

- simplex* Schmidt, 1870
simplicissima Burton, 1931
sphaerooides (Vosmaer, 1882)
= *Isops sphaerooides*
spherastrella Topsent, 1904
(*Stellogeodia*) *stellosa* Czerniavsky, 1880
tuber Lendenfeld, 1894
tuberosa Schmidt, 1862
Isops
anceps (Vosmaer, 1894)
= *Synops anceps*
canaliculata (Schmidt, 1868)
intuta (Topsent, 1892)
= *Cydonium intuta*
maculosus Vosmaer, 1894
pachydermata Sollas, 1888
pallida Vosmaer, 1882
phlegraei Sollas, 1862
Pachymatisma
intermedia (Schmidt, 1868)
johnstonia (Bowerbank in Johnston, 1842)
= *Halichondria johnstonia*
normani Sollas, 1888
- Family Theneidae**
Thenea
abyssorum Koltun, 1959
levis Von Lendenfeld, 1903
muricata Bowerbank, 1858
schmidti Sollas, 1886
valdiviae Von Lendenfeld, 1903
- Family Thrombidae**
Thrombus
abyssi (Carter, 1873)
= *Corticium abyssi*
abyssi subsp. *niger* Topsent, 1904
- Family Coppatiidae**
Dorypleres
incrustans Topsent, 1892
Holoxea
furtiva Topsent, 1892
Jaspis
dendyi Sollas, 1888
inconditus (Topsent, 1892)
= *Coppatias inconditus*
incrustans (Topsent, 1890)
eudermis Lévi & Vacelet, 1958
johnstoni (Schmidt, 1862)
= *Viaja johnstoni*
johnstoni subsp. *incrustans* Topsent, 1904
- Order Lithistida**
- Family Corallistidae**
Corallistes
elegantior Schmidt, 1870
masoni (Bowerbank, 1869)
nolitangere Schmidt, 1870
Racodiscula
clava Schmidt, 1870
- Family Theonellidae**
Discodermia
polydiscus Du Bocage, 1869
ramifera Topsent, 1892
- Family Siphoniidae**
Leiodermatium
lynceus Schmidt, 1870
pfeifferae (Carter, 1876)
= *Azorica pfeifferae*
Macandrewia
azorica Gray, 1859
robusta Topsent, 1904
Siphonidium
ramosum Schmidt, 1870
- Family Desmantidae**
Desmanthus
- incrustans* (Topsent, 1889)
= *Aciculites incrustans*
Lithobubaris
tenens Vacelet, 1969
- Order Spirophorida**
- Family Tetillidae**
Cantabrina
erecta Ferrer-Hernandez, 1914
Chrotella
amphiacantha Topsent, 1904
Cinachyra
australiensis Carter, 1881
cavernosa (Lamarck, 1815)
tarentina Pulitzer-Finali, 1983
Samus
anonyma Gray, 1867
Tetilla
abyssorum (Carter, 1876)
= *Tethya cranium* subsp. *abyssorum*
= *Tethya cranium* subsp. *infrequens*
azorica (Topsent, 1904)
cranium (Müller, 1776)
= *Alcyonium cranium*
= *Alcyonium globosum*
disigma Topsent, 1904
geniculata Von Marenzeller, 1886
infrequens (Carter, 1876)
longipilis Topsent, 1904
nimia Topsent, 1904
polyura Schmidt, 1870
repens (Sara, 1958)
= *Craniella repens*
sandalina Sollas, 1886
sibirica (Fristedt, 1887)
= *Tethya sibirica*
tethyoides (Schmidt, 1870)
truncata Topsent, 1892
zetlandica (Carter, 1872)
= *Tethya zetlandica*
- Order Hadromerida**
- Family Clionidae**
Alectona
millari Carter, 1879
Cliona
abyssorum Carter, 1874
albicans Volz, 1939
alderi Hancock, 1849
angulata Hancock, 1849
burtoni Topsent, 1932
celata (Grant, 1826)
copiosa Sara, 1959
corallinoides Hancock, 1849
cretensis Pulitzer-Finali, 1983
hystrix (Johnston, 1842)
= *Halichondria hystrix*
levispira Topsent, 1898
northumbrica Hancock, 1849
gorgonioides Hancock, 1849
gracilis Hancock, 1849
howsei Hancock, 1849
janitrix Topsent, 1932
lobata Hancock, 1849
nigricans (Schmidt, 1862)
= *Papillina nigricans*
pontica Czerniavsky, 1880
pruvoti Topsent, 1900
rhodensis Rützler & Bromley, 1981
rovignensis Volz, 1939
sarai Melone, 1965
schmidti (Ridley, 1881)
stationis Nassonow, 1883
topsentii (Von Lendenfeld, 1896)
tremitensis Sara, 1961
typica (Nardo, 1833)

- vastifica* Hancock, 1849
vermifera Hancock, 1867
viridis (Schmidt, 1862)
 = *Vioa viridis*
Cliothosa
hancocki (Schmidt, 1862)
 = *Vioa hancocki*
Delectona
alboransis Rosell, 1996
ciconiae Bavestrello *et al.*, 1996
Dotona
pulchella Carter, 1881
Thoosa
armata Topsent, 1888
mollis Volz, 1939
tortonesei Sarà, 1958
- Family Spirastrellidae**
Spirastrella
aculeata Topsent, 1892
cunctatrix Schmidt, 1868
minax Topsent, 1888
- Spiroxya*
heteroclita Topsent, 1896
- Family Timeidae**
Diplastrella
bistellata (Schmidt, 1862)
 = *Suberites bistellata*
ornata Rützler & Sarà, 1962
- Timea*
acutostellata Hanitsch, 1894
bifidostellata Pulitzer-Finali, 1983
chondrilloides (Topsent, 1904)
 = *Hymedesmia chondrilloides*
crassa (Topsent, 1900)
 = *Hymedesmia crassa*
cumana Pulitzer-Finali, 1978
fasciata Topsent, 1934
geminstellata Pulitzer-Finali, 1978
halzei Topsent, 1894
irregularis Sarà & Sribelli, 1960
mixta (Topsent, 1896)
 = *Hymedesmia mixta*
simplistellata Pulitzer-Finali, 1983
stellata (Bowerbank, 1866)
 = *Hymedesmia stellata*
stellifasciata Sarà & Sribelli, 1960
stelligera (Carter, 1876)
 = *Suberites stelligera*
stellivarians (Carter, 1876)
 = *Suberites stellivarians*
tristellata (Topsent, 1892)
 = *Hymedesmia tristellata*
unistellata (Topsent, 1892)
 = *Hymedesmia unistellata*
unistellata subsp. *aspera* (Topsent, 1904)
- Family Placospongiidae**
Placospongia
decorcicans (Hanitsch, 1895)
 = *Physcaphora decorcicans*
- Family Hemiasterellidae**
Adreus
fascicularis (Bowerbank, 1866)
 = *Dictyocylindrus fascicularis*
Hemasterella
aristoteliana Voultsiadou-Koukoura & VanSoest, 1991
- Paratimea*
constellata (Topsent, 1893)
 = *Halicnemia constellata*
loennbergi (Alander, 1942)
 = *Halicnemia loennbergi*
loricata (Sarà, 1958)
 = *Halicnemia loricata*
oxeata Pulitzer-Finali, 1978
- pierantonii* (Sara, 1958)
 = *Halistella pierantonii*
Stelligera
mutilus Topsent, 1928
nix Von Lendenfeld, 1896
rigida (Montagu, 1818)
 = *Spongia rigida*
stuposa (Montagu, 1818)
 = *Spongia stuposa*
- Family Latrunculiidae**
Latrunculia
biannullata Topsent, 1892
citharistae Vacelet, 1969
cratera Du Bocage, 1869
insignis Topsent, 1892
normani Stephens, 1915
tarentina Pulitzer-Finali, 1983
tricincta Hentschel, 1929
triloba (Schmidt, 1875)
 = *Sceptrelia triloba*
- Microstyliifer*
rugosus Vacelet, 1969
- Podospongia*
loveni Du Bocage, 1869
- Sceptrinus*
richardi Topsent, 1898
- Family Tethyidae**
Halicmetes
elongata Boury-Esnault *et al.*, 1994
- Tethya*
aurantium (Pallas, 1766)
 = *Alcyonium aurantium*
citrina Sarà & Melone, 1965
norvegica Bowerbank, 1872
- Family Suberitidae**
Aaptos
aaptos (Schmidt, 1864)
 = *Anchorina aaptos*
papillatus (Keller, 1880)
 = *Tuberella papillatus*
- Atergia*
corticata Stephens, 1915
- Laxosuberites*
durus Stephens, 1915
ectyoninus Topsent, 1900
ferrerhernandezi Boury-Enault & Lopes, 1985
incrustans (Hansen, 1880)
 = *Suberites incrustans*
rugosus (Schmidt, 1868)
- Prosuberites*
brevispinus De Laubenfels, 1951
epiphytum (Lamarck, 1815)
 = *Alcyonium epiphytum*
longispinus Topsent, 1893
modestus Pulitzer-Finali, 1978
rugosus Topsent, 1893
- Pseudosuberites*
hyalinus (Ridley & Dendy, 1886)
 = *Hymeniacidon hyalinus*
mollis Topsent, 1925
montiniger (Carter, 1880)
 = *Suberites montiniger*
sadko Koltun, 1966
simplicima (Bowerbank, 1874)
 = *Hymedesmia simplicima*
sulphureus (Bean *in* Bowerbank, 1866)
 = *Hymeniacidon sulphureus*
- Pseudotrachya*
hystrix (Topsent, 1892)
 = *Trachya hystrix*
oxystyla Sarà, 1959
- Rhizaxinella*
biseta Topsent, 1904
elongata (Ridley & Dendy, 1886)

- = *Suberites elongata*
gracilis (Von Lendenfeld, 1896)
= *Suberites gracilis*
pyrifera (Delle Chiaje, 1828)
schaudini Hentschel, 1929
unisetia Topsent, 1904
- Suberites**
baffini Brøndsted, 1933
bursa Schmidt, 1862
caminatus Ridley & Dendy, 1886
capillitium Topsent, 1904
carnosus (Johnston, 1842)
= *Halichondria carnosus*
carnosus subsp. *incrustans* Topsent, 1900
concinna Lambe, 1900
creolloides Marenzeller, 1877
crustula (Bowerbank, 1866)
= *Hymeniacidon crustula*
distortus Schmidt, 1870
domuncula (Oliv, 1792)
= *Alcyonium domuncula*
elongatum Ridley & Dendy, 1886
ficus (Esper, 1794)
= *Alcyonium ficus*
gibbosiceps Topsent, 1904
glasenapii Merejkowski, 1879
hirsutus (Fleming, 1828)
= *Halichondria hirsutus*
hystrix Schmidt, 1868
laticeps Topsent, 1904
lobatus (Renier in Lieberkühn, 1859)
= *Lithumen lobatus*
luetkeni Schmidt, 1870
maculans (Johnston, 1842)
= *Halichondria maculans*
massa (Nardo, 1847)
montalbidus Carter, 1880
prototypus Czerniavsky, 1880
sevus (Johnston, 1842)
= *Halichondria sevus*
suberia (Montagu, 1818)
= *Spongia suberia*
syringella (Schmidt, 1868)
virgultosus (Johnston, 1842)
= *Halichondria virgultosus*
- Terpios**
coerulea (Carter, 1882)
= *T. fugax* auctt.
- Family Stylocordylidae**
Oxycordyla
pellita Topsent, 1904
- Stylocordyla**
borealis (Loven, 1868)
borealis subsp. *typica* Burton, 1931
- Family Polymastiidae**
Polymastia
agglutinans Ridley & Dendy, 1886, 1887
azorica Lévi & Vacelet, 1958
bartletti De Laubenfels, 1942
boletiformis (Lamarck, 1813)
= *Alcyonium boletiformis*
capitata Vosmaer, 1885
contigera Bowerbank, 1874
corticata Ridley & Dendy, 1886
ectofibrosa Boury-Esnault et al., 1994
grimaldi (Vosmaer, 1885)
= *P. penicillus*
hispida (Bowerbank, 1866)
infata Cabioch, 1968
laganoides Lambe, 1900
mamillaris (Müller, 1806)
= *Spongia mamillaris*
mamillaris subsp. *hyperborea* Hentschel, 1916
martae Boury-Esnault et al., 1994
mespilus Schmidt, 1873
- paupera* Fristedt, 1887
polytylota Vacelet, 1969
radiosa Bowerbank, 1866
spinula Bowerbank, 1866
sola Pulitzer-Finali, 1983
tissieri (Vacelet, 1961)
uberrima (Schmidt, 1870)
= *Rinalda uberrima*
- Quasillina**
brevis (Bowerbank, 1861)
= *Euplectella brevis*
intermedia Boury-Esnault et al., 1994
richardi Topsent, 1913
- Rhaphidorus**
setosus Topsent, 1898
- Ridleia**
oviformis Dendy, 1916
- Sphaerotylus**
borealis (Schwartshevsky, 1906)
schoenus (Sollas, 1882)
- Spirularia**
spinularia (Bowerbank, 1866)
= *Tethea spinularia*
- Tentorium**
semisuberites (Schmidt, 1870)
= *Thecophora semisuberites*
- Polymastia**
hemisphaerica (Sars, 1872)
- Trichostemma**
sarsi Ridley & Dendy, 1886, 1887
- Tylexocladus**
joubini Topsent, 1898
- Vosmaeria**
crustacea Fristedt, 1885
laevigata Topsent, 1896
- Weberella**
bursa Vosmaer, 1885
verrucosa Vacelet, 1960
- Order Chondrosida**
- Family Cbondrillidae**
Chondrilla
nucula Schmidt, 1862
- Chondrosia**
plebeja Schmidt, 1868
reniformis (Nardo, 1847)
- Thymosia**
guernei Topsent, 1895
- Thymosiopsis**
cuticulatus Vacelet & Perez, 1998
- Subclass Ceractinomorpha**
- Order Halichondrida**
- Family Axinellidae**
Acanthella
acuta Schmidt, 1862
annulata Sara, 1958
mastophora (Schmidt, 1870)
= *Axinella mastophora*
multiformis Vosmaer, 1882, 1885
- Auleta**
grantioides Lévi & Vacelet, 1958
pedunculata (Topsent, 1896)
sessilis Topsent, 1904
sycinularia Schmidt, 1870
- Axinella**
arctica (Vosmaer, 1885)
= *Phakellia arctica*
arcticasetosa Hentschel, 1929
babici Vacelet, 1961
blanca Koltun, 1959
calyciformis (Lamarck, 1813)
cannabina (Esper, 1794)
= *Spongia cannabina*
cinnamomea sensu Schmidt, 1873

- damicornis* (Esper, 1794)
 = *Spongia damicornis*
 = *Spongia lactuca*
dissimilis (Bowerbank, 1866)
 = *Isodictya dissimilis*
egregia Ridley, 1881
estacioi Carballo & Garcia-Gomez, 1995
flustra (Topsent, 1892)
 = *Tragorgia flustra*
guiteli Topsent, 1896
hirondelli (Topsent, 1892)
 = *Tragorgia hirondelli*
hispida Koltun, 1959
infundibuliformis (Linnaeus, 1767)
 = *Spongia infundibuliformis*
macrostyla Babic, 1922
mahonensis Ferrer-Hernandez, 1916
mammillata (Hanitsch, 1890)
minuta Lévi, 1957
pocillum (Bowerbank, 1866, 1874)
 = *Isodictya pocillum*
polypoides Schmidt, 1862
pseudominuta Bibiloni, 1993
pumila Babic, 1922
pyramidata Stephens, 1916, 1921
rugosa Schmidt, 1875
salicina Schmidt, 1868
setosa Hentschel, 1929
subdola (Bowerbank, 1866)
 = *Dictyocylindrus subdola*
trichophora Hentschel, 1929
variabilis (Vosmaer, 1882)
 = *Cribrochalina variabilis*
vasonuda Topsent, 1904
vellerea Topsent, 1904
verrucosa (Esper, 1794)
 = *Spongia verrucosa*
Axingella
perlucida (Topsent, 1896)
Homaxinella
informis Burton, 1930
Ophiraphidites
tortuosus Carter, 1876
Phakellia
beringensis Hentschel, 1929
bowerbanki Vosmaer, 1885
dalli Lambe, 1900
lambei Topsent, 1913
robusta Bowerbank, 1864
rugosa (Bowerbank, 1866)
ventilabrum (Linnaeus, 1767)
 = *Spongia ventilabrum*
ventilabrumconnexiva Ridley & Dendy, 1887
Pseudaxinella
sulcata Schmidt, 1875
Styliosa
cribrosa Micklugo-Maclay, 1870
- Family Bubaridae**
- Bubaris*
carcisia Vacelet, 1969
simplicima (Bowerbank, 1866)
 = *Microciona simplicima*
sosia Topsent, 1904
subtyla Pulitzer-Finali, 1983
vermiculata (Bowerbank, 1866)
 = *Hymeraphia vermiculata*
vermiculata subsp. *erecta* Carter, 1876
- Cerbaris*
torquata Topsent, 1898, 1904, 1928
- Hymerhabdia*
contracta Sara & Sirebelli, 1960
intermedia Sara & Sirebelli, 1960
oxytrunca Topsent, 1904
papillosa Sara & Sirebelli, 1960
typica Topsent, 1892
- Monocrepidium*
vermiculatum Topsent, 1898
- Rhabdobaris*
implicata Pulitzer-Finali, 1983
- Skeizia*
minuta Cabioch, 1968
- Family Agelasidae**
- Agelas*
ororides (Schmidt, 1864)
 = *Clathria oroides*
- Family Dictyonellidae**
- Dictyonella*
alonsoi Carballo *et al.*, 1996
incisa (Schmidt, 1880)
madeirensis (Topsent, 1928)
marsili (Topsent, 1893)
obtusa (Schmidt, 1862)
 = *Acanthella obtusa*
pelligera (Schmidt, 1862)
plicata (Schmidt, 1864)
- Scopalina*
azurea Bibiloni, 1993
lophyropoda Schmidt, 1862
- Tethyspira*
spinosa (Bowerbank, 1874)
 = *Tethea spinosa*
- Family Desmoxiyidae**
- Didiscus*
pseudodidiscoides (Corriero *et al.*, 1996)
 = *Dendroxea pseudodidiscoides*
styliferus Tsuruhashi, 1968
- Halicnemia*
arbuscula Topsent, 1928
azorica Topsent, 1928
geniculata Sara, 1958
patera Bowerbank, 1862
 = *Hymedesmia inflata*
"Halicortex"
loricata Sara, 1958
- Heteroxya*
corticata Topsent, 1898, 1904
- Higginsia*
mediterranea Pulitzer-Finali, 1978
pyriformis Brøndsted, 1916
strigilata (Lamarck, 1814)
 = *Spongia strigilata*
thielei (Topsent, 1892)
- Myrmekioderma*
spelaea (Pulitzer-Finali, 1983)
 = *Raphisia spelaea*
- Family Halichondriidae**
- Amorphinopsis*
filigrana (Schmidt, 1862)
 = *Reniera filigrana*
pallescens (Topsent, 1892)
 = *Hymeniacidon pallescens*
- Axinyssa*
digitata (Cabioch, 1968)
 = *Pseudaxinysa digitata*
- Ciocalapata*
almae Carballo *et al.*, 1996
- Ciocalypita*
penicillus Bowerbank, 1862, 1864, 1866, 1874
- Coelocalypita*
hyalina Pulitzer-Finali, 1978
- Halichondria*
agglomerans Cabioch, 1968
albescens Johnston, 1842
arenosa Hentschel, 1929
aspera Schwartzschevski, 1905, 1905
aurantiaca (Schmidt, 1864)
axinelloides Schwartzschevski, 1905, 1905
bowerbanki Burton, 1930
bowerbanki subsp. *stellifera* Breton *et al.*, 1995

- brunnea* (Schmidt, 1868)
 = *Raspagella brunnea*
colossea Lundbeck, 1902
contorta (Sarà, 1961)
 = *Topsentia contorta*
convolvens Sarà, 1960
coralloides Bowerbank, 1882
cornuloides Burton, 1954
difflciliis Lundbeck, 1902
diversispiculata Burton, 1930
dubia (Czerniavsky, 1880)
 = *Amorphina dubia*
fibrosa (Fristedt, 1887)
 = *Amorphina fibrosa*
foraminosa (Czerniavsky, 1880)
 = *Pellina foraminosa*
glabra Bowerbank, 1866
grisea (Fristedt, 1887)
 = *Amorphina grisea*
heterorrhaphis Breitfuss, 1912
labiata Hentschel, 1929
lactea (Bowerbank, 1866)
 = *Hymeniacidon lactea*
lambei Brøndsted, 1933
leuconoides Topsent, 1892, 1904
longispicula (Czerniavsky, 1880)
 = *Pellina longispicula*
membrana (Bowerbank, 1866)
 = *Hymeniacidon membrana*
nodosa (Fristedt, 1887)
 = *Amorphina nodosa*
normani Burton, 1930
oblonga (Hansen, 1885)
 = *Reniera oblonga*
osculum Lundbeck, 1909
paciscens (Schmidt, 1875)
 = *Amorphina paciscens*
panicea (Pallas, 1766)
 = *Spongia panicea*
renieroides (Fristedt, 1887)
 = *Amorphina renieroides*
semitubulosa subsp. *pontica* (Czerniavsky, 1880)
sitiens (Schmidt, 1870)
 = *Eumastia sitiens*
tenera (Marenzeller, 1877)
 = *Isodictya tenera*
tenuiderma Lundbeck, 1902
tenuispiculata Brøndsted, 1916
topsenti DeLaubenfels, 1936
velamentosa (Hansen, 1880)
 = *Amorphina velamentosa*
- Hymeniacidon*
assimilis (Levinsen, 1886)
aurea (Montagu, 1818)
 = *Spongia aurea*
fallax Bowerbank, 1866
consimilis Ferrer-Hernandez, 1914
fasciculata (Fristedt, 1887)
 = *Amorphina fasciculata*
gorbunovi Rezvoj, 1932
heliofila (Parker, 1910)
 = *Stylorella heliofila*
kitchingi (Burton, 1935)
 = *Rhaphidostyla kitchingi*
luxurians (Lieberkühn, 1859)
 = *Halichondria luxurians*
perlevis (Montagu, 1818)
 = *Spongia perlevis*
plumiger Ferrer-Hernandez, 1914
simplex Bowerbank, 1882
virgulatum Bowerbank, 1875, 1882)
- Leucophloeus*
ibericus Ferrer-Hernandez, 1914
mixtus Sarà, 1958, 1978
- Petromica*
- grimaldii* Topsent, 1898
Stylinos
fristedti Topsent, 1892
Spongisorites
cavernicola Bibiloni, 1993
difficilis (Lundbeck, 1902)
fibrosa (Fristedt, 1887)
 = *Amorphina fibrosa*
flavens Pulitzer-Finali, 1983
genitrix (Schmidt, 1870)
 = *Amorphina genitrix*
intricatus (Topsent, 1892)
maxima Uriz, 1978
placenta Topsent, 1896
placenta subsp. *rigidior* Topsent, 1904
- Topsentia*
annandalei (Ferrer-Hernandez, 1922)
 = *Oxeostilon annandalei*
fibrosa (Fristedt, 1887)
 = *Halichondria fibrosa*
garciae Bibiloni, 1993
glabra (Topsent, 1898)
 = *Anisoxya glabra*
pachastrelloides (Topsent, 1892)
 = *Halichondria pachastrelloides*
subtilis (Pulitzer-Finali, 1983)
 = *Laminospongia subtilis*
topsenti (Lévi & Vacelet, 1958)
 = *Spongisorites topsenti*
tenuispinosa (Topsent, 1894)
- Order Poecilosclerida**
- Suborder Microcionina**
- Family Raspailiidae**
- Acantheurypon*
pilosella Topsent, 1904
scabiosum Topsent, 1928
- Cyamon*
spinispinosum (Topsent, 1892)
- Dragmatyle*
lictor Topsent, 1928
- Endectyon*
delaubenfelsi Burton, 1930
pilosus Vacelet, 1961
teissieri Cabioch, 1968
- Eurypon*
cinctum Sara, 1960
clavatum (Bowerbank, 1866)
coronula (Bowerbank, 1874)
 = *Hymeraphia coronula*
(Rhabdoploca) curvispiculifera subsp. *geniculatum*
 (Topsent, 1892)
deniseae Vacelet, 1969
ditoxa Stephens, 1921
hispidulum (Topsent, 1904)
(Acantheurypon) incipiens (Topsent, 1928)
lacazei (Topsent, 1891)
longispiculum (Carter, 1874)
 = *Microciona longispiculum*
major Sara & Sribelli, 1960
microchela Stephens, 1921
minimum (Topsent, 1892)
 = *Hymeraphia minimum*
(Acantheurypon) mixtum Topsent, 1928
(Acantheurypon) mucronale (Topsent, 1928)
mucronatum (Topsent, 1904)
obtusum Vacelet, 1969
radiatum (Bowerbank, 1866)
 = *Hymedesmia radiatum*
simplex (Bowerbank, 1866)
 = *Halichondria simplex*
(Acantheurypon) spinispinosum (Topsent, 1904)
spitzbergensis (Fristedt, 1887)
 = *Hymeraphia spitzbergensis*
topsenti Pulitzer-Finali, 1983

- vesicularis* Sara & Sibelli, 1960
Hemectyon
 atlanticus (Lévi & Vacelet, 1958)
Heterectya
 villosa Thiele, 1898
Hymeraphia
 stellifera Bowerbank, 1864, 1866
 verticillata Bowerbank, 1866
Janulum
 spinispiculum (Carter, 1876)
 = *Isodictya spinispiculum*
Raspaciona
 aculeata (Johnston, 1842)
 = *Halichondria aculeata*
 calva Sara, 1958
 robusta Sara, 1958
Raspailia
 agnata (Topsent, 1896)
 = *Axinella agnata*
 bathyalis Boury-Esnault et al., 1993
 (*Syringella*) falcifera Topsent, 1892, 1904
 formidabilis Hanitsch, 1895
 freyeri Schmidt, 1862
 gracillima Topsent, 1894
 hispida (Montagu, 1818)
 = *Spongia hispida*
 howsei (Bowerbank, 1866)
 = *Dictyocylindrus howsei*
 (*Syringella*) humilis Topsent, 1892, 1904
 radiosa (Bowerbank, 1866)
 = *Dictyocylindrus radiosa*
 ramosa (Montagu, 1818)
 = *Spongia ramosa*
 rectangula (Bowerbank, 1874)
 = *Dictyocylindrus rectangula*
 setosa (Bowerbank, 1873)
 = *Dictyocylindrus setosa*
 ventilabrum (Bowerbank, 1866)
 = *Dictyocylindrus ventilabrum*
 viminalis Schmidt, 1862
 virgultosa (Bowerbank, 1866)
 = *Dictyocylindrus virgultosa*
Rhabdeurypon
 spinosum Vacelet, 1969
Rhabdoploca
 curvispiculifera (Carter, 1880)
Tricheurypon
 viride (Topsent, 1889)
 = *Hymeraphia viride*
- Family Rhabderemiidae**
- Rhabderemia*
 fascicularis Topsent, 1927, 1928
 gallica Van Soest & Hooper, 1993
 guernei Topsent, 1890
 intexta (Carter, 1876)
 = *Microciona intexta*
 minutula sensu Solorzano, 1991
 mutans Topsent, 1927, 1928
 ?parasitica (Carter, 1876)
 = *Corticium ?parasitica*
 profunda Boury-Esnault et al., 1994
 spinosa Topsent, 1896
 topsenti Van Soest & Hooper, 1993
 toxigera Topsent, 1892
- Family Microcionidae**
- Antho*
 arctica (Koltun, 1959)
 = *Plocamia arctica*
 brattegardi Van Soest & Stone, 1986
 circonflexa (Lévi, 1960)
 = *Plocamilla circonflexa*
 coriacea (Bowerbank, 1874)
 = *Isodictya coriacea*
 coriacea subsp. elegans (Ridley & Dendy, 1887)
 dichotoma (Esper, 1794)
- erecta* (Ferrer-Hernandez, 1922)
 = *Plocamilla erecta*
fragilis (Koltun, 1959)
 = *Plocamia fragilis*
involvens (Schmidt, 1864)
 = *Myxilla involvens*
involvens subsp. *inconstans* (Topsent, 1925)
lundbecki (Breitfuss, 1897)
novizelanica (Ridley, 1881)
oxeifera (Ferrer-Hernandez, 1921)
 = *Clathria oxeifera*
paradoxa (Babic, 1922)
paucispina Sara & Sibelli, 1962
Artemisia
apollinis Ridley & Dendy, 1886
arcigera (Schmidt, 1870)
 = *Suberites arcigera*
erecta Topsent, 1904
hispanica Ferrer-Hernandez, 1919
transiens Topsent, 1892
Bubaropsis
alborani Boury-Esnault et al., 1994
Clathria
acanthotoxa (Stephens, 1916, 1921)
 = *Eurypon acanthotoxa*
anancora (Topsent, 1904)
anchorata (Carter, 1874)
 = *Dictyocylindrus anchorata*
angularis (Sara & Sibelli, 1960)
 = *Microciona angularis*
armata (Bowerbank, 1866)
 = *Microciona armata*
ascendens (Cabioch, 1968)
 = *Microciona ascendens*
assimilis (Topsent, 1925)
 = *Microciona assimilis*
atrasanguinea (Bowerbank, 1862, 1866)
 = *Microciona atrasanguinea*
barleei (Bowerbank, 1866)
 = *Isodictya barleei*
basifixa (Topsent, 1913)
 = *Ophilitaspongia basifixa*
bitoxa (Burton, 1930)
 = *Hymantho bitoxa*
campecheae Hooper, 1996
cantabrica (Orueta, 1901)
chelifera Lévi, 1960
cleistochela (Topsent, 1925)
compressa Schmidt, 1862
coralloides (Oliv, 1792)
ctenichela (Alander, 1942)
 = *Microciona ctenichela*
delicata Lambe, 1900
depressa Sara & Melone, 1966
dianae (Schmidt, 1875)
 = *Suberites dianae*
ditoxa (Stephens, 1916)
 = *Eurypon ditoxa*
duplex (Sara, 1958)
 = *Microciona duplex*
elliptichela (Alander, 1942)
 = *Microciona elliptichela*
fallax (Bowerbank, 1866)
 = *Microciona fallax*
fregeti (Vacelet, 1969)
 = *Microciona fregeti*
frondiculata (Schmidt, 1864)
gradalis Topsent, 1925
haplotoxa (Topsent, 1928)
 = *Microciona haplotoxa*
heterotoxa (Hentschel, 1929)
 = *Microciona heterotoxa*
hjorti (Arnesen, 1920)
intermedia (Burton, 1930)
 = *Paresperia intermedia*

jolicoeuri (Topsent, 1892)
 = *Rhaphidophlus jolicoeuri*
laevis (Bowerbank, 1866)
 = *Microciona laevis*
levii (Sara & Siribelli, 1960)
 = *Microciona levii*
macrochela (Lévi, 1960)
marissuperi Pulitzer-Finali, 1983
microchela (Stephens, 1916, 1921)
normani (Burton, 1930)
 = *Hymantho normani*
osismica (Cabioch, 1968)
 = *Microciona osismica*
plurityla Pulitzer-Finali, 1983
palmatichela (Topsent, 1928)
 = *Hymedesmia palmatichela*
poecilosclera (Sara & Siribelli, 1960)
 = *Microciona poecilosclera*
primitiva (Koltun, 1956)
schmitti (De Laubenfels, 1942)
 = *Echinocladria schmitti*
seriata (Grant, 1826)
spinarcus (Carter & Hope, 1889)
 = *Microciona spinarcus*
strepsitoxa (Carter & Hope, 1889)
 = *Microciona strepsitoxa*
tenuissima (Stephens, 1916, 1921)
 = *Eurypon tenuissima*
toximajor (Topsent, 1925)
toxirecta (Sara & Siribelli, 1960)
 = *Microciona toxirecta*
toxistricta Topsent, 1925
toxityla (Sara, 1959)
 = *Microciona toxityla*
toxitenuis Topsent, 1925
toxivaria (Sara, 1959)
 = *Microciona toxivaria*
translata (Pulitzer-Finali, 1978)
 = *Ophilitaspongia translata*
tricurvata Carter, 1876
Melonchela
clathriata Koltun, 1955, 1959
Ophilitaspongia
papilla Ferrer-Hernandez, 1914, 1921

Family Iophonidae

Acarnus
levii Vacelet, 1960
tortilis Topsent, 1892

Acheliderma
lemniscata Topsent, 1892

Anisotylacantha
curvata Vacelet, 1969

Astylinifer
planus Topsent, 1904

Cornulotrocha
cheliradians Topsent, 1927

Cornulum
enteromorphoides Fristedt, 1887
textile Carter, 1876
ascidioides Fristedt, 1887

Iophon
chelifer Ridley & Dendy, 1886
doglieli Koltun, 1955
dubius (Hansen, 1880)
frigidus Lundbeck, 1909
frigidus subsp. *gracilis* Hentschel, 1929
funis Topsent, 1892
hyndmani (Bowerbank, 1864, 1866)
 = *Halichondria hyndmani*
nigricans (Bowerbank, 1866)
 = *Halichondria nigricans*
piceus (Vosmaer, 1882, 1885)
 = *Alebion piceus*
pommeraniae Thiele, 1903
scandens (Bowerbank, 1866)

= *Halichondria scandens*
spinulentum (Bowerbank, 1866)
 = *Microciona spinulentum*
variocillatus Alander, 1942
Quintoxilla
arcitenens (Topsent, 1892)
 = *Myxilla arcitenens*

Suborder Myxillina**Family Tedaniidae**

Tedania
anhelans (Lieberkühn, 1859)
 = *Halichondria anhelans*
commixta Ridley & Dendy, 1886
conuligera Topsent, 1892, 1904
flexistronyla Koltun, 1959
fragilis Lambe, 1900
phacellina Topsent, 1928
suctoria Schmidt, 1870

Family Myxillidae

Hymenancora
conjugens Lundbeck, 1910
duplicata (Lundbeck, 1910)
 = *Hymedesmia duplicata*
interjecta Lundbeck, 1910
pecqueryi (Topsent, 1892, 1904)
 = *Myxilla pecqueryi*
pectinata (Topsent, 1892)
 = *Dendoryx pectinata*
tenuisclera (Lundbeck, 1910)
 = *Hymedesmia tenuisclera*

Melonanchora

elliptica Carter, 1874
emphysema (Schmidt, 1875)
 = *Desmacidon emphysema*

Myxilla

acribaria De Laubenfels, 1942
barentsi Vosmaer, 1885
behringensis Lambe, 1900
brunnea (Hansen, 1880)
dentata Topsent, 1928
diversiancorata Lundbeck, 1905
fasciculata (Pallas, 1766)
fibrosa Levinson, 1893
fimbriata (Bowerbank, 1866)
 = *Isodictya fimbriata*

fimbriata subsp. *flexitornata* Rezvoi, 1925

(*Stelodoryx*) *flabellata* Burton, 1959

hastatiplicata Schwartschevski, 1905

incrustans (Johnston, 1842)

= *Halichondria incrustans*

incrustans subsp. *reses* (Topsent, 1892)

incrustans subsp. *viscosa* (Topsent, 1892, 1904)

inequitornata Burton, 1931

iophonoides Schwartschevski, 1905

iotrochotina (Topsent, 1892)

= *Dendoryx iotrochotina*

macrosigma Bourty-Esnault, 1971

mariana subsp. *polysigma* Topsent, 1904

parasitica (Lambe, 1900)

pedunculata Lundbeck, 1905

perspinosa Lundbeck, 1909

(*Stelodoryx*) *pluridentata* (Lundbeck, 1905)

(*Stelodoryx*) *procera* Topsent, 1904

prouhoi (Topsent, 1892)

= *Damiria prouhoi*

rosacea (Lieberkühn, 1859)

= *Halichondria rosacea*

septentrionalis Fristedt, 1887

tridens Schmidt, 1864

veneta Schmidt, 1862

Paramyxilla

mediterranea Sara & Siribelli, 1960

Plocamiancra

arndti Alander, 1942

- Styloptilon*
anchoratum Cabioch, 1968
- Family Iotrochotidae**
- Iotroota*
abyssi (Carter, 1874)
= *Halichondria abyssi*
acanthostylifera (Stephens, 1916, 1921)
= *Iotrochota acanthostylifera*
affinis (Lundbeck, 1909)
= *Iotrochota affinis*
magna (Lambe, 1900)
= *Iotrochota magna*
oxeata (Lundbeck, 1909)
= *Iotrochota oxeata*
polydentata (Lundbeck, 1905)
= *Iotrochota polydentata*
rotula (Topsent, 1904)
= *Hymetrochota rotula*
rotulancora (Lundbeck, 1909)
= *Iotrochota rotulancora*
spinosa (Lundbeck, 1905)
= *Iotrochota spinosa*
topsenti (Burton, 1930)
= *Hymenotrocha topsenti*
varidens (Lundbeck, 1905)
= *Iotrochota varidens*
- Family Crambeidae**
- Crambe*
crambe (Schmidt, 1862)
= *Suberites crambe*
tailliezi Vacelet & Boury-Esnault, 1982
tuberosa Maldonado & Benito, 1991
- Discorhabdella*
hindeli Boury-Esnault *et al.*, 1992
tuberosocapitatum (Topsent, 1892)
= *Hymeraphia tuberosocapitatum*
- Family Desmacididae**
- Desmacidon*
adriaticum Sara, 1969
anceps Schmidt, 1874
fruticosum (Montagu, 1818)
= *Spongia fruticosum*
funicatus Schmidt, 1870
peltatus Topsent, 1904
- Family Dendoricellidae**
- Dendoricella*
abyssi (Topsent, 1904)
= *Desmacidon abyssi*
flabelliformis (Hansen, 1885)
= *Myxilla flabelliformis*
rhopalum Lundbeck, 1905
- Family Coelosphaeridae**
- Chaetodoryx*
insinuans (Topsent, 1936)
= *Colectys insinuans*
- Coelosphaera*
appendiculatum (Carter, 1874)
= *Histoderma appendiculatum*
(*Histodermiton*) *cryosi* (Boury-Esnault *et al.*, 1994)
macrosigma (Topsent, 1892)
= *Gellius macrosigma*
phlyctenodes (Carter, 1876)
= *Halichondria phlyctenodes*
physa (Schmidt, 1875)
= *Desmacidon physa*
picoensis Topsent, 1928
tunicata (Schmidt, 1870)
= *Desmacidon tunicata*
- Echinostylinos*
reticulatus Topsent, 1927
- Ectyodoryx*
atlanticus Stephens, 1916, 1921
balanoides Koltun, 1959
clavigera (Levinsen, 1887)
- derjugini* (Breitfuss, 1897)
foliatus (Fristedt, 1887)
= *Hastatus foliatus*
indistinctus (Fristedt, 1887)
= *Hastatus indistinctus*
kovdaicum (Rezvoj, 1932)
loynigi Burton, 1934
multiformis Brøndsted, 1932
olgae Hentschel, 1929
oligacantha Hentschel, 1929
- Forcepia*
assimilis (Lundbeck, 1905)
= *Leptolabis assimilis*
brunnea (Topsent, 1904))
= *F. forcipula* subsp. *brunnea*
bulbosa (Carter, 1876)
= *Halichondria forcipis* subsp. *bulbosa*
covergens (Topsent, 1928)
= *Leptolabis covergens*
dentifera (Topsent, 1928)
= *Clinolabis dentifera*
exilis (Topsent, 1892)
= *Leptosia exilis*
fabricans (Schmidt, 1874)
= *Esperia fabricans*
forcipis (Bowerbank, 1866)
forcipula Topsent, 1904
fragilis Stephens, 1917
groenlandica Fristedt, 1887
imperfecta Topsent, 1904
luciensis (Topsent, 1888)
= *Dendoryx luciensis*
megachela (Maldonado, 1992)
= *Leptolabis megachela*
psammophila (Cabioch, 1968)
= *Ectoforcepia psammophila*
pustula (Fristedt, 1887)
= *Esperia pustula*
thielei Lundbeck, 1905
topsenti Lundbeck, 1905
versatilis Topsent, 1904
- Histodermella*
apuliae Sara, 1969
ingolfi Lundbeck, 1905
- Inflatella*
pellicula Schmidt, 1875
rhodus (Hentschel, 1929)
= *Hymedesmia rhodus*
tubulosa (Topsent, 1904)
viridis (Topsent, 1892)
= *Joyeuxia viridis*
- Lissodendoryx*
amaknakensis (Lambe, 1900)
basispinosa Sara, 1958
caduca (Schmidt, 1868)
(*Anomodoryx*) *cavernosa* (Topsent, 1892)
= *Damiria cavernosa*
certa (Topsent, 1892)
= *Dendoryx certa*
complicata (Hansen, 1885)
corallorhizoides (Fristedt, 1887)
= *Clathria corallorhizoides*
diversichela Lundbeck, 1905
fertilior Topsent, 1904
firma (Lambe, 1900)
fragilis (Fristedt, 1885)
= *Hastatus fragilis*
isodictyalis (Carter, 1882)
isodictyalis subsp. *paucispinosa* Topsent, 1928
laxa De Laubenfels, 1935
lobosa Lundbeck, 1905
lundbecki Topsent, 1913
(*Waldoschmittia*) *mediterranea* (Sara & Siribelli, 1960)
= *Paramyxilla mediterranea*

<i>polymorpha</i> Topsent, 1928	<i>Crellastrina</i>
(<i>Waldoschmittia</i>) <i>schmidtii</i> (Ridley, 1884)	<i>alecto</i> (Topsent, 1898)
<i>simplex</i> (Topsent, 1928)	= <i>Yvesia alecto</i>
<i>sophia</i> (Fristedt, 1887)	<i>Crellomima</i>
= <i>Esperia sophia</i>	<i>derma</i> Hentschel, 1929
<i>stipitata</i> (Arnesen, 1903)	<i>imparidens</i> Rezvoy, 1923
<i>vicina</i> Lundbeck, 1905	<i>incrustans</i> Hentschel, 1929
Family Crellidae	Family Hymedesmiidae
<i>Anisocrella</i>	<i>Hamigera</i>
<i>hymedesmina</i> Topsent, 1927	<i>hamigera</i> (Schmidt, 1862)
<i>Crella</i>	= <i>Cribrella hamigera</i>
<i>akraleitae</i> (Brøndsted, 1932)	<i>stipitata</i> Arnesen, 1903
= <i>Grayella akraleitae</i>	<i>Hemimycale</i>
<i>alba</i> (Vacelet, 1969)	<i>columella</i> (Bowerbank, 1866)
= <i>Pytheas alba</i>	= <i>Desmacidon columella</i>
<i>albula</i> (Bowerbank, 1866)	<i>Herceus</i>
= <i>Halichondria albula</i>	<i>orientalis</i> Koltun, 1959
<i>atra</i> (Topsent, 1892, 1904)	<i>Hymedesmia</i>
= <i>Pytheas atra</i>	<i>acerata</i> (Topsent, 1904)
<i>basispinosa</i> Burton, 1931	<i>aequatus</i> Lundbeck, 1910
<i>carnosa</i> (Topsent, 1904)	<i>baculifera</i> (Topsent, 1901)
= <i>Yvesia carnosa</i>	<i>basispinosa</i> Lundbeck, 1910
<i>digitifera</i> (Lévi, 1959)	<i>biscutella</i> Topsent, 1928
= <i>Pytheas digitifera</i>	<i>bocki</i> Alander, 1942
<i>dispar</i> (Topsent, 1904)	<i>bowerbanki</i> Lundbeck, 1910
= <i>Yvesia dispar</i>	<i>bractea</i> Lundbeck, 1910
<i>donsi</i> Burton, 1931	<i>castanea</i> Sara, 1964
<i>elegans</i> (Schmidt, 1862)	<i>chlorosa</i> Alander, 1942
= <i>Cribrella elegans</i>	<i>clavigera</i> (Levinsen, 1887)
<i>fallax</i> (Topsent, 1892, 1904)	= <i>Esperella clavigera</i>
= <i>Yvesia fallax</i>	<i>consanguinea</i> Lundbeck, 1910
<i>fusifera</i> Sara, 1969	<i>constellata</i> (Topsent, 1904)
<i>gelida</i> Lundbeck, 1910	= <i>Leptosastra constellata</i>
<i>gergensteini</i> (Schwartschevski, 1880)	<i>cordicela</i> Alander, 1942
= <i>Amphilectus gergensteini</i>	<i>coriacea</i> (Fristedt, 1885)
<i>gracilis</i> (Alander, 1942)	= <i>Stylopus coriacea</i>
= <i>Yvesia gracilis</i>	<i>coronula</i> (Bowerbank, 1866)
<i>guernei</i> (Topsent, 1892, 1904)	= <i>Hymeraphia coronula</i>
= <i>Yvesia guernei</i>	<i>crux</i> (Schmidt, 1875)
<i>hansenii</i> (Topsent, 1892, 1904)	= <i>Desmacidon crux</i>
= <i>Yvesia hansenii</i>	<i>curvihela</i> Lundbeck, 1910
<i>hospitalis</i> (Schmidt, 1870)	<i>depressa</i> Topsent, 1928
<i>jaegerskioeldi</i> Alander, 1937	<i>derjugini</i> (Breitfuss, 1897)
<i>linguifera</i> (Topsent, 1892, 1904)	<i>dermata</i> Lundbeck, 1910
= <i>Yvesia linguifera</i>	<i>digitata</i> Lundbeck, 1910
<i>lobata</i> (Arnesen, 1903)	<i>donsi</i> Alander, 1937
= <i>Yvesia lobata</i>	<i>dubia</i> (Lundbeck, 1910)
<i>mamillata</i> (Arnesen, 1903)	<i>ebria</i> Alander, 1937
= <i>Yvesia mamillata</i>	<i>siliifera</i> (Schmidt, 1870 also)
<i>mollior</i> Topsent, 1925	= <i>Desmacidon tunicatum</i>
<i>nodulosa</i> Sara, 1969	<i>flaccida</i> Topsent, 1928
<i>papillosa</i> (Schmidt, 1870)	<i>gisleni</i> Alander, 1942
= <i>Cribrella papillosa</i>	<i>gracilisigma</i> Topsent, 1928
<i>pedunculata</i> (Topsent, 1892)	<i>grandis</i> Lundbeck, 1910
= <i>Yvesia pedunculata</i>	<i>gunhildae</i> Alander, 1942
<i>pertusa</i> (Topsent, 1892, 1904)	<i>gustafsoni</i> Alander, 1942
= <i>Yvesia pertusa</i>	<i>hallmanni</i> Topsent, 1928
<i>polymastia</i> (Thiele, 1903)	<i>helgae</i> Stephens, 1916, 1921
= <i>Grayella polymastia</i>	<i>hibernica</i> Stephens, 1916, 1921
<i>pulvinar</i> (Schmidt, 1868)	<i>inflata</i> (Alander, 1937)
= <i>Myxilla pulvinar</i>	= <i>Hymoxenia inflata</i>
<i>pyrula</i> (Carter, 1876)	<i>inflata</i> Vacelet, 1969
= <i>Cometella pyrula</i>	<i>irregularis</i> Lundbeck, 1910
<i>richardi</i> (Topsent, 1892, 1904)	<i>jecusculum</i> (Bowerbank, 1866)
= <i>Yvesia richardi</i>	= <i>Hymeniacidon jecusculum</i>
<i>ridleyi</i> (Topsent, 1892, 1904)	<i>jugalis</i> Topsent, 1928
= <i>Yvesia ridleyi</i>	<i>koehlerii</i> (Topsent, 1896)
<i>rosea</i> (Topsent, 1892)	<i>lacera</i> Lundbeck, 1910
= <i>Yvesia rosea</i>	<i>lamina</i> Lundbeck, 1910
<i>rubiginosa</i> (Schmidt, 1862)	<i>latrunculoides</i> Lundbeck, 1910
= <i>Myxilla rubiginosa</i>	<i>lenta</i> Descatoire, 1966
<i>schottlanderi</i> Arndt, 1913	<i>levis</i> Lundbeck, 1910
<i>sigmata</i> (Topsent, 1925)	<i>lobichela</i> Topsent, 1928
<i>topsenti</i> (Babić, 1922)	<i>longistylus</i> Lundbeck, 1910
= <i>Grayella topsenti</i>	<i>longurius</i> Lundbeck, 1910

- macrosigma* Lundbeck, 1910
mamillaris (Fristedt, 1885)
= *Hastatus mamillaris*
manubriata (Topsent, 1928)
matthesii Arndt, 1941
mertonii? sensu Boury-Esnault & Lopes, 1985
microchela Alander, 1942
minuta Alander, 1935
mollis Lundbeck, 1910
mucronata (Topsent, 1913)
mucronella Lundbeck, 1910
mutabilis (Topsent, 1904)
mutabilis subsp. *costata* Topsent, 1928
nigrescens (Topsent, 1925)
norvegica Thiele, 1903
nummulus Lundbeck, 1910
obtusata Topsent, 1928
occulta Bowerbank, 1874
odhneri (Alander, 1942)
= *Stylopus odhneri*
omissa Topsent, 1938
oxneri Topsent, 1928
pachychela Topsent, 1928
pana Bowerbank, 1882
paupertas (Bowerbank, 1866)
= *Hymeniacidon paupertas*
peachi Bowerbank, 1882
pennata Brøndsted, 1932
perforata Lundbeck, 1910
pilata Bowerbank, 1882
planca Lundbeck, 1910
planisigma Topsent, 1928
plicata Topsent, 1928
poicilacantha Alander, 1942
primitiva Lundbeck, 1910
procumbens Lundbeck, 1910
proteida (Schmidt, 1875)
proxima Lundbeck, 1910
pugio Lundbeck, 1910
pulposa (Topsent, 1925)
rhabigena (Topsent, 1904)
= *Leptosia rhabigena*
rissoi (Topsent, 1936)
rugosa Lundbeck, 1910
senegalensis Lévi, 1952
serrulata Vacelet, 1969
simillima Lundbeck, 1910
similis Lundbeck, 1910
sirventi (Topsent, 1928)
= *Leptosia sirventi*
spinosa Stephens, 1916, 1921
splenium Lundbeck, 1910
stephensi (Burton, 1930)
= *Ectyndoryx stephensi*
storea Lundbeck, 1910
stylata Lundbeck, 1910
stylifer (Alander, 1942)
= *Stylopus stylifer*
tenuicula Lundbeck, 1910
tenulsigma Lundbeck, 1910
thielei Alander, 1942
trichoma Lundbeck, 1910
truncata Lundbeck, 1910
umbellifera (Topsent, 1904)
veneta (Schmidt, 1862)
versicolor (Topsent, 1893)
verrucosa Lundbeck, 1910
vidua (Schmidt, 1875)
= *Spirastrella vidua*
vomerula Topsent, 1928
zettlandica Bowerbank, 1866
- Phorbas*
arnesenii Topsent, 1913
baffini (Brøndsted, 1933)
= *Stylostichon baffini*
- bihamigerum* (Waller, 1878)
demonstrans (Topsent, 1892)
= *Suberotelites demonstrans*
dendyi (Topsent, 1892)
= *Stylostichon dendyi*
dives (Topsent, 1891)
= *Microciona dives*
equiosculatum (Pansini, 1987)
= *Stylostichon equiosculatum*
fibulatum Topsent, 1893
fictitius (Bowerbank, 1866)
= *Microciona fictitius*
kovdaicum (Rezvoi, 1932)
lieberkuehni (Burton, 1930)
mercator (Schmidt, 1868)
microchelifer (Cabioch, 1968)
= *Pronax microchelifer*
microcionides (Carter, 1876)
= *Plumohalichondria microcionides*
papillata (Arnesen, 1903)
= *Plumohalichondria papillata*
paupertas? sensu Boury-Esnault, 1971
perarmatus (Bowerbank, 1866)
= *Hymeniacidon perarmatus*
plumosus (Montagu, 1818)
= *Spongia plumosus*
posidoni Voultiadou & Van Soest, 1991
repens (Topsent, 1904)
= *Pronax repens*
roemeri (Hentschel, 1929)
= *Anchinoe roemeri*
aff. stylata Burton, 1930
tenacior (Topsent, 1925)
Plocamionida
ambigua (Bowerbank, 1866)
= *Microciona ambigua*
Pseudohalichondria
clavatancora (Topsent, 1928)
= *Acanthancora clavatancora*
schmidti (Topsent, 1898)
= *Leptosia schmidti*
Spanioplton
armaturum (Bowerbank, 1866)
= *Hymeniacidon armaturum*
fertile Topsent, 1892, 1904
osculosum (Topsent, 1892)
= *Amphilectus osculosum*
- Suborder Mycalina**
- Family Mycalidae**
- Esperiopsis*
artemisiae Brøndsted, 1932
columnata Topsent, 1904
decora Topsent, 1904
digitata Miklugo-Maclay, 1870
forcipula Lundbeck, 1905
fucorum (Esper, 1794)
glomeris Topsent, 1904
incognita Stephens, 1916, 1921
lobata (Montagu, 1818)
= *Spongia lobata*
macrosigma Stephens, 1916, 1921
pedicellata Lundbeck, 1905
polymorpha Topsent, 1892, 1904
praedita Topsent, 1892, 1904
rigida Lambe, 1900
schmidti Arnesen, 1903
strongylatus (Alander, 1942)
= *Amphilectus strongylatus*
strongylophora Vacelet, 1969
typicella Lundbeck, 1909
villosa (Carter, 1874)
= *Esperia villosa*
Mycale
adhaerens (Lambe, 1893)
adhaerens subsp. *fibrosa* Koltun, 1959

- albicans* (Topsent, 1892)
 = *Esperella albicans*
arctica (Fristedt, 1887)
 = *Esperella lingua*
(Raphidiotheca) arctica Hentschel, 1929
(Paresperella) atlantica Stephens, 1917
bolivari Ferrer-Hernandez, 1914
borassus (Carter, 1876)
 = *Esperia borassus*
bowerbanki (Schmidt, 1862)
 = *Esperia bowerbanki*
contareni (Martens, 1824)
dentata Sara, 1958
dubia (Czerniavsky, 1880)
 = *Esperia dubia*
fascibula (Topsent, 1904)
foraminosa (Schmidt, 1862)
 = *Esperia foraminosa*
hamata (Topsent, 1892)
 = *Esperella hamata*
helios (Fristedt, 1887)
 = *Esperia helios*
(Oxymycale) intermedia (Schmidt, 1874)
 = *Esperia intermedia*
irregularis (Czerniavsky, 1880)
 = *Esperia irregularis*
lanugo (Schmidt, 1875)
 = *Esperia lanugo*
lindbergi Koltun, 1958, 1959
lingua (Bowerbank, 1866)
 = *Raphiodesma lingua*
lobimana (Czerniavsky, 1880)
 = *Protoesperia lobimana*
longistyla Koltun, 1958, 1959
(Gomphostegia) loricata Topsent, 1896, 1904
loveni (Esper, 1887)
 = *Esperia loveni*
macilenta (Bowerbank, 1866)
 = *Hymeniacidon floreum*
 = *Hymeniacidon macilenta*
marshallhalli (Kent, 1870)
 = *Raphidiotheca marshallhalli*
massa (Schmidt, 1862)
 = *Esperia massa*
micracanthoxea Buizer & Van Soest, 1977
minima (Waller, 1880)
 = *Raphiodesma minima*
mollis (Topsent, 1892)
 = *Esperella mollis*
muscoidea (Czerniavsky, 1880)
 = *Esperia muscoidea*
papillosa Koltun, 1959
papillosa subsp. *dulkeiti* Koltun, 1959
papillosa subsp. *ochotensis* Koltun, 1959
retifera Topsent, 1924
rotalis (Bowerbank, 1874)
 = *Desmacidon rotalis*
sanguinea Tsuruhashi, 1969
serrulata Sara & Sribelli, 1960
similaris (Bowerbank, 1874)
 = *Raphiodesma similaris*
 = *Raphiodesma sordida*
strelnicovi Rezvoi, 1932
syrinx (Schmidt, 1862)
 = *Esperia lorenzi*
 = *Esperia syrinx*
titubans (Schmidt, 1870)
 = *Desmacidon titubans*
toporoki Koltun, 1958, 1959
tuberosa (Schmidt, 1868)
 = *Esperia tuberosa*
tunicata (Schmidt, 1862)
 = *Esperia tunicata*
tylota Koltun, 1958, 1959
varpachowski Schwartshevski, 1905
- velutata* (Schmidt, 1862)
 = *Esperia velutata*
vosmaeri (Levinsen, 1886)
 = *Esperella vosmaeri*
Phlyctaenopora
bitorquis Topsent, 1904
Ulosa
stuposa (Esper, 1794)
 = *Spongia stuposa*
tenellula Pulitzer-Finali, 1983
tubulata Pulitzer-Finali, 1983
- Family Cladorhizidae**
- Asbestopluma*
bihamatifera (Carter, 1876)
 = *Esperia cupressiformis* subsp. *bihamatifera*
cupressiformis (Carter, 1874)
 = *Esperia cupressiformis*
furcata Lundbeck, 1905
gracilis Koltun, 1955
hydra Lundbeck, 1905
hypogea Vacelet & Boury-Esnault, 1996
infundibulum (Levinsen, 1886)
lycopodium (Levinsen, 1886)
minuta (Lambe, 1900)
pennatula (Schmidt, 1875)
 = *Cladorhiza pennatula*
quadriserialis Tendal, 1973
ramosa Koltun, 1958
versatilis Topsent, 1904
- Chondrocladia*
alaskensis Lambe, 1900
burtoni Tendal, 1973
gigantea (Hansen, 1880)
pulchra Lambe, 1900
virgata Thomson, 1872
- Cladorhiza*
abyssicola Sars, 1872
arctica Burton, 1946
bathycrinoides Koltun, 1955
corticocancellata Carter, 1876
 = *C. abyssicola* subsp. *corticocancellata*
gelida Lundbeck, 1905
nobilis Fristedt, 1887
nordenskioldi Fristedt, 1887
ocea Lundbeck, 1905
tenuisigma Lundbeck, 1905
- Euchelipluma*
pristina Topsent, 1919
- Family Desmacellidae**
- Bienna*
dautzenbergi Topsent, 1892, 1904
gemmulifera Breitfuss, 1897
parthenopea Pulitzer-Finali, 1978
rosea (Fristedt, 1887)
 = *Desmacella rosea*
tenuisigma Pulitzer-Finali, 1978
variantia (Bowerbank, 1858)
 = *Desmacella variantia*
variantia subsp. *peracuta* (Topsent, 1927, 1928)
- Desmacella*
annexa Schmidt, 1870, 1875
campechiiana (Topsent, 1889)
corrugata (Bowerbank, 1866)
grimaldi (Topsent, 1892, 1904)
 = *Bienna grimaldi*
groenlandica Fristedt, 1887
 = *D. peachi* subsp. *groenlandica*
hamifera Lundbeck, 1909
informis Stephens, 1916, 1921
inornata (Bowerbank, 1866)
 = *Halichondria inornata*
meliorata Wiedenmayer, 1978
peachi sensu Ferrer-Hernandez, 1914
pumilio Schmidt, 1870
suberea (Schmidt, 1870)

- = *Desmacodes suberea*
vagabunda Schmidt, 1870
velutata (Renier)
= *Spongia velutata*
Dragmatella
aberrans Topsent, 1892
- Family Hamacanthidae**
Hamacantha
carteri Topsent, 1904
falcula (Bowerbank, 1866)
= *Halichondria falcula*
integra Topsent, 1904
johsoni (Bowerbank, 1864)
johsoni subsp. *complanata* Topsent, 1892
johsoni subsp. *inelegans* Topsent, 1892
lundbecki Topsent, 1904
meganistra Pulitzer-Finali, 1978
papillata Vosmaer, 1885
schmidti (Carter, 1874)
- Merlia*
normani/deficiens Kirkpatrick, 1908
- Pozziella*
clavisaepa Topsent, 1896, 1904
- Family Guitarridae**
Guitarra
imbriata Carter, 1874
laplani Boury-Esnault *et al.*, 1993
solorzanoi Cristobo, 1998
voluta Topsent, 1904
- Family Isodictyidae**
Isodictya
histodermella De Laubenfels, 1942
palmata (Ellis & Solander, 1786)
= *Spongia palmata*
- Order Haplosclerida**
- Family Petrosiidae**
Petrosia
clavata (Esper, 1794)
= *Spongia clavata*
crassa (Carter, 1876)
= *Reniera crassa*
ficiformis (Poiret, 1798)
intermedia (Czerniavsky, 1880)
= *Schmidtia intermedia*
raphida Boury-Esnault *et al.*, 1994
vanoestii Boury-Esnault *et al.*, 1994
- Xestospongia*
friabilis (Topsent, 1892)
= *Petrosia friabilis*
magna (Vacelet, 1969)
= *Reniera magna*
plana (Topsent, 1892)
= *Reniera plana*
variabilis (*sensu* Topsent, 1892)
= *Petrosia variabilis*
- Family Phloeodictyidae**
Aka
infesta (Johnson, 1889)
- Calyx*
nicaensis (Risso, 1826)
tufa (Ridley & Dendy, 1886)
- Oceanapia*
aereum (Topsent, 1904)
constructa (Rützler, 1965)
= *Phloeodictyon constructa*
coriacea (Topsent, 1904)
decipiens (Sara, 1958, 1964)
= *Phloeodictyon decipiens*
elongata (Topsent, 1892, 1904)
= *Rhizochalina elongata*
fistulosa (Bowerbank, 1873)
isodictyiformis (Carter, 1882)
= *Phloeodictyon isodictyiformis*
- minor* (Sarà, 1958)
= *Phloeodictyon minor*
nodulosa (Topsent, 1927)
perforata (Sarà, 1960)
putridosa (Lamarck, 1813)
= *Spongia putridosa*
reticulata Topsent, 1904
robusta (Bowerbank, 1866)
= *Isodictya robusta*
tuber (Lundbeck, 1909)
vacua (Sarà, 1961)
= *Phloeodictyon vacua*
- Family Callyspongiidae**
Callyspongia
septimaniensis Griessinger, 1971
subcornea Griessinger, 1971
- Chalinopsis*
tuba Von Lendenfeld, 1889
- Siphonochalina*
balearica Ferrer-Hernandez, 1916
coriacea Schmidt, 1868
expansa Sarà, 1960
- Family Niphatidae**
Gelliodes
bifacialis Topsent, 1904
fayalensis Topsent, 1892
- Hemigellius*
arciferus (Vosmaer, 1885)
= *Gellius arciferus*
hartlaubi (Hentschel, 1929)
= *Gellius hartlaubi*
plexus (Lundbeck, 1909)
= *Gelliodes plexus*
porosus (Fristedt, 1887)
= *Desmacella porosus*
pumiceus (Fristedt, 1885)
= *Desmacella pumiceus*
styliferus (Von Lendenfeld, 1897)
= *Gellius styliferus*
- Pachychalina*
arctica Miklucho-Maclay
fracta Hentschel, 1929
gracilenta Ferrer-Hernandez, 1914
gracilis Miklucho-Maclay
rustica Schmidt, 1868
schmidti Lundbeck, 1902
- Family Chalinidae**
Chalinula
fertilis Keller, 1880
limbata (Montagu, 1818)
= *Spongia limbata*
loosanoffi (Hartman, 1958)
= *Haliclona loosanoffi*
nigra Boury-Esnault & Lopes, 1985
renieroides (Schmidt, 1868)
- Cladocroce*
fibrosa Topsent, 1892, 1904
spathiformis Topsent, 1904
- Dendroxea*
adumbrata Corriero *et al.*, 1996
lenis (Topsent, 1892)
= *Reniera lenis*
- Haliclona*
accomodata (Schmidt, 1868)
= *Reniera accomodata*
alba (Schmidt, 1862)
= *Reniera alba*
albaponica (Czerniavsky, 1880)
= *Reniera albaponica*
angulata (Bowerbank, 1866)
= *Halichondria angulata*
aperta (Sarà, 1960)
= *Reniera aperta*
aquaeductus (Schmidt, 1862)

= <i>Reniera aqueductus</i>	= <i>Adocia laevis</i>
<i>aquaeductus</i> subsp. <i>taurica</i> (Czerniavsky, 1880)	<i>laurentina</i> (Lambe, 1900)
<i>arctica</i> (Fristedt, 1887)	<i>laxa</i> (Topsent, 1892)
= <i>Reniera arctica</i>	= <i>Rhaphisia laxa</i>
<i>arctica</i> (Hentschel, 1916)	<i>mamillata</i> (Griessinger, 1971)
= <i>Gellius arctica</i>	= <i>Reniera mamillata</i>
<i>arenata</i> Griessinger, 1971	<i>marismedi</i> (Pulitzer-Finali, 1978)
<i>binaria</i> (Topsent, 1927)	= <i>Gellius marismedi</i>
= <i>Gellius binaria</i>	<i>mediterranea</i> Griessinger, 1971
<i>bioxeata</i> Boury-Esnault et al., 1994	<i>membranacea</i> (Schmidt, 1868)
<i>bowerbanki</i> Ferrer-Hernandez	= <i>Reniera membranacea</i>
<i>bulbifera</i> (Schwartzchevski, 1905)	<i>merejkowskii</i> Schwartzchevski, 1905
<i>calanus</i> Lundbeck, 1909	<i>microsigma</i> (Babic, 1922)
<i>calyx</i> (Ridley & Dendy, 1886)	= <i>Gellius microsigma</i>
= <i>Gellius calyx</i>	<i>microtoxa</i> (Lundbeck, 1909)
<i>caspia</i> (Grimm, 1885)	<i>microxifer</i> (Topsent, 1925)
<i>cinerea</i> (Grant, 1826)	= <i>Reniera microxifer</i>
= <i>Spongia cinerea</i>	<i>mollicula</i> (Lundbeck, 1902)
<i>citrina</i> (Topsent, 1892)	<i>mollis</i> (Schmidt, 1868)
= <i>Reniera citrina</i>	= <i>Siphonochalina mollis</i>
<i>clavata</i> (Levinson, 1886)	<i>mollis</i> (Lambe, 1900)
= <i>Reniera clavata</i>	<i>mucosa</i> (Griessinger, 1971)
<i>columbae</i> (Walker, 18xx)	= <i>Reniera mucosa</i>
= <i>Spongia columbae</i>	<i>neens</i> (Topsent, 1918)
<i>cratera</i> (Schmidt, 1862)	<i>nigricans</i> (Czerniavsky, 1880)
= <i>Reniera cratera</i>	= <i>Reniera nigricans</i>
<i>cibrata</i> (Pulitzer-Finali, 1983)	<i>oculata</i> (Pallas, 1766)
= <i>Adocia cibrata</i>	= <i>Spongia oculata</i>
<i>cribrosa</i> (Czerniavsky, 1880)	<i>omissa</i> (Griessinger, 1971)
= <i>Pellinula cribrosa</i>	= <i>Reniera omissa</i>
<i>cucurbitiformis</i> (Kirkpatrick, 1907)	<i>palmata</i> (?Ellis & Solander, 1791)
<i>cylindrigera</i> (Czerniavsky, 1880)	= <i>Spongia palmata</i>
= <i>Tedaniella cylindrigera</i>	<i>palmensis</i> Carballo & Garcia-Gomez, 1995
<i>digitata</i> subsp. <i>pontica</i> (Czerniavsky, 1880)	<i>papillifera</i> (Schwartzchevski, 1905)
<i>dubia</i> (Babic, 1922)	<i>parasitica</i> (Bowerbank, 1866)
<i>esperi</i> (Arnesen, 1903)	= <i>Isodictya parasitica</i>
= <i>Gellius esperi</i>	<i>parenchyma</i> Lundbeck, 1902
<i>fibulata</i> (Schmidt, 1862)	<i>parietalis</i> (Topsent, 1893)
= <i>Reniera fibulata</i>	= <i>Reniera parietalis</i>
<i>fistulosa</i> (Bowerbank, 1866)	<i>perlucida</i> (Griessinger, 1971)
= <i>Isodictya fistulosa</i>	= <i>Reniera perlucida</i>
<i>fistulosa</i> subsp. <i>latens</i> (Topsent, 1892)	<i>pocilliformis</i> (Griessinger, 1971)
<i>flagellifera</i> (Ridley & Dendy, 1886)	= <i>Reniera pocilliformis</i>
= <i>Gellius flagellifera</i>	<i>poecillastroides</i> (Vacelet, 1969)
<i>flavescens</i> (Topsent, 1893)	= <i>Reniera poecillastroides</i>
<i>folium</i> (Schmidt, 1870)	<i>pons</i> (Schmidt, 1870)
= <i>Chalinula folium</i>	= <i>Reniera pons</i>
<i>foraminosa</i> (Czerniavsky, 1880)	<i>primitiva</i> (Lundbeck, 1902)
= <i>Protoschmidia foraminosa</i>	<i>protochalina</i> (Czerniavsky, 1880)
<i>fulva</i> (Topsent, 1893)	= <i>Amorphina protochalina</i>
= <i>Reniera fulva</i>	<i>proxima</i> (Lundbeck, 1909)
<i>glacialis</i> Hentschel, 1916	<i>pulcherrima</i> (Fristedt, 1885)
<i>gracilis</i> (Miklugo-Maclay)	<i>pulchra</i> (Schwartzchevski, 1905)
<i>griessingeri</i> Van Lent & De Weerdt, 1987	<i>pyrrhi</i> (Hanitsch, 1895)
<i>grimmi</i> (Czerniavsky, 1880)	= <i>Gellius pyrrhi</i>
= <i>Protoschmidia grimmii</i>	<i>rava</i> (Stephens, 1912, 1917)
<i>groenlandica</i> (Fristedt, 1887)	= <i>Gellius rava</i>
= <i>Chalina groenlandica</i>	<i>rectangularis</i> (Ridley & Dendy, 1886)
<i>grossa</i> (Schmidt, 1864)	<i>reptans</i> (Griessinger, 1971)
= <i>Reniera grossa</i>	= <i>Adocia reptans</i>
<i>hirsuta</i> Schwartzchevski, 1905	<i>rhopidiophora</i> (Brøndsted, 1933)
<i>hyalina</i> (Schmidt, 1873)	= <i>Gellius rhopidiophora</i>
= <i>Esperia hyalina</i>	<i>rosea</i> (Bowerbank, 1866)
<i>implexa</i> (Schmidt, 1868)	= <i>Isodictya rosea</i>
= <i>Reniera implexa</i>	<i>rossica</i> (Hentschel, 1929)
<i>indistincta</i> (Bowerbank, 1866)	= <i>Reniera rossica</i>
= <i>Isodictya indistincta</i>	<i>sarai</i> (Pulitzer-Finali, 1969)
<i>inflata</i> (Schmidt, 1862)	= <i>Reniera sarai</i>
= <i>Reniera inflata</i>	<i>schmidti</i> (Czerniavsky, 1880)
<i>inflata</i> subsp. <i>taurica</i> (Czerniavsky, 1880)	= <i>Pellinula schmidti</i>
<i>informis</i> subsp. <i>taurica</i> (Czerniavsky, 1880)	<i>semitubulosa</i> sensu Lieberkühn, 1859
<i>irregularis</i> (Czerniavsky, 1880)	= <i>Halichondria semitubulosa</i>
= <i>Cacochalina irregularis</i>	<i>simplex</i> (Czerniavsky, 1880)
<i>lacazei</i> (Topsent, 1892)	= <i>Protoschmidia simplex</i>
<i>laevis</i> (Griessinger, 1971)	<i>simulans</i> (Johnston, 1842)

= *Halichondria simulans*
sоловецкая (Hentschel, 1929)
= *Reniera solowetzkaja*
spatula (Lundbeck, 1902)
= *Chalina spatula*
spissa (Topsent, 1892)
= *Thrinacophora spissa*
spitzbergensis Hentschel, 1916
spongiosa (Topsent, 1904)
steueri Burton, 1936
stirpescens (Topsent, 1925)
= *Reniera stirpescens*
subtilis Griessinger, 1971
tenera (Marenzeller)
tenuisigma (Sarà & Sibelli, 1960)
= *Gellius tenuisigma*
transitans (Czerniavsky, 1880)
= *Protoschmidtia transitans*
tromsoica (Hentschel, 1929)
= *Reniera tromsoica*
uncinata (Topsent, 1892)
= *Gellius uncinata*
urceola (Rathke & Vahl, 1806)
= *Spongia urceola*
utricula (Topsent, 1904)
valliculata Griessinger, 1971
varia (Bowerbank, 1875)
varia (Sarà, 1958)
= *Adocia varia*
variatabulata (Sarà, 1958)
= *Adocia variatabulata*
vega (Fristedt, 1887)
= *Chalina vega*
venata (Sarà, 1960)
= *Adocia venata*
ventilabrum (Fristedt, 1887)
= *Reniera ventilabrum*
viridis (Keller, 1881)
viscosa (Topsent, 1887)
= *Reniera viscosa*
voeringii Lundbeck, 1909
xena De Weerdt, 1986
Metschnikowia
intermedia Grimm, 1885
tuberculata Grimm, 1885
Rhizoniera
pedunculata Boury-Esnault *et al.*, 1994
rhizophora (Vacelet, 1969)
= *Reniera rhizophora*

Order Dictyoceratida

Family Spongiidae
Coscinoderma
confragosum Polejaeff, 1884
sporadense Voultsiadou-Koukoura *et al.*, 1991
Hippospongia
communis (Lamarck, 1814)
= *Spongia communis*
Hyattella
intestinalis (Lamarck, 1814)
= *Spongia intestinalis*
Spongia
agaricina Pallas, 1766
aurea Montagu, 1818
conus Montagu, 1818
irregularis subsp. *mollior* (Von Lendenfeld, 1889)
laevigata Montagu, 1818
niolissima Schmidt, 1862
nitens (Schmidt, 1862)
= *Ditela nitens*
officinalis Linnaeus, 1759
officinalis subsp. *adriatica* Schmidt, 1862
quarnerensis Schmidt, 1862
virgulosa Schmidt, 1868
zimocca Schmidt, 1862

Family Thorectidae
Aplysinopsis
schnidi (Marenzeller, 1877)
Cacospongia
carduelis Schmidt, 1864
mollior Schmidt, 1862
proficeps Pulitzer-Finali & Pronzato, 1980
scalaris Schmidt, 1862
Fasciospongia
asperillum (Von Lendenfeld, 1889)
= *Stenospongia asperillum*
cavernosa (Schmidt, 1862)
= *Cacospongia cavernosa*
coerulea Vacelet, 1959
Hyrtios
collectrix (Schulze, 1879)
= *Oligoceras collectrix*
erecta (Keller, 1889)
= *Heteronema erecta*
dendyi (Ferrer-Hernandez, 1922)
= *Collospongia dendyi*
Thorectandra
diversispiculifera Ferrer-Hernandez, 1918
Family Irciniidae
Ircinia
chevreuxi (Topsent, 1894)
= *Hircinia chevreuxi*
collectrix (Von Lendenfeld, 1889)
= *Hircinia collectrix*
condensa (Topsent, 1891)
= *Hircinia condensa*
dendroides Schmidt, 1862
digitata (Topsent, 1894)
= *Hircinia digitata*
fasciculata (Esper, 1794)
= *Spongia fasciculata*
flavescens Schmidt, 1862
foetida (Schmidt, 1862)
= *Sarcotragus foetida*
hebes Schmidt, 1862
hirsuta Schmidt, 1862
lingua Schmidt, 1868
muscarum (Schmidt, 1862)
= *Sarcotragus muscarum*
oros (Schmidt, 1864)
= *Hircinia oros*
panicea Schmidt, 1862
paucifilamentosa Vacelet, 1961
pipetta (Schmidt, 1868)
procumbens (Polejaeff, 1884)
= *Cacospongia procumbens*
retidermata Pulitzer-Finali & Pronzato, 1980
spinosa (Schmidt, 1862)
= *Sarcotragus spinosula*
stipitata (Topsent, 1894)
= *Hircinia stipitata*
variabilis (Pallas, 1766)
= *Spongia variabilis*
variabilis subsp. *fistulata* Szymanski, 1904
verrucosa (Ferrer-Hernandez, 1914)
= *Hircinia verrucosa*
vestibulata Szymanski, 1904

Order Dendroceratida

Family Dysideidae
Dysidea
avara (Schmidt, 1862)
= *Spongelia avara*
elastica (Von Lendenfeld, 1889)
= *Spongelia elastica*
elegans subsp. *pontica* (Czerniavsky, 1880)
fragilis (Montagu, 1818)
= *Spongelia fragilis*
incrassata (Schmidt, 1862)
= *Spongelia incrassata*

- incrustans* (Czerniavsky, 1880)
 = *Spongelia incrustans*
pallescens (Schmidt, 1862)
 = *Spongelia pallescens*
pallescens subsp. *pontica* (Czerniavsky, 1880)
papillosa Johnston, 1842
perfistulata Pulitzer-Finali & Pronzato, 1980
tupha (Martens, 1824)
 = *Spongia tupha*
- Spongionella*
gracilis (Vosmaer, 1883)
pulchella (Sowerby, 1804)
 = *Spongia pulchella*
ramodigitata (Topsent, 1901)
repens (Topsent, 1928)
zellandica (Bowerbank, 1866)
 = *Verongia zellandica*
- Family Darwinellidae**
- Aplysilla*
glacialis (Merejkowski, 1877)
rosea (Barrois, 1876)
sulfurea Schulze, 1878
- Chelonaplysilla*
arenosa Hentschel, 1929
erecta (Row, 1911)
noevus (Carter, 1874)
 = *Aplysina noevus*
psammophila (Topsent, 1928)
- Darwinella*
aurea (Von Lendenfeld, 1889)
australiensis Carter, 1885
corneostellata (Carter, 1872)
 = *Aplysina corneostellata*
dalmatica Topsent, 1905
gardineri Topsent, 1905
duplex Topsent, 1905
intermedia Topsent, 1893
simplex Topsent, 1892
viscosa Boury-Esnault, 1971
- Dendrilla*
acantha Vacelet, 1958
cirsoides Topsent, 1893
- Hexadella*
detritifera Topsent, 1904
racovitzai Topsent, 1896
- Pleraplysilla*
minchini Topsent, 1905
spinifera (Schulze, 1878)
 = *Aplysilla spinifera*
- Psammopemma*
commune Carter, 1876
finmarchicum Hentschel, 1929
finmarchicum subsp. *spiculosum* Hentschel, 1929
nicaense Pulitzer-Finali & Pronzato, 1980
- Order Halisarcida**
- Family Halisarcidae**
- Halisarca*
dujardini Johnston, 1842
guttula Schmidt, 1864
metschnikowi Lévi, 1953
sputum Topsent, 1893
- Myceliospongia*
araneosa Vacelet & Perez, 1998?
- Order Verongida**
- Family Aplysinidae**
- Aplysina*
aerophoba Nardo, 1843
carnosa Schmidt, 1862
cavernicola (Vacelet, 1959)
 = *Verongia cavernicola*
massa (Szymanski, 1904)
 = *Aplysinopsis massa*
tuberosa (Szymanski, 1904)
- = *Aplysinopsis tuberosa*
- Class Calcarea**
- Subclass Calcinea**
- Order Clathrinida**
- Family Clathrinidae**
- Clathrina*
ascandroides Borojevic, 1971
biscayae Borojevic & Boury-Esnault, 1987
canariensis (Miklucho-Maclay, 1870)
 = *Leucosolenia canariensis*
cancellata (Verrill, 1873)
 = *Leucosolenia cancellata*
cerebrum (Haeckel, 1870)
 = *Ascalitis cerebrum*
clathrus (Schmidt, 1864)
 = *Grantia clathrus*
contorta (Bowerbank, 1866)
 = *Leucosolenia contorta*
olynthus Borojevic & Boury-Esnault, 1987
coriacea (Montagu, 1818)
 = *Spongia coriacea*
multiformis (Breitfuss, 1898)
 = *Leucosolenia multiformis*
nansenii (Breitfuss, 1896)
 = *Leucosolenia nansenii*
reticulum (Schmidt, 1862)
 = *Nardoa reticulum*
rubra Sarà, 1958
 = *C. coriacea* var. *rubra*
- Guancha*
blanca Miklucho-Maclay, 1868
lacunosa (Bean in Johnston, 1842)
 = *Grantia lacunosa*
- Leucetta*
nausicaa (Schuffner, 1877)
solida (Schmidt, 1862)
- Family Leucaltidae**
- Ascandra*
falcata Haeckel, 1870
- Leucaltis*
clathria Haeckel, 1870
- Leuclathrina*
asconoides Borojevic & Boury-Esnault, 1987
- Family Leucascidae**
- Ascalitis*
lamarcki Haeckel, 1870
- Family Levinellidae**
- Levinella*
thalassae Borojevic & Boury-Esnault, 1987
- Subclass Calcaronea**
- Order Leucosolenida**
- Family Amorphiscidae**
- Amorphiscus*
buccichi Von Ebner, 1887
cylindricus (Haeckel, 1870)
gregorii (Von Lendenfeld, 1891)
oviparus (Haeckel, 1870)
 = *Sycalitis oviparus*
 = *Sycurus oviparus*
- Leucilla*
echinus (Haeckel, 1870)
 = *Leuculmis echinus*
- Family Grantiidae**
- Amphiute*
paulini Hanitsch, 1894
- Aphroceras*
ensata (Bowerbank, 1858)
 = *Grantia ensata*
corticata (Von Lendenfeld, 1891)
- Ebnerella*
schulzei Breitfuss, 1896

- Grantia**
- canadensis* Lambe, 1896
 - capillosa* (Schmidt, 1862)
 - = *Ute capillosa*
 - compressa* (Fabricius, 1780)
 - = *Spongia compressa*
 - singularis* (Breitfuss, 1896)
 - = *Sphenopora singularis*
- Leuconia**
- ananas* (Montagu, 1818)
 - = *Spongia ananas*
 - aspera* (Schmidt, 1862)
 - = *Sycon aspera*
 - balearica* (Lakschewitz, 1886)
 - = *Leucandra balearica*
 - bolivari* Ferrer-Hernandez, 1916
 - bulbosa* (Hanitsch, 1895)
 - = *Leucandra bulbosa*
 - caespitosa* (Haeckel, 1870)
 - caminus* (Haeckel, 1872)
 - fistulosa* (Johnston, 1842)
 - = *Grantia fistulosa*
 - gossei* (Bowerbank, 1862, 1864, 1866, 1874, 1882)
 - = *Leucogypsa gossei*
 - johnstoni* Carter, 1871
 - nivea* (Grant, 1826)
 - = *Leucandra nivea*
 - polejaevi* (Breitfuss, 1896)
 - = *Pericharax polejaevi*
 - pumila* Bowerbank, 1866
 - rodriguezi* (Lakschewitz, 1886)
 - = *Leucandra rodriguezi*
- Scyssa**
- huxleyi* Haeckel, 1870
- Trichogypsia**
- villosa* Carter, 1871
- Ute**
- gladiata* Borojevic, 1967
- Family Heteropiidæ**
- Heteropia**
- ramosa* (Carter, 1886 in Higgin)
 - = *Aphroceras ramosa*
- Family Leucosoleniidæ**
- Ascyssa**
- acufera* Haeckel, 1870
 - troglodytes* Haeckel, 1870
- Leucosolenia**
- botryoides* (Ellis & Solander, 1786)
 - = *Spongia botryoides*
 - complicata* (Montagu, 1818)
 - = *Spongia complicata*
 - darvini* (Haeckel, 1870)
 - echinoides* (Haeckel, 1870)
 - fragilis* (Haeckel, 1870)
 - = *Ascortis fragilis*
 - gegenbauri* (Haeckel, 1870)
 - goethei* (Haeckel, 1870)
 - hermesi* (Haeckel, 1870)
 - lieberkuehni* (Schmidt, 1862)
 - minoricensis* Lakschewitz, 1886
 - primordialis* (Haeckel, 1870)
 - variabilis* (Haeckel, 1870)
 - = *Ascandra variabilis*
- Family Sycettidae**
- Grantessa**
- glacialis* (Haeckel, 1870)
 - = *Syclitis glacialis*
 - kuekenthali* (Breitfuss, 1896)
 - = *Ebnerella kuekenthali*
 - lanceolata* (Breitfuss, 1898)
 - = *Ebnerella lanceolata*
 - murmanensis* (Breitfuss, 1898)
 - = *Amphoriscus murmanensis*
 - nitida* (Arnesen, 1900)
- = *Ebnerella nitida*
- Sycandra**
- utriculus* (Schmidt, 1869)
 - = *Ute utriculus*
- Sycetta**
- asconoides* Breitfuss, 1896
 - confusa* (Haeckel, 1870)
- Sycodorus**
- hystrix* Haeckel, 1870
- Sycon**
- abyssale* Borojevic & Graat-Kleeton, 1965
 - arcticum* (Haeckel, 1870)
 - = *Sycom arcticum*
 - boreale* (Schuffner, 1877)
 - ciliatum* (Fabricius, 1780)
 - = *Spongia ciliatum*
 - elegans* (Bowerbank, 1845)
 - = *Dunstervillia elegans*
 - helleri* (Von Lendenfeld, 1891)
 - humboldti* Risso, 1826
 - lingua* (Haeckel, 1870)
 - = *Sycoites lingua*
 - quadrangulata* (Schmidt, 1868)
 - = *Syconella quadrangulata*
 - raphanus* Schmidt, 1862
 - scaldiense* (Van Koolwijk, 1982)
 - = *Scypha scaldiense*
 - schmidti* (Haeckel, 1870)
 - schuffneri* Dendy & Row, 1913
 - sycandra* (Von Lendenfeld, 1895)
 - = *Homoderma sycandra*
 - tenellum* (Von Lendenfeld, 1891)
 - tuba* (Von Lendenfeld, 1891)
 - villosum* (Haeckel, 1870)
 - = *Syconda villosum*
- Vosmaeropsis*
- gardineri* Ferrer-Hernandez, 1916
- Order Lithonida**
- Family Minchinellidae**
- Plectroninia**
- norvegica* Könnecker, 1989
- Class Hexactinellida**
- Subclass Amphidiscophora**
- Family Pheronematidae**
- Pheronema**
- carpenteri* (Thomson, 1869)
 - = *Holtenia carpenteri*
- Family Hyalonematidae**
- Hyalonema**
- infundibulum* Topsent, 1896
 - lusitanicum* Du Bocage, 1864
 - thomsoni* Marshall, 1875
- Subclass Hexasterophora**
- Order Hexactinosa**
- Family Farreidae**
- Farrea**
- foliascens* Topsent, 1904
 - laminaris* Topsent, 1904
 - occa* (Bowerbank, 1862)
 - weltneri* Topsent, 1901
- Sarostegia**
- oculata* Topsent, 1904
- Family Euretidae**
- Chonetasma**
- ijimai* Topsent, 1901
- Dactylocalyx**
- potatorum* Schmidt, 1880
- Eurete**
- alicei* Topsent, 1901
- Periphragella**
- lusitanica* Topsent, 1890

Pityrete
azoricum (Topsent, 1904)
 = *Aphrocallistes azoricum*

Family Tretodictyidae

Auloplax
filholi (Topsent, 1904)
Hexactinella
grimaldii Topsent, 1890
Tretodictyum
tubulosum (Schulze, 1886)
 = *Hexactinella tubulosum*

Family Aulocalycidae

Fieldingia
lagettoides Kent, 1870
Rhabdodictyum
delicatum Schmidt, 1880

Family Aphrocallistidae

Aphrocallistes
bearrix Gray, 1858
bocagei Schulze, 1886

Order Lyssacinos

Family Euplectellidae

Euplectella
suberea Thomson, 1877
Hertwiga
falcifera Schmidt, 1880
Malacosaccus
floricomatus Topsent, 1901
unguiculatus Schulze, 1886
Regadrella
phoenix Schmidt, 1880
Rhabdoplectella
tintinnus Schmidt, 1880

Family Caulophacidae

Caulophacus
arcticus (Hansen, 1885)
 = *Hyalonema arcticus*
Sympagella
nux Schmidt, 1870

Family Rossellidae

Anoxycalyx
laceratus Koltun, 1967
Asconema
setubalense Kent, 1870
Bathydorus
servatus Topsent, 1927, 1928
Doconesthes
sessilis Topsent, 1928
Lanuginella
pupa Scgmidt, 1870
Mellonympa
velata (Thomson, 1873)
 = *Rossella velata*
Rossella
nodastrella Topsent, 1915
Schaudinnia
arctica Schulze, 1900
rosea (Fristedt, 1887)
Scyphidium
septentrionale Schulze, 1900
Trichasterina
borealis Schulze, 1900
sagittaria Topsent, 1913

Family Leucopsacidae

Leucopsacus
scoliodocus subsp. *retroscissus* Topsent, 1903
Oopsacas
minuta Topsent, 1927, 1928

PLACOZOA

Compiled by Jacob van der Land

The European list was derived from URMO.

References

Harrison, F.W. and J.A. Westfall (eds), 1991. *Microscopic anatomy of invertebrates 2: Placozoa, Porifera, Cnidaria and Ctenophora*: i-xiv, 1-436. Wiley-Liss, New York

PHYLUM PLACOZOA

Family Trichoplacidae

<i>Trichoplax</i>	AM
<i>adhærens</i> Schulze, 1883	
<i>Treptoplas</i>	
<i>reptans</i> Monticelli, 1896	M

MESOZOA

Compiled by Jacob van der Land and J. Hallan

The European list was derived from URMO with names provided by J. Hallan (Austin, Texas).

PHYLUM MESOZOA

Class Rhombozoa

Family Dicyemidae

<i>Dicyema</i>	
<i>macrocephalum</i> (van Beneden, 1878)	M
<i>megalcephalum</i> Nouvel, 1934	A
<i>microcephalum</i> Whitman, 1883	M
<i>monodi</i> Nouvel, 1934	A
<i>moschatum</i> Whitman, 1883	M
<i>paradoxum</i> von Kölliker, 1849	AM
<i>rondeletiolae</i> Nouvel, 1944	M
<i>schulzianum</i> van Beneden, 1876	M
<i>typus</i> van Beneden, 1876	AM
<i>Dicyemennaea</i>	
<i>eleodenes</i> (Wagener, 1857)	AM
<i>gracile</i> (Wagener, 1857)	M
<i>lameerei</i> Nouvel, 1932	AM
<i>Pleodicyema</i>	
<i>delamarei</i> Nouvel, 1961	M
<i>Pseudocyema</i>	
<i>truncatum</i> (Whitman, 1883)	AM
<h5>Family Conocyemidae</h5>	
<i>Conocyema</i>	
<i>polymorpha</i> van Beneden, 1882	M
<i>Microcyema</i>	
<i>vespa</i> van Beneden, 1882	AM

Class Orthonectida

Family Rhopaluridae

<i>Ciliocincta</i>	
<i>julini</i> (Caullery and Mesnil, 1899)	
<i>Intoschia</i>	
<i>leptoplanae</i> (Giard, 1877)	A
<i>linei</i> (Giard, 1877)	A
= <i>Rhopalura linei</i>	A
<i>major</i> Shtein, 1953	A
= <i>Rhopalura major</i>	A
<i>paraphanostomae</i> (Westblad, 1942)	
<i>Rhopalura</i>	
<i>elongata</i> Shtein, 1953	A
<i>gigas</i> (Giard, 1877)	A
<i>granosa</i> Atkins, 1933	A
<i>intoshi</i> Metchnikoff	M
<i>litoralis</i> Shtein, 1954	A
<i>metschnikowi</i> (Caullery and Mesnil, 1901)	A
<i>murmanica</i> Shtein, 1953	A
<i>ophiocomae</i> Giard, 1877	A
<i>pelseeneri</i> Caullery and Mesnil, 1901	A
<i>philinae</i> Lang, 1951	A
<i>pterocirri de Saint-Joseph</i> , 1896	A
<i>variabilis</i> Alexandrov and Sljusarev, 1992	A
<i>Stoecharthrum</i>	
<i>giardi</i> Caullery and Mesnil, 1899	A
<i>monnati</i> Kozloff, 1993	A

Family Pelmatosphaeridae

<i>Pelmatosphaera</i>	
<i>polycirri</i> Caullery and Mesnil, 1904	A

CNIDARIA

The Cnidaria (formerly called Coelenterata) comprises a much-varied phylum including for example such disparate forms as corals, medusae, siphonophores, hydroids, gorgonians, sea-anemones and soft corals. Study of each requires a different approach, and each has a distinct literature. A different author, therefore, has done the list of each of these groups.

Anthozoa

Octocorallia

Compiled by Jacob van der Land, L. van Ofwegen and M. Grasshoff

The European list was derived from URMO which was originally compiled by L. van Ofwegen (Leiden) and further corrected by M. Grasshoff.

Subclass Octocorallia

Order Alcyonacea

<i>Family Acanthogorgiidae</i>	
<i>Acanthogorgia</i>	
<i>armata</i> Verrill, 1878	
<i>hirsuta</i> Gray, 1857	
<i>pico</i> Grasshoff, 1973	

Family Alcyoniidae

<i>Alcyonium</i>	
<i>acaule</i> Marion, 1878	
<i>coralloides</i> (Pallas, 1766)	
<i>digitatum</i> Linnaeus, 1758	
<i>glomeratum</i> (Hassall, 1843)	
<i>palmatum</i> Pallas, 1766	A

Family Anthomastidae

<i>Anthomastus</i>	
<i>grandiflorus</i> Verrill, 1878	A
= <i>Anthomastus purpureus</i>	A

<i>Parerythropodium</i>	
<i>coralloides</i> (Pallas, 1766)	

Family Anthothelidae

<i>Anthothela</i>	
<i>grandiflora</i> (M. Sars, 1856)	A

Family Chrysogorgiidae

<i>Chrysogorgia</i>	
<i>agassizii</i> (Verrill, 1883)	A
<i>campanula</i> Madsen, 1944	A

<i>quadruplex</i> Thomson, 1927	
---------------------------------	--

Iridogorgia

<i>pourtalesii</i> Verrill, 1883	
----------------------------------	--

Metallogorgia

<i>melanotrichos</i> (Wright & Studer, 1889)	
--	--

Radicipes

<i>challengeri</i> (Wright, 1885)	A
-----------------------------------	---

<i>gracilis</i> (Verrill, 1884)	A
---------------------------------	---

Family Clavulariidae

<i>Clavularia</i>	
<i>arctica</i> (Sars, 1860)	A
<i>crassa</i> (Milne Edwards, 1848)	M

<i>marionii</i> (Von Koch, 1891)	A M
----------------------------------	-----

Sarcodictyon

<i>roseum</i> (Philippi, 1842)	A M
--------------------------------	-----

Scleranthelia

<i>microsclera</i> Lopez, Ocaña & Garcia, 1995	A
--	---

<i>rugosa</i> (Pourtales, 1867)	A M
---------------------------------	-----

Scyphopodium

<i>ingolfii</i> (Madsen, 1944)	A
--------------------------------	---

Telestula

<i>septentrionalis</i> Madsen, 1944	A
-------------------------------------	---

Family Coralliidae

<i>Corallium</i>	
------------------	--

<i>rubrum</i> (Linnaeus, 1758)	
--------------------------------	--

<i>niobe</i> Bayer, 1964		
<i>tricolor</i> (Johnson, 1899)		
Family Cornulariidae		
<i>Cerverta</i>		
<i>atlantica</i> (Johnson, 1861)	A	
<i>Cornularia</i>		
<i>cornucopiae</i> (Pallas, 1766)	A M	
Family Ellisellidae		
<i>Ellisella</i>		
<i>paraplexauroides</i> Stiasny, 1936		
<i>Nicella</i>		
<i>granifera</i> (Kölliker, 1865)		
<i>Viminella</i>		
<i>flagellum</i> (Johnson, 1863)		
Family Gorgoniidae		
<i>Eunicella</i>	M	
<i>cavolini</i> (Koch, 1887)		
<i>filiformis</i> (Studer, 1879)		
<i>gazella</i> Studer, 1878		
<i>labiata</i> Thomson, 1927		
<i>singularis</i> (Esper, 1791)	M	
<i>verrucosa</i> (Pallas, 1766)		
<i>Leptogorgia</i>		
<i>albipunctata</i> Stiasny, 1936		
<i>purpureoviolacea</i> (Stiasny, 1936)		
<i>sarmentosa</i> (Esper, 1791)		
<i>viminalis</i> (Pallas, 1766)		
Family Isididae		
<i>Acanella</i>	A	
<i>arbuscula</i> (Johnson, 1862)		
<i>Chelidonisis</i>		
<i>aurantiaca</i> Studer, 1890		
<i>Isidella</i>		
<i>elongata</i> (Esper, 1788)		
<i>losotensis</i> Sars, 1868	A	
<i>Keratoisis</i>		
<i>flexibilis</i> (Pourtales, 1868)		
<i>grayi</i> (Wright, 1869)		
<i>Lepidisis</i>		
<i>caryophyllia</i> Verrill, 1883		
<i>longiflora</i> Verrill, 1883		
<i>macropsiculata</i> (Kükenthal, 1919)		
Family Paragorgiidae		
<i>Paragorgia</i>	A	
<i>arborea</i> (Linnaeus, 1758)		
<i>johsoni</i> Gray, 1862		
Family Paralcyoniidae		
<i>Maasella</i>	M	
<i>edwardsi</i> (de Lacaze-Duthiers, 1888)		
<i>Paralcyonium</i>	A M	
<i>spinulosum</i> Delle Chiaje, 1822		
Family Plexauridae		
<i>Bebryce</i>		
<i>mollis</i> Molippi, 1842		
<i>Dentomuricea</i>		
<i>meteor</i> Grasshoff, 1977		
<i>Muriceides</i>	A M	
<i>kuekenthali</i> (Broch, 1912)		
<i>lepidia</i> Carpine & Grasshoff, 1975		
<i>paucituberculata</i> (Marion, 1882)		
<i>Paramuricea</i>		
<i>biscaya</i> Grasshoff, 1977	M	
<i>candida</i> Grasshoff, 1977		
<i>clavata</i> (Risso, 1826)	A	
<i>grayi</i> (Johnson, 1861)	M	
<i>macrospina</i> (Koch, 1882)	A M	
<i>placomus</i> (Linnaeus, 1758)		
<i>Placogorgia</i>		
<i>becena</i> Grasshoff, 1977		
<i>coronata</i> Carpine & Grasshoff, 1975	A M	
<i>graciosa</i> Tixier-Durivault & d'Hond, 1975		
<i>intermedia</i> (Thomson, 1927)		
<i>massiliensis</i> Carpine & Grasshoff, 1975	M	
<i>terceira</i> Grasshoff, 1977		
<i>Spinimuricea</i>		
<i>atlantica</i> (Johnson, 1862)		
<i>klaverenii</i> (Carpine & Grasshoff, 1975)	M	
<i>Swiftia</i>		
<i>borealis</i> (Kramp, 1930)		
<i>dubia</i> (Thomson, 1929)	A M	
<i>pallida</i> Madsen, 1970	A	
<i>rosea</i> (Grieg, 1887)	A	
<i>Thesea</i>		
<i>talismani</i> Grasshoff, 1986		
<i>Villogorgia</i>		
<i>bebrycoides</i> (Koch, 1887)	A M	
Family Primnoidae		
<i>Callogorgia</i>		
<i>verticillata</i> (Pallas, 1766)	M	
<i>Candidella</i>		
<i>imbricata</i> (Johnson, 1862)		
<i>Narella</i>		
<i>laxa</i> Deichmann, 1936		
<i>regularis</i> (Duchassaing & Michelotti, 1860)		
<i>versluysi</i> (Hickson, 1909)		
<i>Paracalyptrophora</i>		
<i>josephinae</i> (Lindström, 1877)		
<i>Primnoa</i>		
<i>resedaeformis</i> (Gunnerus, 1763)	A	
<i>Primnoella</i>		
<i>jungersenii</i> Madsen, 1944	A	
<i>Thouarella</i>		
<i>hilgendorfi</i> (Studer, 1879)		
Family Xeniidae		
<i>Anthelia</i>		
<i>borealis</i> (Koren & Danielsen, 1883)	A	
<i>fallax</i> Broch, 1912	A	
<i>Ceratocaulon</i>		
<i>wandeli</i> Jungersen, 1892	A	
Pennatulacea		
Compiled by Jacob van der Land and G. C. Williams		
The European list was derived from URMO which was edited by G. C. Williams. The later also checked the ERMS list.		
Order Pennatulacea		
Family Anthoptilidae		
<i>Anthoptilum</i>		
<i>grandiflorum</i> (Verrill, 1879)	A M	
<i>murrayi</i> Kölliker, 1880	A	
Family Funiculinidae		
<i>Funiculina</i>		
<i>quadrangularis</i> (Pallas, 1766)	A	
Family Halipteridae		
<i>Halipteris</i>		
<i>africana</i> (Studer, 1879)	A	
<i>christi</i> (Koren & Danielssen, 1847)	A	
<i>simarchica</i> (Sars, 1851)	A	
Family Kophobelemnidae		
<i>Kophobelemnion</i>		
<i>leucharti</i> Cecchini, 1917	M	
<i>macrospinosum</i> Thomson, 1927	A	
<i>stelliferum</i> (Müller, 1776)	A	
Family Pennatulidae		
<i>Crassophyllum</i>		
<i>thessalonicae</i> Vafidis & Koukouras, 1991	M	
<i>Gyrophyllum</i>		
<i>hirondellei</i> Studer, 1891	A	
<i>Pennatula</i>		
<i>aculeata</i> Danielssen, 1860	A	
<i>grandis</i> Ehrenberg, 1834	A	
<i>phosphorea</i> Linnaeus, 1758	A M	
<i>prolifera</i> Jungersen, 1904	A	

<i>rubra</i> (Ellis, 1764)	M	<i>biscayensis</i> (Fischer, 1874)	A
<i>Pteroeides</i>		<i>verrucosa</i> (Pennant, 1777)	A M
<i>lusitanicum</i> Broch, 1910	A	<i>Bunodosoma</i>	
<i>morbosus</i> Tixier-Durivault, 1961	A	<i>biscayensis</i> (Fischer, 1874)	A M
<i>spinosum</i> (Ellis, 1764)	M	<i>rubripunctata</i> (Grube, 1840)	M
<i>tenerum</i> Kölliker, 1869		<i>Condylostylis</i>	
Family Protoptilidae		<i>aurantiaca</i> (Delle Chiaje, 1825)	M
<i>Distichoptilum</i>		<i>Cribrinopsis</i>	
<i>gracile</i> Verrill, 1882	A M	<i>crassa</i> (Andrés, 1880)	M
<i>Protoptilum</i>		= <i>Anthopleura crassa</i> (Andrés, 1880)	
<i>carpenteri</i> Kölliker, 1872	A	<i>similis</i> Carlgren, 1921	A
<i>denticulatum</i> Jürgensen, 1904	A	<i>Epiactis</i>	
<i>smitii</i> Kölliker, 1872	A	<i>arctica</i> (Verrill, 1868)	A
<i>thomsoni</i> Kölliker, 1872	A	<i>secunda</i> (Verrill, 1899)	A
Family Umbellulidae		<i>incerta</i> Carlgren, 1921	A
<i>Umbellula</i>		<i>marsupialis</i> Carlgren, 1901	A
<i>encrinus</i> Linnaeus, 1758	A	<i>Onubactus</i>	
<i>huxleyi</i> Kölliker, 1880	A	<i>rocioi</i> Lopez, den Hartog & Garcia, 1995	A
<i>lindahli</i> Kölliker, 1875	A M	<i>Paranemoria</i>	
<i>monocephalus</i> Pasternak, 1964	A	<i>cinerea</i> (Contarini, 1844)	M
<i>thomsoni</i> Kölliker, 1874	A M	<i>vouliagmeniensis</i> Chintiroglou, Doumenc & England, 1987	M
Family Veretillidae		<i>Urticina</i>	
<i>Cavernularia</i>		<i>crassicornis</i> (Müller, 1776)	A
<i>elegans</i> (Herklotz, 1858)	A	= <i>Tealia crassicornis</i>	
<i>pusilla</i> (Philippi, 1835)	A M	<i>eques</i> (Gosse, 1858)	A
<i>Veretillum</i>		= <i>Tealia lofotensis</i>	
<i>cynomorium</i> (Pallas, 1766)	A	<i>felina</i> (Linnaeus, 1767)	A
Family Virgulariidae		= <i>Tealia felina</i>	
<i>Stylatula</i>		Family Actinoscyphiidae	
<i>elegans</i> (Danielssen, 1860)	A	<i>Actinoscyphia</i>	
<i>Virgularia</i>		<i>aurelia</i> (Stephenson, 1918)	A
<i>glacialis</i> Kölliker, 1870	A	<i>saginata</i> (Verrill, 1882)	A
<i>mirabilis</i> (Müller, 1776)	A	<i>Epiparactis</i>	
<i>tuberculata</i> Marshall, 1883	A	<i>dubia</i> Carlgren, 1921	A
Hexacorallia		Family Actinostolidae	
Actiniaria		<i>Actinostola</i>	
Compiled by J. van der Land and J. H. den Hartog		<i>callosa</i> (Verrill, 1882)	A
The ERMS list was derived from a preliminary version of the URMO. With the exception of the zooanthids checked by J. Ryland, this list must still be considered preliminary. K. Riemann-Zürneck assisted in the compilation of this list.		= <i>A. abyssorum</i>	
Order Actiniaria		<i>Anthosactis</i>	
Family Actinernidae		<i>ingolfi</i> Carlgren, 1921	A
<i>Actinernus</i>		<i>janmayeni</i> Danielssen, 1890	A
<i>michaelsarsi</i> Carlgren, 1918	A	<i>Glandulactis</i>	
Family Actiniidae		<i>spetsbergensis</i> (Carlgren, 1893)	A
<i>Actinia</i>		<i>Paranthus</i>	
<i>cari</i> Delle Chiaje, 1825	M	<i>rugosus</i> Andrés, 1880	M
<i>elongata</i> Delle Chiaje, 1825	M	<i>Parasicyonis</i>	
<i>equina</i> (Linnaeus, 1758)	A M	<i>ingolfi</i> Carlgren, 1942	A
<i>filiformis</i> Rapp, 1829	A	<i>sarsi</i> Carlgren, 1921	A
<i>fragacea</i> Tugwell, 1856	A	<i>Pycnanthus</i>	
<i>prasina</i> Gosse, 1860	A	<i>densus</i> Carlgren, 1921	A
<i>striata</i> (Rizzi, 1967)	M	<i>laevis</i> Carlgren, 1921	A
<i>Anemonia</i>		<i>Sicyon</i>	
<i>viridis</i> (Forskål, 1775)	A M	<i>biotrans</i> Riemann-Zürneck, 1991	A
<i>Anthopleura</i>		<i>gossei</i> (Stephenson, 1918)	A
<i>ballii</i> (Cocks, 1849)	A M	<i>hemisphaerica</i> Carlgren, 1934	A
<i>rubripunctata</i> (Grube, 1840)	M	<i>ingolfi</i> Carlgren, 1921	A
<i>thallia</i> (Gosse, 1854)	A M	<i>tuberculata</i> Carlgren, 1921	A
<i>Aulactinia</i>		<i>variabilis</i> Carlgren, 1921	A
<i>spetsbergensis</i> (Carlgren, 1921)	A	<i>Stomphia</i>	
<i>stella</i> (Verrill, 1864)	A	<i>coccinea</i> (Müller, 1776)	A
<i>verrucosa</i> (Pennant, 1777)	A M	<i>polaris</i> (Danielssen, 1890)	A
<i>Bolocera</i>		<i>Telidium</i>	
<i>maxima</i> Carlgren, 1921	A	<i>jungersoni</i> Carlgren, 1921	A
<i>tuediae</i> (Johnston, 1832)	A	Family Aiptasiidae	
<i>Bunodactis</i>		<i>Aiptasia</i>	
		<i>diaphana</i> (Rapp, 1829)	M
		<i>mutabilis</i> (Gravenhorst, 1831)	A M
		= <i>A. couchii</i>	
		<i>Aiptasiogonet</i>	
		<i>pellucidus</i> (Hollard, 1848)	A M
		Family Aliciidae	
		<i>Alicia</i>	
		<i>mirabilis</i> Johnson, 1861	A M

= <i>A. costae</i>			
Family Andrenidae			
<i>Andrenia</i>			
<i>partenopea</i> (Andrés, 1884)	M		
Family Andvakiidae			
<i>Andvaka</i>			
<i>mirabilis</i> Danielssen, 1890	A		
<i>parva</i> Carlgren, 1940	A		
Family Aureliidae			
<i>Aureliana</i>			
<i>heterocera</i> (Thompson, 1853)	A M		
Family Bathypelliidae			
<i>Bathypellia</i>			
<i>margaritacea</i> (Danielssen, 1890)	A		
<i>Daontesia</i>			
<i>porcupina</i> Riemann-Zürneck, 1997	A		
<i>praelonga</i> (Carlgren, 1928)	A		
Family Boloceroididae			
<i>Bunodeopsis</i>			
<i>strumosa</i> Andrès, 1880	M		
Family Condylanthidae			
<i>Segonzacis</i>			
<i>platypus</i> Riemann-Zürneck, 1979	A		
Family Diadumenidae			
<i>Diadumene</i>			
<i>cincta</i> Stephenson, 1925	A		
<i>Haliplanella</i>			
<i>luciae</i> (Verrill, 1899)	A M		
Family Edwardsiidae			
<i>Edwardsia</i>			
<i>allmanni</i> McIntosh, 1865	A		
<i>andresi</i> Danielssen, 1890	A		
<i>arctica</i> Carlgren, 1921	A		
<i>beautempsi</i> Quatrefages, 1842	A		
<i>carlgreni</i> Williams, 1981	A		
<i>claparedii</i> (Panceri, 1869)	A M		
<i>costata</i> Danielssen, 1890	A		
<i>danica</i> Carlgren, 1921	A		
<i>delapiae</i> Carlgren & Stephenson, 1928	A		
<i>finmarchica</i> Carlgren, 1921	A		
<i>fusca</i> Danielssen, 1890	A		
<i>islandica</i> Carlgren, 1921	A		
<i>ivelli</i> Manuel, 1975	A		
<i>longicornis</i> Carlgren, 1921	A		
<i>maroccana</i> Carlgren, 1931	A		
<i>norvegica</i> Carlgren, 1942	A		
<i>timida</i> Quatrefages, 1842	A		
= <i>E. callianthus</i>			
= <i>E. harassei</i>			
<i>timida</i> Quatrefages, 1842	A		
= <i>Edwardsiella dixoni</i>			
<i>tuberculata</i> Duben & Koren, 1847	A		
<i>vegae</i> Carlgren, 1921	A		
<i>vitrea</i> (Danielssen, 1890)	A		
<i>Fagesia</i>			
<i>carnea</i> (Gosse, 1856)	A M		
<i>janthina</i> (Andrés, 1880)	M		
<i>loveni</i> (Carlgren, 1892)	A M		
<i>Isoedwardsia</i>			
<i>nidarosiensis</i> Carlgren, 1942	A		
<i>Nematostella</i>			
<i>nathorsti</i> (Carlgren, 1921)	A		
<i>polaris</i> (Carlgren, 1921)	A		
<i>vectensis</i> Stephenson, 1935	A		
= <i>N. pellucida</i>			
<i>Paraedwardsia</i>			
<i>abyssorum</i> Carlgren, 1951	A		
<i>arenaria</i> Carlgren in Nordgaard, 1905	A		
<i>sarsi</i> (Duben & Koren, 1847)	A		
<i>Scolanthus</i>			
<i>callimorphus</i> Gosse, 1853	A M		
= <i>Isoedwardsia mediterranea</i>			
<i>ingolfi</i> Carlgren, 1921	A		
		Family Gonactiniidae	
		<i>Gonactinia</i>	
		<i>prolifera</i> (Sars, 1835)	A M
		<i>Protanthea</i>	
		<i>simplex</i> Carlgren, 1891	A
		Family Halcampidae	
		<i>Cactosoma</i>	
		<i>abyssorum</i> Danielssen, 1890	A
		<i>Halcampia</i>	
		<i>arctica</i> Carlgren, 1893	A
		<i>chrysanthellum</i> (Peach in Johnston, 1847)	A
		<i>duodecimcirrata</i> (Sars, 1851)	A
		Family Halcampoididae	
		<i>Acthelmis</i>	
		<i>intestinalis</i> (Fabricius, 1780)	A
		<i>Halcampella</i>	
		<i>endromitata</i> (Andrés, 1880)	M
		<i>Halcampoides</i>	
		<i>abyssorum</i> Danielssen, 1890	A M
		<i>Siphonactinopsis</i>	
		<i>laevis</i> Carlgren, 1921	A
		Family Halcuriidae	
		<i>Carlgrenia</i>	
		<i>desiderata</i> Stephenson, 1918	A
		Family Haliactiidae	
		<i>Haliactis</i>	
		<i>arctica</i> Carlgren, 1921	A
		Family Haloclavidae	
		<i>Anemonactis</i>	
		<i>mazeli</i> (Jourdan, 1880)	A M
		<i>Mesacmaea</i>	
		<i>mitchelli</i> (Gosse, 1853)	A M
		<i>Peachia</i>	
		<i>boekii</i> (Danielssen & Koren, 1856)	A
		<i>cylindrica</i> (Reid, 1848)	A M
		= <i>P. hastata</i> Gosse, 1855	
		Family Hormathiidae	
		<i>Actinauge</i>	
		<i>abyssorum</i> (Gravier, 1918)	A
		<i>richardi</i> (Marion, 1882)	A M
		<i>Adamsia</i>	
		<i>carcinopodus</i> (Otto, 1823)	A M
		= <i>A. palliata</i>	
		<i>Allantactis</i>	
		<i>parasitica</i> Danielssen, 1890	A
		<i>Amphianthus</i>	
		<i>bathybium</i> (Hertwig, 1882)	A
		<i>crassus</i> (Gravier, 1918)	A
		<i>dohrni</i> (von Koch, 1878)	A M
		<i>impeditus</i> (Gravier, 1918)	A
		<i>ingolfi</i> Carlgren, 1942	A
		<i>inornatus</i> (Gravier, 1918)	A
		<i>islandicus</i> Carlgren, 1942	A
		<i>margaritaceus</i> (Danielssen, 1890)	A
		<i>michaelsarsi</i> Carlgren, 1934	A
		<i>mopsiae</i> (Danielssen, 1890)	A
		<i>norvegicus</i> Carlgren, 1942	A
		<i>verruculatus</i> Carlgren, 1942	A
		<i>Calliactis</i>	
		<i>parasitica</i> (Couch, 1838)	A M
		<i>Cataphellia</i>	
		<i>brodrickii</i> (Gosse, 1859)	A
		<i>Chondrophellia</i>	
		<i>coronata</i> (Verrill, 1883)	A
		<i>Hormathia</i>	
		<i>alba</i> (Andrés, 1880)	A M
		= <i>H. mediterranea</i>	
		<i>coronata</i> (Gosse, 1858)	A M
		<i>digitata</i> (O. F. Müller, 1776)	A
		<i>marioni</i> (Haddon, 1889)	A
		<i>nodosa</i> (Fabricius, 1780)	A
		<i>Monactis</i>	
		<i>vestita</i> Gravier, 1918	A

<i>Paracallistiactis</i>		
<i>azorica</i> Doumenc, 1975	A	
<i>mediterranea</i> Ross & Zamponi, 1982	M	
<i>michaelsarsi</i> Carlgren, 1928	A	
<i>stephensonii</i> Carlgren, 1928	A	
<i>Paractinia</i>		
<i>striata</i> (Risso, 1826)	M	
= <i>Parastepphanauge paxi</i>		
<i>Paraphellia</i>		
<i>expansa</i> (Haddon, 1886)	A	
<i>Phelliactis</i>		
<i>hertwigi</i> Simon, 1892	A	
= <i>P. incerta</i>		
<i>michaelsarsi</i> (Carlgren, 1934)	A	
<i>pulchra</i> (Stephenson, 1918)	A	
<i>robusta</i> Carlgren, 1928	A	
<i>spinosa</i> (Carlgren, 1928)	A	
<i>Stephanauge</i>		
<i>acanellae</i> (Verrill, 1883)	A	
Family Iosactiidae		
<i>Iosactis</i>		
<i>vagabunda</i> Riemann-Zürneck, 1997	A	
Family Isophellidae		
<i>Euphellia</i>		
<i>cinctidifera</i> Pax, 1908	A	
<i>Telmatactis</i>		
<i>cricoides</i> (Duchassaing, 1850)	A M	
<i>forskali</i> (Ehrenberg, 1834)	A M	
= <i>T. limicola</i>		
<i>solidago</i> (Duchassaing & Michelotti, 1860)	A	
Family Kadostactidae		
<i>Kadosactis</i>		
<i>abyssicola</i> (Danielssen, 1877)	A	
<i>commensalis</i> (Gravier, 1918)	A	
<i>rosea</i> Danielssen, 1890	A	
<i>spitsbergensis</i> (Danielssen, 1890)	A	
<i>sulcata</i> Carlgren, 1934	A	
Family Limnactiniidae		
<i>Limnactinia</i>		
<i>laevis</i> Carlgren, 1921	A	
Family Liponematidae		
<i>Liponema</i>		
<i>multicornis</i> (Verrill, 1879)	A	
Family Metridiidae		
<i>Metridium</i>		
<i>senile</i> (Linnaeus, 1767)	A	
Family Octineonidae		
<i>Octineon</i>		
<i>lindahli</i> (Carpenter, 1871)	A	
<i>suecicum</i> Carlgren, 1940	A	
Family Phymantidae		
<i>Phymanthus</i>		
<i>pulcher</i> (Andrés, 1884)	M	
Family Sagartiidae		
<i>Actinothoe</i>		
<i>sphyrodetta</i> (Gosse, 1858)	A	
<i>Cereus</i>		
<i>pedunculatus</i> (Pennant, 1777)	A M	
<i>Phellia</i>		
<i>elongata</i> (Delle Chiaje, 1825)	A M	
<i>gausapata</i> Gosse, 1858	A	
<i>vestita</i> (Johnson, 1861)	A	
<i>Sagartia</i>		
<i>elegans</i> (Dalyell, 1848)	A M	
<i>luciae</i> Verrill, 1898	A M	
<i>ornata</i> Holdsworth, 1858	A	
<i>troglodytes</i> (Price in Johnston, 1847)	A M	
<i>Sagartiogiton</i>		
<i>abyssorum</i> Carlgren, 1942	A	
<i>entellae</i> Schmidt, 1972	M	
<i>flexibilis</i> (Danielssen, 1890)	A	
<i>ingolfi</i> Carlgren, 1928	A	
<i>laceratus</i> (Dalyell, 1848)		
<i>robustus</i> Carlgren, 1924		
<i>tubicolus</i> (Koren & Danielssen, 1877)		
<i>undatus</i> (Müller, 1788)		
<i>verrilli</i> Carlgren, 1942		
<i>viduatus</i> (Müller, 1776)		
Order Ceriantharia		
Family Arachnactidae		
<i>Arachnanthus</i>		
<i>nocturnus</i> den Hartog, 1977		
<i>sarsi</i> Carlgren, 1912		
<i>Isarachnanthus</i>		
<i>cruzi</i> Brito, 1986		
<i>Arachnactis</i>		
<i>albida</i> Sars, 1846		
Family Botrucnidiferidae		
<i>Botracnidifer</i>		
<i>norvegicus</i> Carlgren, 1912		
Family Cerianthidae		
<i>Cerianthus</i>		
<i>lloydii</i> Gosse, 1859		
<i>membranacea</i> (Spallanzani, 1784)		
<i>vogti</i> Danielssen, 1890		
<i>Pachycerianthus</i>		
<i>multiplicatus</i> Carlgren, 1912		
Order Corallimorpharia		
Family Corallimorphidae		
<i>Corallimorphus</i>		
<i>atlanticus</i> Carlgren, 1934		
<i>ingens</i> Gravier, 1918		
<i>Corynactis</i>		
<i>viridis</i> Allman, 1846		
<i>Pseudocorynactis</i>		
<i>caribbeorum</i> Den Hartog, 1980		
Family Sideractiidae		
<i>Nectactis</i>		
<i>singularis</i> Gravier, 1918		
<i>Sideractis</i>		
<i>glacialis</i> Danielssen, 1890		
Order Ptychodactiaria		
Family Ptychodactiidae		
<i>Ptychodactis</i>		
<i>patula</i> Appellöf, 1893		
Order Zoanthidea		
Family Epizoanthidae		
<i>Epizoanthus</i>		
<i>abyssorum</i> Verrill, 1885		
<i>arenaceus</i> delle Chiaje, 1823	M	
<i>beerenislandicus</i> Carlgren, 1913	A	
<i>couchii</i> (Johnston in Couch, 1844)	A	
<i>danielsseni</i> Carlgren, 1913	A	
<i>erdmanni</i> (Danielssen, 1890)	A	
<i>frenzeli</i> Pax, 1937	M	
<i>glacialis</i> Danielssen, 1890	A	
<i>koreni</i> Carlgren, 1913	A	
<i>lindahli</i> Carlgren, 1913	A	
<i>macintoshi</i> Haddon & Shackleton, 1891	A	
<i>mediterraneus</i> Carlgren, 1935	M	
<i>norvegicus</i> (Koren & Danielssen, 1877)	A	
<i>paguricola</i> (Roule, 1900)	M	
<i>paguriphilus</i> Verrill, 1883	A	
<i>papillosum</i> (Johnston, 1842)	A	
= <i>E. americanus</i>		
<i>paxii</i> Abel, 1955	M	
<i>roseus</i> Danielssen, 1890	A	
<i>steueri</i> Pax, 1937	M	
<i>tergestinus</i> Pax, 1937	M	
<i>univittatus</i> (Lorenz, 1860)	M	
<i>vagus</i> Herbersts, 1972	M	
<i>vatovai</i> Pax & Lochter, 1935	M	
Family Parazoanthidae		

<i>Gerardia</i>		A M	References	
<i>savalia</i> (Bertholoni, 1819)			Cairns, S.D., B.W. Hoeksema & J. van der Land, 1999. List of extant stony corals. - <i>Atoll Research Bull.</i> : 459: 13-46.	
<i>Isozoanthus</i>		A		
<i>arborensens</i> (Danielssen, 1890)		A		
<i>bulbosus</i> Carlgren, 1913		A		
<i>dubius</i> Carlgren, 1913		A		
<i>ingolfi</i> Carlgren, 1913		A		
<i>islandicus</i> Carlgren, 1913		A		
<i>magnisulosus</i> Carlgren, 1913		A		
<i>multinsulosus</i> Carlgren, 1913		A		
<i>sulcatus</i> (Gosse, 1859)		A		
<i>Parazoanthus</i>		A		
<i>anguicomus</i> (Norman, 1868)		A		
<i>axinellae</i> (Schmidt, 1862)		A M		
Family Zoanthidae				
<i>Isaurus</i>		A M		
<i>tuberculatus</i> Gray, 1828				
<i>Protopalythoa</i>		A		
<i>canariensis</i> (Haddon & Duerden, 1896)				
Antipatharia				
Compiled by Jacob van der Land and D. M. Opresco				
The ERMS list was derived from URMO which was edited by D. M. Opresco.				
Order Antipatharia				
Family Antipathidae				
<i>Antipathes</i>				
<i>dichotoma</i> Pallas, 1766		A M		
<i>erinaceus</i> (Roule, 1905)		A		
<i>fragilis</i> Gravier, 1918		A		
<i>furcata</i> Gray, 1857		A		
<i>mediterranea</i> Brook, 1889		M		
<i>punctata</i> (Roule, 1905)		A		
<i>spinescens</i> Gray, 1857		A		
<i>subpinnata</i> Ellis & Solander, 1786		A M		
<i>wollastoni</i> Gray, 1857		A		
<i>Aphanipathes</i>		A		
<i>abietina</i> (Pourtalès, 1874)				
<i>Bathyphathes</i>				
<i>alternata</i> Brook, 1889		A M		
<i>arctica</i> (Lütken, 1871)		A		
<i>heterorhodzos</i> (Cooper, 1909)		A M		
<i>lyra</i> Brook, 1889		A M		
<i>patula</i> Brook, 1889		A M		
<i>tenuis</i> Brook, 1889		A M		
<i>Leiopathes</i>		A		
<i>expansa</i> Johnson, 1899		A M		
<i>glaberrima</i> (Esper, 1788)				
= <i>Antipathes glaberrima</i>				
<i>grimaldi</i> Roule, 1902		A		
<i>Parantipathes</i>		A M		
<i>larix</i> (Esper, 1788)				
<i>Schizopathes</i>		A M		
<i>crassa</i> Brook, 1889				
<i>Stichopathes</i>				
<i>abyssicola</i> Roule, 1902		A		
<i>dissimilis</i> Roule, 1902		A		
<i>euplos</i> Schultze, 1903		A		
<i>flagellum</i> Roule, 1902		A		
<i>gracilis</i> Gray, 1857		A M		
<i>richardi</i> Roule, 1902		A		
<i>robusta</i> Gravier, 1918		A		
Scleractinia				
Compiled by S. D. Cairns, B. W. Hoeksema and Jacob van der Land				
The ERMS list was derived from URMO which was recently published (Cairns <i>et al.</i> 1999). The list was checked and amended by H. Zibrowius.				
Order Scleractinia				
Family Caryophylliidae				
<i>Anomocora</i>				
<i>fecunda</i> (Pourtalès, 1871)			A	
<i>Asterosmilia</i>			A	
<i>prolifera</i> (Pourtalès, 1871)			A	
<i>Aulocyathus</i>				
<i>atlanticus</i> Zibrowius, 1980			A	
<i>Caryophyllia</i>				
<i>abyssorum</i> Duncan, 1873			A	
<i>alberti</i> Zibrowius, 1980			A	
<i>ambrosia</i> Alcock, 1898			A M	
<i>atlantica</i> (Duncan, 1873)			A	
<i>calveri</i> Duncan, 1873			A M	
<i>cyathus</i> (Ellis & Solander, 1786)			A M	
<i>diomedae</i> Marenzeller, 1904			A M	
<i>foresti</i> Zibrowius, 1980			A	
<i>inornata</i> (Duncan, 1878)			A M	
<i>sarsiae</i> Zibrowius, 1974			A M	
<i>seguenzae</i> Duncan, 1873			A	
<i>smithii</i> Stokes & Broderip, 1828			A M	
<i>Ceratotrochus</i>				
<i>magnaghii</i> Cecchini, 1914			A M	
<i>Cladocora</i>				
<i>caespitosa</i> (Linnaeus, 1758)			A M	
<i>debilis</i> Milne Edwards & Haime, 1849			A M	
<i>Coenocyathus</i>				
<i>anthophyllites</i> Milne Edwards & Haime, 1848			A M	
<i>cylindricus</i> Milne Edwards & Haime, 1848			A M	
<i>Coenosmilia</i>				
<i>arbuscula</i> Pourtalès, 1874			A	
<i>Concentrotheca</i>				
<i>laevigata</i> (Pourtalès, 1871)			A	
<i>Dasmosmilia</i>				
<i>lymani</i> (Pourtalès, 1871)			A	
<i>variegata</i> (Pourtalès, 1871)			A	
<i>Deltocyathus</i>				
<i>italicus</i> (Michelotti, 1838)			A	
<i>eccentricus</i> Cairns, 1979			A	
<i>moseleyi</i> Cairns, 1979			A	
<i>Desmophyllum</i>				
<i>dianthus</i> (Esper, 1794)			A M	
<i>Hoplangia</i>				
<i>durotrix</i> Gosse, 1860			A M	
<i>Lophelia</i>				
<i>pertusa</i> (Linnaeus, 1758)			A M	
<i>Paracyathus</i>				
<i>arcuatus</i> Lindström, 1877			A	
<i>pulchellus</i> (Philippi, 1842)			A M	
<i>Phyllangia</i>				
<i>mouchezii</i> (Lacaze-Duthiers, 1897)			A M	
<i>Polycyathus</i>				
<i>muellerae</i> (Abel, 1959)			A M	
<i>Pourtalosmilia</i>				
<i>anthophyllites</i> (Ellis & Solander, 1786)			A M	
<i>Premocyathus</i>				
<i>cornuformis</i> (Pourtalès, 1868)			A	
<i>Solenosmilia</i>				
<i>variabilis</i> Duncan, 1873			A M	
<i>Stephanocyathus</i>				
<i>campaniformis</i> (Marenzeller, 1904)			A	
<i>crassus</i> (Jourdan, 1895)			A	
<i>moseleyanus</i> (Sclater, 1886)			A	
<i>nobilis</i> (Moseley, 1873)			A	
<i>Tethocyathus</i>				
<i>variabilis</i> Cairns, 1979			A	
<i>Thalamophyllia</i>				
<i>gasti</i> (Döderlein, 1913)			A M	
<i>Trochocyathus</i>				
<i>spinosocostatus</i> Zibrowius, 1980			A	
<i>mediterraneus</i> Zibrowius, 1980			M	

<i>Vaughanella</i>	
<i>concinna</i> Gravier, 1915	A
Family Dendrophylliidae	
<i>Astroides</i>	
<i>calycularis</i> (Pallas, 1766)	A M
<i>Balanophyllia</i>	
<i>cellulosa</i> Duncan, 1873	A M
<i>europaea</i> (Risso, 1826)	A M
<i>regia</i> Gosse, 1860	A M
<i>thalassae</i> Zibrowius, 1980	A
<i>Cladopsammia</i>	
<i>rolandi</i> Lacaze-Duthiers, 1897	M
<i>Dendrophyllia</i>	
<i>alternata</i> Pourtalès, 1880	A
<i>cornigera</i> (Lamarck, 1816)	A M
<i>laboreli</i> Zibrowius & Brito, 1984	A
<i>ramea</i> (Linnaeus, 1758)	A M
<i>Eguchipsammia</i>	
<i>cornucopia</i> (Pourtalès, 1871)	A
<i>gaditana</i> (Duncan, 1873)	A M
<i>Enallopssammia</i>	
<i>rostrata</i> (Pourtalès, 1878)	A M
<i>Leptopsammia</i>	
<i>britannica</i> (Duncan, 1870)	A
<i>formosa</i> (Gravier, 1915)	A
<i>pruvoti</i> Lacaze-Duthiers, 1897	A
Family Flabellidae	
<i>Flabellum</i>	
<i>alabastrum</i> Moseley, 1876	A
<i>angulare</i> Moseley, 1876	A
<i>chunii</i> Marenzeller, 1904	A
<i>macandrewi</i> Gray, 1849	A
<i>Javania</i>	
<i>cailleti</i> (Duchassaing & Michelotti, 1864)	
<i>pseudoalabastra</i> Zibrowius, 1974	A
<i>Monomyces</i>	
<i>pygmaea</i> (Risso, 1826)	A M
<i>Placotrochides</i>	
<i>frustum</i> Cairns, 1979	A
<i>Truncatoflabellum</i>	
<i>stabile</i> (Marenzeller, 1904)	A
Family Fungiacyathidae	
<i>Fungiacyathus</i>	
<i>crispus</i> (Pourtalès, 1871)	A
<i>fragilis</i> Sars, 1872	A M
<i>marenzelleri</i> (Vaughan, 1906)	A M
Family Guyniidae	
<i>Guynia</i>	
<i>annulata</i> Duncan, 1872	A
<i>Schizocyathus</i>	
<i>fissilis</i> Pourtalès, 1874	A
<i>Stenocyathus</i>	
<i>vermiformis</i> (Pourtalès, 1868)	A M
Family Oculinidae	
<i>Madrepora</i>	
<i>oculata</i> Linnaeus, 1758	A M
<i>Oculina</i>	
<i>patagonica</i> de Angelis, 1908	A M
Family Pocilloporidae	
<i>Madracis</i>	
<i>asperula</i> Milne Edwards & Haime, 1849	A
<i>pharensis</i> (Heller, 1868)	A M
<i>profunda</i> Zibrowius, 1980	A
Family Rhizangiidae	
<i>Astrangia</i>	
<i>macrodentata</i> Thiel, 1940	A
Family Turbinoliidae	
<i>Deltocyathoides</i>	
<i>stimpsonii</i> (Pourtalès, 1871)	A
<i>Peponocyathus</i>	
<i>folliculus</i> (Pourtalès, 1868)	A
<i>Sphenotrochus</i>	
<i>andrewianus</i> Milne Edwards & Haime, 1848	A M

Scyphozoa

Compiled by Paul F. S. Cornelius, Gerhard Jarms, and Yakko Hirano.

The Scyphozoa are a somewhat uniform but ancient group, comprising probably fewer than 200 species world-wide. Many of the extant genera, and in some cases even the species, are markedly distinct from their closest relatives. There are only a few genera of several confusingly similar species. Nevertheless, because many species are large and conspicuous, occurring primarily near the coast, repeated redescription and consequent taxonomic confusion has hampered taxonomic understanding. What follows is probably the first reliable list of European species. Within the present lists, the Stauromedusae were checked by Dr Yakko Hirano, the Coronatae by Dr Gerhard Jarms, and the remainder by Dr Paul F. S. Cornelius.

The most helpful monograph is F. S. Russell's (1970) superb treatise. A key to world genera was published more recently (Cornelius, 1997). Most, but not all, are shallow-water coastal species. However, *Pelagia* is open-ocean planktonic, and *Stygiomedusa* and *Deepstaria* are deep-water forms. Scyphomedusae are most speciose in the tropics. All major divisions are represented in Europe though few rhizostomes extend north of the Mediterranean Sea. One, Europe's second largest invertebrate, *Catostylus tagi*, occurs in the Tajo estuary near Lisbon and from there southwards along the African coast, though the almost as huge *Rhizostoma octopus* is common at least as far north as southern Scotland and may exceptionally occur in Norwegian waters. The semaeostome *Cyanea capillata* may grow to approximately 2m diameter in continuous arctic daylight. A useful paper by Galil, Spanier & Ferguson (1990) describes and illustrates some immigrant species occurring in the Eastern Mediterranean. Of the Stauromedusae there is perhaps no European guide. NW European species were nicely summarised by J. S. Ryland (in Hayward & Ryland, 1990 (detailed), 1995 (abridged)). The European list that follows here was extracted by J. van der Land from a world list prepared by Dr Y. Hirano. A manuscript translation into English of the Stauromedusae section of the important book by Naumov (1961) is held in the library of the Natural History Museum, London.

References

- Cornelius, P. F. S., 1997. Keys to the genera of Cubomedusae and Scyphomedusae (Cnidaria). *Proceedings of the 6th International Conference on Coelenterate Biology (1995)*. J. C. den Hartog, (ed.), Leiden, National Natuurhistorisch Museum, 109-122.
- Cornelius, P. F. S., Manuel, R. L., & Ryland, J. S., 1990. Cnidaria. In: Hayward, P. J. & Ryland, J. S. (eds) *The Marine Fauna of the British Isles and North-West Europe*. Volume 1. Introduction and Protozoans to Arthropods. Oxford, Clarendon Press, pp. 101-180. [includes stauromedusae, no other scyphozoans]
- Cornelius, P. F. S., Manuel, R. L., & Ryland, J. S., 1995. Hydroids, sea anemones, jellyfish, and comb jellies. In: Hayward, P. J. & Ryland, J. S. (eds) *Handbook of the Marine fauna of North-West Europe*. Oxford, Oxford University Press, pp. 62-135. [abridged edition; includes both stauromedusae and scyphomedusae].
- Galil, B. S. Spanier, E. & Ferguson, W. W., 1990. The Scyphomedusae of the Mediterranean coast of Israel, including two Lessepsian migrants new to the Mediterranean. *Zoologisches Mededelingen* 64, 5-105.
- Naumov, D. V., 1961. Stsifoidnye meduzy morei S.S.S.R. [Scyphomedusae of the seas of the U.S.S.R.] *Fauna S.S.S.R.* 75, 1-98.
- Russell, F. S., 1970. *The medusae of the British Isles. II. Pelagic Scyphozoa with a supplement to the first volume on hydromedusae*. Cambridge, Cambridge University Press, pp. xii, 284.

Class Cubozoa

Order Cubomedusae

Family Carybdeidae

Carybdea marsupialis (Linnaeus, 1758)

Class Scyphozoa

Order Coronatae

Family Atollidae

Atolla parva Russell, 1958
vanhoeffeni Russell, 1957
wyvillei Haeckel, 1880

Family Atorellidae

Atorella subglobosa Vanhoffen, 1902

Family Nausithoidae

Nausithoe atlantica Broch, 1914
eumedusoides (Werner, 1974)
globifera Broch, 1914
limpida Hartlaub, 1909
marginata Köllicker, 1853
planulophora (Werner, 1971)
punctata Köllicker, 1853
rubra Vanhoffen, 1902
werneri Jarms, 1990

Family Paraphyllinidae

Paraphyllina ransonni Russell, 1956

Family Periphyllidae

Periphylla periphylla (Péron & Lesueur, 1810)

Order Semaeostomeae

Family Pelagiidae

Chrysaora hysoscella (Linnaeus, 1767)
Pelagia noctiluca (Forsskål, 1775)

Family Cyaneidae

Cyanea capillata (Linnaeus, 1758)
lamarckii Péron & Lesueur, 1810
Drymonema dalmatinum Haeckel, 1880

Family Ulmaridae

Aurelia aurita (Linnaeus, 1758)

Deepstaria

enigmatica Russell, 1967

Discomedusa

lobata Claus, 1877

Phacellophora

camtschatica Brandt, 1835

Poralia

rufescens Vanhoffen, 1902

Stygiomedusa

gigantea (Browne, 1910)

Order Rhizostomeae

Family Cassiopeidae

Cassiopea andromeda (Forsskål, 1775)

Family Cepheidae

Cotylorhiza
tuberculata (Macri, 1778)

Family Mastigiidae

Phyllorhiza
punctata von Lendenfeld, 1884

Family Catostylidae

Catostylus
tagi (Haeckel, 1869)

Family Rhizostomatidae

Rhizostoma
pulmo (Macri, 1778)
Rhopilema
rhopalophorum Haeckel, 1880

Order Stauromedusae

Family Depastridae

Craterolophus
convolvulus (Johnson, 1835)
Depastrum
cyathiforme (M. Sars, 1846)
Manania
auricula (Fabricius, 1780)
= *Halimocyathus lagena*
hexaradiata (Broch, 1907)
uchidai (Naumov, 1961)

Family Kishinouyeidae

Lucernariopsis
campanulata (Lamouroux, 1815)
cruxmelitensis Corbin, 1978

Family Lipkeidae

Lipkea
ruspoliana Vogt, 1886
sturdzi (Antipa, 1893)

Family Lucernariidae

Haliclystus
auricula (Rathke, 1806)
octoradiatus (Lamarck, 1816)
salpinx Clark, 1863
Lucernaria
bathyphila Haeckel, 1880
haeckeli (Antipa, 1892)
infundibulum Haeckel, 1880
quadricornis O.F. Müller, 1776
sainthilarei (Radicorzev.)
walteri (Antipa, 1892)
Stylocoronella
riedli Salvini-Plawen, 1966
variabilis Salvini-Plawen, 1987

Class incertae sedis

Family Tetraplatidae

Tetraplatia
volitans Busch, 1851

Hydrozoa

Hydroida

Compiled by Jacob van der Land, W. Vervoort, S.D. Cairns and P. Schuchert

The European list was derived from URMO which was edited by W. Vervoort. S.D. Cairns provided the list of hydrocorals and P. Schuchert provided additions to the list of hydroids.

Class Hydrozoa

Order Actinulida

Family Halammohydridae

<i>Halammohydra</i>	
<i>adherens</i> Swedmark & Teissier, 1958	A
<i>coronata</i> Clausen, 1967	A
<i>intermedia</i> Clausen, 1967	A
<i>neglecta</i> Clausen	A
<i>octopodides</i> Remane, 1927	A M
<i>schulzei</i> Remane, 1927	A M
<i>vermiformis</i> Swedmark & Teissier, 1857	A

Family Otohydridae

<i>Otohydra</i>	
<i>tremulans</i> Lacassagne, 1973	A M
<i>vagans</i> Swedmark & Teissier, 1958	A M

Order Capitata

Family Acaulidae

<i>Acaulis</i>	
<i>primarius</i> Stimpson, 1854	A
<i>Acauloides</i>	
<i>ammisatum</i> Bouillon, 1965	A
<i>ilonae</i> (Brinckmann-Voss, 1966)	A M

Family Boreohydridae

<i>Boreohydra</i>	
<i>simplex</i> Westblad, 1937	
<i>Meiorhopalon</i>	
<i>arenicolum</i> Von Salvini-Plawen, 1984	A M
<i>Psammonhydra</i>	
<i>nana</i> Schulz, 1950	A M

Family Candelabridae

<i>Candelabrum</i>	
<i>cocksii</i> (Vigurs, 1849)	A
<i>minutum</i> (Bonnevie, 1898)	A
<i>mitra</i> (Bonnevie, 1898)	A
<i>phrygium</i> (Fabricius, 1780)	A
<i>serpentarii</i> Segonzac & Vervoort, 1995	
<i>verrucosum</i> (Bonnevie, 1898)	A

Monocoryne

<i>gigantea</i> (Bonnevie, 1898)	A
----------------------------------	---

Family Cladocorynidiae

<i>Cladocoryne</i>	
<i>floccosa</i> Rotch, 1871	A

Family Cladonematidae

<i>Cladonema</i>	
<i>radiatum</i> Dujardin, 1843	A

Family Corymorphidae

<i>Amathaea</i>	
<i>sarsi</i> (Steenstrup, 1854)	A
<i>Branchiocerianthus</i>	
<i>italicus</i> (Stechow, 1921)	M
<i>norvegicus</i> Brattström, 1957	A
<i>Corymorpha</i>	
<i>abyssalis</i> Broch, 1909	A
<i>annulicornis</i> Sars, 1859	
<i>appelloefi</i> Bonnevie, 1901	A
<i>glacialis</i> Sars, 1859	A
<i>groenlandica</i> (Allman, 1876)	A
<i>nana</i> Alder, 1857	A
<i>nutans</i> Sars, 1835	A M

<i>Eucodonium</i>			
brownii Hartlaub, 1907	A M		
<i>Eugotoea</i>	M		
petalina Margulis, 1989			A
<i>Euphysona</i>	M		
annulata Kramp, 1928			
= <i>Corymorpha annulata</i>			
bigelowi Maas, 1905			
= <i>Corymorpha bigelowi</i>			
<i>Paragotoea</i>	A M		
bathybia Kramp, 1942			
<i>Plotocnide</i>	A M		
borealis Wagner, 1885			
<i>Vannuccia</i>	A		
forbesi (Mayer, 1894)			
Family Corynidae			
<i>Actigia</i>	A M		
vanbenedeni (Hincks, 1869)			
<i>Coryne</i>	A		
brevicornis Bonnevie, 1898			
caespes Allman, 1871	M		
clavata Graeffe, 1883	M		
epizoica Stechow, 1921	M		
filiformis Rees, 1936	A		
fucicola (de Filipi, 1866)	M		
graeffei Jickeli, 1883	M		
hincksi Bonnevie, 1898	A		
muscoidea (Linnaeus, 1761)	A M		
pintneri Schneider, 1897	A M		
pusilla (Gaertner, 1774)	A M		
<i>Dicodonium</i>	M		
adriaticum Graeffe, 1884			
ocellatum (Busch, 1851)	M		
<i>Dipurena</i>	M		
dolichogaster (Haeckel, 1864)			
halterata (Forbes, 1846)	A		
ophiogaster (Haeckel, 1877)	A		
reesi Vannucci, 1956	A M		
simulans Bouillon, 1965	A		
spongicola Anger, 1972	A		
<i>Sarsia</i>	A		
barentsi Linko, 1905			
eximia (Allman, 1859)	A		
geminifera Forbes, 1848	A M		
lovenii (M. Sars, 1846)	A		
occulta Edwards, 1978	A		
pattersoni Haddon, 1886	A		
piriformis Edwards, 1983	A		
princeps (Haeckel, 1879)	A		
producta (Wright, 1858)	A M		
prolifera Forbes, 1848	A M		
striata Edwards, 1983	A		
tubulosa (M. Sars, 1835)	A M		
<i>Sarsiella</i>	A M		
dinema Hartlaub, 1907			
Family Eleutheriidae			
<i>Eleutheria</i>	A M		
claparedei Hartlaub, 1889			
dichotoma De Quatrefages, 1842	A M		
<i>Staurocladia</i>	M		
portmanni Brinckmann, 1964			
Family Euphysonidae			
<i>Euphysona</i>	A M		
aurata Forbes, 1848			
flammea (Linko, 1905)	A M		
obvoluta (Kramp, 1933)	A		
tentaculata Linko, 1905	A		
<i>Gymnognos</i>	A		
crassicornis Bonnevie, 1898			
<i>Siphonohydra</i>	A M		
adriatica Salvini-Plawen, 1966			
Family Halocynthiaidae			
<i>Halocynthia</i>	M		
epizoica Hadzi, 1917			
Family Margelopsidae			
<i>Margelopsis</i>			
haeckeli Hartlaub, 1897			A
hartlaubi Browne, 1903			A
Family Moerisiidae			
<i>Moerisia</i>			
lyonsi Boulenger, 1908			M
pallasi (Derzhavin, 1912)			
<i>Odessa</i>			
maeotica (Ostroumoff, 1896)			
<i>Ostromovia</i>			
inkermanica (Paltschikowa-Ostromova, 1925)			
= <i>Moerisia inkermanica</i>			
Family Paracorynidae			
<i>Paracoryne</i>			
huvei Picard, 1957			M
Family Pennariidae			
<i>Pennaria</i>			
disticha (Goldfuss, 1820)			A
Family Porpitidae			
<i>Porpita</i>			
porpita (Linnaeus, 1758)			A
<i>Velella</i>			
velella (Linnaeus, 1758)			A
Family Protohydridae			
<i>Protohydra</i>			
leuckarti Greeff, 1870			A M
Family Rosalindidae			
<i>Rosalinda</i>			
incrustans (Kramp, 1947)			A M
Family Sphaerocorynidiae			
<i>Sphaerocoryne</i>			
bedoti Pictet, 1893			A
Family Tubulariidae			
<i>Ectopleura</i>			
dumortieri (Van Beneden, 1844)			A
minerva Mayer, 1900			A
petalina (Margulis, 1989)			M
= <i>Egothoea petalina</i>			
sacculifera Kramp, 1957			A
wrighti Petersen, 1979			M
<i>Hybocodon</i>			
prolifer Agassiz, 1862			A M
<i>Rhabdoon</i>			
singulare Keferstein & Ehlers, 1861			M
<i>Tubularia</i>			
bellis Allman, 1865			A
ceratogyné Pérez, 1920			A
cornucopia Bonnevie, 1898			A
crocea (Agassiz, 1862)			A
= <i>Ectopleura crocea</i>			
indivisa Linnaeus, 1758			A M
larynx Ellis & Solander, 1786			A M
mesembryanthemum Allman, 1871			A
regalis Boeck, 1860			
Family Zancleidae			
<i>Zanclea</i>			
costata Gegenbaur, 1857			A M
sessilis (Gosse, 1853)			A M
Order Conida			
Family Aequoreidae			
<i>Aequorea</i>			
aequorea (Forskål, 1775)			M
conica Browne, 1905			M
forskalea (Forskål, 1775)			A
macrodactyla (Brandt, 1835)			A
pensilis (Eschscholtz, 1829)			A M
vitrina Gosse, 1853			A
Family Aglaopheniidae			
<i>Aglaophenia</i>			
acacia Allman, 1883			A M

<i>elongata</i> Meneghini, 1845	A M	<i>huxleyi</i> (Haeckel, 1879)	A
<i>harpago</i> Schenck, 1965	M	<i>Dipleurosoma</i>	
<i>kirchenpaueri</i> (Heller, 1868)	A M	<i>typicum</i> Boeck, 1866	A
<i>latecarinata</i> Allman, 1877	A		
<i>lophocarpa</i> Allman, 1877	A M	Family Eirenidae	
<i>octodonta</i> Heller, 1868	A M	<i>Campanopsis</i>	
<i>parvula</i> Bale, 1882	A	<i>adriaticus</i> (Hadzi, 1915)	M
<i>picardi</i> Svoboda, 1979	A M	= <i>Halanthus adriaticus</i>	
<i>pluma</i> (Linnaeus, 1767)	A M		
<i>tubiformis</i> Marktanner-Turneretscher, 1890	A M	<i>Eirene</i>	
<i>tubulifera</i> (Hincks, 1861)	A M	<i>viridula</i> (Péron & Lesueur, 1809)	A M
Cladocarpus		<i>Eugymnanthea</i>	
<i>bicuspis</i> (Sars, 1873)	A	<i>inquilina</i> Palombi, 1935	M
<i>bonneviaei</i> Jäderholm, 1909	A		
<i>boucheti</i> Ramil & Vervoort, 1992	A	<i>Eutima</i>	
<i>cartieri</i> Bedot, 1921	A	<i>gegenbauri</i> (Haeckel, 1864)	A
<i>cornutus</i> (Verrill, 1879)	A	<i>gracilis</i> (Forbes & Goodsir, 1853)	A M
<i>diana</i> Broch, 1918	A	<i>mira</i> McCrady, 1859	A
<i>formosus</i> Allman, 1877	A		
<i>integer</i> (Sars, 1873)	A	<i>Eutonina</i>	
<i>multiseptatus</i> (Bale, 1915)	A	<i>indicans</i> (Romanes, 1876)	A
<i>paraventricosus</i> Ramil & Vervoort, 1992	A	<i>scintillans</i> (Bigelow, 1909)	A M
<i>sigma</i> (Allman, 1877)	A	<i>Helgicirrrha</i>	
<i>sinuosos</i> Vervoort, 1966		<i>cari</i> (Haeckel, 1864)	A M
Gymnangium		<i>schulzei</i> Hartlaub, 1909	A M
<i>montagui</i> (Billard, 1912)	A M		
<i>sinosum</i> (Fraser, 1925)	A	<i>Irenium</i>	
Lytocarpia		<i>quadrigatum</i> Haeckel, 1879	A
<i>distans</i> (Allman, 1877)	A M		
<i>myriophyllum</i> (Linnaeus, 1758)	A M	Neotima	
<i>phytuma</i> (Kirchenpauer, 1876)	A M	<i>lucullana</i> Petersen	M
Macrorhynchia		Phialopsis	
<i>philippina</i> Kirchenpauer, 1872	A	<i>diegensis</i> Torrey, 1909	A
Nematocarpus			
<i>ramuliferus</i> (Allman, 1874)	A	Tima	
Streptocaulus		<i>bairdii</i> (Johnston, 1833)	A
<i>corneliusi</i> (Ramil & Vervoort, 1992)	A M	<i>flavilabris</i> Eschscholtz, 1829	A M
<i>dollfusi</i> (Billard, 1924)	A M	<i>lucullana</i> (Delle Chiaje, 1822)	M
= <i>Cladocarpus dollfusi</i>		= <i>Neotima lucullana</i>	
<i>pectiniferus</i> (Allman, 1883)	A		
<i>sinuosus</i> (Vervoort, 1966)	A	Family Eucheilotidae	
Family Barcinididae		<i>Eucheilotria</i>	
Barcino		<i>flevensis</i> Van Kampen, 1922	A
<i>foixensis</i> Gili, Bouillon, Pagès, Palanques & Puig, 1999	M	<i>maasi</i> Neppi & Stiasny, 1911	M
Family Blackfordiidae		<i>maculata</i> Hartlaub, 1894	A
Blackfordia		<i>paradoxa</i> Mayer, 1900	M
<i>virginica</i> Mayer, 1910	A M	<i>ventricularis</i> McCrady, 1859	A M
Family Campanulinidae			
Calycella		Family Haleciidae	
<i>gracilis</i> Hartlaub, 1897	A	<i>Campalecium</i>	
<i>syringa</i> (Linnaeus, 1758)	A M	<i>cirratum</i> (Haeckel, 1879)	M
Campomma		<i>medusiferum</i> Torrey, 1902	M
<i>hincksi</i> (Hartlaub, 1897)	A M	<i>microtheca</i> (Hadzi, 1914)	M
Cuspidella		= <i>Haleciella microtheca</i>	
<i>costata</i> Hincks, 1868	A M		
<i>grandis</i> Hincks, 1868		Halecium	
<i>humilis</i> (Alder, 1862)	A M	<i>articulosum</i> Clark, 1875	A
<i>mertensi</i> (Brandt, 1835)	M	<i>banyulense</i> Motz-Kossowska, 1912	M
<i>procumbens</i> Kramp, 1911	A	<i>beanei</i> (Johnston, 1838)	A M
Egmundella		<i>birulai</i> Spassky, 1929	A
<i>amirantensis</i> Millard & Bouillon, 1973	A M	<i>conicum</i> Stechow, 1919	M
<i>grimaldii</i> Leloup, 1940	A	<i>corrugatum</i> Nutting, 1899	A
Lafoeina		<i>curvicaule</i> Lorenz, 1886	A
<i>maxima</i> (Levinsen, 1893)	A	<i>delicatulum</i> Coughtrey, 1876	A
<i>tenuis</i> Sars, 1874	A M	<i>groenlandicum</i> Kramp, 1911	A
Opercularella		<i>halecium</i> (Linnaeus, 1758)	A M
<i>lacerata</i> (Johnston, 1847)	A M	<i>labrosum</i> Alder, 1859	A M
<i>panicula</i> (Sars, 1873)	A M	<i>lankesteri</i> (Bourne, 1890)	A M
<i>pumila</i> Clark, 1875	A	= <i>H. robustum</i> Pieper, 1884	
Tetrapoma		<i>liouvillei</i> Billard, 1934	A
<i>quadridentatum</i> (Hincks, 1874)	A	<i>mediterraneum</i> Weismann, 1883	A M
Family Dipleurosomatidae		<i>minimum</i> Schneider, 1898	M
Cuvieria		<i>minutum</i> Broch, 1903	A
		<i>muricatum</i> (Ellis & Solander, 1786)	A M
		<i>nanum</i> Alder, 1859	A M
		<i>ornatum</i> Nutting, 1901	A
		<i>petrosum</i> Stechow, 1919	A M
		<i>profundum</i> Calder & Vervoort, 1998	A
		<i>pusillum</i> Sars, 1856	A M
		<i>reflexum</i> Stechow, 1919	M
		<i>sessile</i> Norman, 1866	A M
		<i>tenellum</i> Hincks, 1861	A M
		<i>undulatum</i> Billard, 1921	A
		Hydranthea	

<i>aloysii</i> (Zoja, 1893)	M	<i>fruticosa</i> Sars, 1851	A M
<i>diaphana</i> (Hadzi, 1912)	M	= <i>L. dumosa</i>	
<i>margarica</i> (Hincks, 1863)	A M	<i>grandis</i> Hincks, 1874	A M
<i>Hydrodendron</i>		<i>pocillum</i> Hincks, 1868	
<i>gorgonoide</i> (Sars, 1873)	A	<i>Scandia</i>	
<i>mirabile</i> (Hincks, 1866)	A	<i>gigas</i> (Pieper, 1884)	A M
= <i>H. caciniforme</i>		<i>michaelsarsi</i> (Leloup, 1935)	A M
= <i>Ophiodissa mirabile</i>		<i>Zygophylax</i>	
Family Halopterididae		<i>bathypnila</i> Leloup, 1940	A
<i>Antennella</i>		<i>biarmata</i> Billard, 1905	A M
<i>secundaria</i> (Gmelin, 1791)	A M	<i>brownii</i> Billard, 1924	A M
= <i>A. paucinoda</i>		<i>echinata</i> Calder & Vervoort, 1998	
<i>siliquosa</i> (Hincks, 1877)	A M	<i>elegantula</i> Leloup, 1940	A
<i>Halopteris</i>		<i>leloupi</i> Ramil & Vervoort, 1992	A
<i>alternata</i> (Nutting, 1900)	A	<i>levinseni</i> (Saemundsson, 1911)	A
<i>catharina</i> (Johnston, 1833)	A M	= <i>Z. elegantula</i>	
<i>diaphana</i> (Heller, 1868)	A	<i>pinnata</i> (Sars, 1873)	A
<i>glutinosa</i> (Lamouroux, 1816)	A	<i>sibogae</i> Billard, 1918	A
<i>liechtensternii</i> (Marktanner-Turneretscher, 1890)	A M		
<i>Monostacachas</i>		Family Laodiceidae	
<i>quadridens</i> (McCrady, 1859)	A	<i>Krampella</i>	
<i>Schizotricha</i>		<i>dubia</i> Russell, 1957	A M
<i>frutescens</i> (Ellis & Solander, 1786)	A M	<i>tardenti</i> Gili, Bouillon & Pagès, 1998	M
<i>polaris</i> Naumov, 1960	A	<i>Laodicea</i>	
<i>variabilis</i> Bonnevie, 1899	A	<i>bigelowi</i> Neppi & Stiasny, 1911	M
Family Kirchenpaueriidae		<i>chapmani</i> Günther, 1903	A
<i>Kirchenpaueria</i>		<i>neptuna</i> Mayer, 1900	M
<i>echinulata</i> (Lamarck, 1816)	A M	<i>ocellata</i> Babnik, 1948	M
<i>pinnata</i> (Linnaeus, 1758)	A M	<i>undulata</i> (Forbes & Goodsir, 1853)	A M
= <i>K. similis</i>		<i>Melicertissa</i>	
<i>triangulata</i> (Totton, 1930)		<i>adriatica</i> Neppi, 1915	M
= <i>K. bonnevieveae</i>		<i>Ptychogena</i>	
<i>Ophinella</i>		<i>crocea</i> Kramp & Damas, 1925	A
<i>parasitica</i> (Sars, 1873)	A	<i>lactea</i> Agassiz, 1865	A
<i>Ventromma</i>		<i>Staurodiscus</i>	
<i>halecioides</i> (Alder, 1859)	A M	<i>tetraстaurus</i> Haeckel, 1879	
= <i>Kirchenpaueria halecioides</i>		<i>Staurostoma</i>	
Family Lafocidae		<i>mertensi</i> (Brandt, 1838)	A M
<i>Acryptolaria</i>		= <i>Staurophora mertensi</i>	
<i>conferta</i> (Allman, 1877)	A M	<i>Teclalia</i>	
<i>crassicaulis</i> (Allman, 1888)	A	<i>recincola</i> Gili, Bouillon, Pagès, Palanques & Puig, 1999	M
<i>longitheca</i> (Allman, 1877)	A M	<i>Toxorchis</i>	
= <i>A. crassicaulis</i>		<i>arcuatus</i> Haeckel, 1879	
<i>Anthohebella</i>		Family Lovenellidae	
<i>parasitica</i> (Ciamician, 1880)	A M	<i>Lovenella</i>	
= <i>Hebella parasitica</i>		<i>chiquitita</i> Millard, 1959	M
<i>Bedotella</i>		<i>cirrata</i> (Haeckel, 1879)	A M
<i>armata</i> (Pictet & Bedot, 1900)	A	= <i>Campaleicum? cirrata</i>	
<i>Cryptolarella</i>		<i>clausa</i> (Lovén, 1836)	A M
<i>abyssicola</i> (Allman, 1888)	A M	<i>gracilis</i> Clarke, 1882	A M
= <i>C. diffusa</i>		<i>paniculata</i> (Sars, 1873)	A M
= <i>C. humilis</i>		<i>producta</i> (Sars, 1873)	A
<i>Cryptolaria</i>		Family Malagazziidae	
<i>exserta</i> Busk, 1858	A	<i>Octophialucium</i>	
<i>pectinata</i> (Allman, 1888)	A	<i>funerarium</i> (Quoy & Gaimard, 1827)	A M
<i>Filellum</i>		Family Meliceridae	
<i>disaggregatum</i> Peña-Cantero, García-Carrascosa & Vervoort, 1998	M	<i>Melicertum</i>	
<i>serpens</i> (Hassall, 1848)	A M	<i>octocostatum</i> (Sars, 1835)	A
<i>serratum</i> (Clarke, 1879)	A M	<i>Orchistomella</i>	
<i>Grammaria</i>		<i>graeffei</i> (Neppi & Stiasny, 1911)	M
<i>abietina</i> (Sars, 1851)	A	Family Mitrocomidae	
<i>immersa</i> Nutting, 1901	A	<i>Cosmetira</i>	
<i>Hebella</i>		<i>pilosella</i> Forbes, 1848	A
<i>brochi</i> (Hadzi, 1913)	A M	<i>Cyclocanna</i>	
<i>scandens</i> (Bale, 1888)	A M	<i>welshi</i> Bigelow, 1918	A
= <i>H. besnardi</i>		<i>Foersteria</i>	
= <i>H. eximia</i>		<i>antoniae</i> Gili, Bouillon, Pagès, Palanques, Puig & Heussner, 1998	M
= <i>H. michaelseni</i>		<i>ariaiae</i> Gili, Bouillon, Pagès, Palanquea & Puig, 1999.	M
= <i>H. sinuosa</i>		<i>Halopsis</i>	
= <i>H. urceolata</i>		<i>ocellata</i> Agassiz, 1863	A
<i>Lafoea</i>		<i>Mitrocoma</i>	
<i>dumosa</i> (Fleming, 1820)	A M	<i>annae</i> Haeckel, 1864	M
= <i>L. tenellula</i>		<i>Mitrocomella</i>	

<i>brownei</i> (Kramp, 1930)	A M	= <i>D. cornicina</i> McCrady, 1859	
<i>polydiademata</i> (Romanes, 1876)	A	<i>pumila</i> (Linnaeus, 1758)	A M
<i>Mitrocomium</i>		<i>quadridentata</i> (Ellis & Solander, 1786)	A M
<i>cirratum</i> Haeckel, 1879	A M	<i>Hydrallmania</i>	
<i>Octogonade</i>	M	<i>falcata</i> (Linnaeus, 1758)	A
<i>mediterranea</i> Zofja, 1896		<i>Salacia</i>	
= <i>Melicerta adriatica</i> Neppi, 1915	M	<i>desmoides</i> (Torrey, 1902)	A
<i>Tiaropsisidium</i>	A	<i>dubia</i> (Billard, 1922)	A M
<i>atlanticum</i> Russell, 1956	M	<i>Selaginopsis</i>	
<i>mediterraneum</i> (Metschnikoff, 1886)	M	<i>fusca</i> (Johnston, 1847)	A
<i>vilaevelebiti</i> (Hadzi, 1915)	M	<i>Sertularella</i>	
= <i>Halanthus vilaevelebiti</i>		<i>arbuscula</i> (Lamouroux, 1816)	
Family Orchistomidae		<i>crassicaulis</i> (Heller, 1868)	M
<i>Orchistoma</i>	M	<i>cubica</i> García Aguirre & Gonzalez, 1980	M
<i>agariciforme</i> Keller, 1884		<i>cylindritheca</i> (Allman, 1888)	A M
Family Phialellidae		= <i>Cnidoseyphus macrotheca</i>	
<i>Phialella</i>	A M	<i>ellisii</i> (Deshayes & Milne Edwards, 1836)	A M
<i>quadraata</i> (Forbes, 1848)		= <i>S. fusiformis</i>	
Family Plumulariidae		<i>fusiformis</i> (Hincks, 1861)	A M
<i>Monotheca</i>		<i>gayi</i> (Lamouroux, 1821)	A M
<i>margareta</i> Nutting, 1900	A M	<i>mediterranea</i> Hartlaub, 1901	A M
<i>obliqua</i> (Johnston, 1847)	A M	= <i>S. gaudichaudii</i>	
<i>pulchella</i> (Bale, 1882)	A M	<i>ornata</i> Broch, 1933	M
<i>Nematophorus</i>	A	<i>picta</i> (Meyen, 1834)	A M
<i>clarkei</i> Nutting, 1900		<i>polyzonias</i> (Linnaeus, 1758)	A M
= <i>Macrorhynchia clarkei</i>		<i>rugosa</i> (Linnaeus, 1758)	A
<i>Nemertesia</i>		<i>simplex</i> (Hutton, 1873)	A
<i>antennina</i> (Linnaeus, 1758)	A M	<i>tenella</i> (Alder, 1857)	A M
<i>belini</i> Bedot, 1916	A	<i>unituba</i> Calder, 1991	A
<i>falcicula</i> Ramil & Vervoort, 1992	A	<i>Sertularia</i>	
= <i>Plumularia falcicula</i>		<i>albimaris</i> Merezhkovskii, 1878	A
<i>irregularis</i> (Quelch, 1885)	A M	<i>argentea</i> Linnaeus, 1758	A
<i>norvegica</i> (Sars, 1873)	A	<i>cupressina</i> Linnaeus, 1758	A
<i>perrieri</i> (Billard, 1901)	A M	<i>distans</i> Lamouroux, 1816	A M
<i>ramosa</i> (Lamarck, 1816)	A M	<i>fabricii</i> Levinsen, 1893	A
<i>tetrasistica</i> Meneghini, 1845	M	<i>gracilis</i> Hassall, 1848	
<i>ventriculiformis</i> (Marktanner-Turneretscher, 1890)	A M	<i>inflata</i> (Versluys, 1899)	M
<i>Plumularia</i>		<i>marginata</i> (Kirchenpauer, 1864)	A M
<i>canariensis</i> Izquierdo, García-Corrales & Bacallado, 1986		<i>mediterranea</i> (Marktanner-Turneretscher, 1890)	M
<i>filicula</i> Allman, 1877	A M	<i>mirabilis</i> (Verrill, 1872)	A
<i>floridana</i> Nutting, 1900	A	<i>perpusilla</i> Stechow, 1919	M
<i>linkoi</i> Naumov, 1960	M	<i>plumosa</i> (Clark, 1876)	A
<i>obliqua</i> (Johnston, 1847)	A	<i>robusta</i> (Clark, 1876)	A
<i>setacea</i> (Linnaeus, 1758)	A M	<i>schmidti</i> Kudelin, 1914	A
<i>syriaca</i> Billard, 1931	M	<i>tenera</i> Sars, 1874	A
<i>Polyplumaria</i>		<i>turbinata</i> (Lamouroux, 1816)	A
<i>flabellata</i> (Sars, 1873)	A M	<i>Symplectoscyphus</i>	
= <i>P. billardi</i>		<i>amphorifera</i> (Allman, 1877)	A
<i>gracillima</i> (Sars, 1873)	A	<i>bathyalis</i> Vervoort, 1972	A
<i>Pseudoplumaria</i>		<i>tricuspidatus</i> (Alder, 1856)	A
<i>marocana</i> (Billard, 1930)	A M	<i>Tamarisca</i>	
<i>sabinae</i> Ramil & Vervoort, 1992	A	<i>tamarisca</i> (Linnaeus, 1758)	A M
Family Sertulariidae		<i>Thuiaria</i>	
<i>Abietinaria</i>		<i>alternitheca</i> Levinsen, 1893	A
<i>abietina</i> (Linnaeus, 1758)	A M	<i>articulata</i> (Pallas, 1766)	A
<i>filicula</i> (Ellis & Solander, 1786)	A	= <i>Salacia articulata</i>	
<i>interversa</i> (Pictet & Bedot, 1900)	A	<i>breitfussi</i> (Kudelin, 1914)	A
<i>pulchra</i> (Nutting, 1904)	A	<i>carica</i> Levinsen, 1893	A
<i>Amphisbeta</i>		<i>cupressoides</i> (Lepechin, 1783)	A
<i>operculata</i> (Linnaeus, 1758)	A M	<i>decemserialis</i> (Merezhkovskii, 1878)	A
<i>Diphasia</i>		<i>hippuris</i> Allman, 1874	A
<i>alata</i> (Hincks, 1855)	A	<i>lichenastrum</i> (Pallas, 1766)	A
<i>attenuata</i> (Hincks, 1866)	A M	<i>obsoleta</i> (Lepechin, 1781)	A
<i>delagei</i> (Billard, 1912)	A	<i>thuja</i> (Linnaeus, 1758)	A
<i>fallax</i> (Johnston, 1847)	A	<i>variabilis</i> (Broch, 1918)	A
<i>margareta</i> (Hassall, 1841)	A M	Family Syntheciidae	
<i>nigra</i> (Pallas, 1766)	A	<i>Synthecium</i>	
<i>pinastrum</i> (Cuvier, 1830)	A M	<i>evansi</i> (Ellis & Solander, 1786)	A M
<i>rosacea</i> (Linnaeus, 1758)	A M	Family Tiaranidae	
<i>Dynamena</i>		<i>Modeeria</i>	
<i>crisiodes</i> Lamouroux, 1824	A	<i>rotunda</i> (Quoy & Gaimard, 1827)	A M
<i>dalmasi</i> (Versluys, 1899)	A	= <i>Stegopoma fastigiata</i> (Alder, 1860)	
<i>disticha</i> (Bosc, 1802)	A	<i>Stegolaria</i>	
		<i>geniculata</i> (Allman, 1888)	
		<i>Stegopoma</i>	A M

<i>bathyale</i> Vervoort, 1966	A	<i>multicornis</i> (Forskål, 1775)	A M
<i>giganteum</i> Ramil & Vervoort, 1992	A	= <i>C. squamata</i>	
<i>plicatile</i> (Sars, 1863)	A	<i>Cordylophora</i>	
<i>Tiaranna</i>		<i>caspia</i> (Pallas, 1771)	A
<i>ducalis</i> (Forbes & Goodsir, 1853)		<i>inkermanica</i> Marfenin, 1983	M
<i>globulosa</i> (Forbes, 1848)		<i>pusilla</i> Motz-Kossowska, 1905	M
Family Tiaropsidae		<i>Corydendrium</i>	
<i>Tiaropsis</i>	A	<i>dispar</i> Kramp, 1935	A
<i>multicirrata</i> (Sars, 1835)		<i>parasiticum</i> (Linnaeus, 1767)	A
Order Filifera		<i>Merona</i>	
Family Bougainvilliidae		<i>cornucopiae</i> Norman, 1864	A
<i>Aselomaris</i>	A M	<i>ibera</i> Medel, García-Gómez & Bouillon, 1993	M
<i>arenosa</i> (Alder, 1863)	A M	<i>nematophorum</i> (Antsulevich & Polteva, 1986)	A
<i>michaeli</i> Berrill, 1948		<i>Oceania</i>	
= <i>Clavopsella michaeli</i>		<i>armata</i> Källiker, 1853	A
<i>Bimeria</i>	A M	<i>Tubiclava</i>	
<i>vestita</i> Wright, 1859		<i>fruticosa</i> Allman, 1872	A
<i>Bougainvilia</i>	A M	<i>lucerna</i> Allman, 1863	A M
<i>britannica</i> (Forbes, 1841)		<i>Turritopsis</i>	
<i>charcoti</i> Le Danois, 1913		<i>nutricula</i> McCrady, 1857	A
<i>flavida</i> Hartlaub, 1887		= <i>Corydendrium nutricula</i>	
<i>macloviana</i> Lesson, 1836	A	Family Cytaeididae	
<i>maniculata</i> Haeckel, 1864	A M	<i>Cytaeis</i>	
<i>multicilia</i> (Haeckel, 1879)	M	<i>pusilla</i> Gegenbaur, 1856	M
<i>muscoidea</i> (Sars, 1846)	A	<i>tetrastyla</i> Eschscholtz, 1829	A
<i>muscus</i> (Van Beneden, 1844)	A	<i>Perarella</i>	
= <i>B. ramosa</i>	A	<i>propagulata</i> Bavestrello, 1987	M
<i>principis</i> (Steenstrup & Lütken, 1850)	A	<i>schneideri</i> (Motz-Kossowska, 1905)	M
<i>pyramidata</i> (Forbes & Goodsir, 1853)	A	Family Eudendriidae	
<i>rugosa</i> Clarke, 1882	A	<i>Eudendrium</i>	
<i>supercliaris</i> (Agassiz, 1849)	A	<i>album</i> Nutting, 1898	A
<i>Clavopsella</i>	A	<i>annulatum</i> Norman, 1864	A
<i>navis</i> (Millard, 1959)		<i>arbusculum</i> Wright, 1859	A M
<i>Dicoryne</i>	A M	<i>armatum</i> Tichomiroff, 1890	A M
<i>conferta</i> (Alder, 1856)	A M	<i>calceolatum</i> Motz-Kossowska, 1905	M
<i>corybearei</i> (Browne, 1907)	A	<i>capillare</i> Alder, 1856	A M
<i>flexuosa</i> Sars, 1873	A	<i>carneum</i> Clarke, 1882	A
<i>Garveia</i>	A	<i>cunninghami</i> Kirkpatrick, 1910	A M
<i>biscayana</i> (Browne, 1907)	A	<i>fragile</i> Motz-Kossowska, 1905	M
<i>franciscana</i> (Torrey, 1902)	A	<i>glomeratum</i> Picard, 1951	A
<i>grisea</i> (Motz-Kossowska, 1905)	M	<i>insigne</i> Hincks, 1861	A M
<i>nutans</i> Wright, 1859	A M	<i>merulum</i> Watson, 1985	M
<i>Koellikerina</i>	A M	<i>motzkossowskiae</i> Picard, 1951	M
<i>fasciculata</i> (Péron & Lesueur, 1809)		<i>planum</i> Bonnevie, 1898	A
<i>Lizzella</i>	A M	<i>racemosum</i> (Cavolini, 1785)	A M
<i>hyalina</i> (Van Beneden, 1866)	A	<i>rameum</i> (Pallas, 1766)	A M
<i>Lizzia</i>	A M	<i>ramosum</i> (Linnaeus, 1758)	A M
<i>blondina</i> Forbes, 1848	A M	<i>teissieri</i> Cabioch, 1970	A
<i>fulgorans</i> (Agassiz, 1865)	A M	<i>tenellum</i> Allman, 1877	A
<i>octostyla</i> (Haeckel, 1879)	M	<i>vaginatum</i> Allman, 1863	A
<i>Nemopsis</i>	A	= <i>E. annulatum</i>	
<i>bachei</i> Agassiz, 1849	A	<i>vervoorti</i> Marques & Migotto, 1998	A
<i>crucifera</i> (Forbes & Goodsir, 1853)	A	<i>Myrionema</i>	
<i>Nubella</i>	M	<i>amboinense</i> Pictet, 1893	A
<i>mitra</i> Bouillon, 1980		Family Hydractiniidae	
<i>Pachycordyle</i>	A M	<i>Hydractinia</i>	
<i>napolitana</i> Weismann, 1883		<i>aculeata</i> (Wagner, 1833)	M
<i>Rhizorhagium</i>	A	<i>allmani</i> Bonnevie, 1898	A
<i>album</i> Rees, 1938	A M	<i>arctica</i> (Jäderholm, 1902)	A
<i>formosum</i> (Fewkes, 1889)	A	= <i>H. ingolfi</i>	
<i>roseum</i> Sars, 1874	A	= <i>Stylocarpia arctica</i>	
<i>sarsii</i> (Bonnevie, 1898)	A	<i>calderi</i> Bouillon, Medel & Peña-Cantero, 1997	M
<i>Thamnostoma</i>	M	<i>carica</i> Berg, 1887	A
<i>dibalium</i> (Busch, 1851)	A	<i>echinata</i> Fleming, 1823	A M
<i>russelli</i> Rees, 1939		<i>fucicola</i> (Sars, 1857)	A M
Family Calycopsidae		<i>humilis</i> Bonnevie, 1898	A
<i>Bythiotira</i>	A	<i>monocarpa</i> Allman, 1847	A
<i>murrayi</i> Günther, 1903		<i>ornata</i> Bonnevie, 1899	A M
<i>Calycopsis</i>	A	<i>serrata</i> Kramp, 1943	A
<i>gara</i> Petersen, 1957	A	<i>Podocoryna</i>	
<i>krampi</i> Petersen, 1957	A	<i>areolata</i> (Alder, 1862)	A M
<i>simplex</i> Kramp & Damas, 1925	A M	= <i>P. harilaubi</i>	
Family Clavidae		<i>borealis</i> (Mayer, 1900)	A M
<i>Clava</i>		<i>carnea</i> Sars, 1846	A M

<i>exigua</i> (Haeckel, 1879)	M	<i>affinis</i> Moseley, 1879	A
<i>minima</i> (Trinci, 1903)	A M	<i>medioatlantica</i> Zibrowius & Cairns, 1992	A
<i>minuta</i> (Mayer, 1900)	M	<i>tenuiseptata</i> Cairns, 1986	A
<i>proboscidea</i> Hincks, 1869	A	<i>vascomarquesi</i> Zibrowius & Cairns, 1992	A
<i>sarsi</i> Steenstrup, 1850	A	<i>Errina</i>	
<i>Stylocystis</i>		<i>aspera</i> (Linnaeus, 1767)	A M
<i>claviformis</i> Bouillon, 1971	A	<i>atlantica</i> Hickson, 1912	A
<i>fuciola</i> (Sars, 1857)	M	<i>dabneyi</i> (Pourtalès, 1871)	A
= <i>Hydractinia fuciola</i>		<i>Lepidopora</i>	
<i>inermis</i> (Allman, 1872)	M	<i>eburnea</i> (Calvet, 1903)	A
<i>pruvoti</i> (Motz-Kossowska, 1905)	A M	<i>Pliobothrus</i>	
= <i>Hydractomma pruvoti</i>		<i>gracilis</i> Zibrowius & Cairns, 1992	A
<i>Tregoubovia</i>		<i>symmetricus</i> Pourtalès, 1868	A
<i>atentaculata</i> Picard, 1958	M	<i>Stenohelia</i>	
Family Hydrichthyidae		<i>maderensis</i> (Johnston, 1862)	A
<i>Hydrichthys</i>		<i>Stylaster</i>	
<i>cyclothonis</i> Damas, 1934	A	<i>erubescens</i> subsp. <i>britanicus</i> Zibrowius & Cairns, 1992	A
<i>sarcotretis</i> (Jungersen, 1911)	A	<i>erubescens</i> subsp. <i>groenlandicus</i> Zibrowius & Cairns, 1992	A
= <i>Ichthyocodium sarcotretis</i>		<i>erubescens</i> subsp. <i>meterorensis</i> Zibrowius & Cairns, 1992	A
Family Niobiidae		<i>gemmascens</i> (Esper, 1794)	A
<i>Niobia</i>		<i>ibericus</i> Zibrowius & Cairns, 1992	A
<i>dendrotentaculata</i> Mayer, 1900	A	<i>maroccanus</i> Zibrowius & Cairns, 1992	A
Family Pandidae		<i>norvegicus</i> (Gunnerus, 1768)	A
<i>Amphinema</i>		Family Trichydridae	
<i>dinema</i> (Péron & Lesueur, 1810)	A	<i>Trichydra</i>	
<i>krampi</i> Russell, 1956	A	<i>pudica</i> Wright, 1858	A
<i>rubrum</i> (Kramp, 1957)	M	Order Laingiomedusae	
<i>rugosum</i> (Mayer, 1900)	A	Family Laingiidae	
<i>turridum</i> (Mayer, 1900)	A	<i>Kantiella</i>	
<i>Annatiara</i>		<i>enigmatica</i> Bouillon, 1978	M
<i>affinis</i> (Hartlaub, 1914)	A	Order Limnopolypae	
<i>Catablema</i>		Family Armorhydridae	
<i>vesicarium</i> (A. Agassiz, 1862)	A	<i>Armorhydra</i>	
<i>Codonorchis</i>		<i>janowiczi</i> Swedmark & Teissier, 1958	A M
<i>octaedrus</i> Haeckel, 1879	A M	Family Microhydrulidae	
<i>Halitholus</i>		<i>Microhydrula</i>	
<i>pauper</i> Hartlaub, 1913	A	<i>pontica</i> Valkanov, 1965	A M
<i>yoldiaearticatae</i> (Birula, 1897)	A	<i>Rhaptapagis</i>	
<i>Leuckartiara</i>		<i>cantacuzenei</i> Bouillon & Deroux, 1967	A
<i>abyssi</i> (G.O. Sars, 1874)	A	Family Olindiasidae	
<i>breviconis</i> (Murbach & Shearer, 1902)	A	<i>Craspedacusta</i>	
<i>nobilis</i> Hartlaub, 1913	A M	<i>sowerbyi</i> Lankester, 1880	A M
<i>octona</i> (Fleming, 1823)	A	<i>Gonionemus</i>	
<i>Merga</i>		<i>vertens</i> A. Agassiz, 1862	A
<i>galleri</i> Brinckmann, 1962	M	<i>vindobonensis</i> Joseph, 1918	M
<i>reesi</i> Russell, 1956	A	<i>Gossea</i>	
<i>tergesina</i> (Neppi & Stiasny, 1912)	M	<i>corynetes</i> (Gosse, 1853)	A M
<i>tregoubovii</i> Picard, 1960	M	<i>faureae</i> Picard, 1952	A
<i>Neoturris</i>		<i>Maeotias</i>	
<i>pileata</i> (Forskål, 1775)	A M	<i>inexpectata</i> Ostroumov, 1896	A M
<i>Octotiara</i>		<i>Monobrachium</i>	
<i>violacea</i> Kramp, 1959	A	<i>parasiticum</i> Mereschkowsky, 1877	A M
<i>Pandea</i>		<i>Olindias</i>	
<i>conica</i> (Quoy & Gaimard, 1827)	A	<i>phosphorica</i> (Delle Chiaje, 1841)	A M
<i>rubra</i> Bigelow, 1913	A	Family Proboscidactylidae	
<i>Stomotoca</i>		<i>Fabienna</i>	
<i>georginae</i> (Hadzi, 1913)	M	<i>oligonema</i> (Kramp, 1953)	A M
Family Protiaridae		= <i>Trichydra oligonema</i>	
<i>Halitiara</i>		<i>Proboscidactyla</i>	
<i>formosa</i> Fewkes, 1882	A	<i>flavicirrata</i> Brandt, 1835	A
<i>Paratiara</i>		<i>ornata</i> (McCrady, 1859)	A
<i>digitalis</i> Kramp & Damus, 1925	A	<i>stellata</i> (Forbes, 1846)	A
<i>Protiera</i>		Order Narcomedusae	
<i>tetranema</i> (Péron & Lesueur, 1809)	A M	Family Aeginidae	
Family Ptilocodiidae		<i>Aegina</i>	
<i>Thecocodium</i>		<i>citrea</i> Eschscholtz, 1829	A M
<i>brieni</i> Bouillon, 1967	A M	= <i>A. rosea</i>	
<i>penicillatum</i> Jarms, 1987	A	<i>Aeginopsis</i>	
Family Rathkeidae		<i>laurentii</i> Brandt, 1838	A
<i>Rathkea</i>			
<i>octopunctata</i> (Sars, 1835)	A M		
Family Stylasteridae			
<i>Cryptelia</i>			

<i>Aeginura</i>			
<i>grimaldii</i> Maas, 1904	A	<i>everta</i> (Clark, 1876)	M
<i>Solmundella</i>		<i>integra</i> (MacGillivray, 1842)	A M
<i>bitentaculata</i> (Quoy & Gaimard, 1833)	A	<i>Phialidium</i>	
Family Cuninidae		<i>discoideum</i> (Mayer, 1900)	M
<i>Cunina</i>		= <i>Clytia discoideum</i>	
<i>fowleri</i> (Browne, 1906)		<i>folleatum</i> McCrady, 1859	A M
= <i>Cunoctantha fowleri</i>		= <i>P. pentata</i>	
<i>frugifera</i> Kramp, 1948	A M	<i>islandicum</i> Kramp, 1919	A
<i>globosa</i> Eschscholtz, 1829	A	<i>mccradyi</i> (Brooks, 1888)	M
<i>lativentris</i> Gegenbaur, 1856	A	<i>Rhizocaulus</i>	
<i>octonaria</i> McCrady, 1857	A	<i>verticillatus</i> (Linnaeus, 1758)	A
<i>polygonia</i> (Haeckel, 1879)	M	<i>Tulpa</i>	
<i>proboscidea</i> E. & L. Metchnikoff, 1871	M	<i>speciosa</i> (Clark, 1876)	A
<i>simplex</i> Gili, Bouillon, Pagès, Palanques, Puig & Heussner, 1998	M		
<i>vitrea</i> Gegenbaur, 1856	M	Order Trachymedusae	
<i>Solmissus</i>	A M	Family Geryoniidae	
<i>albescens</i> (Gegenbaur, 1856)	A	<i>Geryonia</i>	
<i>incisus</i> (Fewkes, 1886)	A	<i>proboscidalis</i> (Huxley, 1859)	A
Family Solmarisidae		<i>Liriope</i>	
<i>Pegantha</i>		<i>tetraphylla</i> (Otto, 1823)	A
<i>clara</i> Bigelow, 1909	A	Family Halicreatidae	
<i>molicina</i> (Forskål, 1775)	M	<i>Botrynema</i>	
<i>rubiginosa</i> (Kölliker, 1853)	A M	<i>brucei</i> Browne, 1908	A
<i>triloba</i> Haeckel, 1879	A	<i>ellinorae</i> (Hartlaub, 1909)	A
<i>zonaria</i> (Haeckel, 1879)	M	<i>Halicreas</i>	
<i>Solmaris</i>	A	<i>minimum</i> Fewkes, 1882	A
<i>corona</i> (Keferstein & Ehlers, 1861)	M	<i>papillosum</i> Vanhöffen, 1902	A
<i>flavescens</i> (Kölliker, 1853)	M	<i>Haliscera</i>	
<i>leucostyla</i> (Will, 1844)	M	<i>bigelowi</i> Kramp, 1947	A M
<i>solmaris</i> (Gegenbaur, 1856)	M	<i>conica</i> Vanhöffen, 1902	A
<i>vanhoeffeni</i> Neppi & Stiasny, 1911	M	<i>racovitzae</i> (Maas, 1906)	M
		= <i>Homoeonema racovitzae</i>	
Order Proboscoida		<i>Haliotrephes</i>	
		<i>maasi</i> Bigelow, 1909	A
Family Bonneviellidae		<i>Homoeonema</i>	
<i>Bonneviella</i>		<i>platygonum</i> Browne, 1903	A
<i>grandis</i> (Allman, 1876)	A		
Family Campanulariidae		Family Petasidae	
<i>Campanularia</i>		<i>Petasus</i>	
<i>crenata</i> (Hartlaub, 1901)	M	<i>atavus</i> Haeckel, 1879	A M
<i>hinchsii</i> Alder, 1856	A M		
<i>volubilis</i> (Linnaeus, 1758)	A M	Family Ptychogastriidae	
<i>Clytia</i>		<i>Ptychogastria</i>	
<i>gracilis</i> (Sars, 1850)	A M	<i>asteroides</i> (Haeckel, 1879)	M
<i>hemisphaerica</i> (Linnaeus, 1767)	A M	<i>polaris</i> Allman, 1878	A
<i>islandicum</i> Kramp, 1919	A	<i>Tesserogastria</i>	
<i>linearis</i> (Thorneley, 1900)	A	<i>musculosa</i> Beyer, 1959	A
<i>maccradyi</i> (Brooks, 1888)	A		
<i>noliformis</i> (McCrady, 1859)	A M	Family Rhopalonestidae	
<i>paulensis</i> (Vanhöffen, 1910)	A M	<i>Aglantha</i>	
<i>pentata</i> (Mayer, 1900)	A	<i>digitale</i> (Müller, 1776)	A M
<i>raridentata</i> (Alder, 1862)	M	<i>elata</i> (Haeckel, 1879)	M
<i>serrulata</i> (Bale, 1888)	M	<i>Aglaura</i>	
<i>Gonothyraea</i>	A M	<i>hemistoma</i> Péron & Le Sueur, 1809	A
<i>loveni</i> (Allman, 1859)	A M	<i>Arctapodema</i>	
<i>Hartlaubella</i>	A M	<i>ampla</i> (Vanhöffen, 1902)	A
<i>gelatinosa</i> (Pallas, 1766)	A M	<i>australe</i> (Vanhöffen, 1912)	M
<i>Laomedea</i>	A M	<i>Colobonema</i>	
<i>angulata</i> Hincks, 1861	A M	<i>sericeum</i> Vanhöffen, 1902	A
<i>calceolifera</i> (Hincks, 1871)	A M	<i>Crossota</i>	
<i>exigua</i> Sars, 1857	A M	<i>norvegica</i> Vanhöffen, 1902	A
<i>flexuosa</i> Alder, 1857	A M	<i>rufobrunnea</i> (Kramp, 1913)	A
<i>neglecta</i> Alder, 1856	A M	<i>Homeonema</i>	
<i>pseudodichotoma</i> Vervoort, 1959	A	<i>platygonon</i> Browne, 1903	A M
<i>Obelia</i>	A M	<i>Pantachogon</i>	
<i>bidentata</i> Clark, 1875	A M	<i>haeckeli</i> Maas, 1893	A M
<i>dichotoma</i> (Linnaeus, 1758)	A M	<i>militare</i> (Maas, 1893)	A M
<i>fimbriata</i> (Dalyell, 1848)	A M	<i>Persa</i>	
<i>geniculata</i> (Linnaeus, 1758)	A M	<i>incolorata</i> McCrady, 1857	A
<i>longissima</i> (Pallas, 1766)	A M	<i>Ransonnia</i>	
<i>Orthopyxis</i>	M	<i>krampi</i> (Ranson, 1932)	A M
<i>asymmaterica</i> Stechow, 1919	A	<i>Rhopalonema</i>	
<i>crenata</i> Hartlaub, 1901	A	<i>funerarium</i> Vanhöffen, 1902	A
= <i>Campanularia crenata</i>		<i>velatum</i> Gegenbaur, 1856	A
		<i>Sminthea</i>	
		<i>arctica</i> Hartlaub, 1909	A
		<i>eurygaster</i> Gegenbaur, 1856	A

Tetrorchis
erythrogaster Bigelow, 1909

A

Siphonophora

Compiled by G. M. Mapstone

This list was initially drawn up from 'Siphonophores and Velellids' by P. A. Kirkpatrick and P. R. Pugh (1984), and additional species added from other publications covering species recorded within the ERMS geographic area. The reference list includes all taxonomic papers covering any new and redescribed species found in the area, but only those distributional papers published from 1990 onwards. A more comprehensive literature list will be obtainable from a monograph entitled 'Siphonophora (Cnidaria, Hydrozoa) of Canadian Pacific waters.' by G. M. Mapstone and M. N. Arai which is due to be published shortly.

Since the species list is a reflection of sampling effort, it is likely that more species will be added in the future; many Siphonophora species have a cosmopolitan distribution.

P. R. Pugh kindly advised on the validity of species included in a world list from which this ERMS was derived.

References

- Andersen, V., Sardou, J. and Nival, P. 1992. The diel migrations and vertical distributions of zooplankton and micronekton in the Northwestern Mediterranean Sea. 2. Siphonophores, hydromedusae and pyrosomids. *J. Plankton Res.* 8, 1155-1169.
- Bamstedt, U., Fossa, J. H., Martinussen, M. B. and Fossagen, A. 1998. Mass occurrence of the physonect siphonophore *Apolemia uvaria* (Lesueur) in Norwegian waters. *Sarsia* 83, 79-85.
- Buecher, E. 1999. Appearance of *Chelophyses appendiculata* and *Abylopsis tetragona* (Cnidaria, Siphonophora) in the Bay of Villefranche, northwestern Mediterranean. *Journal of Sea Research* 41, 295-307.
- Biggs, D. C., Pugh P. R. and Carré, C. 1978. *Rosacea flaccida* n. sp., a new species of siphonophore (Calycophorae Prayinae) from the North Atlantic Ocean. *Beaufortia* 27: 207-218.
- Gamulin, T. and Krsinic, F. 1993. Distribution and abundance of Calycophores (Siphonophora, Calycophorae) in the Mediterranean and Adriatic Sea. *Mar. Ecol. (Pubbl. Stn. Zool. Napoli I)*, 14 (2): 97-111.
- Gili, J-M., Pagès, F. and Fusté, X. 1991. Mesoscale coupling between spatial distribution of planktonic cnidarians and hydrographic features along the Galician Coast (Northwestern Iberian Peninsula). *Sci. Mar.* 55(2): 419-426.
- Mapstone, G. M. 1998. *Bargmannia lata*, An undescribed species of physonect siphonophore (Cnidaria, Hydrozoa) from Canadian Pacific waters. 141-147, figs 1-3. In *Commemorative volume for the 80th birthday of Willem Vervoort in 1997*. Edited by: J. C. den Hartog, A. C. van Bruggen, P. F. S. Cornelius and L. P. van Ofwegen. *Zool. Verh. Leiden*: 323: i-xii, 1-448.
- Mills, C. E., Pugh, P. R., Harbison, G. R., and Haddock, S. H. D. 1996. Medusae, siphonophores and ctenophores of the Alborán Sea, south western Mediterranean. In *Advances in Hydrozoan Biology*. Edited by S. Piraino., F. Boero, J. Bouillon, P. F. S. Cornelius and J.-M. Gili. *Sci. Mar.* 60 (1): 145-163.
- Kirkpatrick, P. A. and Pugh, P. R. 1984. Siphonophores and velellids. *Synopsis of the British Fauna* (New Series). Leiden, E. J. BrillNumber 29, Pp. 154.
- Pagès, F., González, H. E., and González, S. R. 1996. Diet of the gelatinous zooplankton in Hardangerfjord (Norway) and potential predatory impact by *Aglantha digitale* (Trachymedusae). *Mar. Ecol. Prog. Ser.* 139: 69-77.
- Pinca, S., and Dallot, S. 1995. Meso- and macrozooplankton composition patterns related to hydrodynamic structures in

the Ligurian Sea (Trophos-2 experiment, April-June 1986). <i>Mar. Ecol. Prog. Ser.</i> 126: 49-65.		
Pugh, P. R. 1983. Benthic siphonophores: a review of the family Rhodaliidae (Siphonophora, Physonectae). <i>Philos. Trans. R. Soc. Lon. B Biol. Sci.</i> 301: 165-300.		
Pugh, P. R. 1992a. A revision of the sub-family Nectopyramidinae Siphonophora, Prayidae. <i>Philos. Trans. R. Soc. Lon. B Biol. Sci.</i> 335: 281-322.		
Pugh, P. R. 1992b. The status of the genus <i>Prayoides</i> (Siphonophora, Prayidae). <i>J. Mar. Biol. Asso. U.K.</i> 72 (4): 895-909.		
Pugh, P. R. 1992c. <i>Desmophyes haematogaster</i> , a new species of prayine siphonophore (Calycophorae, Prayidae). <i>Bull. Mar. Sci.</i> 50 (1): 89-96.		
Pugh, P. R. 1998. A re-description of <i>Frillagalma vitiazi</i> Daniel 1966 (Siphonophorae, Agalmatidae). <i>Sci. Mar.</i> 62 (3): 233-245.		
Pugh, P. R. 1999. A review of the genus <i>Bargmannia</i> Totton, 1954 (Siphonophorae, Physonecta, Pyrostephidae). <i>Bull. Mar. Hist. Mus. Lond. (Zool.)</i> 65 (1): 51-72.		
Pugh, P. R. and Harbison, G. R. 1987. Three new species of prayine siphonophore (Calycophorae, Prayidae) collected by a submersible, with notes on related species. <i>Bull. Mar. Sci.</i> 41 (1): 68-91.		
Pugh, P. R. and Pagès, F. 1993. A new species of <i>Clausophyes</i> (Siphonophorae, Clausophyidae), with a redescription of <i>C. galeata</i> and <i>C. moserae</i> . <i>J. Mar. Biol. Assoc. U.K.</i> 73 (3): 595-608.		
Pugh, P. R. and Pagès, F. 1995. Is <i>Lensia reticulata</i> a diphyine species (Siphonophorae, Calycophora, Diphyidae)? A re-description. <i>Sci. Mar.</i> 59 (2): 181-192.		
Pugh, P. R. and Youngbluth, M. J. 1988a. A new species of <i>Halistemma</i> (Siphonophora: Physonectae: Agalmidae) collected by submersible. <i>J. Mar. Biol. Asso. U.K.</i> 68: 1-14.		
Pugh, P. R. and Youngbluth, M. J. 1988b. Two new species of prayine siphonophore (Calycophorae, Prayidae) collected by the submersibles Johnson-Sea-Link I and II. <i>J. Plankton Res.</i> 10 (4): 637-657.		
Totton, A. K. 1965. <i>A Synopsis of the Siphonophora</i> . London, British Museum (Natural History). Pp. 230.		
Subclass Siphonophora		
Order Calycophorae		
Family Abylidiae		
<i>Abyla</i>		
bicarinata Moser, 1925	A	
trigona Quoy & Gaimard, 1827	A	
<i>Abylopsis</i>		
tetragona (Otto, 1823)	A M	
<i>Bassia</i>		
bassensis (Quoy & Gaimard, 1833)	A M	
<i>Ceratocymba</i>		
dentata (Bigelow, 1918)	A M	
sagittata (Quoy & Gaimard, 1827)	A	
<i>Enneagonum</i>		
hyalinum Quoy & Gaimard, 1827	A M	
Family Clausophyidae		
<i>Chuniphyes</i>		
moserae Totton, 1954	A M	
multidentata Lens & van Riemsdijk, 1908	A M	
<i>Clausophyes</i>		
galeata Lens & van Riemsdijk, 1908		
moserae Margulis, 1988	A M	
ovata (Keferstein & Ehlers, 1860)	A M	
= <i>Clausophyes massiliiana</i>		
<i>Crystallophyses</i>		
amygdalina Moser, 1925	A	
<i>Heteropyramis</i>		
crystallina (Moser, 1925)	A	
maculata Moser, 1925	A	
Family Diphyidae		
<i>Chelophyses</i>		
appendiculata (Eschscholtz, 1829)	A M	
<i>Dimophyes</i>		
<i>arctica</i> (Chun, 1897)		A M
<i>Diphyes</i>		
bojani (Eschscholtz, 1829)		A M
dispar Chamisso & Eysenhardt, 1821		A M
<i>Eudoxia</i>		
macra Totton, 1954		A M
tenuis Patriti, 1965		A
<i>Eodoxoides</i>		
mitra (Huxley, 1859)		A M
spiralis (Bigelow, 1911)		A M
<i>Lensia</i>		
achilles Totton, 1941		A
conoidea (Keferstein & Ehlers, 1860)		A M
cossack Totton, 1941		A M
exeter Totton, 1941		A
fowleri (Bigelow, 1911)		A M
grimaldi Leloup, 1933		A
havock Totton, 1941		A M
hostile Totton, 1941		A M
hotspur Totton, 1941		A
lelouvetean Totton, 1941		A
meteori (Leloup, 1934)		A M
multicristata (Moser, 1925)		A M
patritii Alekseyev, 1984		A
reticulata Totton, 1954		A M
subtilis (Chun, 1886)		A M
subtiloides (Lens & van Riemsdijk, 1908)		A M
<i>Muggiaeaa</i>		
atlantica Cunningham, 1892		A M
bargmannae Totton, 1954		
cantabrica Alcazar, 1982		A
kochi (Will, 1844)		A
<i>Sulculeolaria</i>		
biloba (Sars, 1846)		A
chuni (Lens & van Riemsdijk, 1908)		A
quadrivalvis Blainville, 1834		A
turgida (Gegenbaur, 1853)		A M
Family Hippopodiidae		
<i>Hippopodus</i>		
hippopodus (Forskål, 1776)		A
<i>Voglia</i>		
glabra Bigelow, 1918		A M
pentacantha Kölliker, 1853		A M
serrata (Moser, 1925)		A M
spinosa Keferstein & Ehlers, 1861		A
Family Prayidae		
<i>Amphicaryon</i>		
acaule Chun, 1888		A
ernesti Totton, 1954		A M
peltifera (Haeckel, 1888)		A
<i>Desmophyes</i>		
annectens Haeckel, 1888		A M
villafrancae (Carré, 1969)		M
= <i>Rosacea villafrancae</i>		
<i>Lilyopsis</i>		
rosea Chun, 1885		A M
<i>Maresearsia</i>		
praeclara Totton, 1954		A M
<i>Nectadamas</i>		
diomedae (Bigelow, 1911)		A M
richardi Pugh, 1992		A
<i>Nectopyramis</i>		
natans (Bigelow, 1911)		A M
= <i>Nectopyramis spinosa</i>		
thetis Bigelow, 1911		A
= <i>Nectopyramis spinosa</i>		
<i>Praya</i>		
dubia (Quoy & Gaimard in de Blainville, 1830)		A M
= <i>Prayoides intermedia</i>		
reticulata (Bigelow, 1911)		A M
<i>Prayola</i>		
tottoni Carré, 1969		M
<i>Rosacea</i>		
cymbiformis (delle Chiaje, 1830)		A M
plicata sensu Bigelow, 1911		A M

CTENOPHORA	
A	Compiled by Jacob van der Land
M	The European list was derived from URMO mainly based on Harbison <i>et al.</i> (1978) and Harbison & Madin (1982) with additions from the Zoological Record. There does not appear to be an expert on the European species.
A M	References
M	Harbison, G. R., L. P. Madin, and N. R. Swanberg, 1978. On the natural history and distribution of oceanic ctenophores. <i>Deep-Sea Research</i> 25: 233-256. Harbison, G.R. and L.P. Madin, 1982. Ctenophora. In: S.P. Parker (ed.), <i>Synopsis and Classification of Living Organisms</i> , vol. I: 707-715. McGraw-Hill, New York, etc.
PHYLUM CTENOPHORA	
A	Order Beroida
A M	Family Beroidae
A M	<i>Beroe</i> <i>cucumis</i> Fabricius, 1780 <i>forskali</i> Milne Edwards, 1841 <i>gracilis</i> Künne, 1939 <i>ovata</i> Bruguière, 1789
A M	Order Cestida
A M	Family Cestidae
A M	<i>Cestum</i> <i>veneris</i> Lesueur, 1813
A M	<i>Velamen</i> <i>parallelum</i> (Fol, 1869)
A M	Order Cydippida
A M	Family Ctenellidae
A M	<i>Ctenella</i> <i>aurantia</i> Carré & Carré, 1993
A M	Family Dryodoridae
A M	<i>Dryodora</i> <i>glandiformis</i> (Mertens, 1833)
A M	Family Haeckeliidae
A M	<i>Haeckelia</i> <i>beehleri</i> (Mayer, 1912) <i>bimaculata</i> Carré & Carré, 1989 <i>filigera</i> (Chun, 1880) <i>rubra</i> (Gegenbaur, Kölliker & Müller, 1853)
A M	Family Lampeidae
A M	<i>Lampea</i> <i>pancerina</i> (Chun, 1879) = <i>Gastrodres parasiticum</i> (juv.)
A M	Family Mertensiidae
A M	<i>Callianira</i> <i>bialata</i> Delle Chiaje, 1841 <i>ficalbi</i> Curreri, 1900
A M	<i>Charistephane</i> <i>fugiens</i> Chun, 1880
A M	<i>Mertensia</i> <i>ovum</i> (Fabricius, 1780) = <i>Mertensia cucullus</i> L. Agassiz, 1860
A M	Family Pleurobrachiidae
A M	<i>Ceroctena</i> <i>bicornis</i> Carré & Carré, 1991
A M	<i>Euplokamis</i> <i>crinita</i> (Moser, 1909) = <i>Pleurobrachia crinita</i>
A M	<i>octoplera</i> (Mertens, 1833) <i>stationis</i> Chun, 1880
A M	<i>Hormiphora</i> <i>cucumis</i> (Mertens, 1833) = <i>Beroe cucumis</i> Fabricius, 1789 p.p. <i>plumosa</i> L. Agassiz, 1860
A M	<i>Minicetena</i> <i>luteola</i> Carré & Carré, 1993
A M	<i>Pleurobrachia</i>

<i>pileus</i> (Müller, 1776)	A
<i>rhododactyla</i> L. Agassiz, 1860	M
<i>rhodopis</i> Chun, 1880	
<i>Sabaudia</i>	
<i>liguriae</i> Ghighi, 1909	
<i>Tinere</i>	
<i>cyanæa</i> (Chun, 1889)	
Order Lobata	
Family Bolinopsidae	
<i>Bolinopsis</i>	
<i>infundibulum</i> (Müller, 1776)	A
= <i>Bolinopsis alata</i>	
= <i>Bolinopsis microptera</i>	
= <i>Bolinopsis septentrionalis</i>	
<i>vitrea</i> (L. Agassiz, 1860)	M
= <i>Bolinia hydatina</i>	
<i>Lesueuria</i>	
<i>vitrea</i> Milne Edwards, 1841	
<i>Leucothea</i>	
<i>multicornis</i> (Quoy & Gaimard, 1824)	A
Family Deiopidae	
<i>Deiopea</i>	
<i>kaloktenota</i> Chun, 1879	M
Family Eurhamphaeidae	
<i>Eurhamphaea</i>	
<i>vexilligera</i> Gegenbaur, 1856	A M
Family Mnemiidae	
<i>Mnemiopsis</i>	
<i>leidyi</i> A. Agassiz, 1865	
Family Ocyropsidae	
<i>Ocyropsis</i>	
<i>maculata</i> (Rang, 1826)	
Order Platycetenida	
Family Platycetenidae	
<i>Vallicula</i>	
<i>multiformis</i> Rankin, 1956	A

A M

PLATYHELMINTHES

Turbellaria

Compiled by A. Faubel & C. Noreña

Turbellarians are ubiquitous specimens depending on more or less water saturated habitats of marine, brackish, freshwater, and moist terrestrial areas. Dominant areas of distribution are marine and freshwater ones but there are many species living in brackish water areas and transition zones - the ecotones - between land and sea. In the Turbellaria list the focus is set only on species having its dominant distribution in marine habitats and species being immigrants into poly- and mesohaline habitats. Therefore, all species reported from estuaries and the Baltic Proper, in particular, from the Gulf of Finland and the Bothnian Sea, are excluded from the species list if they are immigrants from freshwater habitats and are not distributed in areas with salinities of greater than 5.0.

The first complete review of Turbellaria of the world was by von Graff (1904-08). More recent comprehensive accounts on marine Turbellaria were published by Bock 1913 (Polycladida), Meixner 1938 (Proseriata), Westblad 1942, 1945, 1946, 1948 (Acoela), Luther 1955, 1960, 1962, 1963 (Acoela, Macrostomida, Rhabdocoela), Karling 1931, 1947, 1949, 1950, 1952, 1963 (Kalyptorhynchia), 1958 (Proseriata), 1940, 1968 (Prolecithophora, Rhabdocoela), 1974 (overview of the Turbellaria of the Baltic Proper), Ax 1956a (Proseriata), de Beauchamp 1961 (Turbellaria), Dörjes 1968, 1971 (Acoela), Dörjes & Karling 1975 (Acoela), Rieger 1971a,b (Macrostomida), Faubel 1983, 1984 (Polycladida), and Prudhoe 1985 (Polycladida). In addition, comprehensive studies with focus on taxonomy and faunistic aspects of species on European coasts are as following: North Sea inclusive English Channel (Ax 1960, Den Hartog 1964a, b, 1968, Schilke 1970a,b, Sopott 1972, Ehlers 1974, Faubel 1974a,b, 1977, Hoxhold 1974, Martens 1983), the Irish Sea (Boaden 1963a, b, 1966), the French Atlantic coast (Ehlers 1980, Sopott-Ehlers 1976, L'Hardy 1963, 1965), the Mediterranean Sea (Riedl 1954, 1956, 1959a, b, c, Ax 1956b, Brunet 1965, 1966, 1970, 1972, 1979), the Black Sea (Ax 1959, Mack-Fira 1973, 1974), and the Canary Islands (Ehlers & Ehlers 1979, Sopott-Ehlers 1980).

Up to 1999 a total of 3224 valid marine and brackish-water species of Turbellaria are known world-wide. On European coasts and sublittoral areas 1137 (35%) valid species have been described.

References

- Ax, P., 1956a: *Monographie der Otoplanidae (Turbellaria), Morphologie und Systematik*. Akademie der Wissenschaft und der Literatur Mainz, Abhandlungen der mathematisch-naturwissenschaftlichen Klasse 13, 499-796.
- Ax, P., 1956b: Les Turbellariés des étangs côtiers du littoral méditerranéen de la France méridionale. *Vie et Milieu*, Supplément 5, 1-152.
- Ax, P., 1959: Zur Systematik, Ökologie und Tiergeographie der Turbellarienfauna in den ponto-kaspischen Brackwassermeeren. *Zoologisches Jahrbuch, Abteilung für Systematik, Ökologie und Geographie der Tiere* 87, 43-184.
- Ax, P., 1960: Turbellarien aus salzdurchtränkten Wiesenböden der deutschen Meeresküsten. *Zeitschrift für wissenschaftliche Zoologie* 163, 210-235.
- Beauchamp, P. de, 1961: Classe des Turbellariés, Turbellaria (Ehrenberg, 1832). In: *Traité de Zoologie*, Grasse, P.-P. (ed.), Masson et Cie, Paris, 4 (1), 35-212.

- Boaden, P.J.S., 1963a: The interstitial fauna of some North Wales beaches. *Journal of the marine biological Association of the United Kingdom* 43, 79-96.
- Boaden, P.J.S., 1963b: The interstitial Turbellaria Kalyptorhynchia from some North Wales beaches. *Proceedings of the Zoological Society of London* 141, 173-205.
- Boaden, P.J.S., 1966: Interstitial fauna from Northern Ireland. *Veröffentlichungen des Institutes für Meeresforschung Bremerhaven*, Sonderband 2, 125-130.
- Bock, S., 1913: Studien über Polycladen. *Zoologiska bidrag från Uppsala* 2, 31-344.
- Brunet, M., 1965: Turbellariés Calyptorhynques de substrats meubles de la région de Marseille. *Recueil des Travaux de la Station Marine d'Endoume / Bulletin* 39, 127-219.
- Brunet, M., 1966: Sur quelque Turbellariés Calyptorhynques de sables fins de la région de Marseille. *Bulletin de la Société zoologique de France* 91, 721-730.
- Brunet, M., 1970: Turbellariés Schizorhynchidae de sables infralittoraux de la côte marseillaise. *Cahiers de Biologie marine* 11, 279-306.
- Brunet, M., 1972: Koinocystididae de la région de Marseille (Turbellaria, Kalyptorhynchia). *Zoologica Scripta* 1, 157-174.
- Brunet, M., 1979: Turbellariés Calyptorhynques du Golfe de Marseille. *Revue de Biologie et Ecologie méditerranéenne* 6, 101-120.
- Den Hartog, C., 1964a: Proseriate flatworms from the deltaic area of the rivers Rhine, Meuse and Scheldt. II. *Proceedings. Koninklijke Nederlands Akademie van Wetenschappen* C67, 23-34.
- Den Hartog, C., 1964b: Proseriate flatworms from the deltaic area of the rivers Rhine, Meuse, and Scheldt I. *Proceedings. Koninklijke Nederlands Akademie van Wetenschappen* C67, 10-22.
- Den Hartog, C., 1968: Marine triclad from the Plymouth area. *Journal of the marine biological Association of the United Kingdom* 48, 209-223.
- Dörjes, J., 1968: Die Acoela (Turbellaria) der deutschen Nordseeküste und ein neues System der Ordnung. *Zeitschrift für zoologische Systematik und Evolutionsforschung* 6, 56-452.
- Dörjes, J., 1971: Monographie der Proporidae und Solenifilomorphidae (Turbellaria, Acoela). *Senckenbergiana biologica* 52, 113-137.
- Dörjes, J. & Karling, T.G., 1975: Species of Turbellaria Acoela in the Swedish Museum of Natural History, with remarks on their anatomy, taxonomy and distribution. *Zoologica Scripta* 4, 175-189.
- Ehlers, U., 1974: Interstitialle Typhloplanoida (Turbellaria) aus dem Litoral der Nordseeinsel Sylt. *Mikrofauna des Meeresboden* 49, 425-527.
- Ehlers, U., 1980: Interstitial Typhloplanoida (Turbellaria) from the area of Roscoff. *Cahiers de Biologie marine* 21, 155-167.
- Ehlers, B. & Ehlers, U., 1979: Zur Systematik und geographischen Verbreitung interstitieller Turbellarien der Kanarischen Inseln. *Mikrofauna des Meeresboden* 80, 571-593.
- Faibl, A., 1974a: Die Acoela (Turbellaria) eines Sandstrandes der Nordseeinsel Sylt. *Mikrofauna des Meeresboden* 32, 17-74.
- Faibl, A., 1974b: Macrostomida (Turbellaria) von einem Sandstrand der Nordseeinsel Sylt. *Mikrofauna des Meeresboden* 45, 339-370.
- Faibl, A., 1977: The distribution of Acoela and Macrostomida (Turbellaria) in the Litoral of the North Frisian Islands Sylt, Römö, Jordsand and Amrum (North Sea). *Senckenbergiana maritima* 9, 59-74.
- Faibl, A., 1983: The Polycladida, Turbellaria. Proposal and establishment of a new system. Part. I. The Acotylea. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 80, 17-121.
- Faibl, A., 1984: The Polycladida, Turbellaria. Proposal and establishment of a new system. Part II. The Cotylea.
- Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 81, 189-259.
- Graff, L. v., 1904-08: Turbellaria. In: Dr. H.G. Bronns *Klassen und Ordnungen des Tierreichs*. Bd IV. Vermes. Abt. I c. Turbellaria I. Abteilung: Acoela und Rhabdocoelida. Winter'sche Verlagsbuchhandlung Leipzig, 1733-2599pp.
- Hoxhold, S., 1974: Zur Populationsstruktur und Abundanzdynamik interstitieller Kalyptorhynchia (Turbellaria, Neorhabdocoela). *Mikrofauna des Meeresboden* 41, 1-134.
- Karling, T.G., 1931: Untersuchungen über Kalyptorhynchia (Turbellaria, Rhabdocoela) aus dem Brackwasser des Finnischen Meerbusens. *Acta Zoologica Fennica* 11, 1-66.
- Karling, T.G., 1940: Zur Morphologie und Systematik der Alloeocoela cumulata und Rhabdocoela Lecithophora (Turbellaria). *Acta Zoologica Fennica* 26, 1-260.
- Karling, T.G., 1947: Studien über Kalyptorhynchien (Turbellaria). I. Die Familien Placorhynchidae und Gnathorhynchidae. *Acta Zoologica Fennica* 50, 1-64.
- Karling, T.G., 1949: Studien über Kalyptorhynchien (Turbellaria). II. Die Familien Karkinorhynchidae und Diascorhynchidae. *Acta Zoologica Fennica* 58, 1-42.
- Karling, T.G., 1950: Studien über Kalyptorhynchien (Turbellaria). III. Die Familie Schizorhynchidae. *Acta Zoologica Fennica* 59, 1-33.
- Karling, T.G., 1952: Studien über Kalyptorhynchien (Turbellaria). IV. Einige Eukalyptorhynchia. *Acta Zoologica Fennica* 69, 1-49.
- Karling, T.G., 1958: Zur Kenntnis der Gattung *Coelogynopora* Steinböck (Turbellaria, Proseriata). *Arkiv för Zoologi* 11, 559-567.
- Karling, T.G., 1963: Die Turbellarien Ostfennoskandiens. V. Neorhabdocoela 3. Kalyptorhynchia. *Fauna Fennica* 17, 1-59.
- Karling, T.G., 1968: On the genus *Gnosonesima* Reisinger (Turbellaria). *Sarsia* 33, 81-108.
- Karling, T.G., 1974: Turbellarian fauna of the Baltic proper. Identification, ecology and biogeography. *Fauna Fennica* 27, 1-101.
- L'Hardy, J.-P., 1963: Turbellariés Schizorhynchidae des sables de Roscoff. I. Les genres *Carcharodorhynchus*, *Schizochilus* et *Schizorhynchides*. *Cahiers de Biologie marine* 4, 459-472.
- L'Hardy, J.-P., 1965: Turbellariés Schizorhynchidae des sables de Roscoff. II. Le genre *Proschizorhynchus*. *Cahiers de Biologie marine* 6, 135-161.
- Luther, A., 1955: Die Dalyelliiden (Turbellaria Neorhabdocoela). *Acta Zoologica Fennica* 87, 1-337.
- Luther, A., 1960: Die Turbellarien Ostfennoskandiens I. Acoela, Catenulida, Macrostomida, Lecithoepitheliata, Prolecithophora, Proseriata. *Fauna Fennica* 7, 1-155.
- Luther, A., 1962: Die Turbellarien Ostfennoskandiens III. Neorhabdocoela I. Dalyelliida, Typhloplanida: Byrsophlebiidae, Trigonostomidae. *Fauna Fennica* 12, 1-71.
- Luther, A., 1963: Die Turbellarien Ostfennoskandiens. IV. Neorhabdocoela 2. Typhloplanida: Typhloplanidae, Selenopharyngidae, Carcharodopharyngidae. *Fauna Fennica* 16, 1-163.
- Mack-Fira, V., 1973: Donnees nouvelles sur la faune de Turbellaries du littoral Roumain (Mer Noire). *Rapport et Procès-verbaux des Réunions, Commission internationale pour l'exploration scientifique de la mer Méditerranée*, Monaco 22, 1-79.
- Mack-Fira, V., 1974: The turbellarian fauna of the Romanian littoral waters of the Black Sea and its annexes. In: *Biology of the Turbellaria*, Riser, N.W. & M.P. Morse (eds.). McGraw Hill Book Co., New York, St.Louis, San Francisco, Düsseldorf, Johannesburg, Kuala Lumpur, 248-290pp.
- Martens, P.M., 1983: Three new species of Minoninae (Turbellaria, Proseriata, Monocelididae) from the North Sea, with remarks on the taxonomy of the subfamily. *Zoologica Scripta* 12, 153-160.

Meixner, J., 1938: Turbellaria (Strudelwürmer). <i>Die Tierwelt der Nord- und Ostsee</i> , Grimpe & Wagler Verlag, Leipzig, VIB, 1-146.		A
Prudhoe, S., 1985: A monograph on polyclad Turbellaria. British Museum (Natural History), Oxford University Press, Oxford, 259 pp.		M
Riedl, R., 1954: Neue Turbellarien aus dem mediterranen Felslitoral. Ergebnisse der Unterwasser-Expedition-Austria 1948-49. <i>Zoologisches Jahrbuch, Abteilung für Systematik, Ökologie und Geographie der Tiere</i> 82, 157-404.		M
Riedl, R., 1956: Zur Kenntnis der Turbellarien adriatischer Schlammböden sowie ihre geographische und faunistische Verbreitung. <i>Thalassia Jugoslavia</i> 1, 69-184.		
Riedl, R., 1959a: Turbellarien aus submarinen Höhlen, 3. Seriata und Neorhabdocoela. <i>Pubblicazioni della Stazioni zoologica di Napoli</i> , Suppl. 30, 305-332.		
Riedl, R., 1959b: Turbellarien aus submarinen Höhlen. I. Archophora. Ergebnisse der Österreichischen Tyrrenia-Expedition 1952, Teil VII. <i>Pubblicazioni della Stazioni zoologica di Napoli</i> , Suppl. 30, 178-208.		
Riedl, R., 1959c: Turbellarien aus submarinen Höhlen. 2. Prolecithophora. <i>Pubblicazioni della Stazioni zoologica di Napoli</i> , Suppl. 30, 209-251.		
Rieger, R.M., 1971b: Die Turbellarienfamilie Dolichomacrostomidae Rieger. II. Teil. Dolichomacrostominae I. <i>Zoologisches Jahrbuch, Abteilung für Systematik, Ökologie und Geographie der Tiere</i> 98, 569-703.		
Rieger, R.M., 1971a: Die Turbellarienfamilie Dolichomacrostomidae nov. fam. (Macrostomida). I. Teil. Vorbemerkungen und Karlinginae nov. subfam. I. <i>Zoologisches Jahrbuch, Abteilung für Systematik, Ökologie und Geographie der Tiere</i> 98, 236-314.		
Schilke, K., 1970a: Morphologie und Phylogenie der Schizorhynchia (Turbellaria, Kalyptorhynchia). <i>Zeitschrift für Morphologie der Tiere</i> 67, 118-171.		
Schilke, K., 1970b: Kalyptorhynchia (Turbellaria) aus dem Eulitoral der deutschen Nordseeküste. <i>Helgoländer wissenschaftliche Meeresuntersuchungen</i> 21, 143-265.		
Sopott, B., 1972: Systematik und Ökologie von Proseriaten (Turbellaria) der deutschen Nordseeküste. <i>Mikrofauna des Meeresboden</i> 13, 1-72.		
Sopott-Ehlers, B., 1976: Interstitielle Macrostomida und Proseriata (Turbellaria) von der französischen Atlantikküste und den Kanarischen Inseln. <i>Mikrofauna des Meeresboden</i> 60, 1-35.		
Sopott-Ehlers, B. & Ehlers, U., 1980: Zur Systematik und geographischen Verbreitung interstitieller Turbellarien der Kanarischen Inseln. <i>Mikrofauna des Meeresboden</i> 80, 1-23.		
Westblad, E., 1942: Studien über skandinavische Turbellarien. Acoela II. <i>Arkiv för Zoologi</i> 33A, 1-48.		
Westblad, E., 1945: Studien über skandinavische Turbellarien. Acoela III. <i>Arkiv för Zoologi</i> 36A, 1-56.		
Westblad, E., 1946: Studien über skandinavische Turbellarien. Acoela IV. <i>Arkiv för Zoologi</i> 38A, 1-56.		
Westblad, E., 1948: Studien über skandinavische Turbellarien. Acoela. V. <i>Arkiv för Zoologi</i> 41 A, 1-82.		
Class Turbellaria		
Order Acoela		
Family Anaperidae		
<i>Achoerus</i>		
<i>pachycaudatus</i> Dörjes, 1968	A	
<i>Anaperus</i>		
<i>biaculeatus</i> Boguta, 1970	A	
<i>rubellus</i> Westblad, 1945	A M	
<i>tvaerminnensis</i> Luther, 1912 Westblad, 1955	A	
<i>Paranaperus</i>		
<i>pellucidus</i> Westblad, 1942	A M	
<i>Philactochoerus</i>		
<i>johanni</i> Dörjes, 1968	A	
<i>Praeanaperus</i>		
<i>tetraposthis</i> Faubel & Regier, 1983	A	
Family Antigonariidae		
<i>Antigonaria</i>		
<i>arenaria</i> Dörjes, 1968		A
Family Antroposthidae		
<i>Adenopea</i>		
<i>chuni</i> (Brauner, 1920) Antonius, 1968		M
<i>illardata</i> Löhner & Micoletzky, 1911		M
<i>Antroposthia</i>		
<i>axi</i> Faubel, 1974		A
<i>unipora</i> Faubel, 1974		A
<i>Convoluella</i>		
<i>brunea</i> Faubel, 1974		A
<i>Unantra</i>		
<i>polyvacuola</i> Faubel, 1976		A
Family Childiidae		
<i>Actinoposthia</i>		
<i>beklemishevi</i> Mamkaev, 1965		A
<i>biaculeata</i> Faubel, 1974		A
<i>haplovata</i> Dörjes, 1968		A
<i>longa</i> Faubel, 1976		A
<i>pigmentea</i> Faubel, 1976		A
<i>Adeloposthia</i>		
<i>elegans</i> An Der Lan, 1936		A
<i>Archactinoposthia</i>		
<i>pelophila</i> Dörjes, 1968		A
<i>Atriofronta</i>		
<i>polyvacuola</i> Dörjes, 1968		A
<i>Childia</i>		
<i>groenlandica</i> (Levinsen, 1879) Meixner, 1925		A M
<i>Microposthia</i>		
<i>listensis</i> Faubel, 1974		A
<i>Paraphanostoma</i>		
<i>brachypodium</i> Westblad, 1942		A M
<i>crassum</i> Westblad, 1942		M
<i>cyclopodium</i> Westblad, 1942		A M
<i>dubium</i> Westblad, 1942		A M
<i>gracilis</i> Westblad, 1945		A
<i>leptopodium</i> Riedl, 1956		M
<i>macropodium</i> (Steinböck, 1931) Westblad, 1942		A M
<i>submaculatum</i> Westblad, 1942		A M
<i>trianguliferum</i> Westblad, 1942		A M
<i>Paraproporus</i>		
<i>diovatus</i> Dörjes, 1968		A
<i>elegans</i> (An Der Lan, 1936) Dörjes, 1968		A
<i>rosettiformis</i> Faubel, 1974		A
<i>rubescens</i> Westblad, 1945		A M
<i>Pelophila</i>		
<i>lutheri</i> (Westblad, 1946) Dörjes & Karling, 1975		A M
<i>pachymorpha</i> Dörjes, 1968		A
<i>Philactinoposthia</i>		
<i>adenogonoria</i> Dörjes, 1968		A
<i>diplopodium</i> Dörjes, 1968		A
<i>helgolandica</i> Dörjes, 1968		A
<i>pusilla</i> (Westblad, 1946) Dörjes, 1968		A
<i>rhammifera</i> (Westblad, 1946) Dörjes, 1968		A
<i>saliens</i> (v. Graff, 1882) Dörjes, 1968		A
<i>stylifera</i> (Westblad, 1946) Dörjes & Karling, 1975		A
<i>viridipunctata</i> (Westblad, 1946) Dörjes & Karling, 1975		A M
<i>viridis</i> Dörjes, 1968		A
<i>viridorhabditi</i> Dörjes, 1968		A
<i>Pseudactinoposthia</i>		
<i>granaria</i> Dörjes, 1968		A
<i>saltans</i> Dörjes, 1968		A
Family Convolutidae		
<i>Alluna</i>		
<i>sublitoralis</i> Faubel & Regier, 1983		A
<i>Amphiscolops</i>		
<i>cinereus</i> (v. Graff, 1874) v. Graff, 1905		A M
<i>fulgineus</i> Peebles, 1915		M
<i>langerhansi</i> (v. Graff, 1882) v. Graff, 1905		A M
<i>mosaicus</i> Kozloff, 1998		M
<i>castellanensis</i> Steinböck, 1954		M
<i>gerundensis</i> Steinböck, 1954		M
<i>zeii</i> Riedl, 1956		M
<i>Aphanostoma</i>		

<i>adherens</i> Meixner, 1938	A	<i>Diopisthoporus</i>	
<i>album</i> Dörjes, 1968	A	<i>brachypharyngeus</i> Dörjes, 1968	A
<i>cavernosum</i> Meixner, 1938	A	<i>longitubus</i> Westblad, 1940	A M
<i>elegans</i> Jensen, 1878	A	<i>psammophilus</i> Dörjes, 1968	A
<i>fuscum</i> Meixner, 1938	A		
<i>latissimum</i> Levinsen, 1879	A		
<i>macrobursalis</i> Westblad, 1946	A		
<i>virescens</i> Örsted, 1845	A		
Archaphanostoma			
<i>agile</i> (Jensen, 1878) Dörjes, 1968	A M		
<i>histobursalis</i> Dörjes, 1968	A		
<i>macrospiriferum</i> (Westblad, 1946) Dörjes, 1968	A		
Avagina			
<i>glandulifera</i> Westblad, 1953	A		
<i>incola</i> Leiper, 1902	A		
<i>sublitoralis</i> Faubel, 1976	A		
Baltalimania	M		
<i>kosswigi</i> Ax, 1959			
Bursosaphia			
<i>baltalimaniaformis</i> Dörjes, 1968	A		
Conaperta			
<i>norvegica</i> (Westblad, 1946) Antonius, 1968	A		
<i>variomorpha</i> (Dörjes, 1968) Antonius, 1968	A		
Convoluta			
<i>albomaculata</i> (Pereyaslawzewska, 1893) v. Graff, 1904	M		
<i>baltica</i> Meixner, 1938	A		
<i>böhmigi</i> (Brauner, 1920) Westblad, 1948	M		
<i>borealis</i> Sabussow, 1900	A		
<i>confusa</i> v. Graff, 1904	M		
<i>convoluta</i> (Abildgaard, 1806) v. Graff, 1905	A M		
<i>diplopisthia</i> Steinböck, 1931	A		
<i>elegans</i> Pereyaslawzewska, 1893	M		
<i>flavibacillum</i> Jensen, 1878	A M		
<i>helgolandica</i> Dörjes, 1966	A		
<i>hipparchia</i> Pereyaslawzewska, 1893	M		
<i>lacazi</i> v. Graff, 1891	A		
<i>pelagica</i> Löhner & Micoletzky, 1911	M		
<i>sordida</i> v. Graff, 1882	A M		
<i>subtilis</i> (v. Graff, 1882) v. Graff, 1891	M		
<i>uljanini</i> v. Graff, 1904	M		
<i>variabilis</i> (Pereyaslawzewska, 1893) v. Graff, 1904	M		
Faerlea			
<i>echinocardii</i> Dörjes, 1972	A		
<i>fragilis</i> Westblad, 1945	A		
<i>glomerata</i> Westblad, 1945	A		
Haplodiscus	M		
<i>ussovi</i> Sabussow, 1896			
Postaphanostoma			
<i>atriomagnatum</i> Dörjes, 1968	A		
<i>filum</i> Dörjes, 1968	A		
<i>glandulosum</i> Dörjes, 1968	A		
Praeaphanostoma			
<i>brevifrons</i> Dörjes, 1968	A		
<i>chaetocaudatum</i> Dörjes, 1968	A		
<i>longum</i> Dörjes, 1968	A		
<i>parvum</i> Rieger & Ott, 1971	M		
<i>rubrum</i> Dörjes, 1968	A		
<i>sizilianum</i> Riedl, 1954	M		
Praeconvoluta			
<i>karinae</i> Dörjes, 1968	A		
<i>minor</i> Faubel, 1974	A		
<i>schmidti</i> Faubel, 1977	A		
<i>stephania</i> Faubel & Regier, 1983	A		
Proaphanostoma			
<i>tenuissimum</i> (Westblad, 1946) Dörjes, 1972	A M		
Proconvoluta			
<i>primitiva</i> Dörjes, 1968	A		
Pseudaphanostoma			
<i>brevicaudatum</i> Dörjes, 1968	A		
<i>pelophilum</i> Dörjes, 1968	A		
<i>psammophilum</i> Dörjes, 1968	A		
<i>variabilis</i> Westblad, 1946	A M		
Pseudopisthia			
<i>macrogonopora</i> Westblad, 1946	A		
Family Diopisthoporidae			
		<i>Diopisthoporus</i>	
		<i>brachypharyngeus</i> Dörjes, 1968	A
		<i>longitubus</i> Westblad, 1940	A M
		<i>psammophilus</i> Dörjes, 1968	A
		Family Hallangidae	
		<i>Hallangia</i>	
		<i>proporoides</i> Westblad, 1946	A M
		Family Haploposthiidae	
		<i>Adenocauda</i>	
		<i>helgolandica</i> Dörjes, 1968	A
		<i>Afronta</i>	
		<i>rubra</i> Faubel, 1974	A
		<i>Haplogonaria</i>	
		<i>arenaria</i> (Ax, 1959) Dörjes, 1968	M
		<i>elegans</i> Faubel, 1974	A
		<i>glandulifera</i> Dörjes, 1968	A
		<i>macrobursalis</i> Dörjes, 1968	A
		<i>minima</i> (Westblad, 1946) Dörjes, 1968	A M
		<i>psammalia</i> Faubel, 1974	A
		<i>simplex</i> Dörjes, 1968	A
		<i>sinubursalia</i> Dörjes, 1968	A
		<i>syltensis</i> Dörjes, 1968	A
		<i>viridis</i> Dörjes, 1968	A
		Haplopisthia	
		<i>brunea</i> An Der Lan, 1936	A
		<i>monogonophora</i> Westblad, 1945	A
		<i>rubra</i> (An Der Lan, 1936) Westblad, 1945	A M
		<i>rubropunctata</i> Westblad, 1945	A M
		Kuma	
		<i>monogonophora</i> (Westblad, 1946) Faubel, 1976	A
		<i>viridis</i> (An Der Lan, 1936) Faubel, 1976	A M
		Parahaplogonaria	
		<i>maxima</i> Dörjes, 1968	A
		Parahaplopisthia	
		<i>avesicula</i> Dörjes, 1968	A
		<i>brunea</i> Faubel, 1974	A
		<i>cerebroepitheliala</i> Dörjes, 1968	A
		Pseudohaplogonaria	
		<i>vacua</i> Dörjes, 1968	A
		<i>viridipunctata</i> (Westblad, 1946) Dörjes, 1968	A M
		Simplicomorpha	
		<i>gigantorhabditis</i> Dörjes, 1968	A
		Family Hofsteniidae	
		<i>Hofsteniola</i>	
		<i>pardii</i> Papi, 1957	M
		Family Mecynostomidae	
		<i>Eumecynostomum</i>	
		<i>altitudi</i> Faubel & Regier, 1983	A
		<i>bathycolum</i> (Westblad, 1946) Faubel & Regier, 1983	A
		<i>boreale</i> (Faubel, 1977) Faubel & Regier, 1983	A
		<i>flavescens</i> (Westblad, 1946) Faubel & Regier, 1983	A
		<i>fragilis</i> (Dörjes, 1968) Faubel & Regier, 1983	A
		<i>juistensis</i> (Dörjes, 1968) Faubel & Regier, 1983	A
		<i>macrobursarium</i> (Westblad, 1946) Faubel & Regier, 1983	A
		<i>maritimum</i> (Dörjes, 1968) Faubel & Regier, 1983	A
		<i>pallidum</i> (Beklemishev, 1915) Faubel & Regier, 1983	A
		<i>papillosum</i> (Faubel, 1974) Faubel & Regier, 1983	A
		<i>sanguineum</i> (Beklemishev, 1915) Faubel & Regier, 1983	A
		<i>westbladi</i> (Dörjes, 1968) Faubel & Regier, 1983	A
		Mecynostomum	
		<i>auritum</i> (Schultze, 1851) Karling, 1974	A M
		<i>haplovarium</i> Dörjes, 1968	A
		<i>predatum</i> Faubel, 1976	A
		<i>sicilianum</i> Riedl, 1954	M
		Neomecynostomum	
		<i>granulum</i> (Dörjes, 1968) Faubel & Regier, 1983	A
		Paedomecynostomum	
		<i>bruneum</i> Dörjes, 1968	A
		Paramecynostomum	
		<i>carchedonium</i> Kozloff, 1998	M
		<i>diversicolor</i> (Örsted, 1845) Dörjes, 1968	A M
		<i>Philomecynostomum</i>	

<i>lapillum</i> Dörjes, 1968	A	Order Haplopharyngida
<i>Postmecynostomum</i>		
<i>glandulosum</i> (Westblad, 1946) Karling & Dörjes, 1975	A	
<i>pictum</i> Dörjes, 1968	A	
<i>Pseudmecynostomum</i>		
<i>boreale</i> Faubel, 1977	A	
<i>bruneophilum</i> Faubel, 1974	A	
<i>bruneum</i> Dörjes, 1968	A	
<i>pelophilum</i> Dörjes, 1968	A	
Family Nadinidae		
<i>Nadina</i>		
<i>pulchella</i> Uljanin, 1870	M	
Family Otocelididae		
<i>Archocelis</i>		
<i>macrorhabdites</i> Dörjes, 1968	A	
<i>Haplotes</i>		
<i>curvitubus</i> Dörjes, 1968	A	
<i>Notocelis</i>		
<i>gullmarense</i> (Westblad, 1946) Dörjes, 1968	A M	
<i>subsalina</i> (Ax, 1959) Dörjes, 1968	M	
<i>Otocelis</i>		
<i>rubropunctata</i> (Schmidt, 1852) Diesing, 1862	A M	
<i>westbladi</i> Ax, 1959	A	
<i>Philocelis</i>		
<i>cellata</i> Dörjes, 1968	A	
<i>karlingi</i> (Westblad, 1946) Dörjes & Karling, 1979	A	
Family Paratomellidae		
<i>Paratomella</i>		
<i>rubra</i> Rieger & Ott, 1971	A M	
<i>unichaeta</i> Dörjes, 1966	A	
Family Proporidae		
<i>Proporus</i>		
<i>brochi</i> Westblad, 1945	A M	
<i>lonchitis</i> Dörjes, 1971	A	
<i>minimus</i> (An Der Lan, 1936) Westblad, 1945	A	
<i>venenosus</i> (Schmidt, 1852) v. Graff, 1882	M	
Family Sagittiferidae		
<i>Simsagittifera</i>		
<i>bifoveolata</i> (Mamkaev, 1971) Mamkaev & Kostenko, 1990	M	
<i>poenicea</i> Kozloff, 1998	M	
<i>psammophila</i> (Beklemishev, 1957) Kostenko & Mamkaev, 1990	M	
<i>roscoffensis</i> (v. Graff, 1891) Kostenko & Mamkaev, 1990	A	
<i>schultzei</i> (Schmidt, 1852) Kostenko & Mamkaev, 1990	A M	
Family Solenofilomorphidae		
<i>Oligofilomorpha</i>		
<i>interstitioniphilum</i> Faubel, 1974	A	
<i>karlingi</i> Dörjes, 1971	A	
<i>Solenofilomorpha</i>		
<i>longissima</i> Dörjes, 1968	A	
Family Taurididae		
<i>Taurida</i>		
<i>fulvomaculata</i> (Ax, 1959) Kostenko, 1989	M	
Order Catenulida		
Family Retronectidae		
<i>Retronectes</i>		
<i>clio</i> Sterrer & Rieger, 1974	A	
<i>melpomene</i> Sterrer & Rieger, 1974	A	
<i>sterreri</i> Faubel, 1976	A	
<i>terpsychore</i> Sterrer & Rieger, 1974	A	
Family Stenostomidae		
<i>Stenostomum</i>		
<i>karlingi</i> Luther, 1960	A	
<i>sieboldi</i> v. Graff, 1878	M	
Family Tyrrenhiellidae		
<i>Tyrrenhiella</i>		
<i>sigillata</i> Riedl, 1959	M	
Order Haplopharyngida		
Family Haplopharyngidae		
<i>Haplopharynx</i>		
<i>rostratus</i> Meixner, 1938	A	
Order Lecithoepitheliata		
Family Gnosenesimidae		
<i>Gnosenesimila</i>		
<i>borealis</i> (Westblad, 1952) Karling, 1968	A	
<i>brattstroemi</i> Karling, 1968	A	
<i>mediterranea</i> Martens & Schockaert, 1985	M	
<i>reisingeri</i> Karling, 1968	A	
Order Macrostomida		
Family Dolichomacrostomidae		
<i>Acanthomacrostomum</i>		
<i>spiculiferum</i> Papi & Swedmark, 1959	A	
<i>Bathymacrostomum</i>		
<i>spirale</i> Faubel, 1977	A	
<i>Cylindromacrostomum</i>		
<i>mediterraneum</i> (Ax, 1955) Rieger, 1971	M	
<i>notandum</i> (Ax, 1951) Rieger, 1971	A	
<i>Dolichomacrostomum</i>		
<i>uniporum</i> Luther, 1947	A	
<i>Megamorion</i>		
<i>brevicauda</i> Rieger & Sterrer, 1968	A	
<i>Meiocheta</i>		
<i>spiralis</i> Rieger, 1971	A	
<i>Myozonaria</i>		
<i>arcassonensis</i> Rieger, 1971	A M	
<i>bistylifera</i> Rieger, 1968	M	
<i>falcis</i> Sopott-Ehlers, 1976	A	
<i>Paromalostomum</i>		
<i>atratum</i> Rieger, 1971	M	
<i>dubium</i> (de Beauchamp, 1927) Ax, 1952	A M	
<i>fusculum</i> Ax, 1952	A	
<i>massiliensis</i> Rieger, 1971	M	
<i>minutum</i> Rieger, 1971	M	
<i>parvum</i> Rieger, 1971	M	
<i>proceroacuda</i> Pawlak, 1969	A	
<i>tricladoides</i> Riedl, 1932	A	
Family Macrostomidae		
<i>Antromacrostomum</i>		
<i>armatum</i> Faubel, 1976	A	
<i>Bradynectes</i>		
<i>sterreri</i> Rieger, 1971	A	
<i>Macrostomum</i>		
<i>balticum</i> Luther, 1947	A M	
<i>balticum</i> subsp. <i>meridionale</i> Papi, 1953	M	
<i>bicurvistyla</i> Ax & Armonies, 1990	A	
<i>boreale</i> Riedl, 1932	A	
<i>brevituba</i> Armonies & Hellwig, 1987	A	
<i>caudatum</i> Uljanin, 1870	M	
<i>curvituba</i> Luther, 1947	A	
<i>ermini</i> Ax, 1959	M	
<i>gracile</i> Pereyaslawzewska, 1893	A M	
<i>graffi</i> Ferguson, 1939	M	
<i>hamatum</i> Luther, 1947	A	
<i>hystricinum</i> Beklemishev, 1951	A M	
<i>lineare</i> Uljanin, 1870	A M	
<i>longistylerum</i> Ax, 1956	M	
<i>magnacurvitura</i> Ax, 1994	A	
<i>mediterraneum</i> Ax, 1956	M	
<i>minutum</i> (Luther, 1947) Beklemishev, 1951	A	
<i>peteraxi</i> Mack-Fira, 1971	M	
<i>pseudoobtusum</i> Beklemishev, 1927	M	
<i>pusillum</i> Ax, 1951	A M	
<i>rostratum</i> (Papi, 1951) Ferguson, 1954	A	
<i>rubrocinctum</i> Ax, 1951	A M	
<i>spirale</i> Ax, 1951	A M	
<i>tenuicauda</i> Luther, 1947	A	
<i>timavi</i> v. Graff, 1905	A M	
<i>Myozona</i>		
<i>lutheri</i> Papi, 1953	M	
<i>purpurea</i> Faubel, 1974	A	

<i>stylifera</i> Ax, 1956	A	<i>Stylostomum</i>	
<i>Omalostomum</i>		<i>ellipse</i> (Dalyell, 1853) Lang, 1884	A M
<i>claparedei</i> van Beneden, 1870	A	<i>Stygolepta</i>	
<i>groelandicum</i> (Steinböck, 1935) Faubel, Blome & Cannon, 1994	A	<i>hjalmarii</i> Faubel, 1984	A
<i>schultzei</i> (Claparede, 1863) van Beneden, 1870	A	Family Gnesiocerotidae	
<i>Psammomacrostomum</i>		<i>Echinoplana</i>	
<i>equicaudum</i> Ax, 1966	A	<i>celerrima</i> Haswell, 1907	M
<i>turbanelloides</i> Karling, 1974	A	Family Leptoplanidae	
Family Microstomidae		<i>Hoploplana</i>	
<i>Alaurina</i>		<i>insignis</i> (Lang, 1884) Laidlaw, 1902	M
<i>alba</i> v. Attems, 1896	A M	<i>papillosa</i> (Lang, 1884) Laidlaw, 1902	M
<i>composita</i> Metschnikow, 1865	A	<i>villosa</i> (Lang, 1884) Laidlaw, 1902	M
<i>prolifera</i> Busch, 1851	M	<i>Leptoplana</i>	
<i>Microstomum</i>		<i>tremellaris</i> (Müller, 1774) Örsted, 1843	A M
<i>bioculatum</i> Faubel, 1984	A	Family Notocirridae	
<i>crildenisi</i> Faubel, 1984	A	<i>Neoplanocera</i>	
<i>dermophthalmum</i> Riedel, 1932	A	<i>steueri</i> (Steinböck, 1937) Marcus, 1947	M
<i>groelandicum</i> Levinsen, 1979	A	Family Notoplanidae	
<i>hamatum</i> Westblad, 1952	A	<i>Notoplana</i>	
<i>jensenii</i> Riedel, 1932	A	<i>alcinoi</i> (Schmidt, 1861) Bock, 1913	M
<i>lineare</i> (Müller, 1773) Örsted, 1843	A M	<i>cotylifera</i> Meixner, 1907	M
<i>lucidum</i> Fuhrmann, 196	A	<i>kükenthali</i> (Plehn, 1896) Bock, 1913	A
<i>mundum</i> v. Graff, 1905	M	<i>vitrea</i> (Lang, 1884) Bock, 1913	M
<i>ornatum</i> Uljanin, 1870	M	Family Planoceridae	
<i>papillosum</i> v. Graff, 1882	A M	<i>Planocera</i>	
<i>rubromaculatum</i> v. Graff, 1882	A M	<i>folia</i> (Grube, 1840) Örsted, 1844	A M
<i>septentrionale</i> Sabussow, 1899	A	<i>graffii</i> Lang, 1879	M
<i>spiculifer</i> Faubel, 1974	A	Family Plehniidae	
<i>spiriferum</i> Westblad, 1952	A	<i>Discocelides</i>	
Order Nemertodermatida		<i>langi</i> Bergendal, 1893	A
Family Nemertodermatidae		<i>Plehnia</i>	
<i>Flagellophora</i>		<i>arctica</i> (Plehn, 1896) Bock, 1913	A
<i>apelti</i> Faubel & Dörjes, 1978	A	Family Pleioplanidae	
<i>Meara</i>		<i>Pleiopla</i>	
<i>stichoppi</i> (Bock, 1926) Westblad, 1948	A	<i>atomata</i> (Müller, 1776) Faubel, 1983	A M
<i>Nemertoderma</i>		Family Polyposthiidae	
<i>bathycola</i> Steinböck, 1930	A M	<i>Cryptocelides</i>	
<i>rubra</i> Faubel, 1976	A	<i>loveni</i> Bergendal, 1890	A
Order Polycladida		<i>Polyposthia</i>	
Family Anonymidae		<i>similis</i> Bergendal, 1893	A
<i>Anonymus</i>		Family Prosthiostomidae	
<i>virilis</i> Lang, 1884	M	<i>Euprosthiostomum</i>	
Family Callioplanidae		<i>viscosum</i> Palombi, 1936	M
<i>Trigonoporus</i>		<i>Prosthiostomum</i>	
<i>cephalophthalmus</i> Lang, 1884	M	<i>dohrnii</i> Lang, 1884	M
Family Cestoplanidae		<i>siphunculus</i> (Delle Chiaje, 1818) Lang, 1884	M
<i>Acestopla</i>		Family Pseudocerotidae	
<i>raffaelei</i> (Ranzi, 1927) Faubel, 1983	M	<i>Monobiceros</i>	
<i>Cestopla</i>		<i>langi</i> Faubel, 1984	M
<i>rubocincta</i> (Grube, 1840) Lang, 1884	A M	<i>Thysanozoon</i>	
Family Cryptocelididae		<i>brocchii</i> (Risso, 1818) Grube, 1840	
<i>Cryptocelis</i>		<i>Pseudoceros</i>	
<i>alba</i> (Schmidlein, 1880) Lang, 1884	M	<i>maximum</i> Lang, 1884	M
<i>compacta</i> Lang, 1884	M	<i>velutinus</i> (Blanchard, 1847) Lang, 1884	M
<i>glandulata</i> Jakubova, 1909	M	<i>violaceus</i> (Kelaart, 1858) Hyman, 1959	M
Family Enantiidae		<i>Yungia</i>	
<i>Enantia</i>		<i>aurantiaca</i> (Delle Chiaje, 1822) Lang, 1884	M
<i>spinifera</i> v. Graff, 1889	M	Family Stylochidae	
Family Euryleptidae		<i>Imogene</i>	
<i>Acerotisa</i>		<i>mediiterraneus</i> (Gallen, 1976) Jennings & Newman, 1996	M
<i>typhla</i> (Bock, 1913) Strand, 1928	A	<i>Stylochus</i>	
<i>Cycloporus</i>		<i>alexandrinus</i> Steinböck, 1937	M
<i>papillous</i> (Sars in Jensen, 1878) Lang, 1884	A M	<i>flevensis</i> Hofker, 1930	A
<i>Eurylepta</i>		<i>neapolitanus</i> (Delle Chiaje, 1841) Lang, 1884	M
<i>cornuta</i> (Müller, 1776) Ehrenberg, 1831	A M	<i>pillidum</i> (Goette, 1881) Lang, 1884	M
<i>Oligocladus</i>		<i>plessissi</i> Lang, 1884	M
<i>sanguinolentus</i> (Quatrefages, 1845) Lang, 1884	A M	<i>tauricus</i> Jakubova, 1909	M
<i>Prostheceraeus</i>		<i>vesiculosus</i> Jakubova, 1909	M
<i>albocinctus</i> Lang, 1884	M	Family Stylochoplaniidae	
<i>maculosus</i> (Verrill, 1892) Hyman, 1952	A	<i>Armatopla</i>	
<i>vittatus</i> (Montagu, 1815) Lang, 1884	A M		

<i>taurica</i> (Jakubova, 1909) Faubel, 1983	M	<i>paradoxum</i> An Der Lan, 1936	M
<i>Comoplana</i>		<i>productum</i> (Riedl, 1959)	M
<i>agilis</i> (Lang, 1884) Faubel, 1983	A M	<i>pseudomaculatum</i> Gamble, 1893	A
<i>palmula</i> (Quatrefages, 1845) Faubel, 1983	M	<i>reticulatum</i> (Schmidt, 1852) v. Graff, 1882	A M
<i>Emprostopharynx</i>		<i>rovignense</i> An Der Lan, 1936	M
<i>pallida</i> (Quatrefages, 1845) Faubel, 1983	M	<i>siphonophorum</i> (Schmidt, 1852) v. Graff, 1882	A M
<i>Stylochoplana</i>		<i>sorrentium</i> Riedl, 1954	M
<i>maculata</i> (Quatrefages, 1845) Stimpson, 1857	A	<i>spadix</i> (Riedl, 1959)	M
Order Prolecithophora		<i>striatum</i> Westblad, 1956	A M
Family Cylindrostomidae		<i>sulphureum</i> v. Graff, 1882	A M
<i>Allostoma</i>		<i>violaceum</i> Fuhrmann, 1898	A
<i>austriacum</i> (v. Graff, 1882) v. Graff, 1911	A M	<i>vittatum</i> (Frey & Leuckart, 1847) Jensen, 1883	A
<i>capitatum</i> (Uljanin, 1870) v. Graff, 1882	M	<i>whitmani</i> v. Graff, 1911	A M
<i>catinosum</i> (Beklemishev, 1927) Karling, 1993	M	<i>xenophtalmum</i> (Riedl, 1959)	M
<i>singalianum</i> (Claparede, 1861) Reisinger, 1926	A	<i>Plicastoma</i>	
<i>flavibacillum</i> (Jensen, 1878) Reisinger, 1926	A	<i>bimaculatum</i> (v. Graff, 1882) v. Graff, 1908	M
<i>neostiliferum</i> Karling, 1993	A	<i>Plagiostomum</i>	
<i>pallidum</i> van Beneden, 1861	A M	<i>cuticulatum</i> (Brandtner, 1934) Kulinich, 1980	A
<i>uterinum</i> (Westblad, 1952) Karling, 1993	A	<i>Plicastoma</i>	
<i>Cylindrostoma</i>		<i>glaucum</i> Riedl, 1959	M
<i>elongatum</i> Levinsen, 1879	A	<i>micropharynx</i> (An Dcr Lan, 1936) Karling, 1940	M
<i>gracilis</i> Westblad, 1955	A	<i>norvegium</i> Karling, 1940	A
<i>laetum</i> Riedl, 1959	M	<i>spongophilum</i> (Riedl, 1959) Sluys, 1992	M
<i>luridum</i> Riedl, 1959	M	<i>Vorticeros</i>	
<i>lutheri</i> (Westblad, 1955) Karling, 1993	A	<i>auriculatum</i> (Müller, 1788) v. Graff, 1882	A M
<i>monotrochum</i> (v. Graff, 1882) Westblad, 1955	A M	<i>luteus</i> Hallez, 1979	A M
<i>paralutheri</i> Karling, 1993	A	<i>rudis</i> Riedl, 1959	M
<i>tubiferum</i> Westblad, 1954	A	Family Protomonotresidae	
<i>zooxanthellum</i> (v. Graff, 1886) Reisinger, 1926	M	<i>Archimonotresis</i>	
<i>Einarhelmins</i>		<i>limophila</i> Meixner, 1938	A M
<i>stilifera</i> (Westblad, 1955) Karling, 1993	A	<i>perspicua</i> Riedl, 1954	M
<i>Allostoma</i>		Family Pseudostomidae	
<i>flavibacillum</i> (Jensen, 1878) Reisinger, 1926	A	<i>Gonostomula</i>	
<i>Enterostomula</i>		<i>anocelis</i> Westblad, 1955	A
<i>catinosa</i> (Beklemishev, 1927)	M	<i>Pseudostomum</i>	
<i>dura</i> (Fuhrmann, 1898) Reisinger, 1926	A	<i>arenarium</i> Meixner, 1938	A
<i>graffi</i> (de Beauchamp, 1913) Reisinger, 1926	M	<i>dubium</i> v. Graff, 1911	M
<i>Euxinia</i>		<i>elegans</i> (Pereyaslawzewa, 1893) v. Graff, 1913	M
<i>baltica</i> Meixner, 1938	A	<i>gracilis</i> Westblad, 1955	A
<i>corniculata</i> v. Graff, 1911	M	<i>inerme</i> (Hallez, 1879) v. Graff, 1913	A M
<i>Monoophorum</i>		<i>klostermani</i> (v. Graff, 1874)	A M
<i>catinosum</i> Beklemishev, 1927	M	<i>ponticum</i> (Uljanin, 1870) v. Graff, 1913	M
<i>granulosum</i> Beklemishev, 1927	M	<i>quadrioculatum</i> (Leuckart, 1847) Uljanin, 1870	A M
<i>pleiocelis</i> (v. Graff, 1882) v. Graff, 1911	A	<i>Reisingeria</i>	
<i>striatum</i> (v. Graff, 1878) Böhmig, 1890	A M	<i>hexoculata</i> Westblad, 1955	A
<i>tubiferum</i> (Westblad, 1955) Karling, 1993	A	Family Scleraulophoridae	
Family Hypotrichinidae		<i>Scleraulophorus</i>	
<i>Genostoma</i>		<i>cephalatus</i> Karling, 1940	A
<i>marsiliensis</i> (Calandruccio, 1897) Dörler, 1900	M	Family Ulianiniidae	
<i>tergestina</i> (Calandruccio, 1897) Dörler, 1900	M	<i>Gastropharynx</i>	
Family Plagiostomidae		<i>limophilus</i> Riedl, 1956	A M
<i>Acmostomum</i>		<i>Ulianinia</i>	
<i>dioicum</i> (Mecznikov, 1865) Westblad, 1956	A M	<i>contractilis</i> (Meixner, 1938) Karling, 1963	A
<i>rovignense</i> Steinböck, 1933	M	<i>mollissima</i> Levinsen, 1879	A M
<i>Multipenitata</i>		<i>westbladi</i> Karling, 1963	A
<i>lemani</i> (Forel & Du Plessis, 1874) Kulinich, 1974	A M	Family Urastomidae	
<i>Plagiostomula</i>		<i>Urastoma</i>	
<i>vivipara</i> Westblad, 1955	A	<i>cyprinæ</i> (v. Graff, 1882) Westblad, 1955	A M
<i>Plagiostomum</i>		Order Rhabdocoela	
<i>anocelis</i> Brandtner, 1934	A		
<i>vanbenedeni</i> (Schmidt, 1852) v. Graff, 1882	A M	Suborder Dalyelliida	
<i>cavernae</i> (Riedl, 1959)	M	Family Dalyelliidae	
<i>chromogastrum</i> v. Graff, 1908	M	<i>Alexlutheria</i>	
<i>cinctum</i> Meixner, 1938	A M	<i>acrosiphoniae</i> Karling, 1957	A
<i>cuticulatum</i> (Brandtner, 1934) Kulinich, 1980	A	<i>Axia</i>	
<i>fabrei</i> Fuhrmann, 1898	A	<i>luetjohanni</i> (Ax, 1952) Luther, 1955	A
<i>filicauda</i> Westblad, 1956	A M	<i>remanei</i> (Ax, 1952) Luther, 1955	A
<i>flavum</i> Brandtner, 1934	A	<i>Debeaufchampiola</i>	
<i>girardi</i> (Schmidt, 1857) v. Graff, 1882	A M	<i>artica</i> Ehlers & Franke, 1994	A
<i>koreni</i> Jensen, 1878	A	<i>oculifer</i> (de Beauchamp, 1927) Luther, 1957	A M
<i>maculatum</i> v. Graff, 1882	A M	<i>Halammonvortex</i>	
<i>melenadum</i> v. Graff, 1911	M	<i>macropharynx</i> (Meixner, 1938) Karling, 1943	A M
<i>nostrum</i> Westblad, 1956	A	<i>nigrifrons</i> (Karling, 1935) Karling, 1943	A
<i>ochroleucum</i> v. Graff, 1882	A	<i>Jensenia</i>	

<i>angulata</i> (Jensen, 1878) v. Graff, 1882	A	<i>Vejdovskya</i>	
<i>Microdalyellia</i>		<i>adriatica</i> (Dörler, 1900) Micoletzky, 1910	M
<i>fusca</i> (Fuhrmann, 1894) Ruebush & Hayes, 1939	A M	<i>halileimonia</i> Ax, 1960	A
<i>rossi</i> subsp. <i>maxima</i> Ruebush, 1937		<i>helictos</i> Ax, 1956	M
<i>Sergia</i>		<i>ignava</i> Ax, 1951	A M
<i>sergia</i> (Beklemishev, 1918) Nasonov, 1923	A	<i>mesostyla</i> Ax, 1954	A
Family Fecampiidae		<i>mesostyla</i> subsp. <i>hemicycla</i> Ax, 1959	A
<i>Fecampia</i>		<i>pellucida</i> (Schultze, 1851) v. Graff, 1913	A M
<i>abyssicola</i> Christensen, 1981	A	<i>simrisiensis</i> Karling, 1957	A
<i>erythrocephala</i> Giard, 1886	A M	<i>suecica</i> Luther, 1948	A M
<i>xanthocephala</i> Caulery & Mensil, 1903	A M		
<i>Kronborgia</i>			
<i>amphipodcola</i> Christensen & Kanneworff, 1964	A	Family Pterastericolidae	
<i>caridicola</i> Kanneworff & Christensen, 1966	A	<i>Pterastericola</i>	
Family Graffillidae		<i>astropsychinis</i> (Bashiruddin & Karling, 1970)	
<i>Gieysztoria</i>		<i>Jondelius</i> , 1992	A
<i>ornata</i> subsp. <i>maritima</i> Luther, 1955	A	<i>fedotovi</i> Beklemishev, 1916	A
<i>subsalsa</i> Luther, 1955	M	<i>psilastericola</i> (Jespersen & Lützen, 1972) Jondelius, 1992	A
<i>Graffilla</i>			
<i>brauni</i> Schmidt, 1886	M	Family Umagillidae	
<i>buccincola</i> Jameson, 1897	A	<i>Anoplodiera</i>	
<i>curiosa</i> Westblad, 1954	A	<i>voluta</i> Westblad, 1930	A
<i>graffi</i> Mitin, 1970	A	<i>Anoplodiopsis</i>	
<i>muricicola</i> Ihering, 1880	M	<i>gracilis</i> (Wahl, 1906) Westblad, 1953	M
<i>parasitica</i> (Czerniavsky, 1881) v. Graff, 1882	M	<i>Anoplodium</i>	
<i>Paravortex</i>		<i>chirodotae</i> Sabussow, 1900	A
<i>cardii</i> Hallez, 1908	A	<i>graffi</i> Monticelli, 1892	M
<i>gemellipara</i> Linton, 1910	A	<i>parasita</i> Schneider, 1858	M
<i>karlingi</i> Pike & Burt, 1981	A	<i>stichoppi</i> Bock, 1925	A
<i>scrobiculariae</i> (v. Graff, 1882) Wahl, 1906	A M	<i>Collastoma</i>	
<i>Pseudograffilla</i>		<i>eremita</i> Beklemishev, 1916	A
<i>arenicola</i> Meixner, 1938	A M	<i>minutum</i> Wahl, 1906	M
<i>hymanae</i> Mack-Fira, 1974	M	<i>monorchis</i> Dörler, 1900	A
<i>Bresslaulla</i>		<i>Desmote</i>	
<i>relicta</i> Reisinger, 1929	A M	<i>vorax</i> Beklemishev, 1916	A
Family Provorticidae		<i>Seritia</i>	
<i>Baicallelia</i>		<i>elegans</i> (Westblad, 1953) Cannon, 1982	A
<i>brevituba</i> (Luther, 1921) Nasonov, 1930	A	<i>Syndesmis</i>	
<i>subsalina</i> Ax, 1954	A	<i>aethopharynx</i> Westervelt & Kozloff, 1990	M
<i>Balgetia</i>		<i>albida</i> Kozloff & Westervelt, 1990	A
<i>hyalina</i> Luther, 1962	A	<i>echiniacuti</i> Kozloff, 1997	M
<i>semicirculifera</i> Luther, 1962	A	<i>echinorum</i> Francois, 1886	A M
<i>Canetellia</i>		<i>rubida</i> Kozloff & Westervelt, 1990	A
<i>debeauchampi</i> Ax, 1956	A M	<i>Syndisyrinx</i>	
<i>Coronopharynx</i>		<i>atriovillosa</i> (Westblad, 1953) Cannon, 1982	A
<i>pusillus</i> Luther, 1962	A	<i>Umagilla</i>	
<i>Drepanilla</i>		<i>elegans</i> Westblad, 1953	A
<i>limophila</i> Ehlers, 1979	A	<i>forskalensis</i> Wahl, 1909	M
<i>Hangethelia</i>		<i>Wahlia</i>	
<i>calceifera</i> Karling, 1940	A M	<i>macrostylifera</i> Westblad, 1930	A
<i>Haploveydovskya</i>		Suborder Kalyptorhynchia	
<i>subterranea</i> Ax, 1954	A	Family Acrumenidae	
<i>Luriculus</i>		<i>Acrumena</i>	
<i>minos</i> (Sterrer, 1992) Faubel, Rohde & Watson, 1994	M	<i>massiliensis</i> Brunet, 1965	A M
<i>Pogaina</i>		Family Aculeorhynchidae	
<i>adriatica</i> (Dörler, 1900) Marcus, 1954	M	<i>Aculeorhynchus</i>	
<i>annulata</i> Ax, 1970	M	<i>glandulis</i> Schilke, 1969	A
<i>arcassoniensis</i> Ehlers & Sopott-Ehlers, 1989	A	Family Bertiliellidae	
<i>kinnei</i> Ax, 1970	A	<i>Bertiliella</i>	
<i>muranica</i> (v. Graff, 1905) Marcus, 1954	A	<i>bathycola</i> Rieger & Sterrer, 1975	A
<i>natans</i> (Ax, 1951) Marcus, 1954	A	Family Cicerinidae	
<i>suecica</i> (Luther, 1948) Marcus, 1954	A	<i>Cicerina</i>	
<i>Provortex</i>		<i>brevicirrus</i> Meixner, 1928	A
<i>affinis</i> (Jensen, 1878) v. Graff, 1882	A	<i>eucentrota</i> Ax, 1959	M
<i>balticus</i> (Schultze, 1851) v. Graff, 1882	A	<i>remanei</i> Meixner, 1928	A
<i>hispidus</i> (Claparede, 1863) v. Graff, 1882	A	<i>tetradactyla</i> Giard, 1904	A
<i>karlingi</i> Ax, 1951	A	<i>Didiadema</i>	
<i>littoralis</i> (Ørsted, 1844) v. Graff, 1882	A	<i>picardi</i> Brunet, 1965	M
<i>pallidus</i> Luther, 1948	A	<i>Ethmorrhynchus</i>	
<i>psammophilus</i> Ax, 1951	A	<i>anophthalmus</i> Meixner, 1938	A
<i>tubiferus</i> Luther, 1948	A	<i>Nannorhynchides</i>	
<i>Selimia</i>		<i>coeneus</i> Brunet, 1965	M
<i>similis</i> Ax, 1959	M	<i>harparius</i> Brunet, 1965	M
<i>vivida</i> Ax, 1959	M	<i>herdlaensis</i> (Karling, 1956) Karling, 1964	A
		<i>vividus</i> Brunet, 1965	M

<i>Paraciccerina</i>			
<i>deltoides</i> Martens & Schockaert, 1981	A	<i>typhlus</i> L'Hardy, 1964	A M
<i>globulosa</i> Brunet, 1973	M	<i>Psittacorhynchus</i>	
<i>laboeica</i> Meixner in Karling, 1952	A	<i>verweyi</i> Den Hartog, 1968	A
<i>maristoi</i> Karling, 1952	A	<i>Pterognathus</i>	
<i>Pocillorhynchus</i>		<i>longostilo</i> (Schilke, 1970) Evdonin, 1977	A
<i>agilis</i> Brunet, 1973	M	<i>Uncinorhynchus</i>	
<i>minutus</i> Brunet, 1973	M	<i>flavidus</i> Karling, 1947	A M
<i>Ptyalorhynchus</i>		<i>westibladi</i> Karling, 1952	A
<i>coecus</i> Meixner in Ax, 1951	A	Family Karkinorhynchidae	
<i>piger</i> Brunet, 1973	M	<i>Ballopiana</i>	
<i>Toia</i>		<i>magna</i> Karling, 1949	A M
<i>calceformis</i> Brunet, 1973	M	<i>valkanovi</i> Ax, 1959	M
<i>ycia</i> Marcus, 1952	A	<i>Cheliplana</i>	
<i>Torkarlingia</i>		<i>ascica</i> subsp. <i>terminalis</i> Brunet, 1968	M
<i>euxinica</i> Mack-Fira, 1971	M	<i>boadeni</i> Schilke, 1970	A
<i>Xenocicerina</i>		<i>curvocirro</i> Schilke, 1970	A
<i>gracilis</i> Karling, 1956	A	<i>eugeinos</i> Ax, 1959	M
<i>Zonorhynchus</i>		<i>evdonini</i> Karling, 1983	A
<i>pipetiferus</i> Armonies & Hellwig, 1987	A	<i>firmata</i> Brunet, 1968	M
<i>salinus</i> Karling, 1952	A	<i>gemmifera</i> Noldt, 1989	A
<i>seminascatus</i> Karling, 1956	A	<i>glandulosa</i> de Beauchamp, 1927	A
<i>tvaermannensis</i> (Karling, 1931) Karling, 1952	A	<i>hiemalis</i> Brunet, 1968	M
Family Cystiplanidae		<i>hypergyna</i> Boaden, 1965	M
<i>Cystiplana</i>		<i>mamaevi</i> Evdonin, 1977	A
<i>karlingi</i> Schilke, 1970	A	<i>marcusi</i> (Karling, 1956) Karling, 1983	A
<i>paradoxa</i> Karling, 1964	A M	<i>microcirrus</i> Noldt, 1989	A
<i>rubra</i> Dean, 1977	A	<i>orthocirra</i> Ax, 1959	M
<i>Cystiplex</i>		<i>paradoxa</i> Noldt, 1989	A
<i>axi</i> Karling, 1964	A M	<i>piriformis</i> Brunet, 1968	M
<i>Cystirete</i>		<i>pusilla</i> Brunet, 1968	M
<i>graefei</i> Brunet, 1965	M	<i>remanei</i> (Meixner, 1928) Karling, 1983	A
<i>Nigerrhynchus</i>		<i>rubescens</i> Brunet, 1966	M
<i>opisthoporus</i> Schilke, 1970	A	<i>schilkei</i> Noldt, 1989	A
Family Cytocystidae		<i>stylifera</i> Karling, 1949	A
<i>Cytocystis</i>		<i>vaginalis</i> Karling, 1983	A
<i>clitellatus</i> Karling, 1953	A M	<i>varicauda</i> Brunet, 1971	M
Family Diaseorhynchidae		<i>vestibularis</i> de Beauchamp, 1927	A M
<i>Diascorhynchides</i>		<i>Cheliplanilla</i>	
<i>arenaria</i> Schilke, 1970	A	<i>caudata</i> Meixner, 1938	A M
<i>Diascorhynchus</i>		<i>implicata</i> Brunet, 1968	M
<i>borealis</i> Meixner, 1928	A M	<i>karlingi</i> Boaden, 1963	A
<i>bucina</i> Ehlers & Ehlers, 1980	A	<i>rubra</i> Schilke, 1970	A
<i>caligatus</i> Ax, 1959	M	<i>Karkinorhynchides</i>	
<i>glandulosus</i> (de Beauchamp, 1927)	A	<i>purpureus</i> Schilke, 1970	A
<i>lappvikensis</i> Karling, 1963	A	<i>Karkinorhynchus</i>	
<i>rubrus</i> Boaden, 1963	A	<i>bruneti</i> Schilke, 1970	A
<i>serpens</i> Karling, 1949	A M	<i>l'harydi</i> Schilke, 1970	A
Family Gnathorhynchidae		<i>listensis</i> Schilke, 1970	A
<i>Ancistrorhynchus</i>		<i>primitivus</i> Meixner, 1928	A
<i>ischnurus</i> L'Hardy, 1963	A M	<i>purpureus</i> Schilke, 1970	A
<i>Drepanorhynchides</i>		<i>Lehardya</i>	
<i>diodontus</i> L'Hardy, 1966	A M	<i>megalopharynx</i> (L'Hardy, 1966) Karling, 1983	A
<i>hastatus</i> (Meixner, 1929) L'Hardy, 1966	A	<i>tetragnatha</i> (Ax & Schilke, 1971) Karling, 1983	M
<i>Gnathorhynchus</i>		Family Koinoeystididae	
<i>conocaudatus</i> Meixner, 1929	A M	<i>Brunetia</i>	
<i>inermis</i> Schilke, 1970	A	<i>camarguensis</i> (Brunet, 1965) Karling, 1980	M
<i>krogeni</i> Karling, 1947	A	<i>Leguta</i>	
<i>Neognathorhynchus</i>		<i>chelifera</i> (Karling, 1954) Karling, 1980	A M
<i>lobatus</i> (Ax, 1952) Karling, 1956	A	<i>Itaipusa</i>	
<i>suecicus</i> Karling, 1956	A	<i>scotica</i> (Karling, 1954) Karling, 1978	A
<i>Odontorhynchus</i>		<i>similis</i> (Brunet, 1972) Karling, 1978	M
<i>aculeatus</i> Karling, 1956	A	<i>sophiae</i> (v. Graff, 1905) Karling, 1978	M
<i>lonchiferus</i> Karling, 1947	A	<i>Itaipusina</i>	
<i>Orbiculorhynchus</i>		<i>graefei</i> Karling, 1980	A
<i>luebbeni</i> Noldt, 1989	A	<i>Axiutelga</i>	
<i>Paragnathorhynchus</i>		<i>aculeata</i> (Ax, 1959) Karling, 1980	A M
<i>subterraneus</i> Meixner, 1938	A	<i>Gnorimorhynchus</i>	
<i>Prognathorhynchus</i>		<i>dividuus</i> Brunet, 1972	M
<i>campylostylus</i> Karling, 1947	A	<i>Itaipusa</i>	
<i>canaliculatus</i> Karling, 1947	A	<i>acerosa</i> (Brunet, 1972) Karling, 1978	M
<i>dubius</i> Meixner, 1929	A	<i>karlingi</i> Mack-Fira, 1968	M
<i>karlingi</i> Ax, 1953	A	<i>Marirhynchus</i>	
<i>stilofér</i> Schilke, 1970	A	<i>longasaeta</i> Schilke, 1970	A
		<i>Paratenerrhynchus</i>	
		<i>triplex</i> Brunet, 1972	M

<i>Parauelga</i>		<i>helgolandicus</i> (Mecznikoff, 1865) v. Graff, 1905	A
<i>bilioi</i> Karling, 1964	A	<i>pernix</i> Ax, 1959	M
<i>Pontalaria</i>		<i>Polycystis</i>	
<i>beklemishevi</i> Mack-Fira, 1968	M	<i>contorta</i> Schockaert & Karling, 1975	A M
<i>Tenerrhynchus</i>		<i>crocea</i> (Fabricius, 1826) v. Graff, 1905	A M
<i>magnus</i> Brunet, 1972	M	<i>dolichocephala</i> (Pereyaslawzewska, 1892) v. Graff, 1905	M
<i>Utelga</i>		<i>georgii</i> v. Graff, 1905	M
<i>heinckeii</i> (v. Attems, 1897) Karling, 1954	A M	<i>groenlandica</i> (Levinsen, 1879) v. Graff, 1913	A M
<i>pseudoheinckeii</i> Karling, 1980	A M	<i>intubata</i> subsp. <i>erythrea</i> Beklemishev, 1927	M
<i>spinosa</i> (Beklemishev, 1927) Karling, 1954	M	<i>intubata</i> v. Graff, 1905	M
<i>vesicatula</i> Brunet, 1972	M	<i>mamertina</i> (v. Graff, 1874) v. Graff, 1905	A M
Family Nematorhynchidae		<i>naegellii</i> Kölleiker, 1845	A M
<i>Nematorhynchus</i>		<i>riedli</i> Karling, 1956	M
<i>parvoacumine</i> Schilke, 1969	A	<i>Psammopolycystis</i>	
Family Placorhynchidae		<i>bidentata</i> Meixner, 1938	A
<i>Clyporhynchus</i>		<i>bondensis</i> Karling, 1956	A
<i>monolentis</i> Karling, 1947	A	<i>bredungensis</i> Karling, 1956	A
<i>Placorhynchus</i>		<i>falcata</i> Karling, 1956	A
<i>dimorphis</i> Karling, 1947	A	<i>forcipiens</i> Brunet, 1979	M
<i>echinulatus</i> Karling, 1947	A	<i>riegeri</i> Brunet, 1979	M
<i>octaculeatus</i> Karling, 1931	A M	<i>trilobata</i> Brunet, 1979	M
<i>tetraculeatus</i> Armonies & Hellwig, 1987	A	<i>Rogneda</i>	
Family Polycystididae		<i>acuta</i> Brunet, 1979	M
<i>Papia</i>		<i>anglica</i> Karling, 1953	A
<i>bifida</i> Karling, 1956	M	<i>capulata</i> Karling, 1953	M
<i>Acrorhynchides</i>		<i>cincta</i> Brunet, 1969	M
<i>caledonicus</i> (Claparede, 1861) Karling, 1963	A	<i>exilis</i> Brunet, 1979	M
<i>robustus</i> (Karling, 1931) Karling, 1963	A	<i>falcata</i> Brunet, 1965	M
<i>stylifera</i> Schockaert & Karling, 1975	A	<i>hibernica</i> (Southern, 1936) Karling, 1953	A
<i>Acrorhynchus</i>		<i>minuta</i> Uljanin, 1870	M
<i>bivittatus</i> (Uljanin, 1870) v. Graff, 1882	A M	<i>palula</i> Brunet, 1969	M
<i>dolichocephalus</i> (Pereyaslawzewska, 1892) v. Graff, 1905	M	<i>polyrhabdoa</i> Ax, 1959	M
<i>reprobatus</i> v. Graff, 1905	M	<i>reticulata</i> Brunet, 1969	M
<i>spiralis</i> (Pereyaslawzewska, 1893) v. Graff, 1913	M	<i>steueri</i> (Steinböck, 1933) Karling, 1953	M
<i>Albertorhynchus</i>		<i>tripalmata</i> (Beklemishev, 1927) Karling, 1953	M
<i>amai</i> Schockaert, 1976	M	<i>westbladi</i> subsp. <i>gallica</i> Ax, 1956	M
<i>Annularhynchus</i>		<i>westbladi</i> subsp. <i>westbladi</i> Karling, 1953	M
<i>adriaticus</i> Karling, 1956	M	<i>Scanorhynchus</i>	
<i>Austrorhynchus</i>		<i>forcipatus</i> Karling, 1955	A
<i>bruneti</i> Karling, 1977	M	<i>limophilus</i> Karling, 1955	A
<i>karlingi</i> Brunet, 1965	M	<i>Syltorhynchus</i>	
<i>scoparius</i> Brunet, 1965	M	<i>schockaerti</i> Noldt, 1989	A
<i>Danorhynchus</i>		<i>Typhlopolydystis</i>	
<i>duplostylis</i> Karling, 1955	A	<i>coeca</i> Karling, 1956	A
<i>goesoeensis</i> Karling, 1955	A	<i>coomansi</i> Schockaert & Karling, 1975	A M
<i>Gyratricella</i>		<i>mediterranea</i> Brunet, 1965	A M
<i>attensi</i> (v. Graff, 1913) Karling, 1955	A M	Family Psammorhynchidae	
<i>Gyratrix</i>		<i>Psammorhynchus</i>	
<i>hermaphroditus</i> Ehrenberg, 1831	A M	<i>tubulipenis</i> Meixner, 1938	A
<i>proavus</i> Meixner, 1929	A	Family Schizorhynchidae	
<i>Lagenorhynchus</i>		<i>Amphirhynchus</i>	
<i>peresi</i> Brunet, 1965	M	<i>caudatus</i> Schilke, 1970	A
<i>Lekanorhynchus</i>		<i>Carcharodorhynchus</i>	
<i>remanei</i> Meixner, 1938	A	<i>ambronensis</i> Schilke, 1970	A
<i>Limipolydystis</i>		<i>flavidus</i> Brunet, 1967	M
<i>curvituba</i> Schilke, 1970	A	<i>infundibulatus</i> Brunet, 1970	M
<i>Polycystis</i>		<i>isolatus</i> Schilke, 1970	A
<i>immundus</i> (Schmidt, 1857) v. Graff, 1905	M	<i>listensis</i> Schilke, 1970	A
<i>leucophraeus</i> (Fabricius, 1823)	A	<i>multidentatus</i> Brunet, 1979	M
<i>papillatus</i> (Mereschkowsky, 1878)	A	<i>polyorchis</i> L'Hardy, 1963	A
<i>Mesorhynchus</i>		<i>subterraneus</i> Ax, 1951	A M
<i>terminostylis</i> Karling, 1956	A	<i>tenuis</i> Brunet, 1979	M
<i>Neopolydystis</i>		<i>Limirhynchus</i>	
<i>tridentata</i> Karling, 1955	A M	<i>curvituba</i> Schilke, 1970	A
<i>Opisthocystis</i>		<i>danicus</i> Schilke, 1970	A
<i>goettei</i> (Bresslau, 1906) Sekera, 1912	A	<i>Neoschizorhynchus</i>	
<i>Parachlororhynchus</i>		<i>brevipharyngeus</i> Schilke, 1970	A
<i>axi</i> Karling, 1956	A	<i>longipharyngeus</i> Schilke, 1970	A
<i>bergensis</i> Karling, 1956	A	<i>parvorostro</i> Ax & Heller, 1970	A
<i>Phonorhynchella</i>		<i>Paraschizorhynchoides</i>	
<i>biarciata</i> Karling, 1956	A	<i>glandulis</i> Schilke, 1970	A
<i>Phonorhynchus</i>		<i>Proschizorhynchella</i>	
<i>bitubatus</i> Meixner, 1938	A	<i>bivaginata</i> (Schilke, 1970) Noldt, 1985	A
		<i>echinulata</i> (L'Hardy, 1965) Noldt, 1985	A

<i>saeroensis</i> (Meixner, 1938) Noldt, 1985	A	<i>intermedia</i> Ax, 1952	A
<i>helgolandica</i> (L'Hardy, 1965) Noldt, 1985	A M	Family Promesostomidae	
<i>oculata</i> (Meixner, 1928) Noldt, 1985	A	<i>Adenorhynchus</i>	
<i>robusta</i> Noldt, 1989	A	<i>balticus</i> Meixner, 1938	A
<i>spiracirro</i> (Schilke, 1970) Noldt, 1985	A	<i>Brinkmanniella</i>	
<i>Proschizorhynchoides</i>		<i>macrostomoides</i> Luther, 1948	A M
<i>macrostylus</i> (L'Hardy, 1963) Noldt, 1989	A	<i>obtusa</i> Luther, 1943	A
<i>Proschizorhynchus</i>		<i>palmata</i> Karling, 1986	A M
<i>anophthalmus</i> L'Hardy, 1965	A	<i>procerostyla</i> Ehlers, 1974	A
<i>arenarius</i> de Beauchamp, 1927	A M	<i>Coronhelmis</i>	
<i>gullmarenensis</i> Karling, 1950	A M	<i>inornatus</i> Ehlers, 1974	A
<i>gullmarenensis</i> subsp. <i>tricingulatus</i> Ax, 1959	M	<i>lutheri</i> Ax, 1951	A
<i>lunatus</i> Brunet, 1970	M	<i>multispinosus</i> Luther, 1948	A
<i>pectinatus</i> L'Hardy, 1965	A M	<i>tripartitus</i> Ehlers, 1974	A
<i>reniformis</i> Brunet, 1970	M	<i>Einarella</i>	
<i>triductibus</i> Schilke, 1970	A	<i>argillophila</i> Luther, 1948	A M
<i>Pseudoschizorhynchoides</i>		<i>Listea</i>	
<i>karlingi</i> (Martens & Schockaert, 1981) Noldt, 1989	A	<i>simplex</i> Ax & Heller, 1970	A
<i>ruber</i> Schilke, 1970	A	<i>Litucivis</i>	
<i>Schizochilus</i>		<i>serpens</i> Ax & Heller, 1970	A
<i>caecus</i> L'Hardy, 1963	A	<i>Moevenbergia</i>	
<i>choriurus</i> Boaden, 1963	A	<i>una</i> Armonies & Hellwig, 1987	A
<i>marcusi</i> Boaden, 1963	A	<i>Promesostoma</i>	
<i>parvulus</i> Brunet, 1970	M	<i>balticum</i> Luther, 1918	A
<i>spiniferus</i> Brunet, 1970	M	<i>bilineatum</i> Pereyaslawzewska, 1892	A M
<i>tubulatus</i> Brunet, 1970	M	<i>bipartitum</i> Ax, 1956	A
<i>Schizorhynchoides</i>		<i>caligulatum</i> Ax, 1952	A
<i>aculeatus</i> L'Hardy, 1963	A	<i>cochlearis</i> Karling, 1935	A M
<i>brevistili</i> Noldt, 1989	A	<i>dennisallenii</i> Ax, 1995	A
<i>canaliculatus</i> L'Hardy, 1963	A	<i>digitosa</i> Ax, 1995	A
<i>coronostylus</i> Boaden, 1963	A	<i>ensifer</i> (Uljanin, 1870) Pereyaslawzewska, 1893	M
<i>diplochilus</i> Meixner, 1928	A	<i>ericae</i> Faubel, 1980	A
<i>globulosus</i> Brunet, 1970	M	<i>gallicum</i> Ax, 1956	M
<i>karlingi</i> Martens & Schockaert, 1981	A	<i>gracilis</i> Ax, 1951	A
<i>macrostylus</i> L'Hardy, 1963	A	<i>karlingi</i> Ehlers, 1974	A
<i>meixneri</i> Boaden, 1963	A	<i>kergroixense</i> Ehlers & Soppott-Ehlers, 1989	A
<i>rarus</i> Ax, 1951	A M	<i>maculosum</i> Ax, 1956	A M
<i>spirostylus</i> Boaden, 1963	A	<i>marmoratum</i> (Schultze, 1851) v. Graff, 1882	A M
<i>symmetricus</i> Martens & Schockaert, 1981	A	<i>meixneri</i> Ax, 1951	A
<i>Schizorhynchus</i>		<i>meixneri</i> subsp. <i>roscoffense</i> Ehlers, 1980	A
<i>arenarius</i> de Beauchamp, 1927	A	<i>minutum</i> Ax, 1956	A
<i>coecus</i> Hallez, 1894	A	<i>murmanicum</i> v. Graff, 1905	A
<i>tataricus</i> v. Graff, 1905	M	<i>negletum</i> Karling, 1967	A
<i>Serratorhynchus</i>		<i>norvegicum</i> Karling, 1967	A
<i>stellatus</i> Noldt, 1988	A	<i>nynaeiensis</i> Karling, 1957	A
<i>Thylacorhynchus</i>		<i>ovoideum</i> (Schmidt, 1852) v. Graff, 1882	A M
<i>ambronensis</i> Schilke, 1970	A	<i>paracochlearis</i> Ax, 1952	A
<i>arcassonensis</i> de Beauchamp, 1927	A	<i>rostratum</i> Ax, 1951	A
<i>caudatus</i> Meixner, 1928	A	<i>serpentistylum</i> Ax, 1952	A
<i>conglobatus</i> Meixner, 1928	A M	<i>spiriferum</i> Karling, 1967	A
<i>filostylis</i> Karling, 1956	A	<i>Protandrella</i>	
<i>longicirrus</i> Noldt, 1989	A	<i>cirriera</i> Ehlers, Müller & Franke, 1994	A
<i>pyriferus</i> Karling, 1950	A	<i>Scolipharyngia</i>	
<i>vicarus</i> Boaden, 1963	A	<i>arenicola</i> Ehlers, Reise & Armonies, 1995	A
<i>Typhlorhynchus</i>		<i>listensis</i> Ehlers, Müller & Franke, 1994	A
<i>nanus</i> Laidlaw, 1902	M	<i>Subulagera</i>	
<i>sylensis</i> Noldt, 1985	A	<i>mucronata</i> Ehlers, 1974	A
Suborder Typhloplanida		<i>rubra</i> Martens & Schockaert, 1981	A
Family Byrsophlebidae		<i>Tvaerminnea</i>	
<i>Byrsophlebs</i>		<i>direceptacula</i> Ehlers, Müller & Franke, 1994	A
<i>delamarei</i> (Ax, 1956) Karling, 1985	M	<i>karlingi</i> Luther, 1943	A M
<i>dubia</i> (Ax, 1956) Karling, 1985	A M	<i>Westbladiella</i>	
<i>graffi</i> Jensen, 1878	A	<i>angulifera</i> Riedl, 1954	M
<i>simplex</i> (Ax, 1959) Karling, 1985	M	<i>falcifera</i> Riedl, 1954	M
<i>uncinata</i> (Ax, 1959) Karling, 1985	M	<i>obliquepharynx</i> Luther, 1943	A
<i>Maehrenthalia</i>		<i>tubifera</i> Riedl, 1956	M
<i>agilis</i> (Levinsen, 1879) v. Graff, 1905	A	Family Solenopharyngidae	
<i>americana</i> Ax & Armonies, 1990	A	<i>Adenopharynx</i>	
<i>intermedia</i> (v. Graff, 1882) v. Graff, 1905	A	<i>mitrabursalis</i> Ehlers, 1972	A
<i>Parabyrsophlebs</i>		<i>Anthopharynx</i>	
<i>coeca</i> (Luther, 1948) Karling, 1985	A	<i>sacculipenis</i> Ehlers, 1972	A
Family Ciliopharyngiellidae		<i>vaginatus</i> Karling, 1940	A
<i>Ciliopharyngiella</i>		<i>Aulopharynx</i>	
<i>constricta</i> Martens & Schockaert, 1981	A	<i>aestuarius</i> Ehlers, 1972	A
		<i>Culleopharynx</i>	

<i>armatus</i> (Riedl, 1956) Ehlers, 1972	M	<i>lutheri</i> Den Hartog, 1966	A
<i>Doliopharynx</i>		<i>minimus</i> Den Hartog, 1966	A
<i>geminocirro</i> Ehlers, 1972	A	<i>monotubulus</i> Den Hartog, 1966	A
<i>Espiegrendia</i>		<i>multidentatus</i> Ax., 1971	A
<i>norvegica</i> Westblad, 1954	A	<i>pratensis</i> Ax, 1960	A
<i>Lenopharynx</i>		<i>puccinellicola</i> Ax, 1960	A
<i>languidus</i> Beklemishev, 1929	M	<i>quadrispinosus</i> Den Hartog, 1966	A
<i>Proceropharynx</i>		<i>quinquespinus</i> Ax, 1971	A
<i>anophthalmus</i> (Meixner, 1929) Ehlers, 1972	A	<i>segmentatus</i> Den Hartog, 1966	A
<i>litoralis</i> Ehlers, 1972	A	<i>simplex</i> Luther, 1948	A M
<i>Solenopharynx</i>		<i>tenuispinosus</i> Ehlers, 1974	A
<i>flavidus</i> v. Graff, 1882	M	<i>trigonous</i> Ax, 1960	A
<i>Sophar ynx</i>		<i>tuberculatus</i> v. Graff, 1882	A
<i>oculatus</i> (Pereyaslawzewska, 1892) Beklemishev, 1929	M	<i>uncinatus</i> (v. Graff, 1882) Den Hartog, 1965	A
<i>typhlops</i> Beklemishev, 1929	M	<i>unidentatus</i> Den Hartog, 1965	A
<i>Tensopharynx</i>			
<i>inermis</i> Ehlers, 1972	A		
<i>Trisaccopharynx</i>			
<i>roscoffensis</i> Ehlers, 1980	A		
<i>spiculatus</i> Ehlers, 1972	A		
<i>westbladi</i> Karling, 1940	A		
Family Trigonostomidae			
<i>Astrotothynchus</i>			
<i>bifidus</i> (Mc'Intosh, 1874) v. Graff, 1905	A		
<i>Beklemishevella</i>			
<i>angustior</i> Luther, 1943	A		
<i>contorta</i> (Beklemishev, 1927) Luther, 1943	A		
<i>Brederveldia</i>			
<i>bidentata</i> Van Der Velde & Van De Winkel, 1975	A		
<i>Ceratopera</i>			
<i>axi</i> (Riedl, 1954) Den Hartog, 1964	M		
<i>bifida</i> Ehlers & Ax, 1974	M		
<i>gracilis</i> (v. Graff, 1882) Den Hartog, 1964	A M		
<i>paradoxa</i> (Pereyaslawzewska, 1892) Den Hartog, 1964	M		
<i>reisingeri</i> (Riedl, 1959) Den Hartog, 1964	M		
<i>sellai</i> (Steinböck, 1933) Den Hartog, 1964	M		
<i>steinböcki</i> (Riedl, 1959) Den Hartog, 1964	M		
<i>Lonchoplarella</i>			
<i>axi</i> Ehlers, 1974	A		
<i>Lutheriella</i>			
<i>diplostyla</i> Den Hartog, 1966	A		
<i>Marinella</i>			
<i>lingulifera</i> Riedl, 1954	M		
<i>Mariplanella</i>			
<i>frisia</i> Ax & Heller, 1970	A		
<i>Messoplana</i>			
<i>aremorica</i> Ehlers & Sopott-Ehlers, 1989	A		
<i>canariensis</i> Ehlers & Ehlers, 1980	A		
<i>elegans</i> (Luther, 1948) Den Hartog, 1966	A M		
<i>falcata</i> subsp. <i>falcata</i> (Ax, 1953) Den Hartog, 1966	A M		
<i>floralis</i> Ehlers, 1974	A		
<i>geminata</i> Den Hartog, 1966	A		
<i>helgolandica</i> Ax, 1971	A		
<i>roscoffensis</i> Ehlers, 1980	A		
<i>rugata</i> Ehlers, 1974	A		
<i>Paramesostoma</i>			
<i>helgolandicum</i> Ax, 1959	A		
<i>neapolitanum</i> (v. Graff, 1882) v. Attems, 1897	A M		
<i>Petaliella</i>			
<i>spiracula</i> Ehlers, 1974	A		
<i>Proxenetes</i>			
<i>ampullatus</i> Ax, 1971	A		
<i>bilioi</i> Den Hartog, 1966	A		
<i>britannicus</i> Den Hartog, 1966	A		
<i>chlorosticus</i> (Schmidt, 1857) v. Graff, 1882	M		
<i>cimbricus</i> Ax, 1971	A		
<i>cisorius</i> Den Hartog, 1966	A		
<i>cochlear</i> v. Graff, 1882	A		
<i>deltaoides</i> Den Hartog, 1965	A		
<i>fasciger</i> Ehlers, 1974	A		
<i>flabellifer</i> Jensen, 1878	A M		
<i>flexus</i> Ax, 1971	A		
<i>inflatus</i> Den Hartog, 1966	A		
<i>intermedius</i> Den Hartog, 1966,	A		
<i>karlingi</i> Luther, 1943	A		
<i>Ptychopera</i>			
<i>avicularis</i> Karling, 1974	A		
<i>ehlersi</i> Ax, 1971	A		
<i>hartogi</i> Ax, 1971	A		
<i>plebeia</i> (Beklemishev, 1927) Den Hartog, 1964	M		
<i>purasjokii</i> Ax, 1971	A		
<i>spinifera</i> Den Hartog, 1966	A		
<i>subterranea</i> Ax, 1971	M		
<i>wesibladi</i> (Luther, 1943) Den Hartog, 1964	A M		
<i>zostericola</i> Ax, 1971	A		
<i>Trigonostomum</i>			
<i>armatum</i> (Jensen, 1878) Southem, 1912	A		
<i>breitfussi</i> (v. Graff, 1905)	A		
<i>brunchorsti</i> v. Graff, 1905	A		
<i>coronatum</i> (v. Graff, 1882) v. Graff, 1913	A		
<i>intermedium</i> (v. Attems, 1896) v. Graff, 1913	A		
<i>mirabile</i> (Pereyaslawzewska, 1892) v. Graff, 1913	M		
<i>penicillatum</i> (Schmidt, 1857) Micoletzky, 1910	A M		
<i>piriforme</i> (Pereyaslawzewska, 1893) v. Graff, 1905	M		
<i>quadrifolium</i> Riedl, 1954	M		
<i>seigermani</i> Schmidt, 1852	A M		
<i>venenosum</i> (Ulanin, 1870)	A M		
Family Typhloplanidae			
<i>Castrada</i>			
<i>hofmanni</i> Braun, 1885	A		
<i>intermedia</i> (Volz, 1898) Luther, 1904	A		
<i>lanceola</i> (Braun, 1885) Luther, 1904	A		
<i>subsalsa</i> Luther, 1946	A		
<i>viridis</i> Volz, 1898	A		
<i>Dochmiotrema</i>			
<i>limicola</i> Hofsten, 1907	A		
<i>Gullmariella</i>			
<i>vivipara</i> Luther, 1948	A		
<i>Haloplanella</i>			
<i>curvistyla</i> Luther, 1946	A		
<i>hamulata</i> Ehlers, 1974	A		
<i>longatuba</i> Ax & Heller, 1970	A		
<i>minuta</i> Luther, 1946	A		
<i>multifida</i> Riedl, 1954	M		
<i>obtusituba</i> Luther, 1946	A		
<i>pusilla</i> Riedl, 1959	M		
<i>Hartogia</i>			
<i>pontica</i> Mack-Fira, 1969	M		
<i>Hoplopera</i>			
<i>littoralis</i> Karling, 1957	A		
<i>pusilla</i> Ehlers, 1974	A		
<i>Lioniella</i>			
<i>petiti</i> Riedl, 1954	M		
<i>Notomonoporum</i>			
<i>coecum</i> Luther, 1948	A M		
<i>Opistomum</i>			
<i>immigrans</i> Ax, 1956	A M		
<i>Phaeocora</i>			
<i>rufodorsata</i> (Sekera, 1903) v. Graff, 1909	A		
<i>salinarum</i> subsp. <i>subsalsa</i> Beklemishev, 1927	M		
<i>subsalina</i> (Luther, 1921) Luther, 1963	A		
<i>Pratoplaena</i>			
<i>galeata</i> Ehlers, 1974	A		
<i>salsa</i> Ax, 1960	A		
<i>Stygotplanellina</i>			

<i>halophila</i> Ax, 1954	A	<i>longicirrus</i> Martens, 1984	M
<i>Thalassoplanella</i>	A	<i>paucispina</i> Martens, 1984	M
<i>collaris</i> Luther, 1946		<i>septentrionalis</i> Martens, 1983	A
<i>Thalassoplanina</i>	M	<i>Duploperaclistus</i>	
<i>geniculata</i> Ax, 1959		<i>circocirrus</i> Martens, 1983	A
Order Seriata		<i>Ectocotyla</i>	
Suborder Proseriata		<i>hirudo</i> (Levinsen, 1879) Hyman, 1944	A
Family Coelogynoporidae		<i>Inaloa</i>	
<i>Carencoilia</i>		<i>cirrifera</i> (Meixner, 1943) Martens & Curini-Galletti, 1994	M
<i>bidentata</i> Sopott, 1972	A	<i>Minona</i>	
<i>biforamen</i> Sopott, 1972	A	<i>baltica</i> Karling & Kinnander, 1953	A
<i>Cirrifera</i>		<i>degadli</i> Martens, 1983	A
<i>aculeata</i> (Ax, 1951) Sopott, 1972	A	<i>trigonopora</i> Ax, 1956	M
<i>boletiformae</i> Sopott, 1972	A	<i>Monocelis</i>	
<i>cirrifera</i> Sopott, 1972	A	<i>fuhrmanni</i> Midelburg, 1908	M
<i>dumosa</i> Sopott, 1972	A	<i>fusca</i> Örsted, 1843	A M
<i>Coelogynopora</i>		<i>gracilis</i> Meixner, 1938	A
<i>axi</i> Sopott, 1972	A	<i>lineata</i> (Müller, 1774) Ehrenberg, 1831	A M
<i>biarmata</i> Steinböck, 1924	A M	<i>longistyla</i> Martens & Curini-Galletti, 1987	M
<i>bresslaui</i> Steinböck, 1924	A	<i>nitida</i> Riedl, 1959	M
<i>cassida</i> Sopott-Ehlers, 1985	A	<i>viridorostris</i> (Sabussow, 1900) v. Graff, 1893	A
<i>distortofolio</i> Sopott, 1972	A	<i>Monocelopsis</i>	
<i>faenofurca</i> Sopott-Ehlers, 1992	A	<i>otoplanoides</i> Ax, 1951	A
<i>forcipis</i> Sopott-Ehlers, 1976	A	<i>septentrionalis</i> (Sopott, 1972) Martens & Curini-Galletti, 1994	A
<i>gallica</i> Sopott-Ehlers, 1976	A M	<i>Paramonotus</i>	
<i>gynocotyla</i> Steinböck, 1924	A	<i>hamatus</i> (Jensen, 1878) Meixner, 1938	A
<i>hamulis</i> Sopott-Ehlers, 1980	A	<i>Preminona</i>	
<i>hangensis</i> Karling, 1953	A	<i>insularis</i> (Meixner, 1928) Karling, 1966	A
<i>schultzi</i> Meixner, 1938	A	<i>Promonotus</i>	
<i>sequana</i> Sopott-Ehlers, 1992	A	<i>arcassonensis</i> Ax, 1959	A
<i>solifer</i> Sopott, 1972	A	<i>marci</i> Ax, 1954	A
<i>steinböckii</i> Sopott-Ehlers, 1980	A	<i>ponticus</i> Ax, 1959	M
<i>tenuis</i> Meixner, 1938	A	<i>schultzei</i> Meixner, 1943	A M
<i>Coleogynopora</i>		<i>Pseudomonocelis</i>	
<i>visurgis</i> Sopott-Ehlers, 1989	A	<i>agilis</i> (Schultze, 1851) Ax, 1959	A M
<i>Invenusta</i>		<i>cetinae</i> Meixner, 1943	A M
<i>aestus</i> Sopott-Ehlers, 1976	A	<i>ophiocephala</i> (Schmidt, 1861) Meixner, 1943	M
<i>Vannuccia</i>		Family Monotoplanidae	
<i>campana</i> Ehlers & Ehlers, 1980	A M	<i>Monotoplana</i>	
<i>umbilica</i> Sopott, 1972	A	<i>diorchis</i> Meixner, 1938	A M
Family Monocelididae		<i>rufifrons</i> (Westblad, 1952) Ax, 1958	A
<i>Archilina</i>		Family Nematopланidae	
<i>biselinifera</i> Martens & Curini-Galletti, 1994	M	<i>Nematoplana</i>	
<i>brachycirrus</i> Martens & Curini-Galletti, 1994	M	<i>asita</i> Marcus, 1950	A
<i>caliban</i> Martens & Curini-Galletti, 1994	M	<i>caesarea</i> Curini-Galletti & Martens, 1992	M
<i>deceptoria</i> Martens & Curini-Galletti, 1994	M	<i>coelogynoporoidea</i> Meixner, 1938	A M
<i>endostyla</i> Ax, 1959	M	<i>corsicana</i> Curini-Galletti & Martens, 1992	M
<i>etrusca</i> Martens & Curini-Galletti, 1994	M	<i>riegeri</i> Curini-Galletti & Martens, 1992	M
<i>selenifera</i> Martens & Curini-Galletti, 1994	M	Family Otomesostomidae	
<i>Archiloa</i>		<i>Otomesostoma</i>	
<i>petiti</i> Ax, 1956	A M	<i>auditivum</i> (Du Plessis, 1874) v. Graff, 1882	A
<i>rivularis</i> de Beauchamp, 1910	A	Family Otoplanidae	
<i>westbladi</i> Ax, 1954	A	<i>Archotoplana</i>	
<i>Archilopsis</i>		<i>holotricha</i> Ax, 1956	A M
<i>arenaria</i> Martens, Curini-Galletti & Puccinelli, 1989	A	<i>Bothriomolus</i>	
<i>marifuga</i> Martens, Curini-Galletti & Puccinelli, 1989	A	<i>balticus</i> Meixner, 1938	A
<i>spinosa</i> (Jensen, 1878) Martens, Curini-Galletti & Puccinelli, 1989	A M	<i>constrictus</i> Hallez, 1909	A
<i>unipunctata</i> (Fabricius, 1826) de Beauchamp, 1910	A M	<i>Bulbotoplana</i>	
<i>Archimonocelis</i>		<i>acephala</i> Ax, 1956	A
<i>bathycola</i> (Westblad, 1952) Karling, 1966	A	<i>Dicoelandropora</i>	
<i>carmelitana</i> Martens & Curini-Galletti, 1993	M	<i>atriopapillata</i> Ax, 1956	A
<i>crucifera</i> Martens & Curini-Galletti, 1993	M	<i>Itaspilla</i>	
<i>koinocystis</i> Karling, 1966	A	<i>helgolandica</i> subsp. <i>helgolandica</i> (Meixner, 1938)	
<i>mediterranea</i> Meixner, 1938	M	Sopott, 1972	A
<i>meixneri</i> Martens & Curini-Galletti, 1993	M	<i>Kataplana</i>	
<i>oostendensis</i> Martens & Schockaert, 1981	A	<i>arcuata</i> Sopott-Ehlers, 1976	A
<i>Boreocelis</i>		<i>germanica</i> (Meixner in Ax, 1951) Ax, 1956	A
<i>silicula</i> Westblad, 1952	A	<i>mesopharynx</i> Ax, 1956	A
<i>urodasysoides</i> Ax, 1963		<i>Monostichoplana</i>	
<i>Duplominona</i>		<i>filum</i> (Meixner, 1938) Ax, 1956	A
<i>canariensis</i> subsp. <i>canariensis</i> Ehlers & Ehlers, 1980	A	<i>tenuissima</i> Ax, 1956	A
<i>corsicana</i> Martens, 1984	M	<i>Napoliplana</i>	
<i>istambulensis</i> (Ax, 1959) Karling, 1966	M		

<i>cinctata</i> Ax, Weidemann & Ehlers, 1978	M	Family Bdellouridae
<i>Notocaryoplana</i>	A M	<i>Pentacoelum</i>
<i>arctica</i> Steinböck, 1935		<i>fucoidem</i> Westblad, 1935
<i>Notocaryoturbella</i>	M	<i>punctatum</i> (Brandtner, 1935) Sluys & Bush, 1988
<i>bigermaria</i> Lanfranchi, 1969		A
<i>Orthoplana</i>	A	Family Cercyridae
<i>borealis</i> (Steinböck, 1931) Steinböck, 1933	A	<i>Sabussowia</i>
<i>filum</i> (Meixner, 1938) Ax, 1956	A	<i>dioica</i> (Claparede, 1863) Böhmig, 1906
<i>soliacea</i> (Meixner, 1938) Ax, 1956	A	<i>Cerbussovia</i>
<i>mediterranea</i> Ax, 1956	M	<i>cerruti</i> Wilhelmi, 1909
<i>Otoplana</i>	M	<i>Cercyra</i>
<i>bosporana</i> Ax, 1959	M	<i>hastata</i> Schmidt, 1862
<i>cincinnata</i> (Calendruccio, 1897) Wilhelmi, 1908	M	<i>teissieri</i> Steinmann, 1930
<i>intermedia</i> Du Plessis, 1889	A M	A
<i>maculata</i> Steinböck, 1931	A	Family Procerodidae
<i>setosa</i> (Du Plessis, 1889) Wilhelmi, 1908	M	<i>Procerodes</i>
<i>truncaspina</i> Lanfranchi, 1969	M	<i>dahli</i> Marcus & Marcus, 1959
<i>Otoplanella</i>	A	<i>dohrnii</i> Wilhelmi, 1909
<i>baltica</i> (Meixner, 1938) Ax, 1956	A	<i>litoralis</i> (Ström, 1768) Hallez, 1893
<i>schultzi</i> (Ax, 1951) Ax, 1956	A	<i>lobata</i> (Schmidt, 1861) Hallez, 1893
<i>Otoplaniida</i>	A	<i>plebeia</i> (Schmidt, 1861) Hallez, 1893
<i>endocystis</i> Meixner, 1938		
<i>Paradoxoplana</i>	A	Family Uteriporidae
<i>solida</i> Ax, 1956	M	<i>Foviella</i>
<i>Paraplana</i>	A	<i>affinis</i> (Örsted, 1843) Bock, 1925
<i>biscayensis</i> Ax, 1956		<i>Uteriporus</i>
<i>Parotoplana</i>	A	<i>vulgaris</i> Bergendal, 1890
<i>bicuspa</i> Sopott-Ehlers, 1976		A
<i>capitata</i> Meixner, 1938	A	
<i>macrostyla</i> Lanfranchi, 1978	M	Aspidogastrea
<i>multispinosa</i> Ax, 1956	M	Compiled by David I. Gibson
<i>papii</i> Ax, 1956	A	The list of aspidogastrean species was derived from the Host-Parasite Data-base and Catalogue maintained by the Parasitic Worms Division, Department of Zoology, The Natural History Museum.
<i>primitiva</i> Ax, 1956	A	
<i>procerostyla</i> Ax, 1956	M	Subclass Aspidogastrea
<i>progermaria</i> Ax, 1956	A	Order Aspidogastrida
<i>renatae</i> Ax, 1956	M	<i>Multicalyx</i>
<i>uncinata</i> Lanfranchi, 1978	M	<i>elegans</i> (Olsson, 1869)
<i>Parotoplanella</i>	M	<i>Lophotaspis</i>
<i>heterorhabditica</i> Lanfranchi, 1969	M	<i>vallei</i> (Stossich, 1899)
<i>progermaria</i> Ax, 1956	M	<i>Cotylogaster</i>
<i>Parotoplanina</i>	A	<i>michaelis</i> Monticelli, 1892
<i>geminoducta</i> Ax, 1956		Order Stichocotylida
<i>Philosyrtis</i>	A	<i>Stichocotyle</i>
<i>armoricae</i> Sopott-Ehlers, 1985	A	<i>nephropis</i> Cunningham, 1884
<i>coomansi</i> Martens & Schockaert, 1981	A	
<i>jennica</i> Ax, 1954	A	
<i>monotooides</i> Giard, 1904	A	
<i>rotundicephala</i> Sopott, 1972	A	
<i>rutilata</i> Sopott-Ehlers, 1976	A	
<i>Postbursoplana</i>	M	Digenea
<i>fibulata</i> Ax, 1956	M	Compiled by David I. Gibson
<i>macromystax</i> Lanfranchi, 1969	M	The list of digenetic species was derived from the Host-Parasite Data-base and Catalogue maintained by the Parasitic Worms Division, Department of Zoology, The Natural History Museum. Some attempt has been made to restrict the list to valid taxa and to use that latest nomenclature. Larval forms, e.g. <i>Cercaria</i> spp., have been omitted.
<i>minima</i> Ax, 1956	A	
<i>pontica</i> Ax, 1959	M	The list was kindly commented upon by Dr M. Køie and Prof. P. Bartoli.
<i>propontica</i> Ax, 1959	A	
<i>tyrrhenica</i> Lanfranchi, 1969	A	
<i>Praebursoplana</i>	A	Class Digenea
<i>reisingeri</i> Ax, 1956	A	<i>Acanthostomum</i>
<i>steinböcki</i> Ax, 1956	A	<i>absconditum</i> (Looss, 1901)
<i>subsalsa</i> Ax, 1956	A	<i>inerme</i> Stossich, 1905
<i>Pseudorthoplana</i>	A M	<i>saoudi</i> Hassan, El-Aziz, Khidr & Abu Samak, 1990
<i>soliacea</i> (Meixner, 1939) Ax, Weidemann & Ehlers, 1978		<i>Acanthotrema</i>
<i>subterranea</i> (Ax, 1951) Ax, 1956	A M	<i>sp.</i>
<i>Triporoplana</i>	A	<i>Accacladocoelium</i>
<i>synsiphonioides</i> Ax, 1956	M	<i>alveolatum</i> Robinson, 1934
<i>Xenotoplana</i>	M	<i>macrocotyle</i> (Diesing, 1858)
<i>acus</i> Ax, Weidemann & Ehlers, 1978	M	<i>nigroflavum</i> (Rudolphi, 1819)
<i>thyrrhenica</i> Ax, Weidemann & Ehlers, 1978	M	
Family Polystyliphoridae	A	
<i>Polystyliphora</i>		
<i>filum</i> Ax, 1958		
Suborder Tricladida		

- petasiporum* Odhner, 1928
Accacoelium
contortum (Rudolphi, 1819)
Adinosoma
robusta (Manter, 1934)
Allopodocotyle
jaffensis (Fischthal, 1980)
margolisi Gibson, 1995
pedicellata (Stossich, 1887)
Allopseudocolocytotrema
alioshkini Pozdnyakov, 1994
Anacetabulitrema
samarae Deblock & Rose, 1964
Ancylocoelium
typicum Nicoll, 1912
Anisocladium
fallax (Rudolphi, 1819)
Anisocoelium
capitellatum (Rudolphi, 1819)
Anisocladium
gracile (Looss, 1901)
Anoiktostoma
coronatum (Wagener, 1852)
Anomalotrema
koiae Gibson & Bray, 1984
Apatemon
gracilis (Rudolphi, 1819)
Aphalloides
coelomicola Dollfus, Chabaud & Golvan, 1957
Aphallus
tubarium (Rudolphi, 1819)
Aphanurus
stossichi (Monticelli, 1891)
Apharyngostrigea
intermedia (Tubangui, 1932)
Aponurus
laguncula Looss, 1907
tschugunovi Issatschikov, 1928
Apophallus
brevis Ransom, 1920
muehlingi (Jagerskiold, 1899)
Aporchis
massiliensis Timon-David, 1955
rugosus Linton, 1928
Aporocotyle
simplex Odhner, 1900
spinosicanalis Williams, 1958
Arnola
microcirrus (Vlasenko, 1931)
Ascocotyle
coleostoma (Looss, 1896)
Asymphylodora
demeli Markowski, 1935
Bacciger
bacciger (Rudolphi, 1819)
grandispinatus Naidenova, 1970
israelensis Fischthal, 1980
minutus Gaevskaja & Naidenova, 1996
Bathycreadium
biscayense Bray, 1973
elongata (Maillard, 1976)
flexicollis Kabata, 1961
Bathymonorchis
polyipni (Reimer, 1985)
Botulus
microporus (Mont., 1889)
Brachyenteron
campbelli Bray & Gibson, 1986
pycnorganum (Rees, 1953)
risssoanum Bray & Campbell, 1995
Brachylaima
fuscatus (Rudolphi, 1819)
Brachylecithum
microtesticulatum Timon-David, 1955
Brachyphallus
- crenatus* (Rudolphi, 1802)
Braunina
cordiformis Wolf, 1903
Bucephalopsis
callicotyle Kohn, 1962
Bucephalus
baeri Maillard, 1976
haiameanus (Lacaze-Duthiers, 1854)
labracis Paggi & Orecchia, 1965
marinus Vlasenko, 1931
mytili Cole, 1935
polymorphus Von Baer, 1827
Bunocotyle
cingulata Odhner, 1928
progenetica Chabaud & Buttner, 1959
Bunoderia
luciopercae (Müller, 1776)
Cainocreadium
labracis (Dujardin, 1845)
Calycodes
anthos (Braun, 1899)
Campula
oblonga Cobbald, 1858
Cardiocephalus
longicollis (Rudolphi, 1819)
Centroderma
spinosissima (Stossich, 1883)
Cercariooides
aharonii Witenburg, 1929
Chimaerohemecus
trondheimensis Van der Land, 1967
Chrisomon
tropicus (Manter, 1940)
Cirkennedyia
porlockensis Gibson & Bray, 1979
Ciureana
cryptocotyloides Issatschikoff, 1923
Coeliotrema
thynni Yamaguti, 1938
Complexobursa
sp.
Condylocotyla
pilodora Pearson & Prévôt, 1985
Copiateses
filiferus (Leuckart in Sars, 1885)
Crepidostomum
metoecus (Braun, 1900)
Crowcrocaecum
skrjabini (Iwanitzky, 1928)
Cryptocotyle
concava (Creplin, 1825)
jejuna (Nicoll, 1907)
lingua (Creplin, 1825)
Cymatocarpus
sp.
Derecrema
scorpaenicola Bartoli & Bray, 1990
sebastodis (Yamaguti, 1934)
Deroenes
adriaicus Nikolaeva, 1966
affine (Rudolphi, 1819)
crassus Manter, 1934
latus Janiszewska, 1953
macrouri Szukla, 1981
minor (Looss, 1901)
ruber Luhe, 1900
varicus (Müller, 1784)
Deroenoides
ovacutus Nicoll, 1913
Deroprtis
hispida (Rudolphi, 1819)
inflata (Molin, 1859)
Deuterobaris
proteus (Brandes, 1891)
Dexiogonimus

- ciureanus* Witenburg, 1929
Diaschistorchis
 pandus (Braun, 1901)
Dicrogaster
 contractus Looss, 1902
 perpusilla Looss, 1902
Didymocystis
 lanceolata Guiart, 1938
 macrorchis Guiart, 1938
 philobranchia Yamaguti, 1970
 reniformis Ariola, 1902
 thynni (Taschenberg, 1879)
 xiphadios (MacCallum & MacCallum, 1916)
Didymodichinus
 faciale (Baylis, 1938)
Didymosulcus
 wedli (Ariola, 1902)
Didymozoon
 auxis Taschenberg, 1879
 pretiosum Ariola, 1902
 serrani Monticelli, 1889
 sphyraenae Taschenberg, 1878
 taenioides Monticelli, 1883
Dihemistephanus
 lydiae (Stossich, 1896)
 sturionis Little, 1930
Dinosoma
 hawaiiense Yamaguti, 1970
 thoria Nicolaeva, 1966
 triangulata Campbell & Munroe, 1977
 ventrovesicularis Gibson & Bray, 1986
Dinurus
 breviductus Looss, 1907
 sombri Yamaguti, 1934
 tornatus Rudolphi, 1819
Diphterostomum
 betencourtii (Monticelli, 1893)
 brusinae (Stossich, 1889)
 israelense Fischthal, 1980
 luteum (van Beneden, 1870)
 vividum (Nicoll, 1912)
Diplostomum
 baeri Dubois, 1937
 chromatophorum (Brown, 1931)
 commutatum (Diesing, 1850)
 gasterostei Williams, 1966
 helveticum (Dubois, 1929)
 huronense (La Rue, 1927)
 nordmanni Shigin & Shapirov, 1986
 paraspatheraceum Shigin, 1965
 phoxini (Faust, 1918)
 pseudospathaceum Niewiadomska, 1984
 rutili Razmashkin, 1969
 shigini Zhatkanbaeva, 1978
 spathaceum (Rudolphi, 1819)
 spathaceum (Rudolphi, 1819)
 volvens Nordmann, 1832
Dolichoenterum
 congeri Yamaguti, 1970
 longissimum Ozaki, 1924
Echeneidocoelium
 indicum Simha & Pershad, 1964
Echinochasmus
 mordax (Looss, 1899)
Echinoparyphium
 paracinctum Bychowskaja-Pawlowskaja, 1953
 recurvatum (Linstow, 1873)
Echinostephilla
 patellae (Lebour, 1911)
Echinostoma
 nordiana (Baschirova, 1941)
 parvulum Dietz, 1909
 revolutum (Froehlich, 1802)
Ectenurus
 lepidocybii Yamaguti, 1970
- lepidus* Looss, 1907
trachuri Nicolaeva & Kovaleva, 1966
virgulus Looss, 1910
Elstia
 stossichianum (Monticelli, 1892)
Enodiotrema
 carettae (Blair & Limpus, 1982)
 instar (Looss, 1901)
 megachondrus Looss, 1901
Eucole
 wehri Price, 1930
Fasciola
 skrjabini Delyamure, 1955
Felodistomum
 agnotus (Nicoll, 1909)
 fellis (Olsson, 1868)
Forticulcita
 glabra Overstreet, 1982
Gaevskajatrema
 halosauropsi Bray & Campbell, 1996
 perezi (Mathias, 1926)
 pontica (Koval, 1966)
Galactosomum
 baylisi (Gohar, 1930)
 dolfusi Pearson, 1973
 erinaceum (Poirier, 1886)
 lacteum (Jägerskiöld, 1896)
 phalacrococaris Yamaguti, 1939
 sp. metacerc.
 timondavidi Pearson & Prévot, 1971
Genitocotyle
 mediterranea Bartoli, Gibson & Riutort, 1994
Genolinea
 laticauda Manter, 1925
Gigantobilharzia
 acotylea Odhner, 1910
Glomericirrus
 macrouri (Gaevskaja, 1973)
Gonapodasmius
 branchialis Yamaguti, 1970
 spilonotopteri Yamaguti, 1970
Gonocerca
 macrouri Gaevskaya, 1975
 phycidis Manter, 1925
Gonocercella
 sp.
Gymnophallus
 choledochus Odhner, 1900
 deliciosus (Olsson, 1893)
 gibberosus Loos-Frank, 1971
 rostratus Bartoli, 1982
 somateriae (Levinsen, 1881)
Gynaecotyla
 longiintestinata Leonov, 1958
Hadwenius
 delamarei (Raga & Balbuena, 1988)
 subtilis (Skrjabin, 1959)
 tursionis (Marchi, 1873)
Halvorsenius
 exilis Gibson, MacKenzie & Cottle, 1981
Hapalotrema
 flecterotestis Zhukov, 1971
 loossii Price, 1934
 mistroides (Monticelli, 1896)
Haploporus
 benedenii (Stossich, 1887)
Haplosplanchus
 bivitellosus Zhukov, 1971
 caudatus (Srivastava, 1937)
 pachysomus (Eysenhardt, 1829)
Helicometra
 fasciata (Rudolphi, 1819)
 flava Stossich, 1903
 gobii (Stossich, 1883)
 labri Stossich, 1886

- markewitschi* Pogoreltseva, 1954
mutabilis (Stosich, 1902)
pulchella (Rudolphi, 1819)
sinuata (Rudolphi, 1819)
- Hemipera*
ovocaudata Nicoll, 1913
sharpei Jones, 1933
- Hemiuirus*
appendiculatus (Rudolphi, 1802)
communis Odhner, 1905
levinseri Odhner, 1905
luehei Odhner, 1905
- Heterophyes*
aqualis Looss, 1902
heterophyes (Siebold, 1853)
- Heterotestophyes*
sobolevi Leonov, 1957
- Himasthla*
continua Loos-Frank, 1967
elongata (Mehlis, 1831)
leptosoma (Creplin, 1829)
militaris (Rudolphi, 1802)
secunda (Nicoll, 1906)
- Hirudinella*
ventricosa (Pallas, 1774)
- Holorchis*
legendrei Dollfus, 1946
micracanthum (Stosich, 1888)
pycnoporus Stosich, 1901
- Holostephanus*
dubini Vojtek & Vojtкова, 1968
- Homalometron*
galacicus (Sanmartin, Alvarez, Quinteiro & Paniagua, 1995)
- Hymenocotta*
mugilis Wang & Wang, 1993
- Hyperandrotrema*
cetorhini Maillard & Ktari, 1978
- Hypertrema*
ambovatum Manter, 1960
- Hypoderaeum*
conoideum (Bloch, 1782)
- Hysterolecitha*
nahaensis Yamaguti, 1942
- Hysteromorpha*
triloba (Rudolphi, 1819)
- Ichthyocotylurus*
erraticus (Rudolphi, 1809)
pileatus (Rudolphi, 1802)
platycephalus (Creplin, 1825)
variegatus (Creplin, 1825)
- Kenmackenzia*
gigas (Nardo, 1827)
- Knipowitschia*
tremula (Issatchikow, 1927)
- Koellikeria*
benedenii (Monticelli, 1893)
biaperta (Wedl, 1855)
exocoeti (Parona & Perugia, 1893)
filiicollis (Rudolphi, 1819)
micropterygis (Richardi, 1902)
- Koellikeroides*
orientalis (Yamaguti, 1934)
- Labratrema*
minimus (Stosich, 1887)
- Lacunovermis*
macomae (Lebour, 1908)
- Lampritrema*
meischeri (Zschokke, 1890)
- Lasiotocus*
mulli (Stosich, 1883)
- Learedius*
europaeus Price, 1934
- Lecithaster*
atherinae Orecchia, Paggi & Radujkovic, 1988
- confusus* Odhner, 1905
gibbosus (Rudolphi, 1802)
stellatus Looss, 1906
- Lecithobotrys*
aegyptiacus Hassan, El-Aziz, Khidr Abu & Samak, 1990
putrescens Looss, 1902
- Lecithochirium*
branchialis (Stunkard & Nigrelli, 1934)
furcolabiatum (Jones, 1933)
fusiforme Luhe, 1901
gravidum Looss, 1907
microstomum Chandler, 1935
musculus (Looss, 1907)
physcon Lühe, 1901
rufoviride (Rudolphi, 1819)
- Lecithocladium*
aegyptense Fischthal & Kuntz, 1963
cristatum (Rudolphi, 1819)
excisum (Rudolphi, 1819)
unibulabolabrum Fischthal & Thomas, 1971
- Lecithodesmus*
delphini (Poirier, 1886)
goliath (van Beneden, 1858)
palliatus (Looss, 1885)
- Lecithophyllum*
botryophorum (Olsson, 1868)
magnaprostatica Gaevskaya & Kovaljova in Gaevskaja, 1989
- Lecithostaphylus*
retroflexus Molin, 1859
- Lepidapedon*
arlenae Bray & Gibson, 1995
beveridgei Campbell & Bray, 1993
cambrensis Srivastava, 1966
carcini Mordvinova, 1985
desclerseae Bray & Gibson, 1995
discoveryi Bray & Gibson, 1995
elongatum (Lebour, 1908)
gaevskiae Campbell & Bray, 1993
guevarai Lopez-Roman & Maillard, 1973
mariannae Bray & Gibson, 1995
rachion (Cobbold, 1858)
sommervilleae Bray & Gibson, 1995
zubchenkoi Campbell & Bray, 1993
- Lepidauchen*
stenostoma Nicoll, 1913
- Lepidophyllum*
appyi Bray & Gibson, 1986
steenstrupi Odhner, 1902
- Lepocreadium*
album (Stosich, 1890)
floridanus Sogandares-Bernal & Hutton, 1959
pegorchis (Stosich, 1901)
trullaforme Linton, 1940
- Leucochloridium*
perturbatum Pojmanska, 1969
- Levinsemiella*
brachysoma (Creplin, 1837)
propinquum Jaggerskiold, 1907
- Lintonium*
vibex (Linton, 1905)
- Lomasoma*
gracilis (Manter, 1934)
wardi (Manter, 1934)
- Longiductotrema*
scandolensis Deblock & Bartoli, 1988
- Macvicaria*
alacris (Looss, 1901)
crassigula (Linton, 1910)
dubia (Stosich, 1905)
mailliardi Bartoli, Bray & Gibson, 1989
mormyri (Stosich, 1885)
obovata (Molin, 1859)
soleae (Dujardin, 1845)

<i>Magnibursatus</i>	<i>oculatus</i> (Levinsen, 1881)
<i>caudofilamentosa</i> (Reimer, 1971)	
<i>skrjabini</i> (Vlasenko, 1931)	
<i>Maritrema</i>	
<i>arenaria</i> Hadley & Castle, 1940	
<i>echinocirrata</i> Leonov, 1958	
<i>misenensis</i> (Palombi, 1940)	
<i>oocysta</i> Lebour, 1907	
<i>opisthometra</i> Leonov, 1958	
<i>subdolum</i> Jagerskiold, 1909	
<i>syntomocyclos</i> Deblock & Ky, 1966	
<i>Megalophallus</i>	
<i>carcini</i> Prévôt & Deblock, 1970	
<i>Meiogymnophallus</i>	
<i>fossarum</i> (Bartoli, 1965)	
<i>jamesoni</i> Bowers, 1965	
<i>minutus</i> (Cobbald, 1859)	
<i>rebecqui</i> (Bartoli, 1983)	
<i>strigatus</i> (Lebour, 1908)	
<i>Merlucciotrema</i>	
<i>praeclarum</i> (Manter, 1934)	
<i>Mesometra</i>	
<i>brachycoelia</i> Luhe, 1901	
<i>orbicularis</i> (Rudolphi, 1819)	
<i>Mesorchis</i>	
<i>denticulatus</i> (Rudolphi, 1802)	
<i>pseudoechinatus</i> (Olsson, 1876)	
<i>reynoldi</i> Bhalerao, 1926	
<i>Metadena</i>	
<i>caballeroi</i> Nahhas & Krupin, 1977	
<i>depressa</i> (Stossich, 1883)	
<i>pauli</i> (Vlassenko, 1931)	
<i>Metanematobothrium</i>	
<i>guernei</i> (Moniez, 1891)	
<i>Metorchis</i>	
<i>xanthostomus</i> (Creplin, 1846)	
<i>Microphallus</i>	
<i>abortivus</i> Deblock, 1974	
<i>claviformes</i> (Brandes, 1888)	
<i>fusiformis</i> Reimer, 1963	
<i>hoffmanni</i> Rebecq, 1964	
<i>papillorobustus</i> (Rankin, 1940)	
<i>piriformis</i> (Odhner, 1905)	
<i>pirum</i> (Lebour, 1907)	
<i>primas</i> (Jagerskiold, 1908)	
<i>pygmaeus</i> (Levinsen, 1881)	
<i>similis</i> (Jagerskiold, 1900)	
<i>Microscaphidium</i>	
<i>reticulare</i> (van Beneden, 1859)	
<i>Monascus</i>	
<i>filiformis</i> (Rudolphi, 1819)	
<i>Monodhelmis</i>	
<i>torpedinis</i> Dollfus, 1937	
<i>Monorchis</i>	
<i>monorchis</i> (Stossich, 1890)	
<i>parvus</i> Looss, 1902	
<i>Nematobothrium</i>	
<i>faciale</i> (Baylis, 1938)	
<i>filarina</i> van Beneden, 1858	
<i>filiforme</i> Yamaguti, 1934	
<i>latum</i> Guiart, 1938	
<i>molae</i> (Maclarens, 1904)	
<i>robustum</i> Yamaguti, 1970	
<i>scombri</i> (Taschenberg, 1879)	
<i>Neodiplostomum</i>	
sp.	
<i>Neolepidapedon</i>	
<i>israelense</i> Fischthal, 1980	
<i>smithi</i> Bray & Gibson, 1989	
<i>Neolepocreadium</i>	
<i>caballeroi</i> Thomas, 1960	
<i>Neonotoporus</i>	
<i>trachuri</i> (Yamaguti, 1938)	
<i>Neophasis</i>	
<i>anarrhichiae</i> (Nicoll, 1909)	
<i>Neosteganoderma</i>	
<i>glandulosum</i> Byrd, 1964	
<i>Nephromonorchia</i>	
<i>lari</i> (Timon-David, 1933)	
<i>Nicolla</i>	
<i>gallica</i> (Dollfus, 1941)	
<i>Notocotylus</i>	
<i>attenuatus</i> (Rudolphi, 1809)	
<i>seineti</i> Fuhrmann, 1919	
<i>skrjabini</i> Ablasov, 1953	
<i>Odhneriella</i>	
<i>elongata</i> (Ozaki, 1935)	
<i>Odhnerium</i>	
<i>calyptocotyle</i> (Monticelli, 1893)	
<i>Oesophagocystis</i>	
<i>lydiae</i> Pozdnyakov, 1994	
<i>Ogmogaster</i>	
<i>antarcticus</i> Johnston, 1931	
<i>plicatus</i> (Creplin, 1829)	
<i>Olssonium</i>	
<i>turneri</i> Bray & Gibson, 1980	
<i>Opechona</i>	
<i>bacillaris</i> (Molin, 1859)	
<i>Opecoelina</i>	
<i>scorpaenae</i> Manter, 1934	
<i>Opecoeloides</i>	
<i>furcatus</i> (Bremser in Rudolphi, 1819)	
<i>vitellus</i> (Linton, 1900)	
<i>Opisthadena</i>	
<i>karachii</i> (Srivastava, 1941)	
<i>kyphosi</i> Yamaguti, 1970	
<i>Opisthorchis</i>	
<i>gomtii</i> Mehra, 1941	
<i>Orchidasma</i>	
<i>amphiorchis</i> (Braun, 1899)	
<i>Ornithobilharzia</i>	
<i>canaliculata</i> (Rudolphi, 1819)	
<i>Orophocotyle</i>	
<i>divergens</i> Looss, 1902	
<i>planci</i> (Stossich, 1899)	
<i>Orthosplanchnus</i>	
<i>arcticus</i> Odhner, 1905	
<i>Oschmarinella</i>	
<i>albamarina</i> (Treshchev, 1968)	
<i>mascomai</i> Raga, 1986	
<i>rochebruni</i> (Poirier, 1886)	
<i>Otodistomum</i>	
<i>cestoides</i> (van Beneden, 1871)	
<i>plunketi</i> Fyfe, 1953	
<i>veliporum</i> Dollfus, 1937	
<i>Pachycreadium</i>	
<i>carnosum</i> (Rudolphi, 1819)	
<i>Pachypsopus</i>	
<i>irroratus</i> (Rudolphi, 1819)	
<i>Pachytrema</i>	
<i>calculus</i> Looss, 1907	
<i>paniceum</i> Brinkmann, 1942	
<i>Panopula</i>	
<i>bridgeri</i> Bray & Gibson, 1986	
<i>Paracacacladium</i>	
<i>jamiesoni</i> Bray & Gibson, 1977	
<i>Paracryptogonimus</i>	
<i>aloysiae</i> (Stossich, 1885)	
<i>Paragyliachen</i>	
<i>nagatyi</i> of Rizk, Tanha & El-Rahman, 1996	
<i>Parahemius</i>	
<i>merus</i> (Linton, 1910)	
<i>Paralepidapedon</i>	
<i>williamsi</i> Bray & Gibson, 1988	
<i>Paramonostomum</i>	
<i>alveatum</i> (Mehlis, 1846)	
<i>anatis</i> Garkawi, 1965	
<i>bucephala</i> Yamaguti, 1935	
<i>chabaudi</i> van Strydonck, 1965	

- pseudoalveatum* Price, 1931
Parasaccocoelium
mugili Zhukov, 1971
Parascocotyle
 sp.
Paratimonia
gobii Prévôt & Bartoli, 1967
Parvatrema
affine (Jameson & Nicoll, 1913)
duboi (Dollfus, 1923)
homoeotecnum James, 1964
ovoplemum (Jameson & Nicoll, 1913)
timondavidi Bartoli, 1983
Paryphostomum
radiatum (Dujardin, 1845)
Peracreadium
characis (Stossich, 1886)
commune Olsson
genu Nicoll, 1909
idoneum (Nicoll, 1909)
Petasiger
exaeretus Dietz, 1909
hosptiale (Mendheim, 1940)
phalacrocoracis (Yamaguti, 1939)
Phagicola
longus (Ransom, 1920)
septentrionalis van den Broek, 1963
Philophthalmus
lacrymosus Braun, 1902
lucipetus (Rud., 1819)
nocturnus Looss, 1907
Pholeter
gastrophilus (Kossack, 1910)
Phyllodistomum
acceptum Looss, 1901
crenilabri Dolgikh & Naidenova, 1968
thunni Baudin-Laurencin & Richard, 1973
Plagioporos
congeri Yamaguti, 1970
dogieli Pogoreltseva, 1975
trachuri Pogoreltseva, 1954
tumidulus (Rudolphi, 1819)
tyrrhenicus Paggi & Orecchia, 1976
Plagiorchis
elegans (Rudolphi, 1802)
laricola Skrjabin, 1924
moravicus Sitko, 1993
mutationis Panova, 1927
nanus (Rudolphi, 1802)
Platocystis
alalongae Yamaguti, 1938
 sp.
Pleorchis
polyorchis (Stossich, 1889)
Plerurus
digitatus (Looss, 1899)
Plesiochorus
cymbiformis (Rudolphi, 1819)
Pleurogonius
longiusculus Looss, 1901
trigonocephalus (Rudolphi, 1809)
Podocotyle
alexandrina of Rizk, Tanha & El-Rahman, 1996
angulata (Dujardin, 1845)
atherinae Nicoll, 1914
atomon (Rudolphi, 1802)
olssoni Odhner, 1905
reflexa (Creplin, 1825)
scorpaenae (Rudolphi, 1919)
staffordi Miller, 1941
syngnathi Nicoll, 1913
Polylekithum
trachuri of Rizk, Tanha & El-Rahman, 1996
Poracanthium
furcatum Dollfus, 1948
Posthodiplostomum
brevicaudatum (Nordmann, 1832)
Probolitrema
richardii (Lopez, 1888)
Proctoeces
maculatus (Looss, 1901)
major Yamaguti, 1938
Proctophantastes
abyssorum Odhner, 1900
Proctotrema
bacilliovatum (Odhner, 1911)
Prodistomum
orientalis (Layman, 1930)
polonii (Molin, 1859)
priedei Bray & Merrett, 1998
Profundivermis
intercalarius Bray & Gibson, 1991
Progonus
muelleri (Levinsen, 1881)
Pronoprymna
ventricosum (Rudolphi, 1819)
Proparipyrum
israelense Fischthal, 1980
Prosorhynchoides
arcuatus (Linton, 1900)
gracilescens (Rudolphi, 1819)
Prosorhynchus
aculeatus Odhner, 1905
crucibulum Rudolphi, 1819
squamatus Odhner, 1905
Prosthognomus
ovatus (Rudolphi, 1803)
Prudhoeus
nicholsi Bray & Gibson, 1980
Pseudaeephnidiogenes
africanus (Fischthal & Thomas, 1972)
Pseudamphistomum
truncatum (Rudolphi, 1819)
Pseudobacciger
harengulae (Yamaguti, 1938)
manteri Nahhas & Cable, 1964
Pseudomaritrema
innae Leonov, 1958
Pseudopycnadema
fischthali Saad-Fares & Maillard, 1986
Pseudozoogonoides
subaequiporus (Odhner, 1911)
Psilochasmus
longicirratus Skrjabin, 1913
oxyurus (Creplin, 1825)
skrjabini Gnedina, 1946
Psilostomum
brevicolle (Creplin, 1829)
brevicolle (Creplin, 1829)
Ptychogonimus
megastomus (Rudolphi, 1819)
Pycnadenoides
ghanensis Fischthal & Thomas, 1968
senegalensis Fischthal & Thomas, 1972
umbrinae (Stossich, 1885)
Pyelosomum
renicapite (Leidy, 1856)
Pygidiosis
ardeae Krie, 1990
genata Looss, 1907
Renicola
lari Timon-David, 1933
paraquinata Rajewsky, 1937
roscovita (Stunkard, 1932)
sudarikovi Leonov, 1958
Rhipidocotyle
galeata (Rudolphi, 1819)
genovi Dimitrov, Kostadinova & Gibson, 1996
illense (Ziegler, 1883)
Rhynchopharynx

- paradoxa* Odhner, 1928
Rhytidodes
gelatinosus (Rudolphi, 1819)
Robphildolfusium
fractum (Rudolphi, 1819)
martinezgomezi Lopez- Roman, Gijon-Botella, Kim & Vilca-Choque, 1992
Rossicotrema
lari Leonov, 1957
Saccocoelium
obesum Looss, 1902
tensum Looss, 1902
Saturnius
papernai Overstreet, 1977
Schikhobalotrema
acutum (Linton, 1910)
longivesiculatum Orecchia & Paggi, 1975
Sclerodistomum
italicum (Stossich, 1893)
saoudi of Rizk, Tanha & El-Rahman, 1996
Skrjabinolecithum
spaskii Belous, 1954
Skrjabinopslus
semiaromaticus (Molin, 1858)
Spelotrema
excellens Nicoll, 1907
magnipapillatum Leonov, 1958
simile (Jagerskiold, 1900)
Sphaerostoma
 sp.
Spinoplagioporus
minutus Polyansky, 1952
Steganoherma
retroflexum (Molin, 1859)
Steganodermatooides
agassizi (Campbell, 1975)
maceri Bray & Gibson, 1986
Stenakron
vetustum Stafford, 1904
Stephanostomum
baccatum (Nicoll, 1907)
bicornutum (Stossich, 1883)
caducum (Looss, 1901)
cesticillum (Molin, 1858)
lebourae Caballero, 1952
longisomum Manter, 1940
lophii Quinteiro, Tojo, Nunez, Santamarina & Sammartin, 1993
minutum (Looss, 1901)
pristis (Deslongchamps, 1824)
rhombispinosum (Lebour, 1908)
seriolae Yamaguti, 1970
triglae (Lebour, 1908)
valdeinflatum (Stossich, 1883)
Steringophorus
blackeri Bray, 1973
dorsolineatum (Reimer, 1985)
furciger (Olsson, 1867)
haedrichi Bray & Campbell, 1995
margolisi Bray, 1995
pritchardae (Campbell, 1975)
thulini Bray & Gibson, 1980
Steringotrema
divergens (Rudolphi, 1809)
ovacutum (Lebour, 1908)
pagelli (van Beneden, 1871)
Stictodora
lari Yamaguti, 1939
morosovi Leonov, 1957
savakinensis Looss, 1899
Synaptobothrium
caudiporum (Rudolphi, 1819)
Tanaisia
fedtschenkoi Skrjabin, 1924
Tellervotrema
- armstrongi* Gibson & Bray, 1982
Telorhuis
aculeatus Stunkard, 1915
parvus Braun, 1901
poirieri (Stossich, 1895)
stossichi Goldberger, 1911
Tergestia
acanthocephala (Stossich, 1887)
laticollis (Rudolphi, 1819)
Tetracladium
sternae Kulackowa, 1952
Tetrochetus
raynerii (Nardo, 1833)
sombresoxis Nicolaeva, 1966
Thulinia
microrchis (Yamaguti, 1934)
Timonia
mediterranea Bartoli & Prevot, 1966
Timoniella
imbutiforme (Molin, 1859)
praeteritum (Looss, 1901)
Tormopsisulus
oscultatus (Looss, 1901)
orientalis (Yamaguti, 1934)
Tricharrhen
okenii (Koelliker, 1849)
Trichobilharzia
ocellata (La Valette, 1855)
Tylodelphys
clavata (von Nordmann, 1832)
Tyrrhenia
blennii Paggi & Orecchia, 1975
Unitubulotestis
pelamydis (Taschenburg, 1879)
Wardula
capitellata (Rudolphi, 1819)
sarguicola Bartoli & Gibson, 1989
Zalophotrema
atlanticum Abril, Balbuena & Raga, 1991
Zoogonoides
viviparus (Olsson, 1868)
Zoogonus
rubellus (Olsson, 1868)

Monogenea

Compiled by Rodney A. Bray

The list of monogenean species was derived from the Host-Parasite Data-base and Catalogue maintained by the Parasitic Worms Division, Department of Zoology, The Natural History Museum. Some attempt has been made to restrict the list to valid taxa and to use that latest nomenclature.

The list was kindly commented upon by Prof. L. Euzet and Dr G. Kearn.

Class Monogenea

- Acanthocotyle*
elegans Monticelli, 1890
borealis Brinkmann, 1940
lobianchi Monticelli, 1888
monticellii Scott, 1902
oligoterus (Monticelli, 1899)
Allopseudaxine
 sp.
Amphibdella
flavolineata MacCallum, 1916
paronaperugiae Llewellyn, 1960
torpedinis Chatin, 1874
Amphibdelloides
maccallumi (Johnson & Tiegs, 1922)
vallei Llewellyn, 1960
Ancyrocephalus

- cobitis* Ergens, 1963
paradoxus Creplin, 1839
vanbenedenii (Parona & Perugia, 1890)
- Ancyrocotyle*
vallei (Parona & Perugia, 1895)
Anoplodiscus
richardi (Sonsino, 1890)
- Anthocotyle*
merluccii (Van Beneden & Hesse, 1863)
- Aspinatrium*
euzeti Ktari, 1971
gallieni Euzet & Ktari, 1971
spari (Yamaguti, 1937)
trachini (Parona & Perugia, 1889)
- Atriaster*
heterodus Lebedev & Parukhin, 1969
maillardi Lopez-Roman & De Armas Hernandez, 1989
- Atrispinum*
acarne Maillard & Noisy, 1979
labracis (Van Beneden & Hesse, 1863)
mormyri (Lorenz, 1878)
salpae (Parona & Perugia, 1890)
sargi (Parona & Perugia, 1890)
seminalis (Euzet & Maillard, 1973)
- Axine*
belones Abildgaard, 1794
- Benedenia*
innobilitata Burnheim, Gomes & Varela, 1973
leucanthemum Euzet & Maillard, 1967
micracantha Euzet & Maillard, 1967
monticellii (Parona & Perugia, 1890)
pompatica Burnheim, Gomes & Varela, 1973
sciaenae (Beneden, 1856)
- Bivagina*
alcedinis (Parona & Perugia, 1889)
- Bothitrema*
bothi (MacCallum, 1913)
- Caballerocotyla*
abidjani Bussieras & Baudin-Laurencin, 1970
pelamydis (Taschenberg, 1878)
pseudomagronum Busserias, 1972
verrucosa Busserias, 1972
- Cadenia*
polytestis Euzet & Maillard, 1967
- Calceostoma*
calceostoma (Wagener, 1857)
herculanea Euzet & Vala, 1976
paronai (Parona & Perugia, 1892)
- Calceostomella*
inermis (Parona & Perugia, 1889)
- Calicotyle*
affinis Scott, 1911
kroyeri Diesing, 1850
palombi Euzet & Williams, 1970
stossichi Braun, 1899
- Capsala*
martinieri Bosc, 1811
pelamydis (Taschenberg, 1878)
thynni (Guia, 1938)
- Capsaloides*
perugiae (Setti, 1898)
- Cathariotrema*
 sp.
- Cemocotyle*
trachuri Dillon & Hargis, 1965
- Chauhanea*
mediterranea Euzet & Trilles, 1960
- Chimaericola*
leptogaster (Leuckart, 1830)
- Choricotyle*
chrysophryi (van Beneden & Hesse, 1863)
- Cleithranticus*
 sp.
- Cotyloatlantica*
- mediterranea* (Euzet & Trilles, 1960)
- Cyclocotyla*
charcoti (Doilfus, 1922)
- Cycloplectanum*
beverleyburtonae Oliver, 1984
echinophallus (Euzet & Oliver, 1965)
riouxi Oliver, 1986
- Dactylogyrus*
 sp.
- Dasybatotrema*
dasyatis (MacCallum, 1916)
- Dendromonocotyle*
octodiscus Hargis, 1955
taeniurae Euzet & Maillard, 1967
- Dermophthirius*
carchariphini MacCallum, 1926
- Diclidophora*
bellones (Otto, 1821)
denticulata (Olsson, 1876)
esmarkii (Scott, 1901)
luscae (van Beneden & Hesse, 1864)
macruri (Brinkmann, 1942)
merlangi (Kuhn In Nordmann, 1832)
minor (Olson, 1868)
minuti Tirard, Berrebi, Raibaut & Frenaud, 1992
pagelli Gallien, 1937
palmata (Leuckart, 1830)
phycidis (Parona & Perugia, 1889)
pollachii (van Beneden & Hesse, 1863)
- Diclidophoropsis*
taschenbergii (Parona & Perugia, 1899)
tissieri Gallien, 1937
- Dicybothrium*
armatum Leuckart, 1835
- Dicrunenia*
bichowskyi Mamaev, 1969
- Dictyocotyle*
coeliaca Nybelin, 1941
- Dionchus*
agassizi Goto, 1899
remorae (MacCallum, 1916)
- Diplectanotrema*
 sp.
- Diplectanum*
aculeatum Parona & Perugia, 1889
aequans (Wagener, 1857)
banyulense Oliver, 1968
bilobatum Hargis, 1955
bocqueti Oliver, 1980
chabaudi Oliver, 1980
dolfusi Oliver, 1980
elongatum Obiekezie, 1988
enyenihii Obiekezie, 1988
grassei Oliver, 1974
hargisi Oliver & Paperna, 1984
jamestownense Obiekezie, 1988
labourgi Oliver, 1973
laubieri Lambert & Maillard, 1974
melvillei Oliver & Paperna, 1984
nagibinae Oliver & Paperna, 1984
sciaenae van Beneden & Hesse, 1863
simile Bychowsky, 1957
- Diplozoon*
paradoxum von Nordmann, 1832
- Discocotyle*
sagittata (Leuckart, 1842)
- Empruthotrema*
raiae (MacCallum, 1916)
torpedinis Kearn, 1976
- Encystylabe*
nordmanni Diesing, 1850
pagelli van Beneden & Hesse, 1863
paronae Monticelli, 1907
spari Yamaguti, 1934
vallei Monticelli, 1907

- Enoplocotyle*
minima Tagliani, 1909
- Entobdella*
aegyptiacus Amer, 1990
apiocolpos Euzet & Maillard, 1967
diadema (Monticelli, 1902)
hippoglossi (Mueller, 1776)
soleae (van Beneden & Hesse, 1864)
- Epicotyle*
torpedinis (Price, 1942)
- Ergenstrema*
labrosi Anderson, 1981
mugilis Paperma, 1964
- Erpocotyle*
catenulata (Guberlet, 1933)
abbreviata (Olsson, 1876)
borealis (van Beneden, 1853)
canis (Cerfontaine, 1899)
dubium Euzet & Maillard, 1967
grisea (Cerfontaine, 1899)
laevis van Beneden & Hesse, 1863
maccallumi (Price, 1942)
sphyrae (MacCallum, 1931)
- Flexophora*
ophidii Euzet & Prost, 1962
- Furnestinia*
echenensis (Wagener, 1857)
- Gastrocotyle*
trachuri van Beneden & Hesse, 1863
- Gempylitrema*
longipedunculatum Yamaguti, 1968
- Gotocotyla*
acanthura (Parona & Perugia, 1891)
- Grubea*
cochlear Diesing, 1858
pneumatophori Price, 1961
- Gyrodactyloides*
bychowskii Albova, 1948
- Gyrodactylus*
alviga Gaevskaya & Dmitrieva, 1967
atherinae Bychowsky, 1933
blennii Gaevskaya & Dmitrieva, 1967
bubyri Osmanov, 1965
callariatis Malmberg, 1956
crenilabri Zaika, 1966
cryptarum Malmberg, 1970
cyclopteri Scyborskaya, 1948
derjavini Mikailov, 1975
emembranatus Malmberg, 1970
errabundus Malmberg, 1970
flesi Malmberg, 1956
flexibiliradix Malmberg, 1970
harengi Malmberg, 1956
lavareti Malmberg, 1957
longidactylus Geets, Malmberg & Ollevier, 1998
marinus Bychowsky & Polyansky, 1953
mugili Zhukov, 1970
najdenovi Malmberg, 1970
pharyngicus Malmberg, 1964
proterorhini Ergens, 1964
ptyrgialis Bychowsky & Polyansky, 1953
salaris Malmberg, 1957
syngnathi Appleby, 1996
zhukovi Ling, 1962
- Haliotrema*
cupensis Sasal, Pages & Euzet, 1998
kritskyi Vala, Mailard & Overstreet, 1982
- Heliocotyle*
kartasi Neifar, Euzet & Ben Hassine, 1999
- Heteraxine*
louisae Lopez-Roman & De Armas Hernandez, 1989
seriolae (Ishii, 1936)
- Heteraxinoides*
atlanticus Gaevskaya & Kovaliova, 1979
- hannibali* Euzet & Ktari, 1970
- Heterocotyle*
pastinaceae Scott, 1904
scotti Neifar, Euzet & Ben Hassine, 1998
similis Neifar, Euzet & Ben Hassine, 1998
- Heteronchocotyle*
hypoproni Brooks, 1934
- Hexabothrium*
appendiculatum (Kuhn, 1829)
musteli (MacCallum, 1931)
canicula (Cerfontaine, 1899)
- Hexostoma*
auxisi Palombi, 1943
thynni (Delaroche, 1811)
- Ktariella*
polyorchis Euzet & Valat, 1977
- Kuhnia*
scomtri (Kuhn, 1829)
sprostonae Price, 1961
- Lamellodiscus*
baeri Oliver, 1974
bidens Euzet, 1984
corallinus Paperma, 1965
coronatus Euzet & Oliver, 1966
drummondi Euzet & Oliver, 1967
elegans Bychowsky, 1957
ergensi Euzet & Oliver, 1966
erythrini Euzet & Oliver, 1967
fraternus Bychowsky, 1957
furcosus Euzet & Oliver, 1966
gracilis Euzet & Oliver, 1966
hilii Euzet, 1984
ignoratus Palombi, 1943
impervius Euzet, 1984
knoeffleri Oliver, 1969
mirandus Euzet & Oliver, 1966
mormyri Euzet & Oliver, 1967
obeliae Oliver, 1973
parisi Oliver, 1969
verberis Euzet & Oliver, 1967
virgula Euzet & Oliver, 1967
- Leptobothrium*
pristiuri Gallien, 1937
- Leptocotyle*
minor (Monticelli, 1888)
- Ligophorus*
acuminatus Euzet & Suriano, 1977
angustus Euzet & Suriano, 1977
chabaudi Euzet & Suriano, 1977
confusus Euzet & Suriano, 1977
euzeti Gaevskaya & Dmitrieva, 1997
heteronchus Euzet & Suriano, 1977
imitans Euzet & Suriano, 1977
kaohsianghsieni Gussev, 1962
macrocolpus Euzet & Suriano, 1977
minimus Euzet & Suriano, 1977
mugilinus (Hargis, 1955)
parvicirrus Euzet & Sanfilippo, 1983
szidati Euzet & Suriano, 1977
vanbenedenii (Parona & Perugia, 1890)
- Linguadactyla*
molvae Brinkmann, 1940
- Loimosina*
wilsoni Manter, 1944
- Macrouridophora*
attenuata (Mamaev & Zubchenko, 1979)
caudata (Mamaev & Zubchenko, 1984)
halargyrea Rubec & Dronen, 1994
macruri (Brinkmann, 1942)
nezumiae (Munroe, Campbell & Zwerner, 1981)
- Mazocraeoides*
georgei Price, 1936
- Mazocraes*
alosae (Hermann, 1782)
harengi (van Beneden & Hesse, 1863)

- pilchardi* (van Beneden & Hesse, 1863)
- Megalocotyle*
hexacantha (Parona & Perugia, 1889)
- rhombi* (van Beneden & Hesse, 1863)
- zschorkei* (Mola, 1912)
- Megalocotyloides*
grandiloba (Paperna & Kohn, 1964)
- Merizocotyle*
diaphana Cefontaine, 1894
- Metamicrocotyla*
cephalus (Azim, 1939)
- Microbothrium*
apiculatum Olsson, 1869
- centrophori* Brinkmann, 1940
- Microcotyle*
acanthurum Parona & Perugia, 1896
- canthari* van Beneden & Hesse, 1863
- caudata* Goto, 1894
- centrodonti* Brown, 1929
- cephalus* Azim, 1939
- chrysophrii* van Beneden & Hesse, 1863
- donavini* van Beneden & Hesse, 1863
- draconis* Briot, 1904
- erythrinii* van Beneden & Hesse, 1863
- fusiformis* Goto, 1894
- lichiae* Ariola, 1899
- mormyri* Lorenz, 1878
- mugilis* Vogt, 1878
- pancerii* Sonsino, 1891
- pomatomi* Goto, 1899
- pontica* Pogorel'tseva, 1964
- sebastis* Goto, 1894
- Monocotyle*
ijimae Goto, 1894
- myliobatis* Taschenberg, 1878
- Mycteronastes*
icopae (Beverley-Burton & Williams, 1989)
- undulatae* Kearn & Beverley-Burton, 1990
- Nasicola*
klawei (Stunkard, 1962)
- Neohexostoma*
extensicaudatum (Dawes, 1940)
- thunninae* (Parona & Perugia, 1889)
- Neonchocotyle*
pastinacae Ktari & Maillard, 1972
- Nitzschia*
elongata (Nitzsch, 1826)
- sturionis* (Abildgaard, 1794)
- Octodactylus*
minus (Olsson, 1876)
- morrhuiae* (van Beneden & Hesse, 1863)
- Octolabea*
turchinii Euzet & Trilles, 1961
- Onchocotyle*
borealis van Beneden, 1853
- Ophicotyle*
fintae van Beneden & Hesse, 1863
- Paracyclocotyla*
cherbonnieri Dollfus, 1970
- Paradiplectanotrema*
trachuri (Kovaleva, 1970)
- Paradiplozoon*
bliccae (Reichenbach-Klinke, 1961)
- rutili* (Gläser, 1967)
- Plectanocotyle*
gurnardi (van Beneden & Hesse, 1863)
- Plectanocotyloides*
obscurum Euzet & Suriano, 1974
- Polyciphora*
nezumiae Lambert & Euzet, 1980
- Polyclithrum*
gerasevi Gaevskaya & Dimitrieva, 1997
- ponticum* Dimitrieva *et al.*
- rodgersi* Dimitrieva *et al.*
- Polylabris*
- diplodi* Euzet & Cauwet, 1967
- tubicirrus* (Paperna & Kohn, 1964)
- Protocotyle*
grisea (Cerfontaine, 1899)
- taschenbergi* Maillard & Oliver, 1966
- Protolamellodiscus*
raibauti Oliver & Radujkovic, 1987
- serranelli* (Euzet & Oliver, 1965)
- Protomicrocotyle*
ivoirensis Wahl, 1972
- Pseudanthocotyloides*
heterocotyle (van Beneden, 1871)
- Pseudaxine*
bivaginalis Dillon & Hargis, 1965
- trachuri* Parona & Perugia, 1889
- Pseudoanthocotyle*
markewitschii Nikolaeva & Pogorel'tseva, 1965
- Pseudocotyle*
squatinae van Beneden & Hesse, 1865
- lepidorhini* Guiart, 1938
- Pseudodactylogyrus*
anguillae (Yin & Sproston, 1948)
- Pseudodiplectanum*
gibsoni Oliver, 1987
- kearni* Vala, Lopez-Roman & Boudaoud, 1980
- Pseudokuhnia*
minor (Goto, 1894)
- Pseudomazocraes*
monsivaisae Caballero & Bravo Hollis, 1955
- Pyragraphorus*
hollisiae Euzet & Ktari, 1970
- Rajonchocotyle*
batis Cerfontaine, 1899
- alba* Cerfontaine, 1899
- emarginata* (Olsson, 1876)
- miraletus* Rees & Llewellyn, 1941
- prenanti* (St. Remy, 1890)
- pristium* Maillard, 1970
- Rhinecotyle*
crepitacula Euzet & Trilles, 1960
- Sciaenocotyle*
panceri (Sonsino, 1891)
- Serranicotyle*
labracis (van Beneden & Hesse, 1863)
- Sibitrema*
poonui Yamaguti, 1966
- Solostamenides*
mugilis (Vogt, 1879)
- Sparicotide*
chrysophrii (van Beneden & Hesse, 1863)
- Squalonchocotyle*
centrophori Maillard, 1970
- pontica* Pogorel'tseva, 1964
- spinaci* (Goto, 1894)
- squali* MacCallum, 1931
- tropai* Teindeiro & Valdez, 1955
- Tetraonchoides*
paradoxus Bychowsky, 1951
- Tetraonchus*
monenteron Diesing, 1858
- Thaumatocotyle*
concinna Scott, 1904
- Tricotyla*
cutanea Guiart, 1938
- molae* (Blanchard, 1847)
- Trimusculotrema*
leucanthemum (Euzet & Maillard, 1967)
- micracantha* (Euzet & Maillard, 1967)
- Tristoma*
coccineum (Cuvier, 1817)
- integrum* (Diesing, 1850)
- perugiae* (Setti, 1898)
- Tristomella*
grimaldii Guiart, 1938
- interrupta* (Monticelli, 1891)

- nozawae* (Goto, 1894)
onchidiocotyle (Setti, 1899)
Trochopella
candida Euzet & Trilles, 1962
Trochopus
brauni Mola, 1912
differens Sonsino, 1891
diplocaelus Massa, 1903
gallimhe Little, 1929
heteracanthus Massa, 1903
lineatus Scott, 1901
micracanthus Massa, 1903
oncacanthus Massa, 1906
pini (van Beneden & Hesse, 1863)
tubiporus (Diesing, 1836)
Udonella
caligorum Johnston, 1835
lupi van Beneden & Hesse, 1863
Vallisia
striata Parona & Perugia, 1890
Winkenthughesia
australis Robinson, 1961
bramae Parona & Perugia, 1896
- granulum* Rudolphi, 1819
Aploparaksis
larina Fuhrmann, 1921
Aporhynchus
norvegicus (Olsson, 1868)
Armadoskrjabinia
medici (Stossich, 1890)
parvuncinata (Meggitt, 1927)
Bothrimonus
fallax Lühe, 1900
nylandicus Schneider, 1902
Bothriocephalus
alessandrinii Condorelli-Francaviglia, 1898
andresi Porta, 1911
angustatus Rudolphi, 1819
atherinae Chernyshenko, 1949
auriculatus Rudolphi, 1819
barbatus Renaud, Gabrion & Pasteur, 1983
bramae Ariola, 1899
centrolophopomphi (Wagener, 1854)
clavibothrium Ariola, 1899
claviceps (Goeze, 1782)
coronatus Rudolphi, 1819
funiculus Renaud & Gabrion, 1984
gregarius Renaud, Gabrion & Pasteur, 1983
lophii Rudolphi, 1819
macrophallus Linstow, 1905
microcephalus Rudolphi, 1819
minutus Ariola, 1896
monticelli Ariola, 1899
proboscideus (Batsch, 1786)
punctatus (Rudolphi, 1802)
scorpii (Mueller, 1776)
squali Ariola, 1900
trachypteri Ariola, 1896
trachypteriis Ariola, 1896
uncinatus Rudolphi, 1819
vallei Stossich, 1899
Calliobothrium
coronatum (Rudolphi, 1819)
eschrichtii (van Beneden, 1849)
leuckarti van Beneden, 1850
lintoni Euzet, 1954
verticillatum (Rudolphi, 1819)
Callitetrarhynchus
gracilis (Rudolphi, 1819)
Caryophyllaeus
trisignatus Molin, 1858
Ceratobothrium
xanthocephalum Monticelli, 1892
Chimaerarhynchus
rougetae Beveridge & Campbell, 1989
Chimaerocestos
prudhoei Williams & Bray, 1984
Christianella
minuta (van Beneden, 1849)
Clestobothrium
crassiceps (Rudolphi, 1819)
Cloacotaenia
megalops (Nitsch in Creplin, 1829)
Colymbilepis
multistriata (Rudolphi, 1810)
Confluaria
furcifera (Krabbe, 1869)
podicipina (Szymanski, 1905)
Crossobothrium
angustum (Linton, 1889)
dohrnii (Oerley, 1885)
Dasyrhynchus
giganteus (Diesing, 1850)
talismani Dollfus, 1935
Dendroterina
karajasicus (Kurashvili, 1957)
mahdiaensis (Joyeux, 1923)
Dibothriocephalus

Cestoda

Compiled by Rodney A. Bray

The list of cestode species was derived from the Host-Parasite Data-base and Catalogue maintained by the Parasitic Worms Division, Department of Zoology, The Natural History Museum. Some attempt has been made to restrict the list to valid taxa and to use that latest nomenclature.

The list was kindly commented upon by Dr B.B. Georgiev and Prof. L. Euzet.

Class Cestoda

- Abothrium*
gadi van Beneden, 1871
Acanthobothrium
benedeni (Lönnberg, 1889)
crassicolle Wedl, 1855
dujardinii van Beneden, 1849
filicolle (Zschokke, 1887)
floridensis Goldstein, 1964
musculosum (Baer, 1948)
ponticum Borcea, 1934
quadripartitum Williams, 1968
rajaebatis (Rudolphi, 1810)
tripartitum Williams, 1969
zschorkei Baer, 1948
Alcataenia
atlantiensis Hoberg, 1991
Amphicotyle
heteropleura (Diesing, 1850)
Amphilina
folacea (Rudolphi, 1819)
Archistrocephalus
microcephalus (Rudolphi, 1819)
Anomotaenia
dubininae (Spassky, 1968)
hydrochelidonis (Dubinina, 1953)
micracantha (Krabbe, 1869)
microrhyncha (Krabbe, 1869)
Anthobothrium
auriculatum (Rudolphi, 1819)
cornucopia van Beneden, 1850
parvum Stossich, 1895
variabile (Linton, 1889)
Anthocephalum
gracile (Wedl, 1855)
Anthocephalus
gracilis Rudolphi, 1819

- attenuatus* Guiant, 1935
Dicranotaenia
spasskii (Schigin, 1961)
Dictymetra
laevigata (Rudolphi, 1819)
Diesingium
lomentaceum (Diesing, 1850)
Dilepis
undula (Schrank, 1788)
Dinobothrium
plicatum (Linton, 1922)
septaria van Beneden, 1889
spinosum Baylis, 1950
Diorchis
inflata (Rudolphi, 1819)
Diphyllobothrium
cordatum (Leuckart, 1863)
dendriticum (Nitzsch, 1824)
ditremum (Creplin, 1825)
elegans (Krabbe, 1865)
fuhrmanni Hsu, 1935
lanceolatum (Krabbe, 1865)
latum (Linnaeus, 1758)
polyrugosum Delyamure & Skrjabin, 1966
poniticum Delyamure, 1971
stemmacephalum Cobbold, 1858
Diplocotyle
olrikii (Krabbe, 1874)
Diplogonoporus
balaenopterae (Lonnberg, 1892)
Diploposthe
laevis (Bloch, 1782)
Discobothisrum
sp.
Ditrichybothrium
macrocephalum Rees, 1959
Dubininolepis
rostellata (Abildgaard, 1790)
swiderskii (Gasowska, 1932)
Echeneibothrium
acanthinophyllum Rees, 1961
affine Olsson, 1867
dubium Beneden, 1858
elongatum Williams, 1966
fallax van Beneden, 1870
gracile Zschokke, 1889
julievansum Woodland, 1927
maculatum Woodland, 1927
minimum van Beneden, 1850
minutum Williams, 1966
tumidulum (Rudolphi, 1819)
variabile van Beneden, 1850
Echinobothrium
affine Diesing, 1863
brachysoma Pintner, 1889
clavatum Probert & Stobart, 1989
harfordi McVicar, 1976
musteli Pintner, 1889
typus van Beneden, 1849
Echinocotyle
druzniensis Jarecka, 1958
multiglandularis (Baczynska, 1914)
oweni (Moghe, 1933)
verschureni (Baer, 1959)
vajteki Koubek, 1982
Echinophallus
settii (Ariola, 1895)
wageneri (Monticelli, 1890)
Eubothrium
crassum (Bloch, 1779)
fragile (Rudolphi, 1802)
rugosum (Batsch, 1786)
Euterarhynchus
ruficollis (Eysenhardt, 1829)
Fimbriaria
fasciolaris (Pallas, 1781)
Fimbriariooides
intermedia (Fuhrmann, 1913)
Fistulicola
dalmatinus (Stossich, 1897)
plicatus (Rudolphi, 1819)
Flaminolepis
liguloides (Gervais, 1847)
Floriceps
saccatus Cuvier, 1817
Gastrolecithus
planus (Linton, 1922)
Gilquinia
squali (Fabricius, 1794)
Grillotia
acanthoscolex Rees, 1944
angeli Dollfus, 1969
erinaceus (Beneden, 1858)
heptanchi (Vaullegeard, 1899)
institata (Pintner, 1913)
scolecina (Rudolphi, 1819)
smarisgora (Wagener, 1854)
Grillotia (Paragrillotia)
epinepheli Scholz, Garippa & Scala, 1993
rowei Campbell, 1977
Gryporhynchus
cheilancistratus (Wedl, 1855)
pusillus Nordmann, 1832
Gymnorhynchus
gigas (Cuvier, 1817)
reptans Rudolphi, 1819
Gyrocotyle
confusa Van der Land & Dienske, 1968
urna (Wagener, 1852)
Gyrocotyloides
nybelini Fuhrmann, 1931
Hepatoxylon
attenuatus (Rudolphi, 1819)
megacephalum (Rudolphi, 1819)
squali Bosc, 1811
stenocephala (Guillet, 1935)
trichiuri Holten, 1802
Heteronybelinia
alloiotica Dollfus, 1960
cadenati Dollfus, 1960
estigmene Dollfus, 1960
eureia Dollfus, 1960
punctatissima Dollfus, 1960
rougetcampanae Dollfus, 1960
senegalensis Dollfus, 1960
yamagutii Dollfus, 1960
Kotorella
pronosoma (Stossich, 1901)
Lacistorhynchus
tenuis (van Beneden, 1858)
Lateriporus
clerci (Johnston, 1912)
teres (Krabbe, 1869)
Ligula
intestinalis (Linnaeus, 1758)
Microsomacanthus
abortiva (Linstow, 1904)
fausti (Tseng-Shen, 1932)
heterospinus Spassky & Jurpalova, 1964
microsphabina Spassky & Jurpalova, 1964
microsoma (Creplin, 1829)
Molicola
horridus (Goodsir, 1841)
Monorygma
? delphini (Gervais, 1870)
chlamydoselachi Lönning, 1898
grimaldii (Moniez, 1899)
hyperapolitica (Obersteiner, 1914)
perfectum (van Beneden, 1853)
Nybelinia

- africana* Dollfus, 1960
dakari Dollfus, 1960
edwinlintoni Dollfus, 1960
goreensis Dollfus, 1960
lingualis Cuvier, 1817
oodes Dollfus, 1960
strongyla Dollfus, 1960
Oncobothrium
ganfini Mola, 1934
lintoni Mola, 1934
pseudouncinatum de Beauchamp, 1905
uncinatum (Rudolphi, 1819)
Ophryocotyle
proteus Friis, 1870
Orygmatobothrium
musteli (van Beneden, 1850)
versatile Diesing, 1854
Otobothrium
cysticum (Mayer, 1842)
dipsacus Linton, 1897
pronosomum (Stossich, 1901)
Parabothriocephalus
macruri Campbell, Correia & Haedrich, 1982
Parabothrium
bulbiferum Nybelin, 1922
gadipollachii (Rudolphi, 1810)
Parachristianella
trygonis Dollfus, 1946
Paradilepis
scolecina (Rudolphi, 1819)
Paricterotaenia
dodecacantha (Krabbe, 1869)
inversa (Rudolphi, 1819)
porosa (Rudolphi, 1810)
ransomi (Linton, 1927)
sternina (Krabbe, 1869)
Parvitaenia
ardeolae Burt, 1940
Passerilepis
crenata (Goeze, 1782)
Pelichnibothrium
speciosum Monticelli, 1899
Philobrythos
 sp.
Phyllobothrium
britannicum Williams, 1968
centrurum Southwell
delphini (Bosc, 1802)
foliatum Linton, 1891
gracile Wedl, 1855
laciniatum (Linton, 1889)
lactuca (van Beneden, 1850)
minutum Williams, 1968
piriei Williams, 1968
riggi (Monticelli, 1893)
rotundum (Klaptocz, 1906)
sinuosiceps Williams, 1959
squali Yamaguti, 1952
thridax (van Beneden, 1849)
tumidum Linton, 1922
unilaterale Southwell, 1925
Pistana
 sp.
Platybothrium
auriculatum Yamaguti, 1952
baeri Euzet, 1952
Platyscolex
ciliata (Fuhrmann, 1931)
Plerocercoides
lonchophorus Guiart, 1935
Plicobothrium
globicephala Rausch & Margolis, 1969
Poecilancistrum
caryophyllum (Diesing, 1850)
Priapococephalus
grandis Nybelin, 1922
minor Nybelin, 1928
Prosobothrium
armigerum Cohn, 1902
Proteocephalus
exiguus La Rue, 1911
filicollis (Rudolphi, 1810)
longicollis (Zeder, 1800)
macrocephalus (Creplin, 1825)
percae (Müller, 1780)
tetrastromus (Rudolphi, 1810)
Pseudanomotaenia
larina (Krabbe, 1869)
Pseudanthobothrium
hanseni Baer, 1956
Pseudobothrium
grimaldii Guiart
Pseudogrillozia
perelica (Shuler, 1938)
Pterobothrium
fragile (Diesing, 1850)
senegalense Campbell & Beveridge, 1996
Ptychobothrium
belones (Dujardin, 1845)
Retinometra
skrjabini (Mathevossian, 1945)
Rhinebothrium
polombi Baer, 1948
Rhynchobothrium
adenoplusius Pintner, 1903
Schistocephalus
solidus (Müller, 1776)
Scolex
pleuronectis Müller, 1788
pleuronectis subsp. *bitocularis*
polymorphus Rudolphi, 1819
Sobolevianthus
gracilis (Zeder, 1803)
octacanthoides (Fuhrmann, 1906)
Sphaerobothrium
lubeti Euzet, 1959
Sphyrioccephalus
dollfusi Bussieras & Aldrin, 1968
tergestinus Pintner, 1913
viridis (Wagener, 1854)
Strobilocephalus
triangularis (Diesing, 1850)
Taenia
aurita Rudolphi, 1819
Tatria
biremis Kowalewski, 1904
decacantha Fuhrmann, 1913
Tentacularia
bicolor (Bartels in Nordmann, 1832)
coryphaenae Bosc, 1802
Tetrabothrius
affinis (Lönnberg, 1891)
cylindraceus (Rudolphi, 1819)
diplosoma Guiart, 1935
erostris (Lönnberg, 1889)
forsleri (Krefft, 1871)
immerinus (Abildung, 1790)
macrocephalus (Rudolphi, 1810)
norvegicum Olsson, 1868
perfidus Joyeux & Baer, 1934
phalacrocoracis Burt, 1977
wilsoni (Leiper & Atkinson, 1914)
Tetrahynchobothrium
setiense Dollfus, 1969
Tetraphynchus
discophorus Rudolphi, 1819
megabothrius Rudolphi, 1819
smaridum Pintner, 1893
tenuicollis Rudolphi, 1819
Triaenophorus

crassus Forel, 1868
nodulosus (Pallas, 1781)
Trigonocotyle
globicephala Baer, 1954
lintoni Guiart, 1935
monticellii (Linton, 1923)
Trilocularia
acanthiae vulgaris Olsson, 1867
gracilis Olsson, 1869
Tritaphros
retzii Lönnberg, 1889
Tylocephalum
sp.
Valipora
ardeolae (Singh, 1952)
mutabilis Linton, 1927
unilateralis (Rudolphi, 1819)
Variolepis
farciminoso (Goeze, 1782)
Wardium
cirrosa (Krabbe, 1869)
filamentoovatum Shigin, 1961
fusa (Krabbe, 1869)
spasskii Shigin, 1961
stellorae (Deblock, Biguet & Capron, 1960)

ROTIFERA

Compiled by Myles O'Reilly

Rotifers are minute aquatic organisms generally less than 1mm in length. They have an external cuticle which is often ornamented or sculptured and they are characterised by a conspicuous crown, or "wheel", of long cilia. From this they derive their common name of "wheel animalcules". The phylum comprises over 1500 species mostly from freshwater habitats though a number of species also occur in brackish and marine waters. The genera *Seison* and *Zelinkielia* are exclusively marine and live as commensals on leptostracan crustacea and holothurians respectively (see Holloday, 1949). Most species, however, have a free-living planktonic mode of life though benthic or periphytic forms also occur.

The list provided comprises 139 European species from marine or brackish waters. The present compilation is biased towards the coastal waters of the North East Atlantic and does not, as yet, represent the southern European Atlantic nor Mediterranean fauna. Inclusion of these would probably add a significant number of species.

Pontin (1978) reviewed British freshwater planktonic species, about 30 of which are also recorded from brackish waters. Berzins (1960) summarises the records of about 60 marine planktonic species from the North East Atlantic. Thane-Fenchel (1968a) discussed the ecology and distribution of 34 species of non-planktonic rotifers from brackish waters of Scandinavia. Thane-Fenchel (1968b) also provided a key to marine and brackish-water genera of the world, citing further references where descriptions and illustrations of individual species can be located. The marine interstitial fauna of the North Sea island of Sylt was studied by Tzschaschel (1979) who recorded 21 species, 13 of which were new science.

References

- Berzins, B. 1960. Zooplankton Sheets. Rotatoria I-VI. No.84, Synchaetidae, Genus *Synchaeta*, 7pp. *Fiches d'Identification du Zooplancton*. Conseil International pour l'Exploration de la Mer.
- Berzins, B. 1960. Zooplankton Sheets. Rotatoria I-VI. No.85, Trichocercidae, 3pp. *Fiches d'Identification du Zooplancton*. Conseil International pour l'Exploration de la Mer.
- Berzins, B. 1960. Zooplankton Sheets. Rotatoria I-VI. No.86, Brachionidae, 4pp. *Fiches d'Identification du Zooplancton*. Conseil International pour l'Exploration de la Mer.
- Berzins, B. 1960. Zooplankton Sheets. Rotatoria I-VI. No.87, Brachionidae (Contd.), 5pp. *Fiches d'Identification du Zooplancton*. Conseil International pour l'Exploration de la Mer.
- Berzins, B. 1960. Zooplankton Sheets. Rotatoria I-VI. No.88, Asplanchnidae and Synchaetidae, Genera *Ploesoma* and *Polyarthra*, 4pp. *Fiches d'Identification du Zooplancton*. Conseil International pour l'Exploration de la Mer.
- Berzins, B. 1960. Zooplankton Sheets. Rotatoria I-VI. No.89, Testudinellidae, Conochilidae, and Collothecidae, 4pp. *Fiches d'Identification du Zooplancton*. Conseil International pour l'Exploration de la Mer.
- Holloday, E.D., 1949. A preliminary report on the Plymouth marine and brackish-water Rotifera. *Journal of the Marine Biological Association of the United Kingdom*. 28: 239-253.
- Pontin, R.M. 1978. *A key to British Freshwater Planktonic Rotifera*. Freshwater Biological Association. Scientific Publication No.38, 178pp.

Thane-Fenchel, A. 1968a. Distribution and Ecology of non-planktonic brackish-water rotifers from Scandinavian waters. *Ophelia* 5:273-297.

Thane-Fenchel, A. 1968b. A simple key to the genera of marine and brackish-water rotifers. *Ophelia* 5:299-311.

Tzschaschel, G. 1979. Marine Rotatoria aus dem Interstitial der Nordseeinsel Sylt. *Mikrofauna des Meeresbodens* 71:1-64.

PHYLUM ROTIFERA

Order Monogononta

Suborder Ploima

Family Brachionidae

Argonotholca Gillard, 1948
foliacea (Ehrenberg, 1838)

Brachionis Pallas, 1766

angularis Gosse, 1851
calyciflorus Pallas, 1766
leydigi Cohn, 1862
plicatilis Müller, 1786
quadridentatus Hermann, 1783
rubens Ehrenberg, 1838
urceolaris Müller, 1773

Colerella Bory de St. Vincent, 1824
colurus (Ehrenberg, 1830)

dicentra (Gosse, 1887)

Euchlanis Ehrenberg, 1832

dilitata Ehrenberg, 1832

Kelicottia Ahlstrom, 1938

longispina (Kelicot, 1879)

Keratella Bory de St. Vincent, 1822
cochlearis (Gosse, 1851)

cruciformis (Thompson, 1892)

testudo Ehrenberg, 1838

quadraata (Müller, 1786)

Lepadella Bory de St. Vincent, 1826

psammophila (Tzschaschel, 1979)

Mytilina

videns Levander

Notholca Gosse, 1886

acuminata (Ehrenberg, 1832)

labis Gosse, 1887

psammarina Buchholz & Ruhmann

striata (Müller, 1786)

Pseudonotholca Marakawa 1928

japonica Marakawa 1928

Trichotria

pocillum (Müller)

Tripleuchlanis Myers 1930

plicata (Levander, 1894)

Family Lecanidae

Lecane

cornuta (Müller)

grandis (Murray)

Proales Gosse, 1886

germanica Tzschaschel, 1979.

halophila Remane, 1929

longipes Remane,

oculata Tzschaschel, 1979

paguri Thane-Fenchel, 1966

reinhardti (Ehrenberg, 1833)

syltensis Tzschaschel, 1979

Family Notommatidae

Cephalodella

gibba (Ehrenberg)

Family Lindiidae

Lindia Dujardin, 1841

tecsua Harring & Myers, 1922

Family Trichocercidae

Trichocerca Lamarck, 1801

capucina Wierzejski & Zacharias, 1893

marina Daday, 1890

pusilla (Jennings, 1903)

Family Gastropodidae

Ascomorpha Perty, 1850

ecaudatus Perty, 1850

ovalis Carlin, 1943

Family Dicranophoridae

Aspelta

baltica Remane, 1929

clydona Harring & Myers, 1927

harrangi Remane

reibischii (Remane, 1949)

Encentrum Ehrenberg, 1838

ambiguum Tzschaschel, 1979

axi Tzschaschel, 1979

crenatum Harring & Myers, 1927

eulitorale Tzschaschel, 1979

kosteii Tzschaschel, 1979

limicola Otto, 1937

listense Tzschaschel, 1979

longirostrum Tzschaschel, 1979

marinum (Dujardin, 1841)

matthesi Remane, 1949

obesum Tzschaschel, 1979

permundum Tzschaschel, 1979

psammophilum Tzschaschel, 1979

rousseleti Lie Petersen

sacculiforme Tzschaschel, 1979

simillimum Remane, 1929

striatum Althaus, 1957

villosum Harring & Myers, 1927

wierzejskii (Remane, 1949)

=*Wierzejskella marina* Remane, 1949

Paradicranophorus

hudsoni (Glascott)

Family Asplanchnidae

Asplanchna Gosse, 1850

brightwelli Gosse, 1850

priodata Gosse, 1850

Family Synchaetidae

Ploesoma Herrick 1885

hudsoni (Imhof, 1891)

truncatum (Levander, 1894)

Polyarthra Ehrenberg, 1834

dolichoptera Idelson, 1925

remata Skorikov, 1896

vulgaris Carlin, 1943

Synchaeta Ehrenberg, 1832

atlantica Zelinka, 1907

baccilifera Smirnov, 1933

baltica Ehrenberg, 1838

bicornis Smith, 1904

cecilia Rousselet, 1902

curvata Lie-Petersen, 1905

fennica Rousselet, 1909

glacialis Smirnov, 1932

grimpei Remane, 1929

gyrina Hood, 1887

hyperborea Smirnov, 1932

johanseni Harring, 1921

littoralis Rousselet, 1902

longipes Gosse, 1887

monopus Plate, 1889

neapolitana Rousselet, 1902

pectinata Ehrenberg, 1838

stylata Wierzejski, 1893

tamara Smirnov, 1932

tavina Hood, 1893

tremula Muller, 1786

triophthalma Lauterborn, 1894

vorax Rousselet, 1902

Suborder Flosculariaceae

Family Testudinellidae

Filinia Bory de St. Vincent, 1824

brachiata (Rousselet, 1901)

cornuta (Weisse, 1847)

longiseta (Ehrenberg, 1884)

terminalis (Plate, 1886)
Hexarthra Schmarda, 1854
fennica (Levander, 1892)
mira (Hudson, 1871)
Pompholyx Gosse, 1851
sulcata Hudson, 1855
Testudinella Bory de St. Vincent, 1826
clypeata (Müller, 1786)
patina (Hermann, 1783)

Family Conochiliidae
Conochilus Ehrenberg, 1834
hippocrepis (Schrank, 1830)
unicornis (Rousselet, 1892)

Family Philodinidae
Rotaria
citrina (Ehrenberg)

Suborder Collotheceae

Family Collotheceidae
Collotheca Harring, 1913
mutabilis (Hudson, 1885)
pelagica (Rousselet, 1893)

Order Bdelloidea

Family Philodinidae
Zelinkiella Harring, 1913
synaptae (Lankester, 1868)

Order Seisonidea

Family Seisonidae
Seison Grube, 1859
annulatus Claus, 1876
nebaliae Grube, 1859

GNATHOSTOMULIDA

Compiled by Jacob van der Land

The European list was derived from URMO mainly based on Sterrer (1982) with additions from Zoological Record.

References

Sterrer, W.E., 1982. Gnathostomulida. In: S.P. Parker (ed.), *Synopsis and Classification of Living Organisms*, vol. 1: 847-851. McGraw-Hill, New York, etc.

PHYLUM GNATHOSTOMULIDA

Order Bursovaginoidea

Family Agnathiellidae

Paragnathiella
trifoliceps Sterrer, 1997

A

Family Austrognathiidae

Austrognatharia
boadeni Sterrer, 1969
Austrognathia
clavigera Sterrer, 1997
riedli Sterrer, 1965
Triplignatha
adriatica Sterrer, 1991

A

M

M

Family Gnathostomariidae

Gnathostomaria
lutheri Ax, 1956

A M

Family Gnathostomulidae

Gnathostomula
axi Kirsteuer, 1964
mediterranea Sterrer, 1970
murmanica Mamkaeva, 1962
paradoxa Ax, 1956

M

M

A

A M

Family Mesognathiidae

Labidognathia
longicollis Riedl, 1970
Mesognatharia
remanei Sterrer, 1966

A

A M

Family Onychognathiidae

Valvognathia
pogonostoma Kristensen & Nørrevang, 1978

A

Family Rastrognathiidae

Rastrognathia
macrostoma Kristensen & Nørrevang, 1978

A

Order Filospermoidea

Family Haplognathiidae

Haplognathia
filum (Sterrer, 1966)
gubbarorum (Sterrer, 1969)
lunulifera (Sterrer, 1969)
rosea (Sterrer, 1969)
ruberrima (Sterrer, 1966)
rubromaculata (Sterrer, 1969)
simplex (Sterrer, 1966)

A

A

A

A

A M

A

A

Family Pterognathiidae

Pterognathia
atrox Sterrer, 1969
meixneri Sterrer, 1969
sorex Sterrer, 1969
swedmarki Sterrer, 1966

A

A

A

A

A

NEMERTINI

Compiled by Ray Gibson

The higher classification used in the checklist of European nemertean species and genera follows the proposal given by Sundberg (1991). This was modified to include Sundberg & Hylbom's (1994) conclusion, based on their phylogenetic study of the anopla subclasses Archi- and Palaeonemertea, that the retention of the Archinemertea as a separate taxon could no longer be supported. Nemertean genera listed as archinemerteans in Gibson's (1995) checklist of the nemertean genera and species of the world are included below in the Palaeonemertea. Since 1995 several species of nemerteans have been subjected to taxonomic revision and a number of new genera and species have been established. The present checklist has to a large extent been compiled from information given by Gibson (1994, 1995), but is supplemented by the work of Berg & Gibson (1996), Cantell (1996, 1998), Chernyshev (1995), Envall (1996), Frutos *et al.* (1998) and Senz (1995, 1996a, b, 1997).

References

- Berg, G. & Gibson, R. 1996. A redescription of *Nemertoscolex parasiticus* Greeff, 1879, an apparently endoparasitic heteronemertean from the coelomic fluid of the echiurid *Echiurus echiurus* (Pallas). *Journal of Natural History* 30: 163-173.
- Cantell, C.-E. 1996. Redescription of *Leucocephalonemertes aurantiaca* (Grube, 1855) (Nemertini) n. gen. from the Mediterranean. *Ophelia* 45: 115-125.
- Cantell, C.-E. 1998. The anatomy and taxonomy of *Amorphonemertes kubergensis* gen. et sp. nov. (Nemertini) from the Tromsø area (Norway). *Sarsia* 83: 293-300.
- Chernyshev, A. V. 1995. O vuissikh taksonakh nemertin s obzorom sistemui podklassa Anopla. *Zoologicheskii zhurnal* 74: 7-18.
- Envall, M. 1996. *Ototyphlonemertes correiae* sp. nov. and a redescription of *O. duplex* (Nemertea: Monostilifera: Ototyphlonemertidae), with a phylogenetic consideration of the genus. *Journal of Zoology, London* 238: 253-277.
- Frutos, I., Montalvo, S. & Junoy, J. 1998. A new species of *Prosorhochmus* (Hoplonemertea, Monostilifera) from the Chafarinas Islands (western Mediterranean). *Journal of Zoology, London* 245: 293-298.
- Gibson, R. 1994. *Nemerteans*. Shrewsbury, Field Studies Council, 224 pp. (No. 24 Second Edition in *Synopses of the British Fauna (New Series)* eds R. S. K. Barnes & J. H. Crothers).
- Gibson, R. 1995. Nemertean genera and species of the world: an annotated checklist of original names and description citations, synonyms, current taxonomic status, habitats and recorded zoogeographic distribution. *Journal of Natural History* 29: 271-561.
- Senz, W. 1995. *Lineus rovinjensis* nom. n. (olim *Lineus areniculus* Senz, 1993) and *Tetrastemma kirsteueri* nom. n. (olim *Tetrastemma cruciatum* Senz, 1993) (Nemertinea: Heteronemertinea, Hoplonemertinea). *Annalen des Naturhistorischen Museums in Wien, Serie B Botanik und Zoologie* 97: 55-56.
- Senz, W. 1996a. Eine Untersuchung der Probleme der Heteronemertinen-Systematik, dargestellt anhand von *Lineus bergendali* n. sp., einer neuen heteronemertine aus dem Golf von Triest (Nemertini: Anopla: Lineidae). *Senckenbergiana Biologica* 76: 197-3208.
- Senz, W. 1996b. On the genus *Minutanemertes* Senz, 1993 (Nemertini: Hoplonemertini). *Annalen des Naturhistorischen Museums in Wien, Serie B Botanik und Zoologie* 98: 45-55.
- Senz, W. 1997. Morphologie und klassifikatorische Position einiger anopler Nemertinen (Nemertini: Anopla). *Annalen des Naturhistorischen Museums in Wien, Serie B Botanik und Zoologie* 99: 423-496.
- Sundberg, P. 1991. A proposal for renaming the higher taxonomic categories in the phylum Nemertea. *Journal of Natural History* 25: 45-48.
- Sundberg, P. & Hylbom, R. 1994. Phylogeny of the nemertean subclass Palaeonemertea (Anopla, Nemertea). *Cladistics* 10: 347-402.

PHYLUM NEMERTINI

Class Anopla

Subclass Palaeonemertini

Family Carinomidae

<i>Carinoma armandi</i> (McIntosh, 1875)	A
--	---

Family Cephalothricidae

<i>Cephalothrix arenaria</i> Hylbom, 1957	A
<i>atlantica</i> Gerner, 1969	A
<i>bipunctata</i> Bürger, 1892	M
<i>buergeri</i> Wijnhoff, 1913	M
<i>germanica</i> Gerner, 1969	A
<i>lactea</i> Senz, 1993	M
<i>linearis</i> (Rathke, 1799)	A M
<i>mediterranea</i> Gerner, 1969	M
<i>paragermanica</i> Senz, 1993	M
<i>rufifrons</i> (Johnston, 1837)	A M
<i>Cephalotrichella signata</i> (Hubrecht, 1879)	M
<i>Procephalothrix adriatica</i> Senz, 1993	M
<i>filiformis</i> (Johnston, 1828)	A
<i>kiliensis</i> Friedrich, 1935	A
<i>oestrymnica</i> Junoy & Gibson, 1991	A

Family Hubrechtidae

<i>Hubrechtella atypica</i> Senz, 1992	A
<i>combinata</i> Senz, 1993	M
<i>dubia</i> Bergendal, 1902	A
<i>globocystica</i> Senz, 1993	M
<i>Hubrechtia desiderata</i> (Kennel, 1891)	M

Family Tubulanidae

<i>Callinera buergeri</i> Bergendal, 1900	A
<i>grandis</i> Bergendal, 1903	A
<i>monensis</i> Rogers, Gibson & Thorpe, 1992	A
<i>Carinesta anglica</i> Wijnhoff, 1912	A
<i>Carinina arenaria</i> Hylbom, 1957	A
<i>atavia</i> (Bergendal, 1902)	A
<i>buddenbrocki</i> (Friedrich, 1935)	A
<i>burgeri</i> Joubin, 1902	A
<i>coei</i> Hylbom, 1957	A
<i>heterosoma</i> Müller, 1965	M
<i>poseidonis</i> Friedrich, 1935	A
<i>remanei</i> (Nawitzki, 1931)	A
<i>wijnhoffae</i> Kulikova, 1984	A
<i>Protubulanus theeli</i> (Bergendal, 1902)	A
<i>Tubulanus albocapitatus</i> Wijnhoff, 1912	A
<i>ambiguus</i> (Punnett, 1903)	A
<i>annulatus</i> (Montagu, 1804)	A M
<i>banyulensis</i> (Joubin, 1890)	A M
<i>borealis</i> Friedrich, 1936	A
<i>groenlandicus</i> (Bergendal, 1902)	A
<i>holorhynchocoelomicus</i> Friedrich, 1958	A
<i>inexpectatus</i> (Hubrecht, 1880)	A M
<i>linearis</i> (McIntosh, 1873/74)	A M
<i>miniatus</i> (Bürger, 1892)	A M
<i>norvegicus</i> Senz, 1993	A

<i>nothus</i> (Bürger, 1892)	A M	<i>immaculata</i> (Bürger, 1892)	M
<i>panormitanus</i> Monastero, 1930	M	<i>obscura</i> (Friedrich, 1958)	A
<i>polymorphus</i> Renier, 1804	A	<i>thori</i> Friedrich, 1958	A
<i>rubicundus</i> (Bürger, 1892)	M	<i>Lineopsella</i>	
<i>superbus</i> (Kölliker, 1845)	A M	<i>islandicus</i> (Friedrich, 1958)	A
<i>tubicola</i> (Kennel, 1891)	M	<i>Lineus</i>	
Subclass Heteronemertini			
Family Cerebratulidae			
<i>Amorphonemertes</i>		<i>acutifrons</i> Southern, 1913	A
<i>kubergensis</i> Cantell, 1998	A	<i>alienus</i> Bürger, 1895	A M
<i>Cerebratulus</i>		<i>bergendali</i> Senz, 1996	A
<i>acus</i> Nardo, 1847	M	<i>bilineatus</i> (Renier, 1804)	A M
<i>aerugatus</i> Bürger, 1892	M	<i>cinereus</i> Punnett, 1903	A
<i>alleni</i> Wijnhoff, 1912	A	<i>coccinus</i> Bürger, 1892	A M
<i>anguillula</i> Bürger, 1892	M	<i>dohrnii</i> (Hubrecht, 1879)	M
<i>annellatus</i> (Leuckart, 1849)	A	<i>fischeri</i> Senz, 1997	A
<i>aureolus</i> Bürger, 1892	M	<i>frauenseldi</i> Senz, 1997	A
<i>barentsi</i> Bürger, 1895	A	<i>gilbus</i> Bürger, 1892	M
<i>bivittatus</i> Ulyanina, 1870	M	<i>grubei</i> (Hubrecht, 1879)	M
<i>borealis</i> (Diesing, 1862)	A	<i>gurjanovae</i> Korotkevich, 1977	A
<i>brevis</i> Ushakov, 1926	A	<i>hubrechti</i> (Langerhans, 1880)	A M
<i>cestoides</i> Bürger, 1895	M	<i>insignis</i> Senz, 1993	M
<i>complanatus</i> (Kölliker, 1845)	M	<i>iota</i> Joubin, 1902	M
<i>croceus</i> Grube, 1864	M	<i>islandicus</i> Friedrich, 1958	A
<i>depressus</i> Quatrefages, 1846	M	<i>kennelii</i> Bürger, 1892	M
<i>eisigii</i> Hubrecht, 1880	M	<i>kolaensis</i> Ushakov, 1928	A
<i>epsilon</i> Joubin, 1902	A	<i>kristinebergensis</i> Gering, 1912	A
<i>erythrorochma</i> Joubin, 1902	A	<i>lobianki</i> Bürger, 1892	M
<i>ferrugineus</i> Bürger, 1892	M	<i>longissimus</i> (Gunnerus, 1770)	A
<i>fissuralis</i> Friedrich, 1958	A	<i>marisalbi</i> Ushakov, 1926	A
<i>flavifrons</i> Grube, 1864	M	<i>mcintoshii</i> (Langerhans, 1880)	A
<i>fuscoides</i> Bürger, 1892	M	<i>molochinus</i> Bürger, 1892	M
<i>fuscus</i> (McIntosh, 1873/74)	A M	<i>nigrobrunneus</i> Bergendal, 1903	A
<i>gamma</i> Joubin, 1902	A	<i>parvulus</i> Bürger, 1892	M
<i>greenlandicus</i> Punnett, 1901	A	<i>pseudoruber</i> (Friedrich, 1935)	A
<i>hepaticus</i> Hubrecht, 1879	A M	<i>rovinjensis</i> Senz, 1995	M
<i>joubini</i> Bürger, 1895	M	<i>ruber</i> (Müller, 1774)	A M
<i>liguricus</i> Blanchard, 1849	M	<i>rufocaudatus</i> Bürger, 1892	M
<i>lividus</i> Bürger, 1892	M	<i>sainthilairi</i> Ushakov, 1926	A
<i>marginatus</i> Renier, 1804	A	<i>scandinaviensis</i> Punnett, 1903	A
<i>melanorhynchus</i> Bürger, 1895	M	<i>uschakovi</i> Korotkevich, 1977	A
<i>modestus</i> Chapuis, 1886	A	<i>variegatus</i> Chapuis, 1886	A
<i>niveus</i> (Punnett, 1903)	A	<i>versicolor</i> Bürger, 1892	M
<i>norvegicus</i> Punnett, 1903	A	<i>viridis</i> (Müller, 1774)	A
<i>notabilis</i> Bürger, 1892	M	<i>Micrella</i>	
<i>pantherinus</i> Hubrecht, 1879	A M	<i>rufa</i> Punnett, 1901	A
<i>rigidus</i> Isler, 1900	A	<i>Micrura</i>	
<i>roseus</i> (Delle Chiaje, 1841)	A M	<i>albida</i> Verrill, 1879	A
<i>simulans</i> Bürger, 1892	M	<i>albifrons</i> Timofeeva, 1912	A
<i>urticans</i> (Müller, 1854)	M	<i>ambigua</i> Friedrich, 1958	A
<i>ventriporis</i> Friedrich, 1958	A	<i>atra</i> Punnett, 1903	A
<i>ventrosulcatus</i> Bürger, 1892	M	<i>baltica</i> Cantell, 1975	A
<i>zachsi</i> Ushakov, 1926	A	<i>bergenicola</i> Punnett, 1903	A
<i>Leucocephalonemertes</i>		<i>candida</i> Bürger, 1892	A M
<i>aurantiaca</i> (Grube, 1855)	A M	<i>corallifila</i> Cantell, 1975	A
<i>Micrurides</i>		<i>dellechiaiei</i> (Hubrecht, 1879)	M
<i>albopunctatus</i> Cantell, 1988	A	<i>elegans</i> Senz, 1993	A
<i>islandicus</i> Friedrich, 1960	A	<i>fasciolata</i> Ehrenberg, 1828	A
<i>Oxypolella</i>		<i>filaris</i> (Müller, 1788)	A
<i>alba</i> Bergendal, 1903	A	<i>lithotamni</i> Ushakov, 1928	A
<i>bergendali</i> Cantell, 1972	A	<i>pardalis</i> Haddon, 1886	A
<i>histriana</i> Senz, 1993	M	<i>pseudovaricolor</i> Senz, 1993	A
<i>punnetti</i> Bergendal, 1902	A	<i>purpurea</i> (Dalyell, 1853)	A M
<i>Oxypolia</i>		<i>rockalliensis</i> Dollfus, 1924	A
<i>beaumontiana</i> Punnett, 1901	A	<i>rovinjensis</i> Senz, 1993	M
<i>Paracerebratulus</i>		<i>scotica</i> Stephenson, 1911	A
<i>adriaticus</i> Senz, 1997	M	<i>tristis</i> (Hubrecht, 1879)	M
<i>Valencinura</i>		<i>varicolor</i> Punnett, 1903	A
<i>bahusiensis</i> Bergendal, 1902	A	<i>Micrurinella</i>	
Family Lineidae		<i>antondohrni</i> Friedrich, 1960	A
<i>Aetheolineus</i>		<i>Nemertoscolex</i>	
<i>pulcherrimus</i> Senz, 1993	M	<i>parasiticus</i> Greeff, 1879	A
<i>Euborlasia</i>		<i>Paralineus</i>	
<i>elizabethae</i> (McIntosh, 1873/74)	A M	<i>elisabethae</i> Schütz, 1911	M
		<i>Pontolineus</i>	
		<i>arenarius</i> Müller & Scripcariu, 1964	M
		<i>Pussylineus</i>	

<i>gabriellae</i> Corrêa, 1956	M	<i>pellucidus</i> (Ørsted, 1843)	A
<i>Riseriellus</i>		<i>polyommatus</i> Bürger, 1895	M
<i>occultus</i> Rogers, Junoy, Gibson & Thorpe, 1993	A	<i>pugnax</i> Hubrecht, 1879	M
<i>Tarrhomynos</i>		<i>punnetti</i> Friedrich, 1957	A
<i>praealbescens</i> (Cantell, 1982)	A	<i>reticulatus</i> Bürger, 1895	M
<i>Tenuilineus</i>		<i>roseus</i> (Müller, 1774)	A
<i>albocinctus</i> (Bergendal, 1903)	A	<i>rufostriatus</i> Bergendal, 1903	A
Family Mixolineidae		<i>septentrionalis</i> Friedrich, 1957	A
<i>Mixolineus</i>		<i>sipunculus</i> Saint-Loup, 1886	M
<i>levitrontosus</i> Senz, 1992	A	<i>superbus</i> (Stimpson, 1854)	A
<i>tauricus</i> Müller & Scripcariu, 1971	M	<i>thompsoni</i> Punnett, 1901	A
Family Valenciniidae		<i>validissimus</i> Bürger, 1895	M
<i>Baseodiscus</i>		<i>virgatus</i> Bürger, 1895	M
<i>abyssorum</i> (Joubin, 1902)	A	<i>vitae</i> Monastero, 1930	M
<i>delineatus</i> (Delle Chiaje, 1825)	A	<i>Arctonemertes</i>	
<i>filholi</i> (Joubin, 1902)	M	<i>thori</i> Friedrich, 1957	A
<i>minor</i> (Hubrecht, 1879)	M	<i>Communoporus</i>	
<i>pellucidus</i> (Kennel, 1891)	M	<i>cephalonephridialis</i> (Friedrich, 1940)	A
<i>Poliopsis</i>		<i>hagmeieri</i> (Friedrich, 1940)	A
<i>lacazei</i> Joubin, 1890	AM	<i>rhynchocoelomicus</i> (Friedrich, 1940)	A
<i>Valencinia</i>		<i>Dananemertes</i>	
<i>blanca</i> Bürger, 1892	M	<i>saemundssonii</i> Friedrich, 1957	A
<i>longirostris</i> Quatrefages, 1846	AM	<i>Duosnemertes</i>	
Family incertae sedis		<i>marmorata</i> (Bürger, 1895)	AM
<i>Diplopleura</i>		<i>Gurjanovella</i>	
<i>formosa</i> (Hubrecht, 1879)	M	<i>litoralis</i> Ushakov, 1926	A
<i>Huilkalineus</i>		<i>murmanica</i> Ushakov, 1926	A
<i>inexpectatus</i> Senz, 1993	M	<i>Proneurotes</i>	
<i>Notospermus</i>		<i>baltica</i> Friedrich, 1940	A
<i>geniculatus</i> (Delle Chiaje, 1828)	AM	<i>Psammamphiaporus</i>	
<i>Pseudobaseodiscus</i>		<i>elongatus</i> (Stephenson, 1911)	A
<i>nonsulcatus</i> Senz, 1993	M	<i>Valdivianemertes</i>	
<i>Utolineides</i>		<i>stannii</i> (Grube, 1840)	M
<i>alba</i> Senz, 1997	M	<i>Zygonemertes</i>	
Class Enopla		<i>algensis</i> (Bürger, 1895)	M
Subclass Hoplonemertini		<i>maslovskyi</i> (Czerniavsky 1880)	M
Superorder Monostilifera		Family Carcinonemertidae	
Family Amphiporidae		<i>Carcinonemertes</i>	
<i>Amphiporella</i>		<i>carcinophila</i> (Kölliker, 1845)	A
<i>baltica</i> Friedrich, 1940	A	Family Cratenemertidae	
<i>Amphiporus</i>		<i>Nippornemertes</i>	
<i>adriaticus</i> (Ehrenberg, 1828)	M	<i>danae</i> (Friedrich, 1957)	A
<i>allucens</i> Bürger, 1895	AM	<i>magna</i> (Punnett, 1903)	A
<i>angulatus</i> (Müller, 1774)	A	<i>pulchra</i> (Johnston, 1837)	A
<i>appendiculatus</i> Friedrich, 1957	A	Family Embletonematidae	
<i>arcticus</i> Punnett, 1901	A	<i>Cryptonemertes</i>	
<i>atypicus</i> Friedrich, 1935	A	<i>actinophila</i> (Bürger, 1904)	A
<i>bicolor</i> Bürger, 1895	A	<i>Embletonema</i>	
<i>binocellatus</i> Senz, 1993	M	<i>bocki</i> Brunberg, 1957	A
<i>bioculatus</i> McIntosh, 1873/74	A	<i>derjugini</i> Ushakov, 1928	A
<i>canescens</i> (Leuckart, 1849)	A	<i>duoni</i> (Joubin, 1890)	A
<i>carinelloides</i> Bürger, 1895	M	<i>echinoderma</i> (Marion, 1873)	AM
<i>dissimulans</i> Riches, 1893	A	<i>gracile</i> (Johnston, 1837)	AM
<i>dubius</i> Hubrecht, 1879	M	<i>intestinalis</i> Friedrich, 1958	A
<i>fabricii</i> Levinsen, 1879	A	<i>marioni</i> (Hubrecht, 1879)	M
<i>flavens</i> Monastero, 1930	M	<i>nordgaardi</i> (Punnett, 1903)	A
<i>giardinae</i> Monastero, 1930	M	<i>spongicola</i> (Bergendal, 1903)	A
<i>glandulosus</i> Bürger, 1895	M	<i>Nemertopsis</i>	
<i>groenlandicus</i> Ørsted, 1843	A	<i>bivittata</i> (Delle Chiaje, 1841)	A
<i>grubei</i> (Diesing, 1850)	M	<i>capitulata</i> Timofeeva, 1912	AM
<i>hastatus</i> McIntosh, 1873/74	A	<i>flavida</i> (McIntosh, 1873/74)	AM
<i>islandicus</i> Friedrich, 1957	A	<i>Paranemertes</i>	
<i>juli</i> Giard, 1890	A	<i>neesii</i> (Ørsted, 1843)	AM
<i>korschelti</i> Friedrich, 1940	AM	<i>Tetranemertes</i>	
<i>lactifloreus</i> (Johnston, 1828)	M	<i>antonina</i> (Quatrefages, 1846)	M
<i>langiaegeginus</i> Bürger, 1895	AM	Family Ototyphlonemertidae	
<i>marmoratus</i> Hubrecht, 1879	A	<i>Ototyphlonemertes</i>	
<i>martyi</i> Oxner, 1907	M	<i>antipai</i> Müller, 1968	M
<i>multiculatus</i> (Kölliker, 1845)	A	<i>aurantiaca</i> (du Plessis, 1891)	M
<i>murmanicum</i> Ushakov, 1928	A	<i>aurita</i> (Ulyanina, 1870)	M
<i>nigrostriatus</i> Timofeeva, 1912	A	<i>brunnea</i> Bürger, 1895	AM
<i>oligommatus</i> Bürger, 1895	M	<i>claparedii</i> (du Plessis, 1891)	M
		<i>correae</i> Enwall, 1996	A
		<i>duplex</i> Bürger, 1895	M

<i>esulcata</i> Senz, 1993	M	<i>lophopheliae</i> Bergendal, 1903	A
<i>macintoshii</i> Bürger, 1895	AM	<i>marionis</i> Joubin, 1890	AM
<i>pallida</i> (Keferstein, 1862)	AM	<i>melanocephalum</i> (Johnston, 1837)	AM
Family Prosorhochmidae		<i>mixtum</i> (Timofeeva, 1912)	M
<i>Gononemertes</i>		<i>nimbatum</i> Bürger, 1895	M
<i>parasita</i> Bergendal, 1900	A	<i>peltatum</i> Bürger, 1895	AM
<i>Prosovorchmus</i>		<i>portus</i> Bürger, 1895	M
<i>adriatica</i> Senz, 1993	M	<i>quadristriatum</i> Langerhans, 1880	AM
<i>chafarinensis</i> Frutos, Montalvo & Junoy, 1998	M	<i>quatreagesi</i> (Bürger, 1904)	AM
<i>clapareddii</i> Keferstein, 1862	AM	<i>robertianae</i> McIntosh, 1873/74	A
<i>subterraneus</i> Friedrich, 1949	A	<i>scutelliferum</i> Bürger, 1895	M
Family Tetrastemmatidae		<i>simplex</i> (Timofeeva, 1912)	M
<i>Arenonemertes</i>		<i>subpellucidum</i> Ørsted, 1843	A
<i>arenicola</i> Hylbom, 1991	A	<i>timofeevai</i> Gibson, 1995	M
<i>microps</i> Friedrich, 1933	AM	<i>tutus</i> (Monastero, 1930)	M
<i>minuta</i> Friedrich, 1949	A	<i>unicolor</i> (Hubrecht, 1879)	AM
<i>Cyanophthalma</i>		<i>vastum</i> Bürger, 1895	M
<i>cordiceps</i> (Friedrich, 1933)	A	<i>vermiculus</i> (Quatrefages, 1846)	AM
<i>obscura</i> (Schultze, 1851)	AM	<i>viera</i> (Timofeeva, 1912)	M
<i>Nemertellina</i>		<i>viperula</i> (Timofeeva, 1912)	M
<i>canea</i> Friedrich, 1935	A	<i>virgatum</i> Kirsteuer, 1963	M
<i>minuta</i> Friedrich, 1935	A	<i>vittigerum</i> Bürger, 1904	AM
<i>oculata</i> Friedrich, 1935	A		
<i>Nemertellopsis</i>		Family incertae sedis	
<i>cephalotrichiformis</i> Friedrich, 1935	A	<i>Albanemertes</i>	
<i>macrodasys</i> Friedrich, 1935	A	<i>rovinjensis</i> Senz, 1993	M
<i>typica</i> Friedrich, 1936	A	<i>Annulonemertes</i>	
<i>Paraminutanemertes</i>		<i>minusculus</i> Berg, 1985	A
<i>minuta</i> (Friedrich, 1935)	A	<i>Atrionemertes</i>	
<i>Prostomatella</i>		<i>greenlandica</i> Senz, 1993	A
<i>arenicola</i> Friedrich, 1935	A	<i>Minutanemertes</i>	
<i>Prostomiopsis</i>		<i>advertisculata</i> Senz, 1993	M
<i>alba</i> Friedrich, 1936	A	<i>alba</i> Senz, 1993	M
<i>Sacconemertes</i>		<i>schiffkoi</i> Senz, 1993	M
<i>arenosa</i> Karl Ing, 1933	A	<i>Oerstedia</i>	
<i>Tetrastemma</i>		<i>crassa</i> (Senz, 1993)	A
<i>albolineatum</i> (Timofeeva, 1912)	M	<i>dorsalis</i> (Abildgaard, 1806)	A
<i>ambiguum</i> Riches, 1893	A	<i>immutabilis</i> (Riches, 1893)	A
<i>angulatus</i> Senz, 1993	A	<i>laminariae</i> Friedrich, 1936	A
<i>arcticum</i> Ushakov, 1926	A	<i>nigra</i> (Riches, 1893)	A
<i>aseptatum</i> (Friedrich, 1935)	A	<i>rustica</i> (Joubin, 1890)	AM
<i>assimile</i> Ørsted, 1843	A	<i>similiformis</i> (Friedrich, 1935)	A
<i>bacescui</i> Müller, 1962	M	<i>striata</i> Sundberg, 1988	A
<i>baculum</i> (Quatrefages, 1846)	M	<i>tenuicollis</i> (Kirsteuer, 1963)	M
<i>beaumonti</i> (Southern, 1913)	A	<i>wijnhoffae</i> Friedrich, 1935	A
<i>bioculatum</i> Ørsted, 1844	A		
<i>bipeltatum</i> (Timofeeva, 1912)	M	Superorder Polystilifera	
<i>bistriatum</i> (Timofeeva, 1912)	M	Order Reptantia	
<i>brunnea</i> (Friedrich, 1935)	A	Family Brinkmanniidae	
<i>buxeyum</i> Bürger, 1895	M	<i>Brinkmannia</i>	
<i>candidum</i> (Müller, 1774)	A	<i>mediterranea</i> Stiasny-Wijnhoff, 1926	M
<i>cassidens</i> (Marenzeller, 1886)	A		
<i>cephalophorum</i> Bürger, 1895	AM	Family Drepanogigantidae	
<i>cerasinum</i> Bürger, 1895	M	<i>Drepanogigas</i>	
<i>coronatum</i> (Quatrefages, 1846)	AM	<i>albolineatus</i> (Bürger, 1895)	M
<i>cruciatum</i> Bürger, 1895	M		
<i>diadema</i> Hubrecht, 1879	M	Family Drepanophoridae	
<i>dubium</i> Ørsted, 1845	A	<i>Drepanophoriella</i>	
<i>falsum</i> Bürger, 1895	M	<i>histriana</i> Senz, 1993	M
<i>flavidum</i> Ehrenberg, 1828	A M	<i>Drepanophorus</i>	
<i>fozenensis</i> Gibson & Junoy, 1991	A	<i>massiliensis</i> Joubin, 1894	M
<i>fulvum</i> Kirsteuer, 1963	M	<i>rubrostriatus</i> Hubrecht, 1874	M
<i>glanduliferum</i> Bürger, 1895	M	<i>Punnertia</i>	
<i>graefei</i> (Diesing, 1863)	M	<i>hubrichti</i> Stiasny-Wijnhoff, 1926	M
<i>helvolum</i> Bürger, 1895	AM	<i>splendida</i> (Keferstein, 1862)	A
<i>hermafroditicum</i> (Keferstein, 1868)	A	<i>Wijnhoffella</i>	
<i>herouardi</i> (Oxner, 1908)	A	<i>alexandrinensis</i> Friedrich, 1940	M
<i>humilis</i> (Quatrefages, 1846)	M		
<i>interruptum</i> Bürger, 1895	M	Family Paradrepanophoridae	
<i>kefersteinii</i> (Marion, 1869)	M	<i>Paradrepanophorus</i>	
<i>kirsteueri</i> Senz, 1995	A	<i>corallincola</i> Stiasny-Wijnhoff, 1926	M
<i>knochii</i> (Kölliker, 1845)	M	<i>crassus</i> (Quatrefages, 1846)	AM
<i>laminariae</i> Ushakov, 1928	A	<i>nisidensis</i> (Hubrecht, 1874)	M
<i>leonillae</i> (Oxner, 1908)	A		
<i>longissimum</i> Bürger, 1895	AM	Family Uniporidae	
		<i>Uniporus</i>	
		<i>acutocaudatus</i> (Brinkmann, 1914/15)	A
		<i>borealis</i> (Punnett, 1901)	A
		<i>hyalinus</i> Brinkmann, 1914/15	A

Order Pelagica**Family Armaueriidae**

Armaueria
rubra Brinkmann, 1917

A

Family Balaenamertidae

Balaenamertes
chavesi (Joubin, 1906)
grandis Brinkmann, 1917
hjorti Brinkmann, 1917
lata Brinkmann, 1917
lobata (Joubin, 1906)
musculocaudata Brinkmann, 1917

A A A A A A

Family Buergeriellidae

Buergeriella
notabilis Brinkmann, 1917

A

Family Chuniellidae

Chuniella
elongata (Joubin, 1906)
lanceolata Brinkmann, 1917

A A

Family Dinonemertidae

Dinonemertes
alberti (Joubin, 1906)
arctica Korotkevich, 1977
grimaldii (Joubin, 1906)

A A A

Family Nectonemertidae

Nectonemertes
minima Brinkmann, 1915/16
mirabilis Verrill, 1892
primitiva Brinkmann, 1917

A A M A

Family Pelagonemertidae

Gelanemertes
richardi (Joubin, 1906)
Natonemertes
acutocaudata Brinkmann, 1917
Parabalaenamertes
fusca Brinkmann, 1917
zonata (Joubin, 1906)
Probalaenamertes
wijnhoffae Brinkmann, 1917

A A A A A A A

Family Phallonemertidae

Phallonemertes
murrayi (Brinkmann, 1912)

A

Family Planktonemertidae

Crassonemertes
robusta Brinkmann, 1917

A M

Family Protopelagonemertidae

Calonemertes
hardyi (Wheeler, 1934)
Pendonemertes
levinseini Brinkmann, 1917
Plotonemertes
adhaerens Brinkmann, 1917
Protopelagonemertes
hubrechti (Brinkmann, 1917)

A A A M A

Subclass Bdellonemertini**Family Malacobdelliidae**

Malacobdella
grossa (Müller, 1776)

A M

GASTROTRICHA

Compiled by Jean-Loup d'Hondt and Jacob van der Land

The European list was derived from URMO with additions from Zoological Record.

References

Hondt, J.L. d', 1971. Gastrotricha. *Ocean. Mar. Biol. Ann. Rev.* 9: 141-192.
Hondt, J.L. d', 1974. Clés tabulaires de détermination des genres marins de Gastrotriches. *Bull. Soc. zool. Fr.* 99: 645-665.
[In french]

PHYLUM GASTROTRICHA**Order Chaetonotoidea****Family Chaetonotidae**

Aspidiophorus
bisquamosus Mock, 1979
marinus Remane, 1926
mediterraneus Remane, 1927
ornatus Mock, 1979
plystictos Balsamo and Todaro, 1987
tentacularis Wilke, 1954

A A M A M M

Chaetonotus

aegilonensis Balsamo, Fregni and Tongiorgi, 1994
aequispinosus Schrom, 1972
angustus Schrom, 1972
antipai Rodewald, 1938
apechochaetus Hummon, Balsamo and Todaro, 1992
apolemmus Hummon, Balsamo and Todaro, 1992

M M M M M M

Chaetonotus

atrox Wilke, 1954
balticus Remane, 1926
condensus Mock, 1979
decipiens Remane, 1926
dispar Wilke, 1954
hilarus Schrom, 1972
hystrix Metschnikoff, 1865
inaequidens Kisielewski, 1988
lacunosus Mock, 1979
littoralis d'Hondt, 1970
mariae Todaro, 1992
maximus Ehrenberg, 1831
modestus Schrom, 1972

A A M M M M

Chaetonotus

napoleonicus Hummon, Todaro and Tongiorgi, 1993
neptuni Wilke, 1954
oceanioides d'Hondt, 1971
paradoxus Remane, 1929
parthenopeius Wilke, 1954
parvus Wilke, 1954
persetosus Zelinka, 1889

A M A M M M

schultzei Metschnikoff, 1865
serenus Schrom, 1972
siciliensis Hummon, Balsamo and Todaro, 1992
similis Zelinka, 1889
somniculosus Mock, 1979
tempestivus Mock, 1979
variosquamatus Mock, 1979

M M A M A A

Halichaetonotus

aculifer (Gerlach, 1953)
arenarius (d'Hondt, 1971)
balticus Kisielewski, 1975
batillifer (Luporini and Tongiorgi, 1972)
clavicornis Balsamo, Fregni and Tongiorgi, 1995
dubius (Remane, 1926)
etrolomus Hummon, Balsamo and Todaro, 1992
genatus Balsamo, Fregni and Tongiorgi, 1995
jucundus (d'Hondt, 1971)
lamellatus Kisielewski, 1975
littoralis (d'Hondt, 1971)
margaretae Hummon, Balsamo and Todaro, 1992
marivagus Balsamo, Todaro and Tongiorgi, 1992
paradoxus (Remane, 1927)
parvus (Wilke, 1954)

M M A M M M M M M M M

<i>pleuracanthus</i> (Remane, 1926)	A	<i>ponticus</i> Valkanov, 1957	M
<i>riedli</i> Schrom, 1972	M	<i>Dendropodola</i>	
<i>schromi</i> Kisielewski, 1975	A	<i>transitionalis</i> Hummon, Todaro and Tongiorgi, 1993	M
<i>spinosus</i> Mock, 1979	A M		
<i>swedmarki</i> Schrom, 1972	M		
<i>tentaculatus</i> (d'Hondt, 1971)	A	Family Lepidodasyidae	
<i>thalassopais</i> Hummon, Balsamo and Todaro, 1992	M	<i>Cephalodasys</i>	
<i>atlanticus</i> Kisielewski, 1988	A	<i>hadrosomus</i> Hummon, Todaro and Tongiorgi, 1993	M
<i>Hemichaetonotus</i>		<i>maximus</i> Remane, 1926	A
<i>clipeatus</i> (Schrom, 1972)	M	<i>palavensis</i> Fize, 1963	M
<i>Heterolepidoderma</i>		<i>Dolichodasys</i>	
<i>arenosum</i> Kisielewski, 1988	A	<i>carolinensis</i> Ruppert and Shaw, 1977	A
<i>armatum</i> Schrom, 1966	M	<i>delicatus</i> Ruppert and Shaw, 1977	A
<i>axi</i> Mock, 1979	A	<i>elongatus</i> Gagne, 1977	A
<i>brevituberculatum</i> Kisielewski, 1981	A	<i>Lepidodasys</i>	
<i>contextum</i> Schrom, 1972	M	<i>martini</i> Remane, 1926	A
<i>solidatum</i> Renaud-Mornant, 1967	A	<i>platyurus</i> Remane, 1927	A
<i>grandiculum</i> Mock, 1979	A	<i>unicarenatus</i> Balsamo, Fregni and Tongiorgi, 1994	M
<i>hermaphroditum</i> Wilke, 1954	A	<i>Megadasys</i>	
<i>istrianum</i> Schrom, 1972	M	<i>minor</i> Kisielewski, 1987	A
<i>loricatum</i> Schrom, 1972	M	<i>sterrei</i> (Boaden, 1974)	A
<i>marinum</i> Remane, 1926	A M	<i>Mesodasys</i>	
<i>Ichthydium</i>		<i>adenotubulatus</i> Hummon, Todaro and Tongiorgi, 1993	M
<i>cyclocephalum</i> Grünspan, 1908	M	<i>ischiensis</i> Hummon, Todaro and Tongiorgi, 1993	M
<i>rupperti</i> Mock, 1979	A	<i>laticaudatus</i> Remane, 1951	A M
<i>supralitorale</i> Mock, 1979	A	<i>litoralis</i> Remane, 1951	A M
<i>tergestinum</i> Grünspan, 1908	M	<i>lobocercus</i> (Boaden, 1960)	A
<i>Lepidodermella</i>		<i>Paradasys</i>	
<i>limogenum</i> Schrom, 1972	M	<i>hexadactylus</i> Karling, 1954	A
<i>Musellifer</i>		<i>subterraneus</i> Remane, 1934	A M
<i>delamarei</i> (Renaud-Mornant, 1968)	M	<i>Pleurodasys</i>	
<i>profundus</i> Vivier, 1974	M	<i>helgolandicus</i> Remane, 1927	A
<i>Polymerurus</i>		<i>megasoma</i> Boaden, 1963	A
<i>nodicaudus</i> (Voigt, 1901)	A M	<i>Psammodasys</i>	
<i>tesselatus</i> Renaud-Mornant, 1968	M	<i>cambiensis</i> (Boaden, 1963)	A
Family Neodasyidae		<i>turbanelloides</i> (Boaden, 1960)	A
<i>Neodasys</i>		Family Macrodasyidae	
<i>chaetonotheideus</i> Remane, 1927	A M	<i>Macrodasys</i>	
<i>uchidai</i> Remane, 1961	A M	<i>affinis</i> Remane, 1936	A
Family Xenotrichulidae		<i>balticus</i> Roszczak, 1939	A M
<i>Draculiciteria</i>		<i>buddenbrocki</i> Remane, 1924	A M
<i>tesselata</i> (Renaud-Mornant, 1968)	A M	<i>caudatus</i> Remane, 1927	A M
<i>Heteroxenotrichula</i>		<i>cephalatus</i> Remane, 1927	A
<i>affinis</i> (Remane, 1934)	A	<i>gerlachi</i> Papi, 1957	M
<i>arcassonensis</i> Ruppert, 1979	A	<i>neapolitanus</i> Papi, 1957	M
<i>flandrensis</i> (d'Hondt, 1968)	A	<i>remanei</i> Boaden, 1963	A
<i>pygmaea</i> Remane, 1934	A M	<i>thucus</i> Luporini, Magagnini and Tongiorgi, 1970	M
<i>squamosa</i> Wilke, 1954	A M	<i>Urodasys</i>	
<i>subterranea</i> (Remane, 1934)	A M	<i>mirabilis</i> Remane, 1926	A
<i>variocirrata</i> (d'Hondt, 1966)	A M	<i>roscoffensis</i> Kisielewski, 1987	A
<i>Xenotrichula</i>		<i>viviparus</i> Wilke, 1954	M
<i>beauchampi</i> Lévi, 1950	A M	Family Planodasyidae	
<i>bispina</i> Roszczak, 1939	A M	<i>Crasiella</i>	
<i>cornuta</i> Wilke, 1954	A M	<i>diplura</i> Clausen, 1968	A
<i>intermedia</i> Remane, 1934	A M	<i>oceanica</i> d'Hondt, 1974	A
<i>lineata</i> Schrom, 1972	A M	Family Thaumastodermatidae	
<i>micracantha</i> (Remane, 1926)	A	<i>Acanthodasys</i>	
<i>mirabilis</i> d'Hondt, 1967	A	<i>aculeatus</i> Remane, 1927	A M
<i>punctata</i> Wilke, 1954	A M	<i>arcassonensis</i> Kisielewski, 1987	A
<i>quadriflatala</i> Kisielewski, 1988	A	<i>Diplodasys</i>	
<i>simplex</i> Mock, 1979	A	<i>ankeli</i> Wilke, 1954	A M
<i>soikai</i> Schrom, 1966	M	<i>caudatus</i> Kisielewski, 1987	A
<i>velox</i> Remane, 1927	A M	<i>meloriae</i> Todaro, Balsamo and Tongiorgi, 1992	M
Order Macrodoasyoidea		<i>minor</i> Remane, 1936	A M
Family Dactylopodolidae		<i>platydasyoides</i> Remane, 1927	M
<i>Dactylopodola</i>		<i>swedmarki</i> Kisielewski, 1987	A
<i>baltica</i> (Remane, 1926)	A	<i>Hemidasys</i>	
<i>cornuta</i> (Swedmark, 1956)	A	<i>agaso</i> Claparède, 1867	M
<i>roscovita</i> (Swedmark, 1967)	A	<i>Platydasys</i>	
<i>typhle</i> (Remane, 1929)	A M	<i>brachycephalus</i> Lévi, 1954	A
<i>weilli</i> (d'Hondt, 1965)	A	<i>mastiurus</i> Clausen, 1965	A
<i>Dendrodasys</i>		<i>maximus</i> Remane, 1927	A
<i>affinis</i> Wilke, 1954	M	<i>ocellatus</i> Clausen, 1965	A
<i>gracilis</i> Wilke, 1954	M	<i>phacellatus</i> Clausen, 1965	M
		<i>rarus</i> Forneris, 1961	A

<i>ruber</i> Swedmark, 1956	M	<i>ambronensis</i> Remane, 1943	A M
<i>styliferus</i> Boaden, 1965	M	<i>bocqueti</i> Kaplan, 1958	A M
<i>tentaculatus</i> Swedmark, 1956	M	<i>cornuta</i> Remane, 1925	A M
<i>Pseudostomella</i>		<i>hyalina</i> Schultze, 1853	A M
<i>etrusca</i> Hummon, Todaro and Tongiorgi, 1993	M	<i>italica</i> Gerlach, 1953	M
<i>roscovita</i> Swedmark, 1956	A	<i>remanei</i> Forneris, 1961	A
<i>Ptychostomella</i>		<i>pontica</i> Valkanov, 1957	M
<i>helena</i> Roszczak, 1936	A	<i>thiophila</i> Boaden, 1974	A M
<i>mediterranea</i> Remane, 1927	M	<i>plana</i> (Giard, 1904)	A
<i>ommatophora</i> Remane, 1927	A	<i>lutheri</i> Remane, 1952	A
<i>pectinata</i> Remane, 1936	A	<i>subterrena</i> Remane, 1934	A
<i>tyrrhenica</i> Hummon, Todaro and Tongiorgi, 1993	M	<i>multidigitata</i> Kisielewski, 1987	A
<i>Tetranchyroderma</i>		<i>reducta</i> Boaden, 1974	A
<i>antennatum</i> Luporini, Magagnini and Tongiorgi, 1970	A M	<i>varians</i> Maguire, 1976	A
<i>apus</i> Remane, 1927	A	<i>otti</i> Schrom in Riedl, 1970	M
<i>boadeni</i> Schrom in Riedl, 1970	M	<i>veneziana</i> Schrom, 1972	M
<i>cirrophora</i> Lévi, 1950	A M	<i>petiti</i> Remane, 1952	M
<i>coelopodium</i> Boaden, 1963	A		
<i>dragescoi</i> Swedmark, 1967	A		
<i>esarabdophorum</i> Tongiorgi and Balsamo, 1984	M		
<i>heterotubulatum</i> Hummon, Todaro and Tongiorgi, 1993	M		
<i>hirtum</i> Luporini, Magagnini and Tongiorgi, 1970	M		
<i>hypopsilancrum</i> Hummon, Todaro and Tongiorgi, 1993	M		
<i>hystrix</i> Remane, 1926	A		
<i>insulare</i> Balsamo, Fregni and Tongiorgi, 1994	M		
<i>massiliense</i> Swedmark, 1956	A M		
<i>megastoma</i> (Remane, 1927)	A M		
<i>pachysomum</i> Hummon, Todaro and Tongiorgi, 1993	M		
<i>pacificum</i> Schmidt, 1974	M		
<i>papii</i> Gerlach, 1953	A M		
<i>polyacanthus</i> (Remane, 1927)	A		
<i>polypodium</i> Luporini, Magagnini and Tongiorgi, 1971	M		
<i>quadrilentaculatum</i> Todaro, Balsamo and Tongiorgi, 1992	M		
<i>sanctaecaterinae</i> Todaro, Balsamo and Tongiorgi, 1992	M		
<i>sarokem</i> Todaro, Balsamo and Tongiorgi, 1977	A M		
<i>suecica</i> Boaden, 1960	A		
<i>thysanogaster</i> Boaden, 1965	M		
<i>thysanophorum</i> Hummon, Todaro and Tongiorgi, 1993	M		
<i>tribolosum</i> Clausen, 1965	A		
<i>vera</i> Wilke, 1954	A		
<i>Thaumastoderma</i>			
<i>arcassonense</i> d'Hondt, 1965	A		
<i>bifurcatum</i> Clausen, 1991	A		
<i>cantacuzeni</i> Lévi, 1959	A		
<i>heideri</i> Remane, 1926	A		
<i>mediterraneum</i> Remane, 1927	A M		
<i>ramuliferum</i> Clausen, 1965	A M		
<i>renaudi</i> Kisielewski, 1987	A		
<i>swedmarki</i> Lévi, 1950	A M		
<i>truncatum</i> Clausen, 1991	A		
Family Chordodasyidae			
<i>Xenodasys</i>			
<i>sanctigouveni</i> Swedmark, 1967	A		
Family Turhanellidae			
<i>Desmodasys</i>			
<i>phocoides</i> Clausen, 1965	A		
<i>Dinodasys</i>			
<i>mirabilis</i> Remane, 1927	A		
<i>Paraturbanella</i>			
<i>armoricana</i> (Swedmark, 1954)	A		
<i>cuaneensis</i> Maguire, 1976	A		
<i>dohrni</i> Remane, 1927	A M		
<i>eireanna</i> Maguire, 1976	A		
<i>intermedia</i> Wieser, 1957	A		
<i>microptera</i> Wilke, 1954	M		
<i>pallida</i> Luporini, Magagnini and Tongiorgi, 1970	M		
<i>teissieri</i> Swedmark, 1954	A M		
<i>Pseudoturbanella</i>			
<i>stylifera</i> d'Hondt, 1968	A		
<i>Turbanella</i>			

CEPHALORHYNCHA

Kinorhyncha

Compiled by Birger Neuhaus and Jacob van der Land

The ERMS list for Kinorhyncha and Loricifera was compiled by J. van der Land and Birger Neuhaus. The list of valid species of Kinorhyncha is based on an unpublished list of synonymies by Robert P. Higgins made available to B. Neuhaus.

The Kinorhyncha are a group of microscopic marine animals that comprise some 130 valid species (Pardos *et al.* 1998). Few species live exclusively in eulittoral sediments (e.g., *Echinoderes coulli* Higgins, 1977, *E. caribensis* Kirsteuer, 1964) or in beach sands (e.g., *Cateria styx* Gerlach, 1956). However, most species inhabit the eulittoral and sublittoral including the deep-sea. Kinorhyncha live in the interstitial cavities and crevices of coarse or fine sandy substrate or of muddy sediments (Gerlach 1956, Higgins 1990, Vanhove *et al.* 1995, Zelinka 1928). Only the upper 1-10 cm of oxygen-rich substratum contain Kinorhyncha both in the eulittoral and sublittoral (Horn 1978, Thistle *et al.* 1985). Kinorhyncha feed on diatoms or bacteria (Higgins 1990, own observations). Kinorhyncha can be extracted from the sediment by the bubble and blot technique (Higgins & Thiel 1988). Higgins & Thiel (1988) describe additional suggestions for specimen processing.

Close to nothing is known about the biogeography of kinorhynch species. Every scientist trying to identify Kinorhyncha from the area covered by ERMS is, therefore, well advised to consider species described from neighbouring areas as well, such as the following recorded from other parts of the Arctic Ocean: *Echinoderes angustus* Higgins & Kristensen, 1988, *Echinoderes aquilonius* Higgins & Kristensen, 1988, *Echinoderes arlis* Higgins, 1966, *Echinoderes eximus* Higgins & Kristensen, 1988, *Echinoderes peterseni* Higgins & Kristensen, 1988, *Echinoderes tubilak* Higgins & Kristensen, 1988, *Pycnophyes borealis* Higgins & Korczynski, 1989, *Pycnophyes canadensis* Higgins & Korczynski, 1989, *Pycnophyes chukchiensis* Higgins, 1991, *Pycnophyes cryopygus* Higgins & Kristensen, 1988, *Pycnophyes greenlandicus* Higgins & Kristensen, 1988, *Pycnophyes mokievskii* Adrianov, 1995, *Pycnophyes spitsbergensis* Adrianov, 1995

It is also not impossible that species described from the North American coast may occur in European waters or species new to science may turn up. More recent keys for identification include Adrianov (1995), Higgins (1983), Higgins & Kristensen (1988), Huys & Coomans (1989), Pardos *et al.* (1998). Valuable information is also found in Adrianov & Malakhov (1994), Higgins (1977, 1978, 1985, 1990), Moore (1973), Nebelsick (1990), and Zelinka (1928).

References

- Adrianov, A.V. & V.V. Malakhov 1994: *Kinorhyncha: structure, development, phylogeny and taxonomy*. - Nauka Publ., Moscow, pp. 260. [in Russian]
- Adrianov, A. V. 1995: The first description of kinorhynchs from the Spitsbergen Archipelago (Greenland Sea), with a key to the genus *Pycnophyes* (Homalorhagida, Kinorhyncha). *Canadian Journal of Zoology* 73, 1554-1566.
- Gerlach, S. 1956 Über einen aberranten Vertreter der Kinorhynchen aus dem Küstengrundwasser. *Kieler Meeresforschungen* 12: 120-124.
- Higgins, R. P. 1977: Redescription of *Echinoderes dujardinii* (Kinorhyncha) with descriptions of closely related species. *Smithsonian Contributions to Zoology* 248, 1-26.
- Higgins, R. P. 1978: *Echinoderes gerardi* n. sp. and *E. riedli* (Kinorhyncha) from the Gulf of Tunis. *Transactions of the American Microscopical Society* 97: 171-180.
- Higgins, R. P. 1983: The Atlantic barrier reef ecosystem at Carrie Bow Cay, Belize, II: Kinorhyncha. *Smithsonian Contributions to Marine Science* 18: 1-131.
- Higgins, R. P. 1985: The genus *Echinoderes* (Kinorhyncha: Cyclorhagida) from the English channel. *Journal of the Marine biological Association of the United Kingdom* 65: 785-800.
- Higgins, R. P. 1990: Zelinkaderidae, a new family of cyclorhagid Kinorhyncha. *Smithsonian Contributions to Zoology* 500: 1-26.
- Higgins, R. P. & R. M. Kristensen 1988: Kinorhyncha from Disko Island, West Greenland. *Smithsonian Contributions to Zoology* 458: 1-56.
- Higgins, R. P. & H. Thiel 1988: Introduction to the study of meiofauna. - Smithsonian Institution press, Washington, D. C., 488 pp.
- Horn, T. D. 1978: The distribution of *Echinoderes coulli* (Kinorhyncha) along an interstitial salinity gradient. *Transactions of the American Microscopical Society* 97: 586-589.
- Huys, R. & A. Coomans 1989: *Echinoderes higginsi* sp. n. (Kinorhyncha, Cyclorhagida) from the southern North Sea with a key to the genus *Echinoderes* Claparede. *Zoologica Scripta* 18: 211-221.
- Moore, P. G. 1973: *Campyloderes macquariae* Johnston, 1938 (Kinorhyncha: Cyclorhagida) from the northern hemisphere. *Journal of Natural History* 7: 341-354.
- Nebelsick, M. 1990: *Antygomonas incomitata* gen. et sp. n. (Cyclorhagida, Kinorhyncha) and its phylogenetic relationships. *Zoologica Scripta* 19: 143-152.
- Pardos, F., R. P. Higgins & J. Benito 1998: *Echinoderes cantabricus* sp. n. and *E. hispanicus* sp. n. (Kinorhyncha, Cyclorhagida) from the north coast of Spain. *Zoologischer Anzeiger* 237: 195-208.
- Rachor, E. 1975: Quantitative Untersuchungen über das meiobenthos der nordostatlantischen Tiefsee. "Meteor" Forschungsergebnisse, Reihe D, 21: 1-10.
- Thistle, D., J. Y. Yingst & K. Fauchald 1985: A deep-sea benthic community exposed to strong near-bottom currents on the Scotian Rise (Western Atlantic). *Marine Geology* 66: 91-112.
- Vanhove, S., J. Wittoeck, G. Desmet, B. van den Berghe, R. L. Herman, R. P. M. Bak, G. Nieuwland, J. H. Vosjan, A. Boldrin, S. Rabitti & M. Vincx 1995: Deep-sea meiofauna communities in Antarctica: Structural analysis and relation with the environment. *Marine Ecology Progress Series* 127: 65-76.
- Zelinka, K. 1928: *Monographie der Echinodera*. Verlag W. Engelmann, Leipzig, 396 S.

PHYLUM CEPHALORHYNCHA

Class Kinorhyncha

Order Cyclorhagida

Family Antygomonidae

Antygomonas incomitata Nebelsick, 1990

M

Family Centroderidae

Campyloderes

adherens Nyholm, 1947
macquariae Johnston, 1938

A

Centroderes

eisigii Zelinka, 1928
spinosis (Reinhard, 1881)

M

Condyloderes

multispinosus (McIntyre, 1962)

A

Family Echinoderidae

Echinoderes

agigens Bacescu, 1968

M

<i>cantabricus</i> Pardos, Higgins & Benito, 1998	A	Nanalicida, Nanalicidae) from the Mediterranean Sea. <i>Italian Journal of Zoology</i> 65: 219-226.
<i>capitatus</i> (Zelinka, 1928)	M	
<i>citrinus</i> Zelinka, 1928	M	
<i>druxi</i> d'Hondt, 1973	M	
<i>dujardini</i> Claparède, 1863	A	
<i>elongatus</i> Nyholm, 1947	A	
<i>ferrugineus</i> Zelinka, 1928	M	
<i>gerardi</i> Higgins, 1978	M	
<i>higginsi</i> Huys & Coomans, 1989		
<i>hispanicus</i> Pardos, Higgins & Benito, 1998	A	
<i>krishnaswamyi</i> Higgins, 1985	A	
<i>kristensenii</i> Higgins, 1985	A	
<i>levanderi</i> Karling, 1954	A	
<i>riedli</i> Higgins, 1966	M	
<i>seliger</i> Greef, 1869		
<i>worthingii</i> Southern, 1914	A	
Family Semnoderidae		
<i>Semnoderes</i>		
<i>armiger</i> Zelinka, 1928	A M	
<i>ponticus</i> Bacescu & Bacescu, 1956	M	
Family Zelinkaderidae		
<i>Zelinkaderes</i>		
<i>submersus</i> (Gerlach, 1969)	A	
Order Homalorhagida		
Family Neocentrophyidae		
<i>Paracentrophyes</i>		
<i>quadridentatus</i> (Zelinka, 1928)	A M	
Family Pycnophyidae		
<i>Kinorhynchus</i>		
<i>giganteus</i> (Zelinka, 1928)	A M	
<i>paraneapolitanus</i> Sheremetevskij, 1974	M	
<i>Pycnophyes</i>		
<i>calmani</i> Southern, 1914	A	
<i>carinatus</i> Zelinka, 1928	M	
<i>communis</i> Zelinka, 1908	A M	
<i>dentatus</i> (Reinhard, 1881)	A M	
<i>flaveolatus</i> Zelinka, 1928	A M	
<i>kielensis</i> Zelinka, 1928	A M	
<i>maximus</i> Reimer, 1963	A	
<i>ponticus</i> (Reinhard, 1881)	M	
<i>robustus</i> Zelinka, 1928	M	
<i>rugosus</i> Zelinka, 1928	M	
<i>zelinkaei</i> Southern, 1914	A	

Loricifera

Compiled by Jacob van der Land and Birger Neuhaus

The marine Loricifera are now known with 11 species from North America, Europe, and the Japanese deep-sea (Higgins & Kristensen 1986, Kristensen 1983, Todaro & Kristensen 1998). Only 2 species have been reported from Europe (Kristensen 1983, Todaro & Kristensen 1998). Loricifera inhabit various kinds of sediment from coarse sand or shelly gravel to red clay with silt and sand. They may be extracted from the substratum either by freshwater shock (Kristensen 1983) or by multiple decantations (Todaro & Kristensen 1998). It is certain that species new to science inhabit the area covered by the ERMS programme.

References

- Higgins, R. P. & R. M. Kristensen 1986: New Loricifera from Southeastern United States coastal waters. *Smithsonian Contributions to Zoology* 438: 1-70.
- Kristensen, P. M. 1983: Loricifera, a new phylum with Aschelminthes characters from the meiobenthos. *Zeitschrift für zoologische Systematik und Evolutionsforschung* 21: 163-180.
- Todaro, M. A. & R. M. Kristensen 1998: A new species and first report of the genus *Nanalicus* (Loricifera,

Class Loricifera

Family Nanalicidae

<i>Nanalicus</i>	
<i>mysticus</i> Kristensen, 1983	A
<i>khaitatus</i> Todaro & Kristensen, 1998	M

Nematomorpha and Priapulida

Compiled by Jacob van der Land

The ERMS lists for Priapulida and Nematomorpha were based on Adrianov & Malakhov (1996) and on Malakhov & Adrianov (1995).

References

- Adrianov, A.V. & V.V. Malakhov (1996): Priapiliida [sic] (Priapulida): structure, development, phylogeny, and classification. - KMK Scientific Press, Moscow, pp. 266. [in Russian]
- Malakhov, V.V. & A.V. Adrianov (1995): Cephalorhyncha - A new phylum of the animal kingdom. - KMK Scientific Press, Moscow, pp. 202. [in Russian]

Class Nematomorpha

Order Nectonematida

Family Nectonematidae	
<i>Nectonema</i>	
<i>agile</i> Verril, 1879	A M
<i>svensksundi</i> Bock, 1908	A
<i>munidae</i> Brinkmann, 1930	A

Class Priapulida

Family Chaetostephanidae

<i>Maccabeus</i>	
<i>tentaculatus</i> Por, 1973	M

Family Priapulidae

<i>Halicryptus</i>	
<i>spinulosus</i> von Siebold, 1849	A
<i>Priapulopsis</i>	
<i>bicaudatus</i> (Koren & Danielssen, 1868)	A
<i>cnidophorus</i> Salvini-Plawen, 1973	M
<i>Priapulus</i>	
<i>abyssorum</i> Menzies, 1959	A M
<i>caudatus</i> Lamarck, 1816	A

Family Tubiluchidae

<i>Tubiluchus</i>	
<i>arcticus</i> Adrianov et al., 1989	A

NEMATODA

Nematoda (free-living)

Compiled by Guy De Smet, Magda Vincx, Ann Vanreusel, Sandra Vanhove, Jan Vanaverbeke and Maaike Steyaert

The list was compiled by Guy De Smet, a non-specialist, who was assisted by nematologists Magda Vincx, Ann Vanreusel, Sandra Vanhove, Jan Vanaverbeke and Maaike Steyaert. Dr. F. Riemann gave useful advice for the genus *Daptonema*. The classification in the list is based upon Lorenzen (1979, 1994). Information on the geographic distribution in the list is based upon Gerlach and Riemann (1973, 1974), Platt and Warwick (1983, 1988) and Warwick *et al.* (1998).

References

- Lorenzen, S. *Entwurf eines Phylogenetischen Systems der Freilebenden Nematoden*, Kiel, 1979.
 Lorenzen, S. *The Phylogenetic Systematics of Freeliving Nematodes*, The Ray Society, 1994.
 Gerlach, S. A. and Riemann, F. The Bremerhaven Checklist of Aquatic Nematodes. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven*, Supplement 4, (Heft 1 1973, Heft 2 1974).
 Platt, H. M. and Warwick, R. M. Free-living Marine Nematodes (Part I British Enoplids) *Synopses of the British Fauna* (New series) No. 28, 1983.
 Platt, H. M. and Warwick, R. M. Free-living Marine Nematodes (Part II British Chromadorids) *Synopses of the British Fauna* (New series) No. 38, 1988.
 Warwick, R. M., Platt, H. M. and Somerfield, P. J. Free-living Marine Nematodes (Part III Monhystrerids) *Synopses of the British Fauna* (New series) No. 53, 1998.

NEMATODA, FREE-LIVING

- Acantholaimus*
setosus Vitiello, 1970
ewensis Platt & Zhang, 1982
polydentatus Gerlach, 1951
longisetosus Allgén, 1933
- Acanthonchus*
gracilis (Ditlevsen, 1918) Wieser, 1955
viviparus Cobb, 1920
arcuatus (Kreis, 1928) Wieser, 1955
pugionatus Vitiello, 1970
- Acanthopharyngoides*
tyrrhenicus Wieser, 1954
- Acanthopharynx*
setosissima Schuurmans Stekhoven, 1943
micrampsis Schuurmans Stekhoven, 1942
perarmata Marion, 1870
rigida Schuurmans Stekhoven, 1950
micans (Eberth, 1863) Marion, 1870
affinis Marion, 1870
- Actinonema*
longicaudatum (Steiner, 1918) Wieser, 1954
pachydermatum Cobb, 1920
sidatum Vitiello, 1970
celtica Boucher, 1976
- Adeuchromadora*
megamphida Boucher & Bovée, 1972
- Adoncholaimus*
thalassophysgas (de Man, 1876) Filipjev, 1918
panicus Cobb, 1930
derjugini (Saweljew, 1912) Filipjev, 1924
fuscus (Bastian, 1865) Filipjev, 1918
lepidus (de Man, 1889) Filipjev, 1918
islandicus Kreis, 1963
- Aegialoalaimus*
- punctatus* (Allgén, 1929) Schuurmans Stekhoven, 1931
elegans de Man, 1907
tenuis Kreis, 1928
tenuicaudatus Allgén, 1932
- Alaimella*
cincta Cobb, 1920
- Amphidelus*
puccinelliae Lorenzen, 1966
- Amphimonhystera*
anechma (Southern, 1914) Allgén, 1929
helgolandica Riemann, 1967
paranechma Allgén, 1935
subtilis Lorenzen, 1972
- Anoplostoma*
elegans Kreis, 1929
blanchardi de Man, 1888
exceptum Schulz, 1935
viviparum (Bastian, 1865) Bütschli, 1874
- Anticoma*
strandii Allgén, 1940
stekhoveni Wieser, 1953
acuminata (Eberth, 1863) Bastian, 1865
pellucida Bastian, 1865
murmanica Filipjev, 1927
minor Filipjev, 1927
insulaealbae Filipjev, 1927
arctica Steiner, 1916
longicaudata Kreis, 1928
brevisetosa Platonova, 1967
brevicaudata Kreis, 1963
eberti Bastian, 1865
- Anticomopsis*
typica Micoletzky, 1930
- Anticyathus*
primitivus (Allgén, 1933) Gerlach, 1963
- Anticyclus*
gigas Vitiello, 1969
exilis Cobb, 1920
pachyamphis (Wieser, 1956) Gerlach, 1963
- Antomicron*
pratense Lorenzen, 1966
profundum Vitiello, 1971
elegans (de Man, 1922) De Coninck, 1965
pellucidum Cobb, 1920
- Apomema*
torosa (Lorenzen, 1973) Jensen, 1978
- Araeolaimus*
paucisetosus (Wieser, 1951) Lorenzen, 1981
penelope Moore, 1977
septentrionalis Kreis, 1963
microphthalmus (de Man, 1893) Lorenzen, 1981
macrocirculus Kreis, 1928
longisetosus Wieser, 1954
filipjevi Schuurmans Stekhoven & Adam, 1931
demani (Schuurmans Stekhoven, 1950) Wieser, 1956
bioculatus (de Man, 1876) de Man, 1888
gajevskii Paramonov, 1929
mediterraneus (de Man, 1876) de Man, 1888
ponticus Filipjev, 1922
elegans de Man, 1888
longicauda Allgén, 1929
zosterae (Filipjev, 1918) Chitwood & Chitwood, 1937
tenuicaudatus Allgén, 1935
supralitoralis Wieser, 1960
steineri Filipjev, 1922
- Ascolaimus*
elongatus (Bütschli, 1874) Schuurmans Stekhoven, 1932
- Astomonema*
southwardorum Austen *et al.*, 1993
- Atrochromadora*
parva (de Man, 1893) Wieser, 1954
microlaima (de Man, 1889) Wieser, 1954
- Austranema*

- shirleyae* (Coles, 1965) Inglis, 1969
- Axonolaimus*
- spinosus* (Bütschli, 1874) de Man, 1889
 - helgolandicus* Lorenzen, 1972
 - hexapilus* Wieser & Hopper, 1967
 - paraspinosus* Schuurmans Stekhoven & Adam, 1931
 - filiiformis* de Man, 1889
 - drachi* Luc & De Coninck, 1959
 - arcuatus* Schuurmans Stekhoven, 1950
 - orcombensis* Warwick, 1970
 - schuurmansstekhoveni* Allgén, 1935
 - limalis* Ssweljev, 1912
 - tirrenicus* Brunetti, 1941
 - typicus* de Man, 1922
 - setosus* Filipjev, 1918
 - ponticus* Filipjev, 1918
 - villosus* Skwarra, 1922
 - demanii* De Coninck & Schuurmans Stekhoven, 1933
- Barbonema*
- setiferum* Filipjev, 1927
 - horridum* Gerlach, 1956
- Bathyepsilonema*
- pustulatum* Gerlach, 1952
- Bathylaimus*
- parafilicaudatus* Allgén, 1935
 - arcticus* Kreis, 1963
 - septentrionalis* (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
 - denticaudatus* Allgén, 1930
 - zostericola* (Allgén, 1933) De Coninck & Schuurmans Stekhoven, 1933
 - macramphis* (Schuurmans Stekhoven & De Coninck, 1933) De Coninck & Schuurmans Stekhoven, 1933
 - longisetosus* (Allgén, 1929) De Coninck & Schuurmans Stekhoven, 1933
 - filicaudatus* (Schuurmans Stekhoven & Adam, 1931) De Coninck & Schuurmans Stekhoven, 1933
 - dentatus* Kreis, 1963
 - capacosus* Hopper, 1962
 - cobbi* Filipjev, 1922
 - australis* Cobb, 1894
 - inermis* (Ditlevsen, 1918) de Man, 1922
 - paralongisetosus* Schuurmans Stekhoven & De Coninck, 1933
 - stenolaimus* Schuurmans Stekhoven & De Coninck, 1933
 - tenuicaudatus* (Allgén, 1933) Allgén, 1947
- Bellbolla*
- teissieri* (Luc & De Coninck, 1959) Andrassy, 1973
 - galanachmorae* (Inglis, 1961) Andrassy, 1973
- Boreomicrolaimoides*
- problematicus* (Ditlevsen, 1918) Allgén, 1954
- Bradybucca*
- rhopalura* (Schuurmans Stekhoven, 1950) Schuurmans Stekhoven, 1956
- Calligyrus*
- gerlachi* Lorenzen, 1969
- Calyptronema*
- maxweberi* (de Man, 1922) Wieser, 1953
 - klatti* (Allgén, 1953) Wieser, 1953
 - setifer* (Gerlach, 1953) Wieser, 1953
 - acuminatum* (Eberth, 1863) Wieser, 1953
 - longisetosum* (Schuurmans Stekhoven, 1943) Wieser, 1953
 - eberthi* (de Man, 1876) Schuurmans Stekhoven, 1950
 - sabulicola* (Filipjev, 1918) Kreis, 1926
- Camacolaimoides*
- praedator* (de Man, 1922) De Coninck & Schuurmans Stekhoven, 1933
- Camacolaimus*
- barbatus* Warwick, 1970
 - zostericola* (Filipjev, 1918) Allgén, 1932
 - tardus* de Man, 1889
 - longicauda* de Man, 1922
 - propinquus* Allgén, 1929
- pyrtherchi* Chitwood, 1935
- glauxicola* Allgén, 1951
- dolichocercus* Filipjev, 1922
- bathycola* Filipjev, 1922
- Campylaimus*
- cylindricus* Gerlach, 1956
 - gerlachi* Timm, 1961
 - mirus* Gerlach, 1956
 - inaequalis* Cobb, 1920
 - leseverei* Gerlach, 1956
- Ceramonema*
- undulatum* De Coninck, 1942
 - filipjevi* De Coninck, 1942
 - pisanum* Gerlach, 1953
 - chitwoodi* De Coninck, 1942
 - reticulatum* Chitwood, 1936
 - yunfengi* Platt & Zhang, 1982
 - salsicum* Gerlach, 1956
- Cervonema*
- tenuicaudatum* (Wieser, 1954) Schuurmans Stekhoven, 1950
- Chaetonema*
- vicinum* Gerlach, 1954
 - longisetum* (Steiner, 1916) Filipjev, 1927
 - riemannii* Platt, 1973
- Cheironchus*
- vorax* Cobb, 1917
- Chitwoodia*
- tripapillata* Jayasree, 1976
 - warwicki* Jayasree, 1976
 - menora* Gerlach, 1956
 - falcata* Gerlach, 1956
- Chloronemella*
- filiformis* Allgén, 1929
- Choanolaimus*
- psammophilus* de Man, 1880
- Choniolaimus*
- papillatus* Ditlevsen, 1918
 - panicus* Gerlach, 1956
 - novempapillatus* Schuurmans Stekhoven, 1946
 - crassus* Vitiello, 1971
- Chromadora*
- brevipapillata* Micoletzky, 1924
 - macrolaima* de Man, 1889
 - nudicapitata* Bastian, 1865
 - caeca* Bastian, 1865
 - papillata* Bastian, 1865
 - sabelloides* Bastian, 1865
 - bipapillata* Micoletzky, 1924
 - buesumensis* Kreis, 1924
 - kreisi* Schuurmans Stekhoven & Adam, 1931
 - undecimpapillata* Wieser, 1959
 - axi* Gerlach, 1951
- Chromadorella*
- membranata* (Micoletzky, 1922) Wieser & Hopper, 1967
 - duopapillata* Platt, 1973
 - acriminiensis* (Gerlach, 1953) Lorenzen, 1972
 - parapoecilosoma* (Micoletzky, 1922) Wieser & Hopper, 1967
 - filiformis* (Bastian, 1865) Filipjev, 1918
- Chromadorina*
- inversa* Wieser, 1955
 - germanica* (Bütschli, 1874) Wieser, 1954
 - armata* (Allgén, 1933) Wieser, 1954
 - bergensis* (Allgén, 1932) Wieser, 1954
 - cervix* (Wieser, 1951) Wieser, 1954
 - bioculata* (Carus, 1857) Wieser, 1954
 - demanii* Inglis, 1962
 - erythrophthalma* (Schneider, 1906) Wieser, 1954
 - incurvata* Wieser, 1956
 - laeta* (de Man, 1876) Micoletzky, 1924
 - longisetosa* (De Coninck & Schuurmans Stekhoven, 1933) Wieser, 1954
 - ocellata* (Paramonov, 1929) Wieser, 1954

- viridis* (Linstow, 1876) Wieser, 1954
supralitoralis Lorenzen, 1969
granulopigmentata (Wieser, 1951) Wieser, 1954
rognoensis (Allgén, 1932) Wieser, 1954
macropunctata (Wieser, 1954) Wieser, 1954
obtusa Filipjev, 1918
nuda Wieser, 1954
micoletzkyi Inglis, 1962
majae Wieser, 1968
Chromadorissa
strandii Kreis, 1928
Chromadorita
tenuis (Schneider, 1906) Filipjev, 1922
obliqua (Gerlach, 1953) Wieser, 1954
leuckarti (de Man, 1876) Filipjev, 1929
norvegica (Allgén, 1946) Wieser, 1954
minor (Allgén, 1927) Wieser, 1954
gracilis (Filipjev, 1922) Wieser, 1954
guidoschneideri (Filipjev, 1929) Wieser, 1954
nana Lorenzen, 1973
brevisetosa Gerlach, 1953
minima (Kreis, 1929) Wieser, 1954
hyalocephala (Steiner, 1916) Filipjev, 1922
mucrodonta (Steiner, 1916) Wieser, 1954
abnormis (Kreis, 1928) Wieser, 1954
pachydema (Schneider, 1926) Wieser, 1954
demaniana Filipjev, 1922
heterophya (Steiner, 1916) Filipjev, 1922
brachypharynx (Allgén, 1932) Wieser, 1954
Chromaspirina
lunatica Gerlach, 1965
paucispira Schuurmans Stekhoven, 1950
denticulata (Gerlach, 1953) Luc & De Coninck, 1959
multipapillata Jayasree & Warwick, 1977
thieryi De Coninck, 1943
parapontica Luc & De Coninck, 1959
pellita Gerlach, 1954
crinita Gerlach, 1952
amabilis (de Man, 1922) Luc & De Coninck, 1959
inglisi Warwick, 1970
pontica Filipjev, 1918
Cobia
tresusiaeformis de Man, 1907
caledonia Warwick & Platt, 1973
simplex Allgén, 1929
dentata Gerlach, 1953
trioidonta Filipjev, 1918
Cobionema
acrocerca Filipjev, 1922
cylindrilaimoides Schuurmans Stekhoven, 1950
Comesoma
profundi Bastian, 1865
vulgare Bastian, 1865
stenocephalum Filipjev, 1918
Cominckia
macrospirifera Zhang, 1983
circularis Gerlach, 1956
Corythostoma
filipjevi (Kreis, 1928) Wieser, 1956
Crassolaimus
conicaudatus Kreis, 1929
bipinnatus Wieser, 1951
Crenopharynx
paralepturus (Schuurmans Stekhoven, 1950) Wieser, 1953
gracilis (Linstow, 1900) Filipjev, 1934
marioni (Southern, 1914) Filipjev, 1934
Cricolaimus
elongatus Southern, 1914
Cryptenoplus
gerlachi Riemann, 1966
Curvolaimus
uteratus Vitiello, 1970
Cyartonema
zosterae (Allgén, 1929)
tenuicauda (Allgén, 1933) Juario, 1972
roervikensis (Allgén, 1946) Juario, 1972
elegans Jayasree & Warwick, 1977
germanicum Juario, 1972
Cyatholaimus
savaljevi Allgén, 1933
nordgaardi Allgén, 1933
microsetosus Wieser, 1954
gracilis (Eberth, 1863) Bastian, 1865
minutus Allgén, 1933
paucispira Schuurmans Stekhoven, 1950
canariensis Steiner, 1921
scarratti Inglis, 1961
simulatus Kreis, 1924
prinzi (Marion, 1870) Inglis, 1961
papilliferus Allgén, 1929
ocellatus Bastian, 1865
Cylicolaimus
edentulus Platonova, 1970
obtusidens Ditlevsen, 1926
jaegerskioeldi Türk, 1903
magnus (Villot, 1875) de Man, 1889
Cynura
uniformis Cobb, 1920
Dagda
bipapillata Southern, 1914
asymmetrica Gerlach, 1953
Daptionema
pseudotortum (Vitiello, 1971) Lorenzen, 1977
kornoeense (Allgén, 1929) Lorenzen, 1977
trabeculosum (Schneider, 1906) Lorenzen, 1977
svalbardense (Gerlach, 1965) Lorenzen, 1977
planiere (Vitiello, 1971) Lorenzen, 1977
paradonsi (Allgén, 1949) Lorenzen, 1977
groenlandicum (Ditlevsen, 1928) Lorenzen, 1977
fistulatum (Wieser & Hopper, 1967) Lorenzen, 1977
prominens (Vitiello, 1971) Lorenzen, 1977
curvispiculum (Gerlach, 1953) Lorenzen, 1977
hirsutum (Vitiello, 1967) Lorenzen, 1977
calceolatum (De Coninck & Schuurmans Stekhoven, 1933) Lorenzen, 1977
procerum (Gerlach, 1951) Lorenzen, 1977
psammoides (Warwick, 1970) Warwick *et al.*, 1998
setifer (Gerlach, 1952) Lorenzen, 1977
donsi (Allgén, 1948) Lorenzen, 1977
nanum (Lorenzen, 1972) Lorenzen, 1977
proprium (Lorenzen, 1972) Lorenzen, 1977
normandicum (de Man, 1890) Lorenzen, 1977
trichospiculum (Allgén, 1933) Lorenzen, 1977
elegans (Kreis, 1929) Lorenzen, 1977
levis Rieger & Ott, 1971
curvatus Gerlach, 1956
biggi (Gerlach, 1965) Lorenzen, 1977
oxyicerca (de Man, 1888) Lorenzen, 1977
furcatum (Juario, 1974) Lorenzen, 1977
acrilabiatum (De Coninck & Schuurmans Stekhoven, 1933) Lorenzen, 1977
invagiferum (Platt, 1973) Lorenzen, 1977
setosum (Bütschli, 1874) Lorenzen, 1977
conicum (Filipjev, 1922) Lorenzen, 1977
adiectus (Schulz, 1932) Wieser, 1956
spirum (Gerlach, 1959) Lorenzen, 1977
rusticum (Kreis, 1929) Lorenzen, 1977
circulum (Vitiello, 1971) Lorenzen, 1977
fallax (Lorenzen, 1972) Lorenzen, 1977
microscopiculum (Gerlach, 1953) Lorenzen, 1977
vicinum (Riemann, 1966) Lorenzen, 1977
albigensis (Riemann, 1966) Hopper, 1968
buetschlii Schuurmans Stekhoven, 1935
paralorum (Vitiello, 1971) Lorenzen, 1977
tenuispiculum (Ditlevsen, 1918) Lorenzen, 1977
longicaudatum (Filipjev, 1922) Lorenzen, 1977
Dasynemella
conica Gerlach, 1956
albaensis Warwick & Platt, 1973

- pselionemoides* (Gerlach, 1953) De Coninck, 1965
Dasynemoides
longicollis Gerlach, 1952
Deltanema
parvum Kreis, 1929
Demonema
rapax Cobb, 1894
Dentatonema
crassum Kreis, 1928
Denticulella
polydonta (Schulz, 1932) Wieser, 1954
stygia (Gerlach, 1952) Wieser, 1954
Deontolaimus
papillatus de Man, 1880
Deontostoma
arcticum (Ssaweljev, 1912) Filipjev, 1916
lobatum (Steiner, 1916) Platonova, 1962
Desmodora
varioannulata (Kreis, 1928) Lorenzen, 1981
tenuidentata Kreis, 1963
poseidoni Steiner, 1916
cincta (Cobb, 1920) Lorenzen, 1981
polychaeta Allgén, 1929
mediterranea Lorenzen, 1981
coniseta (Schuurmans Stekhoven, 1950) Lorenzen, 1981
microchaeta (Allgén, 1929) Lorenzen, 1981
conica Vitiello, 1971
longiseta (Schuurmans Stekhoven, 1950) Lorenzen, 1981
roscoffensis Luc & De Coninck, 1959
porifera (Wieser, 1951) Lorenzen, 1981
angusticollis Daday, 1901
ditlevensi (Micoletzky, 1922) Lorenzen, 1981
nybellini Allgén, 1954
microchaetoides Allgén, 1933
intermedia Allgén, 1940
scaldensis de Man, 1889
tenuispiculum (Allgén, 1928) Lorenzen, 1976
tenuicauda Allgén, 1932
bullata Steiner, 1916
sanguinea Southern, 1914
papillata Daday, 1901
hirsuta (Chitwood, 1936) Gerlach, 1963
pontica (Filipjev, 1922) Lorenzen, 1981
sphaerica (Kreis, 1928) Lorenzen, 1981
communis (Bütschli, 1874) de Man, 1889
septentrionalis Kreis, 1963
Desmolaimus
bulbulus Lorenzen, 1969
zeelandicus de Man, 1880
zosterae Allgén, 1933
tristis Allgén, 1935
mirabilis Allgén, 1935
longicaudatus Kreis, 1929
Desmoscolex
petalodes Lorenzen, 1972
proboscis Lorenzen, 1972
labiosus Lorenzen, 1969
amaurus Lorenzen, 1972
pusillus Lorenzen, 1969
nanellus (Allgén, 1939) Lorenzen, 1971
vanoysi De Coninck, 1943
adenotrichus Lorenzen, 1969
tenuiseta Filipjev, 1922
fennicus Lorenzen, 1969
minor Schepotieff, 1907
adriaticus Schepotieff, 1907
longisetosus Timm, 1970
lanuginosus Panceri, 1876
italicus Timm, 1970
cosmopolites Timm, 1970
californiensis (Timm, 1970) Gerlach & Riemann, 1973
borealis Kreis, 1963
laevis Kreis, 1928
salcatus Lorenzen, 1972
balticus Lorenzen, 1971
annulatus Schepotieff, 1907
americanus Chitwood, 1936
minutus Claparède, 1863
chaetogaster Greeff, 1869
Dichromadora
microdonta Kreis, 1929
geophila (de Man, 1876) Kreis, 1929
cephalata (Steiner, 1916) Kreis, 1929
hyalocheile De Coninck & Schuurmans Stekhoven, 1933
gracilis (Kreis, 1929) Wieser, 1954
islandica Kreis, 1963
cucullata Lorenzen, 1973
abnormis Gerlach, 1953
strandii Allgén, 1940
scandula Lorenzen, 1966
punctata Schuurmans Stekhoven, 1950
Didelta
scutellatum Vitiello, 1969
longicaudatum Vitiello, 1969
cascudum Gerlach, 1956
Diodontolaimus
sabulosus Southern, 1914
Diplolaimella
ocellata (Bütschli, 1874) Gerlach, 1957
signifera (Paramonov, 1929) Gerlach & Riemann, 1973
stagnosa Lorenzen, 1966
punicea Timm, 1952
monhysterooides Allgén, 1929
Diplolaimelloides
altherri Meyl, 1954
bruciei Hopper, 1970
islandicus (De Coninck, 1943) Meyl, 1954
oschei Meyl, 1954
deconincki (Gerlach, 1951) Meyl, 1954
Diplopeltis
cirrhatus (Eberth, 1863) Cobb, 1891
Diplopeltoides
ornatus (Gerlach, 1950) Gerlach, 1962
Diplopeltula
striata Gerlach, 1956
bulbosa Vitiello, 1972
lucanica Boucher & Hellouët, 1977
minuta Vitiello, 1972
longiceps Gerlach, 1950
laminata Vitiello, 1972
cassidaigniensis Vitiello, 1972
breviceps Gerlach, 1950
asetosa Juario, 1974
incisa (Southern, 1914) Gerlach, 1962
nuda Gerlach, 1956
intermedia Gerlach, 1954
Disconema
sueicum Allgén, 1935
minutum Vitiello, 1969
longicaudatum Vitiello, 1969
longamphida Vitiello, 1969
alaima Filipjev, 1918
Ditlevsenella
danica Filipjev, 1927
inermis Gerlach, 1953
murmanica Filipjev, 1927
Dolicholaimus
marioni de Man, 1888
Dolichosomatum
klatti Allgén, 1951
Doliolaimus
agilis Lorenzen, 1966
Dorylaimopsis
mediterranea Grimaldi-De Zio, 1968
punctata Ditlevsen, 1918

- Dracognomus*
simplex (Gerlach, 1954) Allen & Noffsinger, 1978
- Draconema*
ophicephalum (Claparede, 1863) Hope & Murphy, 1972
micoletzkyi Kreis, 1928
ponticum Filipjev, 1918
solidum Gerlach, 1952
spinosum (Southern, 1914) Kreis, 1938
armatum (Giard & Barrois, 1875) Hope & Murphy, 1972
inarmense (Panceri, 1876) Hope & Murphy, 1972
claparedii (Metschnikoff, 1867) Hope & Murphy, 1972
macrocephalum (Schepotieff, 1907) Wieser, 1954
cephalatum Cobb, 1913
simplex (Gerlach, 1954) Hope & Murphy, 1972
hibernicum (Southern, 1914) Kreis, 1938
arcticum Kreis, 1963
campbelli (Allgén, 1932) Hope & Murphy, 1972
longirostrum (Schepotieff, 1907) Hope & Murphy, 1972
- Echinotheristus*
teutonicus Thun & Riemann, 1967
cimbricus Thun & Riemann, 1967
- Eleutherolaimus*
parastenosoma Allgén, 1933
parasabuliculus (Allgén, 1935) Gerlach, 1958
longus Filipjev, 1922
demanii (Rouville, 1903) Filipjev, 1922
iniquisetosus Schuurmans Stekhoven, 1935
cylindricaudatus (Schuurmans Stekhoven, 1946) Gerlach, 1963
stenosoma (de Man, 1907) Filipjev, 1922
elegans Schuurmans Stekhoven, 1942
duplicatus Schuurmans Stekhoven, 1950
- Elzalia*
mediterranea Vitiello, 1971
- Enchelidiella*
pellucida Allgén, 1954
- Enoploides*
amphioxii Filipjev, 1918
brunetti Gerlach, 1953
tyrrhenicus Brunetti, 1949
cephalophorus (Ditlevsen, 1918) Filipjev, 1927
labiatus (Bütschli, 1874) Filipjev, 1918
caspersi Riemann, 1966
typicus Ssaweljev, 1912
labrostriatus (Southern, 1914) Filipjev, 1921
spiculohamatus Schulz, 1932
longispulosus Vitiello, 1967
brunettii Gerlach, 1953
hirsutus Filipjev, 1918
cirrhatus Filipjev, 1918
- Enoplolaimus*
filiformis (Allgén, 1935) Wieser, 1953
propinquus de Man, 1922
subterraneus Gerlach, 1953
villosus Gerlach, 1953
parapropinquus Allgén, 1949
robustus Gerlach, 1954
zosterae Schulz, 1932
denticulatus Warwick, 1970
regius Hopper, 1962
glaber Brunetti, 1949
conicollis Gerlach, 1952
balensis Skwarra, 1921
attenuatus Brunetti, 1949
psammae Gerlach, 1952
abnormis Kreis, 1928
longicaudatus (Southern, 1914) Filipjev, 1921
litoralis Schulz, 1936
halophilus Ditlevsen, 1928
vulgaris de Man, 1893
- Enoplus*
groenlandicus Ditlevsen, 1926
alatus Ssaweljev, 1912
auriculatus Ssaweljev, 1912
brachyuris Ditlevsen, 1923
constrictus Ditlevsen, 1926
crassus Filipjev, 1916
flagellicaudatus Schuurmans Stekhoven, 1943
littoralis Filipjev, 1918
communis Bastian, 1865
maeoticus Filipjev, 1916
brevis Bastian, 1865
quadridentatus Berlin, 1853
schulzi Gerlach, 1952
obtusocaudatus Eberth, 1863
serratus Ditlevsen, 1926
inermis Bastian, 1865
sphaericus Kreis, 1928
meridionalis Steiner, 1921
michaelseni von Linstow, 1896
paralittoralis Wieser, 1953
striatus Eberth, 1863
- Epacanthion*
mawsoni Warwick, 1977
buetschlii (Southern, 1914) Wieser, 1953
flagellicauda Gerlach, 1956
crassum (Ditlevsen, 1926) Wieser, 1953
durapelle (Kreis, 1929) Wieser, 1953
enoploidiforme (Gerlach, 1953) Warwick, 1970
gorgonocephalum Warwick, 1970
incurvatum (Ditlevsen, 1926) Wieser, 1953
pellucidum (Ssaweljev, 1912) Wieser, 1959
murmanicum (Ssaweljev, 1912) Wieser, 1959
- Epsilonema*
cygnoides (Metschnikoff, 1867) Steiner, 1927
- Ethmolaimus*
multipapillatus Paramonov, 1926
- Eubostrichus*
macintyrei Platt & Zhang, 1982
phalacrus Greeff, 1869
filiformis Greeff, 1869
smo Platt & Zhang, 1982
- Euchromadora*
inflatipiculum Schuurmans Stekhoven, 1943
striata (Eberth, 1863) de Man, 1886
tridentata Allgén, 1929
gaulica Inglis, 1962
strandii Allgén, 1934
vulgaris (Bastian, 1865) de Man, 1886
africana Linstow, 1908
- Eumorpholaimus*
digiticaudatus Schuurmans Stekhoven, 1946
norvegicus Allgén, 1946
sabuliculus Schulz, 1932
- Eurystomina*
ornata (Eberth, 1863) Marion, 1870
cassiterides (Warwick, 1977) Lambshead & Platt, 1979
terricola (de Man, 1907) Filipjev, 1921
stenolaimoides Wieser, 1953
spectabilis (Marion, 1870) Filipjev, 1921
gerlachi Inglis, 1962
filispiculum Gerlach, 1954
filiformis (de Man, 1888) Filipjev, 1921
assimilis (de Man, 1876) de Man, 1888
retrococellata Micoletzky, 1930
pettiti Inglis, 1962
- Eusynonchus*
hirsutus (Cobb, 1894) Platonova, 1970
brevisetosus (Southern, 1914) Platonova, 1970
- Fenestrolaimus*
murmanicus Filipjev, 1927
insulaealbae Filipjev, 1927
profundis Filipjev, 1927
vestitus Gerlach, 1956
anatarcticus (Mawson, 1956) Lorenzen, 1994

- Filipjeva*
parameridionalis Vitiello, 1970
arctica Ditlevsen, 1928
mediterranea Schuurmans Stekhoven, 1950
- Filipjevia*
macrolabiata Kreis, 1928
- Filoncholaimus*
filicaudatus (Ditlevsen, 1926) Filipjev, 1927
- Fusonema*
multilabiatum Kreis, 1928
- Gairleanema*
anagremiae Warwick & Platt, 1973
- Gammannema*
mediterraneum Vitiello, 1970
conicauda Gerlach, 1953
rapax (Ssweljev, 1912) Gerlach, 1964
- Gammarinema*
ligiae Gerlach, 1967
gammati Kinne & Gerlach, 1953
- Gerlachius*
novussetosus Platt & Zhang, 1982
- Gonianchus*
cumbaensis Benwell, 1981
inaequalis Warwick & Platt, 1973
longicaudatus (Ward, 1972)
- Graphonema*
scampae (Coles, 1965) Platt & Warwick, 1988
northumbriæ Warwick & Coles, 1975
- Greeffia*
oxycaudata (Greeff, 1869) Cobb, 1922
beatlei Lorenzen, 1969
comosa Lorenzen, 1969
moppa Schrage, 1972
- Halalaimus*
monstrocaudatus Vitiello, 1970
cirrhatus Gerlach, 1953
delamarei Vitiello, 1970
diplocephalus Filipjev, 1927
horridus Gerlach, 1956
longicollis Allgén, 1932
pachyderma (Filipjev, 1927) Schuurmans Stekhoven, 1935
zenkevitschi Filipjev, 1927
amphistrius Vitiello, 1970
filicorpus Vitiello, 1970
papillifer Gerlach, 1956
lutarus Vitiello, 1970
capitulatus Boucher, 1978
borealis (Steiner, 1916) Lorenzen, 1981
striatus Gerlach, 1956
gracilis de Man, 1888
longicaudatus (Filipjev, 1927) Schneider, 1939
amphidellus Vitiello, 1970
pachyodoroides Vitiello, 1970
ponicus Filipjev, 1922
turbidus Vitiello, 1970
terrestris Gerlach, 1959
isaitshikovi (Filipjev, 1927) Schuurmans Stekhoven, 1935
leptosoma (Southern, 1914) Schuurmans Stekhoven, 1935
- Halanonchus*
papillatus Groza-Rojancovski, 1972
cornutus Vitiello, 1971
zosterae (Allgén, 1933) Gerlach, 1964
- Halaphanolaimus*
longisetosus Allgén, 1928
minutus Schuurmans Stekhoven, 1942
pellucidus Southern, 1914
- Halichoanolaimus*
norvegicus Allgén, 1940
minor Ssweljev, 1912
consimilis Allgén, 1933
robustus (Bastian, 1865) de Man, 1886
lanceolatus Vitiello, 1970
- dolichurus* Ssweljev, 1912
- Halinema*
norwegicum Allgén, 1933
- Haliplectus*
bibulosus (Schulz, 1935) Gerlach, 1957
bickneri Chitwood, 1956
wheeleri Coles, 1965
dorsalis Chitwood, 1956
schulzi (De Coninck, 1943) Gerlach, 1957
- Hapalomus*
terrestris Lorenzen, 1969
- Heterocyatholaimus*
macrolaimus Allgén, 1935
- Hoffmaenneria*
niddensis (Skwarra, 1921) Schneider, 1940
- Hopperia*
massiliensis Vitiello, 1969
- Hypodontolaimus*
inaequalis (Bastian, 1865) de Man, 1886
schuurmansstekhoveni Gerlach, 1951
balticus (Schneider, 1906) Filipjev, 1918
abyssalis Allgén, 1933
heymonsi (Steiner, 1921) Filipjev, 1930
longiseta (Allgén, 1933) Wieser, 1954
mediterraneus Brunetti, 1949
sivertseni Allgén, 1951
setosus (Bütschli, 1874) Wieser, 1954
colesi Inglis, 1962
- Innocuonema*
tentabunda (de Man, 1890) Inglis, 1969
- Ionema*
cobbi (Steiner, 1916) Micoletzky, 1924
coecum Kreis, 1963
boreale Kreis, 1963
- Ixonema*
sordidum Lorenzen, 1971
- Klugea*
pigmentata (Ditlevsen, 1926) Filipjev, 1927
trilabia Filipjev, 1927
filiformis (Allgén, 1929) Filipjev, 1934
polaris (Steiner, 1916) Filipjev, 1927
stenolaima (Wieser, 1954) Wieser, 1954
- Krampia*
opaca (Allgén, 1930) Kreis, 1934
longicollis Allgén, 1939
acropora Ditlevsen, 1921
- Kraspedonema*
octogonata Gerlach, 1954
- Laimella*
minuta Vitiello, 1970
vera Vitiello, 1971
longicauda Cobb, 1920
- Latronema*
aberrans (Allgén, 1934) Gerlach, 1964
orcinum (Gerlach, 1952) Wieser, 1954
- Lauratonema*
adriaticum Gerlach, 1953
reductum Gerlach, 1953
- Lauratonemoides*
originalis (Gerlach, 1956) De Coninck, 1965
- Laxus*
contortus Cobb, 1894
- Ledovitia*
obtusidens (Schuurmans Stekhoven, 1956) Vitiello, 1970
longiseta (Micoletzky, 1930) Wieser, 1953
hirsuta Filipjev, 1927
- Leptolaimoides*
tubulosus Vitiello, 1971
thermastris (Lorenzen, 1966) Vitiello, 1971
- Leptolaimus*
macer Lorenzen, 1972
turridus Timm, 1963
longiseta (Allgén, 1934) De Coninck, 1965
puccinelliae Gerlach, 1959

- leptaleus* Lorenzen, 1972
membranatus (Wieser, 1951) De Coninck, 1965
minutus Vitiello, 1971
mixtus Lorenzen, 1972
nobilis Gerlach, 1956
papilliger de Man, 1876
pumicosus Vitiello, 1971
steineri (Filipjev, 1922) De Coninck, 1965
scotlandicus Jayasree & Warwick, 1977
limiculus Lorenzen, 1972
elegans (Schuurmans Stekhoven & De Coninck, 1933)
 Gerlach, 1958
ampullaceus Warwick, 1970
trichodes (Kreis, 1929) De Coninck, 1965
venustus Lorenzen, 1972
alatus Vitiello, 1971
acicula Lorenzen, 1966
ditlevenseni (Steiner, 1916) De Coninck, 1965
setiger Schuurmans Stekhoven & De Coninck, 1933
timmi Vitiello, 1971
cupulatus Lorenzen, 1972
tenuis Gerlach, 1956
- Leptonemella*
gorgo Gerlach, 1950
parabullata (Allgén, 1929) Gerlach, 1963
aphanothecae Gerlach, 1950
froeyensis (Allgén, 1946) Gerlach, 1963
- Leptosomatides*
steineri Filipjev, 1922
euxinus Filipjev, 1918
- Leptosomatina*
conicaudata Vitiello, 1970
appendixocaudata Allgén, 1957
- Leptosomatum*
bacillatum (Eberth, 1863) Bastian, 1865
arcticum Filipjev, 1916
tetraphthalmum Ssaweljev, 1912
sabangense Steiner, 1915
elongatum Bastian, 1865
longisetosum Schuurmans Stekhoven, 1943
filipjevi Schuurmans Stekhoven, 1950
breviceps Platonova, 1967
gracile Bastian, 1865
subulatum (Eberth, 1863) Bastian, 1865
punctatum (Eberth, 1863) Bastian, 1865
bathybium Allgén, 1954
caecum Ditlevsen, 1923
- Leptosomella*
acrocerca Filipjev, 1927
- Lineolia*
sieboldii (Kölliker, 1845) Gerlach & Riemann, 1974
- Linhomoeus*
paralineatus Allgén, 1933
timmi Inglis, 1963
elongatus Bastian, 1865
ponticus Filipjev, 1922
pilosus (Schuurmans Stekhoven, 1950) Wieser, 1956
paralongicaudatus (Allgén, 1934) Wieser, 1956
macramphis Schuurmans Stekhoven, 1942
hirsutus Bastian, 1865
kornoensis Allgén, 1929
gracilisetosus (Allgén, 1946) Wieser, 1956
brevisetosus Schuurmans Stekhoven, 1946
- Litinium*
parmatum Wieser, 1954
bananum Gerlach, 1956
simplex (Bird, 1995) Lorenzen, 1981
- Longicyatholaimus*
cervoides Vitiello, 1970
longicaudatus (de Man, 1876) Micoletzky, 1924
salciatus Vitiello, 1971
filicaudatus Schuurmans Stekhoven, 1950
capsulatus Vitiello, 1970
- Manunema*
proboscidis Gerlach, 1957
- annulatum* (Vitiello & De Coninck, 1968) Riemann et al., 1971
- Margonema*
longisetosum Allgén, 1935
- Marylynna*
bellula (Vitiello, 1970) Hopper, 1977
choanolaimoides (Schuurmans Stekhoven, 1942) Hopper, 1977
- effilata* (Schuurmans Stekhoven, 1946) Hopper, 1977
- stekhoveni* (Wieser, 1954) Hopper, 1977
complexa (Warwick, 1971) Hopper, 1977
- Megadesmolaimus*
incisus Wieser, 1954
- Megeurystomina*
combesi Luc & De Coninck, 1959
- Mesacanthion*
audax (Ditlevsen, 1918) Filipjev, 1927
longispiculum Gerlach, 1954
tenuicaudatum (Ssaweljev, 1912) De Coninck & Schuurmans Stekhoven, 1933
- ditlevenseni* (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
conicum (Filipjev, 1918) Filipjev, 1927
banale (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
lucifer (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
agubernatus Vitiello, 1971
africantiforme Warwick, 1970
infantile (Ditlevsen, 1930) De Coninck & Schuurmans Stekhoven, 1933
breviseta (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
hirsutum Gerlach, 1953
diplechma (Southern, 1914) Filipjev, 1927
virile (Ditlevsen, 1930) De Coninck & Schuurmans Stekhoven, 1933
majus (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
- Mesacanthoides*
latignathus (Ditlevsen, 1918) Wieser, 1953
caputmedusae (Ditlevsen, 1918) Wieser, 1959
- Metachromadora*
cystoseirae Gerlach & Riemann, 1973
quadribulba Gerlach, 1956
macrourera Filipjev, 1918
remanei Gerlach, 1951
vivipara (de Man, 1907) Allgén, 1928
suecica (Allgén, 1929) Schulz, 1938
scotlandica Warwick & Platt, 1973
- Metacyatholaimus*
cylindribucca (Schuurmans Stekhoven, 1950) Wieser, 1954
hirschi Schuurmans Stekhoven, 1942
- Metacylicolaimus*
effilatus Schuurmans Stekhoven, 1946
filicaudatus (Ditlevsen, 1926) Schuurmans Stekhoven, 1946
flagellicaudatus Schuurmans Stekhoven, 1946
- Metadasynemella*
macrophalla De Coninck, 1942
cassidinensis Vitiello & Haspelagh, 1972
falciphalla Vitiello & Haspelagh, 1972
- Metadesmodora*
amphidiscata Schuurmans Stekhoven, 1942
- Metadesmolaimus*
aduncus Lorenzen, 1972
gelana (Warwick & Platt, 1973)
aversivulva Gerlach, 1953
coronatus Schuurmans Stekhoven, 1950
heteroclitus Lorenzen, 1972
pandus Lorenzen, 1972
varians Lorenzen, 1972
labiosetosus Schuurmans Stekhoven, 1935
tersus (Gerlach, 1956) Lorenzen, 1972

- hamatus* (Gerlach, 1956) Lorenzen, 1972
Metalaimus
gracilis Kreis, 1928
Metalinhomoeus
zosterae Filipjev, 1918
effilatus Schuurmans Stekhoven, 1942
paratypicus Vitiello, 1969
musaecauda Lorenzen, 1966
microsetosus Allgén, 1953
maroeensis Allgén, 1946
gracilis (Kreis, 1929) Wieser, 1956
flagellicaudatus Schuurmans Stekhoven, 1935
tenuicaudatus Allgén, 1935
parvasetosus Vitiello, 1969
pareffilatus Wieser, 1956
filicaudatus (Allgén, 1930)
tristis (Allgén, 1933)
breviseta Schuurmans Stekhoven, 1950
biratus Vitiello, 1969
typicus de Man, 1907
longiseta Kreis, 1929
siliformis (de Man, 1907) Schuurmans Stekhoven, 1935
Metaparoncholaimus
campylocercus (de Man, 1876) De Coninck & Schuurmans Stekhoven, 1933
Metaraeolaimoides
oxystoma De Coninck, 1936
Metepsilonema
hagmeieri (Stauffer, 1924) Steiner, 1927
Metoncholaimus
pristiurus (Zur Strassen, 1894) Filipjev, 1918
albidus (Bastian, 1865) Filipjev, 1918
demani (Zur Strassen, 1894) Filipjev, 1918
anthophorus (Saweljev, 1912) Filipjev, 1918
scanicus (Allgén, 1935) Warwick, 1977
Meylia
alata Gerlach, 1956
lissa Gerlach, 1956
spinosa Gerlach, 1956
Micoletzkyia
mucronata Vitiello, 1970
magna Vitiello, 1970
parelegans Allgén, 1954
elegans Ditlevsen, 1926
Microlaimus
criminalis Rieger & Ott, 1971
ostracion Schuurmans Stekhoven, 1935
cyatholaimoides de Man, 1922
acuticaudatus De Coninck & Schuurmans Stekhoven, 1933
crassiceps Gerlach, 1953
compridus Gerlach, 1956
citrus Gerlach, 1959
borealis Steiner, 1916
arenicola Schulz, 1938
allgeni Gerlach, 1950
zosterae Allgén, 1930
spirifer Warwick, 1970
dentatus Allgén, 1935
marinus (Schulz, 1932) Schuurmans Stekhoven & De Coninck, 1933
cuanensis Platt, 1973
conothelis (Lorenzen, 1973) Jensen, 1978
acinaces Warwick & Platt, 1973
robustidens Schuurmans Stekhoven & De Coninck, 1933
teutonicus Riemann, 1966
parahonestus Gerlach, 1950
pygmaeus Meyl, 1954
honestus de Man, 1922
undulatus Gerlach, 1953
turgofrons Lorenzen, 1972
texianus Chitwood, 1951
monstruosus Gerlach, 1953
tenuicollis Gerlach, 1952
donsi Allgén, 1935
punctulatus Gerlach, 1950
paraborealis Allgén, 1940
microseta Gerlach, 1953
inermis Ditlevsen, 1923
tenuispiculum de Man, 1922
honestoides Meyl, 1954
lazonus Vitiello, 1971
macrocirculus Gerlach, 1950
globiceps de Man, 1880
Minolaimus
cervoides Vitiello, 1970
lineatus Vitiello, 1970
Molgolaimus
demani Jensen, 1978
Monhystera
paraambigua Allgén, 1933
macquariensis Allgén, 1929
filiformis Bastian, 1865
disjuncta Bastian, 1865
macrolabiata Kreis, 1928
microphthalma de Man, 1880
minuta Filipjev, 1929
natans (Bastian, 1865) Bütschli, 1874
paramacramphis Meyl, 1954
parasimplex De Coninck, 1943
paradisjuncta De Coninck, 1943
parelegantula De Coninck, 1943
refringens Schuurmans Stekhoven, 1935
rotundicapitata Filipjev, 1922
sabulicola Allgén, 1929
socialis Bütschli, 1874
multisetosa Meyl, 1955
attenuata Filipjev, 1922
collaris Filipjev, 1922
fasciculata Skwarra, 1921
filicaudata Allgén, 1929
anophthalma Lorenzen, 1966
gracilicauda Schuurmans Stekhoven, 1950
heterolaima Schuurmans Stekhoven, 1950
islandica De Coninck, 1943
izhorica Filipjev, 1929
paraambiguoides Allgén, 1932
luisae Schuurmans Stekhoven, 1935
macramphis Filipjev, 1929
barentsi Steiner, 1916
capitata Schuurmans Stekhoven, 1942
cartaromanae Meyl, 1954
ampullocauda Paramonov, 1926
droebachiensis Allgén, 1931
parva (Bastian, 1865) Bütschli, 1874
cephalophora de Man, 1876
tenuis Schuurmans Stekhoven, 1950
longicapitata Filipjev, 1922
trichura Allgén, 1930
vulgaris de Man, 1880
similis Bütschli, 1873
elegantula Schuurmans Stekhoven, 1935
Monoposthia
mirabilis Schulz, 1932
costata (Bastian, 1865) de Man, 1889
mielcki Steiner, 1916
arctica Allgén, 1954
Morlaixia
obscura Vincx & Gourbault, 1988
Nannolaimoides
effilatus (Boucher, 1976) Platt & Warwick, 1988
Nannolaimus
volutus Gerlach, 1956
fusus Gerlach, 1956
Nemanema
rectoresectum Kreis, 1928
sabulicola (Allgén, 1939) Wieser, 1953
cylindratacaudatum (de Man, 1922) Filipjev, 1927

- rotundicaudatum* (Ditlevsen, 1926) Filipjev, 1946
filiforme (Filipjev, 1918) Kreis, 1926
- Nemella*
ocellata Cobb, 1920
- Neochromadora*
izhorica (Filipjev, I., 1929) Schuurmans Stekhoven, 1935
paractecta Blome, 1974
trichophora (Steiner, 1921) Gerlach, 1951
sabulicola (Filipjev, 1918) Wieser, 1954
poecilosomoides (Filipjev, 1918) Micoletzky, 1924
munita Lorenzen, 1971
craspedota (Steiner, 1916) Micoletzky, 1924
coudenhovei Wieser, 1956
brevisetosa Wieser, 1954
amembranata Wieser, 1954
poecilosoma (de Man, 1893) Micoletzky, 1924
angelica Riemann, 1976
paramunita Boucher, 1976
tecta Gerlach, 1951
- Neotonchus*
vittius (Warwick, 1971)
corcundus (Gerlach, 1956) Wieser & Hopper, 1966
cupulatus Vitiello, 1970
meeki Warwick, 1971
pseudocorcundus Vitiello, 1971
boucheri Platt, 1982
warwicki (Platt, 1982)
interruptus (Warwick, 1971)
votadinii (Warwick, 1971)
cuanensis (Platt, 1982)
- Nijhoffia*
longicaudata Allgén, 1935
- Notochaetosoma*
killieri Warwick, 1977
- Nudora*
thorakista (Schulz, 1935) Gerlach & Riemann, 1973
bipapillata Platt, 1973
gourbaultae Vanreusel & Vincx, 1989
steineri (Steiner, 1921) Wieser, 1954
- Octonchus*
marinus (Schulz, 1932) Clark, 1961
- Odontophora*
rectangula Lorenzen, 1972
marina Bütschli, 1874
armata (Ditlevsen, 1918) Allgén, 1929
parangustilaima Wieser, 1956
parasetosa (Allgén, 1929) Allgén, 1929
paratenuicaudata Allgén, 1942
phalarata Lorenzen, 1972
angustilaimus (Filipjev, 1918) Allgén, 1929
longicaudata Schuurmans Stekhoven & De Coninck, 1933
fatisca Vitiello, 1971
tenuicaudata Allgén, 1935
ornata Lorenzen, 1972
wieseri Luc & De Coninck, 1959
villoti Luc & De Coninck, 1959
longisetosa (Allgén, 1928) Allgén, 1929
exharena Warwick & Platt, 1973
setosa (Allgén, 1929) Allgén, 1929
- Odontophoroidea*
paramonhystra Lambshead, 1982
- Omicronema*
truncatum Schuurmans Stekhoven, 1950
nidrosiense Allgén, 1933
- Onchium*
robustum Gerlach, 1965
ocellatum Cobb, 1920
conicaudatum (Allgén, 1935) Wieser, 1956
parocellum (Allgén, 1940) Gerlach & Riemann, 1973
- Oncholaimellus*
mediterraneus Schuurmans Stekhoven, 1942
calvadosicus de Man, 1890
- Oncholaimus*
- oxyuris* Ditlevsen, 1911
marinus Schulz, 1932
vesicarius (Wieser, 1959) Rachor, 1969
onchouris Ditlevsen, 1928
groenlandicus Ditlevsen, 1926
medius Kreis, 1928
brachycercus de Man, 1889
steineri Ditlevsen, 1928
skawensis Ditlevsen, 1921
donsi (Allgén, 1947) Wieser, 1953
attenuatus Dujardin, 1845
viridis Bastian, 1865
dujardinii de Man, 1876
lanceolatus Vitiello, 1970
septentrionalis Filipjev, 1927
brevicaudatus Filipjev, 1918
aegypticus Steiner, 1921
spissus Allgén, 1935
serratus Allgén, 1935
campylocercoides De Coninck & Schuurmans Stekhoven, 1933
scanicus (Allgén, 1935) Wieser, 1953
conicauda Filipjev, 1929
curvicauda Allgén, 1957
steinboecki Ditlevsen, 1928
paroxyuris Schuurmans Stekhoven, 1942
- Onyx*
perfectus Cobb, 1891
sagittarius Gerlach, 1950
- Oxyonchus*
polaris (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
pachylabiatus Schuurmans Stekhoven, 1946
hamatus (Steiner, 1916) Filipjev, 1927
acantholaimus (Ssweljev, 1912) Filipjev, 1927
dentatus (Ditlevsen, 1918) Filipjev, 1927
dubius (Filipjev, 1918) De Coninck & Schuurmans Stekhoven, 1933
- Oxystomina*
filicauda (Kreis, 1929) Wieser, 1953
nidrosiensis Allgén, 1933
alpha Chitwood, 1937
islandica (De Coninck, 1943) Wieser, 1953
novozemelica (Filipjev, 1927) Wieser, 1953
nuda (Filipjev, 1927) Wieser, 1953
pulchella Vitiello, 1970
unguiculata Schuurmans Stekhoven, 1935
brevicaudata Gerlach & Riemann, 1974
alpatovi (Filipjev, 1927) Wieser, 1953
elongata (Bütschli, 1874) Filipjev, 1921
oxycaudata (Ditlevsen, 1926) Filipjev, 1921
asetosa (Southern, 1914) Filipjev, 1921
tenuis Thorne, 1939
clavicauda (Filipjev, 1918) Filipjev, 1921
- Pandolaimus*
kornoensis (Allgén, 1929) Jensen, 1976
viscosa (Allgén, 1935) Jensen, 1976
latilaimus (Allgén, 1929) Schuurmans Stekhoven, 1935
- Paracanthonchus*
multitubifer Timm, 1961
spectabilis Allgén, 1931
macrodon (Ditlevsen, 1918) Micoletzky, 1924
breviseta (Schuurmans Stekhoven, 1950) Hope & Murphy, 1972
longicaudatus Warwick, 1971
thaumasius (Schulz, 1932)
platti Vadhyar, 1980
heterodontus (Schulz, 1932)
caecus Micoletzky, 1924
longus Allgén, 1934
elongatus (de Man, 1907) Micoletzky, 1924
steueri (Micoletzky, 1922) Micoletzky, 1924
sunesoni (Allgén, 1942) Wieser, 1954
tyrrhenicus (Brunetti, 1949) Gerlach, 1953

- uniformis* (Schuurmans Stekhoven, 1950) Wieser, 1954
paradoxus Allgén, 1929
nannodontus (Schulz, 1932) Wieser, 1954
microdon (Ditlevsen, 1918) Wieser, 1954
micoletzkyi Schuurmans Stekhoven, 1943
major (Kreis, 1928) Wieser, 1954
filipjevi Micoletzky, 1924
stekhoveni Wieser, 1954
bipapillatus Kreis, 1928
angulatus (Schuurmans Stekhoven, 1950) Wieser, 1954
inglisi Coles, 1965
barka Inglis, 1962
- Paracomesoma*
dubium (Filipjev, 1918) Schuurmans Stekhoven, 1950
- Paracyatholaimoides*
labiisetosus Riemann, 1966
multispiculatus Gerlach, 1953
- Paracyatholaimus*
proximus (Bütschli, 1874) Micoletzky, 1924
arcticus Kreis, 1963
dubiosus (Bütschli, 1874) Micoletzky, 1921
oistospiculoides (Allgén, 1935) Wieser, 1954
parasaveljevi Allgén, 1935
pentodon Riemann, 1966
separatus Wieser, 1954
occultus Gerlach, 1956
intermedius (de Man, 1880) Micoletzky, 1922
- Paracyclicolaimus*
brevisetosus Platonova, 1970
- Paradesmodora*
cephalata Schuurmans Stekhoven, 1950
- Paradraconema*
spinulosum (Southern, 1914) Allen & Noffsinger, 1978
- Paraegialoalaimus*
silicaudatus Allgén, 1934
- Paralinhomoeus*
strandibrevicaudatus Allgén, 1934
linurus (Ssaweljev, 1912)
viscosus (Allgén, 1929) Allgén, 1934
trichurus (Allgén, 1933)
lineatus (Ditlevsen, 1918) Allgén, 1959
tenuicaudatus (Bütschli, 1874) de Man, 1907
staurensis Allgén, 1940
uniovarium Warwick, 1970
conicaudatus (Allgén, 1930)
lepturus (de Man, 1907) de Man, 1907
brevibucca Schuurmans Stekhoven, 1950
intermedius (Allgén, 1929) Wieser, 1956
amphiliabius Schuurmans Stekhoven, 1942
bocki Schuurmans Stekhoven, 1946
brevicaudatus Schuurmans Stekhoven, 1950
caxinus Vitiello, 1969
clavicaudatus Schuurmans Stekhoven, 1950
filiformis (Filipjev, 1918)
ilenensis (Allgén, 1933) Gerlach, 1963
anteporus Vitiello, 1969
deconincki Groza-Rojancovski, 1972
- Paralongicyatholaimus*
zosterae (Allgén, 1933) Lorenzen, 1972
mastigodes Schuurmans Stekhoven, 1950
minutus Warwick, 1971
macramphis Lorenzen, 1972
- Paramesacanthion*
marei Warwick, 1970
truncum Vitiello, 1971
tricuspidis (Schuurmans Stekhoven, 1950) Wieser, 1953
klugei (Filipjev, 1927) Wieser, 1953
hirsum Warwick, 1970
catellum Boucher, 1970
oxycephalum (Ditlevsen, 1926) Wieser, 1953
- Paramicrolaimus*
acanthus Jayasree & Warwick, 1977
- Paramonohystera*
- micramphis* Schuurmans Stekhoven, 1950
levicula (Lorenzen, 1973) Lorenzen, 1977
paranormandica (Micoletzky, 1922) Wieser, 1956
pellucida (Cobb, 1920) Wieser, 1956
riemanni (Platt, 1973)
megacephala (Steiner, 1916) Filipjev, 1918
elliptica (Filipjev, 1918) Wieser, 1956
- Parapinnanema*
harveyi Warwick & Coles, 1975
- Pararaeolaimus*
nudus (Gerlach, 1951) Timm, 1961
- Parasaveljevia*
clavicauda (Filipjev, 1927) Wieser, 1953
- Paraspheerolaimus*
paradoxus (Ditlevsen, 1918) Wieser, 1956
- Paraterschellingia*
brevicaudata (Kreis, 1924) Schuurmans Stekhoven, 1935
fusiformis Gerlach, 1951
- Parathalassoolaimus*
sabulicola Allgén, 1929
- Paratricoma*
papillifer Gerlach, 1956
pratensis Lorenzen, 1969
- Paratuerkiana*
comes (Tsalolikhin, 1972) Platonova, 1970
- Pareudesmoscolex*
laciniosus Lorenzen, 1969
papillosus (Schulz, 1935) Weischer, 1962
pratensis Lorenzen, 1969
- Pareurystomina*
armorica Luc & De Coninck, 1959
acuminata (de Man, 1889) Gerlach, 1952
silicaudata Allgén, 1934
flagellicaudata Schuurmans Stekhoven, 1946
parapugentis Vitiello, 1970
tenuicauda Schuurmans Stekhoven, 1950
biserialis Schuurmans Stekhoven, 1946
scilloniensis Warwick, 1977
typica Micoletzky, 1930
tenuissima (Filipjev, 1927) Filipjev, 1946
- Parodontophora*
quadristicha (Schuurmans Stekhoven, 1950) Timm, 1963
breviseta (Schuurmans Stekhoven, 1950) Timm, 1963
- Paroxystomina*
asymmetrica Micoletzky, 1924
- Pelagonema*
omalum Vitiello, 1970
obtusicauda Filipjev, 1918
tenue Kreis, 1928
norvegicum Allgén, 1939
propinquum Allgén, 1929
simplex Cobb, 1894
angusticavatum Kreis, 1928
- Perilinhomoeus*
longisetosus Schuurmans Stekhoven, 1950
- Phanoderma*
cocksi Bastian, 1865
etha Gadea, 1960
laticolle (Marion, 1870) Marion, 1875
conicaudatum (Steiner, 1916) Steiner, 1921
annulocaudatum Allgén, 1939
parasiticum Ditlevsen, 1926
islandicum Ditlevsen, 1926
longisetum Allgén, 1939
tuberculatum (Eberth, 1863) Bastian, 1865
campbelli Allgén, 1928
brachyductum Wieser, 1953
rigidum Schuurmans Stekhoven, 1946
steineri Ditlevsen, 1918
nasutum Schuurmans Stekhoven, 1950
albidum Bastian, 1865
robustum Allgén, 1957
setigerum (Marion, 1870) Filipjev, 1918

- Phanodermella*
flagellicaudata Vitiello, 1970
Phanodermopsis
caudata Filipjev, 1927
reducta Filipjev, 1927
longicauda (Schuurmans Stekhoven, 1950) Filipjev, 1927
groenlandica Ditlevsen, 1926
Pierrickia
decasetosa Vitiello, 1970
aequalis (Gerlach, 1956) Vitiello, 1970
Platycoma
cephalata Cobb, 1894
Platycomopsis
nuchapilosa Vitiello, 1970
mesjatzevi Filipjev, 1927
effilata Schuurmans Stekhoven, 1946
cobbii Ditlevsen, 1926
Polygastrophora
alexandriae (Schuurmans Stekhoven, 1943) Wieser, 1953
hexabulba (Filipjev, 1918) Kreis, 1926
maior Schulz, 1932
attenuata de Man, 1922
Polysigma
fuscum Gerlach, 1956
Pomponema
loticum Lorenzen, 1972
tesselatum Wieser & Hopper, 1967
debile Lorenzen, 1972
cotylophorum (Steiner, 1916) Lorenzen, 1972
elegans Lorenzen, 1972
reductum Warwick, 1970
sedecima Platt, 1973
tautraense (Allgén, 1933) Lorenzen, 1972
lineatum (Gerlach, 1953) Wieser, 1959
cyatholaimoides (Allgén, 1929) Hope & Murphy, 1972
multipapillatum (Filipjev, 1922) Wieser, 1954
compactum Lorenzen, 1972
clavicaudatum (Schuurmans Stekhoven, 1935) Lorenzen, 1972
astrodes Lorenzen, 1972
ammophilum Lorenzen, 1972
Pontonema
ditlevensi (Kreis, 1934) Cobb & Steiner, 1934
vulgare (Bastian, 1865) Filipjev, 1916
macrolaimus (Southern, 1914) Filipjev, 1921
simile (Southern, 1914) Filipjev, 1921
brevicaudatooides (Menzel, 1920) Gerlach & Riemann, 1974
parpapilliferum (Micoletzky, 1924) Cobb & Steiner, 1934
arcticum (Kreis, 1928) Cobb & Steiner, 1934
parocellatum Wieser, 1954
teissieri (Vitiello, 1967) Lorenzen, 1981
filicaudatus (Kreis, 1928) Lorenzen, 1981
zernovi (Filipjev, 1916) Cobb & Steiner, 1934
donsi (Allgén, 1932) Cobb & Steiner, 1934
papilliferum (Filipjev, 1916) Cobb & Steiner, 1934
balticum (Schulz, 1932) Cobb & Steiner, 1934
Praeacanthonchus
inglisi (Coles, 1965) Platt & Warwick, 1988
opheliae (Warwick, 1970) Gerlach & Riemann, 1973
kreisi (Allgén, 1929) Gerlach & Riemann, 1973
punctatus (Bastian, 1865) Micoletzky, 1924
Procamacolaimus
acer Gerlach, 1954
dolichostylum Gerlach, 1953
Prochaetosoma
primitivum (Steiner, 1916) Micoletzky, 1922
Prochromadora
exigua (Ditlevsen, 1928) Wieser, 1954
megodonta Filipjev, 1922
orleji (de Man, 1880) Filipjev, 1922
magna (Schulz, 1935) Wieser, 1954
spitzbergensis Gerlach, 1965
Prochromadorella
septempapillata Platt, 1973
neapolitana (de Man, 1876) Micoletzky, 1924
macrocellata Wieser, 1951
ditlevensi (de Man, 1922) Wieser, 1954
acridentata (Schulz, 1932) Wieser, 1954
actuaria Vitiello, 1971
ambigua (Ditlevsen, 1928) Wieser, 1954
brachyura Schuurmans Stekhoven, 1950
norwegica (Allgén, 1932) Wieser, 1951
attenuata (Gerlach, 1952) Lorenzen, 1972
maculata (Ditlevsen, 1918) Wieser, 1954
mediterranea (Micoletzky, 1922) Micoletzky, 1924
paramucroonta (Allgén, 1929) Wieser, 1951
tenuicauda Gerlach, 1954
subterranea Gerlach, 1953
quinquepapillata (Schuurmans Stekhoven, 1935) Wieser, 1954
obtusidens (Schuurmans Stekhoven & Adam, 1931) Wieser, 1951
longicaudata (Kreis, 1929) Lorenzen, 1972
Prooncholaimus
eberthi (Filipjev, 1918) Micoletzky, 1924
megastoma (Eberth, 1863) Micoletzky, 1924
banyulensis Inglis, 1962
Prosphaerolaimus
eurypharynx Filipjev, 1918
Protochromadora
scampae (Coles, 1965) Inglis, 1969
mediterranea (Allgén, 1942) Inglis, 1969
Pselionema
simplex De Coninck, 1942
simile De Coninck, 1942
richardi De Coninck, 1942
minutum Vitiello & Haspeslagh, 1972
parasimplex Vitiello, 1971
annulatum (Filipjev, 1922) Cobb, 1933
deconincki Vitiello & Haspeslagh, 1972
longissimum Gerlach, 1953
longiseta Ward, 1974
Pseudocella
citronicauda (Wieser, 1954) Platonova, 1962
saveljevi (Filipjev, 1927) Platonova, 1962
gracilis Platonova, 1967
coeca (Ssaweljev, 1912) Filipjev, 1927
trichodes (Leuckart, 1849) Filipjev, 1927
cavernicola (Wieser, 1954) Platonova, 1962
elegans (Ditlevsen, 1926) Wieser, 1953
Pseudodella
norvegica Allgén, 1947
Pseudometachromadora
longilaima (Schuurmans Stekhoven, 1950) Timm, 1952
papillata (Schuurmans Stekhoven, 1950) Timm, 1952
Pseudoncholaimus
saveljevi (Filipjev, 1927) Kreis, 1934
Pseudonchonus
deconincki Warwick, 1969
northumbriensis Warwick, 1969
norwegicus Allgén, 1933
donsi Allgén, 1948
gerlachi Warwick, 1969
longus Allgén, 1949
Pseudosteineria
horrida (Steiner, 1916) Wieser, 1956
Pterygonema
platti Zhang, 1983
cambriensis Ward, 1973
Ptycholaimellus
monodon (Schuurmans Stekhoven, 1942) Hopper, 1969
ponticus (Filipjev, 1922) Gerlach, 1955
Punctodora

- ratzeburgensis* (Linstow, 1876) Filipjev, 1929
- Quadricoma*
frontalis (Gerlach, 1952) Timm, 1970
greeffii (Reinhard, 1881) Filipjev, 1922
loricata Filipjev, 1922
media (Reinhard, 1881) Filipjev, 1922
steineri Filipjev, 1922
suecica (Allgén, 1930) Timm, 1970
vittata (Lorenzen, 1969) Timm, 1970
trigintatres Timm, 1970
tenuis (Steiner, 1916) Filipjev, 1922
eurycricus (Filipjev, 1922) Timm, 1970
crassicauda Timm, 1970
cobbi (Steiner, 1916) Filipjev, 1922
scanica (Allgén, 1935) Timm, 1970
- Rhabditis*
marina Bastian, 1865
ehrenbaumi Schuurmans Stekhoven, 1935
- Rhabdocoma*
brevicauda Schuurmans Stekhoven, 1950
macrura Cobb, 1920
riemanni Jayasree & Warwick, 1977
- Rhabdodemania*
minor (Southern, 1914) Baylis & Daubney, 1926
major (Southern, 1914) Baylis & Daubney, 1926
mediterranea Boucher, 1971
obtusicauda (Allgén, 1954) Hope & Murphy, 1972
imer Warwick & Platt, 1973
pontica Platonova, 1965
laticauda (Ditlevsen, 1926) Baylis & Daubney, 1926
gracilis (Ditlevsen, 1918) Baylis & Daubney, 1926
coronata Gerlach, 1952
brevicaudata (Schuurmans Stekhoven, 1942)
 Lorenzen, 1981
- Rhadinema*
flexile Cobb, 1920
- Rhips*
paraornata Platt & Zhang, 1982
- Rhynchonema*
longituba Gerlach, 1953
deconincki Vitiello, 1967
lyngei (Allgén, 1940) Gerlach, 1953
scutatum Lorenzen, 1972
brevituba Gerlach, 1953
- Richtersia*
inaequalis Riemann, 1966
norvegica Allgén, 1940
elongata Schuurmans Stekhoven, 1950
demanii Schuurmans Stekhoven, 1935
tenuis (Kreis, 1929) Schuurmans Stekhoven, 1935
collaris Steiner, 1916
- Sabatieria*
conicauda Vitiello, 1970
pulchra (Schneider, 1906) Riemann, 1970
stekhoveni Vitiello, 1970
strigosa Lorenzen, 1972
demanii Schuurmans Stekhoven, 1935
dodecaspillata (Kreis, 1929) Wieser, 1954
granifer Wieser, 1954
elongata Jayasree & Warwick, 1977
vasicola Vitiello, 1970
lyonessa Warwick, 1977
pisinna Vitiello, 1970
armata Gerlach, 1952
macramphis Lorenzen, 1972
longispinosa Lorenzen, 1972
longisetosa (Kreis, 1929) Wieser, 1954
longicaudata Filipjev, 1922
breviseta Schuurmans Stekhoven, 1935
ornata (Ditlevsen, 1918) Filipjev, 1922
praedatrix de Man, 1907
abyssalis (Filipjev, 1918) Filipjev, 1922
celtica Southern, 1914
- Sarsonia*
leptosoma (de Man, 1893) Gerlach, 1967
- Saveljevia*
curvidens Filipjev, 1927
xiphonchus Filipjev, 1927
spissignatha (Allgén, 1940) Wieser, 1953
kolaensis Filipjev, 1927
cornuta Gerlach, 1956
- Scaprella*
brevicaudata Gerlach, 1953
tenuicaudata Gerlach, 1956
- Setoplectus*
gerlachi Vitiello, 1971
- Setosabatieria*
hilarula (de Man, 1922) Platt, 1985
- Sigmophoranema*
monstrosum (Gerlach, 1956) Hope & Murphy, 1972
litorale (Schulz, 1938) Gerlach, 1951
rufum (Cobb, 1933) Hope & Murphy, 1972
- Siphonolaimus*
niger de Man, 1893
cylindricaudatus Schuurmans Stekhoven, 1942
cobbi Riemann, 1966
elongatus (Schuurmans Stekhoven, 1950) Vitiello, 1971
roervikensis Allgén, 1946
nigricans (Cobb, 1894) Filipjev, 1918
obtusicaudatus Allgén, 1930
weismanni (Zur Strassen, 1904) Steiner, 1916
ewensis Warwick & Platt, 1973
anticomoides (Allgén, 1933) Filipjev, 1918
- Solenolaimus*
obtusus Cobb, 1894
- Southerniella*
allometrica Vitiello, 1971
brevicauda (Schuurmans Stekhoven, 1942) Wieser, 1956
conicauda (Schuurmans Stekhoven, 1950) Wieser, 1956
zostericola (Allgén, 1929) Wieser, 1956
stenolaima (Steiner, 1916) Wieser, 1956
- Sphaerocephalum*
crassicauda Filipjev, 1918
chabaudi Inglis, 1962
bulbiferum Schuurmans Stekhoven, 1943
- Sphaerolaimus*
brevicollis Ssweljev, 1912
abyssorum Allgén, 1933
dispar Filipjev, 1918
glaphyrus Vitiello, 1971
latilaimus Allgén, 1958
macrocirculoides Wieser, 1954
paragracilis Vitiello, 1971
cuneatus Paramonov, 1929
minutus Vitiello, 1971
papillatus Kreis, 1929
balticus Schneider, 1906
islandicus Ditlevsen, 1926
paradoxus (Ditlevsen, 1918) Wieser, 1956
macrocirculus Filipjev, 1918
gracilis de Man, 1876
hirsutus Bastian, 1865
ostreae Filipjev, 1918
uncinatus Freudenhammer, 1970
polaris (Filipjev, 1946) Wieser, 1956
makrolasmus Schulz, 1932
profundus Vitiello, 1971
striatus Allgén, 1935
tenuis (Kreis, 1924) Schuurmans Stekhoven, 1935
maeoticus Filipjev, 1922
pacificus Allgén, 1947
- Spiliiphera*
dolichura de Man, 1893
hirsuta Gerlach, 1956
gracilicauda de Man, 1893
spinicauda Vitiello, 1970
minuta Vitiello, 1972

- elegans* Bastian, 1865
Spilophorella
papillata Kreis, 1929
paradoxa (de Man, 1888) Filipjev, 1917
euxina Filipjev, 1918
candida Gerlach, 1951
Spiramphinema
microcephalon (Schuurmans Stekhoven, 1942)
Wieser, 1956
Spirinia
laevis (Bastian, 1865) Gerlach, 1963
flagellata (Vitiello, 1971) Lorenzen, 1981
schneideri (Villot, 1875) Gerlach, 1963
parasitifera (Bastian, 1865) Gerlach, 1963
granulata (Allgén, 1929) Gerlach, 1963
gerlachi (Luc & De Coninck, 1959) Gerlach, 1963
subulicola (Filipjev, 1918) Gerlach, 1963
Steineria
longicaudata Vitiello, 1971
polychaetoides Gerlach, 1951
pontica Groza-Rojancovski, 1972
pilosa (Cobb, 1914) Micoletzky, 1921
Steineridora
loricata (Steiner, 1916) Inglis, 1969
adriatica (Daday, 1901) Inglis, 1969
Stephanolaimus
paraflevensis Gerlach, 1953
jayasreei Platt, 1983
filicaudatus Schuurmans Stekhoven, 1946
spartinae Lorenzen, 1969
flevensis Schuurmans Stekhoven, 1935
Symplocostoma
reisingeri Ditlevsen, 1928
medium Filipjev, 1927
tenuicolle (Eberth, 1863) Wieser, 1953
leptolaimus Filipjev, 1927
murmanicum Ssweljev, 1912
majus Filipjev, 1927
karense Filipjev, 1927
acutum (Cobb, 1920) Wieser, 1953
ponticum Filipjev, 1918
papillatum Kreis, 1928
Synodontium
impar (Ssweljev, 1912) Wieser, 1956
Synonchiella
micramphis (Schuurmans Stekhoven, 1950) Gerlach, 1964
riemanni Warwick, 1970
minuta Vitiello, 1970
Synonchium
depressum Gerlach, 1954
obtusum Cobb, 1920
Synonchus
armatus (Ditlevsen, 1923) Filipjev, 1927
intermedius Allgén, 1939
longisetosus (Southern, 1914) Filipjev, 1927
fusciculus Cobb, 1894
acuticaudatus (Jägerskjöld, 1901) Filipjev, 1916
dubius (Wieser, 1954) Gerlach & Riemann, 1974
roscovianus (Villot, 1875) Filipjev, 1927
murmanicus Filipjev, 1927
Synonema
triadenophorum (Steiner, 1921) Wieser, 1954
obtusicaudatum (de Man, 1889) Gerlach, 1963
Syringolaimus
filicaudatus Vitiello, 1970
striatocaudatus de Man, 1888
brevicaudatus Micoletzky, 1922
caspersi Gerlach, 1951
Syringonomus
typicus Hope & Murphy, 1969
Tarvaiia
angusta Gerlach, 1953
donsi Allgén, 1934
Terschellingia
communis de Man, 1888
longicaudata de Man, 1907
gourbaultae Austen, 1989
pontica Filipjev, 1918
similis Allgén, 1933
parva Vitiello, 1969
capitata Vitiello, 1969
Thalassironus
britannicus de Man, 1889
Thalassoolaimus
egregius Steiner, 1916
septentrionalis Filipjev, 1927
rotundicaudatus Filipjev, 1927
longicaudatus Vitiello, 1970
brevicaudatus Vitiello, 1970
tenuis Allgén, 1935
oxycauda Filipjev, 1927
paramontemari Vitiello, 1970
mediterraneus Vitiello, 1970
pirum Lorenzen, 1969
filiformis Allgén, 1933
aceratus Vitiello, 1970
tardus de Man, 1893
Thalassomonhystera
venusta (Lorenzen, 1979) Jacobs, 1987
parva (Bastian, 1865) Jacobs, 1987
Theristus
maior Gerlach, 1956
meyli Riemann, 1966
pellucidus Allgén, 1939
monstruosus Gerlach, 1954
norwegicus (Allgén, 1933) Wieser, 1956
pyronasi Gerlach, 1965
velox (Bastian, 1865)
obtusicephalus Schuurmans Stekhoven, 1950
lineatus Gerlach, 1965
scanicus Allgén, 1949
flevensis Schuurmans Stekhoven, 1935
pictus Gerlach, 1951
littoralis Filipjev, 1922
leptosoma Allgén, 1950
heteroscopiculoides Gerlach, 1953
inermis Gerlach, 1953
anisotrichus Lorenzen, 1972
tenuicaudatus Allgén, 1951
strandi Allgén, 1934
roscoffensis Vitiello, 1967
ensifer Gerlach, 1951
gracilis (de Man, 1876) Filipjev, 1918
parasetosus (Allgén, 1928)
latissimus Filipjev, 1922
heteroscopiculum (Allgén, 1932) Gerlach, 1953
interstitialis Warwick, 1970
longus Platt, 1973
otoplanobius Gerlach, 1951
pertenuis Schuurmans Stekhoven, 1935
subulicola (Filipjev, 1918) Wieser, 1956
stichotricha Schuurmans Stekhoven, 1950
bipunctatus (Schneider, 1906) Filipjev, 1929
complexus Jayasree & Warwick, 1977
longisetosus (De Coninck & Schuurmans Stekhoven, 1933) Lorenzen, 1977
aculeatus Schulz, 1935
blandicor Ranchor, 1971
bastiani Gerlach & Riemann, 1973
acer Bastian, 1865
acriformis (de Man, 1922) Gerlach & Riemann, 1973
denticulatus Warwick, 1970
Thoracostoma
steineri Micoletzky, 1922
zolae Marion, 1870
penduli Allgén, 1954
multipapillatum Allgén, 1957
amphacanthus (Grube, 1855) Bütschli, 1874
brachylobatum Allgén, 1954

- sivertensi* Allgén, 1957
siliformis (Eberth, 1863) Filipjev, 1921
hispanicum Platonova, 1970
coronatum (Eberth, 1863) Marion, 1870
Thoracostomopsis
galeata Filipjev, 1927
ditlevenseni Filipjev, 1927
barbata Ditlevsen, 1918
doveae Warwick, 1970
longissima Filipjev, 1927
Trefusia
litoralis (Allgén, 1932) De Coninck & Schuurmans Stekhoven, 1933
silicauda Allgén, 1933
zostericola Allgén, 1933
helgolandica Riemann, 1966
longicauda de Man, 1893
pseudilitoralis Vitiello, 1970
filum Schuurmans Stekhoven, 1942
Trichethermolaimus
hirsutus (Gerlach, 1956) Platt, 1982
Trichotheristus
mirabilis (Schuurmans Stekhoven & De Coninck, 1933) Wieser, 1956
psammooides (Warwick, 1970)
Tricoma
brevirostris (Southern, 1914) Steiner, 1916
longirostris (Southern, 1914) Steiner, 1916
suecica Allgén, 1930
strandii (Allgén, 1939) Gerlach, 1958
steineri de Man, 1922
septuaginta Schuurmans Stekhoven, 1942
profunda (Schepotieff, 1907) Stauffer, 1924
pellucida Allgén, 1935
maxima (Schepotieff, 1907) Allgén, 1942
islandica Kreis, 1963
bacescui (Paladian & Andriescu, 1963) Timm, 1970
bergensis (Schepotieff, 1907) Filipjev, 1922
platycephala Filipjev, 1922
elongata (Panceri, 1876) Stauffer, 1924
polydesma (Southern, 1914) Steiner, 1916
litoralis (Allgén, 1934) Timm, 1970
norvegica (Schepotieff, 1907) Stauffer, 1924
nematooides (Greeff, 1869) Stauffer, 1924
adelpha (Greeff, 1869) Stauffer, 1924
cincta Cobb, 1894
Trileptium
subterraneum (Gerlach, 1952) Cobb, 1933
parisetum Warwick & Platt, 1973
Triodontolaimus
acutus (Villot, 1875) de Man, 1893
Tripyla
cornuta Skwarra, 1921
setifera Bütschli, 1873
glomerans Bastian, 1865
Triploides
gracilis (Ditlevsen, 1918) Filipjev, 1927
omblaica Micoletzky, 1924
marinus (Bütschli, 1874) de Man, 1886
Trissonchulus
benepapillosus (Schulz, 1935) Gerlach & Riemann, 1974
oceanus Cobb, 1920
obtusus (Schuurmans Stekhoven, 1935) Inglis, 1961
Trochanus
carinatus Boucher & Bovée, 1972
Tubolaimoides
tenuicaudatus (Allgén, 1934) Gerlach, 1963
Valvaelaimus
euxinus (Filipjev, 1918) Lorenzen, 1977
Vasculonema
caeruleum (Eberth, 1863) Kreis, 1928
conicaudatum Kreis, 1928
Viscosia
minudonta Vitiello, 1970
separabilis (Wieser, 1953) Lorenzen, 1981
stenolaima Filipjev, 1927
longidentatus (Schuurmans Stekhoven, 1931) Lorenzen, 1981
brevidentatus (Vitiello, 1967) Lorenzen, 1981
elegans (Kreis, 1924) Lorenzen, 1981
parva Kreis, 1929
palmae Schuurmans Stekhoven, 1942
profundus (Vitiello, 1970) Lorenzen, 1981
langrunensis (de Man, 1890) Filipjev, 1918
brachylaima Filipjev, 1927
elongata Filipjev, 1922
rusticus (Kreis, 1929) Lorenzen, 1981
papillosa (Eberth, 1863) Kreis, 1934
viscosa (Bastian, 1865) de Man, 1890
minor Filipjev, 1918
glabra (Bastian, 1865) de Man, 1890
abyssorum (Allgén, 1933) Wieser, 1953
filipjevi Paramonov, 1929
macramphida Chitwood, 1951
macrorhopalocerca Kreis, 1932
megalaaima (Ditlevsen, 1928) Hope & Murphy, 1972
cobbi Filipjev, 1918
diodon (Wieser, 1951) Lorenzen, 1981
Wieseria
glandulosa (Kreis, 1929) Gerlach, 1956
clavata Gerlach, 1956
pica Gerlach, 1956
hispida Vitiello, 1972
inaequalis Gerlach, 1956
leptura Vitiello, 1972
longiseta (Allgén, 1947) Gerlach, 1956
Xenella
suecica Allgén, 1935
Xyala
striata Cobb, 1920
longicaudata Ward, 1972
barbata Ward, 1972
riemannii Boucher & Helléouët, 1977

Nematoda (parasitic)

Compiled by David I. Gibson

The list of parasitic nematode species was derived from the Host-Parasite Data-base and Catalogue maintained by the Parasitic Worms Division, Department of Zoology, The Natural History Museum. Some attempt has been made to restrict the list to valid taxa and to use that latest nomenclature. However, the many, usually unrecognisable and often larval, names listed under *Ascaris* have been listed separately.

The list was kindly commented upon by Dr F. Moravec and Dr H.-P. Fagerholm.

NEMATODA, PARASITIC

- Acanthocheilus*
rotundatus (Rudolphi, 1819)
Agamascaris
septentrionalis Kreis, 1952
Agamoneema
sp.
Amidostomum
acutum (Lundahl, 1848)
Anguillicola
crassus Kuwahara, Niimi & Itagaki, 1974
Anisakis
pegreffii Campana-Rouget & Biocca, 1955
physeteris (Baylis, 1923)
schupakovi Mosgovoi, 1951
simplex (Rudolphi, 1809)
simplex C of Nascetti *et al.* (1986)
typica (Diesing, 1860)
ziphidarum Paggi *et al.*, 1988

<i>Ascarophis</i>	
<i>arctica</i> Poljansky, 1952	<i>spinosissimus</i> von Linstow, 1905
<i>capelanus</i> Nikolaeva & Naidenova, 1964	<i>uncinatus</i> Molin, 1858
<i>crassicollis</i> Dollfus & Campana-Rouget, 1956	<i>Echinomermella</i>
<i>filiformis</i> Poljansky, 1952	<i>matsi</i> Jones & Hagen, 1987
<i>morrhuæ</i> van Beneden, 1871	<i>Eustrongylides</i>
<i>mullus</i> Naidenova & Nikolaeva, 1968	sp.
<i>Bulbocephalus</i>	<i>Fellicola</i>
<i>pseudopenei</i> Vassiliades & Diaw, 1978	<i>longispiculus</i> Petter & Køie, 1993
<i>Camallanus</i>	<i>Goezia</i>
<i>lacustris</i> (Zoega, 1776)	<i>annulata</i> (Molin, 1859)
<i>truncatus</i> Rudolphi, 1814	<i>ascaroides</i> (Goeze, 1782)
<i>Capillaria</i>	<i>kollari</i> (Molin, 1859)
<i>anatis</i> (Schrank, 1790)	<i>tricirrata</i> Osmanov, 1940
<i>gracilis</i> (Bellingsham, 1840)	<i>Halocercus</i>
<i>wickinsi</i> Ogden, 1965	<i>delphini</i> (Baylis & Daubney, 1925)
<i>Contracaecum</i>	<i>kleinenbergi</i> Delamure, 1951
<i>filiforme</i> (Stossich, 1904)	<i>lagenorhynchi</i> Baylis & Daubney, 1925
<i>mulli</i> (Wedl, 1855)	<i>ponticus</i> Delamure, 1946
<i>osculatum</i> (Rudolphi, 1802)	<i>taurica</i> Delyamure, 1942
<i>papilligerum</i> (Creplin, 1846)	<i>Heliconema</i>
<i>pedum</i> (Deslongchamps, 1824)	<i>heliconema</i> Travassos, 1919
<i>rudolphii</i> Hartwich, 1964	<i>Heligmus</i>
<i>septentrionale</i> Kreis, 1955	<i>longicirrus</i> Dujardin, 1845
<i>Cosmocephalus</i>	<i>Hysterothylacium</i>
<i>obvelatus</i> (Creplin, 1825)	<i>aduncum</i> (Rudolphi, 1802)
<i>Crassicauda</i>	<i>arnoglossi</i> Petter & Maillard, 1988
<i>anthonyi</i> Chabaud, 1962	<i>auctum</i> (Rudolphi, 1802)
<i>boopis</i> Baylis, 1920	<i>bifidatum</i> Petter & Maillard, 1988
<i>carbonelli</i> Raga <i>et al.</i> , 1988	<i>collare</i> (Cobb, 1929)
<i>grampicola</i> Johnston & Mawson, 1941	<i>cornutum</i> (Stossich, 1904)
<i>Cristitectus</i>	<i>corrugatum</i> Deardorff & Overstreet, 1981
<i>congeri</i> Petter, 1970	<i>cyclopteri</i> (Kreis, 1952)
<i>Cucullanus</i>	<i>fabri</i> (Rudolphi, 1819)
<i>attenuata</i> (Molin, 1858)	<i>increscens</i> (Molin, 1858)
<i>bioccai</i> Orecchia & Paggi, 1987	<i>incurvum</i> (Rudolphi, 1819)
<i>campanae</i> Lebre & Petter, 1984	<i>legendrei</i> (Dollfus, 1933)
<i>carbonelli</i> Campos, Carbonell & Rodriguez-Babio, 1993	<i>reliquens</i> Norris & Overstreet, 1975
<i>cirratus</i> Müller, 1777	<i>rhacodes</i> (Deardorff & Overstreet, 1978)
<i>dodsworthi</i> Barreto, 1922	<i>rigidum</i> (Rudolphi, 1809)
<i>dogieli</i> Krotas, 1959	<i>Ichthyasakis</i>
<i>elongatus</i> Smedley, 1933	<i>monodi</i> Gendre, 1928
<i>gendrei</i> Campana-Rouget, 1957	<i>Ichthyascaris</i>
<i>heterochrous</i> Rudolphi, 1802	<i>mediterraneus</i> (Lebre & Petter, 1983)
<i>hians</i> (Dujardin, 1845)	<i>Ichthyobronema</i>
<i>longicollis</i> (Stossich, 1899)	<i>hamulatum</i> (Moulton, 1931)
<i>longispiculum</i> De Oliveira Rodrigues <i>et al.</i> , 1973	<i>Ichthyofilaria</i>
<i>lophii</i> Campana-Rouget & Chabaud, 1956	<i>bergenensis</i> (Wulker, 1930)
<i>micropapillatus</i> Tornquist, 1931	<i>Johnstonmawsonia</i>
<i>mugili</i> Belous, 1965	<i>murenophidis</i> (Campana-Rouget, 1955)
<i>murenophidis</i> Campana-Rouget, 1957	<i>Kathlania</i>
<i>orthagorisci</i> (Rudolphi, 1819)	<i>leptura</i> (Rudolphi, 1819)
<i>parvus</i> Tornquist, 1931	<i>Mawsonascaris</i>
<i>praecinctus</i> (Dujardin, 1845)	<i>laymani</i> (Mozgovoy, 1950)
<i>pulcherimus</i> Barreto, 1918	<i>pastinaceae</i> (Rudolphi, 1819)
<i>rotundata</i> (Molin, 1859)	<i>Onchophora</i>
<i>truttae</i> Fabricius, 1794	<i>melanocephala</i> (Rudolphi, 1819)
<i>Cystidicola</i>	<i>Paracapillaria</i>
<i>farionis</i> Fischer, 1798	<i>gibsoni</i> Moravec, 1987
<i>Cystoopsis</i>	<i>Paracuarria</i>
<i>scomber</i> Zlatev, 1936	<i>adunca</i> (Creplin, 1846)
<i>Desmidocerella</i>	<i>tridentata</i> (Linstow, 1877)
sp.	<i>Paranisakiopsis</i>
<i>Dichelyne</i>	<i>coelorhynchi</i> Yamaguti, 1941
<i>abbreviatus</i> (Rudolphi, 1819)	<i>Paranisakis</i>
<i>adriaticus</i> (Tornquist, 1931)	<i>monodi</i> (Gendre, 1928)
<i>elongatus</i> (Tornquist, 1931)	<i>squatinae</i> Baylis, 1923
<i>mauritanicus</i> (Gendre, 1928)	<i>Parascarophis</i>
<i>minutus</i> (Rudolphi, 1819)	<i>sphyrnae</i> Campana-Rouget, 1955
<i>pomadasysi</i> Vassiliades & Petter, 1981	<i>Philometra</i>
<i>tripapillatus</i> (Gendre, 1927)	<i>globiceps</i> (Rudolphi, 1819)
<i>Dipetalonema</i>	<i>obturans</i> (Pennant, 1886)
<i>sirocauda</i> (Leidy, 1858)	<i>onchorhynchi</i> Kuitinen-Ekbaum, 1933
<i>Echinocephalus</i>	<i>ovata</i> (Zeder, 1803)
	<i>riscita</i> Skrjabin, 1917
	<i>rubra</i> (Leidy, 1856)

saltatrix Ramachandran, 1973
sanguinea (Rudolphi, 1819)
sombresoxaxis Nikolaeva & Naidenova, 1964
tauridica Ivashkin, Naidenova, Kovaleva &
 Khromova, 1971
translucida Walton, 1927
Philometroides
oveni Parukhin, 1975
Phocascaris
cystophorae Berland, 1963
phocae Host, 1932
Piscicapillaria
baylisi Moravec, 1987
Porrocaecum
sulcatum (Rudolphi, 1819)
Proleptus
acutus Dujardin, 1845
obtusus Dujardin, 1845
robustus (van Beneden, 1871)
Pseudalius
inflexus (Rudolphi, 1808)
Pseudanisakis
baylisi Gibson, 1973
Pseudocapillaria
adriatica (Nikolaeva & Naidenova, 1964)
bainae (Justine & Radujkovic, 1988)
salvelini (Polyansky, 1952)
Pseudoproleptus
 sp.
Pseudoterranova
bulbosa (Cobb, 1888)
decipiens (Krabbe, 1878)
krabbei Paggi *et al.*, 2000
Raphidascaris
acus (Bloch, 1779)
adelinae (Condorelli-Francaviglia, 1898)
Rhabdochona
 sp.
Skrjabinalius
cryptocephalus Delyamure, 1942
guevarai Gallego & Selva, 1979
Spinitectoides
berlandi Petter, 1969
Spinitectus
oviflagellis Fourment, 1883
Stenurus
globicephala Baylis & Daubney, 1925
minor (Kuhn, 1829)
ovatus (von Linstow, 1910)
Streptocara
crassicauda (Creplin, 1829)
Terranova
chiloscyllii Johnston & Mawson, 1951
galeocerdonis (Thwaite, 1927)
rochalimai (Pereira, 1935)
Tetrameres
fissispina (Diesing, 1861)

ACANTHOCEPHALA

Compiled by David I. Gibson

The list of acanthocephalan species was derived from the Host-Parasite Data-base and Catalogue maintained by the Parasitic Worms Division, Department of Zoology, The Natural History Museum. Some attempt has been made to restrict the list to valid taxa and to use that latest nomenclature.

The list was kindly commented upon by Prof. C.R. Kennedy and Dr Z.M. Dimitrova.

PHYLUM ACANTHOCEPHALA

Acanthocephaloidea
distinctus Golvan, 1969
geneticus (Buron, Renaud & Euzet, 1985)
incrassatus (Molin, 1858)
propinquus (Dujardin, 1845)
Acanthocephalus
anguillae (Müller, 1780)
clavula Dujardin, 1845
lucii (Müller, 1777)
Acanthogyrus
lizae Orecchia, Paggi & Radujkovic, 1988
Andracantha
phalacrococaris (Yamaguti, 1939)
tunitae (Weiss, 1914)
Arhytmorhynchus
longicollis (Villot, 1875)
Aspersantis
johni (Baylis, 1929)
Bolbosoma
capitatum (von Linstow, 1880)
physeteris Cubanov, 1952
vasculosum (Rudolphi, 1819)
Breizacanthus
chabaudi Golvan, 1969
golvani Gaevskaja & Shukhgalter, 1984
irenae Golvan, 1969
ligur Paggi, Orecchia & Della Setta, 1975
Corynosoma
magdaleni Montreuil, 1958
semerme (Forssell, 1904)
strumosum (Rudolphi, 1802)
Echinorhynchus
armoricanus Golvan, 1969
bothniensis Zdzitowiecki & Valtonen, 1987
brayi Wayland, Sommerville & Gibson, 1999
calloti Golvan, 1969
cinctulus Porta, 1905
 = *E. borealis* Linstow, 1901
gadi Zoega in Müller, 1776
gadi sp. I of Väinölä Valtonen & Gibson (1994)
gadi sp. II of Väinölä Valtonen & Gibson (1994)
gadi sp. III of Väinölä Valtonen & Gibson (1994)
trachyrinci Wayland, Gibson & Sommerville, 1997
Euzetacanthus
simplex (Rudolphi, 1810)
Floridosentis
mugilis (Machado, 1951)
Golvanacanthus
blennii Paggi & Orecchia, 1972
problematicus Mordvinova & Parukhin, 1978
Gorgorhynchus
robertdolfusi Golvan, 1956
Illiosentis
africanus (Golvan, 1955)
Leptorhynchoides
plagicephalus (Westrum, 1821)
Metechinorhynchus
salmonis (Müller, 1780)
truttae (Schrank, 1788)
Neoechinorhynchus
agilis (Rudolphi, 1819)

rutili (Müller, 1780)
tylosuri Yamaguti, 1939
Nerhadinorhynchus
atlanticus Gaevskaja & Nigmatulin, 1977
myctophumi Mordinova, 1988
Paracanthocephaloides
chabanaudi (Dollfus, 1951)
Plagiorhynchus
charadrii (Yamaguti, 1939)
charadriicola (Dollfus, 1953)
crassicollis (Villot, 1875)
phippsi Kostylev, 1922
ponticus Lisitsina, 1992
Polymorphus
botulus (Van Cleave, 1916)
contortus (Bremser, 1821)
minutus (Zeder, 1800)
Pomphorhynchus
laevis (Zoega in Müller, 1776)
terreticollis (Rudolphi, 1809)
Rhadinorhynchus
cadenati (Golvan & Houin, 1964)
lintoni Cable & Linderoth, 1963
pristis (Rudolphi, 1802)
saltatrix Troncy & Vassiliades, 1973
tenuicornis (Linton, 1891)
Serrasentis
socialis (Leidy, 1851)
Solearhynchus
solea (Porta, 1905)
Southwellina
hispida (Van Cleave, 1925)
Telosentis
exiguus (von Linstow, 1901)
molini Van Cleave, 1923
Yamagutisentis
sp.

CYCLIOPHORA

Compiled by Chris S. Emblow

Cycliophorans are the most recently discovered animal phyla (Funch & Kristensen 1995). They are minute sessile organisms that live on the mouthparts of the Dublin Bay prawn or Norway lobster *Nephrops norvegicus*.

A lack of morphological data has made the true phylogenetic relationship of Cycliophora difficult to determine, however examination of its ribosomal RNA (Winnepenningckx *et al.* 1998) has revealed a close relationship with acanthocephalans and rotifers.

The phylum is at present represented by one species which has been recorded from Denmark, Sweden, the Faeroes Islands and the Firth of Clyde, UK (O'Reilly 1997). It appears likely its distribution is related to that of its host.

References

- Funch, P. & Kristensen, R. M. 1995. Cycliophora is a new phylum with affinities to Entoprocta and Ectoprocta. *Nature* **378**: 711-714.
O'Reilly, M. G. 1997. Cycliophora. In: *The species directory of the marine fauna and flora of the British Isles and surrounding seas*, Howson, C. & Picton, B. E., (eds). Belfast and Ross-on-Wye, Ulster Museum and the Marine Conservation Society 508 pp.
Winnepenningckx, B. M. H., Backeljau, T. & Kristensen, R. M. 1998. Relations of the new phylum Cycliophora. *Nature* **393**: 636-637.

PHYLUM CYCLIOPHORA

Class Eucyliophora

Order Symbiida

Family Symbibidae

Symbion

pandora Funch & Kristensen 1995

A

ECHIURA

Compiled by Jacob van der Land

The European list was derived from URMO, which was mainly based on Stephen & Edmonds (1972) with additions from Zoological Record. J. I. Saiz-Salinas checked the list.

References

Stephen, A.C. & S.J. Edmonds, 1972. *The phyla Sipuncula and Echiura*. British Museum (Natural History) London: 1-528.

PHYLUM ECHIURA

Family Bonelliidae

<i>Amalosoma</i>	
<i>eddystonense</i> Stephen, 1956	A
<i>Bonellia</i>	
<i>minor</i> Marion, 1886	AM
<i>suhmi</i> Selenka, 1885	A
<i>viridis</i> Rolando, 1821	AM
<i>Hamingia</i>	
<i>arctica</i> Danielssen and Koren, 1881	A
<i>Maxmulleria</i>	
<i>aulacoferum</i> (Herubel, 1924)	A
<i>faex</i> (Selenka, 1885)	A
<i>gigas</i> (M. Müller, 1852)	M
<i>lankesteri</i> (Herdman, 1898)	A

Family Echiuridae

<i>Echiurus</i>	
<i>abyssalis</i> Skorikow, 1906	AM
<i>echiurus</i> (Pallas, 1767)	A

Family incertae sedis

<i>Sactosoma</i>	
<i>vitreum</i> Danielssen & Koren, 1881	A

Family Thalassematidae

<i>Anelassorhynchus</i>	
<i>indivisus</i> (Sluiter, 1900)	A
<i>semoni</i> (Fischer, 1896)	A
<i>Ochetostoma</i>	
<i>baroni</i> (Greeff, 1879)	A
<i>erythrogrammon</i> Leuckart and Ruppell, 1828	M
<i>Thalassema</i>	
<i>arcassonense</i> Cuenot, 1902	A
<i>elapsum</i> Sluiter, 1912	A
<i>thalasseum</i> (Pallas, 1766)	AM

SIPUNCULA

Compiled by Jacob van der Land

The European list was derived from URMO, which was compiled by Jacob van der Land, based on Cutler (1994). J. I. Saiz-Salinas checked the ERMS list.

References

Cutler, E.B., 1994. *The Sipuncula. Their Systematics, Biology, and Evolution*: i-xvii, 1-453. Cornell Univ. Press, Ithaca, etc.

PHYLUM SIPUNCULA

Family Aspidosiphonidae

<i>Aspidosiphon</i>	
<i>albus</i> Murina, 1967	A
<i>elegans</i> (Chamisso & Eysenhardt, 1821)	
<i>laevis</i> de Quatrefages, 1865	A
<i>mexicanus</i> Murina, 1967	AM
= <i>Golfingia mexicanus</i>	
<i>misakiensis</i> Ikeda, 1904	AM
<i>muellieri</i> Diesing, 1851	AM
<i>steenstrupii</i> Diesing, 1859	A
= <i>Paraspidosiphon steenstrupii</i>	
<i>zinni</i> Cutler, 1969	A

Family Golfingiidae

<i>Golfingia</i>	
<i>elongata</i> (Keferstein, 1862)	AM
<i>iniqua</i> (Sluiter, 1912)	A
<i>margaritacea</i> (Sars, 1851)	A
<i>muricata</i> (Southern, 1913)	A
<i>vulgaris</i> (de Blainville, 1827)	AM

Nephasoma

<i>Nephasoma</i>	
<i>abyssorum</i> (Koren & Danielssen, 1875)	A
<i>capilleforme</i> (Murina, 1973)	A
<i>confusa</i> (Sluiter, 1902)	M
<i>constrictum</i> (Southern, 1913)	AM
<i>constricticervix</i> (Cutler, 1969)	A
<i>diaphanes</i> (Gerould, 1913)	AM
<i>eremita</i> (Sars, 1851)	A
<i>flagiferum</i> (Selenka, 1885)	A
<i>lilljeborgi</i> (Koren & Danielssen, 1880)	A
= <i>Golfingia lilljeborgi</i>	
<i>minutum</i> (Keferstein, 1862)	A
= <i>Golfingia minutum</i>	
<i>rimicola</i> (Gibbs, 1973)	A
<i>wodjanizkii</i> (Murina, 1973)	A

Thysanocardia

<i>Thysanocardia</i>	
<i>procera</i> (Mobius, 1875)	AM
= <i>Golfingia procera</i>	

Family Phascolionidae

<i>Onchesoma</i>	
<i>magnibathrum</i> Cutler, 1969	A
<i>squamatum</i> (Koren & Danielssen, 1875)	AM
<i>steenstrupi</i> Koren & Danielssen, 1875	AM

Phascolion

<i>Phascolion</i>	
<i>caupo</i> Hendrix, 1975	A
<i>convestitum</i> Sluiter, 1902	M
<i>lutense</i> Selenka, 1885	AM
<i>pacificum</i> Murina, 1957	AM
<i>strombus</i> (Montagu, 1804)	AM
= <i>Phascolion brotzkajae</i>	
<i>tuberculatum</i> Theel, 1875	A

Family Phascolosomatidae

<i>Antillesoma</i>	
<i>antillarum</i> (Grube & Oersted, 1858)	A

= *Phascolosoma antillarum*

Apionsoma

<i>Apionsoma</i>	
<i>capitata</i> (Gerould, 1913)	A
= <i>Fisherana capitata</i>	
= <i>Phascolosoma capitata</i>	
<i>murinae</i> (Cutler, 1969)	AM
= <i>Golfingia murinae</i>	

Phascolosoma

granulatum Leuckart, 1828
stephensi (Stephen, 1942)

A M
A M

Family Sipunculidae

Siphonosoma
arcassonense (Cuenot, 1902)
Sipunculus
norvegicus Danielssen, 1869
nudus Linnaeus, 1766
phalloides (Pallas, 1774)

A
A
A M
A

ENTOPROCTA (KAMPTOZOA)

Compiled by Peter J. Hayward

The Entoprocta is a small phylum of rather inconspicuous animals. The two genera of colony-forming species, *Barentsia* and *Pedicellina*, are most frequently reported in faunistic surveys, both are common on seaweeds, hydroids, live or dead mollusc shells, some crab species, and on bryozoans. The non-colonial forms are strictly commensal and are only discovered by careful examination of likely hosts, which include sponges, polychaetes, sipunculans and some echinoderms.

There is a limited literature. All European species were described and figured by Nielsen (1989), and no new species, or newly recorded species, have been described subsequently.

References

Nielsen, C. 1989. Entoprocts. *Synopses of the British Fauna* (new series) 41: 1 - 131

PHYLUM ENTOPROCTA

Order Solitaria

Family Loxosomatidae

Loxosoma
annelidicola (van Beneden & Hesse, 1864)
claparedei Bobin & Prenant, 1953
loxalina Assheton, 1912
monensis Eggleston, 1965
pectinaricola (Franzen, 1962)
rhodinicola Franzen, 1962
saltans Assheton, 1912
significans Nielsen, 1964
singulare Keferstein, 1862
Loxosomella
aripes Nielsen, 1964
atkinsae Bobin & Prenant, 1953
brucei Eggleston, 1965
claviformis (Hincks, 1880)
compressa Nielsen & Ryland, 1961
crassicauda (Salensky, 1877)
discopoda Nielsen & Ryland, 1961
elegans Nielsen, 1964
glandulifera Franzen, 1962
jagei Bobin & Prenant, 1953
fauveli Bobin & Prenant, 1953
harmeri (Schults, 1895)
kefersteini (Claparedé, 1867)
marsypos Nielsen & Ryland, 1961
murmanica (Nilus, 1909)
nitschei (Vigilius, 1882)
nordgaardi Ryland, 1961
obesa (Atkins, 1932)
ornata Nielsen, 1964
phascolo-somata (Vogt, 1876)
polita Nielsen, 1964
scaura Nielsen, 1964
similis Nielsen, 1964
teissieri (Bobin & Prenant, 1953)
varians Nielsen, 1964
Loxomes pilon
perezi Bobin & Prenant, 1953

Order Coloniales

Family Pedicellinidae

Pedicellina
cernua (Pallas, 1774)
hispida Ryland, 1965
nutans Dalyell, 1848

Family Barentiidae

Barentsia
benedeni (Foettinger, 1886)

elongata Jullien & Calvet, 1903
gracilis M. Sars, 1935
matsushi-mana Toriumi, 1951
mutabilis Toriumi, 1951
ramosa (Robertson, 1900)

MOLLUSCA

Compiled by Philippe Bouchet, Jacques Le Renard and Serge Gofas

The Mollusca part of ERMS is based on CLEMAM (Checklist of European Marine Molluscs), a database at the website <http://www.mnhn.fr/base/malaco.html>.

The need to compile a check-list of the European marine molluscs was discussed during the 10th International Malacological Congress (Tübingen, 1989) and a resolution was passed to this effect by the General Assembly of Unitas Malacologica, and Anders Warén (Stockholm) and Philippe Bouchet (Paris) were asked to investigate into its feasibility. The acronym CLEMAM (Check List of the European MArine Molluscs) was chosen because of its similarity to CLOFNAME and CLOFETA, the very successful regional checklists of Fishes compiled under the umbrella of UNESCO. In 1994, an *ad hoc* meeting, funded by Unitas Malacologica, brought together in Paris malacologists representing several regional or national lists of marine molluscs:

- (i) Riccardo Giannuzzi-Savelli, representing the *Catalogo annotato dei molluschi marini del Mediterraneo*, compiled by B. Sabelli, R. Giannuzzi-Savelli and D. Bedulli (1992) for Società Italiana di Malacologia, and published by Libreria Naturalistica Bolognese, Bologna.
- (ii) Elizabeth Platts, representing the *Checklist of British Marine Mollusca*, compiled by D. Heppell and S. Smith (1991), and published by the National Museums of Scotland.
- (iii) Jon-Arne Sneli, representing the *Annotated checklist of marine molluscs of the Norwegian coast and adjacent waters*, compiled by T. Hoisaeter (1986) and published in *Sarsia*, 71: 73-145.

In the following year, electronic versions of these three lists, in addition to a card catalogue of post-1960 names, were merged by Serge Gofas into a single list in text format. The resulting list was then transformed into database format by Jacques Le Renard, and the first CLEMAM was presented at the 12th International Malacological Congress (Vigo, 1995).

Considering the data added since 1995, CLEMAM currently contains ca. 15,000 names and over 3,300 valid species. New literature is permanently scanned for new material. Old literature is also being scanned for synonyms and gaps in taxonomical coverage. As of 1999, illustrations of type material of valid species and synonyms are being added to the backbone of taxonomical and nomenclatural information.

All native species, benthic or planktonic, and introduced species that have become established are listed in CLEMAM. Old and/or unconfirmed records of non-native species are excluded.

CLEMAM records published information only and does not make new taxonomical decisions. When different state-of-the-art taxonomical opinions exist in the recent literature, the taxonomic editor records what he thinks best reflects the needs of the users of the database.

The following manage CLEMAM: Jacques Le Renard (Directeur de Recherches, INRA. E-mail: lerenard@mnhn.fr), Database design and development; Serge Gofas (Curator, MNHN and Lecturer, University

of Malaga. E-mail: gofas@mnhn.fr), Taxonomic editor; and Philippe Bouchet (Senior Curator, MNHN. E-mail: bouchet@mnhn.fr), External liaison.

We thank the following persons and institutions who have permitted the development of CLEMAM: Unitas Malacologica and its secretary (E. Gittenberger, Leiden, NL) and treasurer (J. van Goethem, Brussels, B) for seed money in the early phases of the project; Società Italiana di Malacologia and R. Giannuzzi-Savelli (Palermo, I), J.-A. Sneli and T. Hoisaeter (Trondheim and Bergen, N), E. Platts, D. Heppell and S. Smith (Edinburgh, UK), and A. Guerra (Vigo, SP) for making available electronic versions of published checklists; A. Warén (Stockholm, S) for input especially on northern and Arctic taxa; staff of MNHN for input at various levels: P. Lozouet, A. Foubert.

Within the list genera are listed alphabetically within sub-families although subfamily names have not been included. The on-line database contains more details on phylogeny and nomenclature.

PHYLUM MOLLUSCA

Class Aplacophora

Family Chaetodermatidae

Chaetoderma

- intermedium* Knipowitsch, 1896
- luifredi* (Ivanov in Scarlato, 1987)
- marinae* (Ivanov in Scarlato, 1987)
- nitens* Möbius, 1876
- nitidulum* Lovén, 1844
- pellucidum* Ivanov in Scarlato, 1987
- simplex* Salvini-Plawen, 1971
- strigisquamatum* Salvini-Plawen, 1977
- tetradens* (Ivanov, 1981)

Falcidens

- aequabilis* Salvini-Plawen, 1972
- crossotus* Salvini-Plawen, 1968
- gutturosus* (Kowalewsky, 1901)
- profundus* Salvini-Plawen, 1968
- sagittiferus* Salvini-Plawen, 1968
- sterreri* (Salvini-Plawen, 1967)
- thorensis* Salvini-Plawen, 1971
- vasconiensis* Salvini-Plawen, 1996

Family Limifossoridae

Scutopus

- robustus* Salvini-Plawen, 1970
- ventrolineatus* Salvini-Plawen, 1968

Psilodens

- elongatus* (Salvini-Plawen, 1972)
- tenius* Salvini-Plawen, 1977

Family Prochaetodermatidae

Prochaetodermia

- breve* (Salvini-Plawen, 1999)
- iberogalicum* (Salvini-Plawen, 1999)
- raduliferum* (Kowalewsky, 1901)
- yongei* (Scheltema, 1985)

Spathoderma

- clenchi* (Scheltema, 1985)

Chevroderma

- gauson* (Scheltema, 1985)
- turnerae* (Scheltema, 1985)

Family Dondersiidae

Dondersia

- festiva* Hubrecht, 1888

Ichthyomenia

- ichthyooides* (Pruvot, 1890)

Micromenia

- fodiens* (Schwabl, 1955)
- simplex* Leloup, 1948

Nematomenia

- banyulensis* (Pruvot, 1890)

corallophila (Kowalewsky, 1881)

flavens (Pruvot, 1890)

Stylomenia

salvatori Pruvot, 1899

Squamatoherpia

tricuspidata (Büchinger & Handl, 1996)

Helluoherpia

aegiri (Handl & Büchinger, 1996)

Family Gymnomeniidae

Gymnomenia

pellucida Odhner, 1921

Genitoconia

atriolonga Salvini-Plawen, 1967

rosea Salvini-Plawen, 1967

Wirenia

argentea Odhner, 1921

gonoconata (Salvini-Plawen, 1988)

Family Lepidomeniidae

Lepidomenia

hystrix Marion & Kowalewsky in Fischer, 1885

swedmarki Salvini-Plawen, 1985

Tegulaherpia

myodoryata Salvini-Plawen, 1988

stimulosa Salvini-Plawen, 1983

Family Macellomeniidae

Macellomenia

palifera (Pruvot, 1890)

Family Neomeniidae

Neomenia

carinata Tullberg, 1875

dalyelli (Koren & Danielssen, 1877)

Meromenia

hirondellei (Leloup, 1949)

Family Amphimeniidae

Amphimenia

neapolitana Thiele, 1889

Alexandromenia

crassa Odhner, 1921

grimaldii Leloup, 1946

Paragymnomenia

richardi Leloup, 1947

Family Drepanomeniidae

Drepanomenia

incrusted (Koren & Danielssen, 1877)

Family Pruvotinidae

Pararrhopalia

pruvoti Simroth, 1893

Eleutheromenia

sierra (Pruvot, 1890)

Hypomenia

nierstraszi Lummel, 1930

Pruvotina

impexa (Pruvot, 1890)

Family Proneomeniidae

Proneomenia

desiderata (Kowalewsky & Marion, 1887)

sluiteri Hubrecht, 1880

Dorymenia

sarsii (Koren & Danielssen, 1877)

vagans (Kowalewsky & Marion, 1887)

Entonomenia

atlantica Leloup, 1948

Family Rhopalomeniidae

Rhopalomenia

aglaopheniae (Kowalewsky & Marion, 1887)

Pruvotia

sopita (Pruvot, 1891)

Family Simrothiellidae

Simrothiella

borealis Odhner, 1921

margaritacea (Koren & Danielssen, 1877)

minima (Nierstrasz, 1903)

- Biserramenia**
psammobionta Salvini-Plawen, 1967
- Uncimenia**
neapolitana Nierstrasz, 1903
- Family Strophomeniidae**
- Anamenia*
borealis (Koren & Danielssen, 1877)
gorgonophila (Kowalewsky, 1880)
- Strophomenia**
lacazei Pruvot, 1899
- Class Monoplacophora**
- Family Neopilinidae**
- Laevipilina*
rolani Warén & Bouchet, 1990
- Micropilina*
minuta Warén, 1989
- Rokopella*
brummeri Goud & Gittenberger, 1993
euglypta (Dautzenberg & Fischer H., 1897)
- Veleropilina*
reticulata (Seguenza G., 1876)
zografi (Dautzenberg & Fischer H., 1896)
- Class Polyplacophora**
- Family Leptochitonidae**
- Lepidopleurus*
cajetanus (Poli, 1791)
- Leptochiton*
africanus (Nierstrasz, 1906)
algesirensis (Capellini, 1859)
alveolus (Lovén, 1846)
arcticus (Sars G.O., 1878)
asellus (Gmelin, 1791)
bedullii Dell'Angelo & Palazzi, 1986
cancellatus (Sowerby G.B. II, 1840)
cimicooides (Monterosato, 1879)
compostellanum (Carmona Zalvide & Urgorri, 1999)
gascognensis Kaas & Van Belle, 1985
geronensis Kaas & Van Belle, 1985
leloupi (Kaas, 1979)
odhneri (Bergenhayn, 1931)
pseudogloriosus (Strack, 1991)
rarinotus (Jeffreys, 1883)
sarsi Kaas, 1981
scabridus (Jeffreys, 1880)
tenuis Kaas, 1979
thalattius Kaas & Van Belle, 1985
xanthus (Kaas & Van Belle, 1990)
- Family Hanleyidae**
- Hanleya*
hanleyi (Bean in Thorpe, 1844)
nagelfar (Lovén, 1846)
- Family Ischnochitonidae**
- Ischnochiton*
albus (Linné, 1767)
dolii (Van Belle & Dell'Angelo, 1998)
exaratus (Sars G.O., 1878)
obtusus (Carpenter in Pilsbry, 1893)
rissoi (Payraudeau, 1826)
usticensis (Dell'Angelo & Castriota, 1999)
vanellei Kaas, 1985
- Connexochiton*
platynomenus Kaas, 1979
- Callistochiton*
pachylasmae (Monterosato, 1878)
- Callochiton*
calcatus Dell'Angelo & Palazzi, 1994
septemvalvis (Montagu, 1803)
- Lepidochitona*
canariensis (Thiele, 1909)
cinerea (Linné, 1767)
corrugata (Reeve, 1848)
furtiva (Monterosato, 1879)
iberica Kaas & Van Belle, 1981
- monterosatoi* Kaas & Van Belle, 1981
piceola (Shuttleworth, 1853)
simrothi (Thiele, 1902)
stroemfelti (Bergenhayn, 1931)
- Tonicella*
marmorea (Fabricius O., 1780)
rubra (Linné, 1767)
- Chaetopleura*
angulata (Spengler, 1797)
- Family Mopaliidae**
- Placiophorella*
atlantica (Verrill & Smith, 1882)
- Family Chitonidae**
- Chiton*
corallinus (Risso, 1826)
hululensis (Smith E.A. in Gardiner, 1903)
olivaceus Spengler, 1797
phaseolinus Monterosato, 1879
- Family Acanthochitonidae**
- Acanthochitona*
crinita (Pennant, 1777)
discrepans (Brown, 1827)
fascicularis (Linné, 1767)
- Class Gastropoda**
- Family Patellidae**
- Patella*
caerulea Linné, 1758
cander d'Orbigny, 1840
ferruginea Gmelin, 1791
intermedia Murray in Knapp, 1857
nigra da Costa, 1771
piperata Gould, 1846
rustica Linné, 1758
ulyssiponensis Gmelin, 1791
vulgata Linné, 1758
- Anstes*
pellucida (Linné, 1758)
- Family Nacellidae**
- Cellana*
rota (Gmelin, 1791)
- Family Acmaeidae**
- Acmaea*
virginea (Müller O.F., 1776)
- Pectinodonta*
maxima (Dautzenberg, 1925)
- Family Lepetidae**
- Lepeta*
caeca (Müller O.F., 1776)
- Iothia*
fulva (Müller O.F., 1776)
- Propilidium*
exiguum (Thompson W., 1844)
- Family Lottiidae**
- Erginus*
rubellus (Fabricius O., 1780)
- Tectura*
testudinalis (Müller O.F., 1776)
- Family Cocculinidae**
- Cocculina*
leptoglypta Dautzenberg & Fischer H., 1897
- Coccopigya*
lata (Warén, 1996)
spinigera (Jeffreys, 1883)
viminensis (Rocchini, 1990)
- Family Bathysciadiidae**
- Bathysciadium*
costulatum (Locard, 1898)
xylophagum Warén & Carozza in Warén, 1997
- Xenodonta*
bogasoni Warén, 1993
- Family Lepetellidae**

- Lepetella**
barrajoni Dantart & Luque, 1994
espinosae Dantart & Luque, 1994
laterocompressa (de Rayneval & Ponzi, 1854)
sierrai Dantart & Luque, 1994
- Bogia**
labronica (Bogi, 1984)
- Family Addisoniidae**
Addisonia
excentrica (Tiberi, 1855)
- Family Choristellidae**
Choristella
nofronii McLean, 1992
- Family Osteopeltidae**
Osteopelta
ceticola Warén, 1989
- Family Pseudococculinidae**
Amphiplica
concentrica (Thiele, 1909)
Copulabyschia
corrugata (Jeffreys, 1883)
- Pilus**
conicus (Verrill, 1884)
Kaiparapelta
askewi McLean & Harasewych, 1995
- Family Neritidae**
Nerita
sanguinolenta Menke, 1829
Smaragdia
souverbiana (Montrouzier, 1863)
viridis (Linné, 1758)
- Family Fissurellidae**
Fissurella
nubecula (Linné, 1758)
Diodora
dorsata (Monterosato, 1878)
gibberula (Lamarck, 1822)
graeca (Linné, 1758)
italica (Defrance, 1820)
producta (Monterosato, 1880)
ruppelli (Sowerby G.B. I, 1835)
Emarginula
adriatica Costa O.G., 1829
christiana Piani, 1985
crassa Sowerby J., 1813
divae (van Aartsen & Carrozza, 1995)
fissura (Linné, 1758)
hazardii (Payraudeau, 1826)
multistriata Jeffreys, 1882
octaviana Coen, 1939
punctulum Piani, 1980
pustula Thiele in Küster, 1913
rosea Bell T., 1824
sicula Gray J.E., 1825
solidula Costa O.G., 1829
tenera Locard, 1892
tuberculosa Libassi, 1859
Cranopsis
asturiana (Fischer P., 1882)
Fissurepta
crossei (Dautzenberg & Fischer H., 1896)
granulosa Jeffreys, 1883
microphyma (Dautzenberg & Fischer H., 1896)
Puncturella
alicei (Dautzenberg & Fischer H., 1897)
noachina (Linné, 1771)
profundi (Jeffreys, 1877)
Zeidora
naufragia Watson, 1883
- Family Scissurellidae**
Scissurella
costata d'Orbigny, 1824
Anatoma
- aspera* (Philippi, 1844)
crispata (Fleming, 1828)
richardi (Dautzenberg & Fischer H., 1896)
umbilicata (Jeffreys, 1883)
- Sinezona**
cingulata (Costa O.G., 1861)
semicostata (Burnay & Rolán, 1990)
- Family Haliotidae**
Haliotis
coccinea Reeve, 1846
pustulata subsp. *cruenta* Reeve, 1846
tuberculata Linné, 1758
- Family Trochidae**
Trochus
erythreus Brocchi, 1821
Clanculus
berthelotii (d'Orbigny, 1840)
corallinus (Gmelin, 1791)
cruciatus (Linné, 1758)
jussieui (Payraudeau, 1826)
Clelandella
miliaris (Brocchi, 1814)
Jujubinus
baudoni (Monterosato, 1891)
dispar Curini-Galletti, 1982
exasperatus (Pennant, 1777)
gravinæ (Dautzenberg, 1881)
guanchus Curini-Galletti, 1985
karpathoensis Nordsieck, 1973
montagui (Wood W., 1828)
poppei Curini-Galletti, 1985
pseudogravinae Nordsieck, 1973
rusrurianus (Weinkauff, 1868)
striatus (Linné, 1758)
tumidulus (Aradas, 1846)
unidentatus (Philippi, 1844)
vexationis Curini-Galletti, 1990
- Gibbula**
adansonii (Payraudeau, 1826)
adriatica (Philippi, 1844)
albida (Gmelin, 1791)
ardens (Von Salis, 1793)
candei (d'Orbigny, 1844)
cineraria (Linné, 1758)
delgadensis Nordsieck, 1982
deversa Milaschewitsch, 1916
divaricata (Linné, 1758)
drepanensis (Brugnone, 1873)
fanulum (Gmelin, 1791)
guttadauri (Philippi, 1836)
leucophæa (Philippi, 1836)
magus (Linné, 1758)
nivosa Adams A., 1851
pennanti (Philippi, 1846)
philberti (Récluz, 1843)
racketti (Payraudeau, 1826)
rarielineata (Michaud, 1829)
spratti (Forbes, 1844)
spurca (Gould, 1856)
tingitana Pallary, 1901
tumida (Montagu, 1803)
turbinoïdes (Deshayes, 1835)
umbilicalis (da Costa, 1778)
umbilicaris (Linné, 1758)
varia (Linné, 1758)
vimontiae Monterosato, 1884
- Phorcus**
mutabilis (Philippi, 1846)
richardi (Payraudeau, 1826)
- Osilinus**
articulatus (Lamarck, 1822)
atratus (Wood W., 1828)
lineatus (da Costa, 1778)
sauciatus (Koch, 1845)
turbinatus (Von Born, 1778)

- Calliostoma*
caroli (Dautzenberg, 1927)
cleopatra (Locard, 1896)
conulus (Linné, 1758)
granulatum (Von Born, 1778)
grimaldii (Dautzenberg & Fischer H., 1896)
gualterianum (Philippi, 1848)
gubbiolii Nofroni, 1984
hirondellei (Dautzenberg & Fischer H., 1896)
kochi Pallary, 1902
laugieri (Payraudeau, 1826)
leptophyma (Dautzenberg & Fischer H., 1896)
lithocletum (Dautzenberg, 1925)
milneedwardsi (Locard, 1898)
normani (Dautzenberg & Fischer H., 1897)
obesulum (Locard, 1898)
occidentale (Mighels & Adams, 1842)
virescens Coen, 1933
zizyphinum (Linné, 1758)
- Danilia*
affinis (Dautzenberg & Fischer H., 1896)
otaviana (Cantraine, 1835)
- Turcicula*
alicei (Dautzenberg & Fischer H., 1896)
miranda (Locard, 1898)
- Calliotropis*
ottoi (Philippi, 1844)
- Putzeysia*
wiseri (Calcara, 1842)
- Margarites*
costalis (Gould, 1841)
groenlandicus (Gmelin, 1791)
helicinus (Phipps, 1774)
olivaceus (Brown, 1827)
vahlii (Møller, 1842)
- Solariella*
amabilis (Jeffreys, 1865)
cincta (Philippi, 1836)
cingulima (Locard, 1898)
infundibulum Watson, 1879
lusitanica (Fischer P., 1887)
micans (Dautzenberg & Fischer H., 1896)
mogadorensis (Locard, 1898)
obscura (Couthouy, 1838)
paucivaricosa (Dautzenberg, 1889)
regalis (Verrill & Smith, 1880)
rhina (Watson, 1886)
rudecta (Locard, 1898)
vallanti (Dautzenberg & Fischer H., 1896)
varicosa (Mighels & Adams, 1842)
- Callumbonella*
suturalis (Philippi, 1836)
- Minolia*
nedyma Melvill, 1897
- Family Seguenziidae**
- Seguenzia*
elegans Jeffreys, 1885
ionica Watson, 1878
monocingulata Seguenza G., 1876
- Basilissopsis*
watsoni (Dautzenberg & Fischer H., 1897)
- Carenzia*
carinata (Jeffreys, 1877)
- Asthelys*
mundus (Watson, 1879)
- Ancistrobasis*
reticulata (Philippi, 1844)
- Family Trochaclididae**
- Trochacalis*
islandicus Warén, 1989
versilensis Warén, Carozza & Rocchini in Warén, 1992
- Family Pendromidae**
- Rugulina*
- fragilis* (Sars G.O., 1878)
monterosatoi (van Aartsen & Bogi, 1987)
- Family Skeneidae**
- Skenea*
areolata (Sars G.O., 1878)
basistriata (Jeffreys, 1877)
catenoides (Monterosato, 1877)
ferruginea Warén, 1991
larseni Warén, 1993
nilarum (Engl, 1996)
ossiansarsi Warén, 1991
pelagia Nofroni & Valenti, 1987
peterseni (Friele, 1877)
politá Warén, 1993
ponsonbyi (Dautzenberg & Fischer H., 1897)
profunda (Friele, 1879)
rugulosa (Sars G.O., 1878)
serpuloides (Montagu, 1808)
trochoides (Friele, 1876)
turgida (Odhner, 1912)
valvatoides (Jeffreys, 1883)
- Adeuomphalus*
ammoniformis (Seguenza G., 1876)
- Akritogyra*
conspicua (Monterosato, 1880)
curvilineata Warén, 1992
helicella Warén, 1993
similis (Jeffreys, 1883)
- Anekse*
affinis (Jeffreys, 1883)
giustii Bogi & Nofroni, 1989
infidata Warén, 1992
paucistriata Warén, 1992
sculpturata Warén, 1992
undulisculpta Bouchet & Warén, 1979
- Cirsonella*
ateles (Dautzenberg & Fischer H., 1896)
gaudryi (Dautzenberg & Fischer H., 1896)
romellensis (Granata-Grillo, 1877)
- Dikoleps*
cutleriana (Clark W., 1848)
depressa (Monterosato, 1880)
marianae (Rubio, Dantart & Luque, 1998)
pruinosa (Chaster, 1896)
pusilla (Jeffreys, 1847)
rolani (Rubio, Dantart & Luque, 1998)
umbilicostriata (Gaglini, 1987)
- Eudaronia*
aperta (Sykes, 1925)
- Ganesa*
bujnitzkii (Gorbunov, 1946)
nitidiuscula Jeffreys, 1883
- Granigryra*
arenosa Warén, 1993
granulifera Warén, 1992
pruinosa (Jeffreys, 1883)
tenera (Jeffreys, 1883)
- Lissomphalia*
bithynoides (Monterosato, 1880)
- Lissotesta*
gittenbergeri (van Aartsen & Bogi, 1988)
major Warén, 1992
minima (Seguenza G., 1876)
turrita (Gaglini, 1987)
- Mikro*
globulus Warén, 1996
- Protolira*
thorvaldsoni (Warén, 1996)
- Moelleriopsis*
messanensis (Seguenza G., 1876)
normani (Dautzenberg & Fischer H., 1897)
richardi (Dautzenberg & Fischer H., 1896)
- Palazzia*
ausonia (Palazzi, 1988)
- Parviturbo*

- elegantulus* (Philippi, 1844)
fenestratus (Chaster, 1896)
Pseudorbis
granulum (Brugnone, 1873)
jameoensis (Rubio & Rodriguez Babio, 1991)
Reticyra
granulosa (Sykes, 1925)
iheringi (Dautzenberg & Fischer H., 1897)
millipunctata (Friele, 1886)
Skeneoides
digeronimo (La Perna, 1998)
exilissima (Philippi, 1844)
jeffreysii (Monterosato, 1872)
Xyloskenea
naticiformis (Jeffreys, 1883)
- Family Cyclostrematidae**
Vetulonia
cancellata (Jeffreys, 1883)
- Family Turbinidae**
Bolma
macandrewii (Mørch, 1868)
rugosa (Linné, 1767)
Cantrainea
peloritana (Cantraine, 1835)
Homalopoma
carmelae Oliverio & Buzzurro, 1994
sanguineum (Linné, 1758)
Moelleria
costulata (Møller, 1842)
- Family Tricoliidae**
Tricolia
algoidea (Pallary, 1920)
deschampsi Gofas, 1993
entomocheila Gofas, 1993
miniata (Monterosato, 1884)
nordsiecki (Talavera, 1978)
pullus (Linné, 1758)
punctura Gofas, 1993
speciosa (Von Mühlfeldt, 1824)
tenuis (Michaud, 1829)
tingitana Gofas, 1982
- Family Cerithiidae**
Cerithium
nodulosum Bruguière, 1792
alukastrum (Brocchi, 1814)
lividulum Risso, 1826
nesioticum Pilsbry & Vanatta, 1906
protractum Bivona Ant. in Bivona And., 1838
rupestrum Risso, 1826
scabridum Philippi, 1848
vulgatum Bruguière, 1792
Bittium
depauperatum Watson, 1897
incile Watson, 1897
lacteum (Philippi, 1836)
latreillii (Payraudeau, 1826)
proteinum (Jousseaume, 1930)
reticulatum (da Costa, 1778)
simplex (Jeffreys, 1867)
submamillatum (de Rayneval & Ponzi, 1854)
watsoni (Jeffreys, 1885)
Cassiella
abyensis Gofas, 1987
Clypeomorus
bifasciatus (Sowerby G.B. II, 1855)
Rhinoclavis
kochi (Philippi, 1848)
- Family Dialidae**
Diala
varia Adams A., 1861
- Family Fossaridae**
Fossarus
ambiguus (Linné, 1758)
- Family Litiopidae**
Litiopa
melanostoma (Rang, 1829)
Alaba
mirabilis (Hornung & Mermad, 1926)
- Family Obtortionidae**
Finella
pupoides Adams A., 1860
Clathrofenella
fusca (Adams A., 1860)
- Family Planaxidae**
Planaxis
punctostriata Smith E.A., 1872
savignyi Deshayes, 1844
- Family Potamididae**
Potamides
conicus (de Blainville, 1829)
- Family Scaliolidae**
Scaliola
elata Issel, 1869
- Family Siliquariidae**
Tenagodus
obitusus (Schumacher, 1817)
senegalensis (Récluz in Mørch, 1860)
- Family Turritellidae**
Mesalia
brevialis (Lamarck, 1822)
Turritella
communis Risso, 1826
decipiens Monterosato, 1878
turbona Monterosato, 1877
Tachyrhynchus
spitzbergensis Golikov, 1986
- Family Triphoridae**
Cheirodonta
pallescens (Jeffreys, 1867)
Cosmotriphora
melanura (Adams C.B., 1850)
Marshallora
adversa (Montagu, 1803)
Monophorus
erythrosoma (Bouchet & Guillemot, 1978)
perversus (Linné, 1758)
thiriota Bouchet, 1985
- Obesula*
marisnstri Bouchet, 1985
Pogonodon
pseudocanaricus (Bouchet, 1985)
Nototriphora
canarica Nordsieck & Talavera, 1979
- Similiphora*
similior (Bouchet & Guillemot, 1978)
tricolata (Bouchet, 1997)
- Strobiligera*
brychia (Bouchet & Guillemot, 1978)
flammulata Bouchet & Warén, 1993
lubrica Bouchet & Warén, 1993
- Metaxia*
abrupta (Watson, 1880)
bacillum (Issel, 1869)
metaxae (delle Chiaje, 1828)
- Family Cerithiopsidae**
Cerithiopsis
atalaya Watson, 1885
barleei Jeffreys, 1867
perlata (Monterosato, 1889)
diadema Monterosato, 1874
fayalensis Watson, 1886
horrida (Monterosato, 1874)
jeffreysi Watson, 1885
minima (Brusina, 1865)
nofronii Amati, 1987
pulvis (Issel, 1869)

- scalaris* (Locard, 1892)
tenthrenoides (Melvill, 1896)
tuberculatus (Montagu, 1803)
- Cerithiella**
amblytera (Watson, 1880)
enodis (Watson, 1880)
insignis (Jeffreys, 1885)
metula (Lovén, 1846)
- Triforis**
anel pistos (Bouchet & Fechter, 1981)
superstes (Bouchet & Fechter, 1981)
- Dizoniopsis**
coppolae (Aradas, 1870)
micalii Cecalupo & Villari, 1997
- Ektonus**
turbanilloides (Dautzenberg & Fischer H., 1896)
- Krachia**
cossmanni (Dautzenberg & Fischer H., 1896)
cylindrata (Jeffreys, 1885)
guernei (Dautzenberg & Fischer H., 1896)
obeliscoidea (Jeffreys, 1885)
tiara (Monterosato, 1874)
- Laeocochlis**
sinistratus (Nyst, 1835)
- Onchodia**
valeriae (Giusti Fr., 1987)
- Seila**
trilineata (Philippi, 1836)
- Eumetula**
aliceae (Dautzenberg & Fischer H., 1896)
arctica (Mørch, 1857)
bouvieri (Dautzenberg & Fischer H., 1896)
- Family Janthinidae**
- Janthina**
exigua Lamarck, 1816
janthina (Linné, 1758)
nitens Menke, 1828
pallida Thompson W., 1840
- Family Acididae**
- Aclis**
ascaris (Turton, 1819)
attenuans Jeffreys, 1883
gulsonae (Clark W., 1850)
minor (Brown, 1827)
sarsi Dautzenberg & Fischer H., 1912
trilineata Watson, 1897
verduini van Aartsen, Menhorst & Gittenberger, 1984
vitrea Watson, 1897
walleri Jeffreys, 1867
- Discalis**
canariensis Moolenbeek & Warén, 1987
- Family Epitonidae**
- Epitonium**
aculeatum (Allan, 1818)
algerianum (Weinkauf, 1866)
candidissimum (Monterosato, 1877)
celesti (Aradas, 1854)
clathratulum (Kanmacher, 1798)
commune (Lamarck, 1822)
dallianum (Verrill & Smith, 1880)
delicatulum (Adams H., 1869)
dendrophylliae Bouchet & Warén, 1986
fischeri (Watson, 1897)
fulvovittatum (Dautzenberg, 1890)
greenlandicum (Perry, 1811)
hispidulum (Monterosato, 1874)
jolyi (Monterosato, 1878)
linctum (de Boury & Monterosato, 1890)
nanum (Jeffreys, 1884)
pseudonanum Bouchet & Warén, 1986
pulchellum (Bivona, 1832)
rhips (Watson, 1897)
semidisjunctum (Jeffreys, 1884)
- striatissimum* (Monterosato, 1878)
tiberii (de Boury, 1890)
trevelyanum (Thompson W., 1840)
tryoni (de Boury, 1913)
turtonis (Turton, 1819)
vittatum (Jeffreys, 1884)
webbii (d'Orbigny, 1840)
- Cycloscala**
hyalina (Sowerby G.B. II, 1844)
- Acisra**
coarctata (Jeffreys, 1884)
subdecussata (Cantraine, 1835)
- Acriloscala**
lamyi (de Boury, 1909)
- Alora**
retifera Bouchet & Warén, 1986
tenerrima (Dautzenberg & Fischer H., 1896)
- Cirsorema**
cochlea (Sowerby G.B. II, 1844)
- Claviscala**
richardi (Dautzenberg & de Boury, 1897)
- Cylindriscala**
acus (Watson, 1883)
auricula Bouchet & Warén, 1986
guernei (Dautzenberg & de Boury, 1897)
jeffreysi (Tryon, 1887)
mirifica (Fischer P. in Filhol, 1886)
thalassae Bouchet & Warén, 1986
vicina (Dautzenberg & de Boury, 1897)
- Gregorioiscala**
exfoliata Bouchet & Warén, 1986
pachya (Locard, 1897)
sarsi (Kobelt, 1904)
- Gyroscala**
lamellosa (Lamarck, 1822)
- Opalia**
abbotti Clench & Turner, 1952
crenata (Linné, 1758)
fortunata Bouchet & Warén, 1986
hellenica (Forbes, 1844)
- Periapta**
gracilis (Verrill, 1880)
polygyrella (Fischer P. in Locard, 1897)
- Punctiscala**
cerigottana (Sturany, 1896)
- Eccliseogrya**
exquisita Bouchet & Warén, 1986
folini (Dautzenberg & de Boury, 1897)
formosissima (Jeffreys, 1884)
monnioti Bouchet & Warén, 1986
nitida (Verrill & Smith in Verrill, 1885)
sericea Bouchet & Warén, 1986
- Iphitus**
cancellatus Dautzenberg & Fischer H., 1896
marshalli (Sykes, 1925)
tuberatus Jeffreys, 1883
- Narrimania**
concinna (Sykes, 1925)
- Opaliopsis**
atlantis (Clench & Turner, 1952)
- Papuliscala**
cerithielloides Bouchet & Warén, 1986
elongata (Watson, 1881)
praelonga (Jeffreys, 1877)
tavianii Bouchet & Warén, 1986
- Family Eulimidae**
- Eulima**
altilimirai Nordsieck, 1977
anonyma Bouchet & Warén, 1986
bilineata Alder, 1848
chionea Bouchet & Warén, 1986
confusa Bouchet & Warén, 1986
devistoma Bouchet & Warén, 1986
fuscozonata Bouchet & Warén, 1986
glabra (da Costa, 1778)

- leptozena* Dautzenberg & Fischer H., 1896
strongylostoma Bouchet & Warén, 1986
Acrochalix
callosa Bouchet & Warén, 1986
Auriculigerina
miranda Dautzenberg, 1925
Bathewulma
abbreviata (Jeffreys, 1884)
fuscoapicata (Jeffreys, 1884)
Bathyrycincola
curta (Warén, 1972)
macrapex Bouchet & Warén, 1986
micrapex Bouchet & Warén, 1986
talaena (Dautzenberg & Fischer H., 1897)
Bulimeulima
magna Bouchet & Warén, 1986
Campylorhaphion
famelicum (Watson, 1883)
machaeropse (Dautzenberg & Fischer H., 1896)
Costaclis
mizon (Watson, 1881)
Crinolamia
angustispira Bouchet & Warén, 1986
dahli Bouchet & Warén, 1979
Crinophtheiros
comatuliculus (Graff, 1875)
giustii Gaglini, 1991
junii (de Folin, 1887)
Curveulima
macrophthalmica (Warén, 1972)
marshalli Bouchet & Warén, 1986
obliquistoma Bouchet & Warén, 1986
Enteroxenos
oestergreni Bonnevie, 1902
Entocolax
ludwigi Voigt, 1888
Entoconcha
mirabilis (Müller J., 1852)
Ersilia
mediterranea (Monterosato, 1869)
Eulitoma
arca Bouchet & Warén, 1986
insignis (Dautzenberg & Fischer H., 1896)
josephinae Bouchet & Warén, 1986
obtusiuscula Bouchet & Warén, 1986
Fuscapex
baptocephalus (Dautzenberg & Fischer H., 1896)
cabiochi Bouchet & Warén, 1986
major Bouchet & Warén, 1986
microcostellatus Bouchet & Warén, 1986
talismanni Bouchet & Warén, 1986
Fuseulima
boscheineni (Engl, 1998)
fulva (Watson, 1897)
ingolfiana Bouchet & Warén, 1986
minuta (Jeffreys, 1884)
projectilabrum Bouchet & Warén, 1986
sordida (Watson, 1897)
thalassae Bouchet & Warén, 1986
Haliella
canarica Bouchet & Warén, 1986
stenostoma (Jeffreys, 1858)
Halielloides
fragilis Bouchet & Warén, 1986
nitidus (Verrill, 1884)
Hemiaclis
major Bouchet & Warén, 1986
obtusus Bouchet & Warén, 1986
ventrosus (Friele, 1876)
Megadenus
oneirophantae Bouchet & Lützen, 1980
Melanella
alba (da Costa, 1778)
boscii (Payraudeau, 1826)
charissa (Verrill, 1884)
- compactilis* (Monterosato, 1884)
costellata (Dautzenberg & Fischer H., 1897)
crosseana (Brusina, 1886)
densicostata Bouchet & Warén, 1986
doederleini (Brusina, 1886)
frielei (Jordan, 1895)
gagei Bouchet & Warén, 1986
glypta Bouchet & Warén, 1986
halopora (Dautzenberg & Fischer H., 1896)
inconspicua (Watson, 1897)
insculpta Bouchet & Warén, 1986
jeffreysi (Tryon, 1886)
laurae (Friele, 1886)
lineata (Monterosato, 1869)
lubrica (Monterosato, 1890)
lucida (Verrill, 1884)
martynjordani (Jordan, 1895)
microsculpta Bouchet & Warén, 1986
monterosatoi (Monterosato, 1890)
myriotrochi Bouchet & Warén, 1986
obtusoapicata Bouchet & Warén, 1986
orphanensis Clarke A.H., 1974
petitiana (Brusina, 1869)
planisulis Bouchet & Warén, 1986
polita (Linné, 1758)
praecurta (Pallary, 1904)
pseudoglabra (Dautzenberg & Fischer H., 1912)
reticulata Bouchet & Warén, 1986
richardi (Dautzenberg & Fischer H., 1896)
similior Bouchet & Warén, 1986
spiridioni (Dautzenberg & Fischer H., 1896)
stalioi (Brusina, 1869)
translucens (Monterosato, 1890)
trunca (Watson, 1897)
turbanilloides Bouchet & Warén, 1986
turrita Bouchet & Warén, 1986
Mucronalia
mammillata Dall, 1927
Nanobalcis
nanus (Monterosato, 1878)
Niso
foresti Bouchet & Warén, 1986
Oceanida
confluens Bouchet & Warén, 1986
Ophieulima
minima (Dall, 1927)
Ophiolamia
fragilissima Bouchet & Warén, 1986
Parastilbe
acuta (Jeffreys, 1884)
Parvioris
ibizencus (Nordsieck, 1968)
Pelseneeria
media Koehler & Vaney, 1908
minor Koehler & Vaney, 1908
profunda Koehler & Vaney, 1908
striata Bouchet & Warén, 1986
stylifera (Turton, 1825)
Echineulima
mittrei (Petit de la Saussaye, 1851)
Pisolania
brychia (Watson, 1883)
Rectilabrum
lanceolatum Bouchet & Warén, 1986
Sabinella
pachya (Dautzenberg & Fischer H., 1896)
piriformis Brugnone, 1873
Slichteulima
badia (Watson, 1897)
jeffreysiana (Brusina, 1869)
lata Bouchet & Warén, 1986
lentiginosa (Adams A., 1861)
richteri (Engl, 1997)
wareni (Engl, 1997)
Umbilibalcis

- crassulus* Bouchet & Warén, 1986
subumbilicatus (Jeffreys, 1883)
- Vitreolina**
antiflexa Monterosato, 1884
cionella (Monterosato, 1878)
curva (Monterosato, 1874)
incurva (Bucquoy, Dautzenberg & Dollfus, 1883)
knudseni Bouchet & Warén, 1986
levantina Oliverio, Buzzurro & Villa, 1994
permixta (Jeffreys, 1883)
philippi (de Raynal & Ponzi, 1854)
- Family Littorinidae**
Melarhaphe
neritoidea (Linné, 1758)
Nodilittorina
punctata (Gmelin, 1791)
Littorina
arcana Hannaford-Ellis, 1978
sabalis (Turton, 1825)
littorea (Linné, 1758)
compressa Jeffreys, 1865
obtusata (Linné, 1758)
saxatilis (Olivi, 1792)
striata King & Broderip, 1832
- Family Lacunidae**
Lacuna
crassior (Montagu, 1803)
pallidula (da Costa, 1778)
parva (da Costa, 1778)
vincta (Montagu, 1803)
- Family Skeneopsidae**
Skeneopsis
planorbis (Fabricius O., 1780)
sultanicum Gofas, 1983
- Family Zerotulidae**
Zerotula
incognita Warén & Hain, 1996
- Family Cingulopsidae**
Eatonina
celata (Monterosato, 1884)
cossuriae (Calcaria, 1841)
fulgida (Adams J., 1797)
maiilidae Rubio & Rodriguez Babio, 1996
ochroleuca (Brusina, 1869)
pumila (Monterosato, 1884)
- Tubbrevia**
micrometrica (Aradas & Benoit, 1876)
- Family Rissoidae**
Rissoa
aartseni Verduin, 1985
albugo Watson, 1873
alleryi (Nordsieck, 1972)
angustior (Monterosato, 1917)
auriformis Pallary, 1904
auriscalpium (Linné, 1758)
decorata Philippi, 1846
frauenseldiana Brusina, 1868
gemmula Fischer P. in de Folin, 1869
gomerica (Nordsieck & Talavera, 1979)
guernei Récluz, 1843
italiensis Verduin, 1985
janusi (Nordsieck, 1972)
lilacina Récluz, 1843
lia (Monterosato, 1884)
membranacea (Adams J., 1800)
monodonta Philippi, 1836
multicincta (Smriglio & Mariottini, 1995)
multicostata (Nordsieck & Talavera, 1979)
panhormensis Verduin, 1985
paradoxa (Monterosato, 1884)
parva (da Costa, 1778)
pseudoguerini (Nordsieck & Talavera, 1979)
- rodhensis* Verduin, 1985
scurra (Monterosato, 1917)
similis Scacchi, 1836
splendida Eichwald, 1830
torquilla Pallary, 1912
variabilis (Von Mihlfeldt, 1824)
ventricosa Desmarest, 1814
violacea Desmarest, 1814
- Pusillina**
amblia (Watson, 1886)
benzi (Aradas & Maggiore, 1844)
ehrenbergi (Philippi, 1844)
inconspicua (Alder, 1844)
lineolata (Michaud, 1832)
marginata (Michaud, 1832)
metivieri Bouchet & Warén, 1993
munda (Monterosato, 1884)
philippi (Aradas & Maggiore, 1844)
radiata (Philippi, 1836)
sarsi (Lovén, 1846)
testudae (Verduin, 1979)
- Setia**
aartseni (Verduin, 1984)
amabilis (Locard, 1886)
ambigua (Brugnone, 1873)
anselmoi (van Aartsen & Engl, 1999)
antiopolitana (van der Linden & Wagner, 1987)
bruggenii (Verduin, 1984)
gittenbergeri (Verduin, 1984)
jansseni (Verduin, 1984)
kuiperi (Verduin, 1984)
lacourtii (Verduin, 1984)
latior (Mighels & Adams, 1842)
lidya Verduin, 1988
maculata (Monterosato, 1869)
miae Verduin, 1988
microbia Hoenselaar & Hoenselaar, 1991
pulcherrima (Jeffreys, 1848)
quisquilarum (Watson, 1886)
scillae (Aradas & Benoit, 1876)
slifikorum (Verduin, 1984)
subvaricosa Gofas, 1990
tiberiana (Issel, 1869)
ugesae Verduin, 1988
valvatoides (Milaschewitsch, 1909)
- Rudolphosetia**
fusca (Philippi, 1844)
- Alvania**
aartseni Verduin, 1986
abstersa van der Linden & van Aartsen, 1993
adiaphoros Bouchet & Warén, 1993
adinogramma Bouchet & Warén, 1993
aeoliae Palazzi, 1988
algeriana (Monterosato, 1877)
amatii Oliverio, 1986
angioyi van Aartsen, 1982
angularis (Warén, 1996)
aspera (Philippi, 1844)
aurantiaca (Watson, 1873)
basteriae (Moolenbeek & Faber, 1986)
beani (Hanley in Thorpe, 1844)
canariensis (d'Orbigny, 1840)
cancapae Bouchet & Warén, 1993
cancellata (da Costa, 1778)
carinata (da Costa, 1778)
cimex (Linné, 1758)
cinicoides (Forbes, 1844)
cingulata (Philippi, 1836)
clarae (Nofroni & Pizzini, 1991)
clathrella (Seguenza L., 1903)
colossaliphila Oberling, 1970
consociella Monterosato, 1884
corona Nordsieck, 1972
coeli (Gofas, 1999)
datchaensis Amati & Oliverio, 1987

- dianensis* Oliverio, 1988
dictyophora (Philippi, 1844)
dijkstrai (Hoenselaar & Goud, 1998)
dipacoi Giusti Fr. & Nofroni, 1989
discors (Allan, 1818)
disparilis Monterosato, 1890
dorbignyi (Audouin, 1826)
electa (Monterosato, 1874)
elegantissima (Monterosato, 1875)
euchila (Watson, 1886)
fischeri (Jeffreys, 1884)
formicarum Gofas, 1989
fractospira (Oberling, 1970)
gagliniae Amati, 1985
geryonia (Nardo, 1847)
guancha Moolenbeek & Hoenselaar, 1989
hallgassi Amati & Oliverio, 1985
hirta (Monterosato, 1884)
hispidula (Monterosato, 1884)
imperspicua (Pallary, 1920)
incognita (Warén, 1996)
internodula (Hoenselaar & Goud, 1998)
iunoniae Palazzi, 1988
jeffreysi (Waller, 1864)
johannae (Moolenbeek & Hoenselaar, 1998)
joseae (Hoenselaar & Goud, 1998)
lactea (Michaud, 1832)
lamellata Dautzenberg, 1889
lanciae (Calcaria, 1845)
leacocki (Watson, 1873)
lineata Risso, 1826
litoralis (Nordsieck, 1972)
lucinae Oberling, 1970
macandrewi (Manzoni, 1868)
mamillata Risso, 1826
marchadi (Gofas, 1999)
mediolittoralis Gofas, 1989
microstriata (Hoenselaar & Goud, 1998)
minuta (Golikov & Fedjakov in Scarlato, 1987)
moerchi (Collin, 1886)
moniziana (Watson, 1873)
multiquadrata van der Linden & Wagner, 1989
nestaresi Oliverio & Amati, 1990
nonsculpta (Hoenselaar & Goud, 1998)
obliqua Warén, 1974
pagodula (Bucquoy, Dautzenberg & Dollfus, 1884)
parvula (Jeffreys, 1884)
piersmai Moolenbeek & Hoenselaar, 1989
platycephala Dautzenberg & Fischer H., 1896
porcupinae Gofas & Warén, 1982
poucheti Dautzenberg, 1889
pseudoareolata Warén, 1974
pseudosyngenes Warén, 1973
punctura (Montagu, 1803)
renei (Hoenselaar & Goud, 1998)
rudis (Philippi, 1844)
scabra (Philippi, 1844)
schwartziana Brusina, 1866
scrobiculata (Møller, 1842)
sculptilis (Monterosato, 1877)
settepassii Amati & Nofroni, 1985
simulans Locard, 1886
steuersi (Amati, 1987)
slieringsi (Hoenselaar & Goud, 1998)
sororecula Granata-Grillo, 1877
spinosa (Monterosato, 1890)
stenolopha Bouchet & Warén, 1993
subareolata Monterosato, 1869
subcalathus (Dautzenberg & Fischer H., 1906)
subcrenulata (Bucquoy, Dautzenberg & Dollfus, 1884)
subsoluta (Aradas, 1847)
tarsodes (Watson, 1886)
tenera (Philippi, 1844)
tessellata Weinkauff, 1868
- testae* (Aradas & Maggiore, 1844)
tomentosa (Pallary, 1920)
verrilli (Friele, 1886)
watsoni (Schwartz in Watson, 1873)
weinkauffi Weinkauff, 1868
wyvillethomsoni (Friele, 1877)
zetlandica (Montagu, 1815)
zoderi (Hoenselaar & Goud, 1998)
zylenensis Gofas & Warén, 1982
- Amphirissa*
cyclostomoides Dautzenberg & Fischer H., 1897
- Benthonella*
tenella (Jeffreys, 1869)
- Benthonellania*
agastyches Bouchet & Warén, 1993
fayalensis (Watson, 1886)
oligostigma Bouchet & Warén, 1993
- Boreocingula*
castanea (Møller, 1842)
- Cingula*
trifasciata (Adams J., 1800)
- Crisilla*
amphiglypha Bouchet & Warén, 1993
beniamina (Monterosato, 1884)
callosa (Manzoni, 1868)
chiarelli (Cecalupo & Quadri, 1995)
cristallinula (Manzoni, 1868)
depicta (Manzoni, 1868)
lincta (Watson, 1873)
marioni Fasulo & Gaglini, 1987
permixta (Manzoni, 1868)
picta (Jeffreys, 1867)
postrema (Gofas, 1990)
semistriata (Montagu, 1808)
spadix (Watson, 1897)
- Frigidoalvania*
cruenta (Odhner, 1915)
janmayeni (Friele, 1878)
thalassae Bouchet & Warén, 1993
- Gofasia*
galiciae Bouchet & Warén, 1993
josephinae Bouchet & Warén, 1993
vanderlandi Bouchet & Warén, 1993
- Manzonia*
boogi Moolenbeek & Faber, 1987
boucheti Amati, 1992
castanea Moolenbeek & Faber, 1987
crassa (Kanmacher, 1798)
crispa (Watson, 1873)
darwini Moolenbeek & Faber, 1987
dionisi Rolán, 1987
gibbera (Watson, 1873)
guitiani Rolán, 1987
heroensis Moolenbeek & Hoenselaar, 1992
madeirensis Moolenbeek & Faber, 1987
overdiepi van Aartsen, 1983
pelorum Moolenbeek & Faber, 1987
unifasciata Dautzenberg, 1889
vigoensis (Rolán, 1983)
wilmae Moolenbeek & Faber, 1987
- Obtusella*
intersecta (Wood S., 1857)
macilenta (Monterosato, 1880)
roseotincta (Dautzenberg, 1889)
tumidula (Sars G.O., 1878)
- Onoba*
aculeus (Gould, 1841)
dimassai Amati & Nofroni, 1991
gianninii (Nordsieck, 1974)
guzmani Hoenselaar & Moolenbeek, 1987
improcera (Warén, 1996)
islandica (Friele, 1876)
josae Moolenbeek & Hoenselaar, 1987
leptalea (Verill, 1884)
manzoniana Rolán, 1987

- mighelsi* (Stimpson, 1851)
moreleti Dautzenberg, 1889
obliqua (Warén, 1974)
semicostata (Montagu, 1803)
tarifensis Hoenselaar & Moolenbeek, 1987
torelli (Warén, 1996)
- Plagyostila**
asturiana Fischer P. in de Folin, 1872
- Peringiella**
denticulata Ponder, 1985
elegans (Locard, 1892)
- Botryphallus**
epidauricus (Brusina, 1866)
ovummuscae (Gofas, 1990)
- Pseudosetia**
amydroax Bouchet & Warén, 1993
azorica Bouchet & Warén, 1993
ficariensis (Brugnone, 1876)
semipellucida (Friele, 1879)
turgida (Jeffreys, 1870)
- Rissoina**
bertholleti Issel, 1869
bruguieri (Payraudeau, 1826)
- Zebina**
paivensis (Watson, 1873)
robustior (Gofas, 1999)
- Family Adeorbidae**
Circulus
striatus (Philippi, 1836)
supranitidus (Wood S., 1848)
tricarinatus (Wood S., 1848)
- Teinostoma**
azoricum (Dautzenberg & Fischer H., 1896)
- Family Anabathridae**
Nodulus
contortus (Jeffreys, 1856)
- Pisinna**
glabrata (Von Mühlfeldt, 1824)
- Family Assimineidae**
Assiminea
grayana Fleming, 1828
Paludinella
littorina (delle Chiaje, 1828)
- Family Barleeidae**
Barleeia
gougeti (Michaud, 1830)
seminulum (Monterosato, 1877)
unifasciata (Montagu, 1803)
- Family Caecidae**
Caecum
armoricum de Folin, 1869
atlantidis Watson, 1897
auriculatum de Folin, 1868
clarkii Carpenter, 1859
elegantissimum (Carpenter, 1859)
engli (Nofroni, Pizzini & Oliverio, 1997)
eunoai (Nofroni, Pizzini & Oliverio, 1997)
glabrum (Montagu, 1803)
pollicare Carpenter, 1859
searleswoodi Carpenter, 1859
subannulatum de Folin, 1870
swinneni (Nofroni, Pizzini & Oliverio, 1997)
trachea (Montagu, 1803)
vitreum (Carpenter, 1859)
- Parastrophia**
asturiana de Folin, 1870
- Family Elachisinae**
Laeviphitus
verduini van Aartsen, Bogi & Giusti, 1989
- Elachisina**
canarica Nordsieck & Talavera, 1979
globulooides (Warén, 1972)
- Family Hydrobiidae**
- Hydrobia**
acuta (Draparnaud, 1805)
joossei van Aartsen, Menkhorst & Gittenberger, 1984
neglecta Muus, 1963
ulvae (Pennant, 1777)
ventrosa (Montagu, 1803)
- Heleobia**
stagnorum (Gmelin, 1791)
- Family Iravadiidae**
Ceratia
proxima (Forbes & Hanley, 1850)
- Hyala**
vitrea (Montagu, 1803)
- Family Tornidae**
Tornus
mienisi (van Aartsen, Carrozza & Menkhorst, 1998)
subcarinatus (Montagu, 1803)
- Discopsis**
costulatus (de Folin, 1870)
- Family Truncatellidae**
Truncatella
subcylindrica (Linné, 1767)
- Family Vermetidae**
Vermetus
granulatus (Gravenhorst, 1831)
rugulosus Monterosato, 1878
semisorrectus Bivona Ant., 1832
triquetrus Bivona Ant., 1832
- Dendropoma**
anguliferum (Monterosato, 1884)
petraeum (Monterosato, 1884)
- Petalonconchus**
glomeratus (Linné, 1758)
- Serpulorbis**
arenarius (Linné, 1767)
- Family Strombidae**
Strombus
persicus (Swainson, 1821)
- Family Aporrhaidae**
Aporrhais
pespelecani (Linné, 1758)
serresianus (Michaud, 1828)
- Family Vanikoridae**
Larsenia
scalaroides Warén, 1989
- Macromphalus**
abyensis Warén & Bouchet, 1988
- Megalomphalus**
azonus (Brusina, 1865)
disciformis (Granata-Grillo, 1877)
petitianus (Tiberi, 1869)
- Talassia**
coriacea (Manzoni, 1868)
daguenneti (de Folin, 1873)
tenuisculpta (Watson, 1873)
- Family Hippocoridae**
Hipponix
conicus (Schumacher, 1817)
- Leptonotis**
perplexus (Suter, 1907)
- Family Xenophoridae**
Xenophora
crispa (Koenig, 1825)
- Family Calypttraeidae**
Calypttraea
chinensis (Linné, 1758)
- Crepidula**
calyptraeiformis Deshayes, 1830
fornicata (Linné, 1758)
gibbosa Defrance, 1818
porcellana Linné, 1758
unguiformis Lamarck, 1822

Family Haloceratidae	<i>Pseudosimnia</i>
<i>Haloceras</i>	<i>carnea</i> (Poiret, 1789)
<i>carinatus</i> (Jeffreys, 1883)	
<i>cingulatus</i> (Verrill, 1884)	
<i>laxus</i> (Jeffreys, 1885)	
<i>mediocostatus</i> (Dautzenberg & Fischer H., 1896)	
<i>tricarinatus</i> (Jeffreys, 1885)	
Family Capulidae	<i>Simnia</i>
<i>Capulus</i>	<i>nicaensis</i> Risso, 1826
<i>simplex</i> Locard, 1898	<i>patula</i> (Pennant, 1777)
<i>ungaricus</i> (Linné, 1758)	<i>purpurea</i> Risso, 1826
<i>Torellia</i>	<i>Primovula</i>
<i>delicata</i> (Philippi, 1844)	<i>bellociae</i> (Cardin, 1997)
<i>didyma</i> Bouchet & Warén, 1993	<i>Pedicularia</i>
<i>Trichotropis</i>	<i>sicula</i> Swainson, 1840
<i>borealis</i> Broderip & Sowerby G.B. I, 1829	
<i>conicus</i> Møller, 1842	Family Naticidae
<i>bicarinatus</i> subsp. <i>tenuis</i> Smith E.A., 1877	<i>Natica</i>
Family Lamellariidae	<i>canariensis</i> Odhner, 1932
<i>Lamellaria</i>	<i>dillwynii</i> Payraudeau, 1826
<i>latens</i> (Müller O.F., 1776)	<i>furva</i> Watson, 1897
<i>perspicua</i> (Linné, 1758)	<i>gualteriana</i> Récluz, 1844
<i>Calyptococoncha</i>	<i>hebraea</i> (Martyn, 1784)
<i>pellucida</i> (Verrill, 1880)	<i>prietoi</i> (Hidalgo, 1873)
<i>Marseniella</i>	<i>stercusmuscarum</i> (Gmelin, 1791)
<i>borealis</i> Bergh, 1886	<i>vittata</i> (Gmelin, 1791)
<i>Marsenina</i>	Cryptonatica
<i>glabra</i> (Couthouy, 1839)	<i>affinis</i> (Gmelin, 1791)
Family Velutinidae	<i>bathybii</i> (Friele, 1879)
<i>Velutina</i>	<i>operculata</i> (Jeffreys, 1885)
<i>lanigera</i> Møller, 1842	Tectonatica
<i>plicatilis</i> (Müller O.F., 1776)	<i>filosa</i> (Philippi, 1845)
<i>scheideri</i> Friele, 1886	<i>rizzae</i> (Philippi, 1844)
<i>undata</i> Smith J., 1839	Euspira
<i>velutina</i> (Müller O.F., 1776)	<i>catena</i> (da Costa, 1778)
<i>Onchidiopsis</i>	<i>fusca</i> (de Blainville, 1825)
<i>glacialis</i> (Sars M., 1851)	<i>grossularia</i> (Marche-Marchad, 1957)
<i>latissima</i> Odhner, 1913	<i>guillemini</i> (Payraudeau, 1826)
<i>spitzbergensis</i> Jensen in Thorson, 1944	<i>macilenta</i> (Philippi, 1844)
<i>Piliscus</i>	<i>montagui</i> (Forbes, 1838)
<i>commodus</i> (Middendorff, 1851)	<i>notabilis</i> Jeffreys, 1885
<i>Pseudotorellia</i>	<i>obtusa</i> (Jeffreys, 1885)
<i>fragilis</i> Warén, 1989	<i>pallida</i> (Broderip & Sowerby G.B. I, 1829)
Family Triviidae	<i>phaeocephala</i> (Dautzenberg & Fischer H., 1896)
<i>Trivia</i>	<i>presubplicata</i> (Bouchet & Warén, 1993)
<i>arcica</i> (Pulteney, 1799)	<i>pulchella</i> (Risso, 1826)
<i>bitou</i> Pallary, 1913	<i>subplicata</i> (Jeffreys, 1885)
<i>levantina</i> (Smriglio, Mariottini & Buzzurro, 1998)	Amauroopsis
<i>monacha</i> (da Costa, 1778)	<i>brassiculina</i> (Locard, 1897)
<i>multilirata</i> Sowerby G.B. II, 1870	<i>islandica</i> (Gmelin, 1791)
<i>pulex</i> (Solander in Gray J.E., 1828)	<i>sphaeroides</i> (Jeffreys, 1877)
<i>spongicola</i> (Monterosato, 1923)	Bulbus
Family Eratoidae	<i>globosus</i> (Jeffreys, 1885)
<i>Erato</i>	<i>smithii</i> Brown, 1839
<i>voluta</i> (Montagu, 1803)	Pseudopolinices
Family Cypraeidae	<i>nanus</i> (Møller, 1842)
<i>Erosaria</i>	Neverita
<i>spurca</i> (Linné, 1758)	<i>josephinia</i> Risso, 1826
<i>turdus</i> (Lamarck, 1810)	Payraudeautia
<i>Luria</i>	<i>intricata</i> (Donovan, 1804)
<i>lurida</i> (Linné, 1758)	Sinum
<i>Purpuradusta</i>	<i>bifasciatum</i> (Récluz, 1851)
<i>gracilis</i> (Gaskoin, 1849)	Family Tonnidae
<i>Schilderia</i>	<i>Tonna</i>
<i>achatidea</i> (Gray in Sowerby G.B. II, 1837)	<i>galea</i> (Linné, 1758)
<i>Zonaria</i>	Eudolium
<i>pyrum</i> (Gmelin, 1791)	<i>bairdii</i> (Verrill & Smith, 1881)
Family Ovulidae	Family Cassidae
<i>Aperiowlia</i>	<i>Galeoidea</i>
<i>adriatica</i> (Sowerby G.B. I, 1828)	<i>echinophora</i> (Linné, 1758)
<i>juanjosensis</i> Perez & Gomez, 1987	<i>rugosa</i> (Linné, 1771)
<i>Neosimnia</i>	Oocorys
<i>spelta</i> (Linné, 1758)	<i>sulcata</i> Fischer P., 1883
	Phalium
	<i>granulatum</i> (Von Born, 1778)
	<i>saburon</i> (Bruguière, 1792)
	Family Laubierinidae
	<i>Laubierina</i>
	<i>peregrinator</i> Warén & Bouchet, 1990
	Family Ranellidae

- Ranella**
olearium (Linné, 1758)
- Charonia**
lampas (Linné, 1758)
tritonis subsp. *variegata* Lamarck, 1816
- Cymatium**
aquatile (Reeve, 1844)
corrugatum (Lamarck, 1816)
kobelti (Von Maltzan, 1884)
labiosum (Wood W., 1828)
martinianum (d'Orbigny, 1847)
muricinum (Röding, 1798)
nicobaricum (Röding, 1798)
parthenopeum (Von Salis, 1793)
pharcidum (Dall, 1889)
trambiquecum (Lamarck, 1816)
trigonum (Gmelin, 1791)
- Cabestana**
cutacea (Linné, 1767)
dolaria (Linné, 1767)
- Linatella**
caudata (Gmelin, 1791)
- Family Pisaniuridae**
Pisanianura
grimaldii (Dautzenberg, 1899)
- Family Personidae**
Distorsio
perdistorta (Fulton, 1938)
- Family Bursidae**
Bursa
marginata (Gmelin, 1791)
scrabiculator (Linné, 1758)
ranelloides subsp. *tenuisculpta* Dautzenberg & Fischer H., 1906
thomae (d'Orbigny, 1842)
- Family Carinariidae**
Carinaria
mediterranea de Blainville, 1825
- Family Atlantidae**
Atlanta
fusca Souleyet, 1852
helicinoides Souleyet, 1852
inclinata Souleyet, 1852
infata Souleyet, 1852
lesueurii Souleyet, 1852
peresi Frontier, 1966
peronii Lesueur, 1817
- Oxygyrus*
keraudrenii (Lesueur, 1817)
- Protatlanta*
souleyeti (Smith E.A., 1888)
- Family Fiolidae**
Fioloida
desmarestia Lesueur, 1817
- Pterotrachea*
coronata Forskål, 1775
hippocampus Philippi, 1836
minuta Bonnevie, 1920
scutata Gegenbaur, 1855
- Family Muricidae**
Murex
forskoehlii Röding, 1798
- Aspella*
anceps (Lamarck, 1822)
- Bolinus*
brandaris (Linné, 1758)
cornutus (Linné, 1758)
- Murexiella*
bojadorensis (Locard, 1897)
- Dermomurex*
gosasi Houart, 1996
scalaroides (de Blainville, 1829)
- Hadriana*
- craticuloides* (Vokes, 1964)
- Hexaplex**
trunculus (Linné, 1758)
- Muricopsis**
aradasii (Poirier, 1883)
cristata (Brocchi, 1814)
medicago (Watson, 1897)
- Ocenebra**
chavesi Houart, 1996
erinacea (Linné, 1758)
inordinata Houart & Abreu, 1994
- Ocinebrina**
aciculata (Lamarck, 1822)
edwardsii (Payraudeau, 1826)
hybrida (Aradas & Benoit, 1876)
miscewichi Pallary, 1920
- Houartiella**
alboranensis Smriglio, Mariottini & Bonfitto, 1997
- Pterynotus**
atlantideus Bouchet & Warén, 1985
leucas (Locard, 1897)
- Trophon**
barvicensis (Johnston, 1825)
clathratus (Linné, 1767)
clavatus Sars G.O., 1878
dabneyi Dautzenberg, 1889
drouetti Dautzenberg, 1889
echinatus (Kiener, 1840)
fabricii Beck in Möller, 1842
muricatus (Montagu, 1803)
truncatus (Ström, 1768)
- Poirieria**
actinophora (Dall, 1889)
- Typhinellus**
sowerbyi (Broderip, 1833)
- Ergalatax**
obscura Houart, 1996
- Urosalpinx**
cinerea (Say, 1822)
- Thais**
lacerus (Von Born, 1778)
- Stramonita**
haemastoma (Linné, 1766)
- Nucella**
lapillus (Linné, 1758)
- Orania**
fusulus (Brocchi, 1814)
- Cytheromorula**
grayi (Dall, 1889)
- Rapana**
rapiformis (Von Born, 1778)
venosa (Valenciennes, 1846)
- Family Coralliophilidae**
Coralliophila
basis (Dautzenberg & Fischer H., 1896)
brevis (de Blainville, 1832)
meiyendorffii (Calcara, 1845)
panormitana (Monterosato, 1869)
richardi (Fischer P., 1882)
sofiae (Aradas & Benoit, 1876)
squamosa (Bivona Ant. in Bivona And., 1838)
- Littaxis**
benoiti (Tiberi, 1855)
cariniferus (Sowerby G.B. I, 1834)
sentix (Bayer, 1971)
- Family Fasciolariidae**
Fasciolaria
lignaria (Linné, 1758)
- Fusinus**
agadirensis (Hadorn & Rolán, 1999)
amiantus (Dall, 1889)
bocagei (Fischer P., 1882)
boucheti (Hadorn & Ryall, 1999)
crassus (Pallary, 1901)
marmoratus (Philippi, 1846)

- maroccensis* (Gmelin, 1791)
profetai (Nofroni, 1982)
pulchellus (Philippi, 1844)
rostratus (Olivii, 1792)
rudis (Philippi, 1844)
rusticulus (Monterosato, 1880)
syracusanus (Linné, 1758)
tenerifensis (Hadorn & Rolán, 1999)
- Latirus*
armatus Adams A., 1854
- Family Buccinidae**
- Buccinum*
abyssorum Verrill & Smith, 1884
angulosum Gray J.E., 1839
belcheri Reeve, 1845
ciliatum (Fabricius O., 1780)
conoideum Sars G.O., 1878
cyanum Bruguière, 1792
fimmarchianum Verkrüzen, 1875
glaciale Linné, 1761
humphreysianum Bennet, 1824
hydrophanum Hancock, 1846
kjennerudae Bouchet & Warén, 1985
maltzani Pfeffer, 1886
micropoma Thorson, 1944
nivale Friele, 1882
oblitum Sykes, 1911
parvulum Verkrüzen, 1875
polare Gray J.E., 1839
pulchellum Sars G.O., 1878
scalariforme Möller, 1842
superangulare Thorson & Oskarsson in Oskarsson, 1962
tumidulum Sars G.O., 1878
undatum Linné, 1758
- Belomitra*
quadruplex (Watson, 1882)
- Beringius*
ossiania (Friele, 1879)
turttoni (Bean, 1834)
- Buccinulum*
cornuum (Linné, 1758)
- Chauvetia*
affinis (Monterosato, 1889)
brunnea (Donovan, 1804)
candidissima (Philippi, 1836)
crassior (Odner, 1932)
decorata Monterosato, 1889
gjunchiorum (Micali, 1999)
lefebvrei (Maravigna, 1840)
lineolata (Tiberi, 1868)
obliqua Nordsieck & Talavera, 1979
procerula Monterosato, 1889
recondita (Brugnone, 1873)
retifera (Brugnone, 1880)
submamilata (Bucquoy, Dautzenberg & Dollfus, 1882)
turritellata (Deshayes, 1835)
ventrosa Nordsieck, 1976
- Colubraria*
canariensis Nordsieck & Talavera, 1979
reticulata (de Blainville, 1829)
- Colus*
altus (Wood S., 1848)
frielei (Kantor, 1981)
glaber (Kobelt, 1876)
gracilis (da Costa, 1778)
holboelli (Möller, 1842)
howsei (Marshall, 1902)
islandicus (Mohr, 1786)
jeffreysianus (Fischer P., 1868)
kroeyeri (Möller, 1842)
latericeus (Möller, 1842)
sabini (Gray J.E., 1824)
turgidulus (Friele, 1877)
- verkruzezeni* (Kobelt, 1876)
- Engina*
leucozona (Philippi, 1843)
- Kryptos*
koehleri (Locard, 1896)
- Liomesus*
ovum (Turton, 1825)
- Mohnia*
abyssorum (Fischer P., 1883)
caelata (Verrill & Smith, 1880)
danielsseni (Friele, 1879)
glypta (Verrill, 1882)
mohni (Friele, 1877)
parva (Verrill & Smith, 1882)
simplex (Verrill, 1884)
- Neptunea*
antiqua (Linné, 1758)
brevicauda (Deshayes, 1832)
contraria (Linné, 1771)
despecta (Linné, 1758)
- Pisania*
striata (Gmelin, 1791)
- Pollia*
assimilis (Reeve, 1846)
dorbignyi (Payraudeau, 1826)
scabra Locard, 1886
scacchiana (Philippi, 1844)
viverratoides (d'Orbigny, 1840)
- Troschelia*
berniensis (King, 1846)
- Turrisiphon*
dalli (Friele in Tryon, 1881)
fenestratus (Turton, 1834)
lachesis (Mørch, 1869)
moebii (Dunker & Metzger, 1874)
undulatus (Friele in Tryon, 1881)
voerungi Bouchet & Warén, 1985
- Volutopsis*
deformis (Reeve, 1847)
norwegicus (Gmelin, 1791)
- Euthriostoma*
saharicum (Locard, 1897)
- Family Nassariidae**
- Nassarius*
antiquatus (Watson, 1897)
circumcinctus (Adams A., 1852)
coralligenus (Pallary, 1900)
corniculum (Olivii, 1792)
cuvieri (Payraudeau, 1826)
denticulatus (Adams A., 1852)
elatus (Gould, 1845)
frigens (Von Martens, 1878)
gibbosulus (Linné, 1758)
goreensis (Von Maltzan, 1884)
granum (Lamarck, 1822)
heynemani (Von Maltzan, 1884)
incrassatus (Ström, 1768)
johni (Monterosato, 1889)
lima (Dillwyn, 1817)
louisi (Pallary, 1912)
mutabilis (Linné, 1758)
nitidus (Jeffreys, 1867)
ovoideus (Locard, 1886)
pfeifferi (Philippi, 1844)
arcularius subsp. *plicatus* (Röding, 1798)
pygmæus (Lamarck, 1822)
macrodon subsp. *recidivus* (Von Martens, 1876)
reticulatus (Linné, 1758)
robustus (Monterosato, 1890)
tinei (Maravigna in Guérin, 1840)
tingitanus (Pallary, 1901)
turulosus (Risso, 1826)
unifasciatus (Kiener, 1834)
vaucheri (Pallary, 1906)
- Cyclope*

- neritea* (Linné, 1758)
pellucida Risso, 1826
Demoulia
obtusata (Link, 1807)
Family Columbellidae
Columbella
rustica (Linné, 1758)
adansonii Menke, 1853
Amphissa
acuteostriata (Philippi, 1844)
Anachis
aliceae (Pallary, 1900)
avaroides Nordsieck, 1975
savignyi (Moazzo, 1939)
troglodytes (Souverbie in Souverbie & Montrouzier, 1866)
Mitrella
broderipi (Sowerby G.B. I, 1844)
bruggeni van Aartsen, Menkhorst & Gittenberger, 1984
gervillii (Payraudeau, 1826)
minor (Scacchi, 1836)
nitidulina (Locard, 1897)
ocellata (Gmelin, 1791)
ocellina (Nordsieck, 1975)
pallaryi (Dautzenberg, 1927)
rac (Dautzenberg, 1891)
rosacea (Gould, 1840)
scripta (Linné, 1758)
- Family Costellariidae**
Vexillum
dexenum (Deshayes, 1834)
ebenus (Lamarck, 1811)
granum (Forbes, 1844)
hypatiae (Pallary, 1913)
savignyi (Payraudeau, 1826)
tricolor (Gmelin, 1791)
zebrinum (d'Orbigny, 1840)
- Family Marginellidae**
Marginella
aronnax Bouchet & Warén, 1985
colomborum (Bozzetti, 1995)
glabella (Linné, 1758)
hesperia Sykes, 1905
impudica Fischer P., 1883
limbata Lamarck, 1822
marocana Locard, 1897
subturrita Fischer P., 1883
Volvarina
ampelusica Monterosato, 1906
ingolfi Bouchet & Warén, 1985
mitrella (Risso, 1826)
oceanica Gofas, 1989
olivaeformis (Kiener, 1834)
ptychasthenia Gofas, 1989
roberti Bavay, 1917
Granulina
boucheti Gofas, 1992
gofasi Smriglio & Mariottini, 1996
guancha (d'Orbigny, 1840)
gubbiolii (Smriglio & Mariottini, 1999)
guttula (La Perna, 1999)
marginalata (Bivona, 1832)
melitensis (Smriglio, Mariottini & Rufini, 1998)
minusculina (Locard, 1897)
occulta (Monterosato, 1869)
torosa Gofas, 1992
vanharenii (van Aartsen, Menkhorst & Gittenberger, 1984)
- Family Cystiscidae**
Gibberula
abyssicola Locard, 1897
caelata (Monterosato, 1877)
epigrus (Reeve, 1865)
- jansseni* van Aartsen, Menkhorst & Gittenberger, 1984
lazaroi (Contreras, 1992)
miliaria (Linné, 1758)
oryza (Lamarck, 1822)
philippii (Monterosato, 1878)
recondita Monterosato, 1884
secreta Monterosato, 1889
turgidula (Locard & Caziot, 1900)
vignali (Dautzenberg & Fischer H., 1896)
- Persicula**
canaryensis (Clover, 1972)
- Family Mitridae**
Mitra
corniculum (Linné, 1758)
cornea (Lamarck, 1811)
zonata Marryat, 1818
- Family Olividae**
Olivella
oteroi Bermejo, 1979
- Family Pseudolivididae**
Benthobia
tryonii Dall, 1889
- Family Turbinellidae**
Latiromitra
cryptodon (Fischer P., 1882)
Metzgeria
alba (Wyville-Thompson, 1873)
gagei Bouchet & Warén, 1985
- Family Volutidae**
Cymbium
cucumis Röding, 1798
olla (Linné, 1758)
tritonis (Broderip, 1830)
Ampulla
priamus (Gmelin, 1791)
- Family Volutomitridae**
Volutomitra
groenlandica (Beck in Möller, 1842)
Microvoluta
superstes Bouchet & Warén, 1985
- Family Cancillariidae**
Cancillaria
cancellata (Linné, 1767)
coronata Scacchi, 1835
similis Sowerby G.B. I, 1833
Axerella
minima (Reeve, 1856)
Broccchinia
azorica (Bouchet & Warén, 1985)
clenchi Petit R.E., 1986
cornidei (Altimira, 1978)
nodosa (Vertill & Smith, 1885)
pusilla (Adams H., 1869)
Solatia
piscatoria (Gmelin, 1791)
Admete
contabulata Friele, 1879
sadko Gorbulov, 1946
viridula (Fabricius O., 1780)
- Iphinopsis**
alba Bouchet & Warén, 1985
fuscoapicata Bouchet & Warén, 1985
inflata (Friele, 1879)
- Family Drillidae**
Crassopleura
maravignae (Bivona Ant. in Bivona And., 1838)
Spirotropis
azoricus Bouchet & Warén, 1980
centimatus (Dall, 1889)
modiolus (de Cristofori & Jan, 1832)
monterosatoi Locard, 1897

Family Turridae*Fusiturris*

- similis* (Bivona Ant. in Bivona And., 1838)
undatiruga (Bivona Ant. in Bivona And., 1838)

Haedropleura

- flexicosta* (Monterosato, 1884)
septangularis (Montagu, 1803)

Micropleurotoma

- melvilli* (Sykes, 1906)
spirotropoides (Thiele, 1925)
travailleurii Bouchet & Warén, 1980

Ancistrosyrinx

- clytotropis* (Sykes, 1906)

Irenosyrinx

- hypomela* (Dall, 1889)

Leucosyrinx

- verrillii* (Dall, 1881)

Family Conidae*Borsonia*

- hirondellae* (Dautzenberg, 1891)

Typhlomangelia

- nivalis* (Lovén, 1846)

Comarmondia

- gracilis* (Montagu, 1803)

Corinnaeturris

- leucomata* (Dall, 1881)

Mitrolumna

- crenipicta* Dautzenberg, 1889
dalli (Dautzenberg & Fischer H., 1896)
melitensis (Mifsud, 1993)
olivoidea (Cantraine, 1835)
smithi (Dautzenberg & Fischer H., 1896)
wilheminae van Aartsen, Menhorst & Gittenberger, 1984

Drilliola

- emendata* (Monterosato, 1872)
megalacme (Sykes, 1906)

Microdrillia

- loprestiana* (Calcaria, 1841)
priuina (Watson, 1881)

Oenopota

- angulosa* (Sars G.O., 1878)
assimilis (Sars G.O., 1878)
bergensis (Friele, 1886)
cinerea (Møller, 1842)
conoidea (Sars G.O., 1878)
convexigyrta Bouchet & Warén, 1980
declivis (Lovén, 1846)
dictyophora Bouchet & Warén, 1980
elegans (Møller, 1842)
exarata (Møller, 1842)
graphica (Locard, 1897)
arpa (Dall, 1884)
harpularia (Couthouy, 1838)
impressa (Mørch, 1869)
nobilis (Møller, 1842)
novajasemljensis (Leche, 1878)
obliqua (Sars G.O., 1878)
ovalis (Friele, 1877)
pingelii (Møller, 1842)
pyramidalis (Ström, 1788)
rufa (Montagu, 1803)
rugulata (Troschel, 1866)
scalaris (Møller, 1842)
scalaroides (Sars G.O., 1878)
spitzbergensis (Friele, 1886)
tenuicostata (Sars G.O., 1878)
trevelliiana (Turton, 1834)
turricula (Montagu, 1803)
violacea (Mighels & Adams, 1842)
viridula (Møller, 1842)

Obesotoma

- gigas* (Verkrüzen, 1875)
simplex (Middendorff, 1849)
woodiana (Møller, 1842)

Bela

- brachystoma* (Philippi, 1844)
costulata (Risso, 1826)
cycladensis (Reeve, 1845)
decussata (Locard, 1897)
laevigata (Philippi, 1836)
menhorsti van Aartsen, 1988
nebula (Montagu, 1803)
oceanica (Locard, 1897)
ornata (Locard, 1897)
septenvillei (Dautzenberg, 1913)
zonata (Locard, 1892)
- Benthomangelia*
- antonia* (Dall, 1881)
decapitata Bouchet & Warén, 1980
macra (Watson, 1881)
- Clathromangelia*
- granum* (Philippi, 1844)
loiselieri Oberling, 1970
strigillata Pallary, 1904
- Fehria*
- taprurensis* (Pallary, 1904)
zenetouae van Aartsen, 1988
- Mangelia*
- attenuata* (Montagu, 1803)
barashi (van Aartsen & Fehr-de Wal, 1978)
bertrandii (Payraudeau, 1826)
brusinae van Aartsen & Fehr-de Wal, 1978
caerulans (Philippi, 1844)
callosa (Nordsieck, 1977)
coarctata (Forbes, 1840)
costata (Donovan, 1804)
difficilis (Locard & Caziot, 1900)
fieldeni (van Aartsen & Fehr-de Wal, 1978)
goodallii Reeve, 1846
indistincta (Monterosato, 1875)
jerbaensis (Della Bella & Spada in Chirli, 1997)
multilineolata (Deshayes, 1835)
nuperrima (Tiberi, 1855)
paciniana (Calcaria, 1839)
pallaryi (Nordsieck, 1977)
payraudeaui (Deshayes, 1835)
pontica Milaschewitsch, 1908
sandrii (Brusina, 1865)
scabrida Monterosato, 1890
secreta (van Aartsen & Fehr-de Wal, 1978)
serga (Dall, 1881)
sicula Reeve, 1846
smithii (Forbes, 1840)
stossiciana Brusina, 1869
taeniata (Deshayes, 1835)
tenuicostata Brugnone, 1868
unifasciata Deshayes, 1835
vauquelini (Payraudeau, 1826)
- Raphitoma*
- aequalis* Jeffreys, 1867
alternans (Monterosato, 1884)
arnoldi Pallary, 1904
atropurpurea (Locard & Caziot, 1900)
bosilliana (Sulliotti, 1889)
bracteata (Pallary, 1904)
concinna (Scacchi, 1836)
corbis (Potiez & Michaud, 1838)
cordieri (Payraudeau, 1826)
cylindracea (Locard & Caziot, 1900)
densa (Monterosato, 1884)
divae Carrozza, 1984
echinata (Brocchi, 1814)
erronea (Monterosato, 1884)
histrix Bellardi, 1847
horrida (Monterosato, 1884)
leufroyi (Michaud, 1828)
linearis (Montagu, 1803)
mirabilis (Pallary, 1904)
nivea (Marshall in Sykes, 1906)

- papillosa* (Pallary, 1904)
philberti (Michaud, 1829)
pruinosa (Pallary, 1906)
pseudohystrix Sykes, 1906
pupoidea (Monterosato, 1884)
purpurea (Montagu, 1803)
volutella (Kiener, 1839)
- Nepotilla**
amoena (Sars G.O., 1878)
- Teretia**
teres (Reeve, 1844)
- Tereiopsis**
thaumastopsis (Dautzenberg & Fischer H., 1896)
- Conopleura**
aliena Smriglio, Mariottini & Calascibetta, 1999
- Gymnobela**
abyssorum (Locard, 1897)
agassizi (Verrill & Smith, 1880)
aquilarum (Watson, 1881)
chyla (Watson, 1881)
emertoni (Verrill & Smith, 1884)
engonia Verrill, 1884
frielei (Verrill, 1885)
fulvotincta (Dautzenberg & Fischer H., 1896)
homeotata (Watson, 1886)
lamyi (Dautzenberg, 1925)
leptoglypta (Dautzenberg & Fischer H., 1896)
physanor (Watson, 1886)
pyrrhogamma (Dautzenberg & Fischer H., 1896)
subaraneosa (Dautzenberg & Fischer H., 1896)
watsoni (Dautzenberg, 1889)
- Pleurotomella**
anceyi (Dautzenberg & Fischer H., 1897)
benedicti Verrill & Smith, 1884
bureaui (Dautzenberg & Fischer H., 1897)
coelaropha (Dautzenberg & Fischer H., 1896)
demosia (Dautzenberg & Fischer H., 1896)
eurybroscha (Dautzenberg & Fischer H., 1896)
gibbera Bouchet & Warén, 1980
lottae Verrill, 1885
megalembryon (Dautzenberg & Fischer H., 1896)
obesa Bouchet & Warén, 1980
packardi Verrill, 1872
sandersoni Verrill, 1884
- Neopleurotomoides**
callembryon (Dautzenberg & Fischer H., 1897)
distinctus Bouchet & Warén, 1980
- Thesbia**
nana (Lovén, 1846)
- Theta**
chariessa (Watson, 1881)
lyronuclea (Clarke A.H., 1959)
vayssierei (Dautzenberg, 1925)
- Famelia**
catharinæ (Verrill & Smith, 1884)
mirmidina (Dautzenberg & Fischer H., 1896)
monoceros (Watson, 1881)
monotropis (Dautzenberg & Fischer H., 1896)
- Bathybela**
nudator (Locard, 1897)
tenelluna (Locard, 1897)
- Xanthodaphne**
bruneri (Verrill, 1884)
charcotiana Bouchet & Warén, 1980
dalmasi (Dautzenberg & Fischer H., 1897)
heterogramma (Odhner, 1960)
leptalea (Bush, 1893)
- Aliceia**
aenigmatica Dautzenberg & Fischer H., 1897
- Lusitanops**
blanchardi (Dautzenberg & Fischer H., 1896)
bullioides (Sykes, 1906)
cingulatus Bouchet & Warén, 1980
expansus (Sars G.O., 1878)
hyaloïdes (Dautzenberg, 1925)
- lusitanicus* (Sykes, 1906)
macrapex Bouchet & Warén, 1980
sigmoideus Bouchet & Warén, 1980
- Phymorhynchus**
alberti (Dautzenberg & Fischer H., 1906)
chevreuxi (Dautzenberg & Fischer H., 1897)
sulciferus (Bush, 1893)
- Taranis**
borealis Bouchet & Warén, 1980
laevisculpta Monterosato, 1880
malmi (Dall, 1889)
moerchi (Malm, 1863)
- Conus**
mediterraneus Hwass in Bruguière, 1792
vayssierei Pallary, 1906
- Family Architectonicidae**
- Basisulcata*
lepidia (Bayer, 1942)
- Discotectonica*
discus (Philippi, 1844)
- Heliacus*
canariensis Nordsieck, 1982
contextus (Seguenza L., 1903)
jeffreysianus (Tiberi, 1867)
subvariegatus (d'Orbigny, 1852)
- Philippia*
hybrida (Linné, 1758)
- Pseudomalaxis*
zancleus (Philippi, 1844)
- Pseudotorinia*
architae (Costa O.G., 1841)
- Solatisonax*
alleryi (Seguenza G., 1876)
bannocki (Melone & Taviani, 1980)
certesi (Dautzenberg & Fischer H., 1896)
dolfusi (Dautzenberg & Fischer H., 1896)
- Spirolaxis*
centrifugus (Monterosato, 1890)
clenchi Jaume & Borro, 1916
lamelliferus (Rehder, 1935)
- Family Mathildidae**
- Mathilda*
cochlaeformis Brugnone, 1873
coronata Monterosato, 1875
gemmulata Semper, 1865
quadricarinata (Brocchi, 1814)
retusa Brugnone, 1873
- Tuba*
jeffreysi (Dall, 1889)
pinguis (Jeffreys, 1884)
- Turritellopsis*
stimpsoni (Dall, 1919)
- Brookesena*
turrita (Warén, 1996)
- Family Rissoellidae**
- Rissoella*
diaphana (Alder, 1848)
globularis (Forbes & Hanley, 1853)
opalina (Jeffreys, 1848)
- Family Omalogyridae**
- Omalogyra*
atomus (Philippi, 1841)
discula Palazzi, 1988
simplex (Costa O.G., 1861)
undosa Palazzi, 1988
- Ammonicera*
fischeriana (Monterosato, 1869)
ligneua (Palazzi, 1988)
rota (Forbes & Hanley, 1850)
rotundata (Palazzi, 1988)
- Retrotortina*
fuscata Chaster, 1896
- Family Xylodisculidae**
Xylodiscula

- boucheti* Warén, Carrozza & Rocchini in Warén, 1992
lens Warén, 1992
- Family Cornirostridae**
- Tomura*
depressa (Granata-Grillo, 1877)
- Noerrevangia*
fragilis Warén & Schander in Warén, Gofas & Schander, 1993
- Family Hyalogyrinidae**
- Xenoskenea*
pellucida (Monterosato, 1874)
- Hyalogryra*
zibrowii Warén in Warén, Carrozza & Rocchini, 1997
- Hyalogyrina*
amphorae Warén, Carrozza & Rocchini, 1997
- Family Cimidae**
- Cima*
cuticulata Warén, 1993
cylindrica (Jeffreys, 1856)
inconspicua Warén, 1993
melitensis Mifsud, 1998
minima (Jeffreys, 1858)
- Murchisonella*
columna (Hedley, 1907)
- Graphis*
albida (Kanmacher, 1798)
gracilis (Monterosato, 1874)
striata (Jeffreys, 1884)
- Family Orbitestellidae**
- Orbitestella*
dariae (Liuzzi & Zucchi Stolfa, 1979)
- Family Pyramidellidae**
- Tiberia*
minusculoides (van Aartsen, Gittenberger & Goud, 1998)
minuscula (Monterosato, 1880)
- Atomiscala*
islandica Warén, 1989
- Rissopsetia*
islandica Warén, 1989
- Chrysallida*
angulosa (Monterosato, 1889)
bjoernsoni Warén, 1991
boucheti (Peñas & Rolán, 1999)
brattstroemi Warén, 1991
brevicula (Jeffreys, 1883)
brusinai (Cossmann, 1921)
canariensis Nordsieck & Talavera, 1979
clathrata (Jeffreys, 1848)
decussata (Montagu, 1803)
dollfusi (Kobelt, 1903)
elegans (de Folin, 1870)
emaciata (Brusina, 1866)
excavata (Philippi, 1836)
eximia (Jeffreys, 1849)
fenerata (Jeffreys, 1848)
fischeri (Hornung & Mermod, 1925)
flexuosa (Monterosato, 1874)
ghisotti (van Aartsen, 1984)
hoeisaeteri Warén, 1991
indistincta (Montagu, 1808)
intermixta (Monterosato, 1884)
interstincta (Adams J., 1797)
jeffreysiana (Monterosato, 1884)
juliæ (de Folin, 1872)
limitum (Brusina in de Folin & Périer, 1876)
maiae (Hornung & Mermod, 1924)
marthinae (Nofroni & Schander, 1994)
monozena (Brusina, 1869)
monterosatii (Clessin, 1900)
moolenbeeki Amati, 1987
nivosa (Montagu, 1803)
palazzii Micali, 1984
pellucida (Dillwyn, 1817)
- pirinthella* (Melvill, 1910)
sarsi Nordsieck, 1972
- seamounti* (Peñas & Rolán, 1999)
sigmoidea (Monterosato, 1880)
stefanisi (Jeffreys, 1869)
sublustris (Friele, 1886)
suturalis (Philippi, 1844)
terebellum (Philippi, 1844)
truncatula (Jeffreys, 1850)
undata (Watson, 1897)
- Kejdonia*
cachiai (Mifsud, 1998)
- Pseudoscilla*
bilirata (de Folin, 1870)
pauciemersa Peñas & Rolán, 1999
- Odostomella*
doliolum (Philippi, 1844)
- Clathrella*
clathrata (Philippi, 1844)
- Euparthenia*
bulinea (Lowe, 1841)
humboldti (Risso, 1826)
- Iolaea*
neofelixoides (Nomura, 1936)
- Monotygra*
amoena (Adams A., 1853)
modesta (de Folin, 1870)
- Oscilla*
jocosa Melvill, 1904
- Cingulina*
isseli (Tryon, 1886)
- Styloptygma*
beatrix (Melvill, 1911)
- Eulimella*
acicula (Philippi, 1836)
ataktos Warén, 1991
atlantis (Peñas & Rolán, 1999)
bogii van Aartsen, 1995
carmina (Peñas & Micali, 1999)
cerullii (Cossmann, 1916)
compactilis (Jeffreys, 1867)
cossignani van Aartsen, 1995
neoattenuata (Gaglini, 1992)
phaula (Dautzenberg & Fischer H., 1896)
protofunis (Peñas & Rolán, 1999)
scillae (Scacchi, 1835)
similebala (Peñas & Rolán, 1999)
sudis (Peñas & Rolán, 1999)
unifasciata (Forbes, 1844)
vanhareni (van Aartsen, Gittenberger & Goud, 1998)
ventricosa (Forbes, 1844)
verduini (van Aartsen, Gittenberger & Goud, 1998)
- Syrnola*
cinctella (Adams A., 1860)
fasciata Jickeli, 1882
minuta Adams H., 1869
wenzi Nordsieck, 1972
- Odostomia*
aartseni Nofroni, 1988
acuta Jeffreys, 1848
angusta Jeffreys, 1867
bernardi (van Aartsen, Gittenberger & Goud, 1998)
bismichaelis Sacco, 1892
bulimulus Monterosato, 1874
buzzurroi (Peñas & Rolán, 1999)
carrozzai van Aartsen, 1987
clavulina Fischer P., 1877
conoidea (Brocchi, 1814)
conspicua Alder, 1850
duurenii (van Aartsen, Gittenberger & Goud, 1998)
electa Jeffreys, 1883
erjaveciana Brusina, 1869
eulimoides Hanley, 1844
fehrae (van Aartsen, Gittenberger & Goud, 1998)
franki (Peñas & Rolán, 1999)

- fusulus* Monterosato, 1878
glabrata Forbes & Hanley, 1850
harveyi van Aartsen & Smith, 1996
hieroensis (Peñas & Rolán, 1999)
ignorata (Monterosato, 1917)
kromi van Aartsen, Menkhorst & Gittenberger, 1984
kuiperi (van Aartsen, Gittenberger & Goud, 1998)
lesuroiti (Peñas & Rolán, 1999)
lorellae Micali, 1987
lorioli (Hornung & Mermod, 1924)
lukisi Jeffreys, 1859
marci (van Aartsen, Gittenberger & Goud, 1998)
microeques (Rolán & Templado in Peñas & Rolán, 1999)
nitens Jeffreys, 1870
oblongula Marshall, 1895
odostomella (Peñas & Rolán, 1999)
omphaloessa Watson, 1897
palmaensis (Peñas & Rolán, 1999)
plicata (Montagu, 1803)
prona (Peñas & Rolán, 1999)
restii (Peñas & Rolán, 1999)
rutor (Nofroni & Schander, 1994)
scalaris MacGillivray, 1843
sicula Philippi, 1851
silesui Nofroni, 1988
striolata Forbes & Hanley, 1850
suboblonga Jeffreys, 1884
testicula (Peñas & Rolán, 1999)
turgida Sars G.O., 1878
turriculata Monterosato, 1869
turrita Hanley, 1844
umbilicaris (Malm, 1863)
umbilicatissima (Peñas & Rolán, 1999)
indentata (Montagu, 1803)
verduini van Aartsen, 1987
winfredi (Peñas & Rolán, 1999)
- Aartsenia*
candida (Møller, 1842)
- Liostomia*
afzelii Warén, 1991
clavulus (Lovén, 1846)
eburnea (Stimpson, 1851)
hansgei Warén, 1991
mamoi Mif sud, 1993
- Menestho*
albula (Fabricius O., 1780)
truncatula Odhner, 1915
- Noemiamea*
dolioliformis (Jeffreys, 1848)
- Ordina*
anceps (Gaglini, 1992)
coarctata (Sars G.O., 1878)
crystallina Locard, 1892
diaphana (Jeffreys, 1848)
dilucida (Monterosato, 1884)
divisa (Adams J., 1797)
mosti (van Aartsen, Gittenberger & Goud, 1998)
normani (Friele, 1886)
obliqua (Alder, 1844)
scandens (Monterosato, 1884)
vitrea (Brusina, 1866)
warreni (Thompson W., 1845)
- Turbanilla*
abradi (Fischer-Piette & Nicklès, 1946)
acutissima Monterosato, 1884
acuta (Donovan, 1804)
atlantica (Locard, 1897)
attenuata (Jeffreys, 1884)
amoena (Monterosato, 1878)
crenata (Brown, 1827)
delicata Monterosato, 1874
edgari (Melvill, 1896)
eodem (Peñas & Rolán, 1999)
formosa (Jeffreys, 1848)
- fulgidula* (Jeffreys, 1884)
gofasi (Peñas & Rolán, 1999)
gradata Bucquoy, Dautzenberg & Dollfus, 1883
hamata Nordsieck, 1972
hamonvillei (Dautzenberg & Fischer H., 1896)
hannoni Pallary, 1920
hoecki (Dautzenberg & Fischer H., 1896)
inobservata (Peñas & Rolán, 1999)
internodula (Wood S., 1848)
jéffreysii (Forbes & Hanley, 1851)
lactea (Linné, 1758)
macandreae (Adams H., 1871)
magnifica (Seguenza G., 1879)
mediocris (Peñas & Rolán, 1999)
micans (Monterosato, 1875)
mirifica Pallary, 1904
multilirata (Monterosato, 1875)
nofronii Peñas & Rolán, 1997
obliquata (Philippi, 1844)
pallaryi Dautzenberg, 1910
paucistriata (Jeffreys, 1884)
pauperata Locard, 1897
postacuticostata Sacco, 1892
pumila Seguenza G., 1876
pusilla (Philippi, 1844)
rectogallica Sacco, 1892
rosewateri Corgan & van Aartsen, 1993
rufescens (Forbes, 1846)
rufa (Philippi, 1836)
semicostata (Jeffreys, 1884)
sinuosa (Jeffreys, 1884)
striatula (Linné, 1758)
subulina Monterosato, 1889
syrtensis van Aartsen, 1981
vallanti (Dautzenberg & Fischer H., 1896)
- Family Ebalidae**
Ebala
nitidissima (Montagu, 1803)
pointeli (de Folin, 1868)
striatula (Jeffreys, 1856)
trigonostoma (de Folin, 1872)
- Family Tjaernoiidae**
Tjaernoeia
boucheti Warén, 1991
equisita (Jeffreys, 1883)
unisulcata (Chaster, 1897)
- Family Acteonidae**
Acteon
monterosatoi Dautzenberg, 1889
tornatilis (Linné, 1758)
- Tomlinula*
turrita (Watson, 1886)
- Inopinodon*
azoricus (Locard, 1897)
- Crenilabrum*
exile (Jeffreys, 1870)
- Liocarenus*
globulinus (Forbes, 1844)
- Japonacteon*
pusillus (MacGillivray, 1843)
- Callostracon*
amabile (Watson, 1886)
tyrrhenicum Smriglio & Mariottini, 1996
- Family Hydatinidae**
Hydatina
vesicaria (Lightfoot, 1786)
- Micromelo*
undatus (Bruguière, 1792)
- Family Diaphanidae**
Diaphana
abyssalis (Schiøtte, 1999)
cretica (Forbes, 1844)
flava (Watson, 1897)
glacialis (Odhner, 1907)

- globosa* (Jeffreys, 1864)
hiemalis (Couthouy, 1839)
lactea (Jeffreys, 1877)
makarovi (Gorbunov, 1946)
marshalli (Sykes, 1905)
minuta Brown, 1827
quadrata (Monterosato, 1874)
Colobocephalus
costellatus Sars M., 1870
striatulus (Monterosato, 1874)
Colpodaspis
pusilla Sars M., 1870
Rhinodiaphana
ventricosa (Jeffreys, 1865)
Toledonia
limnaeoides (Odhner, 1913)
Bogasonia
volutoides Warén, 1989
- Family Retusidae**
Retusa
candidula (Locard, 1892)
fourierii (Audouin, 1826)
leptoeneilema (Brusina, 1866)
mammillata (Philippi, 1836)
mariei (Dautzenberg, 1889)
minutissima (Monterosato, 1878)
obtusa (Montagu, 1803)
pellucida (Sars G.O., 1878)
piriformis Monterosato, 1878
tornata (Watson, 1886)
truncatula (Bruguière, 1792)
- Relichna**
simplex (Locard, 1897)
- Cylchnina**
canariensis Nordsieck & Talavera, 1979
crebrisculpta Monterosato, 1884
girardi (Audouin, 1826)
laevisculpta (Granata-Grillo, 1877)
nitidula (Lovén, 1846)
robagliana (Fischer P. in de Folin, 1869)
tenerifensis Nordsieck & Talavera, 1979
umbilicata (Montagu, 1803)
- Pyrunculus**
hoernesii (Weinkauff, 1866)
ovatus (Jeffreys, 1870)
spretus (Watson, 1897)
- Volvulella**
acuminata (Bruguière, 1792)
- Family Ringiculidae**
Ringicula
auriculata (Ménard de la Groye, 1811)
pulchella Morlet, 1880
leptocheila Brugnone, 1873
- Family Runcinidae**
Runcina
adriatica Thompson T., 1980
africana Pruvot-Fol, 1953
aurata Garcia-Gomez, Lopez, Luque & Cervera, 1988
bahiensis Cervera, Garcia-Gomez & Garcia, 1991
brenkoae Thompson T., 1980
capreensis Mazzarelli, 1894
coronata (de Quatrefages, 1844)
ferruginea Kress, 1977
macrodenticulata Garcia-Gomez & Lopez de la Cuadra, 1990
zavodnikii Thompson T., 1980
- Family Bullidae**
Bulla
amygdala Dillwyn, 1817
striata Bruguière, 1792
- Family Haminoeidae**
Haminoea
cymoelia Monterosato, 1917
elegans Leach, 1852
- exigua* Schaefer, 1992
fusari Alvarez, Garcia & Villani, 1983
hydatis (Linné, 1758)
navicula (da Costa, 1778)
orbignyana (de Féussac, 1822)
orteai Talavera, Murillo & Templado, 1987
templanoi Garcia, Perez-Hurtado & Garcia-Gomez, 1991
- Atys**
blainvilliana (Récluz, 1843)
jeffreysi (Weinkauff, 1866)
macandrewi Smith E.A., 1872
- Cylchnium**
africanum (Locard, 1897)
oliviforme (Watson, 1883)
- Smaragdinella**
algirae (Adams A. in Sowerby G.B. II, 1850)
- Weinkauffia**
turgidula (Forbes, 1844)
- Family Philinidae**
Philine
angulata Jeffreys, 1867
aperta (Linné, 1767)
azorica Bouchet, 1975
calva van der Linden, 1995
catena (Montagu, 1803)
complanata Watson, 1897
condensa (van der Linden, 1995)
denticulata (Adams J., 1800)
desmotis Watson, 1897
infornata Pilsbry, 1895
intricata Monterosato, 1875
lima (Brown, 1827)
monilifera Bouchet, 1975
monterosati Monterosato, 1874
monterosatoi (Sykes, 1905)
punctata (Adams J., 1800)
quadrata (Wood S., 1839)
rugulosa (Dautzenberg & Fischer H., 1896)
scabra (Müller, 1784)
trachyostraca Watson, 1897
- Johania**
retifera (Forbes, 1844)
- Laona**
finmarchica (Sars M., 1858)
flexuosa (Sars M., 1870)
pruinosa (Clark W., 1827)
- Family Aglajidae**
Aglaja
berrieri (Dieuzeide, 1935)
maderensis (Watson, 1897)
taila Marcus Ev. & Er., 1966
tricolorata Renier, 1807
- Chelidonura**
fulvipunctata Baba, 1938
italica Sordi, 1980
orchidea Perrone, 1990
- Melanochlamys**
seurati (Vayssièvre, 1926)
- Philinopsis**
depicta (Renier, 1807)
- Family Cylichnidae**
Cylichna
alba (Brown, 1827)
crossei Bucquoy, Dautzenberg & Dollfus, 1886
cylindracea (Pennant, 1777)
lemchei Bouchet & Warén, 1979
occulta (Mighels & Adams, 1842)
parvula Jeffreys, 1883
propecylindracea (de Gregorio, 1890)
- Acteocina**
mucronata (Philippi, 1849)
- Roxania**
monterosatoi Dautzenberg & Fischer H., 1896

- pinguicula* (Seguenza G., 1879)
utriculus (Brocchi, 1814)
- Scaphander**
gracilis Watson, 1886
lignarius (Linné, 1758)
nobilis Verrill, 1884
punctostriatus (Mighels & Adams, 1842)
- Melosaphander**
imperceptus Bouchet, 1975
- Mamillocylichna**
abyssicola Bouchet, 1975
richardi (Dautzenberg, 1889)
- Utriculopsis**
vitrea Sars M., 1870
- Family Gasteropteridae**
Gasteropteron
meckeli Kosse, 1813
- Family Philinoglossidae**
Philinoglossa
helgolandica Hertling, 1932
latosoleata Salvini-Plawen, 1973
praelongata Salvini-Plawen, 1973
remanei Marcus Ev. & Er., 1958
- Family Cavoliniidae**
Cavolinia
gibbosa (d'Orbigny, 1835)
inflexa (Lesueur, 1813)
tridentata (Forskål, 1775)
uncinata (Rang, 1829)
- Diacavolinia*
constricta (van der Spoel, Bleeker & Kobayashi, 1993)
deshayesi (van der Spoel, Bleeker & Kobayashi, 1993)
limbata (d'Orbigny, 1836)
atlantica van der Spoel, Bleeker & Kobayashi, 1993
- Diacria*
atlantica (Dupont in Bontes & van der Spoel, 1998)
quadridentata (Lesueur, 1821)
rampali (Dupont, 1979)
rubecula (Bontes & van der Spoel, 1998)
trispinosa (Lesueur, 1821)
- Clio*
cuspidata (Bosc, 1802)
pyramidata Linné, 1767
- Creesis*
acicula Rang, 1828
virgula Rang, 1828
- Hyalocylis*
obtusa Di Geronimo, 1974
striata (Rang, 1828)
- Styliola*
subula (Quoy & Gaimard, 1827)
- Cuvierina*
columnella (Rang, 1827)
- Family Limacinidae**
Limacina
bulimoides (d'Orbigny, 1836)
helicina Phipps, 1774
helicoides Jeffreys, 1877
inflata (d'Orbigny, 1836)
lesueuri (d'Orbigny, 1836)
retroversa (Fleming, 1823)
trochiformis (d'Orbigny, 1836)
- Family Cymbuliidae**
Cymbulia
parvidentata Pelseneer, 1888
peronii Lamarck, 1819
- Gleba*
chrysosticta (Troschel, 1854)
cordata Forskål, 1776
- Corolla*
spectabilis Dall, 1871
- Family Desmopteridae**
- Desmopterus*
cirropterus Gegenbaur, 1855
papilio Chun, 1889
- Family Peracidae**
Peracle
bispinosa Pelseneer, 1888
diversa (Monterosato, 1875)
reticulata (d'Orbigny, 1836)
triacantha (Fischer P., 1882)
- Family Clionidae**
Clione
limacina (Phipps, 1774)
Fowlerina
punctata (Tesch, 1903)
zetesios Pelseneer, 1906
- Paraclione*
flavescens (Gegenbaur, 1855)
longicaudata (Souleyet, 1852)
- Thalassopterus*
zancleus Kwietniewski, 1910
- Thliptodon*
gegenbauri Boas, 1886
- Family Cliopsidae**
Cliopsis
krohni Troschel, 1854
- Family Notobranchaeidae**
Notobranchaea
bleekerae van der Spoel & Pafort, 1985
mcdonaldi Pelseneer, 1886
- Schleschia*
tetrabranchiata (Bonnevie, 1913)
- Family Pneumodermatidae**
Pneumoderma
mediterraneum van Beneden, 1838
violaceum d'Orbigny, 1836
- Pneumodermopsis*
canephora Pruvot-Fol, 1924
ciliata (Gegenbaur, 1855)
paucidens (Boas, 1886)
pupula Pruvot-Fol, 1926
teschi van der Spoel, 1973
- Family Oxynoidea**
Oxynoe
olivacea Rafinesque, 1814
- Lobiger*
serradifalci (Calcaria, 1840)
- Family Volvatellidae**
Ascobulla
fragilis (Jeffreys, 1856)
- Family Elysiidae**
Elysia
j lava Verrill, 1901
gordanae Thompson & Jaklin, 1988
hetta Perrone, 1990
ornata (Swainson, 1840)
picta Verrill, 1901
timida (Risso, 1818)
translucens Pruvot-Fol, 1957
viridis (Montagu, 1804)
- Thuridilla*
hopei (Vérany, 1853)
- Family Boselliidae**
Bosellia
levis Fernandez-Ovies & Ortea, 1986
mimetica Trinchese, 1891
- Family Hermaeidae**
Polybranchia
borgniini (Trinchese, 1896)
viridis (Deshayes, 1857)
- Aplysiopsis*
elegans Deshayes, 1864
formosa Pruvot-Fol, 1953

- Cyerce**
crystallina (Trinchesse, 1881)
graeca Thompson T., 1988
- Hermaea**
bifida (Montagu, 1815)
boucheti (Cervera, Garcia-Gomez & Ortea, 1991)
paucicirra Pruvot-Fol, 1953
variopicta (Costa A., 1869)
- Caliphylla**
mediterranea Costa A., 1869
- Family Costasiellidae**
Costasiella
virescens Pruvot-Fol, 1951
- Family Stiligeridae**
Alderella
comosa (Costa A., 1867)
Alderia
modesta (Lovén, 1844)
Calliopaea
bellula d'Orbigny, 1837
oophaga Lemche in Gascoigne & Sartory, 1974
rissoana Milne-Edwards, 1842
souleyeti Vérany, 1846
- Ercolania*
coerulea Trinchesse, 1892
funerea (Costa A., 1867)
lozanoi Ortea, 1981
viridis (Costa A., 1866)
- Placida*
cremoniana (Trinchesse, 1892)
dendritica (Alder & Hancock, 1843)
saronica (Thompson T., 1988)
tardyi (Trinchesse, 1873)
verticillata Ortea, 1981
viridis (Trinchesse, 1873)
- Stiliger*
llerai Ortea, 1981
niger Lemche, 1935
- Family Limapontiidae**
Limapontia
capitata (Müller O.F., 1774)
depressa Alder & Hancock, 1862
senestra (de Quatrefages, 1844)
- Family Umbraculidae**
Umbraculum
umbraculum (Röding, 1798)
Spiricella
unguicula Rang, 1828
- Family Tylodinidae**
Anidolya
duebeni (Lovén, 1846)
Tylodina
perversa (Gmelin, 1791)
- Family Pleurobranchidae**
Pleurobranchus
forskali (Rüppell & Leuckart, 1831)
membranaceus (Montagu, 1815)
testudinarius Cantraine, 1835
- Berthella*
aurantiaca (Risso, 1818)
dautzenbergi (Watson, 1897)
elongata (Cantraine, 1835)
lowei (Watson, 1897)
ocellata (delle Chiaje, 1830)
plumula (Montagu, 1803)
sideralis Lovén, 1846
stellata (Risso, 1826)
- Berthellina*
citrina (Rüppell & Leuckart, 1831)
engeli Gardiner, 1936
- Pleurobranchaea*
inconspicua Bergh, 1897
meckelii Meckelin Leue, 1813
- Family Microhedylylidae**
Microhedylyle
glandulifera (Kowalewsky, 1901)
glomerans Salvini-Plawen, 1973
lactea Hertling, 1930
neapolitana (Rankin, 1979)
odhneri (Marcus Ev. & Er., 1955)
- Pontohedyle*
milatschevitchi (Kowalewsky, 1901)
- Parahedylyle*
cryptophtalma (Westheide & Wawra, 1974)
tyrtowii (Kowalewsky, 1901)
- Family Platyhedylylidae**
Platyhedylyle
denudata Salvini-Plawen, 1973
- Family Asperspinidae**
Asperspina
brambelli (Swedmark, 1968)
rhopalotecta (Salvini-Plawen, 1973)
- Antemnella*
loricata (Swedmark, 1968)
- Family Hedylopsidae**
Hedylopsis
brambelli Swedmark, 1968
spiculifera (Kowalewsky, 1901)
suecica Odhner, 1937
- Family Minicheviellidae**
Minicheviella
murmanica (Kuchinskaja & Minichev, 1978)
- Family Rhopopidae**
Rhodope
crucispiculata Salvini-Plawen, 1988
veranyi Koelliker, 1847
- Family Akeridae**
Akera
bullata Müller O.F., 1776
- Family Aplysiidae**
Aplysia
dactylomela Rang, 1828
depilans Gmelin, 1791
fasciata Poiret, 1789
juliana Quoy & Gaimard, 1832
parvula Guilding in Mørch, 1863
punctata (Cuvier, 1803)
- Bursatella*
leachii de Blainville, 1817
- Family Dolabriferidae**
Dolabrifera
holboelli Bergh, 1872
- Notarchus*
indicus Schweigger, 1820
punctatus Philippi, 1836
- Petalifera*
gravieri (Vayssiére, 1906)
petalifera (Rang, 1828)
- Phyllaplysia*
depressa (Cantraine, 1835)
lafonti (Fischer P., 1870)
- Family Bathydorididae**
Bathydoris
ingolfiana Bergh, 1899
- Family Aegiretidae**
Aegires
leuckarti Vérany, 1853
palensis Ortea, Luque & Templado, 1990
punctilucens (d'Orbigny, 1837)
sublaevis Odhner, 1932
- Triopella*
incisa (Sars G.O., 1872)
- Family Corambidae**
Corambe
obscura (Verrill, 1870)

- testudinaria* Fischer P., 1889
Echinocorambe
brattegardi (Valdés & Bouchet, 1998)
- Family Goniodorididae**
Goniodoris
barroisi Vayssiére, 1901
castanea Alder & Hancock, 1845
nodosa (Montagu, 1808)
Lophorodoris
danielsseni (Friele & Hansen, 1876)
Okenia
aspersa (Alder & Hancock, 1845)
elegans (Leuckart, 1828)
hispanica (Valdés & Ortea, 1995)
cupella (Vogel & Schultz, 1970)
leachii (Alder & Hancock, 1854)
mediterranea (Von Ihering, 1886)
Ancula
gibbosa (Risso, 1818)
Trapania
fusca (Lafont, 1874)
hispalensis Cervera & Garcia-Gomez, 1989
lineata Haefelfinger, 1960
maculata Haefelfinger, 1960
orteai Garcia-Gomez & Cervera in Cervera & Garcia-Gomez, 1989
pallida Kress, 1968
tartanella (Von Ihering, 1886)
Bermudella
polycerelloides Ortea & Bouchet, 1983
- Family Onchidorididae**
Onchidoris
albonigra (Pruvot-Fol, 1951)
aspera (Alder & Hancock, 1842)
bouvieri (Vayssiére, 1919)
cervinoi Ortea & Urgorri, 1979
depressa (Alder & Hancock, 1842)
fusca (Müller O.F., 1776)
inconspicua (Alder & Hancock, 1851)
muricata (Müller O.F., 1776)
neapolitana (delle Chiaje, 1841)
oblonga (Alder & Hancock, 1845)
pusilla (Alder & Hancock, 1845)
reticulata Ortea, 1979
sparsa (Alder & Hancock, 1846)
tridactila Ortea & Ballesteros, 1982
- Acanthodoris*
pilosa (Abildgaard in Müller, 1789)
Adalaria
loveni Alder & Hancock, 1862
proxima (Alder & Hancock, 1854)
Diaphorodoris
luteocincta (Sars M., 1870)
papillata Portmann & Sandmeier, 1960
- Doridunculus*
echinulatus Sars G.O., 1878
- Family Triophidae**
Crimora
papillata Alder & Hancock, 1862
Kaloplocamus
aureus (Odhner, 1932)
filosus Cattaneo-Vietti & Sordi, 1988
ramosus (Cantraine, 1835)
Plocamopherus
maderae (Lowe, 1842)
ocellatus (Rüppell & Leuckart, 1831)
- Family Dorididae**
Doris
alboranica Bouchet, 1977
bertheloti (d'Orbigny, 1839)
bicolor (Bergh, 1884)
marmorata Risso, 1818
nobilis Odhner, 1907
ocelligera (Bergh, 1881)
- pseudoverrucosa* (Von Ihering, 1886)
sticta (Iredale & O'Donoghue, 1923)
verrucosa Linné, 1758
Thorybopus
lophatus Bouchet, 1977
- Family Aldisidae**
Aldisa
banyulensis Pruvot-Fol, 1951
barlettae Ortea & Ballesteros, 1989
berghi Vayssiére, 1901
binotata Pruvot-Fol, 1953
zealandica (Alder & Hancock, 1854)
- Family Archidorididae**
Archidoris
pseudoargus (Rapp, 1827)
Anisodoris
marmorata (Bergh, 1881)
Atagema
gibba Pruvot-Fol, 1951
rugosa Pruvot-Fol, 1951
Sclerodoris
minor Eliot, 1903
- Family Chromodorididae**
Chromodoris
britoi Ortea & Perez, 1983
elegantula (Philippi, 1844)
goslineri Ortea & Valdés in Ortea, Valdés & Garcia-Gómez, 1996
krohni (Vérany, 1846)
luteopunctata (Gantès, 1962)
luteorosea (Rapp, 1827)
purpurea (Risso in Guérin, 1831)
quadricolor (Rüppell & Leuckart, 1831)
rodomaculata Ortea & Valdés, 1991
Cadlina
excavata (Pruvot-Fol, 1951)
glabra (Friele & Hansen, 1876)
laevis (Linné, 1767)
pellucida (Risso, 1826)
Glossodoris
edmundsi Cervera, Garcia-Gomez & Ortea, 1989
limbata Vicente, 1967
Hypsodoris
bilineata (Pruvot-Fol, 1953)
cantabrica Bouchet & Ortea, 1980
orsini (Vérany, 1846)
fontanraui (Pruvot-Fol, 1951)
gasconi Ortea, Valdés & Garcia-Gómez, 1996
infuscata (Rüppell & Leuckart, 1831)
malacitana Luque, 1986
tricolor (Cantraine, 1835)
villafranca (Risso, 1818)
picta (Schultz in Philippi, 1836)
- Family Discodorididae**
Discodoris
atromaculata (Bergh, 1880)
concinna (Alder & Hancock, 1864)
confusa Ballesteros, Llera & Ortea, 1985
edwardsi Vayssiére, 1902
erubescens Bergh, 1884
fragilis (Alder & Hancock, 1864)
maculosa Bergh, 1884
millegrana (Alder & Hancock, 1854)
patriziae Perrone, 1991
porri (Vérany, 1846)
punctifera (Abraham, 1877)
rosi Ortea, 1979
rubens Vayssiére, 1919
sordii (Perrone, 1990)
stellifera (Ihering in Vayssiére, 1904)
Geitodoris
bacalladoi Ortea, 1990
bonosi Ortea & Ballesteros, 1981
joubini (Vayssiére, 1919)

- perfossa* Ortea, 1990
planata (Alder & Hancock, 1846)
portmanni (Schmekel, 1972)
pusae (Marcus Er., 1955)
rubens (Vayssi  re, 1919)
- Paradoris**
ceneris Ortea, 1995
granulata Bergh, 1884
indecora (Bergh, 1881)
inversa Ortea, 1995
mollis Ortea, 1995
- Thordisa**
aurea Pruvot-Fol, 1951
azmani Cervera & Garc  a-Gomez, 1989
filix Pruvot-Fol, 1951
pallida Bergh, 1884
- Family Kentrodorididae**
Jorunna
atypa Bergh, 1881
onubensis Cervera, Garc  a-Gomez & Garc  a, 1986
tomentosa (Cuvier, 1804)
- Family Platydorididae**
Platydoris
argo (Linn  , 1767)
dura Pruvot-Fol, 1951
maculata Bouchet, 1977
philippi Bergh, 1877
stomascuta Bouchet, 1977
- Baptodoris*
cinnabarinus Bergh, 1884
perezi Llera & Ortea in Ortea, Perez & Llera, 1982
- Carminodoris*
boucheti Ortea, 1979
spinobranchialis Ortea & Martinez, 1992
- Taringa*
armata Swennen, 1961
ascitica Ortea, Perez & Llera, 1982
bacalladoi Ortea, Perez & Llera, 1982
faba Ballesteros, Llera & Ortea, 1985
fanabensis Ortea & Mart  nez, 1992
oleica Ortea, Perez & Llera, 1982
pinoi Perrone, 1985
tarifaensis Garc  a-Gomez, Cervera & Garc  a-Martin, 1993
tritorquis Ortea, Perez & Llera, 1982
- Family Rostangidae**
Rostanga
anthelia Perrone, 1991
rubra (Risso, 1818)
- Family Polyceridae**
Polycreta
aurantiomarginata Garc  a-Gomez & Bobo, 1984
dubia Sars M., 1829
elegans (Bergh, 1894)
faeroensis Lemche, 1929
hedgpethi (Marcus Er., 1964)
maculata Pruvot-Fol, 1951
nothus (Johnston, 1838)
quadrilineata (M  ller O.F., 1776)
- Colga*
pacifica (Bergh, 1894)
- Limacia*
clavigera (M  ller O.F., 1776)
- Polycrella*
emertoni Verrill, 1881
- Thecacera*
pennigera (Montagu, 1815)
virescens Forbes & Hanley, 1851
- Family Gymnodorididae**
Robostra
europea Garc  a-Gomez, 1985
- Tambja*
ceutae Garc  a-Gomez & Ortea, 1988
marbellensis (Schick & Cervera, 1998)
- Family Dendrodorididae**
Dendrodoris
grandiflora (Rapp, 1827)
herytra Vald  s & Ortea in Vald  s, Ortea, Avila & Ballesteros, 1996
limbata (Cuvier, 1804)
nigra (Stimpson, 1855)
- Doriopsilla*
areolata Bergh, 1880
pelseneeri d'Oliveira, 1895
- Family Phyllidiidae**
Phyllidia
flava Aradas, 1847
- Phyllidiopsis*
berghi Vayssi  re, 1902
- Fryeria*
bayi Bouchet, 1983
- Family Dendronotidae**
Dendronotus
comteti (Vald  s & Bouchet, 1998)
frondosus (Ascanius, 1774)
lacteus (Thompson W., 1840)
robustus Verrill, 1880
- Family Dotidae**
Doto
acuta Schmekel & Kress, 1977
affinis (d'Orbigny, 1835)
arteoi Ortea, 1978
cerviceniga Ortea & Bouchet, 1989
cindyneutes Bouchet, 1977
confluens Hesse, 1872
coronata (Gmelin, 1791)
crassicornis Sars M., 1870
cuspidata Alder & Hancock, 1862
dunnei Lemche, 1976
eireana Lemche, 1976
escallari (Ortea, Moro & Espinosa, 1998)
floridicola Simroth, 1888
fluctifraga Ortea & Perez, 1982
fragaria Ortea & Bouchet, 1989
fragilis (Forbes, 1838)
furva Garc  a J.C. & Ortea, 1984
hydrallmanniae (Morrow, Thorpe & Picton, 1992)
hystrix Picton & Brown, 1981
koenneckeri Lemche, 1976
lemchei Ortea & Urgorri, 1978
leopardina Vicente, 1967
millbayana Lemche, 1976
oblicua Ortea & Urgorri, 1978
paulinae Trinchese, 1881
pinnatifida (Montagu, 1804)
pontica Swennen, 1961
pygmaea (Bergh, 1871)
rosea Trinchese, 1881
sotilloi (Ortea, Moro & Espinosa, 1998)
tuberculata Lemche, 1976
unguis Ortea & Rodriguez, 1989
verdicioi Ortea & Urgorri, 1978
- Family Hancockiidae**
Hancockia
uncinata (Hesse, 1872)
- Family Lomanotidae**
Lomanotus
barlettai Garc  a-Gomez, Lopez-Gonzalez & Garc  a, 1990
genei V  rany, 1846
marmoratus (Alder & Hancock, 1845)
- Family Phylliroididae**
Phylliroe
atlantica Bergh, 1871
bucephala P  eron & Lesueur, 1810
- Cephalopyge*
trematoides (Chun, 1889)

- Family Scyllaeidae**
- Scyllaea*
 - pelagica* Linné, 1758
- Family Tethydidae**
- Tethys*
 - fimbria* Linné, 1767
 - Melibe*
 - fimbriata* Alder & Hancock, 1864
- Family Tritoniidae**
- Tritonia*
 - episcopalis* Bouchet, 1977
 - griegi* Odhner, 1922
 - hombergi* Cuvier, 1803
 - ingolfiana* (Bergh, 1899)
 - lineata* Alder & Hancock, 1848
 - manicata* Deshayes, 1853
 - nilsodhneri* Marcus Ev., 1983
 - plebeia* Johnston, 1828
 - striata* Haefelfinger, 1963
 - Marionia*
 - blainvillaea* (Risso, 1818)
 - Tritoniopsis*
 - cincta* (Pruvot-Fol, 1937)
- Family Arminidae**
- Armina*
 - loveni* (Bergh, 1860)
 - maculata* Rafinesque, 1814
 - neapolitana* (delle Chiaje, 1824)
 - tigrina* Rafinesque, 1814
 - tricuspidata* Thompson, Cattaneo & Wong, 1990
- Lingella**
- elfortiana* de Blainville, 1823
 - quadrilateralis* (Bergh, 1860)
- Family Heterodorididae**
- Heterodoris*
 - ingolfiana* (Bergh, 1899)
 - robusta* Verrill & Emerton, 1882
- Family Heroidae**
- Hero*
 - blanchardi* Vayssièvre, 1888
 - formosa* (Lovén, 1844)
- Family Gonaeolidiidae**
- Gonaeolis*
 - typica* Sars M., 1861
- Family Madrellidae**
- Madrella*
 - aurantiaca* Vayssièvre, 1902
 - Eliotia*
 - souleyeti* Vayssièvre, 1909
- Family Proctonotidae**
- Janolus*
 - cristatus* (delle Chiaje, 1841)
 - faustoi* Ortea & Llera, 1988
 - hyalinus* (Alder & Hancock, 1854)
 - Proctonotus*
 - mucroniferus* (Alder & Hancock, 1844)
- Family Aeolidiidae**
- Aeolidia*
 - papillosa* (Linné, 1761)
 - Aeolidiella*
 - alderi* (Cocks, 1852)
 - glauca* (Alder & Hancock, 1845)
 - indica* Bergh, 1888
 - rubra* (Cantraine, 1835)
 - sanguinea* (Norman, 1877)
 - Baeolidia*
 - cryoporus* Bouchet, 1977
 - nodosa* (Haefelfinger & Stamm, 1958)
 - Berghia*
 - columbina* (Garcia-Gomez & Thompson, 1990)
 - norvegica* Odhner, 1939
 - Spurilla*
- caeruleescens* Laurillard, 1830
- neapolitana* (delle Chiaje, 1841)
- vayssierei* Garcia J.C. & Cervera, 1985
- verrucicornis* (Costa A., 1867)
- Cerberilla*
- bernadettae* Tardy, 1965
- Phidiana*
- indica* (Bergh, 1896)
- Family Calmidae**
- Calma*
 - glaucooides* (Alder & Hancock, 1854)
- Family Cumanotidae**
- Cumanotus*
 - beaumonti* (Eliot, 1906)
 - cuenoti* (Pruvot-Fol, 1948)
- Family Eubranchidae**
- Eubranchus*
 - arci* Ortea, 1981
 - cingulatus* (Alder & Hancock, 1847)
 - doriae* (Trinchesi, 1874)
 - exiguus* (Alder & Hancock, 1848)
 - farrani* (Alder & Hancock, 1844)
 - limensis* Garcia-Gomez, Cervera & Garcia, 1990
 - pallidus* (Alder & Hancock, 1842)
 - prietoi* Llera & Ortea, 1981
 - rupium* (Møller, 1842)
 - tricolor* Forbes, 1838
 - vittatus* (Alder & Hancock, 1842)
- Family Facelinidae**
- Facelina*
 - annulicornis* (Chamisso & Eysenhardt, 1821)
 - bostoniensis* (Couthouy, 1838)
 - coronata* (Forbes & Goodsir, 1839)
 - dubia* (Pruvot-Fol, 1948)
 - fusca* Schmekel, 1966
 - lugubris* (Bergh, 1882)
 - quatrefoigesi* Vayssièvre, 1888
 - rubrovittata* (Costa A., 1866)
 - rutila* Pruvot-Fol, 1951
 - variegata* d'Oliveira, 1895
 - vicina* (Bergh, 1882)
 - Algarvia*
 - alba* Garcia-Gomez & Cervera, 1989
- Antonietta*
- luteorufa* Schmekel, 1966
- Caloria*
- elegans* (Alder & Hancock, 1845)
- Facelinopsis*
- marioni* (Vayssièvre, 1888)
- Pruvofolia*
- pselliotes* (Labbé, 1923)
- Rolandia*
- hispanica* Pruvot-Fol, 1951
- Family Favorinidae**
- Favorinus*
 - blianus* Lemche & Thompson, 1974
 - branchialis* (Rathke, 1806)
 - vitreus* Ortea, 1982
- Dicata*
- odhneri* Schmekel, 1967
- Dondice*
- banyulensis* Portmann & Sandmeier, 1960
- Family Fionidae**
- Fiona*
 - pinnata* (Eschscholtz, 1831)
- Family Flabellinidae**
- Flabellina*
 - affinis* (Gmelin, 1791)
 - babai* Schmekel, 1972
 - baetica* Garcia-Gomez, 1984
 - insolita* Garcia-Gomez & Cervera, 1989
 - ischitana* Hirano & Thompson, 1990
 - pedata* (Montagu, 1815)

- pellucida* (Alder & Hancock, 1843)
rubrolineata (O'Donoghue, 1929)
- Calmella**
cavolini (Vérany, 1846)
- Coryphella**
borealis Odhner, 1922
browni Picton, 1980
gracilis (Alder & Hancock, 1844)
lineata (Lovén, 1846)
nobilis Verrill, 1880
parva Hadfield, 1963
salmonacea (Couthouy, 1838)
sarsi Friele, 1902
verrucosa (Sars M., 1829)
- Rioselleolis**
anadoni Ortea, 1979
- Paracoryphella**
islandica (Odhner, 1937)
- Family Glaucidae**
Glaucus
atlanticus Forster, 1777
- Family Piseinotecidae**
Piseinotecus
gabinieri (Vicente, 1975)
gaditanus Cervera, Garcia-Gomez & Garcia, 1987
sphaeriferus (Schmekel, 1965)
- Family Pseudovermidae**
Pseudovermis
artabrensis Urgorri, Cobo & Besteiro, 1991
axi Marcus Ev. & Er., 1955
boadeni Salvini-Plawen & Sterrer, 1968
kowalewskyi Salvini-Plawen & Sterrer, 1968
papillifer Kowalewsky, 1901
paradoxus Periaslavzev, 1891
schultzi Marcus Ev. & Er., 1955
setensis Fize, 1961
thompsoni Salvini-Plawen, 1991
- Family Tergipedidae**
Tergipes
edwardsii Nordmann, 1844
tergipes (Forskål, 1775)
- Catriona*
gymnota (Couthouy, 1838)
maua Marcus Ev. & Er., 1960
- Cuthona*
abyssicola (Bergh, 1884)
albopunctata (Schmekel, 1968)
amoena (Alder & Hancock, 1845)
berghi (Friele, 1902)
caerulea (Montagu, 1804)
concinna (Alder & Hancock, 1843)
distans Odhner, 1922
ferruginea (Friele, 1902)
foliata (Forbes & Goodsir, 1839)
genovae (O'Donoghue, 1929)
granosaa (Schmekel, 1966)
ilonae (Schmekel, 1968)
leopardina (Vayssiére, 1888)
miniostriata (Schmekel, 1968)
nana (Alder & Hancock, 1842)
norvegica (Odhner, 1929)
ocellata (Schmekel, 1966)
peregrina (Gmelin, 1791)
pustulata (Alder & Hancock, 1854)
rubescens Picton & Brown, 1978
suecica (Odhner, 1940)
thompsoni Garcia, Lopez-Gonzalez & Garcia-Gomez, 1991
viridis (Forbes, 1840)
willani Cervera, Garcia-Gomez & Lopez-Gonzalez, 1992
- Embletonia*
pulchra (Alder & Hancock, 1844)
- Tenellia*
adspersa (Nordmann, 1845)
- Family Siphonariidae**
Siphonaria
kurracheensis (Reeve, 1856)
pectinata (Linné, 1758)
- Williamia*
gussonii (Costa O.G., 1829)
- Family Trimusculidae**
Trimusculus
afer (Gmelin, 1791)
mammilaris (Linné, 1758)
- Family Onchidiidae**
Onchidella
celtica (Cuvier, 1817)
- Family Ellobiidae**
Pseudomelampus
exiguus (Lowe, 1832)
- Auriculinella*
bidentata (Montagu, 1808)
- Ovatella*
aequalis (Lowe, 1832)
firminii (Payraudeau, 1826)
vulcani (Morelet, 1860)
- Myosotella*
denticulata (Montagu, 1803)
myosotis (Draparnaud, 1801)
- Pedipes*
afer (Gmelin, 1791)
- Family Otinidae**
Otina
ovata (Brown, 1827)
- Class Bivalvia**
- Family Xenoturbellidae**
Xenoturbella
bocki (Westblad, 1949)
westbladi (Israelsson, 1999)
- Family Solemyidae**
Solemya
togata (Poli, 1795)
- Family Nuculidae**
Nucula
atacellana Schenck, 1939
bicornis Gofas & Salas, 1996
decipiens Philippi, 1844
hanleyi Winckworth, 1931
nitidosa Winckworth, 1930
nucleus (Linné, 1758)
recondita Gofas & Salas, 1996
sulcata Bronn, 1831
tumidula Malm, 1861
- Ennucula*
aegaeensis (Forbes, 1844)
bushae (Dollfus, 1898)
corticata (Möller, 1842)
tenuis (Montagu, 1808)
- Brevinucula*
verrilli (Dall, 1886)
- Family Pristiglomidae**
Pristigloma
nitens (Jeffreys, 1876)
- Family Nuculanidae**
Nuculana
commutata (Philippi, 1844)
illirica Carrozza, 1987
minuta (Müller O.F., 1776)
pella (Linné, 1767)
pernula Müller O.F., 1779
- Family Malletidae**
Katadesmia
cuneata (Jeffreys, 1876)

- kolthoffi* (Hägg, 1904)
Pseudomalletia
obtusa (Sars G.O., 1872)
- Family Neilonellidae**
Neilonella
striolata (Brugnone, 1876)
- Family Spinulidae**
Bathyspinula
excisa (Philippi, 1844)
- Family Tindariidae**
Tindaria
derjugini Gorbunov, 1946
sericea (Jeffreys, 1876)
- Family Yoldiidae**
Yoldia
amygdalea Valenciennes, 1846
hyperborea Torell, 1859
micrometrika Seguenza G., 1877
minima Seguenza G., 1877
Yoldiella
annenkovae (Gorbunov, 1946)
dissimilis Verrill & Bush, 1898
expansa (Jeffreys, 1876)
folini Warén, 1978
frami Knudsen, 1985
frigida (Torell, 1859)
insculpta (Jeffreys, 1879)
intermedia (Sars M., 1865)
jeffreysi (Hidalgo, 1877)
lenticula (Møller, 1842)
lucida (Lovén, 1846)
messanensis (Jeffreys, 1870)
nana (Sars M., 1865)
philippiana (Nyst, 1845)
propinqua (Leche, 1878)
pustulosa (Jeffreys, 1876)
seguenzae Bonfitti & Sabelli, 1995
semistriata (Jeffreys, 1879)
solidula Warén, 1989
subaequilatera (Jeffreys, 1879)
tamara (Gorbunov, 1946)
- Megayoldia*
thraciaeformis (Storer, 1838)
- Portlandia*
arctica (Gray J.E., 1824)
- Microgloma*
guilonardi (Hoeksema, 1993)
pusilla (Jeffreys, 1879)
tumidula (Monterosato, 1880)
- Family Arcidae**
- Arcia*
noae Linné, 1758
tetragona Poli, 1795
- Asperarca*
magdalena (La Perna, 1998)
nodulosa (Müller O.F., 1776)
secreta (La Perna, 1998)
- Barbatia*
barbata (Linné, 1758)
clathrata (Defrance, 1816)
plicata (Dillwyn, 1817)
- Anadara*
corbuloides (Monterosato, 1878)
demiri (Piani, 1981)
inaequivalvis (Bruguière, 1789)
natalensis (Krauss, 1848)
polii (Mayer, 1868)
- Bathyarca*
frielei (Friele, 1877)
glacialis (Gray J.E., 1824)
inaequisculpta (Smith E.A., 1885)
pectunculoides (Scacchi, 1834)
philippiana (Nyst, 1848)
- Bentharca*
- asperula* (Dall, 1881)
- Family Noetiidae**
Striarca
lactea (Linné, 1758)
- Family Limopsidae**
Limopsis
angusta Jeffreys, 1879
aurita (Brocchi, 1814)
cristata Jeffreys, 1876
minuta (Philippi, 1836)
multistriata (Forskål, 1775)
friedbergi Glibert & van de Poel, 1965
- Family Glycymerididae**
Glycymeris
bimaculata (Poli, 1795)
glycymeris (Linné, 1758)
violacea (Lamarck, 1819)
- Family Mytilidae**
Mytilus
edulis Linné, 1758
galloprovincialis Lamarck, 1819
trossulus Gould, 1850
- Brachidontes*
pharaonis (Fischer P., 1870)
- Mytilaster*
lineatus (Gmelin, 1791)
marioni (Locard, 1889)
minimus (Poli, 1795)
solidus Monterosato, 1872
- Perna*
perna (Linné, 1758)
- Crenella*
arenaria Monterosato, 1875
canariensis (Odhner, 1932)
decussata (Montagu, 1808)
pellucida (Jeffreys, 1859)
- Gregariella*
petagna (Scacchi, 1832)
semigranata (Reeve, 1858)
- Modiolarca*
subpicta (Cantraine, 1835)
- Musculus*
costulatus (Risso, 1826)
discors (Linné, 1767)
laevigatus (Gray J.E., 1824)
niger (Gray J.E., 1824)
- Rhomboidella*
prideauxi (Leach, 1815)
- Lithophaga*
lithophaga (Linné, 1758)
- Myoforceps*
aristatus (Dillwyn, 1817)
- Modiolus*
adriaticus (Lamarck, 1819)
auriculatus Krauss, 1848
barbatus (Linné, 1758)
lulat (Dautzenberg, 1891)
martorelli (Hidalgo, 1878)
modiolus (Linné, 1758)
stultorum (Jousseaume, 1893)
- Amygdalum*
agglutinans (Cantraine, 1835)
politum (Verrill & Smith, 1880)
- Dacrydium*
abyssorum (Allen, 1999)
balgimi Salas & Gofas, 1997
dauvini Salas & Gofas, 1997
filiferum Salas & Gofas, 1997
hyalinum Monterosato, 1875
ockelmanni Mattson & Warén, 1977
sandersi (Allen, 1999)
vitreum (Møller, 1842)
viviparum Ockelmann, 1983
wareni Salas & Gofas, 1997

- Idas***
argenteus Jeffreys, 1876
ghisotti Warén & Carozza, 1990
modiolaeformis (Sturany, 1896)
pelagicus (Forbes in Woodward, 1854)
simpsoni (Marshall, 1900)
- Bathymodiolus***
azoricus (Cosel & Comtet in Cosel, Comtet & Krylova, 1999)
- Modiolula***
phaseolina (Philippi, 1844)
- Musculista***
perfragilis (Dunker, 1857)
senhousia (Benson in Cantor, 1842)
- Xenostrobus***
securis (Lamarck, 1819)
- Family Pinnidae**
- Pinna***
nobilis Linné, 1758
rudis Linné, 1758
- Atrina***
pectinata (Linné, 1767)
- Family Pteriidae**
- Pteria***
hirundo (Linné, 1758)
- Pinctada***
radiata (Leach, 1814)
- Family Malleidae**
- Malleus***
regulus (Forskål, 1775)
- Family Isognomonidae**
- Isognomon***
isognomon (Linné, 1758)
- Family Pectinidae**
- Pecten***
jacobeus (Linné, 1758)
maximus (Linné, 1758)
- Aequipecten***
commutatus (Monterosato, 1875)
opercularis (Linné, 1758)
- Lissopecten***
hyalinus (Poli, 1795)
- Nodipecten***
corallinoides (d'Orbigny, 1840)
- Palliolum***
furtivum (Lovén, 1846)
incomparabile (Risso, 1826)
striatum (Müller O.F., 1776)
tigerinum (Müller O.F., 1776)
- Arctinula***
greenlandica (Sowerby G.B. II, 1842)
- Propeamussium***
fenestratum (Forbes, 1844)
lucidum (Wyville-Thompson, 1873)
propinquum (Smith E.A., 1885)
- Pseudamussium***
clavatum (Poli, 1795)
peslutrae (Linné, 1771)
- Cyclopecten***
brundisiensis Smriglio & Mariottini, 1990
hoskynsi (Forbes, 1844)
imbrifer (Lovén, 1846)
- Electopecten***
vitreus (Gmelin, 1791)
- Hyalopecten***
frigidus Jensen, 1904
similis (Laskey, 1811)
- Crassadoma***
multistriata (Poli, 1795)
pusio (Linné, 1758)
- Chlamys***
alicei Dautzenberg & Fischer H., 1897
flexuosa (Poli, 1795)
glabra (Linné, 1758)
- islandica*** (Müller O.F., 1776)
pesfelis (Linné, 1758)
protea (Dillwyn, 1817)
sulcata (Müller O.F., 1776)
varia (Linné, 1758)
- Family Spondylidae**
- Spondylus***
gaederopus Linné, 1758
gussonii Costa O.G., 1829
senegalensis (Schreibers, 1793)
spinosus (Schreibers, 1793)
- Family Anomiidae**
- Anomia***
ephippium Linné, 1758
- Isonomia***
alberti (Dautzenberg & Fischer H., 1896)
- Heteranomia***
squamula (Linné, 1758)
- Pododesmus***
patelliformis (Linné, 1761)
squama (Gmelin, 1791)
- Family Limidae**
- Lima***
lima (Linné, 1758)
marioni Fischer P., 1888
- Limaria***
hians (Gmelin, 1791)
tuberculata (Wood S., 1839)
- Limea***
loscombii (Sowerby G.B. I, 1824)
- Acesta***
excavata (Fabricius J.C., 1779)
- Limatula***
gwyni (Sykes, 1903)
hyperborea Jensen, 1905
jeffreysi (Fischer P., 1882)
laminifera (Smith E.A., 1885)
similis (Jensen, 1912)
subauriculata (Montagu, 1808)
subovata (Jeffreys, 1876)
- Notolimea***
clandestina Salas, 1994
crassa (Forbes, 1844)
- Family Ostreidae**
- Ostrea***
edulis Linné, 1758
- Crassostrea***
gigas (Thunberg, 1793)
virginica (Gmelin, 1791)
- Ostreola***
stentina (Payraudeau, 1826)
- Saccostrea***
commercialis (Iredale & Roughley, 1933)
- Family Gryphaeidae**
- Neopycnodonte***
cochlear (Poli, 1795)
- Family Lucinidae**
- Linga***
adansoni (d'Orbigny, 1839)
- Ctena***
decussata (Costa O.G., 1829)
- Loripes***
lacteus (Linné, 1758)
- Megaxinus***
appendiculatus (Locard, 1898)
unguiculinus (Pallary, 1904)
- Divaricella***
angulifera Von Martens, 1880
- Lucinella***
divaricata (Linné, 1758)
- Anodontia***
fragilis (Philippi, 1836)
- Myrtea***

- amorpha* (Sturany, 1896)
spinifera (Montagu, 1803)
- Lucinoma**
borealis (Linné, 1767)
- Family Thyasiridae**
- Axinus*
grandis (Verrill, 1885)
- Axinopsida*
orbiculata (Sars G.O., 1878)
- Thyasira*
allenii Carrozza, 1981
biplicata (Philippi, 1836)
biscayensis Payne & Allen, 1991
brevis (Verrill & Bush, 1898)
croulinensis (Jeffreys, 1847)
equalis (Verrill & Bush, 1898)
eumyaria (Sars M., 1870)
ferruginea (Locard, 1886)
flexuosa (Montagu, 1803)
gouldii (Philippi, 1845)
granulosa (Monterosato, 1874)
incrassata (Jeffreys, 1876)
inflata Payne & Allen, 1991
oblonga (Monterosato, 1878)
obsoleta (Verrill & Bush, 1898)
planata (Jeffreys, 1882)
perplicata (Salas, 1996)
pygmaea (Verrill & Bush, 1898)
sarsi (Philippi, 1845)
striata (Sturany, 1896)
subcircularis Payne & Allen, 1991
subovata (Jeffreys, 1881)
succisa (Jeffreys, 1876)
tortuosa (Jeffreys, 1881)
transversa Payne & Allen, 1991
ultima Payne & Allen, 1991
- Family Ungulinidae**
- Diplodonita*
apicalis Philippi, 1836
brocchii (Deshayes, 1852)
eddystonia (Marshall, 1895)
rotundata (Montagu, 1803)
subrotunda (Issel, 1869)
torelli Jeffreys, 1876
- Ungulina*
cuneata (Spengler, 1782)
- Family Chamidae**
- Chama*
pacifica (Broderip, 1834)
gryphoides Linné, 1758
- Pseudochama*
cornucopia (Reeve, 1846)
gryphina (Lamarck, 1819)
radians (Lamarck, 1819)
- Family Galeommatidae**
- Galeomma*
turtoni Sowerby G.B. I in Turton, 1825
- Vasconiella*
jeffreysiana (Fischer P. in de Folin & Périer, 1873)
- Family Kelliidae**
- Kellia*
suborbicularis (Montagu, 1803)
- Pseudopythina*
macandrewi (Fischer P., 1867)
- Bornia*
geoffroyi (Payraudeau, 1826)
sebetia (Costa O.G., 1829)
- Solecardia*
rotunda (Jeffreys, 1881)
- Family Lasaeidae**
- Lasaea*
rubra (Montagu, 1803)
- Scacchia*
- oblonga* (Philippi, 1836)
zorni van Aartsen & Fehr-de Wal, 1985
- Family Leptonidae**
- Lepton*
lacerum Jeffreys in de Folin & Périer, 1873
squamosum (Montagu, 1803)
subtrigonom Jeffreys in de Folin & Périer, 1873
- Hemilepton*
nitidum (Turton, 1822)
- Arculus*
sykesi (Chaster, 1895)
- Litigiella*
corganii van Aartsen, 1996
glabra (Fischer P. in de Folin & Périer, 1873)
- Family Montacutidae**
- Montacuta*
convexa Monterosato, 1875
donacina Wood S., 1848
ferruginosa (Montagu, 1808)
goudii (van Aartsen, 1996)
maltzani Verkrüzen, 1876
phascolionis Dautzenberg & Fischer H., 1925
pura (Smith E.A., 1885)
semirubra (Gaglini, 1992)
substriata (Montagu, 1808)
tenella Lovén, 1846
voeringi Friese, 1879
- Axinodon*
symmetros (Jeffreys, 1876)
- Kelliopsis*
jozinae (van Aartsen & Carrozza, 1997)
- Mysella*
bidentata (Montagu, 1803)
dawsoni (Jeffreys, 1864)
obliquata (Chaster, 1897)
ovata (Jeffreys, 1881)
pellucida (Jeffreys, 1881)
triangularis (Watson, 1897)
tumidula (Jeffreys, 1866)
- Devonia*
perrieri (Malard, 1904)
- Epilepton*
clarkiae (Clark W., 1852)
parrussetensis (Giribet & Peñas, 1999)
- Family Neoleptonidae**
- Neolepton*
cancellatum (Salas & Gofas, 1998)
guanche (Salas & Gofas, 1998)
sulcatulum (Jeffreys, 1859)
- Family Sportellidae**
- Sportella*
recondita (Fischer P. in de Folin, 1872)
- Family Turtoniidae**
- Turtonia*
minuta (Fabricius O., 1780)
- Family Carditidae**
- Cardita*
calyculata (Linné, 1758)
- Glans*
aculeata (Poli, 1795)
trapezia (Linné, 1767)
- Pteromeris*
jozinae van Aartsen, 1984
minuta (Scacchi, 1836)
- Venericardia*
antiquata (Linné, 1758)
- Family Astartidae**
- Astarte*
acuticostata Friese, 1877
arctica (Gray, 1824)
borealis (Schumacher, 1817)
crebricostata McAndrew & Forbes, 1847
crenata (Gray, 1824)

- elliptica* (Brown, 1827)
fusca (Poli, 1795)
montagui (Dillwyn, 1817)
sulcata (da Costa, 1778)
- Digitaria**
digitaria (Linné, 1758)
- Gonlia**
biplicata (Dall, 1903)
- Goodallia**
gofasi (Giribet & Peñas, 1999)
macandrewi (Smith E.A., 1881)
micalii (Giribet & Peñas, 1999)
pusilla (Forbes, 1844)
triangularis (Montagu, 1803)
- Family Crassatellidae**
Crassatina
planata (Calcaro, 1840)
- Family Cardiidae**
Cardium
indicum Lamarck, 1818
Acanthocardia
aculeata (Linné, 1758)
deshayesii (Payraudeau, 1826)
echinata (Linné, 1758)
paucicostata (Sowerby G.B. II, 1841)
spinosa (Solander, 1786)
tuberculata (Linné, 1758)
- Parvicardium*
elegantulum (Møller, 1842)
exiguum (Gmelin, 1791)
hauniense (Høpner Petersen & Russell, 1971)
minimum (Philippi, 1836)
ovale (Sowerby G.B. II, 1840)
scabrum (Philippi, 1844)
trapezium (Cecalupo & Quadri, 1996)
- Afrocardium*
richardi (Audouin, 1826)
- Plagiocardium*
papillosum (Poli, 1795)
- Laevicardium*
angmagsalikense (Voskuil & Onverwagt, 1992)
crassum (Gmelin, 1791)
flavum (Linné, 1758)
oblongum (Gmelin, 1791)
- Cerastoderma*
edule (Linné, 1758)
glaucum (Poiret, 1789)
- Ciliocardium*
ciliatum (Fabricius O., 1780)
- Serripes*
groenlandicus (Bruguière, 1789)
- Fulvia*
australis (Sowerby G.B. II, 1834)
fragilis (Forskål, 1775)
- Family Mactridae**
Mactra
glauca Von Born, 1778
olorina Philippi, 1846
stultorum (Linné, 1758)
- Spisula*
elliptica (Brown, 1827)
solida (Linné, 1758)
subtruncata (da Costa, 1778)
- Lutraria*
angustior Philippi, 1844
lutraria (Linné, 1758)
magna (da Costa, 1778)
- Eastonia*
rugosa (Helbling, 1779)
- Family Mesodesmatidae**
Atactodea
striata (Gmelin, 1791)
- Donacilla*
cornea (Poli, 1795)
- Monterosatus*
primus (Locard, 1899)
- Family Solenidae**
Solen
marginatus Pulteney, 1799
- Family Pharidae**
Pharus
legumen (Linné, 1758)
- Ensis*
arcuatus (Jeffreys, 1865)
directus (Conrad, 1843)
ensis (Linné, 1758)
minor (Chenu, 1843)
silqua (Linné, 1758)
- Phaxas*
pellucidus (Pennant, 1777)
- Sinupharus*
combieri (Fischer-Piette & Nicklès, 1946)
- Family Tellinidae**
Tellina
compressa Brocchi, 1814
distorta Poli, 1791
donacina Linné, 1758
fabula Gmelin, 1791
incarnata Linné, 1758
nitida Poli, 1791
planata Linné, 1758
pulchella Lamarck, 1818
punicea (Von Born, 1778)
pygmaea Lovén, 1846
serrata Brocchi, 1814
tenuis da Costa, 1778
valtonis Hanley, 1844
- Arcopagia*
balaustina (Linné, 1758)
crassa (Pennant, 1777)
- Macoma*
balthica (Linné, 1758)
calcarea (Gmelin, 1791)
cumana (Costa O.G., 1829)
melo (Sowerby G.B. III, 1866)
- Gastrana*
fragilis (Linné, 1758)
- Strigilla*
pseudocarnaria (Boss, 1969)
- Family Donacidae**
Donax
semistriatus Poli, 1795
trunculus Linné, 1758
venustus Poli, 1795
vittatus (da Costa, 1778)
- Capsella*
variegata (Gmelin, 1791)
- Family Psammobiidae**
Gari
costulata (Turton, 1822)
depressa (Pennant, 1777)
fervens (Gmelin, 1791)
intermedia (Deshayes, 1855)
pseudoweinkauffi Cosel, 1989
tellinella (Lamarck, 1818)
- Soletellina*
rubra (Schroeter, 1788)
subradiata (Deshayes in Reeve, 1857)
- Family Scrobiculariidae**
Scrobicularia
cottardi (Payraudeau, 1826)
plana (da Costa, 1778)
- Family Semelidae**
Semele
striata (Reeve, 1853)
- Ervilia*
castanea (Montagu, 1803)

- Abra**
alba (Wood W., 1802)
longicallus (Scacchi, 1834)
nitida (Müller O.F., 1776)
prismatica (Montagu, 1808)
profundorum (Smith E.A., 1885)
segmentum (Récluz, 1843)
tenuis (Montagu, 1803)
- Family Solecurtidae**
Solecurtus
multistriatus (Scacchi, 1835)
scopula (Turton, 1822)
strigilatus (Linné, 1758)
- Azorinus*
chamasolen (da Costa, 1778)
- Family Arcticidae**
Arctica
islandica (Linné, 1767)
- Family Kelliellidae**
Kelliella
abyssicola (Forbes, 1844)
- Family Trapezidae**
Trapezium
oblongum (Linné, 1758)
Coralliphaga
lithophagella (Lamarck, 1819)
- Isorropodon*
perplexum (Sturany, 1896)
- Family Vesicomyidae**
Vesicomya
atlantica (Smith E.A., 1885)
- Family Glossidae**
Glossus
humanus (Linné, 1758)
- Family Veneridae**
Venus
casina Linné, 1758
nux Gmelin, 1791
verrucosa Linné, 1758
- Mercenaria*
mercenaria (Linné, 1758)
- Globivenus*
effossa (Philippi, 1836)
- Chamelea*
gallina (Linné, 1758)
striatula (da Costa, 1778)
- Liocyma*
fluctuosa (Gould, 1841)
- Clausinella*
fasciata (da Costa, 1778)
- Timoclea*
ovata (Pennant, 1777)
roemeriana (Issel, 1869)
- Gastrarium*
calipygum (Von Born, 1778)
pectinatum (Linné, 1758)
- Gouldia*
minima (Montagu, 1803)
- Clementia*
papyracea (Gmelin, 1791)
- Dosinia*
exoleta (Linné, 1758)
lupinus (Linné, 1758)
- Pitar*
mediterranea (Tiberi, 1855)
rudis (Poli, 1795)
- Callista*
chione (Linné, 1758)
- Tapes*
decussata (Linné, 1758)
philippinarum (Adams & Reeve, 1850)
- Irus*
irus (Linné, 1758)
- Paphia**
aurea (Gmelin, 1791)
dura (Gmelin, 1791)
lucens (Locard, 1886)
rhomboides (Pennant, 1777)
textilis (Gmelin, 1791)
- Venerupis*
geographica (Gmelin, 1791)
pullastra (Montagu, 1803)
saxatilis (Fleuriau de Bellevue, 1802)
- Family Petricolidae**
Petricola
lithophaga (Philippson, 1788)
pholadiformis Lamarck, 1818
- Lajonkairia*
lajonkairii (Payraudeau, 1826)
- Mysia*
undata (Pennant, 1777)
- Family Myidae**
Mya
arenaria Linné, 1758
truncata Linné, 1758
- Sphenia*
binghami Turton, 1822
rueppelli Adams A., 1850
- Family Corbulidae**
Corbula
gibba (Olivi, 1792)
- Lentidium*
mediterraneum (Costa O.G., 1829)
- Family Gastrochaenidae**
Gastrochaena
cymbium Spengler, 1783
dubia (Pennant, 1777)
- Family Hiatellidae**
Hiatella
arctica (Linné, 1767)
rugosa (Linné, 1767)
- Panomya*
norvegica (Spengler, 1793)
- Panopea*
glycimeris (Von Born, 1778)
- Saxicavella*
angulata (Wood S., 1857)
jeffreysi Winckworth, 1930
- Cyrtodaria*
siliqua (Spengler, 1793)
- Family Pholadidae**
Pholas
dactylus Linné, 1758
- Barnea*
candida (Linné, 1758)
parva (Pennant, 1777)
- Pholadidea*
loscombiana Goodall in Turton, 1819
- Zirfaea*
crispata (Linné, 1767)
- Family Teredinidae**
Teredo
bartschi Clapp, 1923
bipartita (Jeffreys, 1860)
navalis Linné, 1758
utriculus Gmelin, 1791
- Lyrodus*
pedicellatus (de Quatrefages, 1849)
- Psiloteredo*
megotara (Hanley in Forbes & Hanley, 1848)
senegalensis (de Blainville, 1824)
- Teredora*
malleola (Turton, 1822)
- Bankia*
bipennata (Turton, 1819)
carinata (Gray J.E., 1827)

- jimbriatula* (Moll & Roch, 1931)
- Nototeredo*
norvegica (Spengler, 1792)
- Teredothyra*
excavata (Jeffreys, 1860)
- Spathoteredo*
spatha (Jeffreys, 1860)
- Family Xylophagidae**
- Xylophaga*
anselli (Harvey, 1996)
dorsalis (Turton, 1819)
gagei (Harvey, 1996)
nidarosiensis Santhakumaran, 1980
noradi Santhakumaran, 1980
praestans Smith E.A., 1903
ricei (Harvey, 1996)
- Xyloredo*
ingolfia (Turner, 1972)
- Family Pholadomyidae**
- Pholadomya*
loveni Jeffreys, 1882
- Family Thraciidae**
- Thracia*
convexa (Wood W., 1815)
corbuloides Deshayes, 1830
devexa Sars G.O., 1878
distorta (Montagu, 1803)
myopsis Beck in Møller, 1842
papyracea (Poli, 1791)
pubescens (Pulteney, 1799)
villosiuscula (MacGillivray, 1827)
- Family Laternulidae**
- Laternula*
subrostrata (Lamarck, 1818)
- Family Periplomatidae**
- Cochlodesma*
praetenua (Pulteney, 1799)
tenerum Fischer P., 1882
- Family Clavagellidae**
- Clavagella*
aperta Sowerby G.B. I., 1824
balanorum Scacchi, 1841
melitensis Broderip, 1835
- Family Pandoridae**
- Pandora*
glacialis Leach, 1819
inaequivalvis (Linné, 1758)
pimma (Montagu, 1803)
- Family Lyonsiidae**
- Lyonsia*
arenosa (Møller, 1842)
norwegica (Gmelin, 1791)
- Allogramma*
formosa (Jeffreys, 1882)
- Family Poromyidae**
- Poromya*
granulata (Nyst & Westendorp, 1839)
neaeroides Seguenza G., 1877
tornata (Jeffreys, 1876)
- Cetoconcha*
braziliensis Allen & Morgan, 1981
- Family Verticordiidae**
- Verticordia*
acuticostata (Philippi, 1844)
angulata (Jeffreys, 1882)
granulata Seguenza G., 1860
quadrata Smith E.A., 1885
sinuosa (Jeffreys, 1882)
trapezoidea Seguenza G., 1876
- Halicardia*
axinoides Seguenza G., 1876
ferruginea Di Geronimo, 1974
- Lyonsiella*
abyssicola Sars G.O., 1878
compressa (Locard, 1898)
frielei Allen & Turner, 1974
smidti (Friele, 1886)
subquadrata (Jeffreys, 1881)
- Policordia*
densicostata (Locard, 1898)
gemma (Verrill, 1880)
jeffreysi (Friele, 1879)
- Family Cuspidariidae**
- Cuspidaria*
arctica (Sars M., 1859)
atlantica Allen & Morgan, 1981
canariensis De Boer, 1985
centobi Bouchet & Warén, 1979
circinata (Jeffreys, 1876)
cuspidata (Olivi, 1792)
elliptica Di Geronimo, 1974
exigua (Jeffreys, 1876)
glacialis (Sars G.O., 1878)
inflata (Jeffreys, 1882)
jeffreysi (Dall, 1881)
Jugosa (Wood S., 1856)
lamellosa (Sars G.O., 1878)
obesa (Lovén, 1846)
obliqua De Boer, 1985
parva Verrill & Bush, 1898
rostrata (Spengler, 1793)
subtorta (Sars G.O., 1878)
teres (Jeffreys, 1882)
truncata (Jeffreys, 1882)
ventricosa Verrill & Bush, 1898
wollastoni (Smith E.A., 1885)
- Tropidomya*
abbreviata (Forbes, 1843)
- Cardiomya*
costellata (Deshayes, 1835)
curta (Jeffreys, 1876)
knudseni Allen & Morgan, 1981
striata (Jeffreys, 1876)
- Rhinoclama*
halimera (Dall, 1886)
notabilis (Jeffreys, 1876)
- Myonera*
angularis (Jeffreys, 1876)
atlantica Allen & Morgan, 1981
paucistriata (Dall, 1881)
tillamookensis Dall, 1916
- Halonympha*
depressa (Jeffreys, 1882)
- Protocuspidaria*
atlantica (Allen & Morgan, 1981)
azorica (Smith E.A., 1885)
colpodes (Dautzenberg & Fischer H., 1897)
simplicis (Allen & Morgan, 1981)
- Class Scaphopoda**
- Family Dentaliidae**
- Dentalium*
agile Sars M. in Sars G.O., 1872
dentalis Linné, 1758
entalis Linné, 1758
inaequicostatum (Bucquoy, Dautzenberg & Dollfus, 1891)
novemcostatum (Lamarck, 1818)
occidentale Stimpson, 1851
panormum Chenu, 1858
rossati Caprotti, 1966
vulgare da Costa, 1778
- Fissidentalium*
candidum (Jeffreys, 1877)
capillosum (Jeffreys, 1877)
ergasticum (Fischer P., 1882)
exuberans (Locard, 1897)

- scamnamum* (Locard, 1897)
Laevidentalium
caudani (Locard, 1896)
- Family Fustiariidae**
Fustiaria
rubescent (Deshayes, 1825)
- Family Gadilinidae**
Episiphon
filum (Sowerby G.B. II, 1860)
- Family Entalinidae**
Entalina
tetragona (Brocchi, 1814)
Bathoxiphus
ensiculus (Jeffreys, 1877)
Heteroschismoides
subterfissum (Jeffreys, 1877)
- Family Pulsellidae**
Pulsellum
affine (Sars G.O., 1878)
lofotense (Sars M., 1865)
teres (Jeffreys, 1883)
- Family Gadiliidae**
Cadulus
amphora (Jeffreys, 1883)
artatus (Locard, 1897)
cylindratus (Jeffreys, 1877)
gibbus (Jeffreys, 1880)
gracilis (Jeffreys, 1877)
jeffreysi (Monterosato, 1875)
monterosatoi (Locard, 1897)
- Gadila*
olivii (Scacchi, 1835)
- Cadulus*
ovulum (Philippi, 1844)
propinquus (Sars G.O., 1878)
strangulatus (Locard, 1897)
subfusiformis (Sars M., 1865)
tumidostus (Jeffreys, 1877)
- Dischides*
politus (Wood S., 1842)
- Siphondentalium*
laubieri Bouchet & Warén, 1979
lobatum (Sowerby G.B. II, 1860)
- Class Cephalopoda**
- Family Spirulidae**
Spirula
spirula (Linné, 1758)
- Family Sepiidae**
Sepia
bertheloti (d'Orbigny, 1839)
elegans de Blainville, 1827
officinalis Linné, 1758
orbigniana de Féussac, 1826
- Family Sepiolidae**
Rossia
macrosoma (Delle Chiaje, 1828)
- Neorossia*
caroli (Joubin, 1902)
- Heteroteuthis*
dispar (Rüppell, 1845)
- Stoloteuthis*
leucophterus (Verrill, 1878)
- Rondeletiola*
minor (Naef, 1912)
- Sepiola*
affinis Naef, 1912
atlantica d'Orbigny, 1839
aurantiaca Jatta, 1896
intermedia Naef, 1912
ligulata Naef, 1912
robusta Naef, 1912
rondeleti Leach, 1817
- steenstrupiana* Levy, 1912
Sepieta
neglecta Naef, 1916
obscura Naef, 1916
oweniana (d'Orbigny, 1839)
- Family Loliginidae**
Loligo
forbesi Steenstrup, 1856
vulgaris Lamarck, 1798
- Alloleuthis*
medius (Linné, 1758)
subulatus (Lamarck, 1798)
- Family Ctenopterygidae**
Ctenopteryx
siculus (Vérany, 1851)
- Family Lycoteuthidae**
Lampadioteuthis
megaleius Berry, 1916
- Family Enoplateuthidae**
Abralia
veranyi (Rüppell, 1844)
- Abraaliopsis*
pfefferi Joubin, 1896
- Family Pyroteuthidae**
Pyroteuthis
margaritifer (Rüppell, 1844)
- Pterygioteuthis*
giardi Fischer, 1896
- Family Ancistrocheiridae**
Ancistrocheirus
lesueurii (de Féussac & d'Orbigny, 1835)
- Family Octopoteuthidae**
Octopoteuthis
siculus Rüppell, 1844
- Taningia*
danae Joubin, 1931
- Family Onychoteuthidae**
Chaunoteuthis
mollis Appellöf, 1890
- Onychoteuthis*
banksii (Leach, 1817)
- Ancistroteuthis*
lichtensteinii (de Féussac & d'Orbigny, 1839)
- Onykia*
carriboea Lesueur, 1821
- Family Cycloteuthidae**
Cycloteuthis
sirventi Joubin, 1919
- Family Gonatidae**
Gonatus
steenstrupi Kristensen, 1981
- Family Lepidoteuthidae**
Lepidoteuthis
grimaldii Joubin, 1895
- Family Histiopteuthidae**
Histioteuthis
bonnellii (de Féussac, 1835)
- corona* (Voss & Voss, 1962)
- doylei* (Pfeffer, 1912)
- elongatus* (Voss & Voss, 1962)
- meleagrotoeuthis* (Chun, 1910)
- reversus* (Verrill, 1880)
- Family Bathytteuthidae**
Bathyteuthis
abyssicola Hoyle, 1885
- Family Brachioteuthidae**
Brachioteuthis
riisei (Steenstrup, 1882)
- Family Ommastrephidae**
Illex

- coindetii* (Vérany, 1839)
Todaropsis
eblanae (Ball, 1841)
Todarodes
sagittatus (Lamarck, 1798)
Ommastrephes
bartrami (Lesueur, 1821)
Sthenoteuthis
pteropus (Steenstrup, 1855)
- Family Thysanoteuthidae**
Thysanoteuthis
rhombus Troschel, 1857
- Family Chiroteuthidae**
Chiroteuthis
picteti Joubin, 1894
veranyi (de Féussac, 1835)
Valbyteuthis
danae Joubin, 1931
- Family Mastigoteuthidae**
Mastigoteuthis
atlanticus Joubin, 1933
flammeus Chun, 1908
grimaldii (Joubin, 1895)
schmidti Degner, 1925
talismani (Fischer & Joubin, 1906)
- Family Cranchiidae**
Leachia
atlantica (Degner, 1925)
Liocranchia
reinhardtii (Steenstrup, 1856)
Helicocranchia
pfefferi Massy, 1907
Bathothauma
lyromma Chun, 1906
Taonius
pavo (Lesueur, 1821)
Galiteuthis
armatus Joubin, 1898
Teuthowenia
megalops (Prosche, 1847)
Megalocranchia
speculator (Chun, 1906)
- Family Vampyroteuthidae**
Vampyroteuthis
infernalis Chun, 1903
- Family Cirroteuthidae**
Cirrothauma
murrayi Chun, 1911
- Family Opisthoteuthidae**
Opisthoteuthis
agassizi Verrill, 1883
grimaldii (Joubin, 1903)
Grimpoteuthis
umbellatus (Fischer, 1883)
- Family Bolitaenidae**
Japetella
diaphana Hoyle, 1885
- Family Vitreledonellidae**
Vitreledonella
richardi Joubin, 1918
- Family Octopodidae**
Octopus
defilippi Vérany, 1851
macropus Risso, 1826
salutii Vérany, 1836
vulgaris Cuvier, 1797
Scaeurgus
unicirrus (Delle Chiaje, 1838)
Pteroctopus
tetricirrus (Delle Chiaje, 1830)
Eledone
cirrhosa (Lamarck, 1798)
- moschata* (Lamarck, 1798)
Bathytopypus
arcticus (Prosche, 1849)
proschi Muus, 1962
sponsalis (Fischer & Fischer, 1892)
Benthocotpus
ergasticus (Fischer & Fischer, 1892)
- Family Tremoctopodidae**
Tremoctopus
violaceus Delle Chiaje, 1830
- Family Ocythoidae**
Ocythoe
tuberculata Rafinesque, 1814
- Family Argonautidae**
Argonauta
argo Linné, 1758
- Family Alloposidae**
Haliphron
atlanticum Steenstrup, 1861

ANNELIDA

Polychaeta

Compiled by Gérard Bellan

The list of polychaetes was compiled from several published and unpublished lists. Among the most important published sources were Castelli *et al.* (1995), Ariño (1987), Mackie and Erséus (1997), and Holte and Brattegard (1997). Publications on the polychaete fauna of Germany (Hartmann-Schröder 1996, Böggemann 1998), and many short papers too numerous to cite here, were also used. Unpublished lists of polychaetes of the Mediterranean and Black Seas established by Dr K. Arvanitidis (Heraklion, Greece), for the English Channel, the French Atlantic, and northern coast of Spain, by Dr J.C. Dauvin and F. Gentil, and Dr G. Bachelet (Bay of Biscay from west of Brittany to the coast of Galicia at 09° W), respectively, have provided unpublished lists. H. G. Hansson provided information concerning Iceland and Greenland. Drs R. Barnich and D. Fiege generously helped in correcting the general list and especially the Aphroditidae, Polynoidae, Acoetidae, Pholoidae, Eulepethidae, Sigalionidae, Magelonidae, Owenidae and the Serpulidae. Dr Mary E. Petersen reviewed the list in general and especially the Chaetopteridae, Cirratulidae, Cossuridae, Ctenodrilidae, Fauveliopsidae, Longosomatidae, Pholoidae and some Spionidae. The compiler is extremely grateful to all that assisted.

The classification of polychaete families into orders have changed significantly in recent years and it is likely that these changes will continue to occur.

The ERMS project has tended to be conservative in relation to classifications to minimise confusing non-specialist readers. Therefore systematic hierarchy has been delimited to family level and families, genera and species have been arranged in alphabetic order. The Pogonophora have been reduced to the rank of Family within the Polychaeta by Rouse and Fauchald (1997), who revert them to the name Siboglinidae Caullery, 1914 within the Order Sabellida. However, few non-specialists will be aware of this change so the Pogonophora have been kept distinct in this volume.

However due to this kind of compilation from very different origins, it was necessary to decide among several nomenclatural denominations of genus and species. In fact, only few families have been recently studied taking into account all the species of the ERMS geographical extension. This list is expected to be improved in terms of taxonomy and also in terms of the validity of some species with future research. Therefore apart from the printed version please check the web site for updated information.

References

- Ariño, A.H. 1987. Bibliografía Iberica de Poliquetos. Base de datos y Catálogo de especies.(Iberian bibliography about Polychaeta. Data Base and Catalogue of Species). *Publicaciones de Biología de la Universidad de Navarra Serie Zoologica*, 16.
- Arvanitidis, C. 2000. Polychaete fauna of the Aegean Sea: Inventory and new information. *Bulletin of Marine Sciences*, 66: 73-96
- Böggemann, M. 1998. Polychaeten aus der Deutschen Bucht. *Courier Forschungsinstitut Senckenberg* 202, 315 p.
- Castelli, A., Abbiati, M., Badalamenti, F., Bianchi, C.N., Cantone, G., Gambi, M.C., Giangrande, A., Gravina, M.F., Lanera, P., Lardicci, C., Somaschini, A. & Sordino, P. 1995. Annelida Polychaeta, Pogonophora, Echiura, Sipunculida. 45 pp. In: Checklist delle specie della fauna italiana,. 19. Calderini, Bologna 1-45.
- Hartmann-Schröder, G. 1996. Annelida, Borstewürmer, Polychaeta. *Die Tierwelt Deutschland* 58, Teil, 648 p. Mit 295 Abbildungen, 2 neuarbeitete Auflage.
- Holte, T. & Brattegard, T. 1997. Class Polychaeta (phylum Annelida). pp. 115-129 In: Brattegard, T. & Holte, T. 1997. *Distribution of marine, benthic macroorganisms in Norway. A tabulated catalogue. Preliminary Edition*. Research Report for Directorate for Nature Management. 409 pp.
- Mackie, A.S.Y. & Erséus, C. 1997. Annelida. In: Howson C. M. and Picton B. E. (eds), *The Species Directory of the Marine Fauna and Flora of the British Isles and surrounding Seas*. Marine Conservation Society, Ross-on-Wye, pp 109-146.
- Rouse, G.W. & Fauchald, K. 1997. Cladistics and polychaetes. *Zoologica Scripta* 28, 139-204.

Class Polychaeta

Family Acoetidae

- Euarche*
tubifex Ehlers, 1887
- Eupanthalis*
kinbergi McIntosh, 1876
- Eupolyodontes*
gulo (Grube, 1855)
- Panthalis*
oerstedi Kinberg, 1856
- Polyodontes*
maxillosus (Ranzani, 1817)

Family Acrocirridae

- Acrocirrus*
frontifilis (Grube, 1860)
- Chaviniella*
arctica Averintsev, 1980
biscayensis Laubier, 1974
- Flabelligella*
biscayensis Kolmer, 1985
mediterranea Kolmer, 1985
- Macrochaeta*
caroli Westheide, 1981
clavicornis (M. Sars, 1835)
helgolandica Friedrich, 1937
natalensis Hartmann-Schröder, 1996
polyonyx Eliason, 1962

Family Aeolosomatidae

- Aeolosoma*
polyonyx Bunke, 1967
litorale Bunke, 1967

Family Alciopidae

- Alciopa*
reynaudi Audouin & Milne-Edwards, 1833
- Alciopina*
parasitica Claparède & Panceri, 1867
- Corynocephalus*
albomaculatus Levinsen, 1885
- Krohnia*
lepidota (Krohn, 1845)
- Naiades*
cantrainii Delle Chiaje, 1928
- Plotolhelmis*
capitata (Greef, 1876)
tenuis (Apstein, 1893)
- Rhynchonerella*
angelinia (Kinberg, 1866)
fulgens Greeff, 1885
gracilis A. Costa, 1864
moebii (Apstein, 1893)
nasuta (Greeff, 1876)
petersii (Langerhans, 1880)
setosa (Greeff, 1885)
- Torrea*

- candida* (Delle Chiaje, 1828)
Vanadis
crystallina Greeff, 1876
formosa Claparède, 1870
longissima (Levinsen, 1885)
tentaculata Langerhans, 1880
- Family Ampharetidae**
- Adercodon*
pleijeli Mackie, 1994
- Alkmaria*
romijni Horst, 1919
- Amage*
adspersa (Grube, 1863)
auricula Malmgren, 1866
gallasii Marion, 1875
scotica Clark, 1952
- Amagopsis*
klugei Pergament & Hlebovic in Chlebovich, 1964
- Ampharete*
baltica Eliason, 1955
acutifrons (Grube, 1860)
falcata Eliason, 1955
finmarchica (M. Sars, 1864)
goesi Malmgren, 1866
gracilis Rioja, 1931
lindstroemi Malmgren, 1867 sensu Hessle, 1917
petersenae Zhirkov, 1997
vega (Wirén, 1883)
- Amphicteis*
gunneri (M. Sars, 1835)
midas (Gosse, 1855)
ninonae Jirkov, 1985
sundevalli Malmgren, 1865
- Amythasides*
macroglossus Eliason, 1955
- Anobothrus*
gracilis (Malmgren, 1866)
- Asabellides*
siberica (Wirén, 1883)
- Auchenoplax*
crinita Ehlers, 1887
- Ectyssipe*
vanelii (Fauvel, 1936)
- Glyphanostomum*
abyssale Day, 1967
pallescens (Théel, 1879)
- Hypania*
antiqua (Ostrooumoff, 1896)
brevispinis (Grube, 1860)
invalida (Grube, 1860)
- Hypaniola*
kowalewskii (Grube, 1877)
- Isolda*
whydahensis Augener, 1918
- Lysippe*
labiata Malmgren, 1866
- Lysippides*
fragilis (Wollebaeck, 1912)
- Melissa*
albincincta Mackie & Pleijel, 1995
collare Detinova, 1985
cristata (M. Sars, 1851)
elisabethae McIntosh, 1885, sensu Mackie
islandica Saemundsson, 1918
monoceroides Fauvel, 1936
palmata Grube, 1870
- Melinnampharete*
laubieri (Desbruyères, 1978)
- Melinopsis*
arctica (Annenkova, 1931)
- Melythasides*
laubieri Desbruyères, 1978
- Mugga*
bathyalis Holthe, 1986
wahrbergi Eliason, 1955
- Neosabellides*
oceanica (Fauvel, 1909)
- Noanelia*
hartmanae Desbruyères & Laubier, 1977
- Paedampharete*
acutiseries Russell, 1987
- Sabellides*
borealis M. Sars, 1856
octocirrata (M. Sars, 1835)
- Samytha*
sexcircrata (M. Sars, 1856)
- Samythella*
neglecta Wollebaek, 1912
- Sosane*
sulcata Malmgren, 1865
- Sosanopsis*
Wiréni Hessle, 1917
- Uschakovius*
enigmaticus Laubier, 1973
- Zatsepinia*
rittichae Jirkov, 1986
- Family Amphinomidae**
- Amphinome*
rostrata (Pallas, 1766)
- Chloeia*
venusta Quatrefages, 1865
viridis Schmarda, 1861
- Chloenopsis*
atlantica (McIntosh, 1885)
- Eurythoe*
complanata (Pallas, 1776)
syriaca Kinberg, 1857
- Hermodice*
carunculata (Pallas, 1776)
- Hipponoe*
gaudichaudii (Audouin & Milne Edwards, 1830)
- Linopherus*
canariensis Langerhans, 1881
hemuli (Fauchald, 1972)
paucibranchiata Fauvel, 1932
- Notopygos*
megalops McIntosh, 1885
- Paramphinoe*
jeffreysii (McIntosh, 1868)
- Pareurythoe*
borealis (M. Sars, 1862)
- Family Antonbruuniidae**
- Antonbruunia*
viridis Hartman & Boss, 1965
- Family Aphroditidae**
- Aphrodita*
aculeata Linnaeus, 1758
alta Kinberg, 1856
perarmata Roule, 1898
- Aphrogenia*
alba Kinberg, 1856
- Laetmonice*
filicornis Kinberg, 1856
hystrix (Savigny, 1820)
producta Grube, 1876
producta subsp. *britannica* McIntosh, 1900
- Pontogenia*
chrysecoma (Baird, 1865)
sericoma Ehlers, 1867
- Family Apistobranchidae**
- Apistobranchus*
tenuis Orrrage, 1962
tullbergi (Théel, 1879)
- Family Arabellidae**
- Arabella*
coeca Fauvel, 1940
geniculata (Claparède, 1868)
iricolor (Monagu, 1804)
- Drilonereis*

- brattstroemi* Fauchald, 1972
filum (Claparède, 1868)
- Haematocheleps*
terrebellidis Wirén, 1886
- Laborostratus*
jonicus Tenerelli, 1961
parasiticus de Saint Joseph, 1888
- Notocirrus*
arcachonis Quatrefages, 1843
scoticus McIntosh, 1869
- Oligognathus*
bonelliae Spengel, 1882
parasiticus Cerruti, 1909
- Family Arenicolidae**
- Abarenicola*
claparedei (Levinsen, 1883)
pusilla (Quatrefages, 1865)
- Arenicola*
cristata Stimpson, 1856
defodiens Cadman & Nelson-Smith, 1993
marina (Linnaeus, 1758)
- Arenicolides*
branchialis (Audouin & Milne Edwards, 1833)
grubii Claparède, 1868
ecaudata (Johnston, 1835)
- Branchiomaldane*
vincenti Langerhans, 1881
- Family Calamyzidae**
- Calamyza*
amphictenicola Arwidsson, 1932
- Family Capitellidae**
- Baldia*
johnstoni Garwood & Bamber, 1988
- Capitella*
capitata (Fabricius, 1780)
giardi (Mesnil, 1897)
hermaphrodita Boletzky & Dohle, 1967
minima Langerhans, 1880
- Capitellethus*
dispar (Ehlers, 1907)
- Dasybranchus*
caducus (Grube, 1846)
carneus Grube, 1870
gajolae Eisig, 1887
- Heteromastus*
filiformis (Claparède, 1864)
- Leiocapitella*
dollfusi (Fauvel, 1936)
glabra Hartman, 1947
- Leiochrides*
norvegicus Fauchald, 1972
- Mastobranchus*
dollfusi Fauvel, 1936
trinchesii Eisig, 1887
- Mediomastus*
capensis Day, 1961
cirripes Ben-Eliah, 1976
fragilis Rasmussen, 1973
- Neopseudocapitella*
brasiliensis Rullier & Amoureaux, 1979
- Notomastus*
aberans Day, 1957
agassizii McIntosh, 1885
exsertilis de Saint Joseph, 1906
formianus Eisig, 1887
latericeus M. Sars, 1851
lineatus Claparède, 1870
profundus Eisig, 1887
rubicundus (Keferstein, 1862)
- Notoproctus*
oculatus Arwidsson, 1907
- Paracapitella*
southwardi Kirkegaard, 1983
- Paraleiocapitella*
- mossambica* Thomassin, 1970
- Peresiella*
clymenoides Harmelin, 1968
- Pseudocapitella*
incerta Fauvel, 1913
incerta subsp. *aberrans* Amoureaux, 1972
- Pseudoleiocapitella*
fauveli Harmelin, 1964
- Pseudomastus*
delticus Capaccioni-Azzati & Martin, 1992
- Pseudonotomastus*
southerni Warren & Parker, 1994
- Family Chaetopteridae**
- Chaetopterus*
norvegicus M. Sars, 1835
sarsi Böck in M. Sars, 1851
variopedatus (Renier, 1804)
- Mesochaetopterus*
minutus (Claparède, 1868)
sagittarius (Claparède, 1868)
- Phyllochaetopterus*
anglicus Potts, 1914
gracilis Grube, 1863
major Claparède, 1870
socialis Claparède, 1870
- Spiochaetopterus*
bergensis Gitay, 1969
solitarius (Rioja, 1917)
costarum (Claparède, 1868)
typicus M. Sars, 1856
- Family Chrysopetalidae**
- Bhawania*
goodei Webster, 1884
reynsi Katzmann, Laubier & Ramos, 1974
- Chrysopetalum*
debile (Grube, 1855)
- Dysponetus*
bipapillatus Dahlgren, 1996
caecus (Langerhans, 1880)
gracilis Hartman, 1965
palaeophorus Hartmann-Schröder, 1974
pygmaeus Levinsen, 1879
- Heteropale*
bellis Johnson, 1897
- Paleanotus*
chrysolepis Schmarda, 1861
schmardai Mileikowski, 1962
- Vigorniella*
zaikai (Kisseleva, 1992)
- Family Cirratulidae**
- Aphelochaeta*
filiformis (Keferstein, 1862)
marioni (de Saint Joseph, 1894)
mcintoshii (Southern, 1914)
serrata (Eliason, 1962)
vivipara (Christie, 1984)
- Caulleriella*
alata (Southern, 1914)
bioculata (Keferstein, 1862)
killianiensis (Southern, 1914)
parva Gilliland, 1979
viridis (Langherans, 1880)
zelandica (McIntosh, 1911)
- Chaetozone*
abranchiata (G.A. Hansen, 1878)
benthaliana McIntosh, 1885
caputesocis (de Saint Joseph, 1894)
carpenteri (McIntosh, 1911)
gibber Woodham & Chambers, 1994
macrophthalma Langherans, 1880
setosa Malmgren, 1867
- Cirratulus*
abyssorum Hansen, 1878
caudatus Levinsen, 1893

- cirratus* (O.F. Müller, 1776)
chrysoderma Claparède, 1868
incertus McIntosh, 1916
glandularis Langerhans, 1884
Cirriformia
semicincta (Ehlers, 1905)
tentaculata (Montagu, 1808)
Cirrhinereis
incertus Fauvel, 1936
Dodecaceria
ater (Quatrefages, 1866)
caulleryi Dehorne, 1933
concharum Oersted, 1843
diceria Hartman, 1951
jimbriata sensu Gibson 1998
saxicola (Grube, 1855)
sextentaculata (Delle Chiaje, 1822-1826)
Monticellina
dorsobranchialis (Kirkegaard, 1959)
heterochaeta (Laubier, 1961)
tesselata (Hartman, 1960)
Protocirrineris
chrysoderma (Claparède, 1868)
Tharyx
multibranchiis (Grube, 1863)
retieri Lechapt, 1994
Timarete
anchylochaeta (Schmarda, 1861)
dasylophius (Marenzeller, 1879)
dolfusi (Fauvel, 1928)
filigera (Delle Chiaje, 1828)
norvegica (Quatrefages, 1866)
- Family Cossuridae**
Cossura
coasta sensu Bogdanos & Fredj, 1983
longocirrata Webster & Benedict, 1887
pygodactylata Jones, 1956
soyeri Laubier, 1963
- Family Ctenodrilidae**
Aphropharynx
heterochaeta Wilfert, 1974
Ctenodrilus
parvulus Scharff, 1887
serratus (Schmidt, 1857)
Raricirrus
berylli Petersen & George, 1991
Raphidrilus
nemasoma (Monticelli, 1910)
Zeppelinia
monostyla (Zeppelin, 1883)
- Family Diurodrilidae**
Diurodrilus
benazzii Gerlach, 1952
dohrni Gerlach, 1953
minimus Remane, 1925
subterraneus Remane, 1934
- Family Dorvilleidae**
Apharyngtus
punicus Westheide, 1971
Arenotrocha
minuta (Lévi, 1954)
Dinophilus
gigas Weldon, 1886
graciliatus O. Schmidt, 1857
taeniatus Harmer, 1889
vorticoides Schmidt, 1848
Dorvillea
albomaculata Åkesson & Rice, 1992
atlantica (McIntosh, 1885)
erucaeformis (Malmgren, 1865)
kastjani Tsetlin, 1980
romeri (Augener, 1912)
rubrovittata (Grube, 1855)
Iphitime
- cuenoti* Fauvel, 1914
hartmanae Kirkegaard, 1977
paguri Fage & Legendre, 1934
Mammiiphitome
cosmetandra (Oug, 1990)
Marycarmenia
lysandrae Nuñez, 1998
Ophryotroca
bacci Parenti, 1961
Ophryotrocha
dubia Hartmann-Schröder, 1974
gerlachi Hartmann-Schröder, 1974
geryonicola (Esmark, 1874)
gracilis Huth, 1933
hartmanni Huth, 1933
hartmani subsp. *medicea* Parenti, 1962
irinae Tsetlin, 1980
labronica La Greca & Bacci, 1962
littoralis (Levinsen, 1879)
longidentata Josefson, 1975
macrovifera Åkesson, 1975
maculata Åkesson, 1973
mediterranea Martin, Abello & Cartes, 1991
puerilis subsp. *puerilis* Claparède & Mecznikov, 1863
puerilis subsp. *siberti* (McIntosh, 1885)
robusta Åkesson, 1975
scarlatoi Averincev, 1989
socialis Ockelmann & Åkesson, 1990
Palpiphitime
lobifera (Oug, 1978)
Paraphryotrocha
isochaeta (Eliason, 1962)
Paraprodilus
psammophilus Westheide, 1965
Parougia
caeca (Webster & Benedict, 1884)
eliasoni (Oug, 1978)
macilenta (Oug, 1978)
nigridentata (Oug, 1978)
Pettiboneia
urciensis Campoy & San Martin, 1980
Protodorvillea
kefersteini (McIntosh, 1869)
Pusillotrocha
akessoni Westheide & von Nordheim, 1985
Schistomerings
caeca (Webster & Benedict, 1884)
neglecta (Fauvel, 1923)
subaequalis Oug, 1978
rudolphi (Delle Chiaje, 1828)
Trilobodrilus
axi Westheide, 1967
heideri Remane, 1925
- Family Eulepetidae**
Grubeulepis
augeneri Pettibone, 1969
katzmanni Pettibone, 1986
- Family Eunicidae**
Amphiro
simplex Langerhans, 1884
Eunice
antennata (Lamarck, 1818)
aphroditoides (Pallas, 1788)
cariboea Grube, 1856
dubitata Fauchald, 1974
floridana (Pourtales, 1867)
harassii Audouin & Milne-Edwards, 1833
norvegica (Linnaeus, 1767)
oerstedi Stimpson, 1853
pennata (O.F. Müller, 1776)
purea Grube, 1866
rubrocincta Ehlers, 1868
rousseauai Quatrefages, 1865
schizobranchia Claparède, 1870
senta (Moore, 1902)

- torquata* Quatrefages, 1865
vittata (Delle Chiaje, 1829)
- Lysibranchia*
paucibranchiata Cantone, 1983
- Lysidice*
collaris Grube, 1870
margaritacea Claparède, 1868
ninetta Audouin & Milne-Edwards, 1833
- Marpissa*
bellii (Audouin & Milne-Edwards, 1833)
fallax Marion & Bobretzky, 1875
kinbergi McIntosh, 1910
sanguinea (Montagu, 1815)
saxicola Langerhans, 1881
- Nematonereis*
unicornis (Grube, 1840)
- Palola*
siciliensis (Grube, 1840)
- Family Euniphysidae**
Paraeuniphysa
italica (Cantone & Gravina, 1991) n. comb.
- Family Euphrasinidae**
Euphrasine
armadillo Sars, 1851
borealis Orsted, 1843
cirratepropinqua Amoureux, 1982
foliosa Audouin & Milne-Edwards, 1833
myrtosa Savigny, 1818
- Euphrasyne*
cirrata M. Sars, 1851
- Palmyreuphrasyne*
paradoxa Fauvel, 1914
- Family Fauveliopsidae**
Fauveliopsis
adriatica Katzmann & Laubier, 1974
brattegardi Fauchald, 1972
brevis (Hartman, 1965)
cabiochi Amoureux, 1982
fauchaldi Katzmann & Laubier, 1974
jameoquensis Nuñez in Nuñez, Ocaña & Brito, 1997
olgae Hartmann-Schröder, 1983
scabra Hartman & Fauchald, 1971
- Family Flabelligeridae**
Brada
granulosa Hansen, 1880
incrassata Støp-Bowitz, 1948
inhabilis (Rathke, 1843)
rugosa (Hansen, 1882)
villosa (Rathke, 1843)
- Diplocirrus*
glaucus (Malmgren, 1867)
hirsutus (Hansen, 1879)
longisetosus (Marenzeller, 1895)
- Flabelligera*
affinis M. Sars, 1829
biscayensis Kolmer, 1985
diplochaitius (Otto, 1820)
infundibularis H.P. Johnson, 1901
- Ilyphagus*
ilyvestris Hartman, 1960
- Pherusa*
arctica Støp-Bowitz, 1948
falcata Støp-Bowitz, 1948
flabellata (M. Sars in G.O. Sars 1872)
fucicola Leach
monilifera (Delle Chiaje, 1841)
plumosa (O.F. Müller, 1776)
- Piromis*
eruca (Claparède, 1870)
- Therochaeta*
collarifera (Ehlers, 1887)
- Family Glyceridae**
Glycera
alba (O.F. Müller, 1776)
- alba* subsp. *adspersa* Fauvel, 1939
capitata Oersted, 1843
celtica O'Connor, 1987
convoluta subsp. *uncinata* Rioja, 1918
dayi O'Connor, 1987
gigantea Quatrefages, 1865
lapidum Quatrefages, 1865
mimica Hartman, 1965
minuta (Bobretzky, 1870)
oxycephala Ehlers, 1887
rouxi Audouin & Milne Edwards, 1833
tesselata Grube, 1863
tesselata subsp. *minor* La Greca, 1946
tridactyla Schmarda, 1861
unicornis Savigny, 1818
- Glycerella*
atlantica Wesenberg-Lund, 1950
magellanica (McIntosh, 1885)
- Hemipodus*
septentrionalis Roule, 1896
- Family Goniadidae**
Glycinde
bonhourei Gravier, 1904
nordmanni (Malmgren, 1865)
- Goniada*
emerita Audouin & Milne-Edwards, 1833
euxina Jakubova, 1930
gallica Rioja, 1923
maculata Oersted, 1843
norvegica Oersted, 1844
pallida Arwidsson, 1898
- Goniadella*
bobrezkii (Annenkova, 1929)
gallica Rioja, 1923
gracilis (Verrill, 1973)
unicirra Agirrezabalaga, 1980
- Progonyada*
regularis Hartman, 1965
- Family Hesionidae**
Amphiduros
fuscescens (Marenzeller, 1875)
- Anoploneurus*
herrmanni Giard, 1882
- Dalhousiella*
carpenteri McIntosh, 1901
- Glyphohesione*
klatti Friedrich, 1950
- Gyptis*
brevipalapa (Hartmann-Schröder, 1959)
golikovi (Averincev, 1990)
mackiei Pleijel, 1993
mediterranea Pleijel, 1993
propinqua Marion & Bobretzky, 1875
rosea (Malm, 1874)
- Hesione*
splendida Savigny, 1818
steenstrupii de Quatrefages, 1886
- Hesionides*
arenaria Friedrich, 1937
gohari Hartmann-Schröder, 1960
maxima Westheide, 1967
- Hesiospina*
similis (Hessle, 1925)
- Kefersteinia*
cirrata (Keferstein, 1862)
cirrata subsp. *hibernica* (Southern, 1914)
- Leocrates*
atlanticus (McIntosh, 1885)
chinensis Kinberg, 1866
- Microphthalmus*
aberrans (Webster & Benedict, 1887)
bifurcatus Hartmann-Schröder, 1974
ephippiophorus Clausen,
fragilis Bobretzky, 1870
listensis Westheide, 1967

- pseudoaberrans* Campoy & Vieitez, 1982
sczelkowii Mecznikow, 1865
sczelkowii subsp. *cantabrica* Rioja, 1925
southerni Westheide, 1967
similis Bobretzky, 1870
tyrrhenicus Zunareli Vandini, 1967
- Nereimyra**
aphroditoides (O. Fabricius, 1780)
artica (Malmgren, 1867)
aurantiaca (M. Sars, 1862)
britannica (Chamberlin, 1920)
longicornis (M. Sars, 1862)
multipapillata (Théel, 1879)
punctata (O.F. Müller, 1788)
- Ophiodromus**
agilis (Ehlers, 1864)
flexuosus (Delle Chiaje, 1825)
longocirratus Tenerelli, 1974
pallidus (Claparède, 1864)
pelagicus (Claparède, 1864)
- Orseis**
pulla Ehlers, 1864
- Periboea**
longocirrata Ehlers, 1864
- Podarke**
obscura Verrill, 1873
- Podarkeopsis**
arenicola (La Greca, 1947)
capensis (Day, 1963)
galangaui Laubier, 1961
helgolandica Hilbig & Dittmer, 1979
- Syllidia**
armata Quatrefages, 1865
assimilis (Pryde, 1914)
- Family Histiobdellidae**
Histiobdella
homari van Beneden, 1858
- Family Ichthyotomidae**
Ichthyotomus
sanguinarius Eisig, 1906
- Family Iopsilidae**
Phalacroforus
pictus Greeff, 1879
uniformis Reibsich, 1875
- Family Lacydoniidae**
Lacydonia
eliasoni Hartmann-Schröder, 1996
laureci Laubier, 1975
miranda Marion & Bobretzky, 1875
- Family Longosomatidae**
Heterospio
longissima Ehlers, 1874
mediterranea Laubier, Picard & Ramos, 1973
reducta Laubier, Picard & Ramos, 1973
- Family Lopadorrhynchidae**
Lopadorrhynchus
appendiculatus Southern, 1909
brevis Grube, 1855
krohnii (Claparède, 1870)
nationalis Reibish, 1895
uncinatus Fauvel, 1915
- Maupasia**
caeca Viguier, 1886
isochaeta (Reibish, 1895)
magna Southern, 1909
- Pedinosoma**
curtum Reibish, 1995
- Pelagobia**
longicirrata Greef, 1879
serrata Southern, 1909
- Family Lumbrineridae**
Abyssiniae
abyssorum (McIntosh, 1885)
- scopa* (Fauchald, 1974) sensu Miura 1980
Abyssiniae
scopa subsp. *aequilobata* (Winsnes, 1881)
- Augeneria**
algida (Wirén, 1901)
tentaculata Monroe, 1930
- Eranio**
ehlersi (MacIntosh, 1885)
- Lumbricalus**
adriatica (Fauvel, 1940)
- Lumbrinerides**
acuta (Verrill, 1875)
amoureuxi Miura, 1980
carpinei (Ramos, 1976)
laubieri Miura, 1980
neogesae Ramos, 1976
- Lumbrineriopsis**
gasconensis Miura, 1980
magnidendata Winsnes, 1981
paradoxa (de Saint Joseph, 1888)
- Lumbrineris**
agastos Fauchald, 1974
albidodata (Ehlers, 1908)
aniara Fauchald, 1974
bifrons (Kinberg, 1865)
cingulata (Ehlers, 1897)
cluthensis (Clark, 1953)
coccinea (Renier, 1804)
flabellicola (Fage, 1936)
gracilis (Ehlers, 1868)
hibernica (McIntosh, 1903)
inflata (Moore, 1911)
labrofimbriata (de Saint Joseph, 1888)
latreilli Audouin & Milne-Edwards, 1834
longipodiata Cantone, 1989
mixochaeta Oug, 1998
nonatoi Ramos, 1976
pseudofragilis Amoureux, 1977
scopa Fauchald, 1974
- Ninoe**
armoricana Glémarec, 1968
nigripes Verrill, 1873
- Paraninoe**
brevipes (McIntosh, 1903)
fusca (Moore, 1911)
minuta (Théel, 1879)
- Scleotoma**
debilis (Grube, 1878)
emandibla subsp. *mabitii* (Ramos, 1976)
fragilis (O.F. Müller, 1776)
funchalensis (Kinberg, 1865)
impatiens (Claparède, 1868)
rovignensis (Fauvel, 1940)
tetraura (Schmarda, 1861)
- Family Magelonidae**
Magelona
allenii Wilson, 1958
equilamellae Harmelin, 1964
filiformis Wilson, 1959
johnstoni Fiege, Licher & Mackie, 2000
minuta Eliason, 1962
mirabilis (Johnston, 1845)
papillicornis Müller, 1858 (sensu Fauvel, 1927)
rosea Moore, 1907
wilsoni Glémarec, 1966
- Family Maldanidae**
Axiothella
catenata (Malmgren, 1865)
constricta (Claparède, 1870)
- Chirimia**
biceps (M. Sars, 1861)
- Clymenella**
cincta (de Saint Joseph, 1894)
torquata (Leidy, 1855)
- Clymenopsis**

- constricta* (Wesenberg-Lund, 1948)
- Clymenura**
- borealis* (Arwidsson, 1906)
 - clypeata* (de Saint Joseph, 1894)
 - johnstoni* (McIntosh, 1915)
 - lankesteri* (McIntosh, 1885)
 - leiopygos* (Grube, 1860)
 - polaris* (Théel, 1879)
 - tenuis* (Day, 1957)
 - tricirrata* (Bellan & Reys, 1967)
- Euclymene**
- affinis* (M. Sars in G.O. Sars, 1872)
 - collaris* (Claparède, 1870)
 - droebrachiensis* (M. Sars in G.O. Sars, 1872)
 - lindrothi* Eliason, 1962
 - lumbricoides* (Quatrefages, 1865)
 - oerstedi* (Claparède, 1863)
 - palermitana* (Grube, 1840)
 - robusta* (Arwidsson, 1906)
- Johnstonia**
- clymenoides* Quatrefages, 1865
- Isocirrus**
- planiceps* (M. Sars in G.O. Sars, 1872)
- Lumbryclymene**
- cylindricaudata* Sars, 1872
 - minor* Arwidsson, 1906
- Macroclymene**
- santandrensis* (Rioja, 1917)
- Maldane**
- arctica* Detinova, 1985
 - glebifex* Grube, 1860
 - malmgreni* McIntosh, 1885
 - sarsi* Malmgren, 1865
- Maldanella**
- harai* (Izuka, 1902)
- Metasychis**
- gotoi* (Izuka, 1902)
- Microclymene**
- acirrata* Arwidsson, 1906
 - tricirrata* Arwidsson 1906
- Micromaldane**
- ornithochaeta* Mesnil, 1897
- Nicomache**
- lumbricalis* (Fabricius, 1780)
 - maculata* Arwidsson, 1911
 - minor* Arwidsson, 1906
 - quadrispinata* Arwidsson, 1906
 - personata* Johnson, 1901
 - trispinata* Arwidsson, 1906
- Notoproctus**
- oculatus* Arwidsson, 1906
- Petaloprotus**
- borealis* Ardwisson, 1906
 - tenuis* (Théel, 1879)
 - terricola* Quatrefages, 1865
- Praxillella**
- affinis* (M. Sars, 1872)
 - challengeriae* (Rioja, 1931)
 - gracilis* (M. Sars, 1861)
 - lophoseta* (Orlandi, 1898)
 - praetermissa* (Malmgren, 1866)
- Praxillura**
- longissima* Arwidsson, 1906
 - quadrilobata* (M. Sars, 1856)
- Proclymene**
- muelleri* (Sars, 1856)
- Rhodine**
- gracilior* Tauber, 1879
 - loveni* Malmgren, 1865
- Family Myzostomidae**
- Myzostoma**
- alatum* von Graff, 1884
 - asteriae* Marenzeller, 1895
 - bucchichii* F. Wagner, 1887
 - carpenteri* (von Graff, 1884)
- cirriferum* Leuckart 1827
- giganteum* Nansen, 1885
- glabrum* Leuckart, 1827
- ingolfi* Jägersten, 1940
- parasiticum* Leuckart, 1827
- pluvinar* von Graff, 1884
- Protomysostomum*
- polynephritis* Fedotov, 1912
- Family Nephtyidae**
- Aglaophamus*
- agilis* (Langerhans, 1879)
 - malmgreni* (Théel, 1870)
 - rubella* (Michaelson, 1896)
- Inermonephys*
- inermis* (Ehlers, 1887)
- Micronephys*
- maryae* San Martin, 1982
 - minuta* (Théel, 1879)
 - neotena* (Noyes, 1980)
 - sphaerocirrata* (Wesenberg-Lund, 1949)
 - stammeri* (Augener, 1932)
- Nephtys*
- assimilis* Oersted, 1843
 - breogani* Laborda & Vieitez, 1985
 - caeca* (Fabricius, 1780)
 - ciliata* (O.F. Müller, 1776)
 - cirrosa* Ehlers, 1868
 - cirrosa* subsp. *longicornis* Jakubova, 1930
 - hombergii* Savigny, 1818
 - hystricis* McIntosh, 1900
 - incisa* Malmgren, 1865
 - incisa* subsp. *bilobata* Heinen, 1911
 - kersivalensis* McIntosh, 1908
 - longosetosa* Oersted, 1842
 - paradoxa* Malm, 1874
 - pente* Rainer, 1984
 - pulchra* Rainer, 1991
- Family Nereididae**
- Ceratocephale*
- loveni* Malmgren, 1867
- Ceratonereis*
- anoculata* Amoureux, 1982
 - brunnea* Langerhans, 1884
 - costae* (Grube, 1840)
 - hircinicola* (Eisig, 1870)
 - mirabilis* Kinberg, 1866
 - vittata* Langerhans, 1884
- Eunereis*
- elittoralis* (Eliason, 1962)
 - longissima* (Johnston, 1840)
- Leonnates*
- decipiens* Fauvel, 1929
 - jousseaumi* Gravier, 1899
 - persica* Wesenberg-Lund, 1949
 - pusillus* Langerhans, 1880
- Micronereis*
- variegata* Claparède, 1863
- Namalyctis*
- brevicornis* (Audouin & Milne-Edwards, 1834)
- Namanereis*
- quadraticeps* Gay in Blanchard, 1849
- Neanthes*
- agulhana* Day, 1963
 - bioculata* (Hartmann-Schröder, 1975)
 - caudata* (Delle Chiaje, 1828)
 - flavipes* Ehlers, 1868
 - fucata* (Savigny, 1818)
 - grandis* (Stimpson, 1853)
 - irrorata* (Malmgren, 1867)
 - kerguelensis* (McIntosh, 1885)
 - succinea* (Frey & Leuckart, 1847)
 - virens* (M. Sars, 1835)
 - willeyi* Day, 1934
- Nereis*
- diversicolor* (O.F. Müller, 1776)

- falcaria* (Willey, 1905)
fallax Quatrefages, 1865
falsa Quatrefages, 1865
funchalensis (Langerhans, 1879)
gilchristi Day, 1960
icosiensis Gravier & Dantan, 1928
jacksoni Kinberg, 1866
lamellosa Ehlers, 1868
moroccensis Amoureaux, 1976
pelagica Linnaeus, 1758
persica Fauvel, 1911
rava Ehlers, 1868
zonata Malmgren, 1867
- Perinereis**
cultifera (Grube, 1840)
macropus (Claparède, 1870)
marionii (Audouin & Milne- Edwards, 1833)
nuntia (Savigny, 1818)
oliveirae (Horst, 1889)
rullieri Pilato, 1974
taurica Langerhans, 1891
tenuisetis (Fauvel, 1915)
- Platynereis**
coccinea (Delle Chiaje, 1841)
dumerilii (Audouin & Milne-Edwards, 1833)
massiliensis (Moquin-Tandon, 1869)
nadiae Abbiati & Castelli, 1992
pulchella Gravier, 1899
- Pseudonereis**
anomala Gravier, 1899
- Rüllierinereis**
anoculata Cantone, 1982
faroensis Kirkegaard, 1998
tenericensis Nuñez, 1984
- Unanereis**
zghal Ben Amor, 1980
- Websterinereis**
glauca (Claparède, 1870)
- Family Nerillidae**
- Bathychaetus**
heptapous Faubel, 1978
- Meganerilla**
clavata Magagnini, 1966
swedmarki Boaden, 1961
- Mesonerilla**
armoricana Swedmark, 1959
biantennata Jouin, 1963
diatomeophaga Nuñez in Nuñez, Ocaña & Brito, 1997
fagei Swedmark, 1959
intermedia Wilke, 1953
roscovita Lévi, 1953
- Micronerilla**
minuta (Swedmark, 1959)
- Nerilla**
antennata O. Schmidt, 1848
marginata Tilzer, 1970
mediterraneum Schlieper, 1925
stygicola Ax, 1957
- Nerillidium**
gracile Remane, 1925
marinum Faubel, 1978
mediterraneum Remane, 1925
simplex Lévi, 1953
troglochaetoides Remane, 1925
- Nerillidopsis**
hyalina Jouin, 1966
- Paranerilla**
limicola Jouin & Swedmark, 1965
- Thalassochaetus**
palpifoliaceus Ax, 1954
- Family Oenonidae**
- Halla**
parthenopeia (Delle Chiaje, 1828)
sulfurea Langerhans, 1880
- Oenone**
- fulgida* (Savigny, 1818)
- Family Onuphidae**
- Aponuphis**
bilineata (Baird, 1870)
bilineata subsp. *grubii* (Marenzeller, 1886)
brementi (Fauvel, 1916)
ornata (Fauvel, 1928)
rigida (Claparède, 1868)
rubra (Langerhans, 1880)
willsii Cantone & Bellan, 1994
- Diopatra**
madeirensis Langerhans, 1880
neapolitana Delle Chiaje, 1841
- Epidiopatra**
huperiana subsp. *monroi* Day, 1957
- Hyalinoecia**
edwardsi Roule, 1906
robusta Southward, 1977
tubicola (O.F. Müller, 1776)
- Longibrachium**
atlanticum (Day, 1973)
- Nothria**
britannica (McIntosh, 1903)
conchylega (M. Sars, 1835)
geophiliformis Moore, 1903
- Nothrya**
hyperborea (Hansen, 1882)
- Onuphis**
eremita Audouin & Milne-Edwards, 1833
falesia Castelli, 1982
iberica Hartmann-Schröder, 1975
opalina (Verrill, 1873)
rullieriana (Amouroux, 1977)
- Parahyalinoecia**
apalpata Hartmann-Schröder, 1975
- Paradiopatra**
calliopae Arvanitidis & Kououras, 1997
fiordica (Fauchald, 1974)
hartmanae (Kirkegaard, 1980)
hispanica Amoureaux, 1972
lepta (Chamberlin, 1919)
quadricuspis (M. Sars in G.O. Sars, 1872)
- Rhamphobrachium**
agassizii Ehlers, 1887
brevibranchiatum (Ehlers, 1875)
- Family Opheliidae**
- Ammotrypanella**
arctica McIntosh, 1879
- Armandia**
cirrhosa Filippi, 1861
polyophtalma Kukenthal, 1887
- Euzonus**
flabelligerus (Ziegelmeier, 1955)
- Kesun**
gravieri (McIntosh, 1908)
- Ophelia**
amoureuxi Bellan & Costa, 1987
bicornis Savigny, 1818
borealis Quatrefages, 1866
celtica Amoureaux & Dauvin, 1981
laubieri Bellan & Costa, 1987
limacina (Rathke, 1843)
neglecta Schneider, 1887
rathkei McIntosh, 1908
roscoffensis Augener, 1910
translucens (Katzmann, 1973)
- Ophelina**
abranchiata Støp-Bowitz, 1948
acuminata Oersted, 1843
aulogaster (Rathke, 1843)
breviata (Pettibone, 1954)
cylindricaudata (Hansen, 1878)
cylindricaudata subsp. *minima* Hartmann-Schröder, 1974
helgolandica Augener, 1912

- minima* Hartmann-Schröder, 1974
modesta Støp-Bowitz, 1958
norvegica Støp-Bowitz, 1945
opistobranchia Wirén, 1901
Polyopthalmus
pictus (Dujardin, 1839)
Tachytrypane
jeffreysii McIntosh, 1879
Travisia
forbesii Johnston, 1840
- Family Orbiniidae**
- Leitoscoloplos*
kerguelensis (McIntosh, 1885) sensu Ramos 1976
mammosus Day, 1977
- Naineris*
laevigata (Grube, 1855)
quadraticeps Day, 1965
quadricuspida (Fabricius, 1780)
- Orbinia*
armandi (McIntosh, 1910)
borei (Fauvel, 1919)
cornidei (Rioja, 1934)
cuvieri (Audouin & Milne-Edwards, 1833)
latreillii (Audouin & Milne-Edwards, 1833)
sertulata (Savigny, 1820)
- Phylo*
foetida subsp. *adjimansis* (Fauvel, 1925)
foetida subsp. *atlantica* (Fauvel, 1925)
foetida subsp. *foetida* (Claparède, 1870)
foetida subsp. *imitans* (Eisig, 1914)
foetida subsp. *ligustica* (Orlandi, 1896)
foetida subsp. *typica* Eisig, 1914
grubei (McIntosh, 1910)
kupfferi (Ehlers, 1874)
norvegicus (Sars, 1872)
- Protoarcia*
capsulifera (Bobretzky, 1870)
oerstedi (Claparède, 1864)
- Schroederella*
laubieri Badalamenti & Castelli, 1991
- Scolaricia*
haasi Monro, 1937
typica Eisig, 1914
- Scoloplos*
armiger (O.F. Müller, 1776)
chevalieri (Fauvel, 1901)
chevalieri subsp. *candiensis* Harmelin, 1969
- Family Oweniidae**
- Myriochela*
danielsseni Hansen, 1879
fragilis Nilson & Holthe, 1985
heeri Malmgren, 1867
oculata (Zachs, 1923)
- Owenia*
fusiformis Delle Chiaje, 1842
- Family Paralacydoniidae**
- Paralacydonia*
paradoxa (Fauvel, 1913)
- Family Paraonidae**
- Aedicira*
mediterranea Laubier & Ramos, 1974
- Aricidea*
aberrans Laubier & Ramos, 1974
abbranchiata Hartman, 1965
abyssalis Laubier & Ramos, 1974
albatrossae Pettibone, 1957
anna Laubier, 1967
assimilis Tebble, 1959
capensis subsp. *bansei* Laubier & Ramos, 1974
catherinae Laubier, 1967
cerrutii Laubier, 1967
claudiae Laubier, 1967
fauveli Hartman, 1957
fragilis Webster, 1879
- fragilis* subsp. *mediterranea* Laubier & Ramos, 1974
hartmani Strelzov, 1968
laubieri Hartley, 1981
mariannae Katzmünn & Laubier, 1975
minuta Southward, 1956
monicae Laubier, 1967
neosuecica Hartman, 1965 sensu Laubier & Ramos, 1974
cf. philbinae Brown, 1976
pseudannae Katzmünn & Laubier, 1975
quadrilobata Webster & Benedict, 1887
roberti Hartley, 1983
simonae Laubier & Ramos, 1974
suecica Eliason, 1920
suecica subsp. *meridionalis* Laubier & Ramos, 1974
suecica subsp. *simplex* Day, 1963
trilobata Laubier & Ramos, 1975
wassi Pettibone, 1965
- Cirrophorus*
branchiatus Ehlers, 1908
furcatus (Hartman, 1957)
- Levinseria*
gracilis (Tauber, 1879)
- Paradoneis*
armata Glémarec, 1966
drachi Laubier & Ramos, 1974
eliasoni Mackie, 1991
fulgens (Levinsen, 1884)
ilvana Castelli, 1985
harpagonea (Storch, 1967)
lyra (Southern, 1914)
- Paraonides*
myriamae Katzmünn & Laubier, 1975
neapolitana Cerruti, 1909
- Paraonis*
uncinatus Hartman, 1965
- Family Parergodrilidae**
- Stylocapitella*
subterranea Knöllner, 1934
- Family Pectinariidae**
- Pectinaria*
auricoma (O.F. Müller, 1776)
auricoma subsp. *mediterranea* Nilsson, 1928
belgica (Pallas, 1766)
granulata (Linnaeus, 1767)
hyperborea (Malmgren, 1866)
koreni (Malmgren, 1866)
neapolitana Claparède, 1870
- Petta*
pusilla Malmgren, 1866
- Family Pholoidae**
- Pholoe*
assimilis Oersted, 1843
baltica Oersted, 1843
fauveli Kirkegaard, 1983
inornata Johnston, 1839
longa (O.F. Müller, 1776)
minuta (Fabricius, 1780)
pallida Chambers, 1985
synopthalmica Claparède, 1868
tecta Stimpson, 1854
- Pholoides*
dorsipapillatus (Marenzeller, 1893)
- Family Phyllodocidae**
- Carobia*
dohrnii Langerhans, 1880
lanceoligera Czerniavsky, 1882
- Chaetoparia*
nilssoni Malmgren, 1867
- Eteone*
arctica Malmgren, 1867
barbata (Malmgren, 1865)
flava (Fabricius, 1780)
foliosa Quatrefages, 1865

- fucata* M. Sars, 1872
longa (Fabricius, 1780)
picta Quatrefages, 1865
spetsbergensis Malmgren, 1865
suecica Bergström, 1914
syphonodonta (Delle Chiaje, 1822)
- Eulalia**
trilineata de Saint Joseph, 1888
- Eulalia**
parva de Saint Joseph, 1898
anoculata Hartman & Fauchald, 1971
aurea Gravier, 1896
bilineata (Johnston, 1840)
expusilla Pleijel, 1987
flavescens Bobretzki, 1868
fucescens de Saint Joseph, 1888
hanssoni Pleijel, 1987
impostii Quatrefages, 1865
microoculata Pleijel, 1987
mustela Pleijel, 1987
ornata de Saint Joseph, 1888
quadrilineata de Saint Joseph, 1898
tjalfsiensis Ditlevsen, 1917
tripunctata McIntosh, 1874
venusta de Saint Joseph, 1888
venustissima Banse, 1959
viridis (Linnaeus, 1767)
- Eumida**
ockelmanni Eibye-Jacobsen, 1987
arctica (Annenkova), 1946
bahusiensis Bergström, 1914
punctifera (Grube, 1860)
sanguinea (Oersted, 1843)
- Genetyllis**
nana (de Saint Joseph, 1908)
- Hesionura**
augeneri (Friedrich, 1937)
coineau (Laubier, 1962)
coineau subsp. *longissima* Minichev, 1982
elongata (Southern, 1914)
serrata (Hartmann-Schröder, 1960)
- Lugia**
pterophora (Ehlers, 1864)
- Mystides**
bidentata Langerhans, 1880
borealis Théel, 1879
caeca Langerhans, 1880
southerni Banse, 1954
- Nereiphylla**
lutea (Malmgren, 1865)
paretti Blainville, 1828
pusilla (Claparède, 1870)
rubiginosa (de Saint Joseph, 1888)
- Notophyllum**
americanum Verrill, 1885
foliosum (M. Sars, 1835)
- Paranaitis**
kosteriensis (Malmgren, 1867)
wahlbergi (Malmgren, 1865)
- Pareteone**
abyssalis Hartmann-Schröder, 1975
- Phyllodoce**
albovittata Grube, 1860
bruneoviridis de Saint Joseph, 1898
citrina Malmgren, 1865
gervillei Audouin & Milne-Edwards, 1834
groenlandica Oersted, 1842
lamelligera (Linnaeus, 1791)
laminosa Savigny, 1818
lineata (Claparède, 1870)
longifrons Ben-Elijah, 1972
longipes Kinberg, 1866
macropapillosa de Saint Joseph, 1895
macrophthalma Schmarda, 1861
maculata (Linnaeus, 1767)
- madeirensis* Langerhans, 1880
malmgreni Gravier, 1900
mucosa Oersted, 1843
rosea (McIntosh, 1877)
- Protomystides**
brunnea Hartmann-Schröder, 1963
exigua (Eliason, 1862)
- Pseudomystides**
limbata de Saint Joseph, 1888
spinachia Petersen & Pleijel, 1993
- Pterocirrus**
slastnikovi Annenkova, 1946
macroceros (Grube, 1860)
microcephala Claparède 1870
limbata (de Saint Joseph, 1888)
- Pterocirrus**
nidarosiensis Pleijel, 1987
- Sige**
fusigera Malmgren, 1865
oliveri Pleijel, 1990
- Family Pilargidae**
- Ancistargis**
hamata (Hartman, 1960)
- Ancistrosyllis**
cingulata (Korschelt, 1893)
fioronii Fiege & Böggemann
groenlandica McIntosh, 1879
- Glypohesione**
klatti Friedrich, 1950
- Litocorsa**
stremma Pearson, 1970
- Otopsis**
chardyi Katzmann, Laubier & Ramos, 1974
- Pilargis**
papillata Rasmussen, 1973
verrucosa de Saint Joseph, 1899
- Sigambra**
constricta (Southern, 1921)
parva (Day, 1963)
tentaculata (Treadwell, 1941)
- Synelmis**
albini (Langerhans, 1881)
dineti Katzmann, Laubier & Ramos, 1974
- Family Pisionidae**
- Pisone**
puzae Siewing, 1953
remota (Southern, 1914)
- Family Poecilochaetidae**
- Elicodasia**
mirabilis Laubier & Ramos, 1973
- Poecilochaetus**
fauchaldi Pilato & Cantone, 1976
fulgoris Claparède in Ehlers, 1875
serpens Allen, 1904
- Family Polygordiidae**
- Polygordius**
appendiculatus Fraipont, 1887
erythrophthalmus (Giard, 1880)
lacteus Schneider, 1868
neapolitanus Fraipont, 1884
schnederi Langerhans, 1881
triestinus Woltereck in Hemplemann, 1906
villoti Perrier, 1875
- Family Polynoidae**
- Acanthicolpis**
asperrima (M. Sars, 1861)
- Acholoe**
astericola (Delle Chiaje, 1841)
- Adyle**
assimilis (McIntosh, 1876)
- Alentia**
gelatinosa (M. Sars, 1835)
- Alentiana**
aurantiaca (Verrill, 1885)

- Almaniella*
setubalensis McIntosh, 1885
- Austrolaenilla*
mollis (Sars, 1863)
- Bathyeliasona*
abyssicola (Fauvel, 1913)
kirkegaardi (Uschakov, 1971)
- Bathyfauelia*
affinis Fauvel, 1914
- Bathynoe*
nodulosa Ditlevsen, 1917
- Bylgides*
acutisetis Loshann, 1981
annenkovae Pettibone, 1993
elegans (Théel, 1879)
groenlandicus (Malmgren, 1867)
promamme (Malmgren, 1867)
sarsi (Kinberg in Malmgren, 1865)
- Diplaconotum*
paucidendatum (Eliason, 1962)
- Enipo*
elisabethae McIntosh, 1900
kinbergi (Malmgren, 1865)
torelli Malmgren, 1865
- Eucranta*
villosa Malmgren, 1865
- Eunoë*
anoculata Hartmann-Schröder, 1975
hubrechti (McIntosh, 1900)
laetmogonensis Kirkegaard & Billett, 1980
mamilloba Czerniavsky, 1882
nodosa (M. Sars, 1861)
oerstedi Malmgren, 1865
senta (Moore, 1902)
- Gattyana*
admonseni (Malmgren, 1867)
cirrhosa (Pallas, 1766)
- Gesiella*
jameensis (Hartmann-Schröder, 1974)
- Gorgoniaplynœ*
caeciliae (Fauvel, 1913)
- Harmothoe*
rarispina (M. Sars, 1861)
abyssicola Bidenkap, 1894
alba (Malmgren, 1865)
antilopes McIntosh, 1876
areolata (Grube, 1860)
aspera (Hansen, 1879)
bellani Barnich & Fiege, 2000
borealis (Théel, 1879)
casabullicola Brito, Nuñez & Bacallado, 1991
clavigera M. Sars, 1863
evei Kirkegaard, 1980
extenuata (Grube, 1840)
fragilis Moore, 1910
fraserithomsoni McIntosh, 1897
gilchristi Day, 1960
globifera (G.O. Sars, 1873)
goreensis Augener, 1918
imbricata (Linnaeus, 1767)
imbricata subsp. *incerta* (Bobretzky, 1881)
impar (Johnston, 1839)
ingolfiana Ditlevsen, 1917
joubini Fauvel, 1913
lagiscoïdes subsp. *serrata* Day, 1963
longisetis (Grube, 1863)
micraspis Eliason, 1962
oculinarum (Storm, 1879)
pagenstecheri Michaelsen, 1896
pokoui Intes & Le Loeuff, 1975
reticulata (Claparedé, 1870)
spinifera (Ehlers, 1864)
tenuisetis (McIntosh, 1885)
villosa Malmgren, 1865
violacea (Storm, 1876)
- viridis* Loshann, 1981
- Hermationella*
truncata (Moore, 1902)
- Heteroplynœ*
nordgaardi Bidenkap, 1907
- Lagisca*
aequespina Langerhans, 1884
elisabethae McIntosh, 1900
floccosa (Savigny, 1818)
peracuta McIntosh, 1885
talismanni (Roule, 1898)
taurica (Czerniavsky, 1882)
- Lepidasthenia*
argus Hodgson, 1900
brunnea Day, 1960
elegans (Grube, 1840)
maculata Potts, 1910
- Lepidonotus*
brevicornis Quatrefages, 1865
carinulatus (Grube, 1870)
clava (Montagu, 1808)
squamatus (Linnaeus, 1758)
tenuisetosus (Gravier, 1902)
- Leucia*
nivea (M. Sars, 1863)
- Macellicephala*
incerta Fauvel, 1914
laubieri Reyss, 1971
longipalpa Uschakov, 1957
mirabilis (McIntosh, 1885)
violacea (Levinse, 1887)
- Malmgreniella*
andreopolis (McIntosh, 1876)
arenicolae (de Saint Joseph, 1888)
castanea McIntosh, 1876
darbouxi Pettibone, 1993
glabra (Malmgren, 1865)
liliacea Pettibone, 1993
ljungmani (Malmgren, 1867)
lunulata (Delle Chiaje, 1830)
marphysae (McIntosh, 1876)
mcintoshii (Tebble & Chambers, 1982)
- Melaenis*
loveni Malmgren, 1865
- Nectochaeta*
grimaldii Marenzeller, 1892
- Neolagisca*
drachi (Reyss, 1961)
- Neoplynœ*
paradoxa (Storm, 1888)
- Paradye*
crinoidicola (Potts, 1910)
- Polaruschakov*
polaris (Uschakov, 1976)
reyssi Pettibone, 1976
- Polynœ*
antillicola (Augener, 1906)
coeliaca Fauvel, 1913
laevis Audouin & Milne Edwards, 1832
longissima Blainville, 1828
scolopendrina Savigny, 1818
- Robertianella*
synopthalma McIntosh, 1885
- Sinantenna*
macrophthalma (Fauvel, 1913)
- Subadyte*
pellucida (Ehlers, 1864)
- Wilsoniella*
furcosexta (Loshann, 1981)
- Family Pomatodrilidae**
- Pomatodrilus*
fluvialis (Lastochkin, 1935)
- Family Pontodoridae**
- Enapteris*
euchaeta (Chun, 1888)

- Pontodora*
pelagica Greef, 1879
- Family Protodrilidae**
- Protodriloides*
chaetifer (Remane, 1926)
symbioticus (Giard, 1904)
- Protodrilus*
adhaerens Jägersten, 1952
affinis Jouin, 1968
albicans Jouin, 1970
brevis Jouin, 1970
ciliatus Jägersten, 1952
flavocapitatus (Ulanin, 1877)
gracilis Nordheim, 1989
hatscheki Pierantoni, 1908
helgolandicus von Nordheim, 1983
hypoleucus Armenante, 1903
leuckarti Hatschek, 1880
mirabilis Czerniavsky, 1881
oculifer Pierantoni, 1908
purpureus (Schneider, 1868)
robustus Jägersten, 1952
rubropharyngeus Jägersten, 1940
scheideri Langerhans, 1880
similis Jouin, 1970
spongoides Pierantoni, 1903
- Family Psammodrilidae**
- Psammodrilus*
balanoglossoides Swedmark, 1952
- Psammodriloides*
fauveli Swedmark, 1958
- Family Questidae**
- Questa*
caudicirra Hartman, 1966
- Family Sabellariidae**
- Lygdamis*
muratus (Allen, 1904)
- Phalacrostemma*
cidariophilum Marenzeller, 1895
elegans Fauvel, 1914
norvegicum Strömgren, 1971
- Sabellaria*
alcocki Gravier, 1906
alveolata (Linnaeus, 1767)
spinulosa Leuckart, 1849
taurica (Rathke, 1837)
- Family Sabellidae**
- Amphiglena*
mediterranea (Leydig, 1851)
- Augeneriella*
lagunari Gitay, 1970
- Bispira*
crassicornis (Sars, 1851) com. nov.
fabricii (Kroyer, 1858)
mariae Lo Bianco, 1893
viola (Grube, 1863)
volutacornis (Montagu, 1804)
- Branchiomma*
bahiusense Johansson, 1927
boholense (Grube, 1878)
bombyx (Dalyell, 1853)
inconspicua (M. Sars in G.O. Sars, 1872)
infarctum (Kroyer, 1856)
luctuosum (Grube, 1869)
lucullanum (Delle Chiaje, 1828)
moebii P. Knight-Jones, 1994
spongiarum P. Knight-Jones, 1994
- Chone*
acustica (Claparède, 1870)
arenicola Giangrande, 1992
collaris Langerhans, 1880
duneri Malmgren, 1867
fauveli McIntosh, 1916
filicaudata Southern, 1914
- infundibuliformis* Kroyer, 1856
longiseta Giangrande, 1992
longocirrata Sars, 1872
paucibranchiata (Kroyer, 1856)
- Demonax*
branchyona (Claparède, 1870)
cambrensis Knight-Jones & Walker, 1985
langerhansi (Knight-Jones, 1983)
saxicola (Grube, 1861)
tenuicollaris (Grube, 1870)
tommasi Giangrande, 1994
torulis Knight-Jones & Walker, 1985
- Desdemona*
ornata Banse, 1957
- Euchone*
analis (Kroyer, 1865)
capensis Day, 1961
elegans Verrill, 1873
papillosa (M. Sars, 1851)
rosea Langerhans, 1884
rubella Ehlers,
rubrocincta (M. Sars, 1861)
southerni subsp. *southerni* Banse, 1970
southerni subsp. *incisa* Banse, 1970
- Fabricia*
sabella (Ehrenberg, 1836)
stellaris subsp. *stellaris* (O.F. Müller, 1774)
stellaris subsp. *adriatica* (Banse, 1956)
stellaris subsp. *caspica* (Zenkevitch, 1922)
- Fabriciola*
baltica Friedrich, 1939
tonerella Banse, 1956
- Gorbunovia*
malmgreni (Hansen, 1878)
- Hypsicomus*
stichophthalmos (Grube, 1863)
- Jasmineira*
candela (Grube, 1863)
caudata Langerhans, 1880
elegans de Saint Joseph, 1894
oculata Langerhans, 1884
schaudinni Augener, 1925
- Laonome*
kroegeri Malmgren, 1865
salmacidis Claparède, 1870
- Manayunkia*
aestuarina (Bourne, 1883)
caspica Annenkova, 1929
- Megalomma*
linaresi (Rioja, 1917)
vesiculosum (Montagu, 1815)
vigilans (Claparède, 1870)
- Myxicola*
aesthetica (Claparède, 1870)
infundibulum (Renier, 1804)
sarsi Krøyer, 1856
steenstrupi Krøyer, 1856
- Notaulax*
phaeotaenia (Schmarda, 1861)
rectangulata Levinsen, 1883
- Novaesaria*
bilobata Martin & Giangrande, 1991
- Oriopsis*
alata subsp. *pectinata* Banse, 1957
armandi (Claparède, 1864)
cursoria (Quatrefages, 1865)
eimeri subsp. *eimeri* (Langerhans, 1880)
eimeri subsp. *persicina* Ben-Eliah, 1975
ingeloreae Plate, 1995
hynensis Knight-Jones, 1983
liefdefjordensis Plate, 1995
metchnikowi Caullery & Mesnil, 1896
- Panousea*
africana Amoureux & Rullier, 1974
- Perkinsiana*

- rubra* (Langerhans, 1880)
socialis (Langerhans, 1884)
- Potamethus*
breviuncatus Hartmann-Schröder, 1977
filiformis Hartmann-Schröder, 1977
malmgreni Hansen, 1878
spathiferus (Ehlers, 1887)
- Pothametus*
murrayi (McIntosh, 1916)
- Potamilla*
neglecta (M. Sars, 1851)
torelli Malmgren, 1866
rubra Rioja, 1908
- Pseudofabricia*
aberrans Cantone, 1970
- Pseudofabriciola*
analis Fitzhugh, Giangrande & Simboula, 1994
capensis (Monro, 1937)
longipyga Fitzhugh, Giangrande & Simboula, 1994
- Pseudopotamilla*
ceresina (Grube, 1870)
reniformis (Bruguière, 1789)
- Sabella*
discifera Grube, 1874
fabricii Fauvel, 1927
flabellata Savigny, 1820
pavonina Savigny, 1820
sarsi Kröyer, 1856
spallanzanii (Viviani, 1805)
variabilis Langerhans, 1884
- Family Saccocirridae**
- Saccocirrus*
major Pierantoni, 1907
papillocercus Bobretzky, 1872
parvus Gerlach, 1953
- Family Scalibregmatidae**
- Asclerocheilus*
intermedius (de Saint Joseph, 1894)
- Lipobranchius*
jeffreysi McIntosh, 1869
- Polyphysia*
crassa (Oersted, 1843)
crassa subsp. *fauveti* (Laubier, 1959)
- Pseudoscalibregma*
parvum (Hansen, 1878)
- Scalibregma*
celticum Mackie, 1991
inflatum Rathke, 1843
- Sclerobregma*
branchiata Hartman, 1965
- Sclerocheilus*
minutus Grube, 1863
- Family Serpulidae**
- Apomatus*
ampulliferus Philippi, 1844
similis Marion & Bobretzky, 1875
- Bathyvermilia*
langerhansi Fauvel, 1909
- Chiitnopoma*
serrula (Stimpson, 1854)
- Ditrupa*
arietina (O.F. Müller, 1776)
- Ficopomatus*
enigmaticus (Fauvel, 1923)
- Filograna*
implexa Berkeley, 1828
- Filogramula*
annulata (O.G. Costa, 1861)
calculata (O.G. Costa, 1861)
gracilis Langerhans, 1884
stellata (Southward, 1963)
- Hyalopomatus*
claparadii Marenzeller, 1878
marennzelleri (Langerhans, 1884)
- variorugosus* Ben-Eliahu & Fiege, 1996
- Hydroïdes*
- albiceps* Grube, 1870
azoricus Zibrowius, 1972
brachyacanthus Rioja, 1941
dianthus (Verrill, 1873)
dirampha Mörch, 1863
elegans (Haswell, 1883)
ezoenensis Okuda, 1934
helmatius (Iroso, 1921)
heterocerus (Grube, 1868)
homocerus Pixell, 1913
minax (Grube, 1878)
niger Zibrowius, 1971
norvegicus Gunnerus, 1768
novae-pomeraniae Augener, 1925
opercularis (Treadwell, 1929)
pseudouncinatus subsp. *africanus* Zibrowius, 1971
pseudouncinata subsp. *pseudouncinata* Zibrowius, 1968
steinitzi Ben-Eliahu, 1972
stoichadon Zibrowius, 1971
- Janita*
imbriata (Delle Chiaje, 1822)
spinosa (Langerhans, 1884)
- Josephella*
marennzelleri Caullery & Mesnil, 1896
- Metavermilia*
arctica Kuprianova, 1993
multicristata (Philippi, 1844)
taenia Zibrowius, 1971
- Neovermilia*
salicigera (Roule, 1906)
- Placostegus*
crystallinus sensu Zibrowius, 1968
langerhansi Marenzeller, 1893
tridentatus (Fabricius, 1779)
- Pomatoceros*
lamarckii (Quatrefages, 1865)
triqueter (Linnaeus, 1767)
- Pomatoleios*
kraussii (Baird, 1865)
- Protula*
alberti Fauvel, 1909
- Protis*
arctica (Hansen, 1878)
- Protula*
globifera (Théel, 1876)
intestinum (Savigny, 1818)
lusitanica Rioja, 1917
similis Marion & Bobretzky, 1875
tubularia (Montagu, 1803)
- Pseudovermilia*
occidentalis Zibrowius, 1969
- Rhodopsis*
pusilla Bush, 1905
- Salmacina*
dysteri (Huxley, 1855)
edificatrix Claparède, 1868
incrassata Claparède, 1868
setosa Langerhans, 1884
- Semivermilia*
agglutinata (Marenzeller, 1893)
crenata (O.G. Costa, 1861)
cibrata (O.G. Costa, 1861)
pomatostegoides (Zibrowius, 1969)
torulosa (Delle Chiaje, 1822)
- Serpula*
cavernicola Fassari & Mollica, 1991
concharum Langerhans, 1880
israelitica Amoureux, 1976
lobiancoi Rioja, 1917
vermicularis Linnaeus, 1767
- Spiraserpula*
massiliensis (Zibrowius, 1968)

<i>Spirodiscus</i>	<i>arcticus</i> (M. Sars, 1851)
<i>grimaldii</i> Fauvel, 1909	<i>citrinus</i> (Stimpson, 1854)
<i>Spirobranchus</i>	<i>Sphincter</i>
<i>lima</i> (Grube, 1862)	<i>oniscoides</i> Johnston, 1845
<i>polytrema</i> (Philippi, 1844)	<i>Wiréni</i> Hartman, 1948
<i>tetraceros</i> (Schmarda, 1861)	
<i>Vermiliopsis</i>	Family Spionidae
<i>eliasoni</i> Zibrowius, 1970	<i>Aonides</i>
<i>infundibulum</i> (Philippi, 1844)	<i>oxycephala</i> (Sars, 1862)
<i>labiata</i> (G.O. Costa, 1861)	<i>paucibranchiata</i> Southern, 1914
<i>monodiscus</i> Zibrowius, 1968	<i>Atherospio</i>
<i>striaticeps</i> (Grube, 1862)	<i>disticha</i> Mackie & Duff, 1986
<i>Vitreotubus</i>	<i>Apopronospio</i>
<i>digeronimoi</i> Zibrowius, 1979	<i>pygmaea</i> Hartman, 1955
Family Sigalionidae	<i>Aurospio</i>
<i>Euthalenessa</i>	<i>dibranchiata</i> Maciolek, 1981
<i>oculata</i> (Peters, 1854)	<i>Boccardia</i>
<i>Claparedepolgenia</i>	<i>polybranchia</i> (Haswell, 1885)
<i>inclusa</i> (Claparède, 1868)	<i>redeki</i> (Horst, 1920)
<i>Fimbriosthenelais</i>	<i>semibranchiata</i> Guerin, 1990
<i>minor</i> (Pruvot & Racovitza, 1895)	<i>Boccardiella</i>
<i>zelandica</i> (McIntosh, 1876)	<i>ligerica</i> (Ferronnière, 1898)
<i>Labioleanira</i>	<i>Dipolydora</i>
<i>yhleni</i> (Malmgren, 1867)	<i>armata</i> Langerhans, 1880
<i>Leanira</i>	<i>caulleryi</i> Mesnil, 1897
<i>hystricis</i> Ehlers, 1874	<i>coeca</i> (Oersted, 1843)
<i>Neoleanira</i>	<i>flava</i> Claparède, 1870
<i>tetragona</i> (Oersted, 1845)	<i>giardi</i> Mesnil, 1896
<i>Parasthenelais</i>	<i>langerhansis</i> Mesnil, 1896
<i>hibernica</i> (McIntosh, 1876)	<i>quadrilobata</i> Jacobi, 1883
<i>Pelogenia</i>	<i>rogeri</i> Martin, 1996
<i>arenosa</i> (Delle Chiaje, 1830)	<i>saintjosephi</i> Eliason, 1920
<i>Sigalion</i>	<i>socialis</i> (Schmarda, 1861)
<i>mathildae</i> Audouin & Milne Edwards in Cuvier, 1830	<i>Dispio</i>
<i>squamosus</i> (Delle Chiaje, 1830)	<i>uncinata</i> Hartman, 1951
<i>Sthenelais</i>	<i>Laonice</i>
<i>boa</i> (Johnston, 1833)	<i>appelloefi</i> (Söderström, 1920)
<i>ctenolepis</i> Claparède, 1868	<i>bahusiensis</i> (Söderström, 1920)
<i>jeffreysi</i> McIntosh, 1876	<i>cirrata</i> (M. Sars, 1851)
<i>limicola</i> (Ehlers, 1864)	<i>sarsi</i> Söderström, 1920
Family Sphaerodoridae	<i>Malacoceros</i>
<i>Clavodorum</i>	<i>ciliatus</i> (Keferstein, 1862)
<i>adriaticum</i> Katzmann, 1973	<i>fuliginosus</i> (Claparède, 1868)
<i>fauchaldi</i> Desbruyères, 1980	<i>girardii</i> Quatrefages, 1843
<i>Commensodorum</i>	<i>tetraceros</i> (Schmarda, 1861)
<i>commensalis</i> (Lützen, 1961)	<i>vulgaris</i> (Johnston, 1827)
<i>Ephesiella</i>	<i>Marenzelleria</i>
<i>abyssorum</i> (Hansen, 1878)	<i>viridis</i> (Verrill, 1873)
<i>cantonei</i> Mollica, 1994	<i>Wiréni</i> Augener, 1913
<i>peripatus</i> (Claparède, 1863)	<i>Microspio</i>
<i>ramosae</i> Desbruyères, 1980	<i>atlantica</i> (Langerhans, 1880)
<i>Euritmia</i>	<i>mecznikowianus</i> (Claparède, 1868)
<i>hamulifera</i> Sarda-Borroy, 1987	<i>Minuspio</i>
<i>Sphaerodoridium</i>	<i>cf. multibranchiata</i> (Berkeley, 1927)
<i>claparedii</i> (Grefft, 1866)	<i>Parapronospio</i>
<i>fauchaldi</i> Hartmann-Schröder, 1993	<i>pinnata</i> (Ehlers, 1901)
<i>longiparapodium</i> Katzmann, 1973	<i>Polydora</i>
<i>Sphaerodoropsis</i>	<i>ciliata</i> (Johnston, 1838)
<i>baltica</i> (Reimers, 1933)	<i>colonia</i> Moore, 1907
<i>disticum</i> (Eliason, 1962)	<i>cornuta</i> Bosc, 1802
<i>minuta</i> (Webster & Benedict, 1887)	<i>guillei</i> Laubier & Ramos, 1974
<i>chardyi</i> Desbruyères, 1980	<i>hermaphroditica</i> Hannerz, 1956
<i>laureci</i> Desbruyères, 1980	<i>hoplura</i> Claparède, 1870
<i>longipapillata</i> Desbruyères, 1980	<i>limicola</i> Annenkova, 1934
<i>martinae</i> Desbruyères, 1980	<i>posthamata</i> Jones, 1962
<i>parva</i> Ehlers, 1913	<i>pulchra</i> (Carazzi, 1895)
<i>philippi</i> (Fauvel, 1911)	<i>spongicola</i> E. & C. Berkeley, 1950
<i>sibuetae</i> Desbruyères, 1980	<i>Prionospio</i>
<i>flavum</i> Oersted, 1843	<i>banyulensis</i> Laubier, 1966
<i>Sphaerodorum</i>	<i>caspersi</i> Laubier, 1962
<i>gracilis</i> (Rathke, 1843)	<i>cirrifera</i> Wirén, 1883
<i>ophiurophoretos</i> Martin & Alva, 1988	<i>decipiens</i> Rioja, 1931
Family Spintheridae	<i>dubia</i> Day, 1961
<i>Spinther</i>	<i>ehlersi</i> Fauvel, 1928
	<i>fallax</i> Söderström, 1920
	<i>fauchaldi</i> Maciolek, 1985

- malmgreni* Claparède, 1870
multibranchiata Berkeley & Berkeley, 1927
plumosa (M. Sars, 1872)
saccifera Mackie & Hartley, 1990
salzi Laubier, 1970
sexoculata Augener, 1918
steenstrupi Malmgren, 1867
tripinnata Maciolek, 1985
- Pseudopolydora**
antennata (Claparède, 1868)
kempi (Southern, 1921)
paucibranchiata (Okuda, 1937)
pulchra (Carazzi, 1895)
- Pygospio**
elegans Claparède, 1863
- Scolelepis**
bonnieri (Mesnil, 1896)
cantabra (Rioja, 1918)
- Scolepis**
cirratulus (Delle Chiaje, 1828)
- Scolelepis**
foliosa (Audouin & Milne Edwards, 1833)
korsuni Sikorski, 1994
longirostris (Quatrefages, 1843)
mesnili (Bellan & Lagardère, 1971)
squamata (O.F. Müller, 1789)
tridentata (Southern, 1914)
- Spio**
armata (Thulin, 1957)
crenaticornis Montagu, 1813
decoratus Bobretzky, 1870
filicornis (O.F. Müller, 1776)
gonocephala Thulin, 1957
martinensis Mesnil, 1896
multioculata (Rioja, 1918)
- Spiophanes**
bombyx (Claparède, 1870)
kroyeri subsp. *kroyeri* Grube, 1860
kroyeri subsp. *reissi* Laubier, 1964
wigleyi Pettibone, 1962
- Streblospio**
benedicti Webster, 1979
dekhuyzeni Horst, 1909
shrubsoili (Buchanan, 1890)
- Family Spirorbidae**
- Bushiella**
atlantica (P. Knight-Jones, 1978)
evoluta (Bush, 1905)
kofsiadii (Rzhavsky, 1988)
verruca (O. Fabricius, 1780)
- Circeis**
armoricana (de Saint Joseph, 1894)
armoricana subsp. *fragilis* Knight-Jones & Knight-Jones, 1977
paguri Knight-Jones & Knight-Jones, 1977
spirillum (Linnaeus, 1758)
- Dexiospira**
corrugata (Montagu, 1803)
- Janua**
echinata (Wesenberg-Lund, 1953)
pagenstecheri (Quatrefages, 1865)
- Jugaria**
granulata (Linnaeus, 1767)
quadrangularis (Stimpson, 1854)
similis (Bush, 1905)
- Leodora**
laevis (Quatrefages, 1865)
- Neodexiospira**
brasiliensis (Grube, 1872)
pseudocorrugata (Bush, 1904)
- Neomicrorbis**
azoricus Zibrowius, 1972
- Nidificaria**
chilensis (Blanchard, 1849)
clavus (Harris, 1968)
- Paradexiospira**
cancelata (O. Fabricius, 1780)
violacea (Levinsen, 1884)
- Paralaeospira**
vitrea (Fabricius, 1780)
malardi (Caullery & Mesnil, 1897)
- Pileolaria**
berkeleyana (Rioja, 1942)
heteropoma (Zibrowius, 1968)
militaris (Claparède, 1868)
moerchi (Levinsen, 1883)
- Protolaeospira**
striata (Quievreux, 1963)
- Smilaria**
pseudomilitaris (Thiriot-Quievreux, 1965)
- Spirorbis**
corallinae de Silva & Knight-Jones, 1962
cuneatus Gee, 1964
gesae Knight-Jones & Knight-Jones, 1995
infundibulum Harris & Knight-Jones, 1964
inornatus L'Hardy et Quievreux, 1962
marioni Caullery & Mesnil, 1897
mentosus Bush, 1910
moerchi (Levinsen, 1883)
nudus Bush, 1910
ponicus Eichwald in Mörch, 1863
pusilla Rathke, 1837
rupestris Gee & Knight-Jones, 1962
serratus Bush, 1910
spirorbis (Linnaeus, 1758)
tridentatus Levinsen, 1883
- Vinearia**
endoumensis (Zibrowius, 1968)
koehleri (Caullery & Mesnil, 1897)
- Family Sternaspidae**
- Sternaspis**
fossor Stimpson, 1853
scutata (Ranzani, 1817)
- Family Syllidae**
- Amblyosyllis**
finmarchica (Malmgren, 1867)
dorsigera Claparède, 1864
formosa (Claparède, 1863)
granosa Sarda, 1983
inmaturata Langerhans, 1879
lineolata (A. Costa, 1864)
madeirensis Langerhans, 1879
- Autolitus**
syllisetosus Langerhans, 1884
- Autolytus**
alexandri Malmgren, 1867
benazzii Cognetti, 1953
brachycephalus (Marenzeller, 1874)
convolutus Cognetti, 1953
edwardsii de Saint Joseph, 1887
ebbiensis de Saint Joseph, 1887
groenlandicus Wesenberg-Lund, 1947
inermis de Saint Joseph, 1887
langheransi Gidholm, 1967
longiferiens de Saint Joseph, 1887
longoprimicirratus López, San Martin & Jiménez, 1997
lugens de Saint Joseph, 1887
megodon de Saint Joseph, 1887
neapolitanus Cognetti, 1953
paradoxus de Saint Joseph, 1887
pictus de Saint Joseph, 1887
prismaticus (O. Fabricius, 1780)
prolifer (O.F. Müller, 1788)
punctatus de Saint Joseph, 1887
quindecimdentatus Langerhans, 1884
roseus Claparède, 1864
rubrolineatus Gidholm, 1967
rubropunctatus (Grube, 1860)
rubrovittatus Claparède, 1864

- sardai* San Martin & Alos, 1987
smittiæ Malaquin, 1893
usaensis Imajima, 1966
verrilli Marenzeller, 1872
Branchiosyllis
exilis (Gravier, 1900)
Brania
arminii (Langerhans, 1881)
atokalis (Czerniavsky, 1881)
tenuicirrata (Claparède, 1864)
Calamyzae
amphictenoides Ardwisson, 1932
Dioplosyllis
cirrosa Gidholm, 1962
Ehlersia
cornuta (Rathke, 1843)
ferrugina Langerhans, 1881
Euryssyllis
tuberculata Ehlers, 1864
Eusyllis
assimilis Marenzeller, 1875
blomstrandii Malmgren, 1867
intermedia de Saint Joseph, 1887
lamelligera Marion & Bobretzky, 1875
monilicornis Malmgren, 1867
Exogone
acerata San Martin & Parapar, 1990
brevipes (Claparède, 1864)
cognetti Castelli, Badalamenti & Lardici, 1987
convoluta Campoy, 1982
dispar (Webster, 1879)
fauveti Cognetti, 1961
furcifera Eliason, 1962
gambiae Lanera, Sordino & San Martin, 1994
hebes (Webster & Benedict, 1884)
insdignis (Langerhans, 1879)
naidina Oersted, 1845
parahomoseta subsp. *mediterranea* San Martin, 1984
rostrata Naville, 1933
verugera (Claparède, 1868)
Fauvelia
martinensis Gravier, 1900
Grubeosyllis
clavata (Claparède, 1863)
alvaradoi (San Martin, 1984)
balani (Hartmann-Schröder, 1960)
celiae Parapar & San Martin, 1992
euritmica (Sarda, 1984)
limbata (Claparède, 1868)
neapolitana (Goodrich, 1930)
pusilla (Dujardin, 1839)
swedmarki (Gidholm, 1962)
vieitezi (San Martin, 1984)
yraidae (San Martin, 1984)
Haplosyllis
chamaeleon Laubier, 1960
spongicola (Grube, 1855)
spongicola subsp. *tentaculata* Marion, 1877
Miscellania
dentata Martin, Alos & Sarda, 1990
Myrianida
pinnigera (Montagu, 1808)
Odontosyllis
ctenostoma Claparède, 1868
dugesiana Claparède, 1864
fulgurans (Audouin & Milne-Edward, 1833)
gibba Claparède, 1863
polydonta de Saint Joseph, 1887
Opisthodontia
morena Langerhans, 1879
pterochaeta Southern, 1914
Opisthosyllis
brunnea Langerhans, 1879
viridis Langerhans, 1879
Palposyllis
- prosostoma* Hartman-Schroëder, 1977
prosostoma Hartmann-Schröder, 1977
Parapionosyllis
brevicirra Day, 1954
elegans (Pierantoni, 1903)
gestans (Pierantoni, 1903)
labronica Cognetti, 1965
minuta (Pierantoni, 1903)
papillosa (Pierantoni, 1903)
Paraprocerastea
crocantinae San Martin & Alos, 1989
Parasphaerosyllis
indica Monro, 1937
Paratyposyllis
peresi Laubier, 1968
Petitia
amphophthalma Siewing, 1955
Pionosyllis
serata Southern, 1914
anophthalma Capaccioni & San Martin, 1990
compacta Malmgren, 1867
dionisi Nuñez & San Martin, 1991
divaricata (Keferstein, 1862)
gorringensis Hartmann-Schröder, 1977
lamelligera de Saint Joseph, 1886
longocirrata de Saint Joseph, 1886
malmgreni McIntosh, 1869
morenoae San Martin, 1984
propeweismanni Dauvin & Lee, 1983
pulligera (Krohn, 1852)
serratisetosa López, San Martin & Jiménez, 1997
weismanni Langerhans, 1879
Plakosyllis
brevipes Hartmann-Schröder, 1956
Proceraea
aurantiaca Claparède, 1868
cornuta (Agassiz, 1884)
fasciata Langerhans, 1879
macrophthalmia (Marenzeller, 1874)
picta Ehlers, 1864
prismatica (O.F. Müller, 1776)
Procerastea
halleyiana Malaquin, 1893
nematodes Langerhans, 1884
perieri Gravier, 1900
pori Ben-Elijah, 1977
Pseudobranchia
subterranea Hartmann-Schröder, 1977
Pseudosyllides
balearica San Martin, 1984
Pseudosyllis
brevipennis Grube, 1863
Spermosyllis
torulosa Claparède, 1864
Sphaerosyllis
adelae San Martin, 1984
austriaca Banse, 1959
belizensis Russell, 1989
brevicirra Hartmann-Schröder, 1960
bulbosa Southern, 1914
campoyi San Martin, 1982
capensis Day, 1953
claparedei Ehlers, 1864
cryptica Ben-Elijah, 1977
erinaceus Claparède, 1863
giandoi Somaschini & San Martin, 1994
glandulata Perkins, 1981
gravinae Somaschini & San Martin, 1994
hystrix Claparède, 1863
longicauda Webster & Benedict, 1887
ovigera Langerhans, 1879
papillifera Naville, 1933
pirifera Claparède, 1868
piriferopsis Perkins, 1981
taylori Perkins, 1981

- tetralix* Eliason, 1920
thomasi San Martin, 1984
xarifae Hartmann-Schröder, 1960
- Streptosyllis**
arenae Webster & Benedict, 1884
bidentata Southern, 1914
templadis San Martin, 1984
varians Webster & Benedict, 1887
websteri Southern, 1914
- Syllides**
articulocirrata Gillandt, 1979
bansei Perkins, 1981
benedicti Banse, 1971
convoluta Webster & Benedict, 1884
edentatus Westheide, 1974
edentula Claparède, 1868
fulvus (Marion & Bobretzky, 1875)
japonicus Imajima, 1966
longocirrata Oersted, 1845
pseudoconvolutus San Martin, 1984
- Syllis**
armillaris (O.F. Müller, 1776)
atlantica Cognetti, 1960
brevipennis (Grube, 1963)
fasciata Malmgren, 1867
alternata Moore, 1908
amica Quatrefages, 1865
beneliahui (Campoy, 1982)
binocula Czerniavsky, 1881
bouvieri Gravier, 1900
caeca (Katzmann, 1973)
columbreensis Campoy, 1982
compacta Gravier, 1900
corallicola Verrill, 1900
corallicoloides Augener, 1922
crux Nuñez & San Martin, 1991
cucullata McIntosh, 1908
czerniavskyi (Czerniavsky, 1881) nov comb.
ferrani Alos & San Martin, 1987
garciai (Campoy, 1982)
gerlachi (Hartmann-Schröder, 1960)
gerundensis Alos & Campoy, 1981
goltonoovensis (Hartmann-Schröder, 1962)
gracilis Grube, 1860
hyalina Grube, 1863
incisa (O. Fabricius, 1780)
kabilica (Ben-Eliahu, 1977)
katzmanni (Katzmann, 1973)
krohnii Ehlers, 1864
luquei San Martin, 1984
lutea Hartmann-Schröder, 1960
mediterranea (Ben-Eliahu, 1977)
nepiotosa Caullery & Mesnil, 1916
nidarosiensis Bidenkap, 1907
nigrans Bobretzky, 1870
nigricirris Grube, 1863
nuchalis Hartmann-Schröder, 1960
oerstedi (Malmgren, 1867)
pectinans Haswell, 1920
pontica (Czerniavsky, 1881)
prolifera Krohn, 1852
pulvinata (Langerhans, 1881)
rosea (Langerhans, 1879)
schulzi Hartmann-Schröder, 1960
torquata Marion & Bobretzky, 1875
truncata subsp. *cryptica* (Ben-Eliahu, 1977)
variegata Grube, 1860
vittata (Grube, 1840)
vivipara Krohn, 1869
westheidei San Martin, 1984
zonata (Haswell, 1833)
- Trypanosyllis**
aeolis Langerhans, 1879
coeliaca Claparède, 1868
gigantea (Mcintosh, 1885)
- rosea* (Grube, 1863)
zebra (Grube, 1840)
Umbellisyllis
clavata (Langerhans, 1879)
fasciata M. Sars, 1869
Xenosyllis
scabra (Ehlers, 1864)
- Family Terebellidae**
- Amaeana*
trilobata (M. Sars, 1863)
- Amphitrite**
affinis Malmgren, 1866
alcicornis Fauvel, 1909
cirrata (O.F. Müller, 1771 in 1776)
edwardsi (Quatrefages, 1865)
orotavae Langerhans, 1881
praecox de Saint Joseph, 1899
rubra (Risso, 1828)
variabilis (Risso, 1826)
- Amphitritides**
gracilis (Grube, 1860)
kuehlmanni Arvanitidis & Koukouras, 1994
- Artacama*
proboscidea Malmgren, 1866
- Axonice*
flexuosa (Grube, 1860)
maculata (Dalyell, 1853)
mirabilis (McIntosh, 1885)
- Baffinia*
hesslei (Annenkova, 1924)
- Ehlersiella*
hirsuta Roule, 1896
- Eupistella*
dibranchiata (Fauvel, 1909)
- Eupolymnia*
nebulosa (Montagu, 1818)
nesidensis (Delle Chiaje, 1828)
- Euthelphus*
setubalensis McIntosh, 1885
- Hauchellia*
tribulata (McIntosh, 1869)
- Lanassa*
nordenskioeldi Malmgren, 1866
venusta (Malm, 1874)
- Lanice*
conchilega (Pallas, 1766)
- Laphania*
boecki Malmgren, 1866
- Leanea*
abranchiata Malmgren, 1866
ebranchiata (M. Sars, 1865)
- Loimia*
medusa (Savigny, 1818)
- Lysilla*
loveni Malmgren, 1866
nivea Langerhans, 1884
- Neoamphitrite*
affinis (Malmgren, 1866)
edwardsi (Quatrefages, 1865)
figulus (Dalyell, 1853)
grayi (Malmgren, 1866)
groenlandica Malmgren, 1866
- Nicolea*
venustula (Montagu, 1818)
zostericola Oersted, 1844
- Octobranchus*
floriceps Kingston & Mackie, 1980
- Paramphitrite*
tetrabranchia Holthe, 1976
- Parathelphus*
collaris (Southern, 1914)
- Phisidia*
aurea Southward, 1956
oculata (Langerhans, 1881)
- Pista*

- cretacea* (Grube, 1860)
cristata (O.F. Müller, 1776)
maculata (Dalyell, 1853)
malmgreni Saphronova, 1997
mediterranea de Gaillande, 1970
mirabilis (McIntosh, 1885)
unibranchia Day, 1963
- Pistella**
lornensis (Pearson, 1969)
- Polycirrus**
arcticus M. Sars, 1825
arenivorus Caulery, 1916
aurantiacus Grube, 1860
calendrum Claparède, 1868
denticulatus de Saint Joseph, 1894
haematodes (Claparède, 1864)
jubatus Bobretzky in Annenkova, 1924
latidens Eliason, 1962
medusa Grube, 1850
norvegicus (Wollebaek, 1912)
pallidus (Claparède, 1864)
plumosus Wollebaeck, 1912
tenuisetis Langerhans, 1880
- Proclea**
grafii (Langerhans, 1884)
malmgreni (Ssolowiew, 1899)
- Streblosoma**
bairdi (Malmgren, 1865)
hesslei Day, 1955
intestinale M. Sars, 1872
- Terebella**
ehrenbergi Grube, 1870
lapidaria Linnaeus, 1767
- Thelepus**
cincinnatus (Fabricius, 1780)
setosus (Quatrefages, 1865)
triserialis (Grube, 1855)
- Family Tomopteridae**
- Tomopteris**
apsteini (Rosa, 1908)
biancoi Terio, 1947
cavallii Rosa, 1907
duccii Rosa, 1907
dunckeri Rosa, 1907
elegans Chun, 1888
eschscholtzi Greef, 1879
espansa Caroli, 1932
helgolandica Greef, 1879
kefersteini Greeff, 1879
krampi Stop-Bowitz, 1948
levipes Greeff, 1879
ligulata Rosa, 1908
marginata Caroli, 1932
miriglandulata Terio, 1947
nationalis Apstein, 1900
nisseni Rosa, 1908
partenopea Caroli, 1928
pediculosa Terio, 1947
pierantonii Terio, 1947
planktonis Apstein, 1900
poliglandulata Caroli, 1932
septentrionalis Steenstrup, 1849
tenuis Caroli, 1932
- Family Trichobranchidae**
- Octobranchus**
floriceps Kingston & Mackie, 1980
lingulatus (Grube, 1863)
- Terebellides**
carnea Bobretsky, 1881
stroemii M. Sars, 1835
- Trichobranchus**
glacialis Malmgren, 1865
roseus (Malm, 1874)
- Family Trochochaetidae**

- Trochochaeta**
multisetosa (Oersted, 1844)
watsoni (Fauvel, 1916)
- Family Typhloscolecidae**
- Sagitella**
kowalevskyi Wagner, 1872
- Travisopsis**
lanceolata Southern, 1910
levinseni Southern, 1910
lobifera Levinsen, 1885
- Typhloscolex**
muelleri Busch, 1851
phyloides Reibich, 1895

Oligochaeta

Compiled by C. Erséus and B. Healy

This list was compiled by Brenda Healy (family Enchytraeidae) and C. Erséus (all other families).

Naididae is primarily a freshwater group, but the species included here occur sometimes or exclusively in brackish water. The most comprehensive and still useful review of naidids is Sperber's (1948) monograph, but for the partly marine genus *Paranais*, Brinkhurst and Coates (1985) should also be considered.

The marine members of the Tubificidae have been subject to extensive systematic studies in recent years, in particular by C. Erséus, but since the publication of the key to marine and estuarine European species by Brinkhurst (1982), there is no compiled information other than that to be gathered from the primary taxonomical literature (e.g. Brinkhurst 1985, 1986; Erséus 1982, 1983a-b, 1985, 1987a-b and 1992). Some tubificids are restricted to brackishwater, but the majority of species are indeed marine, some species occurring at bathyal or even abyssal depths.

The single megascolecid included, *Pontodrilus bermudensis*, known from the Mediterranean Sea, is an earthworm (Easton 1984). It is widely distributed in marine beaches throughout the tropics and subtropics of the world, and one of the few, if not the only, earthworm in marine habitats.

The revision and taxonomy of European Enchytraeidae by Nielsen & Christensen (1959, 1961, 1963) provides the basis for the present list of this family, but over 30 European species have been subsequently added and more species await description. Taxonomic difficulties, and the lack of a modern guide to European species, mean that enchytraeids continue to be neglected and are almost never identified, even to genus, in surveys of marine fauna. Faunistic studies have been concentrated in principally the western Baltic, western Scandinavia and continental North Sea coasts, Iceland, Ireland, Wales, and the Atlantic coast of France. There are some records from Svalbard, Barents Sea, Belye More (White Sea), the eastern Baltic, and Scotland, but few from the coasts of the Iberian Peninsula or the Mediterranean.

Marine enchytraeids are mainly associated with habitats in the upper intertidal, and only the genus *Grania* is widespread in offshore benthos. Most records are from intertidal deposits, and tidal debris around high water mark where there is some overlap with the terrestrial fauna. The most commonly reported terrestrial species in saltmarshes and tidal debris are included in the list (six spp.). The lower salinity limit for most brackishwater

species appears to be about 3 ppt, i.e. α -oligohaline according to the Venice System of classification. In the Baltic, some brackishwater enchytraeids have been found to coexist with freshwater and terrestrial species at lower salinities (0.5-3 ppt). However, the salinity of oligohaline waters is rarely recorded by oligochaete specialists and the range of tolerance of the species is unknown. For this reason, four frequently recorded "freshwater" species taken with brackishwater species are included in the list.

Many oligochaetes of the Baltic Sea are included in a recently published guide to the Estonian annelids (Timm 1999), including an illustrated key.

References

- Baker, H. R. & Brinkhurst, R. O. 1981 A revision of the genus *Monopylephorus* and redefinition of the subfamilies *Rhyacodrilinae* and *Branchiurinae* (Tubificidae: Oligochaeta). *Canadian Journal of Zoology* 59: 939-965.
- Brinkhurst, R. O. 1962 A check-list of the British Oligochaeta. *Proceedings of the zoological Society of London* 130: 317-330.
- Brinkhurst, R. O. 1982. British and other marine and estuarine oligochaetes. Keys and notes for the identification of the species. *Synopses of the British Fauna* (New Series). 21. Cambridge: Cambridge University Press. 127 pp.
- Brinkhurst, R. O. 1985. A further contribution to the taxonomy of the genus *Tubificoides* Lastockin (Oligochaeta: Tubificidae). *Canadian Journal of Zoology* 63: 400-410.
- Brinkhurst, R. O. 1986. Taxonomy of the genus *Tubificoides* Lastockin (Oligochaeta: Tubificidae): species with bifid setae. *Canadian Journal of Zoology* 64: 1270-1279.
- Christensen, B. 1962 Oligochaeta 3. Enchytraeidae. *Zoology of Iceland* 2: 1-11.
- Easton, E. G. 1984 Earthworms (Oligochaeta) from islands of the south-western Pacific, and a note on two species from Papua New Guinea. *New Zealand Journal of Zoology* 11: 111-128.
- Erséus, C. 1976a Marine subtidal Tubificidae and Enchytraeidae (Oligochaeta) of the Bergen area, western Norway. *Sarsia* 62: 25-48.
- Erséus, C. 1976b Littoral Oligochaeta (Annelida) from Eyjafjördur, north coast of Iceland. *Zoologica Scripta* 5: 5-11.
- Erséus, C. 1977 Marine Oligochaeta from the Koster area, west coast of Sweden, with descriptions of two new enchytraeid species. *Zoologica Scripta* 6: 293-298.
- Erséus, C. 1979a Taxonomic revision of the marine genera *Bathydrilus* Cook and *Macroseta* Erséus (Oligochaeta, Tubificidae), with descriptions of six new species and subspecies. *Zoologica Scripta* 8: 139-151.
- Erséus, C. 1979b Taxonomic revision of the marine genus *Phallodrilus* Pierantoni (Oligochaeta, Tubificidae), with descriptions of thirteen new species. *Zoologica Scripta* 8: 187-208.
- Erséus, C. 1980a New species of *Phallodrilus* (Oligochaeta, Tubificidae) from the Arctic deep sea and Norwegian fjords. *Sarsia* 65: 57-60.
- Erséus, C. 1980b Taxonomic studies on the marine genera *Aktedrilus* Knöllner and *Bacescuella* Hrabe (Oligochaeta, Tubificidae), with descriptions of seven new species. *Zoologica Scripta* 9: 97-111.
- Erséus, C. 1982 Taxonomic revision of the marine genus *Limnodriloides* (Oligochaeta: Tubificidae). *Verhandlungen des naturwissenschaftlichen Vereins in Hamburg* (Neue Folge) 25: 207-277.
- Erséus, C. 1983a Deep-sea *Phallodrilus* and *Bathydrilus* (Oligochaeta, Tubificidae) from the Atlantic Ocean, with descriptions of ten new species. *Cahiers de Biologie Marine* 24: 125-146.
- Erséus, C. 1983b Three new species of the marine genus *Coralliodrilus* (Oligochaeta, Tubificidae) from Italy. *Bollettino di Zoologia* 49 (1982): 241-247.
- Erséus, C. 1985 Distribution and biogeography of Oligochaeta. In *Peuplements profonds du golfe de Gascogne* (eds. L. Laubier & C. Monniot): 365-367. Brest: Institut Français de Recherche pour l'Exploitation de la Mer.
- Erséus, C. 1987a Seven new marine species of *Phallodrilus* (Oligochaeta; Tubificidae) from various parts of Europe, and a re-examination of the type species *P. parthenopaeus* Pierantoni. *Journal of Natural History* 21: 915-931.
- Erséus, C. 1987b Marine Limnodriloidinae (Oligochaeta, Tubificidae) from Italy, with descriptions of two new species. *Bollettino di Zoologia* 54: 159-164.
- Erséus, C. 1992 A generic revision of the Phallodrilinae (Oligochaeta, Tubificidae). *Zoologica Scripta* 21: 5-48.
- Erséus, C., Grimm, R., Healy, B., Lundberg, S., Rota, E. & Timm, T. 1999 A survey of Clitellata in Nationalstadsparken, an urban national park in Stockholm, Sweden. *Hydrobiologia* 406: 101-110.
- Erséus, C. & Lasserre, P. 1976 Taxonomic status and geographic variation of the marine enchytraeid genus *Grania* Southern (Oligochaeta). *Zoologica Scripta* 5: 121-132.
- Finogenova, N.P. & Timm, T. 1988 Enchytraeids (Oligochaeta, Enchytraeidae) of the littoral of the White Sea. In *Explorations of the fauna of the seas 38(46); benthic ecosystems of the south eastern part of the Kandalaksha Bay and adjacent waters of the White Sea*. Moscow: USSR Academy of Sciences, pp. 91-109.
- Giere, O. 1976 Zur Kenntnis der litoralen Oligochaetenfauna Südfinnlands. *Annales Zoologici Fennici* 13: 156-160.
- Healy, B. 1979 Marine fauna of County Wexford. -I. Littoral and brackishwater Oligochaeta. *Irish Naturalists' Journal* 19: 418-422.
- Healy, B. 1979 Records of Enchytraeidae (Oligochaeta) in Ireland. *Journal of Life Sciences of the Royal Dublin Society* 1: 39-7.
- Kasprzak, K. 1986 *Skapospzczyt wodne i glebowe, II. Rodzina: Wazonkowce (Enchytraeidae)*. Warszawa: Państwowe wydawnictwo naukowe. 366 pp.
- Knöllner, F.H. 1935 Ökologische und systematische Untersuchungen über litorale und marine Oligochäten der Kieler Bucht. *Zoologische Jahrbücher* (Abteilung Systematik.) 66: 425-563.
- Kossmagk-Stephan, K.-J. 1983. Marine Oligochaeta of a sandy beach of the island of Sylt (North Sea) with description of four new enchytraeid species. *Mikrofauna des Meeresbodens* 89: 1-28.
- Lasserre, P. 1966 Oligochètes marins des côtes de France. I. Bassin d'Arcachon: systématique. *Cahiers de Biologie marine* 7: 295-317.
- Lasserre, P. 1967 Oligochètes marins des côtes de France. II. Roscoff, Penpoul, étangs saumâtres de Concarneau: systématique, écologie. *Cahiers de Biologie marine* 8: 273-293.
- Nielsen, C.O. & Christensen, B. 1959 The Enchytraeidae. Critical revision and taxonomy of European species. *Natura Jutlandica* 8-9: 1-160.
- Nielsen, C.O. & Christensen, B. 1961 The Enchytraeidae. Critical revision and taxonomy of European species. Supplement 1. *Natura Jutlandica* 10: 1-23.
- Nielsen, C.O. & Christensen, B. 1963 The Enchytraeidae. Critical revision and taxonomy of European species. Supplement 2. *Natura Jutlandica* 10: 1-19.
- Nurminen, M. 1965 Enchytraeid and lumbricid records (Oligochaeta) from Spitzbergen. *Annales Zoologici Fennici* 2: 1-10.
- Nurminen, M. 1966 Further notes on the enchytraeids (Oligochaeta) from Spitzbergen. *Annales Zoologici Fennici* 3: 68-69.
- Nurminen, M. 1967 Faunistic notes on north European enchytraeids. *Annales Zoologici Fennici* 4: 567-587.
- Nurminen, M. 1973 Enchytraeidae (Oligochaeta) of Iceland. *Annales Zoologici Fennici* 10: 412-413.
- Southern, R. 1909 Contributions towards a monograph of the British and Irish Oligochaeta. *Proceedings of the Royal Irish Academy* 27B: 119-182.

- Southern, R. 1913 Clare Island Survey. Part 48. Oligochaeta. *Proceedings of the Royal Irish Academy* 31B: 1-14.
- Timm, T. 1999 *Eesti rongusside (Annelida) määraja. A guide to the Estonian Annelida*. Tartu-Tallinn: Teaduste Akadeemia Kirjastus/Estonian Academy Publishers. 208 pp.
- Tynen, M.J. 1972 The littoral Enchytraeidae (Oligochaeta) of Anglesey and the Menai Strait with notes on habitats. *Journal of Natural History* 6: 21-29.
- Tynen, M.J. & Nurminen, M. 1969 A key to the European littoral Enchytraeidae (Oligochaeta). *Annales Zoologici Fennici* 6: 150-155.
- Stephenson, J. 1911 On some littoral Oligochaeta of the Clyde. *Transactions of the Royal Society of Edinburgh* 48: 31-65.
- von Bülow, T. 1957 Systematisch-autökologische Studien an eulitoralen Oligochaeten der Kimbrischen Halbinsel. *Kieler Meeresforschung* 13: 69-116.
- Class Oligochaeta**
- Family Naididae**
- Amphichaeta*
sannio Kallstenius, 1892
- Chaetogaster*
diaphanus (Gruithuisen, 1828)
langi Bretscher, 1896
- Nais*
communis Piguet, 1906
elinguis Müller, 1773
variabilis Piguet, 1906
- Paranais*
botniensis Sperber, 1948
frixi Hrabe, 1941
litoralis Müller, 1773
- Stylaria*
lacustris (Linnaeus, 1767)
- Uncinaria*
uncinata (Oersted, 1842)
- Family Tubificidae**
- Subfamily Limnodriloidinae**
- Thalassodrilides*
gurwitschi (Hrabe, 1971)
- Tectidrilus*
pranzoi Erséus, 1987
- Limnodriloides*
agnes Hrabe, 1967
appendiculatus Pierantoni, 1903
hrabetovae Erséus, 1987
maslinicensis (Hrabe, 1971)
monothecus Cook, 1974
pierantoni (Hrabe, 1971)
scandinavicus Erséus, 1982
- Subfamily Rhycodrilinae**
- Heterodrilus*
arenicolus Pierantoni, 1902
subtilis (Pierantoni, 1919)
- Monopylephorus*
irroratus (Verrill, 1873)
parvus Ditlevsen, 1904
rubroniveus Levinsen, 1884
- Subfamily Phallodrilinae**
- Phallodrilus*
parthenopaeus Pierantoni, 1902
- Abyssidrilus*
profundus (Cook, 1970)
- Paraktedrilus*
bakeri (Kossmagk-Stephan & Erséus, 1985)
- Aktedrilus*
arcticus (Erséus, 1978)
cuneus Erséus, 1984
curvipennis Erséus, 1980
magnus Erséus, 1980
mediterraneus (Erséus, 1980)
monospermathecus Knöllner, 1935
parvithecatus (Erséus, 1978)
ponicus (Hrabe, 1973)
- sardus* Erséus, 1987
sphaeropenis Erséus & Kossmagk-Stephan, 1982
svetlovi Finogenova, 1976
- Inermidrilus*
georgei (Erséus, 1987)
- Gianius*
cristolatus (Erséus, 1983)
densespectinis (Erséus, 1987)
monnioti (Erséus, 1987)
- Mexidrilus*
minutissimus (Erséus, 1987)
nidarosiensis (Erséus, 1987)
postspermaticatus (Erséus, 1980)
- Adelodrilus*
borcei (Hrabe, 1973)
cooki Erséus, 1978
kiselevi (Finogenova, 1972)
pusillus Erséus, 1978
- Atlantidrilus*
biparis (Erséus, 1983)
quadrisetis Erséus, 1982
styloatriatus (Erséus, 1979)
vulnus (Erséus, 1983)
- Thalassodrilus*
firmus (Erséus, 1979)
klarae Erséus, 1987
prostatis (Knöllner, 1935)
- Pirodrilus*
messanensis (Erséus, 1987)
minutus (Hrabe, 1973)
pinguis (Erséus, 1983)
- Spiridion*
insigne Knöllner, 1935
modricensis (Hrabe, 1973)
- Pectinodrilus*
duplex (Erséus, 1987)
rectisetosus (Erséus, 1979)
- Coralliodrilus*
giacobbei Erséus, 1983
statutus Erséus, 1983
tyndarensis Erséus, 1983
- Bathydrilus*
adriaticus (Hrabe, 1971)
atlanticus Erséus, 1979
desbruyeresi Erséus, 1983
graciliatriatus Erséus, 1979
medius Erséus, 1983
rarisetis (Erséus, 1975)
sandersi Erséus, 1983
- Inanidrilus*
bonomii Erséus, 1984
- Olavius*
algarvensis Giere, Erséus & Stuhlmacher, 1998
- Subfamily Tubificinae**
- Tubificoides*
amplivasatus (Erséus, 1975)
benedii (d'Udekem, 1855)
brownae Brinkhurst & Baker, 1979
crenacoleus Baker, 1983
diazi Brinkhurst & Baker, 1979
euxinicu (Hrabe, 1966)
galiciensis Martinez-Ansemil & Giani, 1987
heterochaetus (Michaelsen, 1926)
insularis (Stephenson, 1922)
intermedius (Cook, 1969)
kozloffi Baker, 1983
maritimus (Hrabe, 1973)
pseudogaster (Dahl, 1960)
scoticus Brinkhurst, 1985
swirencoides Brinkhurst, 1985
swirenkowi Jaroschencko, 1937
vestibulatus Erséus & Bonomi, 1987
- Clitellio*
arenarius (Müller, 1776)
- Tubifex*

<i>nerthus</i> Michaelsen, 1908	<i>lineatus</i> Müller, 1774
<i>tubifex</i> (Müller, 1774))	<i>macrothecatus</i> Erséus, 1976
<i>Limnodrilus</i>	<i>minutus</i> Müller, 1776
<i>hoffmeisteri</i> Claparède, 1862	<i>murmanicus</i> Finogenova & Strelsov, 1978
<i>Christierius</i>	<i>musciculus</i> (Stephenson, 1924)
<i>litoralis</i> (Erséus, 1976)	<i>nigrofasciatus</i> Nurminen, 1965
<i>Heterochaeta</i>	<i>niger</i> Southern, 1909
<i>costata</i> Claparède, 1863	<i>pagenstecheri</i> (Ratzel, 1869)
<i>Isochaetides</i>	<i>profugus</i> (Eisen, 1878)
<i>michaelseni</i> (Lastockin, 1937)	<i>pseudominutus</i> Timm, 1988
<i>Potamothrix</i>	<i>pumilio</i> Stephenson, 1932
<i>alatus</i> Finogenova, 1972	<i>reynoldsoni</i> Backlund, 1948
<i>bavaricus</i> (Oschmann, 1913)	<i>rivalis</i> Levinsen, 1883
<i>caspicus</i> (Lastockin, 1937)	<i>rubidus</i> Finogenova & Strelsov, 1978
<i>danubialis</i> (Hrabe, 1941)	<i>scoticus</i> Elmhirst & Stephenson, 1926
<i>hammoniensis</i> (Michaelsen, 1901)	<i>semifuscus</i> Stephenson, 1911
<i>heuscheri</i> (Bretschler, 1900)	<i>tuba</i> Stephenson, 1911
<i>moladviensis</i> Vejdovský & Mrázek, 1902	<i>viridis</i> Stephenson, 1911
<i>mrazeki</i> (Hrabe, 1941)	<i>westhei</i> Kossmagk-Stephan, 1983
<i>vejdovskyi</i> (Hrabe, 1941)	<i>Marionina</i>
<i>Psammoryctides</i>	<i>achaeta</i> Lasserre, 1964
<i>deserticola</i> (Grimm, 1877)	<i>appendiculata</i> Nielsen & Christensen, 1959
<i>lastoschkini</i> (Jaroschenko, 1948)	<i>argentea</i> (Michaelsen, 1889)
Family Megascolecidae	<i>elongata</i> Lasserre, 1966
<i>Pontodrilus</i>	<i>glandulifera</i> (Jansson, 1960)
<i>litoralis</i> Grube, 1855	<i>graelei</i> Kossmagk-Stephan, 1983
Family Enchytraeidae	<i>istriæ</i> Giere, 1974
<i>Achaeta</i>	<i>macrathi</i> Healy, 1996
<i>littoralis</i> Lasserre, 1968	<i>mesopsamma</i> Lasserre, 1966
<i>Cernosvitoviella</i>	<i>oligosetosa</i> Kossmagk-Stephan, 1983
<i>atrata</i> Bretscher, 1903	<i>preclitellochaeta</i> Nielsen & Christensen, 1963
<i>immodesta</i> (Knöllner, 1935)	<i>riparia</i> Bretscher, 1899
<i>microtheca</i> Rota & Healy, 1999	<i>sjaelandica</i> Nielsen & Christensen, 1961
<i>minor</i> Dósza-Farkas, 1990	<i>southerni</i> (Cernosvitov, 1937)
<i>Enchytraeus</i>	<i>spicula</i> (Leuckart, 1847)
<i>albidus</i> Henle, 1837	<i>sublitoralis</i> Erséus, 1976
<i>buchholzi</i> Vejdovsky, 1879	<i>subterranea</i> (Knöllner, 1935)
<i>capitatus</i> Bülow, 1957	<i>ulstrupae</i> Healy, 1996
<i>christensenii</i> Dósza-Farkas, 1992	<i>weilli</i> Lasserre, 1966
<i>minutus</i> Nielsen & Christensen, 1961	
<i>lieudeensis</i> Stephenson, 1924	
<i>mediterraneus</i> Michaelsen, 1926	
<i>sabulosus</i> Southern, 1906	
<i>Fridericia</i>	
<i>callosa</i> (Eisen, 1878)	
<i>Grania</i>	
<i>atlantica</i> Coates & Erséus, 1985	
<i>macrochaeta</i> (Pierantoni, 1901)	
<i>maricola</i> Southern, 1913	
<i>mira</i> Locke & Coates, 1998	
<i>ovitheca</i> Erséus, 1977	
<i>postclitellochaeta</i> (Knöllner, 1935)	
<i>pusilla</i> Erséus, 1974	
<i>roscoffensis</i> Lasserre, 1967	
<i>variochaeta</i> Erséus & Lasserre, 1976	
<i>Henlea</i>	
<i>nasuta</i> (Eisen, 1878)	
<i>perpusilla</i> Friend, 1911	
<i>ventriculosa</i> (d'Udekem, 1854)	
<i>Lumbricillus</i>	
<i>algensis</i> Erséus, 1977	
<i>arenarius</i> (Michaelsen, 1888)	
<i>balticus</i> Bülow, 1957	
<i>brunoi</i> Martinez-Ansemil, 1982	
<i>buelowi</i> Nielsen & Christensen, 1959	
<i>cervisiae</i> Kossmagk-Stephan, 1983	
<i>christensenii</i> Tynen, 1966	
<i>dubius</i> (Stephenson, 1911)	
<i>enteromorphae</i> Bülow, 1957	
<i>fennicus</i> Nurminen, 1964	
<i>helgolandicus</i> (Michaelsen, 1934)	
<i>horridus</i> Finogenova, 1988	
<i>immoderatus</i> Finogenova, 1988	
<i>kaloensis</i> Nielsen & Christensen, 1959	
<i>knoellneri</i> Nielsen & Christensen, 1959	
	<i>Austrobella</i>
	<i>anoculata</i> Moore, 1940
	<i>Branchellion</i>
	<i>angeli</i> Sigalas, 1921
	<i>borealis</i> Leigh-Sharpe, 1933
	<i>orbiensis</i> Quatrefages, 1852
	<i>torpedinis</i> Savigny, 1822
	<i>Brumptiana</i>
	<i>lineata</i> Knight-Jones & Llewellyn, 1984
	<i>Calliobdella</i>
	<i>lophii</i> Van Beneden & Hesse, 1863
	= <i>Calliobdella punctata</i>
	= <i>Calliobdella striata</i>
	<i>nodulifera</i> (Malm, 1863)
	<i>Ganymedebdella</i>
	<i>craterae</i> Leigh-Sharpe, 1915
	<i>Hemibdella</i>
	<i>branchiarum</i> (De Silva & Kabata, 1961)
	<i>solae</i> van Beneden & Hesse, 1863
	<i>Johannssonia</i>

<i>kolaensis</i> Selensky, 1914	A	<i>brevicephalum</i> Flugel, 1990	A
<i>Janusion</i>		<i>ekmani</i> Jägersten, 1956	A
<i>brunnea</i> (Johansson, 1929)	A	<i>fiordicum</i> Webb, 1963	A
= <i>Malmiana brunnea</i>		<i>holmei</i> Southward, 1963	A
<i>bubali</i> (Srivastava, 1966)	A	<i>hyperboreum</i> Ivanov, 1960	A
= <i>Malmiana bubali</i>		<i>inerme</i> Southward & Southward, 1958	A
<i>scorpii</i> (Malm, 1863)	A	<i>lacteum</i> Southward, 1963	A
= <i>Malmiana scorpii</i>		<i>leucopleurum</i> Flugel & Callsen-Cencic, 1993	A
<i>yorki</i> (Srivastava, 1966)	A	<i>norvegicum</i> Ivanov, 1960	A
= <i>Malmiana yorki</i>		<i>pholidotum</i> Southward & Brattegard, 1968	A
<i>Mysidobdella</i>		<i>poseidonii</i> Flugel & Langhof, 1983	A
<i>oculata</i> Selensky, 1927	A		
<i>Notostomobdella</i>			
<i>laeve</i> (Levinsen, 1882)	A		
= <i>Notostomum laeve</i>			
<i>Oceanobdella</i>			
<i>blenni</i> (Knight-Jones, 1940)	A		
<i>microstoma</i> (Johansson, 1896)	A		
<i>sexoculata</i> (Malm, 1863)	A		
<i>Oxytonostoma</i>			
<i>arctica</i> Johansson, 1898	A		
= <i>Johanssonia arctica</i>			
<i>typica</i> Malm, 1863	A		
<i>Piscicola</i>			
<i>geometra</i> (Linnaeus, 1758)	A		
<i>Platybdella</i>			
<i>anarrhichae</i> (Diesing, 1859)	A		
<i>fabricii</i> Malm, 1863	A		
= <i>Crangonobdella achmerovi</i>			
= <i>Crangonobdella murmanica</i>			
<i>olriki</i> Malm, 1853	A		
<i>pinnara</i> (De Silva & Jones, 1961)	A		
<i>quadrioculata</i> Malm, 1863	A		
<i>Pontobdella</i>			
<i>muricata</i> (Linnaeus, 1758)	A M		
<i>vosmaeri</i> (Apáthy, 1888)	A M		
<i>Pontobdellina</i>			
<i>macrothela</i> (Schmarda, 1861)	A		
<i>Pterobdella</i>			
<i>jensenii</i> (Bennike & Bruun, 1939)	A		
<i>Trachelobdella</i>			
<i>alcibiades</i> (Leigh-Sharpe, 1933)	M		
<i>lubrica</i> (Grube, 1840)	A M		
Family Ozobranchidae			
<i>Ozobranchus</i>			
<i>margoi</i> (Apáthy, 1890)	A M		

Pogonophora

Compiled by Jacob van der Land and Eve Southward

The ERMS list was derived from URMO, which was based on Ivanov (1963), and scattered literature and the Zoological Record. Corrections and additions were made by Torleiv Brattegard and Eve Southward.

References

Ivanov, A. V., 1963. *Pogonophora*. Consultants Bureau; New York.

Class Pogonophora

Order Athecanephria

Family Oligobrachiidae

Nereilinum

murmanicum Ivanov, 1961

A

Oligobrachia

gracilis Southward, 1978

A

haakonmosbiensis Smirnov, 2000

A

ivanovi Southward, 1959

A

webbi Brattegard, 1966

A

Family Siboglinidae

Siboglinum

angustum Southward & Brattegard, 1968

A

atlanticum Southward & Southward, 1958

A

TARDIGRADA

Compiled by Jacob van der Land

The European list was derived from URMO.

PHYLUM TARDIGRADA

Order Eutardigrada

Family Hypsibiidae

Halobiotus

<i>appeloei</i> (Richters, 1908)	A
<i>arcturulus</i> Crisp & Kristensen, 1983	A
<i>crispae</i> Kristensen, 1982	A
<i>geddesi</i> (Hallas, 1971)	A
<i>stenostomus</i> (Richters, 1908)	A M

Order Heterotardigrada

Family Batillipedidae

<i>Batillipes</i>	
<i>acaudatus</i> Pollock, 1971	A
<i>adriaticus</i> Grimaldi de Zio, Morone de Lucia, D'Addabbo Gallo & Grimaldi, 1979	M
<i>africanus</i> D'Addabbo Gallo, Grimaldi de Zio, Morone de Lucia & Troccoli, 1992	A
<i>annulatus</i> Grimaldi de Zio, 1963	M
<i>bullacaudatus</i> McGinty & Higgins, 1968	A
<i>carnonensis</i> Fize, 1957	M
<i>litoralis</i> Renaud-Debyser, 1959	A M
<i>marcelli</i> Morone de Lucia, D'Addabbo Gallo & Grimaldi de Zio, 1988	M
<i>mirus</i> Richters, 1909	A M
<i>noerrevangi</i> Kristensen, 1978	A
<i>pennaki</i> Marcus, 1946	A M
<i>phreaticus</i> Renaud-Debyser, 1959	A M
<i>roscoffensis</i> Kristensen, 1978	A
<i>similis</i> Schulz, 1955	M
<i>tubernatis</i> Pollock, 1971	A

Family Coronarctidae

<i>Coronarctus</i>	
<i>laubieri</i> Renaud-Mornant, 1987	A
<i>stylisetus</i> Renaud-Mornant, 1987	A
<i>tenellus</i> Renaud-Mornant, 1974	A

<i>Trogloarctus</i>	
<i>trionyches</i> Villora Moreno, 1996	M

Family Echiniscoididae

<i>Echiniscoides</i>	
<i>bruni</i> D'Addabbo Gallo, Grimaldi de Zio, Morone de Lucia & Troccoli, 1992	M
<i>hoepleri</i> Kristensen & Hallas, 1980	A
<i>sigismundi</i> Plate, 1889	A M

Family Halechiniscidae

<i>Actinarcus</i>	
<i>doryphorus</i> Schulz, 1935	A M
<i>neretinus</i> Grimaldi de Zio, D'Addabbo Gallo, Morone de Lucia, Vacoarella & Grimaldi, 1982	M
<i>physophorus</i> Grimaldi de Zio, D'Addabbo Gallo, Morone de Lucia, Vacoarella & Grimaldi, 1982	M

<i>Bathycchiniscus</i>	
<i>hallasi</i> (Kristensen, 1977)	M
= <i>Styraconyx hallasi</i>	

<i>Exoclavarcutus</i>	
<i>dineti</i> Renaud-Mornant, 1983	A

<i>Florarctus</i>	
<i>acer</i> Renaud-Mornant, 1990	A M
<i>asper</i> Renaud-Mornant, 1990	M
<i>cinctus</i> Renaud-Mornant, 1975	A M
<i>hulingsi</i> Renaud-Mornant, 1975	M
<i>salvati</i> Delamare & Renaud-Mornant, 1965	M
<i>stellatus</i> Renaud-Mornant, 1990	M

<i>Halechiniscus</i>	
<i>chafarinensis</i> Grimaldi de Zio & Villora Morone, 1996	M
<i>flabellatus</i> Grimaldi de Zio, D'Addabbo Gallo, Morone de Lucia, Vacoarella & Grimaldi, 1982	M

<i>greveni</i> Renaud-Mornant & Deroux, 1976	A M
<i>guiteli</i> Richters, 1908	A M
<i>macrocephalus</i> Grimaldi de Zio, D'Addabbo Gallo, Morone de Lucia & Daddabbo Gallo, 1988	M
<i>paratuleari</i> Grimaldi de Zio, D'Addabbo Gallo & Morone de Lucia, 1988	M
<i>perfectus</i> Schulz, 1955	A M
<i>remanei</i> Schulz, 1955	A M
<i>Hemitanarctus</i>	
<i>chimaera</i> Grimaldi de Zio, D'Addabbo Gallo, Morone de Lucia & Troccoli, 1996	M
<i>Orzeliscus</i>	
<i>belopus</i> du Bois-Reymond Marcus, 1952	A M
<i>septentrionalis</i> Schulz, 1953	A
<i>Paratanarctus</i>	
<i>kristenseni</i> D'Addabbo Gallo, Grimaldi de Zio, Morone de Lucia & Troccoli, 1992	M
<i>Pleocola</i>	
<i>limnoriae</i> Cantacuzène, 1951	A
<i>Raiarctus</i>	
<i>aureolatus</i> Renaud-Mornant, 1981	A
<i>variabilis</i> D'Abbabbo Gallo, Grimaldi de Zio & Morone de Lucia, 1986	M
<i>Styraconyx</i>	
<i>craiculus</i> (Pollock, 1983)	A
<i>haploceros</i> Thulin, 1942	A
<i>kristenseni</i> Renaud-Mornant, 1981	M
<i>nanoqsunguak</i> Kristensen & Higgins, 1984	A M
<i>paulae</i> Robbotti, 1971	M
<i>qivitoq</i> Kristensen & Higgins, 1984	A M
<i>sardiniae</i> D'Addabbo Gallo, Morone de Lucia & Grimaldi de Zio, 1989	M
<i>sargassii</i> Thulin, 1942	A M
<i>testudo</i> D'Addabbo Gallo, Grimaldi de Zio & Morone de Lucia, 1984	M
<i>tyrrhenus</i> D'Addabbo Gallo, Morone de Lucia & Grimaldi de Zio, 1989	M
<i>Tanarctus</i>	
<i>gracilis</i> Renaud-Mornant, 1980	A
<i>longisetosus</i> Grimaldi de Zio, D'Addabbo Gallo, Morone de Lucia, Vacoarella & Grimaldi, 1982	M
<i>ramazzottii</i> Renaud-Mornant, 1975	A
<i>tauricus</i> Renaud-Debyser, 1959	A M
<i>Tholocarctus</i>	
<i>natans</i> Kristensen & Renaud-Mornant, 1983	A M
<i>Wingstrandarctus</i>	
<i>intermedius</i> (Renaud-Mornant, 1967)	M
= <i>Halechiniscus intermedius</i>	
<i>Family Renaudarctidae</i>	
<i>Neostygarcutus</i>	
<i>acanthophorus</i> Grimaldi de Zio, D'Addabbo Gallo & Morone de Lucia, 1982	M
<i>Family Stygarctidae</i>	
<i>Megastygarcites</i>	
<i>setuloso</i> Morgan & O'Reilly, 1988	A
<i>Mesostygarcutus</i>	
<i>intermedius</i> Renaud-Mornant, 1979	M
<i>Neoarctus</i>	
<i>primigenius</i> Grimaldi de Zio, D'Abbabbo Gallo & Morone de Lucia, 1992	M
<i>Parastygarcutus</i>	
<i>biungulatus</i> Morone de Lucia, Grimaldi de Zio & d'Addabbo Gallo, 1984	M
<i>sterreri</i> Renaud-Mornant, 1970	A M
<i>Stygarcutus</i>	
<i>bradyurus</i> Schulz, 1951	A M

CHELICERATA

Arachnida

Compiled by Anastasios Legakis

The list of the marine arachnids is definitely not complete. There are two reasons for this situation. One is that references to marine arachnids are scattered in various entomological publications some of which are old and not readily available. Second, the definition of a marine arachnid is not commonly agreed. Some authors consider them as arachnids found in marine habitats while others include those that are sporadically associated with the marine environment. The definition proposed by Chang & Frank (1993) for marine insects which could apply to marine arachnids as well, considers as marine species an insect which spends at least one of its developmental stages habitually in a marine habitat. Even in this case there are some who consider the supralittoral zone as terrestrial (Bigot *et al.*, 1984). In this list, the supralittoral zone is excluded as a very large number of species would have to be included.

The list was compiled using Riedl (1970) and Hayward & Ryland (1995).

References

- Bigot L., Picard J., Roman M.L. 1984.- Signification des peuplements d'invertébrés des plages et dunes du delta du Rhône; délimitation des domaines marin et terrestre. *C.R. Acad. Sci. Paris (III)*2980(1): 5-7.
 Chang L. & Frank J.H. 1993.- Marine insects and their reproduction. *Oceanogr. Mar. Biol. Annu. Rev.* 31: 479-506.
 Hayward P.J. & Ryland J.S. 1995.- *Handbook of the Marine Fauna of North-West Europe*. Oxford Univ. Press, pp. 462-476.
 Riedl R. 1970.- *Fauna und Flora der Adria*. Verlag Paul Parey, pp. 371-380.

Class Arachnida

Order Pseudoscorpiones

Family Neobisiidae

Neobisium maritimum (Leach)

Acarina

Halacaridae

Compiled by Ilse Bartsch

The list of species covers halacarid mites, family Halacaridae, Superfamily Halacaroidea, suborder Prostigmata. It is the only mite family which completely is adapted to life in the sea; halacarids are part of the benthos from the upper littoral fringe to the deep sea. The northeastern Atlantic Ocean, with adjacent seas, is an area studied intensely in respect to its halacarid fauna and assumedly half of the species are known. The most relevant descriptive papers are listed below.

Beside halacarids, numerous other mites inhabit tidal and shallow subtidal areas. Seashore mites in tidal areas, belonging to the 'terrestrial moiety', are representatives of the suborders Mesostigmata, Astigmata, Prostigmata, and Cryptostigmata (Oribatei). Another mite family, the Pontarachnidae (Prostigmata) are commonly found in warm waters regions, in a depth of 0-10 m. The Pontarachnidae are most similar and closely related to

freshwater 'Hydrachneilae'. It is intended to include these groups in future editions of ERMS.

References

- Abé, H. & O'Connor, J.P. 1991. A redescription of *Scaptognathus trouessarti* Halbert, 1915 (Acar, Halacaridae) with notes on some taxonomic characters. *Ir. Nat. J.*, 23: 429-472.
 André, M. 1938. Description de six Halacariens de Terre de Feu (2e partie). *Bull. Mus. nat. hist. nat.*, sér. 2, 10: 385-390.
 André, M. 1939a. Sur trois espèces de *Rhombognathus* (Halacariens) des côtes françaises. *Bull. Mus. nat. hist. nat.*, sér. 2, 11: 308-313.
 André, M. 1939b. Description du *Coloboceras longiusculus* Trouessart (Halacarien). *Bull. Mus. nat. hist. nat.*, sér. 2, 11: 402-404.
 André, M. 1940. Description du *Copidognathus* (s. str.) *crassirostris* Trouessart (Halacarien). *Bull. Mus. nat. hist. nat.*, sér. 2, 12: 329-331.
 André, M. 1941. Sur les *Agauopsis hirsuta* Trt. et *microrhyncha* Trt. (Halacariens). *Bull. Soc. zool. Fr.*, Paris, 66: 257-271.
 André, M. 1946. Halacariens marins. *Faune Fr.*, 46: 152 pp.
 André, M. 1961. Description d'une nouvelle espèce du genre *Scaptognathus*, recueillie en Méditerranée (*Scaptognathus sabularius* n. sp.). *Acarologia*, 3: 297-302.
 Angelier, E. 1950. *Halacarus (Halacarellus) petiti* - halacariens nouveau d'étang de Salses (Pyrénées Orientales). *Vie Milieu*, 1 : 214-216.
 Angelier, E. 1954. Halacariens des sables littoraux méditerranéens. *Vie Milieu*, 4 : 281-289.
 Bartsch, I. 1972. Ein Beitrag zur Systematik, Biologie und Ökologie der Halacaridae (Acarini) aus dem Litoral der Nord- und Ostsee. I. Systematik und Biologie. *Abh. Verh. naturw. Ver. Hamburg*, (NF) 16: 155-230.
 Bartsch, I. 1973a. Halacaridae (Acarini) von der Josephinebank und der großen Meteorbank aus dem östlichen Nordatlantik. I. Die Halacaridae aus den Schleppnetzproben. "Meteor" Forsch.-Ergebn., D, 13: 37-46.
 Bartsch, I. 1973b. Halacaridae (Acarini) von der Josephinebank und der großen Meteorbank aus dem östlichen Nordatlantik. II. Die Halacaridae aus den Bodengreiferproben. "Meteor" Forsch.-Ergebn., D, 15: 51-78.
 Bartsch, I. 1973c. *Copidognathus raekor*, eine neue Halacaride (Acarini) von der Großen Meteorbank. "Meteor" Forsch.-Ergebn., D, 16: 65-68.
 Bartsch, I. 1974a. *Simogathus minutus* (Hodge) (Halacaridae, Acari), eine Wiederbeschreibung nebst Bemerkung zur Unterfamilie der Simognathinae. *Cah. Biol. mar.*, 15: 275-284.
 Bartsch, I. 1974b. Über das Auftreten von Epimeralporen besonders bei den Rhombognathinae (Halacaridae, Acari). *Zool. Anz.* 193: 266-268.
 Bartsch, I. 1975a. Ein Beitrag zur Rhombognathinen-Fauna (Halacaridae, Acari) der Bretagne-Küste. *Acarologia*, 17: 53-80.
 Bartsch, I. 1975b. Beitrag zur Halacaridenfauna des Eulitorals bei Banyuls-sur-Mer (Mittelmeer) (Halacaridae, Acari). *Mitt. hamb. zool. Mus. Inst.*, 72: 137-155.
 Bartsch, I. 1976a. Beitrag zur Halacariden-Fauna (Halacaridae, Acari) der Bretagne-Küste. Beschreibung von fünf Arten aus dem Sandlückensystem. *Acarologia*, 17: 652-667.
 Bartsch, I. 1976b. *Copidognathides minutirostris* n. g., n. sp., eine psammophile Halacaride (Halacaridae, Acari). *Cah. Biol. mar.*, 17: 237-241.
 Bartsch, I. 1976c. Zur Systematik und Verbreitung der *Halacarellus*-Arten (Halacaridae, Acari) an der Bretagne-Küste. *Ent. Mitt. Zool. Mus. Hamb.*, 5: 97-109.
 Bartsch, I. 1976d. Ergänzung zur Halacariden-Fauna (Halacaridae, Acari) im Becken von Arcachon. *Vie Milieu*, A, 26: 31-46.
 Bartsch, I. 1976e. *Copidognathus stevcici* n. sp., eine parasitisch lebende Halacaride (Halacaridae, Acari). *Thalass. Jugosl.*, 12: 399-404.

- Bartsch, I. 1977a. Ergänzung zur Halacariden-Fauna (Halacaridae, Acari) der Bretagne-Küste. Zur Gattung *Arhodeoporus* und *Copidognathus*. *Acarologia*, 18: 626-641.
- Bartsch, I. 1977b. New species of *Lohmannella* (Halacaridae, Acari) from the Roscoff area, Brittany. *Cah. Biol. mar.*, 18: 141-153.
- Bartsch, I. 1977c. Zur *oculatus*- und *gibbus*-Gruppe der Gattung *Copidognathus* (Halacaridae, Acari). *Ent. Mitt. Zool. Mus. Hamb.*, 6: 1-12.
- Bartsch, I. 1977d. Eine neue *Actacarus*-Art (Acari, Halacaridae) aus dem Bathyal vor der Küste von North Carolina, USA. *Zool. Scripta*, 6: 323-326.
- Bartsch, I. 1978a. Halacaridae (Acari) aus der Tiefsee des atlantischen Ozeans. *Cah. Biol. mar.*, 19: 47-62.
- Bartsch, I. 1978b. Halacaridae (Acari) von Gezeitenstränden Nordnorwegens. *Mikrofauna Meeresboden*, 70: 1-22.
- Bartsch, I. 1979a. *Copidognathus hartwigi* n. sp. (Halacaridae, Acari), eine Meeresmilbe neu für die Fauna der Bermudas. *Acarologia*, 20: 93-96.
- Bartsch, I. 1979b. Ergänzungen zur *Copidognathus*-Fauna (Halacaridae, Acari) der Bretagne-Küste. *Acarologia*, 20: 217-234.
- Bartsch, I. 1979c. Halacaridae (Acari) von der Atlantikküste Nordamerikas. Beschreibung der Arten. *Mikrofauna Meeresboden*, 79: 1-62.
- Bartsch, I. 1980a. Halacaridae (Acari) aus der Bucht von Morlaix (Bretagne). *Acarologia*, 21: 34-45.
- Bartsch, I. 1980b. Halacariden (Acari) im marinem Mesopsmal der Ostküste Nordamerikas. *Ent. Mitt. Zool. Mus. Hamb.*, 6: 393-407.
- Bartsch, I. 1983a. *Halacarellus coecus* (Trouessart) und *H. princeps* (Trouessart) (Halacaridae, Acari), Wiederbeschreibung der Arten. *Acarologia*, 24: 49-57.
- Bartsch, I. 1983b. *Copidognathus richardi* Trouessart (Halacaridae, Acari), eine Wiederbeschreibung. *Acarologia*, 24: 285-288.
- Bartsch, I. 1984. Ergänzungen zur Halacariden-Fauna der Bretagne-Küste und Beschreibung einer neuen Art (Halacaridae, Acari). *Cah. Biol. mar.*, 25: 113-122.
- Bartsch, I. 1985a. Halacaridae (Acari) from the Strangford Narrows and the Irish Sea. *Proc. R. Ir. Acad.*, 85 (B): 21-35.
- Bartsch, I. 1985b. Notes on the Halacaridae (Acari) from Yorkshire. *Naturalist*, 110: 41-48.
- Bartsch, I. 1986a. Zur Gattung *Agauopsis* (Acari, Halacaridae), Beschreibung zweier Arten und Übersicht über Verwandtschaftsgruppen. *Zool. Scr.*, 15: 165-174.
- Bartsch, I. 1986b. Lohmannellinae (Halacaroidea, Acari) aus dem Mittelmeer. *Entomol. Mitt. zool. Mus. Hamburg*, 8: 231-244.
- Bartsch, I. 1986c. *Halacarellus affinis* (Halacaridae, Acari), eine Wiederbeschreibung. *Acarologia*, 27: 247-250.
- Bartsch, I. 1986d. New species of the genus *Rhombognathus* (Acari, Halacaridae), and a key to Mediterranean *Rhombognathus*. *Mesogée*, 46: 3-7.
- Bartsch, I. 1986e. Zur Gattung *Werthella* Lohmann, *Pelacarus* n. gen. und *Werthelloides* n. gen. (Halacaridae, Acari). *Cah. Biol. mar.*, 27: 211-223.
- Bartsch, I. 1987. *Halacarus griseus* sp. n., eine Halacaride (Acari) aus dem Mittelmeer. *Entomol. Mitt. zool. Mus. Hamburg*, 9: 75-79.
- Bartsch, I. 1988. Halacariden (Acari) im Nordatlantik. Beschreibungen von *Halacarus balginus* sp. n. und *Agauides cryosi* gen. et sp. n. *Cah. Biol. mar.*, 29: 353-359.
- Bartsch, I. 1989. *Copidognathus brifaci*s n. sp. (Halacaridae, Acari) und Bemerkungen zu weiteren Bewohnern des Seeigels *Spatangus purpureus* (Echinoidea). *Mitt. hamb. zool. Mus. Inst.*, 86: 315-328.
- Bartsch, I. 1990. Hydrothermal vent fauna: *Agauopsis auzendei* n. sp. (Acari, Halacaridae). *Bull. Mus. nat. hist. nat.*, Paris, 4e sér. 12, Sect. A: 69-73.
- Bartsch, I. 1991a. Taxonomic notes on halacarids (Acari) from the Skagerrak area. *Helgoländer Meeresunters.*, 45: 97-106.
- Bartsch, I. 1991b. On the identity of some North Atlantic halacarid species (Acari). *J. nat. Hist.*, 25: 1339-1353.
- Bartsch, I. 1992. *Bradyagae stocki* nov. spec., a deep-sea halacarid mite (Halacaridae, Acari) from the Atlantic Ocean. *Bull. Zool. Mus. Amsterdam* 13 (8): 81-84.
- Bartsch, I. 1994. Halacarid mites (Acari) from hydrothermal deep-sea sites. New records. *Cah. Biol. Mar.*, 35: 479-490.
- Bartsch, I. 1996a. Halacarid mites (Acari) from the Mid-Atlantic Ridge. New records. *Cah. Biol. Mar.*, 37: 159-167.
- Bartsch, I. 1996b. *Agauopsis* (Acari, Halacaridae) of the Sevastopol area; supplementary notes on taxonomy and ecology. *Rev. suisse Zool.*, 103: 697-712.
- Bartsch, I. 1996c. Rhombognathines (Acari: Halacaridae) of the Black Sea: A survey. *Mitt. hamb. zool. Mus. Inst.*, 93: 141-160.
- Bartsch, I. 1997a. *Copidognathus nautilei* (Acari: Halacaridae), a new mite from a hydrothermal vent field in the Mid-Atlantic Ridge. *Cah. Biol. Mar.*, 38: 29-33.
- Bartsch, I. 1997b. A new species of the *Copidognathus tricornatus* group (Acari: Halacaridae) from Western Australia with a review of the species-group. *Species Diversity*, 2: 155-166.
- Bartsch, I. 1998a. First record of *Isobactrus* from the Black Sea (Acari, Halacaridae, Rhombognathinae). *Ent. Mitt. zool. Mus. Hamb.*, 12: 269-276.
- Bartsch, I. 1998b. *Colobocerasides koehleri* (Trouessart) und *C. auster* n. sp. (Arachnida: Acari: Halacaridae), Beschreibung der Arten. *Mitt. Mus. Natkd. Berl., Zool. Reihe*, 74: 225-232.
- Bartsch, I. 1998c. Halacrinae (Acari, Halacaroidea) from the northwestern Black Sea: A review. *Mitt. hamb. zool. Mus. Inst.*, 95: 143-178.
- Bartsch, I. 1999. *Actacarus ponticus*, a new marine mite from the Black Sea (Arachnida, Acari, Halacaridae). *Senckenberg. biol.*, 78: 141-151.
- Benard, F. 1962. Sur une nouvelle espèce d'Acariens marin, *Agauopsis tricuspidis* (superfamille des Prostigmata, famille des Halacaridae). *Acarologia*, 4: 215-229.
- Green, J. & MacQuitty, M. 1987. Halacarid Mites. *Synops. British Fauna*, (ns), 36: 178 pp.
- Halbert, J.N. 1915. The Acarina of the seashore. *Proc. R. Ir. Acad.*, (Sect. II) 31: 45-136.
- Laubier, L. 1960. *Parhalixodes travei* n. g., n. sp., un nouvel Halicodinae (Halacariens) ectoparasite de nemerte en Méditerranée occidentale. *Acarologia*, 2: 541-551.
- Konnerth-Ionescu, A. 1970. Nouvelles données sur les halacarides de la zone psamique du littoral roumain de la Mer Noire. *Trav. Mus. hist. Nat. 'Grigore Antipa'* 10: 19-23.
- Lohmann, H. 1889. Die Unterfamilie der Halacaridae MURR. und die Meeresmilben der Ostsee. *Zool. Jb. (Syst.)*, 4: 269-408.
- Lohmann, H. 1893. Die Halacarinen der Plankton-Expedition. *Ergebn. Atlant. Ozean Planktonexp. Humboldt-Stift.*, 2. G. a: 11-95.
- Mari, M., Morselli, I. 1990. Idracnelle ed Alacaridi (Acari) di una Prateria di *Posidonia oceanica* (L.) delile dell'Isola d'Ischia. *Atti Soc. Tosc. Sci. Nat., Mem.*, (B), 96 : 243-256.
- Monniot, F. 1961. *Simognathus andrei*, nouvelle espèce d'halacarien recoltée en Méditerranée. *Acarologia* 3: 585-590.
- Monniot, F. 1962a. Recherches sur les graviers a *Amphioxus* de la région de Banyuls-sur-Mer. *Vie Milieu*, 13 : 231-322.
- Monniot, F. 1962b. Description d'une nouvelle espèces d'acariens, Halacaridae : *Coloboceras drachi*, recoltée en Méditerranée. *Bull. Soc. zool. Fr.*, 87 : 492-497.
- Monniot, F. 1964. Sur deux espèces du genre *Scaptognathus* présentes à Roscoff. *Acarologia*, 6: 491-498.
- Monniot, F. 1967. Deux halacariens endoparasitiques : *Halacarus anomalus* Trouessart 1894 et *Halacarus marcandrei* n. sp. *Cah. Biol. mar.*, 8: 82-98.
- Morselli, I. 1969. Ricerche sugli Alacari di delle coste livornesi. I. - Studio preliminare di alcune specie raccolte su fondi sabbiosi. *Atti Soc. Nat. Mat. Modena*, 100: 1-21.

- Morselli, I. 1970. Ricerche sugli Alacari di delle coste livornesi. II. - *Acaromantis monnioti*, una nuova specie raccolta sui fondi sabbiosi delle Secche della Meloria (Livorno). *Pubbl. Staz. Zool. Napoli*, 38 : 109-119.
- Morselli, I. 1980. Su tre Acari prostigmati di acque salmastre dell'alto Adriatico. *Atti Soc. Tosc. Sc. Nat., Mem.*, (B), 87: 181-195.
- Morselli, I. 1981. Ricerche sugli Alacaridi delle coste livornesi. III. - *Plegadognathus labronicus* n.g., n.sp. (Acari, Prostigmata), un alacaride di sabbie grossolane. *Atti Soc. Tosc. Sci. Nat., Mem.*, (B), 87 : 265-273.
- Morselli, I. & Mari, M. 1979. Alacaridi di acque lagunari e costiere della Toscana meridionale. *Atti Soc. Tosc. Sc. Nat., Mem.*, (B), 85 : 63-91.
- Morselli, I. & Mari, M. 1981. *Anomalohalacarus affinis* n.sp., un alacaride (Halacaridae, Acari) delle coste Toscane. *Atti Soc. Tosc. Sci. Nat., Mem.*, (B) 87 : 275-283.
- Morselli, I. & Mari, M. 1982. Alacaridi (Acari, Prostigmata) di fondi sabbiosi della Costa Ionica del Salento. *Atti Soc. Tosc. Sc. Nat., Mem.*, (B) 88 : 229-247.
- Morselli, I. & Mari, M. 1985. Ricerche sugli alacaridi delle coste Livornesi. IV. - Osservazioni su alcune specie raccolte su fondi sabbiosi della zona di Piombino. *Atti Soc. Tosc. Sc. Nat., Mem.*, (B) 91: 201-220.
- Morselli, I. & Mari, M. 1989. Halacarids (Halacaridae, Acari) of the Leghorn coast. V. *Anomalohalacarus singularis*, a new interstitial infralittoral species. *Bull. Zool.*, 56: 105-108.
- Motas, C. & Soarec, J. 1940. Sur quelques halacarides de la Mer Noire. *Annls scient. Univ. Jassy*, 26: 139-175.
- Newell, I. M. 1947. A systematic and ecological study of the Halacaridae of eastern North America. *Bull. Bingham oceanogr. Coll.*, 10: 1-232.
- Newell, I. M. 1956. *Pachygnathus notops* Gosse 1855 - 100 years later. *Ann. Mag. Nat. Hist.*, ser. 12, 9: 465-475.
- Newell, I. M. & André, M. 1959. Révision des espèces de *Rhombognathus* (Halacariens marins), décrites par Édouard L. Trouessart. *Acarologia*, 1: 124-146.
- Petrova, A. 1972. Sur la présence d'*Halacarellus subterraneus* Schulz, 1933 et *Halacarellus phreaticus* n. sp. (Halacaridae, Acari) en Bulgarie. *Acarologia*, 13: 367-373.
- Petrova, A. 1976. Une nouvelle espèce d'*Agauiopsis* (Prostigmata, Halacaridae) du littoral de la Mer Noire. *Chidrobol., Sofiya*, 4: 67-70.
- Police, G. 1909. Alcune nuove specie di Halacaridae del Golfo di Napoli. *Arch. zool. ital.*, 3: 409-443.
- Schulz, E. 1933. Zur Halacaridenfauna der Kieler Bucht. *Schr. naturw. Ver. Schlesw.-Holst.*, 20: 96-105.
- Schulz, E. 1937. *Actacarus pygmaeus* n.g., n.sp., eine merkwürdige Meeresmilbe aus der Otoplanen-Zone der Nordsee. *Kieler Meeresforschungen*, 1: 327-331.
- Sokolov, I. 1946. New species of Halacaridae from the Arctic Ocean. Trudy dreifujuei ekspedicia glavsevmorputi na ledokolnom parohode 'G. Sedov' 1937-1940 gg.' 3: 302-307.
- Travé, J. 1972. Premières données sur les Halacariens (Acariens) interstitiels de Grèce. *Biol. Gallo-Helen.*, 4: 1972.
- Trouessart, E., 1893. Note sur les acariens recueillis au Spitzberg pendant le voyage de la Manche. *Nouv. Arch. missions scient. littérair.*, Paris, 5: 255-263.
- Trouessart, E., 1894. Note sur les acariens marins (Halacaridae) dragués par M.P. Hallez dans le Pas-de-Calais. *Revue biol. N. Fr.* 6: 154-184.
- Trouessart, E., 1896. Halacariens. Résultats scientifiques de la Campagne du 'Caudan' dans le Golfe de Gascogne. *Ann. Univ. Lyon* 26: 325-353.
- Trouessart, E. 1998. Note sur les acariens marins (Halacaridae) récoltés par M. Henri Gadeau de Kerville sur le littoral du département du Calvados et aux îles Saint-Marcouf (Manche). *Bull. Soc. Amis Sci. nat. Rouen*, (sér 4) 33: 423-433.
- Trouessart, E. 1901. Note sur les acariens marins (Halacaridae) récoltés par M. Henri Gadeau de Kerville dans la région d'Omonville-la-Rogue (Manche) et dans la fosse de la Hague. *Bull. Soc. Amis Sci. nat. Rouen*, (Sér 4) 14: 247-266.
- Trouessart, E. & Neumann, G., 1893. Description d'un nouveau genre et d'une nouvelle espèce de la famille des Halacaridae ou Acariens marins. *Naturaliste*, Paris (s. 2), 15: 207-208.
- Viets, K. 1927a. Die Halacaridae der Nordsee. *Z. wiss. Zool.*, 130: 83-173.
- Viets, K. 1927b. Halacaridae. *Tierwelt N.-u. Ostsee* XIc: 1-72.
- Viets, K. 1928a. Wassermilben aus dem Schwarzen Meer, dem Kaspiischen Meer und dem Aral-See. *Abh. naturw. Ver. Bremen*, 27: 47-80.
- Viets, K. 1928b. Halacaridae von der Murman-Küste, aus dem Weissen Meer und von Nowaja-Semlja. *Issled. Morei SSSR*, 6: 81-88.
- Viets, K. 1933. Kleine Sammlungen in- und ausländischer Wassermilben. *Zool. Anz.*, 104: 261-274.
- Viets, K. 1935. The Fishery Grounds near Alexandria. IV. Some marine mites from Alexandria. *Notes Mem. Fishery Res. Dir.*, Cairo, 11: 1-3.
- Viets, K. 1936. Über eine neue Halacaride (Acari) aus Bulgarien. *Zool. Anz.*, 115: 199-202.
- Viets, K. 1939a. Halacariden (Acari) aus süditalienischen Höhlengewässern. *Arch. Hydrobiol.*, 35: 625-630.
- Viets, K. 1939b. Meeresmilben aus der Adria (Halacaridae und Hydrachnellae, Acari). *Arch. Naturgesch.*, (N.F.) 8: 518-550.
- Viets, K. 1940. Meeresmilben aus der Adria (Halacaridae und Hydrachnellae, Acari). *Arch. Naturgesch.*, (N.F.) 9: 1-135.
- General papers**
- André, M. 1946. Halacariens marins. *Faune Fr.*, 46: 152 pp.
- Bartsch, I. 1997. *Thalassarchna* und *Halacarellus* (Halacaridae: Acari): two separate genera. *J. nat. Hist.*, 31: 1223-1236.
- Green, J. & MacQuitty, M. 1987. Halacarid Mites. *Synops. British Fauna*, (ns), 36: 178 pp.
- Sokolov, I.I. 1952. Vodjanye kleci. II. Halacarae. *Fauna SSSR*, 5: 201 pp.
- Viets, K. 1927b. Halacaridae. *Tierwelt N.- u. Ostsee*, XIc: 1-72.
- Viets, K. 1936. Spinnentiere oder Arachnoidea. VII. Wassermilben oder Hydracarina (Hydrachnellae und Halacaridae). *Tierwelt Dfl.*, 31, 32: 1-574.
- Viets, K. 1956. Die Milben des Süßwassers und des Meeres. Katalog der Halacaridae, Meeresmilben. II. Abschnitt. In Viets, K. (Ed.) *Die Milben des Süßwassers und des Meeres. Hydrachnellae et Halacaridae*. Fischer Verlag Jena: 641-870.
- Family Halacaridae**
- Subfamily Acarochelopodinae**
- Acarochelopodia*
- angelieri* Travé, 1972 M
 - delamarei* Angelier, 1954 A M
 - tarentina* Morselli et al., 1998 M
- Subfamily Actacarinae**
- Actacarus*
- bacescui* Konnerth-Ionescu, 1970 A M
 - obductus* Bartsch, 1977 A
 - ponticus* Bartsch, 1999 M
 - pygmaeus* Schulz, 1937 A M
- Subfamily Anomalohalacarinae**
- Anomalohalacarus*
- acnemus* Bartsch, 1985 A
 - affinis* Morselli & Mari, 1981 M
 - anomalus* (Trouessart, 1894) A M
 - arenarius* Bartsch, 1976 A
 - intermedius* Bartsch, 1976 A
 - marcandrei* Monniot, 1967 A
 - minutus* Bartsch, 1976 A
 - mollis* Morselli & Mari, 1982 M
 - poizati* Bartsch, 1985 A
 - ruffoi* Morselli & Mari, 1979 M
 - septentrionalis* Bartsch, 1991 A
 - similis* Bartsch, 1976 A M
 - singularis* Morselli & Mari, 1989 M

Subfamily Copidognathinae

<i>Copidognathides</i>			
<i>minutirostris</i> subsp. <i>minutirostris</i> Bartsch, 1976	A	<i>panopae</i> (Lohmann, 1893)	A M
<i>minutirostris</i> subsp. <i>magniporus</i> Bartsch, 1976	A	<i>plutonia</i> Bartsch, 1996	A
<i>ampliatus</i> Bartsch, 1999	M	<i>Agauides</i>	
<i>Copidognathus</i>		<i>cryosi</i> Bartsch, 1988	A
<i>alvinus</i> Bartsch, 1994	A	<i>Agauopsis</i>	
<i>arcticus</i> Sokolov, 1946		<i>brevipalpus</i> (Trouessart, 1889)	A M
<i>brachystomus</i> Viets, 1940	M	= <i>A. pontica</i> Chichkoff 1907	
<i>brevipes</i> Viets, 1940	M	= <i>A. spinipes</i> (Police, 1909) ?	
<i>brevirostris</i> Viets, 1927	A M	<i>conjuncta</i> Viets, 1940	M
<i>brifacius</i> Bartsch, 1989	A	<i>ibssi</i> Bartsch, 1996	M
<i>caudani</i> (Trouessart, 1896)	A	<i>marinovi</i> Petrova, 1976	A M
<i>crassirostris</i> (Trouessart, 1901) ?	A	= <i>A. aremorica</i> Bartsch 1984	
<i>cribosoma</i> (Police, 1909)	M	<i>meteoris</i> Bartsch, 1973	A
<i>dentatus</i> Viets, 1940	A M	<i>microrhyncha</i> (Trouessart, 1889)	A M
<i>extensus</i> Viets, 1940	M	<i>minor</i> (Trouessart, 1894)	A
<i>fabricii</i> (Lohmann, 1889)	A M	= <i>A. meteoris</i> Bartsch, 1980	
= <i>C. fabricius</i>		<i>pteropea</i> Bartsch, 1986	M
<i>falcifer</i> Viets, 1940	M	<i>spinosa</i> Bartsch, 1978	A
<i>figeus</i> Bartsch, 1976	A	<i>tricuspis</i> Benard, 1962	A M
<i>gibbus</i> (Trouessart, 1889)	A M	= <i>Halacarellus minutipalpis</i> Viets 1937	
<i>glyptoderma</i> (Trouessart, 1888)	A	<i>Arhodeoporus</i>	
<i>gorbunovi</i> Sokolov, 1946	A	<i>brevocularis</i> Bartsch, 1973	A
<i>granulatus</i> (Hodge, 1863)	A	<i>gracilipes</i> (Trouessart, 1889)	A M
<i>hartwigi</i> Bartsch, 1979	A	<i>labronicus</i> (Morselli, 1981)	M
<i>humerosus</i> (Trouessart, 1896)	A M	= <i>Plegadognathus labronicus</i>	
<i>lamellosus</i> Lohmann, 1893	A M	<i>lineatus</i> Bartsch, 1973	A
<i>latisetus</i> Viets, 1940	A M	<i>minor</i> Bartsch, 1977	A
<i>latus</i> Viets, 1927	A	<i>Atelopsalis</i>	
<i>longipes</i> Bartsch, 1973	A	<i>newelli</i> Bartsch, 1973	A
<i>longirostris</i> (Trouessart, 1896)	M	<i>tricuspis</i> Trouessart, 1896	A
<i>loricifer</i> André, 1946	A	<i>Bathyhalacarus</i>	
<i>majusculus</i> (Trouessart, 1894)	M	sp.	A
<i>magnipalpus</i> subsp. <i>magnipalpus</i> (Police, 1909)	A M	<i>Bradyagae</i>	
= <i>C. magnipalpus</i> subsp. <i>serratisetus</i> Viets 1928		<i>alberti</i> (Trouessart, 1902)	A
<i>magnipalpus</i> subsp. <i>ponticus</i> Viets, 1936	M	<i>meteoris</i> Bartsch, 1991	A
<i>magniporus</i> Bartsch, 1973	A	<i>stocki</i> Bartsch, 1992	A
<i>mucronatus</i> Viets, 1928	M	<i>Camactognathus</i>	
<i>nautilei</i> Bartsch, 1997	A	<i>borealis</i> Bartsch, 1991	A
<i>novus</i> Bartsch, 1980	A	<i>tesselatus</i> (Morselli & Mari, 1982)	M
<i>oculatus</i> sensu Lohmann, 1889	A M	= <i>Arhodeoporus tessellatus</i>	
<i>oxianus</i> Viets, 1928		<i>Caspihalacarus</i>	
<i>perforatus</i> Viets, 1940	M	<i>hyrcanus</i> Viets, 1928	M
<i>ponteuximus</i> subsp. <i>ponteuximus</i> Viets, 1928	M	<i>Coloboceras</i>	
<i>ponteuximus</i> subsp. <i>pectiniger</i> Motas & Soarec, 1940	M	<i>drachi</i> Monniot, 1962	A M
<i>poucheti</i> (Trouessart, 1893)	A	<i>karamani</i> Bartsch, 1973	A
<i>pseudofigueus</i> Bartsch, 1977	A	<i>longiusculus</i> Trouessart, 1889	A M
<i>pulcher</i> (Lohmann, 1893)	M	<i>Colobocerasides</i>	
<i>quadricostatus</i> (Trouessart, 1894)	A M	<i>koehleri</i> (Trouessart, 1896)	A
= <i>C. adriaticus</i> Viets, 1940 ?		<i>Halacarellus</i>	
= <i>C. consimilis</i> Bartsch 1979		<i>auzendei</i> (Bartsch, 1990)	A
<i>raekor</i> Bartsch, 1973	A	= <i>Agauposis auzendei</i>	
<i>remipes</i> (Trouessart, 1894)	A M	<i>balticus</i> (Lohmann, 1889)	A
<i>reticulatus</i> (Trouessart, 1893)	A	= <i>Thalassarachna baltica</i>	
<i>rhodostigma</i> subsp. <i>rondus</i> Bartsch, 1979	A M	<i>capuzinus</i> (Lohmann, 1893) ?	A
<i>rhodostigma</i> subsp. <i>typicus</i> (Gosse, 1855)	A M	= <i>H. areolatus</i> Halbert, 1915 ?	
<i>richardi</i> (Trouessart, 1902)	A	= <i>Thalassarachna capuzina</i>	
<i>sculptus</i> (Police, 1909)	M	<i>chersonesus</i> Bartsch, 1998	M
<i>septentrionalis</i> (Halbert, 1915)	A	<i>discretus</i> Bartsch, 1998	M
= <i>C. graveolus</i> Monniot, 1962 ?		<i>floridarium</i> (Lohmann, 1889)	A
= <i>C. lubricellus</i> Bartsch 1977		= <i>H. bergensis</i> Viets 1927	
<i>stevcici</i> Bartsch, 1976	A M	<i>micropectinatus</i> Bartsch, 1972	A M
<i>tectirostris</i> Bartsch, 1979	A M	<i>petiti</i> Angelier, 1950	A M
<i>tabellio</i> (Trouessart, 1894)	A M	= <i>Thalassarachna petiti</i>	
<i>trouessarti</i> (Voinov, 1896)	A M	<i>phreaticus</i> Petrova, 1972	M
= <i>C. tricornatus</i> Bartsch, 1973, 1979		<i>procerus</i> Viets, 1927	A M
<i>venustus</i> Bartsch, 1977	A	= <i>Thalassarachna procera</i>	
Subfamily Halacarinae		<i>southerni</i> Halbert, 1915	A
<i>Agae</i>		= <i>Thalassarachna southerni</i>	
<i>abyssorum</i> (Trouessart, 1896)	A	<i>subcrispus</i> Bartsch, 1978	A
= <i>Halacarus</i> (<i>Halacarellus</i>) <i>abyssorum</i>		<i>subterraneus</i> Schulz, 1933	A M
<i>adriatica</i> Viets, 1940	A M	= <i>Thalassarachna subterranea</i>	
<i>chevreuxi</i> (Trouessart, 1889)	A M	<i>Halacaropsis</i>	
<i>corollata</i> Bartsch, 1978	A	<i>hirsuta</i> (Trouessart, 1889)	A M
		= <i>Agauposis hirsuta</i>	
		<i>Halacarus</i>	

<i>actenos</i> Trouessart, 1889	A	<i>Metarhombognathus</i>
<i>aegyptus</i> Viets, 1935	M	<i>armatus</i> (Lohmann, 1893)
<i>balgimus</i> Bartsch, 1988	A	<i>nudus</i> (Viets, 1928)
<i>bisulcus</i> Viets, 1927	A M	<i>Rhombognathides</i>
<i>borealis</i> Trouessart, 1893	A	<i>brevipes</i> Bartsch, 1975
<i>ctenopus</i> sensu Lohmann, 1893	A	<i>exoplus</i> (Trouessart, 1901)
<i>griseus</i> Bartsch, 1987	M	<i>mucronatus</i> (Viets, 1927)
<i>longiunguis</i> Police, 1909	M	<i>pascens</i> (Lohmann, 1889)
<i>prolongatus</i> Bartsch, 1996	A	<i>seahami</i> (Hodge, 1860)
<i>rismondoi</i> Viets, 1940	M	<i>spinipes</i> (Viets, 1933)
<i>spiniger</i> Bartsch, 1973	A	<i>trionyx</i> subsp. <i>trionyx</i> (Trouessart, 1899)
<i>subtilis</i> Viets, 1940	M	= <i>R. merrimani</i> subsp. <i>needleri</i> Newell 1947
<i>Parhalixodes</i>	M	<i>trionyx</i> subsp. <i>merrimani</i> Newell, 1947
<i>travel</i> Laubier, 1960	M	= <i>R. merrimani</i> subsp. <i>merrimani</i> Newell 1947
<i>Pelacarus</i>	A M	<i>Rhombognathus</i>
<i>aculeata</i> (Trouessart, 1896)		<i>cetratus</i> Bartsch, 1974
= <i>Werthella aculeata</i>		<i>conjunctus</i> Bartsch, 1986
<i>Thalassarachna</i>	M	<i>karlvietsi</i> Bartsch, 1975
<i>affinis</i> (Trouessart, 1896)		<i>intermedius</i> Schulz, 1933
= <i>Halacarellus affinis</i>		<i>lionyx</i> Trouessart, 1899
<i>alvina</i> (Bartsch, 1994)	A	<i>magnirostris</i> Trouessart, 1889
= <i>Halacarellus alvinus</i>		<i>notops</i> (Gosse, 1855)
<i>basteri</i> (Johnston, 1836)	A M	<i>paranotops</i> Bartsch, 1986
= <i>Halacarellus basteri</i>		<i>parvulus</i> Viets, 1939
<i>caecoides</i> (Bartsch, 1978)	A	<i>peltatus</i> Viets, 1939
= <i>Halacarellus caecoides</i>		<i>ponticus</i> Motas & Soarec, 1940
<i>coeca</i> (Trouessart, 1902)	A	<i>praegracilis</i> Viets, 1939
= <i>Halacarellus coecus</i>		<i>procerus</i> Bartsch, 1975
<i>hexacantha</i> (Viets, 1927)	A	<i>sphaerorhynchus</i> Police, 1909
= <i>Halacarellus hexacanthus</i>		<i>subtilis</i> Bartsch, 1975
<i>inermis</i> (Trouessart, 1888)	A	<i>tonops</i> Bartsch, 1996
= <i>Halacarellus inermis</i>		
<i>litoralis</i> (Bartsch, 1976)	A	Subfamily Simognathinae
= <i>Halacarellus litoralis</i>		<i>Acaromantis</i>
<i>longipes</i> (Trouessart, 1888)	A	<i>minutus</i> Bartsch, 1976
= <i>Halacarellus longipes</i>		<i>monnioti</i> Morselli, 1970
<i>princeps</i> (Trouessart, 1902)	A	<i>squilla</i> Trouessart & Neumann, 1893
= <i>Halacarellus princeps</i>		<i>Simognathus</i>
<i>reticulata</i> (Bartsch, 1978)	A	<i>minutus</i> (Hodge, 1863)
= <i>Halacarellus reticulatus</i>		<i>leiomerus</i> Trouessart, 1894
<i>robusta</i> (Bartsch, 1978)	A	= <i>S. andrei</i> Monniot 1961
= <i>Halacarellus robustus</i>		
<i>striata</i> (Lohmann, 1889)	A	Pycnogonida
= <i>Halacarellus similis</i> Viets 1927		Compiled by Jacob van der Land and Franz Krapp
= <i>Halacarellus striatus</i>		The European list was derived from URMO which was primarily based on Müller's electronic catalogue (1993). Stock, Child, and Bamber contributed many corrections and additions.
Subfamily Lohmannellinae		References
<i>Lohmannella</i>		Müller, H.-G., 1993. <i>Bibliography of the Pycnogonida</i> . Müller Wissenschafts Verlag, Wetzlar. [on floppy disk]
<i>falcata</i> (Hodge, 1863)	A M	
<i>kervillei</i> (Trouessart, 1894)	A	Class Pycnogonida
<i>multisetosa</i> Bartsch, 1986	A M	Family Ammoteidae
<i>nudipes</i> Bartsch, 1977	A	<i>Achelia</i>
<i>norvegica</i> Viets, 1927	A	<i>anomala</i> Arnaud, 1974
<i>rustica</i> Bartsch, 1977	A	<i>borealis</i> (Schimkewitsch, 1895)
<i>stammeri</i> Viets, 1939	M	<i>echinata</i> Hodge, 1864
<i>steueri</i> subsp. <i>steueri</i> Viets, 1940	M	<i>laevis</i> Hodge, 1864
<i>steueri</i> subsp. <i>reducta</i> Bartsch, 1986	M	<i>langi</i> (Dohrn, 1881)
<i>Scaptognathus</i>	A M	<i>neotenica</i> Krapp, 1986
<i>hallezi</i> Trouessart, 1894		<i>simplex</i> Giltay, 1934
= <i>S. neretinus</i> Morselli & Mari 1982		<i>turba</i> Stock, 1990
<i>minutus</i> Bartsch, 1973	A	<i>vulgaris</i> (Costa, 1861)
<i>sabularius</i> André, 1961	M	= <i>A. armata</i>
<i>tereninus</i> Bartsch, 1986	M	<i>Ammothea</i>
<i>tridens</i> Trouessart, 1889	A M	<i>hilgendorfi</i> (Boehm, 1879)
<i>trouessarti</i> Halbert, 1915	A	= <i>A. marginatum</i> (Cole, 1904)
Subfamily Rhombognathinae		<i>Ammothella</i>
<i>Isobactrus</i>		<i>appendiculata</i> (Dohrn, 1881)
<i>cryptorhynchus</i> (Trouessart, 1901)	A	<i>biunguiculata</i> (Dohrn, 1881)
<i>hartmanni</i> Bartsch, 1972	A	<i>gibraltaricensis</i> Munilla, 1993
<i>levis</i> (Viets, 1927)	A	<i>hispida</i> (Hodge, 1864)
<i>peregrinus</i> Bartsch, 1998	M	
<i>pulchellus</i> Bartsch, 1975	A	
<i>rugosus</i> Bartsch, 1975	A	
<i>setosus</i> (Lohmann, 1889)	A M	
<i>ungulatus</i> Bartsch, 1975	A M	
<i>uniscutatus</i> (Viets, 1939)	A M	

<i>longioculata</i> (Faragiana, 1940)	M	= <i>C. spei</i> Pushkin, 1970	
<i>longipes</i> (Hodge, 1864)	A M	<i>minor</i> Schimkewitsch, 1893	A
= <i>A. hispida</i> (Hodge, 1864)		<i>minuta</i> Hoek, 1881	A
= <i>A. magnirostris</i> (Dohrn, 1881)		<i>proboscidea</i> (Sabine)	A
<i>tubicen</i> Stock, 1978	A	<i>Hedgpethia</i>	
<i>uniunguiculata</i> (Dohrn, 1881)	AM	<i>atlantica</i> (Stock, 1970)	A
Ascorhynchus		Family Endeidae	
<i>abyssi</i> Sars, 1877	A	<i>Endeis</i>	
<i>arenicolum</i> (Dohrn, 1881)	M	<i>charybdea</i> (Dohrn, 1881)	A M
<i>armatus</i> (Wilson, 1881)	A	<i>mollis</i> (Carpenter, 1904)	A
<i>castelli</i> (Dohrn, 1881)	AM	<i>spinosa</i> (Montagu, 1808)	A M
<i>endoparasiticus</i> Arnaud, 1978	A	<i>straughani</i> Clark, 1970	A
<i>pudicus</i> Stock, 1970	AM		
<i>simile</i> Fage, 1942	M		
<i>turritus</i> Stock, 1978	A	Family Nymphonidae	
Cilunculus		<i>Boreonymphon</i>	
<i>alcicornis</i> Stock, 1978	A	<i>abyssorum</i> (Norman, 1873)	A
<i>batteneae</i> Bamber & Thurston, 1993	A	<i>compactum</i> Just, 1972	A
<i>europaeus</i> Stock, 1978	A	<i>ossiansarsi</i> Knaben, 1972	A
<i>tubicinus</i> Child, 1982	A	<i>robustum</i> (Bell, 1855)	A
Eurycyde		<i>Heiteronymphon</i>	
<i>hispida</i> (Kroyer, 1844)	A	<i>ponsitor</i> Child, 1982	A
Nymphonella		<i>Nymphon</i>	
<i>tapetis</i> Ohshima, 1927	M	<i>akanthochoeros</i> Bamber & Thurston, 1995	A
Paranymphon		<i>brevitarse</i> Kroyer, 1838	A
<i>spinosum</i> Caullery, 1896	AM	= <i>N. microcollis</i> Hilton, 1942	
Sericosura		<i>caldarium</i> Stock, 1987	A M
<i>heteroscela</i> Child & Ségonzac, 1996	A	<i>chainae</i> Child, 1982	A
Tanystylum		<i>cognatum</i> Loman, 1928	A
<i>conirostre</i> (Dohrn, 1881)	AM	<i>elegans</i> Hansen, 1887	A
<i>orbiculare</i> Wilson, 1878	AM	<i>gracile</i> Leach, 1814	A M
Trygaeus		<i>grossipes</i> (Fabricius, 1780)	A
<i>communis</i> Dohrn, 1881	M	<i>gruveli</i> Bouvier, 1910	A
Family Austrodecidae		<i>hirtipes</i> Bell, 1853	A
Austrodecus		<i>hirtum</i> Kroyer, 1844	A
<i>conifer</i> Stock, 1991	A	<i>inerme</i> Fage, 1956	A
Pantopipetta		<i>laterospinum</i> Stock, 1963	A
<i>armoricana</i> Stock, 1978	A	<i>leptocheles</i> Sars, 1888	A
<i>longituberculata</i> (Turpaeva, 1955)	A	<i>longimanum</i> Sars, 1888	A
= <i>P. brevicauda</i> Stock, 1963		<i>longitarse</i> Kroyer, 1844	A
Family Callipallenidae		<i>macronyx</i> Sars, 1877	A
Callipallene		<i>megalops</i> Sars, 1877	A
<i>acribica</i> Krapp, 1975	M	<i>micronyx</i> Sars, 1888	A
<i>acus</i> (Meinert, 1898)	A	<i>microrhynchum</i> Sars, 1888	A
<i>brevirostris</i> (Johnston, 1837)	AM	<i>parasiticum</i> Merton, 1906	M
<i>emacinata</i> (Dohrn, 1881)	M	<i>procerum</i> Hoek, 1881	A
<i>phantoma</i> (Dohrn, 1881)	AM	<i>puellula</i> Krapp, 973	M
<i>producta</i> (Sars, 1888)	AM	<i>rubrum</i> Hodge, 1865	A
<i>spectrum</i> (Dohrn, 1881)	AM	<i>schimkewitschi</i> Losina-Losinsky, 1929	A
<i>tiberi</i> (Dohrn, 1881)	AM	<i>serratum</i> Sars, 1879	A
Hannonia		<i>sluiteri</i> Hoek, 1901	A
<i>echinata</i> Stock, 1990	A	<i>spinosissimum</i> (Norman, 1894)	A
<i>stocki</i> Munilla, 1993	A	<i>stenocheir</i> Norman, 1908	A
Neopallene		<i>stroemi</i> Kroyer, 1844	A
<i>azorensis</i> Arnaud, 1974	A	<i>tenellum</i> (Sars, 1888)	A
<i>campanellae</i> Dohrn, 1881	M	<i>tubiferum</i> Stock, 1978	A
Pigromoritus		Family Phoxichiliidae	
<i>timsanus</i> Calman, 1927	A	<i>Anoplodactylus</i>	
Pseudopallene		<i>angulatus</i> (Dohrn, 1881)	A M
<i>brevicollis</i> (Sars, 1891)	A	<i>arnaudae</i> Stock, 1978	A
<i>circularis</i> (Goodsir, 1842)	A	<i>californicus</i> Hall, 1912	A M
<i>longicollis</i> (Sars, 1888)	A	= <i>A. projectus</i> Hilton, 1942	
<i>malleolata</i> (Sars, 1879)	A	<i>compositus</i> Chirmens <i>et al.</i> , 1991	M
<i>spinipes</i> (Fabricius, 1780)	A	<i>digitatus</i> (Bohm, 1879)	A
Family Colossendeidae		<i>maritimus</i> Hodgson, 1914	A
Colossendeis		<i>massiliensis</i> Bouvier, 1916	M
<i>angusta</i> Sars, 1877	A M	<i>oculatus</i> Carpenter, 1905	A
= <i>C. gracilis</i> Hoek, 1881		<i>petiolatus</i> (Kroyer, 1844)	A M
<i>arcuata</i> A. Milne-Edwards, 1885	A	<i>pygmaeus</i> (Hodge, 1864)	A M
<i>bicincta</i> Schimkewitsch, 1893	A	<i>robustus</i> (Dohrn, 1881)	A M
<i>clavata</i> Meinert, 1899	A	<i>stocki</i> Bacescu, 1958	M
<i>colossea</i> Wilson, 1881	AM	<i>trispinosus</i> Stock, 1951	A
<i>cucurbita</i> Cole, 1909	A	<i>typhlops</i> Sars, 1888	A
<i>macerrima</i> Wilson, 1881	AM	= <i>A. pelagicus</i> Flynn, 1928	A M
		<i>virescens</i> (Hodge, 1864)	
		<i>Pallenopsis</i>	

<i>arctica</i> Stock, 1956	A
<i>calcanea</i> Stephensen, 1933	AM
<i>guineensis</i> Stock, 1975	A
<i>juttingae</i> Stock, 1964	A
<i>longirostris</i> Wilson, 1881	A
<i>mollissima</i> (Hoek, 1881)	AM
<i>oscitans</i> (Hoek, 1881)	A
<i>scoparia</i> Fage, 1956	AM
<i>tritonis</i> Hoek, 1883	A
<i>tydemani</i> Loman, 1908	A
<i>Phoxichilidium</i>	
<i>femoratum</i> (Rathke, 1799)	A
= <i>P. tubulariae</i> Lebour, 1945	
Family Pycnogonidae	
<i>Pycnogonum</i>	
<i>crassirostrum</i> Sars, 1888	A
<i>littorale</i> (Strom, 1762)	AM
<i>nodulosum</i> Dohrn, 1881	AM
<i>plumipes</i> Stock, 1960	M
<i>pusillum</i> Dohrn, 1881	AM
Family Rhynchothoracidae	
<i>Rhynchothorax</i>	
<i>alcicornis</i> Krapp, 1973	M
<i>mediterraneus</i> Costa, 1861	AM
<i>monnioti</i> Arnaud, 1974	AM
<i>philopsammus</i> Hedgpeth, 1951	M
= <i>R. anophthalmus</i> Arnaud, 1972	
<i>voxorinus</i> Stock, 1966	M

CRUSTACEA

Remipedia

Compiled by Geoff Boxshall

This class of blind centipede-like predators was first discovered in 1981. Only one of the 12 known species, all of which occur in anchialine caves, occurs in Europe (Lanzarote). The classification follows Schram, Yager & Emerson (1986).

References

Schram, F.R., Yager, J. & Emerson, M.J. 1986. Remipedia. Part 1. Systematics. *Mem. San Diego Soc. Nat. Hist.* 15: 60pp.

Class Remipedia

Order Nectiopoda

Family Speleonectidae

Speleonectes

ondinae (Garcia-Valdecasas, 1984)

Branchiura

Compiled by Geoff Boxshall

The branchiuran fish lice are primarily parasites of freshwater fishes but a small number of species is known from marine fishes in European waters. The data are drawn from Yamaguti (1963) and from records kept in the Natural History Museum, London.

References

Yamaguti, S. 1963. *Parasitic Copepoda and Branchiura of fishes*: i-x, 1-1104. Interscience Publishers, New York.

Subclass Branchiura

Family Argulidae

Argulus

arcassonensis Cuenot, 1912

purpureus (Risso, 1826)

Cladocera (Branchiopoda)

Compiled by Geoff Boxshall

The water fleas belong within the crustacean class Branchiopoda but there is considerable debate concerning the higher classification of the class (see Fryer, 1987). The taxon Cladocera may not be monophyletic, representing instead a level of organisation within the diplostracan branchiopods. Most cladocerans are freshwater but a few, mainly belonging to the infraorder Onychopoda, are marine or occur in brackish waters, such as the Baltic Sea. The data are compiled from Rammner (1939) and Alonso (1996).

References

Alonso, M. 1996. Crustacea Branchiopoda. In, *Fauna Iberica*, Museo Nacional de Ciencias Naturales (CSIC), Madrid, 7: 486pp.

Fryer, G. 1987. A new classification of the branchiopod Crustea. *Zool. J. Linn. Soc.* 91: 357-383.

Rammner, W. 1939. Cladocera. *Fiches d'Identification du Zooplankton. Conseil International pour l'Exploration de la Mer*, Sheet 3. 4pp.

Order Cladocera

Suborder Anomopoda

Family Bosminidae

Bosmina

coregoni subsp. *maritima* P.E. Müller, 1867

Suborder Ctenopoda**Family Sididae**

Penilia
 avirostris Dana, 1849

Suborder Onychopoda**Family Podonidae**

Evadne
 nordmanni Lovén, 1836
 spinifera P.E. Müller, 1867

Pleopsis
 polyphemoides (Leuckart, 1859)

Podon
 intermedius Lilljeborg, 1853
 leuckarti G.O. Sars, 1862

Pseudovedadne
 tergestina (Claus, 1877)

Ostracoda

Compiled by David J. Horne, Alasdair Bruce and John E. Whittaker

The Ostracoda list attempts to include representatives of all Ostracoda (orders Podocopida, Myodocopida and Platycopida) found living in marine or brackish waters in the study area. We estimate that the list is now approximately 80% complete and will continue to work to update and complete it. We have attempted to achieve an even geographical coverage; however, the focus of our attention has been on shelf and coastal waters, and we are aware of a considerable body of literature on deep-sea benthonic ostracods from which species lists remain to be abstracted. For the time being we have chosen not to add geographical distribution information to the list, but we are currently developing a GIS application which will enable distribution maps to be produced. A limitation of the list is uncertainty as to whether many of the taxa are actually living today in the areas where they have been recorded, since many have been identified only on the basis of shells (without "soft parts") found in Recent sediments and may therefore be fossil or "subfossil"; we hope to be able to indicate, eventually, which taxa can be verified as "living".

The following Web site offers further information on Ostracoda:

<http://www.gre.ac.uk/directory/earthsci/ostracod/Index.htm>.

References

- Angel, M. 1999 (personal communication). Unpublished checklist of myodocopid Ostracoda.
- Aranki, J.F., 1987. Marine Lower Pliocene Ostracoda of southern Spain with notes on the Recent fauna. *Bull. geol. Inst. Univ. Uppsala*, NS 13, 1-144.
- Athersuch, J., Horne, D.J. & Whittaker, J.E., 1989. Marine and Brackish Water Ostracods. *Synopses of the British Fauna* (New Series) No. 43, 1-343, pls 1-8.
- Bamber, R., Horne, D.J. & Whittaker, J.E. *A checklist of Ostracoda (marine and nonmarine) from British waters, compiled at the Natural History Museum, London.* (Unpublished).
- Bless, M.J.M., 1973. Note on the ecology of Recent ostracods in the Ria de Arousa (Galicia, N.W. Spain). *Zoologische Mededelingen*, 46 (10), 129-147.
- Bonaduce, G. & Masoli, M., 1968. Benthic marine Ostracoda from Malta. *Pubbl. Staz. zool. Napoli*, 38, 47-56.
- Bonaduce, G. & Masoli, M., 1968. Marine Ostracoda of Tunisia. *Pubbl. Staz. zool. Napoli*, 36, 458-470.
- Bonaduce, G. & Pugliese, N., 1973. Ostracoda from Lybia. *Pubbl. Staz. zool. Napoli*, 39, 129-135.
- Bonaduce, G., Ciampo, G. & Masoli, M., 1976. Distribution of Ostracoda in the Adriatic Sea. *Pubbl. Staz. zool. Napoli*, 40 (vol. for 1975) suppl., 1-304, pls 1-73.
- Elofson, O., 1938. Neue und wenig bekannte Cytheriden von der Schwedischen Westküste. *Ark. Zool.*, 30A (21), 1-22, figs 1-33.
- Elofson, O., 1941. Zur Kenntnis der marinen Ostracoden Schwedens mit besonderer Berücksichtigung des Skageraks. *Zool. Bidr. Upps.*, 19, 217-534, figs 1-52.
- Frenzel, P., 1996. Rezente Faunenverteilung in den Oberflächensedimenten des Greifswalder Boddens (südliche Ostsee) unter besonderer Berücksichtigung der Ostrakoden (Crustacea). *Senckenbergiana maritima*, 27, 11-31.
- Hartmann, G., 1953a. Les ostracodes de la zone d'algues de l'eulitoral de Banyuls. *Vie Milieu*, 4, 608-612.
- Hartmann, G., 1953b. Ostracodes des eaux souterraines littorales de la Méditerranée et de Majorque. *Vie Milieu*, 4, 238-253.
- Hartmann, G., 1953c. Les ostracodes du sable à *Amphioxus* de Banyuls. *Vie Milieu*, 4, 648-658.
- Klie, W., 1938. Krebstiere oder Crustacea 3: Ostracoda, Muschelkrebs. In: Dahl, F., *Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise*, 34, 1-230, figs 1-786.
- Kornicker, L.S., 1974. Revision of the Cypridinacea of the Gulf of Naples (Ostracoda). *Smithsonian Contr. Zool.*, 178, 1-64.
- Lachanel, A.M., 1989. Écologie des ostracodes du domaine Méditerranéen. Application au Golfe de Gabès (Tunisie Orientale) les variations du niveau marin depuis 30 000 ans. *Docum. Lab. Géol. Lyon*, 108, 1-239, pls 1-7.
- Lerner-Seggev, R., 1964. Preliminary notes on the Ostracoda of the Mediterranean coast of Israel. *Israel Journal of Zoology*, 13, 145-176.
- Maddocks, R.F., 1969. Revision of Recent Bairdiidae (Ostracoda). *United States National Museum Bulletin* 295, 1-126, figs 1-63.
- Maddocks, R.F., 1990. Living and fossil Macrocyprididae (Ostracoda). *University of Kansas Paleontological Contributions*, Monograph 2, 1-404, figs 1-80, pls 1-114.
- Malz, H. & Jellinek, T., 1994. Podocopida Tiefsee-Ostracoden aus Kastengreifer-Proben im östlichen Mittelmeer ("Meteor"-Fahrt 25/leg 1: Ionisches bis Levantinisches Becken). *Senckenbergiana lethaea*, 74 (1/2), 9-32.
- Marinov, T., 1962. On the ostracod fauna of the Western Black Sea coast. *Izv. tsent. nauchnoizsl. Inst. Rib. Varna*, 2, 81-108 (in Russian).
- Neale, J.W. & Howe, H., 1975. The marine Ostracoda of Russian Harbour, Novaya Zemlya and other high latitude faunas. In: Swain, F.M. (ed.), Biology and palaeobiology of Ostracoda. *Bull. Am. Paleont.*, 65, 381-431, pls 1-7.
- Neale, J.W., 1959. Normanicythere gen. nov. (Pleistocene and Recent) and the division of the ostracod family Trachyleberidae. *Palaeontology*, 2 (1), 72-93.
- Puri, H.S., Bonaduce, G. & Malloy, J., 1964. Ecology of the Gulf of Naples. *Pubbl. Staz. zool. Napoli*, 33 suppl., 87-199.
- Reys, S., 1961. Recherches sur la systématique et la distribution des ostracodes de la région de Marseille. *Rec. Trav. St. Mar. End.*, Bull. 22 (31), 53-109.
- Rome, D.R., 1964. Ostracodes des environs de Monaco, leur distribution en profondeur, nature des fonds marins explorés. *Pubbl. Staz. zool. Napoli*, 33 suppl., 200-212.
- Rudjakov, J.A., 1962. Ostracoda of the shallows of the Kandalaksha Bay (White Sea). *Trudy belom. biol. Sta. M.G.U.*, 1, 130-142 (in Russian).
- Sars, G.O., 1922. *An Account of the Crustacea of Norway*, Volume 9 – Ostracoda. Parts 1, 2 Cypridinidae, Conchoecidae, Polycopidae (part), 1-32, pls 1-16. Bergen Museum.
- Sars, G.O., 1923. *An Account of the Crustacea of Norway*, Volume 9 – Ostracoda. Parts 3, 4 Polycopidae (concluded), Cytherellidae, Cypridae (part), 33-72, pls 17-32. Bergen Museum.

- Sars, G.O., 1925a. *An Account of the Crustacea of Norway*, Volume 9 – *Ostracoda*. Parts 9, 10 *Cypridae (concluded)*, *Cytheridae (part)*, 137-176, pls 65-80. Bergen Museum.
- Sars, G.O., 1925b. *An Account of the Crustacea of Norway*, Volume 9 – *Ostracoda*. Parts 11, 12 *Cytheridae (continued)*, 177-208, pls 81-96. Bergen Museum.
- Sars, G.O., 1926. *An Account of the Crustacea of Norway*, Volume 9 – *Ostracoda*. Parts 13, 14 *Cytheridae (continued)*, 209-240, pls 97-112. Bergen Museum.
- Sars, G.O., 1928. *An Account of the Crustacea of Norway*, Volume 9 – *Ostracoda*. Parts 15, 16 *Cytheridae (concluded)*, 241-277, pls 113-119. Bergen Museum.
- Skaumal, U., 1977. *Zur Kenntnis der Ostracoden des felslitorals von Helgoland*. Dissertation, University of Hamburg, 99 + xii pp, 3 pls.
- Uffenorde, H., 1972. Okologie und jahreszeitliche Verteilung rezenten benthonischer Ostracoden des Limski kanal bei Rovinj (nördliche Adria). *Göttinger Arb. Geol. Paläont.*, 13, 1-121, pls 1-12.
- Vos, A.P.C. de, 1957. Liste annotée des ostracodes marins des environs de Roscoff. *Arch. Zool. Exp. gén.*, 95, 1074, pls 1-29.
- Wagner, C.W., 1957. *Sur les Ostracodes du Quaternaire récent des Pays-Bas et leur utilisation dans l'étude géologique des dépôts Holocènes*. 158 pp, 50 pls, Mouton & Co., The Hague.
- Whatley, R. & Eynon, M.P., 1996. Four new Arctic deepwater ostracod species from East Greenland. In: Keen, M.C. (ed.), *Proceedings of 2nd European Ostracologists' Meeting, Glasgow 1993*, 195-200. British Micropalaeontological Society, London.
- Yassini, I., 1979. The littoral system ostracodes from the Bay of Bou-Ismail, Algiers, Algeria. *Revta Esp. Micropal.*, 11 (3), 353-416.
- ### Subclass Ostracoda
- #### Order Myodocopida
- ##### Family Cylindroleberidae
- Parasterope muelleri* (Skogsberg, 1920)
- Asterope abyssicola* Sars, 1870
mariae (Baird, 1850)
muelleri Skogsberg, 1920
teres (Norman, 1861)
- Cycloleberis lobiancoi* (G.W. Mueller, 1894)
- Parasterope aberrata* (Skogsberg, 1920)
- Polypleberis mackenziei* Kornicker, 1974
- Prionotoleberis gyion* Kornicker, 1974
- Synasterope norvegica* (Sars, 1869)
- ##### Family Cypridinidae
- Azygocypridina imperialis* (Stebbing, 1901)
- Cypridina (Vargula) norvegica* (Baird, 1860)
- Cypridina*
megalops Sars, 1872
- Doloria acorensis* (Granata & Caporiacco, 1949)
sarsi Kornicker, 1987
- Gigantocypris danae* Poulsen, 1962
dracontovalis Cannon, 1940
muelleri Skogsberg, 1920
- Macrocypidina castanea* (Brady, 1897)
- Philomedes (Philomedes) lilljeborgi* (Sars, 1866)
- Philomedes aspera* G.W. Mueller, 1894
- brenda* (Baird, 1850)
globosus (Lilljeborg, 1853)
interpuncta (Baird, 1850)
levis G.W. Mueller, 1894
lilljeborgi (Sars, 1866)
macandrei (Baird, 1850)
- Pseudophilomedes angulata* G.W. Mueller, 1894
foveolata G.W. Mueller, 1894
- Skogsbergia costai* Kornicker, 1974
megalops (Sars, 1872)
squamosa (G.W. Mueller, 1894)
- Vargula mediterranea* (Costa, 1845)
norvegica (Baird, 1860)
- ##### Family Halocyprididae
- Alacia*
alata subsp. *alata* (G.W. Mueller, 1906)
leptostrix (G.W. Mueller, 1906)
valdiviae (G.W. Mueller, 1906)
- Archiconchoecia*
aff. cucullata (Brady, 1902)
bifurcata Deevey, 1978
bimucronata Deevey, 1978
bispicula Deevey, 1978
cucullata (Brady, 1902)
cuneata G.W. Mueller, 1908
fabiformis Deevey, 1978
falcata Deevey, 1978
gastrodes Deevey, 1978
longiseta Deevey, 1978
pilosa Deevey, 1978
simula Deevey, 1982
striata G.W. Mueller, 1894
ventricosa G.W. Mueller, 1906
versicula Deevey, 1978
- Bathyconchoecia*
arctica Angel, 1976
caini Ellis, 1989
darcythompsoni (Scott, 1909)
deeveyae Kornicker, 1970
diacantha Deevey, 1975
foveolata Deevey, 1968
galerita Deevey, 1968
hardingae Deevey, 1975
kornickeri Deevey, 1968
laqueata Deevey, 1968
latrostris Poulsen, 1972
longispinata Ellis, 1987
nodosa Poulsen, 1972
paulula Deevey, 1968
sagittarius Deevey, 1968
septemspinosa Angel, 1970
subrufa Angel, 1970
- Boroecia*
borealis (Sars, 1866)
maxima (Brady & Norman, 1896)
- Conchoecetta*
acuminata Claus, 1890
giesbrechti (G.W. Mueller, 1906)
- Conchoecia*
hyalophyllum Claus, 1890
lophura G.W. Mueller, 1906
macrocheira G.W. Mueller, 1906
magna Claus, 1874
parvidentata G.W. Mueller 906
subarcuata Claus, 1890
- Conchoecilla*
daphnoides subsp. *daphnoides* Claus, 1890
daphnoides subsp. *minor* (G.W. Mueller, 1906)
- Conchoecissa*
ametra (G.W. Mueller, 1906)
imbricata (Brady, 1880)
plinthina (G.W. Mueller, 1906)

- symmetrica* (G.W. Mueller, 1906)
Discoconchoecia
discophora subsp. *discophora* (G.W. Mueller, 1906)
elegans (Sars, 1866)
Discoconchoecia
elegans (Sars, 1866)
Euconchoecia
aculeata (Scott, 1894)
chierchiai G.W. Mueller, 1890
Felia
bicornis (G.W. Mueller, 1906)
cornuta (G.W. Mueller, 1906)
Gaussicia
edentata (G.W. Mueller, 1906)
gaussi (G.W. Mueller, 1908)
incisa (G.W. Mueller, 1906)
subdentata (Gooday, 1976)
Guassicia
gaussi (G.W. Mueller, 1908)
Halocypris
globosa Claus, 1874
Halocypris
inflata Dana, 1849
pelagica Claus, 1890
Loricoecia
acutimarginata Chavturi, 1977
ctenophora (G.W. Mueller, 1906)
loricata (Claus, 1894)
Macroconchoecia
macroreticulata (Ellis, 1984)
reticulata (G.W. Mueller, 1906)
spinoreticulata (Ellis, 1984)
Metaconchoecia
acuta (Gooday, 1981)
arcuata (Deevey, 1978)
discoveryi (Gooday, 1981)
fowleri (Gooday, 1981)
glandulosa (G.W. Mueller, 1906)
inflata (Gooday, 1981)
kyrtophora (G.W. Mueller, 1906)
macromma (G.W. Mueller, 1906)
nasotuberculata (G.W. Mueller, 1906)
obtusa (Gooday, 1981)
pusilla (G.W. Mueller, 1906)
rotundata (G.W. Mueller, 1906)
skogsbergi (Iles, 1953)
subinflata (Gooday, 1981)
teretivalvata (Iles, 1953)
wolfi (Gooday, 1981)
Metaonchoecia
fowleri (Gooday, 1981)
Mikroconchoecia
acuticosta (G.W. Mueller, 1906)
curta (Lubbock, 1860)
curta (Lubbock, 1860)
echinulata Claus, 1891
stigmatica (G.W. Mueller, 1906)
Mollacia
eltaninae (Deevey, 1978)
kampta (G.W. Mueller, 1906)
mollis (G.W. Mueller, 1906)
tyloda (G.W. Mueller, 1906)
Oblusoecia
obtusata (Sars, 1866)
Orthoconchoecia
atlantica (Lubbock, 1856)
bispinosa (Claus, 1890)
haddoni (Brady & Norman, 1896)
secernenda (Vavra, 1906)
Paraconchoecia
aequisepta (G.W. Mueller, 1906)
allotherium (G.W. Mueller, 1906)
cophopyga (G.W. Mueller, 1906)
dasyophtalma (G.W. Mueller, 1906)
dentata (G.W. Mueller, 1906)
- echinata* (G.W. Mueller, 1906)
hirsuta (G.W. Mueller, 1906)
inermis Claus, 1890
mamillata (G.W. Mueller, 1906)
mesadenia (Ellis, 1985)
nanomamillata (Deevey & Brooks, 1980)
oblonga Claus, 1890
spinifera Claus, 1890
spinifera Claus, 1891
vitfazi (Rudjakov, 1962)
Paramollacia
dichotoma (G.W. Mueller, 1906)
major (G.W. Mueller, 1906)
plactolycos (G.W. Mueller, 1906)
rhynchena (G.W. Mueller, 1906)
Porroecia (*Spinoezia*)
hystrix (Angel & Ellis, 1981)
hystrix (Angel & Ellis, 1987)
Porroecia
parthenoda (G.W. Mueller, 1906)
orrecta subsp. *adriatica* (Gooday & Angel, 1977)
orrecta subsp. *orrecta* (Claus, 1890)
pseudoparthenoda (Angel, 1972)
spinirostris (Claus, 1874)
Proceroezia
brachyaskos (G.W. Mueller, 1906)
brachyaskos G.W. Mueller, 1906
convexa (Deevey, 1977)
macroprocera (Angel, 1971)
microprocera (Angel, 1971)
procera (G.W. Mueller, 1894)
Pseudoconchoecia
concentrica (G.W. Mueller, 1906)
- Family Philomedidae**
- Angulorostrum*
segonzaci Kornicker, 1981
- Euphilomedes*
asper G.W. Mueller, 1894
interpuncta (Baird, 1850)
sinister Kornicker, 1974
- Streptoleberis*
rectirostris Brady & Norman, 1896
- Family Polycopidae**
- Metapolycope*
compressa (Brady & Robertson, 1869)
- Polycope*
arenicola Hartmann, 1954
areolata Sars, 1923
clathrata Sars, 1923
demulderi Sissingh, 1972
dentata Brady, 1868
dimorpha Hartmann, 1954
dispar G.W. Mueller, 1894
fragilis G.W. Mueller, 1894
frequens G.W. Mueller, 1894
longipes Hartmann, 1954
maculata G.W. Mueller, 1894
orbicularis Sars, 1866
parareticulata Bonaduce, Ciampo & Masoli, 1976
punctata Sars, 1869
pustulata Sars, 1890
reticulata G.W. Mueller, 1894
rostrata (G.W. Mueller, 1894)
striata G.W. Mueller, 1894
sublaevis Sars, 1923
teneriffa Hartmann, 1959
tholiformis Bonaduce, Ciampo & Masoli, 1976
truncatula Bonaduce, Ciampo & Masoli, 1976
tuberosa G.W. Mueller, 1894
vasiensis Sissingh, 1972
- Polycoptis*
quadridentata Bonaduce, Ciampo & Masoli, 1976
serrata G.W. Mueller, 1894
- Rutiderma*
compressa Brady & Norman

- Family Sarsiellidae**
- Eusarsiella*
 - zostericola* (Cushman, 1906)
 - Sarsiella*
 - capsula* Norman, 1869
 - levis* G.W. Mueller, 1894
 - neapolis* Kornicker, 1974
- Order Platycopida**
- Family Cytherellidae**
- Cytherella*
 - abyssorum* Sars, 1866
 - alvearium* Bonaduce, Ciampo & Masoli, 1976
 - circumpunctata* Ciampo, 1976
 - pori* Lerner-Seggev, 1964
 - serrulata* Brady & Norman, 1896
 - vandenboldi* Sissingh, 1972
 - vulgata* Ruggieri, 1962
 - Cytherelloidea*
 - beckmanni* Barbeito-Gonzalez, 1971
 - sordida* (G.W. Mueller, 1894)
- Order Podocopida**
- Superfamily Bairdioidea**
- Family Bairdiidae**
- Neonesidea*
 - corpulenta* (G.W. Mueller, 1894)
 - decipiens* (G.W. Mueller, 1894)
 - formosa* (Brady, 1868)
 - frequens* (G.W. Mueller, 1894)
 - incognita* (Lerner-Seggev, 1964)
 - inflata* (Norman, 1862)
 - longevaginata* (G.W. Mueller, 1894)
 - mediterranea* (G.W. Mueller, 1894)
 - minor* (G.W. Mueller, 1894)
 - obscura* (G.W. Mueller, 1894)
 - roquebrunensis* (Rome, 1942)
 - Paranesidea?*
 - conformis* (Terquem, 1878)
 - Triebelina*
 - reticulata* (G.W. Mueller, 1894)
 - rariplata* (G.W. Mueller, 1894)
- Family Bythocypriidae**
- Bythocyrpis*
 - bosquetiana* (Brady, 1866)
 - obtusata* (Sars, 1866)
- Superfamily Cypridoidea**
- Family Candonidae**
- Aglaocypris*
 - complanata* (Brady & Robertson, 1869)
 - rara* (G.W. Mueller, 1894)
 - Paracypris*
 - polita* Sars, 1866
- Family Bythocytheridae**
- Bathynocythere*
 - styx* Malz & Jellinek, 1994
 - Bythoceratina*
 - reticulata* Bonaduce, Ciampo & Masoli, 1976
 - Bythocythere*
 - bradleyi* Athersuch, Horne & Whittaker, 1983
 - bradyi* Sars, 1926
 - constricta* Sars, 1866
 - dromedaria* Sars, 1866
 - insignis* Sars, 1869
 - intermedia* Elofson, 1938
 - minima* Bonaduce, Ciampo & Masoli, 1976
 - robinsoni* Athersuch, Horne & Whittaker, 1983
 - turgida* Sars, 1866
 - zetlandica* Athersuch, Horne & Whittaker, 1983
 - Dipseucythere*
 - armata* (Bonaduce *et al.*, 1980)
 - kraspedon* Malz & Jellinek, 1994
 - mediterranea* (Bonaduce *et al.*, 1980)
- Family Sarsiellidae**
- Jonesia*
 - acuminata* (Sars, 1866)
 - Monoceratina*
 - mediterranea* Sissingh, 1971
 - Pseudocythere*
 - britannica* Horne, 1986
 - caudata* Sars, 1866
 - Rostrocythere*
 - hastata* (Bonaduce *et al.*, 1980)
 - Sclerochilus*
 - abbreviatus* Brady & Robertson, 1869
 - aequus* G.W. Mueller, 1894
 - bradyi* Rudjakov, 1962
 - contortus* (Norman, 1861)
 - gewemuelleri* Dubowsky, 1939
 - gewemuelleri* subsp. *dubowskyi* Marinov, 1962
 - hicksi* Athersuch & Horne, 1987
 - levis* G.W. Mueller, 1894
 - rudjakovi* Athersuch & Horne, 1987
 - schornikovi* Athersuch & Horne, 1987
 - truncatus* (Malcomson, 1886)
 - whatleyi* Athersuch & Horne, 1987
- Family Cuneocytheridae**
- Cuneocythere*
 - semipunctata* (Brady, 1868)
- Family Cushmanicidae**
- Pontocythere*
 - elongata* (Brady, 1868)
 - rubra* (G.W. Mueller, 1894)
 - turbida* (G.W. Mueller, 1894)
- Family Cytherettidae**
- Cytherella*
 - adriatica* Ruggieri, 1952
 - judaea* (Brady, 1868)
 - rubra* G.W. Mueller, 1894
 - subradiosa* (Roemer, 1838)
 - Loculicytheretta*
 - pavonia* (Brady, 1866)
- Family Cytheridae**
- Cobancycythere*
 - hartmanni* G.W. Mueller, 1894
 - Cythere*
 - lutea* O.F. Mueller, 1785
 - Cythereis*
 - amnicola* (Sars, 1887)
 - polygonata* Rome, 1942
 - Cytherideis*
 - foveolata* Brady, 1870
 - Eucytherura*
 - gibbera* G.W. Mueller, 1894
 - Palmenella*
 - limicola* (Norman, 1865)
 - Paracytheridea*
 - bovettensis* (Seguenza, 1880)
- Family Cytherideidae**
- Cyprideis*
 - torosa* (Jones, 1850)
 - Cytheridea*
 - angustata* (Baird, 1850)
 - muelleri* Muenster, 1938
 - neapolitana* Kollman, 1960
 - Heterocyprideis*
 - fascis* (Brady & Norman, 1889))
 - sorbyana* (Jones, 1856)
 - Paracyprideis*
 - fennica* (Hirschmann, 1909)
 - Sarsicytheridea*
 - bradii* (Norman, 1865)
 - macrolaminata* (Elofson, 1939)
 - punctillata* (Brady, 1865)
- Family Cytheromatidae**
- Cytheroma*
 - karadagiensis* Dubowsky

- latianennata* Elofson, 1938
variabilis G.W. Mueller, 1894
- Family Cytheruridae**
- Cytheropteron*
alatum Sars, 1866
angulatum Brady & Robertson, 1872
arcticum Neale & Howe, 1973
crassipinnatum Brady & Norman, 1889
depressum (Brady & Norman, 1889)
dorsocostatum Whatley & Masson, 1980
dromedaria (Sars)
garganicum Bonaduce, Ciampo & Masoli, 1976
groenlandicum Whatley & Eynon, 1996
hadriaticum Bonaduce, Ciampo & Masoli, 1976
hamatum Sars, 1869
inflatum Brady, Crosskey & Robertson, 1874
inflatum sensu Sars
inornatum Brady & Robertson, 1872
latissimum (Norman, 1865)
latum G.W. Mueller, 1894
monoceros Bonaduce, Ciampo & Masoli, 1976
nodosoalatum Neale & Howe, 1973
nodosum Brady, 1868
paralatissimum Swain, 1963
pseudoinflatum Whatley & Eynon, 1996
punctatum Brady, 1868
pyramidele (Brady, 1868)
rarum G.W. Mueller, 1894
rotundatum (G.W. Mueller, 1894)
ruggieri Pucci, 1955
scoresbyi Whatley & Eynon, 1996
subcircinatum Sars, 1866
sulcatum Bonaduce, Ciampo & Masoli, 1976
testudo Sars, 1870
venustum Bonaduce, Ciampo & Masoli, 1976
vespertilio (Reuss, 1850)
zinzulusae Bonaduce, Ciampo & Masoli, 1976
- Cytheropteron?*
infelix Bonaduce, Ciampo & Masoli, 1976
- Cytherura**
- areolata* Reys, 1961
atra Sars, 1866
bidens Klie, 1929
brevifistulosa Klie, 1936
cochlearis Klie, 1929
gibba (O.F. Mueller, 1785)
inconspicua Klie, 1934
muelleri Puri, 1963
nasuta Klie, 1934
- Eucytherura**
- alata* G.W. Mueller, 1894
amfibola Barbeito-Gonzalez, 1971
angulata G.W. Mueller, 1894
bulgarica Klie
complexa (Brady, 1866)
delineata Whatley & Eynon, 1996
mistretta Sissingh, 1972
- Hemicytherura**
- cellulosa* (Norman, 1865)
clathrata (Sars, 1866)
defiorei Ruggieri, 1953
gracilicosta Ruggieri, 1953
hoskini Horne, 1981
videns (G.W. Mueller, 1894)
- Kangarina**
- abyssicola* (G.W. Mueller, 1894)
septentrionalis Neale, 1972
- Microcytherura**
- angulosa* (Seguenza, 1880)
fulva (Brady & Robertson, 1874)
inflexa G.W. Mueller, 1894
nigrescens G.W. Mueller, 1894
- Pedicythere**
- mirabilis* Sissingh, 1975
phryne Bonaduce, Ciampo & Masoli, 1976
- Pedicythere?**
- tessellata* Bonaduce, Ciampo & Masoli, 1976
- Pseudocytherura**
- calcarata* (Seguenza, 1880)
Semicytherura
abrita Bonaduce, Ciampo & Masoli, 1976
acuminata (G.W. Mueller, 1894)
acuta (G.W. Mueller, 1912)
acuticostata (Sars, 1866)
acuticostata subsp. *ventricosa* (G.W. Mueller, 1894)
aenariensis Bonaduce, Ciampo & Masoli, 1976
affinis (Sars, 1866)
alifera Ruggieri, 1959
amorpha Bonaduce, Ciampo & Masoli, 1976
angulata (Brady, 1868)
brachyptera (Ruggieri, 1952)
concentrica (Brady, Crosskey & Robertson, 1874)
cornuta (Brady, 1868)
costata (G.W. Mueller, 1894)
cribiformis (G.W. Mueller, 1894)
diaspora Barbeito-Gonzalez, 1971
dispar (G.W. Mueller, 1894)
heinzei (Puri, 1963)
incongruens (G.W. Mueller, 1894)
inversa (Seguenza, 1880)
kaloderma Bonaduce & Pugliese, 1975
mediterranea (G.W. Mueller, 1894)
nigrescens (Baird, 1838)
occulta Bonaduce, Ciampo & Masoli, 1976
paradoxa (G.W. Mueller, 1894)
pontica (Marinov, 1962)
producta (Brady, 1868)
psila Barbeito-Gonzalez, 1971
punctata (G.W. Mueller, 1894)
quadratovolatilis (Hartmann, 1953)
quadridentata (Hartmann, 1953)
rara (G.W. Mueller, 1894)
rarecostata Bonaduce, Ciampo & Masoli, 1976
reticulata (G.W. Mueller, 1894)
robusta Bonaduce, Ciampo & Masoli, 1976
ruggieri (Pucci, 1956)
sella (Sars, 1866)
similis (Sars, 1866)
simplex (Brady & Norman, 1889)
stilifera Bonaduce, Ciampo & Masoli, 1976
striata (Sars, 1866)
sulcata (G.W. Mueller, 1894)
tela Horne & Whittaker, 1980
tergestina Masoli, 1968
trachina Bonaduce, Ciampo & Masoli, 1976
undata (Sars, 1866)
ventricosa (G.W. Mueller, 1894)
- Tetracytherura**
- angulosa* (Seguenza, 1880)
- Family Eucytheridae**
- Eucythere**
- prava* Brady & Robertson, 1869
anglica Brady, 1868
argus (Sars, 1866)
curta Ruggieri, 1975
declivis (Norman, 1865)
prava Brady & Robertson, 1869
pubera Bonaduce, Ciampo & Masoli, 1976
- Family Hemicytheridae**
- Aurila**
- arborescens* (Brady, 1865)
convexa (Baird, 1850)
cymbaeformis (Seguenza, 1883)
glyptica Barbeito-Gonzalez, 1971
interpretis Uliczny, 1969
prasina Barbeito-Gonzalez, 1971
punctata (Muenster, 1830)
speyeri (Brady, 1868)
woutersi Horne, 1986
- Baffinicythere**

- howei* Hazel, 1967
Caudites
calceolatus (Costa, 1853)
Elofsonella
concinna (Jones, 1857)
Finnmarchinella (*Barentsovia*)
barenzovensis (Mandelstam, 1957)
angulata (Sars, 1866)
logani (Brady & Crosskey, 1871)
Finnmarchinella
angulata (Sars, 1866)
finnmarchica (Sars, 1866)
Hemicythere
emarginata (Sars, 1866)
rubida (Brady, 1868)
villosa (Sars, 1866)
Heterocythereis
albomaculata (Baird, 1838)
Muellerina
abyssicola (Sars, 1866)
Normanicythere
leioderma (Norman, 1869)
Tenedocythere
prava (Baird, 1850)
Thaerocythere
crenulata (Sars, 1866)
Urocythereis
britannica Athersuch, 1977
favosa (Roemer, 1838)
flexicauda Bonaduce, Ciampo & Masoli, 1976
margaritifera (G.W. Mueller, 1894)
margaritifera subsp. *alba* Uliczny, 1969
- Family Krithidae**
Krithe
bartonensis subsp. *levantina* Lerner-Seggev, 1964
keyi Breman, 1978
papillosa (Bosquet, 1852)
praetexta (Sars, 1866)
producta Brady, 1880
reniformis (Brady, 1868)
similis (G.W. Mueller, 1894)
Parakrithe
dimorpha Bonaduce, Ciampo & Masoli, 1976
Pseudosammocythere
similis (G.W. Mueller, 1894)
- Family Leptocytheridae**
Callistocythere
adriatica Masoli, 1968
badia (Norman, 1862)
craterigera (Rome, 1942)
crispata (Brady, 1868)
crucifera (Hartmann, 1953)
diffusa (G.W. Mueller, 1894)
discrepans (G.W. Mueller, 1894)
flavidofusca (Ruggieri, 1950)
folliculosa Bonaduce, Ciampo & Masoli, 1976
foveata (Hartmann, 1953)
gilva Bonaduce, Ciampo & Masoli, 1976
intricatoides Ruggieri, 1953
littoralis (G.W. Mueller, 1894)
lobiancoi (G.W. Mueller, 1894)
mediterranea (G.W. Mueller, 1894)
murrayi Whittaker, 1978
pallida (G.W. Mueller, 1894)
praecincta Ciampo, 1976
rastrifera (Ruggieri, 1953)
taeniata (Rome, 1942)
vexata Bonaduce, Ciampo & Masoli, 1976
Cluthia
cluthae (Brady, Crosskey & Robertson, 1874)
keiji Neale, 1975
Leptocythere
bacescoi (Rome, 1942)
baltica Klie, 1929
bituberculata Bonaduce, Ciampo & Masoli, 1976
- castanea* (Sars, 1866)
ciliata Hartmann, 1957
craterigera (Rome, 1942)
crepidula Ruggieri, 1950
crucifera Hartmann, 1953
diffusa (G.W. Mueller, 1894)
sabaeformis (G.W. Mueller, 1894)
lacertosa (Hirschmann, 1912)
lagunae Hartmann, 1958
levis (G.W. Mueller, 1894)
macallana (Brady & Robertson, 1869)
macella Ruggieri, 1975
mediterranea (G.W. Mueller, 1894)
micra (Rome, 1942)
multipunctata (Seguenza, 1942)
pallida (G.W. Mueller, 1894)
pellucida (Baird, 1850)
pellucida? (Baird, 1850)
porcellanea (Brady, 1869)
psammophila Guillaume, 1976
ramosa (Rome, 1942)
rara (G.W. Mueller, 1894)
tenera (Brady, 1868)
- Family Loxoconchidae**
Cytheromorpha
elongata Hartmann, 1953
fuscata (Brady, 1869)
nana Bonaduce, Ciampo & Masoli, 1976
Elofsonia
baltica (Hirschmann, 1909)
pusilla (Brady & Robertson, 1870)
Hirschmannia
viridis (O.F. Mueller, 1785)
Loxocauda
decipiens (G.W. Mueller, 1894)
Loxoconcha
agilis Ruggieri, 1967
alata Brady, 1868
avellana (Brady, 1866)
bairdi G.W. Mueller, 1894
bonaducei Ciampo, 1971
concentrica Bonaduce, Ciampo & Masoli, 1976
dimorpha Hartmann, 1959
elliptica Brady, 1868
exagona Bonaduce, Ciampo & Masoli, 1976
fragilis Sars, 1866
geometrica Bonaduce, Ciampo & Masoli, 1976
gibberosa Terquem, 1878
globosa Schornikov, 1965
helgolandica Klie, 1929
levis G.W. Mueller, 1894
malcomsoni Horne & Robinson, 1985
mediterranea G.W. Mueller, 1894
minima G.W. Mueller, 1894
ovulata (Costa, 1853)
parallela G.W. Mueller, 1894
pontica Klie, 1937
rhomboidea (Fischer, 1855)
rubritincta Ruggieri, 1964
stellifera G.W. Mueller, 1894
subrugosa Ruggieri, 1976
Loxoconchidea
minima Bonaduce, Ciampo & Masoli, 1976
Microloxoconcha
compressa Hartmann, 1953
Nannocythere
pavo (Malcomson, 1886)
Palmococoncha
guttata (Norman, 1865)
laevata (Norman, 1865)
turbida (G.W. Mueller, 1894)
Phlyctocythere
pellucida (G.W. Mueller, 1894)
Pseudoloxoconcha
minima G.W. Mueller, 1894

- Roundstomia*
globulifera (Brady, 1868)
macchesneyi (Brady & Crosskey, 1871)
robertsoni (Brady, 1868)
- Sagmatocythere*
littoralis (G.W. Mueller, 1894)
multifora (Norman, 1865)
napoliana (Puri, 1963)
versicolor (G.W. Mueller, 1894)
- Tuberoloxoconcha*
atlantica Horne, 1989
nana (Marinov, 1962)
tuberosa (Hartmann, 1953)
- Family Microcytheridae**
- Microcythere*
dentata (G.W. Mueller, 1894)
depressa (G.W. Mueller, 1894)
gibba G.W. Mueller, 1894
helgolandica Klie, 1936
hians G.W. Mueller, 1894
inflexa G.W. Mueller, 1894
levis G.W. Mueller, 1894
longiantennata Marinov, 1962
minuta Klie, 1936
nana G.W. Mueller, 1894
navicula (Norman, 1868)
obliqua G.W. Mueller, 1894
parallela G.W. Mueller, 1894
parva Klie, 1936
rara G.W. Mueller, 1894
subterranea Hartmann, 1953
varnensis Marinov, 1962
vitreata Bonaduce, Ciampo & Masoli, 1976
- Family Neocytherideidae**
- Neocytherideis*
complicata (Ruggieri, 1953)
crenulata (Klie, 1929)
fasciata (Brady & Robertson, 1874)
soveolata (Brady, 1870)
subspiralis (Brady, Crosskey & Robertson, 1874)
subulata (Brady, 1868)
- Sahnicythere*
retroflexa (Klie, 1936)
- Family Paracytherideidae**
- Paracytheridea*
cuneiformis (Brady, 1868)
depressa G.W. Mueller, 1894
norvegica Neale, 1972
- Tuberculocythere*
batrachoides Colango & Pasini, 1980
- Family Paradoxostomatidae**
- Acetabulastoma*
hyperboreum(?) (Scott, 1899)
- Cytherois*
arenicola Klie, 1929
fischeri (Sars, 1866)
frequens G.W. Mueller, 1894
incongruens G.W. Mueller, 1894
pusilla Sars, 1928
stephanidesi Klie, 1938
succinea G.W. Mueller, 1894
triangularis Bonaduce, Masoli, Minichelli & Pugliese, 1979
vitreata (Sars, 1866)
- Paracythere*
minima G.W. Mueller, 1894
- Paracytherois*
acuminata G.W. Mueller, 1894
arcuata (Brady, 1868)
flexuosa (Brady, 1867)
mediterranea Bonaduce, Ciampo & Masoli, 1976
oblonga G.W. Mueller, 1894
producta (Brady & Norman, 1889)
rara G.W. Mueller, 1894
- striata* G.W. Mueller, 1894
sulcata G.W. Mueller, 1894
tenera (Brady, Crosskey & Robertson, 1874)
- Paradoxostoma*
abbreviatum Sars, 1866
acuminatum G.W. Mueller, 1894
amygdaloïdes (Brady, 1870)
anglorum Horne & Whittaker, 1985
angustum G.W. Mueller, 1894
arcticum Elofson, 1941
atrum G.W. Mueller, 1894
bradyi Sars, 1928
breve G.W. Mueller, 1894
caecum G.W. Mueller, 1894
chamberlaini Rome, 1939
curtum Hartmann, 1959
cylindricum G.W. Mueller, 1894
ensiforme Brady, 1868
fleetense Horne & Whittaker, 1985
hibernicum Brady, 1868
hodgii Brady, 1868
incongruens G.W. Mueller, 1894
insigne Hartmann, 1959
intermedium G.W. Mueller, 1894
maculatum G.W. Mueller, 1894
mediterraneum G.W. Mueller, 1894
nealei Horne & Whittaker, 1985
normani Brady, 1868
obliquum Sars, 1866
ornata Hartmann, 1953
parallelum G.W. Mueller, 1894
planum G.W. Mueller, 1894
ponticum Klie, 1942
porlockense Horne & Whittaker, 1985
pulchellum Sars, 1866
rarum G.W. Mueller, 1894
robinhoodi Horne & Whittaker, 1985
rostratum(?) Sars
rubrum G.W. Mueller, 1894
sarniense Brady, 1868
simile G.W. Mueller, 1894
striatum G.W. Mueller, 1894
taeniatum G.W. Mueller, 1894
tenuissimum (Norman, 1869)
trieri Horne & Whittaker, 1985
triste G.W. Mueller, 1894
variabile (Baird, 1835)
versicolor G.W. Mueller, 1894
- Redeka*
perpusilla de Vos, 1953
- Family Trachyleberididae**
- Acanthocythereis*
ascolii (Puri, 1963)
dunelmensis (Norman, 1865)
hystrix (Reuss, 1850)
ornata (G.W. Mueller, 1894)
- Basslerites*
berchoni (Brady, 1869)
teres (Brady, 1869)
- Bosquetina*
carinella (Reuss, 1957)
dentata (G.W. Mueller, 1894)
pectinata (Bosquet, 1852)
- Buntonia* (*Rectobuntonia*)
subulata Ruggieri, 1954
- Buntonia*
dertoniensis (Ruggieri, 1954)
giesbrechti (G.W. Mueller, 1894)
sublatissima (Neviani, 1906)
textilis Bonaduce, Ciampo & Masoli, 1976
- Carinocythereis*
carinata (Roemer, 1838)
whitei (Baird, 1850)
- Celtia*
quadridentata (Baird, 1850)

Costa
batei (Brady, 1866)
edwardsii (Roemer, 1838)
hamata (G.W. Mueller, 1894)
runcinata (Baird, 1850)
Echinocythereis
echinata (Sars, 1866)
laticarina (Brady, 1868)
scabra (von Muenster, 1830)
Falsocythere
maccagnoi (Ciampo, 1971)
Henryhowella
sarsi (G.W. Mueller, 1894)
Hiltermannicythere
emaciata (Brady, 1867)
rubra (G.W. Mueller, 1894)
rubra subsp. *pontica* Dubowsky
sphaerulolineata (Jones, 1856)
turbida (G.W. Mueller, 1894)
Occultocythereis
dohrni (Puri, 1963)
Pterygocythereis
ceratoptera (Bosquet, 1852)
jonesii (Baird, 1850)
mucronata (Sars, 1866)
Rectobuntonia
miranda Bonaduce, Ciampo & Masoli, 1976
Robertsonites
tuberculatus (Sars, 1866)
incertae sedis
Cistacythereis
caelatura Uliczny, 1969
Parahemingwayella
tetrapteron (Bonaduce, Ciampo & Masoli, 1976)
Parvocythere
hartmanni Marinov, 1962
Psammocythere
remanei Klie, 1936
Rabilimis
mirabilis (Brady, 1868)
septentrionalis (Brady, 1866)
Typhlocythere
ruggieri Bonaduce, Ciampo & Masoli, 1976
Verrucocythere
bulbuspinata (Uliczny, 1969)

Family Xestoleberididae

Aspidoconcha
humilis (Brady & Norman, 1889)
limnoriae de Vos, 1953
Microxestoleberis
nana G.W. Mueller, 1894
rostrata Klie, 1936
xenomis (Barbieto-Gonzalez, 1971)

Xestoleberis
aurantia (Baird, 1838)
communis G.W. Mueller, 1894
costata Hartmann, 1953
decipiens G.W. Mueller, 1894
delamarei Hartmann, 1953
depressa Sars, 1866
fuscomaculata G.W. Mueller, 1894
gigas Rome, 1942
humilis Klie, 1940
labiata Brady & Robertson, 1874
margaritea (Brady, 1866)
margaritopsis Rome, 1942
nitida (Liljeborg, 1853)
parva G.W. Mueller, 1894
pellucida G.W. Mueller, 1894
perrula Athersuch, 1976
plana G.W. Mueller, 1894
rara G.W. Mueller, 1894
rubens Whittaker, 1978
ventricosa G.W. Mueller, 1894

Superfamily Macrocypridoidea

Family Macrocyprididae

Macrocypris
sarsi (G.W. Mueller, 1912)
Macrocyprina
succinea G.W. Mueller, 1894
Macrocypris
minna (Baird, 1850)
Macromckenziea
ligustica (Bonaduce, Masoli & Pugliese, 1977)

Superfamily Pontocypridoidea

Family Pontocyprididae

Argilloecia (*Robustoargilloecia*)
acuminata (G.W. Mueller, 1894)
Argilloecia
bulbifera (G.W. Mueller, 1894)
caudata (G.W. Mueller, 1894)
conoidea Sars, 1923
cylindrica Sars, 1865
levis (G.W. Mueller, 1894)
micra Bonaduce, Ciampo & Masoli, 1976
minor (G.W. Mueller, 1894)
propinquua Brady, 1904
robusta Bonaduce, Ciampo & Masoli, 1976

Pontocypris
spinosa (G.W. Mueller, 1894)
Pontocypris
acupunctata Brady, 1868
acuta Bonaduce, Ciampo & Masoli, 1976
discrepans (G.W. Mueller, 1894)
frequens (G.W. Mueller, 1894)
hispida (Sars, 1866)
intermedia G.W. Mueller, 1894
levis G.W. Mueller, 1894
mediterranea G.W. Mueller, 1894
monstrosa G.W. Mueller, 1894
mytiloides (Norman, 1862)
obtusa (G.W. Mueller, 1894)
pallida (G.W. Mueller, 1894)
rara (G.W. Mueller, 1894)
subfuscata G.W. Mueller, 1894

Propontocypris
acuminata (G.W. Mueller, 1894)
declivis (G.W. Mueller, 1894)
dispar (G.W. Mueller, 1894)
intermedia (Brady, 1868)
pellucida (G.W. Mueller, 1894)
pirifera (G.W. Mueller, 1894)
serrata (G.W. Mueller, 1894)
setosa (G.W. Mueller, 1894)
subfuscata (G.W. Mueller, 1894)
succinea (G.W. Mueller, 1894)
trigonella (Sars, 1866)

Pentastomida

Compiled by Jacob van der Land

Only a few parasites of sea birds are known.

References

Osche, G., 1963. Die Systematische Stellung u. Phylogenie der Pentastomida: Embryologische u. Vergleichend Anatomische Studien an Reighardia sterna. *Z. Morph. ök. ser. Tier.* 52: 487-596.

Class Pentastomida

Order Cephalobaenida

Family Reighardiidae

<i>Reighardia</i>	
<i>sterna</i> (Diesing, 1864)	A M
<i>lomviae</i> Dyck, 1975	A

Mystacocarida

Compiled by Geoff Boxshall.

The mystacocarids are microscopic members of the meiofauna. Data on the two European species are taken from Boxshall & Defaye (1996).

References

- Boxshall, G.A. & Defaye, D. 1996. Classe des Mystacocarides (Mystacocarida Pennak & Zinn, 1943). In, J.Forest (Ed.) *Traité de Zoologie*, Tome VII. Fasc. II. Crustacés. pp. 409-424.

Subclass Mystacocarida

Family Derocheilocarididae

- Derocheilocaris*
remanei Delamare-Deboutteville & Chappuis, 1951
tehiyae Masry & Por, 1970

Copepoda

Compiled by Geoff Boxshall

The ordinal level classification of the copepods is based on Huys & Boxshall (1991). Data on the pelagic copepods were drawn primarily from the two excellent volumes on the diversity and distribution of pelagic copepods published by Claude Razouls (1995, 1996). These data were supplemented with the new taxa recorded or described from European waters since 1996 (Fosshagen *et al.*, in press; Jaume & Boxshall, 1995, 1996a-d, 1997a-b; Jaume *et al.*, 1999; Jaume *et al.*, in press; Markhaseva, 1996; Zagami *et al.*, 2000) and updated to include recent changes in nomenclature (Belmonte, 1998; Böttger-Schnack, 1999; Boxshall & Jaume, in press; Huys & Böttger-Schnack, 1997). Included in these papers are the many new taxa that have been discovered in anchialine cave habitats in the Mediterranean region and in the Canary Islands.

The parasitic copepods of marine fishes have been compiled primarily from the following major sources: the monograph of Kabata (1979) and the analysis of species richness of parasitic copepods of Mediterranean fishes by Raibaut, Combes & Benoit (1998). Other changes and new species are included (Boxshall, in press; Hogans, 1987). The copepods associated with marine invertebrate hosts in Northern European waters were summarised by Gotto (1993). Data for copepods from invertebrates in Southern European waters and the Mediterranean region were widely scattered through the primary literature and are probably only 90 to 95% complete. Data has been drawn from standard references (e.g. Giesbrecht, 1899; Illg & Dudley, 1961, 1965, 1980) and supplemented by information from subsequent publications (Boxshall & Defaye, 1995; Ho *et al.*, 1998; Holmes, 1998; Humes & Boxshall, 1996; Lamb *et al.*, 1996; López-González, *et al.*, 1998; López-González & Conradi, 1996; Marchenkov & Boxshall, 1995; O'Reilly, 1995). The order Monstrilloidea has been the subject of profound nomenclatural change (see Grygier, 1995 and 1997).

Additional data and corrections were provided by Myles O'Reilly and Dusan Zavodnik.

References

- Apostolov, A. & T.M. Marinov, 1988. Copepoda Harpacticoida (morski kharpaktikoidi). [Copepoda, Harpacticoida (marine harpacticoids)]. - *Fauna Bulgarii [Fauna Bulgarica]*, 18: 1-384. Sofia, Izd. B'Ig. Akad. Nauk. (Bulgarian).

- Belmonte, G. 1998. *Pteriacartia*, a new genus of Acartiidae (Calanoida, Diaptomoidea) for *Acartia josephinae* Crisafi, 1974. *J. Mar. Syst.* 15: 359-368.
- Bodin, P., 1964. Recherches sur la systématique et la distribution des Copépodes Harpacticoides des substrats meubles des environs de Marseille. - *Recl Trav. Stn mar. Endoume*, 51 (=Bull. 35): 107-183.
- Böttger-Schnack, R. 1999. Taxonomy of Oncaeidae (Copepoda, Poecilostomatoidea) from the Red Sea. - I. 11 species of *Triconia* gen nov. and a redescription of *T. similis* (Sars) from Norwegian waters. *Mitt. Hamb. Zool. Mus. Inst.* 96: 37-128.
- Boxshall, G.A. (in press) Parasitic copepods (Copepoda: Siphonostomatoida) from deep-sea and mid-water fishes. *Systematic Parasitology*
- Boxshall, G.A. & Jaume, D. (in press) Discoveries of Cave Misophrioids (Crustacea: Copepoda) Shed New Light on the Origin of Anchialine Faunas. *Zool. Anz.*
- Boxshall, G.A. & Defaye, D. 1995. Copépodes nouveaux (Siphonostomatoida: Nicothoidea) parasites de Cumacés et des Décapodes profonds. *Bull. Mus. natn. Hist. nat.* 4^e s., sect. A, 17(3-4): 283-296.
- Fosshagen, A., Boxshall, G.A. & Iliffe, T.M. (in press) The Epaeteriscidae, a cave-living family of calanoid copepods. *Sarsia*
- Giesbrecht, W. 1899. Die Asterocheriden des Golfs von Neapel und der angrenzenden Meeres-Abschnitte. *Fauna Flora Golf. Neapel* 25: i-vi, 1-217, pls 1-11.
- Gotto, V. 1993. Commensal and Parasitic Copepods associated with marine invertebrates (and whales). In: D.M. Kermack, R.S.K. Barnes & J.H. Crothers eds, *Synopses of the British Fauna* (New Series), No. 46: 1-264.
- Grygier, M.J. 1995. Annotated chronological bibliography of Monstrilloida (Crustacea: Copepoda). *Galaxea* 12: 1-82.
- Grygier, M.J. 1997. *Monstrilla* Dana, 1849 and *Thaumaleus*Bull. zool. Nom. (Opinion 1869) 54 (2): 131-132.
- Ho, J.-S., Conradi, M. & López-González, P.J. 1998. A new family of cyclopoid copepods (Fratiidae) parasitic in the ascidian (*Clavelina dellavallei*) from Cádiz, Spain. *J. Zool. Lond.* 246: 39-48
- Hogans, W.E. 1987. Morphological variation in *Pennella balaeoptera* and *P. filosa* (Copepoda: Pennellidae) with a review of the genus *Pennella* Oken, 1816 parasitic on Cetacea. *Bull. mar. Sci.* 40 (3): 442-453.
- Holmes, J.M.C. 1998. Phenetic relationships among some free-living siphonostomatoid copepods from southwest Ireland, and the description of *Glannapontius maculatus* gen. et sp. nov. *J. Marine Systems* 15: 229-241.
- Humes, A.G. & Boxshall, G.A. 1996. A revision of the lichomolgoid complex (Copepoda: Poecilostomatoidea), with the recognition of six new families. *J. nat. Hist.* 30: 175-227.
- Huys, R., 1996. Superornatiremidae fam. nov. (Copepoda: Harpacticoida): An enigmatic family from North Atlantic anchialine caves. - *Sci. Mar.*, 60(4): 497-542.
- Huys, R. & Böttger-Schnack, R. 1997. On the diphyletic origin of the Oncaeidae Giesbrecht, 1892 (Copepoda: Poecilostomatoidea) with a phylogenetic analysis of the Lubbockiidae fam. nov. *Zool. Anz.* 235 (3-4): 243-261.
- Huys, R. & Boxshall, G.A. 1991. *Copepod Evolution*: 1-468. The Ray Society, London.
- Huys, R., J.M. Gee, C.G. Moore & R. Hamond, 1996. Marine and brackish water harpacticoid copepods. Part I.
- Huys, R. & T.M. Iliffe, 1998. Novocriniidae, a new family of harpacticoid copepods from anchialine caves in Belize. *Zool. Scr.*, 27: 1-15.
- Illg, P.L. & Dudley, P.L. 1961. Notodelphyid copepods from Banyuls-sur-Mer. *Vie Milieu* (Supplement 12): 1-126.
- Illg, P.L. & Dudley, P.L. 1965. Notodelphyid copepods from the vicinity of Naples. *Pubbl. Staz. zool. Napoli* 34: 373-451.
- Illg, P.L. & Dudley, P.L. 1980. The family Ascidiicolidae and its subfamilies (Copepoda, Cyclopoida), with descriptions of

- new species. *Mém. Mus. nat. Hist. nat. Paris* (N.S.) (A) (Zoologie) 117: 1-192.
- Jaume, D. & Boxshall, G.A. 1995 *Stygocyclopia balearica*, a new genus and species of calanoid copepod (Pseudocyclopidae) from anchihaline caves in the Balearic Islands (Mediterranean). *Sarsia* 80: 213-222.
- Jaume, D. & Boxshall, G.A. 1996a A new genus and two new species of cave-dwelling copepods from the Balearic Islands (Mediterranean). *J. nat. Hist.* 30: 989-1006.
- Jaume, D. & Boxshall, G.A. 1996b Rare cyclopoid copepods (Crustacea) from Mediterranean littoral caves. *Bull. nat. Hist. Mus. Lond. (Zool.)* 62: 83-99.
- Jaume, D. & Boxshall, G.A. 1996c Two new genera of cyclopiniid copepods (Crustacea) from anchihaline caves on western Mediterranean Islands. *Zool. J. Linn. Soc.* 117: 283-304.
- Jaume, D. & Boxshall, G.A. 1996d The persistence of an ancient marine fauna in Mediterranean waters: new evidence from misophrioid copepods living in anchihaline caves. *J. nat. Hist.* 30: 1583-1595.
- Jaume, D. & Boxshall, G.A. 1997a A new genus and two new species of misophrioid copepods (Crustacea) from the North Atlantic hyperbenthos. *Sarsia* 82: 39-54.
- Jaume, D. & Boxshall, G.A. 1997b Two new genera of cyclopiniid copepods (Cyclopoida: Cyclopiniidae) from anchihaline caves of the Canary and Balearic islands, with a key to genera of the family. *Zool. J. Linn. Soc.* 120: 79-101.
- Jaume, D., Cartes, J.E. & Boxshall, G.A. (in press) Shallow-water and not deep-sea as most plausible origin for cave-dwelling *Paramisophria* species (Copepoda: Calanoida: Arietellidae), with description of three new species from Mediterranean bathyal hyperbenthos and littoral caves. *Contrib. Zool.*
- Jaume, D., Fosshagen, A. & Iliffe, T.M. 1999. New cave-dwelling pseudocyclopiids (Copepoda, Calanoida, Pseudocyclopidae) from the Balearics, Canary, and Philippine archipelagos. *Sarsia* 84: 391-417.
- Kabata, Z. 1979. *Parasitic Copepoda of British fishes*. No. 152: i-xii, 1-468, figs. 1-2031. The Ray Society, London.
- Lamb, E.J., Boxshall, G.A., Mill, P.J. & Grahame, J. 1996 A new family of endoparasitic copepods (Poecilostomatoidea) from the Dog Whelk *Nucella lapillus*. *J. Crustacean Biol.* 16: 142-148.
- Lang, K., 1948. *Monographie der Harpacticiden*, I: 1-896, figs. 1-361; II: 897-1682, figs. 362-607, maps 1-378.
- López-González, P.J., Bresciani, J. & Conradi, M. 1998. *Tarifiscola bulbosus*, new genus, new species, a highly transformed parasitic copepod, with information on its parasitism and larval development. *J. crust. Biol.* 18 (3): 581-589.
- López-González, P.J. & Conradi, M. 1996. *Mychophilus palmatus*, a new species (Copepoda: Cyclopoida: Ascidiicolidae) associated with the compound ascidian *Botryllus leachi* Savigny, with remarks on congeneric species and related genera. *Hydrobiologia* 330: 67-72.
- Marchenkov, A.V. & Boxshall, G.A. 1995 A new family of copepods associated with ascidiaceans in the White Sea, and an analysis of antennular segmentation and setation patterns in the order Poecilostomatoidea. *Zool. Anz.* 234: 133-144.
- Markhaseva, E.L. 1996. *Calanoid copepods of the family Aetideidae of the World Ocean*: 1-331. St Petersburg. (Trudy Zoologicheskogo Instituta RAN 268).
- Noodt, W., 1955. Marmara denizi Harpacticoid'leri (Crust. Cop.). Marine Harpacticoiden (Crust. Cop.) aus dem Marmara Meer. - *Istanb. Univ. Fen Fak. Mecm.*, (B)20(1-2): 49-94.
- O'Reilly, M.G. 1995. A new genus of copepod (Copepoda: Poecilostomatoidea) commensal with the maldanid polychaete *Rhodine gracilior*, with a review of the family Clausiidae. *J. nat. Hist.* 29: 47-64.
- Por, F.D., 1964. A study of Levantine and Pontic Harpacticoida (Crustacea, Copepoda). - *Zool. Verh.*, Leiden, 64: 1-128.
- Raiabut, A., Combes, C. & Benoit, F. 1998. Analysis of the parasitic copepod species richness among Mediterranean fish. *J. mar. Syst.* 15: 185-206.
- Razouls, C. 1995. Diversité et répartition géographique chez les copépodes pélagiques. 1. Calanoida. *Annal. Inst. Oceanogr.* 71: 81-404.
- Razouls, C. 1995. Diversité et répartition géographique chez les copépodes pélagiques. 2. Platycopioida, Misophrioida, Mormonilloida, Cyclopoida, Poecilostomatoidea, Siphonostomatoidea, Harpacticoida, Monstrilloida. *Annal. Inst. Oceanogr.* 72: 1-149.
- Soyer, J., 1971. Bionomie benthique du plateau continental de la côte catalane française. III. Les peuplements de Copépodes Harpacticoides (Crustacea). - *Vie Milieu*, (B)21(2): 337-511.
- Zagami, G., Campolmi, M. & Costanzo, G. 2000. A new species of *Stephos* T.Scott, 1892 (Copepoda: Calanoida) from coastal waters of Sicily, Italy. *J. plankton Res.* 22: 15-27.
- ### Subclass Copepoda
- #### Order Cyclopoida
- ##### Family Ascidiicolidae
- Ascidicola*
 - rosea* Thorell, 1859
 - Styelicola*
 - bahusia* Lützen, 1968
- ##### Family Botryllophilidae
- Botryllophilus*
 - aspinosus* Schellenberg, 1922
 - banyulensis* Brément, 1909
 - bergensis* Schellenberg, 1921
 - brevipes* Brément, 1909
 - macropus* Canu, 1891
 - norvegicus* Schellenberg, 1921
 - ruber* Hesse, 1864
 - Schizoprocus*
 - inflatus* Aurivillius, 1885
 - vestitus* (Sars, 1921)
- ##### Family Buproridae
- Buprorus*
 - loveni* Thorell, 1859
 - nordgaardi* Sars, 1921
- ##### Family Chordeumiidae
- Chordeumium*
 - obesum* (Jungersen, 1912)
 - Ophioika*
 - appendiculata* Stephensen, 1935
 - Parachordeumium*
 - amphiurae* (Hérouard, 1906)
- ##### Family Cyclopidae
- Euryte*
 - curticornis* Sars, 1913
 - longicauda* Philippi, 1843
 - robusta* Giesbrecht, 1900
 - Halicyclops*
 - brevispinosus* subsp. *brevispinosus* Herbst, 1952
 - brevispinosus* subsp. *meridionalis* Herbst, 1953
 - dalmatinus* Petkovski, 1955
 - magniceps* (Lilljeborg, 1853)
 - neglectus* Kiefer, 1935
 - rotundipes* subsp. *putealis* Kiefer, 1938
 - rotundipes* subsp. *rotundipes* Kiefer, 1935
 - troglodytes* Kiefer, 1954
 - Neocyclops*
 - mediterraneus* (Kiefer, 1960)
 - remanei* (Herbst, 1952)
 - vicus* (Herbst, 1955)
- ##### Family Cyclopiniidae
- Cyclopetta*
 - difficilis* Sars, 1913
 - Cyclopicina*
 - longifurcata* (T. Scott, 1901)

- Cyclopina**
adriatica Petkovski, 1955
brachystylis Sars, 1921
crassisetosa Herbst, 1953
dilatata Sars, 1921
elegans T. Scott, 1894
esilis Brian, 1938
euacantha Sars, 1918
gracilis Claus, 1863
hadzii Petkovski, 1955
kieferi subsp. *elongata* Petkovski, 1955
longicornis Boeck, 1872
norvegica Boeck, 1864
oblivia Monchenko, 1981
pontica Monchenko, 1977
psammophila Steuer, 1940
pygmaea Sars, 1918
rotundipes Herbst, 1952
schneideri T. Scott, 1903
steueri Fruchtl, 1923
- Cyclopinella**
tumidula Sars, 1913
- Cyclopinooides**
littoralis (Brady, 1872)
- Cycloporella**
eximia Monchenko, 1981
- Gnesia**
longicaudata Jaume & Boxshall, 1997
- Glareolina**
kliei (Herbst, 1953)
- Hemicyclopina**
setifera (Herbst, 1952)
- Herbstina**
exigua (Herbst, 1974)
- Metacyclopina**
brevisetosa Herbst, 1974
roscoffensis Bozic, 1953
- Muceddina**
trispinosa Jaume & Boxshall, 1996
- Neocyclopina**
reducta Herbst, 1952
rotundipes subsp. *putealis* Kiefer, 1938
- Oromiina**
fortunata Jaume & Boxshall, 1997
- Pseudocyclopinodes**
dacunhai Lindberg, 1961
neglecta (Herbst, 1952)
- Pterinopsyllus**
egregius Giesbrecht, 1900
illistris Giesbrecht, 1900
insignis (Brady, 1878)
- Troglocyclopina**
balaearica Jaume & Boxshall, 1996
- Family Enterocolidae**
- Enterocola**
beaumonti T. & A. Scott, 1895
bilamellata Sars, 1921
brementi Illg & Dudley, 1980
clavelinae Chatton & Harant, 1924
fertilis Illg & Dudley, 1980
fulgens van Beneden, 1860
gottoi López-González, Conradi & Garcia-Gomez, 1993
hessei Chatton & Harant, 1924
hibernica T. & A. Scott, 1895
mammifera Chatton & Harant, 1922
megalova Gotto, 1962
petiti Guille, 1964
precaria Illg & Dudley, 1980
pterophora Chatton & Brément, 1909
setifera Hansen, 1923
sydnii Chatton & Harant, 1924
- Enterocolides**
ecaudatus Chatton & Harant, 1922
- Lequerrea**
- canui* Illg & Dudley, 1980
perezi Chatton & Harant, 1924
- Family Enterognathidae**
- Enterognathus**
comatulae Giesbrecht, 1900
- Family Enteropsidae**
- Enteropsis**
chattoni Monniot, 1961
roscoffensis Chatton & Brément, 1909
- Mychophilus**
palmatus López-González & Conradi, 1996
roseus Hesse, 1865
- incertae sedis**
- Gomphopodarion**
byssicum Humes, 1974
- Family Fratiidae**
- Fratia**
gaditana Ho, Conradi & López-González, 1998
- Family Haplostomidae**
- Haplostoma**
banyulensis Brément, 1909
brevicauda (Canu, 1886)
canui Chatton & Harant, 1924
eruca (Norman, 1869)
mizoulei Monniot, 1962
- Haplostomides**
hibernicus (Scott, T. & A., 1895)
scotti Chatton & Harant, 1924
- Family Notodelphyidae**
- Bonnierilla**
altera Stock, 1967
arcuata Brément, 1909
armata Schellenberg, 1922
filipes Stock, 1967
longipes (Kerschner, 1879)
similis Illg & Dudley, 1961
- Botachus**
cylindratus Thorell, 1859
- Bremertia**
balneolensis Chatton & Brément, 1915
illgi Laubier & Lafargue, 1974
- Conchodelphys**
delamarei Lafargue & Laubier, 1968
- Demoixys**
chattoni Illg & Dudley, 1961
dialepta Illg & Dudley, 1961
- Doroixys**
uncinata Kerschner, 1879
- Doropygella**
normani (Brady, 1878)
porcicauda (Brady, 1878)
psyllus (Thorell, 1859)
thorelli (Aurivillius, C.W.S., 1882)
- Doropygopsis**
arctica Marchenkova, 1998
longicauda (Aurivillius, C.W.S., 1882)
- Doropygus**
pulex Thorell, 1859
- Goniodelphys**
trigona Buchholz, 1869
- Gunenotophorus**
globularis Buchholz, 1869
- Haplostatus**
incubatrix Illg & Dudley, 1961
- Lobodelphys**
elephas Illg & Dudley, 1961
- Mesoixys**
otaria Illg & Dudley, 1965
- Notodelphys**
acanthomela Illg & Dudley, 1961
agilis Thorell, 1859
allmani Thorell, 1859
caerulea Thorell, 1859

- canui* Roland, 1962
cryptopyge Bocquet & Stock, 1960
elegans Thorell, 1859
haranti Illg & Dudley, 1961
platymera Illg & Dudley, 1961
prasina Thorell, 1859
reducta Illg & Dudley, 1961
rufescens Thorell, 1859
tenera Thorell, 1859
translantica Bocquet & Stock, 1960
- Notopterophorus**
auritus (Thorell, 1859)
dimitus Illg & Dudley, 1961
elatus Giesbrecht, 1892
elongatus Buchholz, 1869
papilio Hesse, 1864
- Ooneides**
amela Chatton & Brément, 1915
- Ophiodelphys**
illgi Bocquet & Stock, 1961
- Ophioseide**
cardiacephalus Gerstaecker, 1871
- Pachypygus**
gibber (Thorell, 1859)
- Prophioseides**
abdominalis (Chatton & Brement, 1915)
delamarei Illg & Dudley, 1961
diplosomae Illg & Dudley, 1961
- Scolecimorpha**
joubini (Chatton, 1909)
- Sicyodelphys**
bocqueti Laubier & Lafargue, 1968
- Family Oithonidae**
- Oithona*
atlantica Farran, 1908
brevicornis Giesbrecht, 1891
decipiens Farran, 1913
fallax Farran, 1913
hamata Rosendorn, 1917
hebes Giesbrecht, 1891
linearis Giesbrecht, 1891
longispina Nishida, 1977
minuta T. Scott, 1894
nana Giesbrecht, 1892
plumifera Baird, 1843
pseudofrigida Rosendorn, 1917
robusta Giesbrecht, 1891
setigera (Dana, 1849)
similis Claus, 1866
simplex Farran, 1913
tenuis Rosendorn, 1917
vivida Farran, 1913
- Paroithona*
parvula Farran, 1908
pulla Farran, 1913
- Family Thaumatopsyllidae**
Thaumatopsyllus
paradoxus Sars, 1913
- Order Poecilostomatoida**
- Family Anomoclausiidae**
Anomoclausia
indrehusae Gotto, 1964
- Family Antheacheridae**
Antheacheres
duebeni M. Sars, 1857
Stauromoma
parasiticum Will, 1844
- Family Anthessiidae**
Anthessius
arenicolus (Brady, 1872)
leptostylis (Sars, 1916)
teissieri Bocquet & Stock, 1958
- Rhinomolgus*
- anomalus* Sars, 1918
- Family Bomolochidae**
Bomolochus
bellones (Burmeister, 1835)
soleae Claus, 1864
unicirrus (Brian, 1902)
Holobomolochus
confusus (Stock, 1959)
Nothobomolochus
cornutus (Claus, 1864)
scomberesocis (Krøyer, 1864)
Pseudoeucanthus
alosae Brian, 1906
- Family Catiniidae**
Catinia
plana Bocquet & Stock, 1957
- Myzomolgus*
stupendus Bocquet & Stock, 1957
- Family Chondracanthidae**
Acanthochondria
clavata (Bassett-Smith, 1896)
cornuta (Müller, 1776)
limandae (Krøyer, 1863)
soleae (Krøyer, 1838)
triglae Herrera-Cubilla & Raibaut, 1990
- Acanthochondrites*
annulatus (Olsson, 1869)
- Chondracanthodes*
deflexus Wilson, 1932
radiatus (Müller, 1776)
- Chondracanthus*
angustatus Heller, 1865
lophii Johnson, 1836
merluccii (Holten, 1802)
neali Leigh-Sharpe, 1930
ninnii Richiardi, 1882
nodosus (Müller, 1776)
ornatus T. Scott, 1900
zei Delaroche, 1811
- Lernentoma*
asellina (Linnaeus, 1758)
- Strabax*
monstruosus (von Nordmann, 1832)
- Family Clausidiidae**
Conchyliurus
cardii subsp. *cardii* Gooding, 1957
cardii subsp. *tapetis* Bocquet & Stock, 1957
solenis Bocquet & Stock, 1957
- Giardella*
callianassae Canu, 1888
thompsoni A. Scott, 1906
- Hemicyclops*
aberdonensis (T. & A. Scott, 1892)
cylindracea Pelseneer, 1929
pureus Boeck, 1872
- Hersiliodes*
latericia (Grube, 1869)
- Leptinogaster*
histrio (Pelseneer, 1929)
pholadis Pelseneer, 1929
- Pseudopsyllus*
elongatus Scott, 1902
- Family Clausiidae**
Clausia
lubbocki Claparède, 1863
uniseta Bocquet & Stock, 1960
- Megaclausia*
mirabilis O'Reilly, 1995
- Mesnilia*
cluthae (T. & A. Scott, 1896)
martinensis Canu, 1898
- Pontoclausia*
tomis Bacescu & Por, 1959
- Pseudoclausia*

- giesbrechti* Bocquet & Stock, 1960
longiseta Bocquet & Stock, 1963
- Rhodinicola*
elongata Levinsen, 1878
gibbosa Bresciani, 1964
rugosum (Giesbrecht, 1895)
- Family Corycaeidae**
- Agetus*
flaccus (Giesbrecht, 1891)
limbatus (Brady, 1883)
typicus (Krøyer, 1849)
- Corycaeus*
clausi F. Dahl, 1894
crassiusculus Dana, 1849
speciosus Dana, 1849
venustus Dana, 1849
- Ditrichocorycaeus*
africanus (F. Dahl, 1894)
agilis (Dana, 1849)
andrewsi (Farran, 1911)
anglicus (Lubbock, 1857)
brehmi (Steuer, 1910)
lubbocki (Giesbrecht, 1891)
minimus (F. Dahl, 1894)
subtilis (M. Dahl, 1912)
tenuis (Giesbrecht, 1891)
- Farranula*
carinata (Giesbrecht, 1891)
curta (Farran, 1911)
gracilis (Dana, 1849)
rostrata (Claus, 1863)
- Monocorycaeus*
robustus (Giesbrecht, 1891)
- Onychocorycaeus*
catus (F. Dahl, 1894)
giesbrechti (F. Dahl, 1894)
latus (Dana, 1849)
ovalis (Claus, 1863)
pumilis (M. Dahl, 1912)
- Urocorrycaeus*
furcifer (Claus, 1863)
lautus (Dana, 1849)
- Family Entobiidae**
- Entobius*
hamondi Gotto, 1966
- Family Ergasilidae**
- Ergasilus*
briani Markevich, 1933
gibbus von Nordmann, 1832
lizae (Krøyer, 1863)
ponticus Markewisch, 1934
sieboldi von Nordmann, 1832
- Neoergasilus*
japonicus (Harada, 1930)
- Nipergasilus*
bora (Yamaguti, 1939)
- Paraergasilus*
longidigitus Yin, 1954
- Thersitina*
gasterostei (Pagenstecher, 1861)
- Family Eunicicolidae**
- Eunicicola*
clausi Kurz, 1877
insolens (T. & A. Scott, 1898)
- Family Gastrodelphyidae**
- Gastrodelphys*
clausii Graeffe, 1883
myxicolae List, 1890
- Sabellacheres*
drachi Laubier
gracilis M. Sars, 1862
- Family Herpilloidae**
- Eurysilenium*
oblongum Hansen, 1887
truncatum Sars, M., 1870
- Herpillobius*
arcticus Steenstrup & Lützen, 1861
polynoes (Krøyer, 1863)
- Phallusiella*
psalliota Leigh-Sharpe, 1926
vera Leigh-Sharpe, 1926
- Family Intramolgidae**
- Intramolgus*
arcticus Marchenkov & Boxshall, 1995
- Family Lamippidae**
- Enalcyonium*
confusum Stock, 1988
forbesi (T. Scott, 1901)
olssoni (De Zulueta, 1908)
rubicundum Olsson, 1869
- Lamippe*
parva De Zulueta, 1908
rubra Bruzelius, 1858
seigera De Zulueta, 1908
- Lamippella*
faurei Bouligand & Delamare Debouteville, 1959
- Lamippina*
proteus (Claparède, 1867)
- Lamippulina*
aciculifera (De Zulueta, 1908)
- Linarea*
mamillifera De Zulueta, 1908
- Family Lichomolgidae**
- Astericola*
clausi Rosoll, 1888
- Epimolgus*
trochi (Canu, 1899)
- Herrmannella*
barneae (Pelseneer, 1929)
duggani Holmes & Minchin, 1991
haploceras (Bocquet & Stock, 1959)
parva Norman & T. Scott, 1905
pecteni (Sowinski, 1884)
rostrata Canu, 1891
valida Sars, 1918
- Heteranthessius*
furcatus Stock, 1971
scotti Bocquet, Stock & Benard, 1959
- Lichomolgella*
isseli Gallienghani, 1952
pusilla Sars, 1918
- Lichomolgides*
cuanensis Gotto, 1954
- Lichomolgidium*
cynthiae (Brian, 1924)
sardum Koßmann, 1877
- Lichomolgus*
albens Thorell, 1859
canui Sars, 1917
diazonae Gotto, 1961
elegantulus Stock, 1960
forficula Thorell, 1859
furcillatus Thorell, 1859
leptodermatus Gooding, 1957
longicauda (Claus, 1860)
marginatus Thorell, 1859
- Modiolicola*
inermis Canu, 1892
insignis Aurivillius, 1882
maximus (Thompson, 1893)
- Stellicola*
hochbergi López-González & Pascual, 1996
- Zygomolgus*
curtiramus (Bocquet & Stock, 1962)
didemni (Gotto, 1956)
poucheti (Canu, 1891)
tenuifurcatus (Sars, 1917)

Family Lubbockiidae	
<i>Atrophia</i>	
<i>minuta</i> (Wolfenden, 1905)	
<i>Homoeognathia</i>	
<i>brevis</i> (Farran, 1908)	
<i>Lubbockia</i>	
<i>aculeatus</i> Giesbrecht, 1891	
<i>squillimana</i> Claus, 1863	
<i>Pseudolubbockia</i>	
<i>dilatata</i> Sars, 1909	
Family Macrochironidae	
<i>Macrochiron</i>	
<i>fucicolum</i> Brady, 1872	
<i>mutatum</i> Stock, 1957	
Family Mesoglicolidae	
<i>Mesoglicola</i>	
<i>delagei</i> Quidor, 1906	
Family Myicolidae	
<i>Pseudomyicola</i>	
<i>spinosus</i> (Raffaele & Monticelli, 1885)	
Family Mytilicolidae	
<i>Mytilicola</i>	
<i>intestinalis</i> Steuer, 1902	
<i>orientalis</i> Mori, 1935	
<i>ostrea</i> Hoshina & Sugiura, 1953	
<i>Trochicola</i>	
<i>entericus</i> Dollfus, 1914	
Family Nereicolidae	
<i>Anomopsyllus</i>	
<i>abyssorum</i> Laubier, 1988	
<i>pranizoides</i> Sars, 1921	
<i>Flabelliphilus</i>	
<i>inersus</i> Bresciani & Lützen, 1962	
<i>Nereicola</i>	
<i>ovatus</i> Keferstein, 1863	
<i>Selioides</i>	
<i>bocqueti</i> Carton, 1963	
<i>bolbroei</i> Levinse, 1878	
<i>Selius</i>	
<i>bilobus</i> Krøyer, 1837	
<i>Sigecheres</i>	
<i>brittae</i> Bresciani, 1964	
<i>Vectoriella</i>	
<i>marinovi</i> Stock, 1968	
<i>ramosae</i> Laubier & Carton, 1973	
Family Nucellicolidae	
<i>Nucellcola</i>	
<i>holmanae</i> Lamb, Boxshall, Mill & Grahame, 1996	
Family Octopicolidae	
<i>Octopicola</i>	
<i>superbus</i> Humes, 1957	
Family Oncaeidae	
<i>Conaea</i>	
<i>gracilis</i> (Dana, 1849)	
<i>rapax</i> Giesbrecht, 1891	
<i>Epicalyptma</i>	
<i>ancora</i> (Gordeeva, 1973)	
<i>brittoni</i> Heron, English & Damkaer, 1984	
<i>exigua</i> (Farran, 1908)	
<i>schmitti</i> Heron, 1977	
<i>umbonata</i> Heron, 1977	
<i>Oncaea</i>	
<i>africana</i> Shmeleva, 1979	
<i>alboranica</i> Shmeleva, 1979	
<i>anglica</i> Brady, 1905	
<i>atlantica</i> Shmeleva, 1967	
<i>bathyialis</i> Shmeleva, 1968	
<i>brodskii</i> Shmeleva, 1968	
<i>curta</i> Sars, 1916	
<i>delicata</i> Heron, English & Damkaer, 1984	
<i>englishi</i> Heron, 1977	
<i>furnestini</i> Shmeleva, 1979	
<i>infantula</i> Gordeeva, 1972	
	<i>ivlevi</i> Shmeleva, 1966
	<i>lacinia</i> Heron, English & Damkaer, 1984
	<i>longipes</i> Shmeleva, 1968
	<i>longiseta</i> Shmeleva, 1968
	<i>media</i> Giesbrecht, 1891
	<i>mediterranea</i> (Claus, 1863)
	<i>minima</i> Shmeleva, 1968
	<i>minor</i> Shmeleva, 1979
	<i>mollicula</i> Gordeeva, 1975
	<i>notopus</i> Giesbrecht, 1891
	<i>ornata</i> Giesbrecht, 1891
	<i>ovalis</i> Shmeleva, 1966
	<i>parila</i> Heron, 1977
	<i>prendeli</i> Shmeleva, 1966
	<i>pumilis</i> Heron, 1977
	<i>rotundata</i> Boxshall, 1977
	<i>scottodiculari</i> Heron & Bradford-Grieve, 1995
	<i>shmelevi</i> Gordeeva, 1972
	<i>subtilis</i> Giesbrecht, 1892
	<i>tenella</i> Sars, 1916
	<i>venusta</i> Philippi, 1843
	<i>zernovi</i> Shmeleva, 1966
	Family Pharodidae
	<i>Pharodes</i>
	<i>banyulensis</i> Delamare Deboutteville, 1951
	<i>clini</i> Brian, 1940
	Family Philichthyidae
	<i>Colobomatooides</i>
	<i>splendidus</i> Essafi & Raibaut, 1980
	<i>Colobomatus</i>
	<i>agassizi</i> (Richiardi, 1877)
	<i>baraldii</i> (Richiardi, 1877)
	<i>benazzii</i> Delamare Deboutteville & Nunes Ruivo, 1958
	<i>canthari</i> Delamare Deboutteville & Nunes Ruivo, 1952
	<i>denticis</i> (Richiardi, 1877)
	<i>doderleini</i> (Richiardi, 1883)
	<i>edwardsi</i> (Richiardi, 1876)
	<i>fiatolae</i> (Richiardi, 1880)
	<i>grubei</i> (Richiardi, 1877)
	<i>haeckeli</i> (Richiardi, 1877)
	<i>labracis</i> Delamare Deboutteville & Nunes Ruivo, 1952
	<i>lichiae</i> (Richiardi, 1877)
	<i>mugilis</i> Raibaut <i>et al.</i> , 1978
	<i>mulli</i> Essafi, Raibaut & Boudadoud-Krissat, 1993
	<i>muraenae</i> (Richiardi, 1877)
	<i>oblatae</i> (Richiardi, 1900)
	<i>pagelli</i> (Richiardi, 1877)
	<i>pagri</i> (Richiardi, 1877)
	<i>sciaenae</i> (Richiardi, 1876)
	<i>sieboldi</i> (Richiardi, 1877)
	<i>sparsi</i> Essafi, 1982
	<i>steenstrupi</i> (Richiardi, 1876)
	<i>vallei</i> Essafi, Cabral & Raibaut, 1984
	<i>Lepophilus</i>
	<i>labrei</i> Hesse, 1866
	<i>Lernaeascus</i>
	<i>nematoxys</i> Claus, 1887
	<i>Philichthys</i>
	<i>xiphiae</i> Steenstrup, 1862
	<i>Sphaerifer</i>
	<i>corvinae</i> Lyedig, 1851
	<i>leydigii</i> Richiardi, 1877
	Family Phyllodocidae
	<i>Cyclorrhiza</i>

- eteonicola* Heegaard, 1942
megalova Gotto & Leahy, 1988
- Phyllodicola*
petiti (Delamare Deboutteville & Laubier, 1960)
- Family Pseudanthessiidae**
- Heteranthessius*
dubius (T. Scott, 1903)
furcatus Stock, 1971
scotti Bocquet, Stock & Benard, 1959
- Pseudanthessius*
assimilis Sars, 1917
dubius Sars, 1918
gracilis Claus, 1889
liber (Brady, 1880)
nemertophilus Gallien, 1936
sauvagei Canu, 1891
thorelli (Brady, 1880)
- Family Rhynchomolgidae**
- Critomolgas*
actiniae (Della Valle, 1880)
bulbipes (Stock & Kleeton, 1963)
- Doridicola*
actiniae (Della Valle, 1880)
agilis Leydig, 1853
botulosus (Stock & Kleeton, 1963)
chlamydias (Humes & Stock, 1973)
hirsutipes (T. Scott, 1893)
longicauda (Claus, 1860)
trispinosus (Stock, 1959)
- Paranthessius*
anemoniae Claus, 1889
- Pennatulicola*
pteroidis (Della Valle, 1880)
- Family Sabelliphilidae**
- Acaenomolgas*
protulae (Stock, 1959)
serpulae (Stock, 1960)
- Eupolymniphilus*
finmarchicus (T. Scott, 1903)
- Myxomolgas*
myxicolae (Bocquet & Stock, 1958)
proximus Humes & Stock, 1973
- Sabelliphilus*
elongatus M. Sars, 1862
sarsi Claparède, 1870
- Family Saccopsidae**
- Melinnacheres*
ergasiloides M. Sars, 1970
steenstrupi Bresciani & Lützen, 1961
terebellidis (Levinsen, 1878)
- Family Sapphirinidae**
- Copilia*
lata Giesbrecht, 1891
mediterranea (Claus, 1863)
mirabilis Dana, 1849
quadrata Dana, 1849
vitrea (Haeckel, 1864)
- Sapphirina*
angusta Dana, 1849
auronitens Claus, 1863
bicuspidata Giesbrecht, 1891
darwini Haeckel, 1864
gastrica Giesbrecht, 1891
gemma Dana, 1849
granulosa Giesbrecht, 1891
intestinata Giesbrecht, 1891
iris Dana, 1849
lactens Giesbrecht, 1892
maculosa Giesbrecht, 1892
metallina Dana, 1849
nigromaculata Claus, 1863
opalina Dana, 1849
ovatolanceolata Dana, 1849
pyrosomatis Giesbrecht, 1892
- sali* Farran, 1929
scarlata Giesbrecht, 1891
sinuicauda Brady, 1883
stellata Giesbrecht, 1891
vorax Giesbrecht, 1891
- Vettoria*
granulosa (Giesbrecht, 1891)
longifurca (Rose & Vaissière, 1952)
parva (Farran, 1936)
- Family Serpulidicolidae**
- Rhabdopus*
salmacinae Southward, 1964
- Serpulidicola*
josephella Humes & Grassle, 1979
omphalopomae Southward, 1964
placostegi Southward, 1964
- Family Spiophanicolidae**
- Spiophanicola*
spinosis Ho, 1984
- Family Splanchnotropidae**
- Splanchnotropus*
brevipes Hancock & Norman, 1863
gracilis Hancock & Norman, 1863
- Family Synapticolidae**
- Calypsina*
changeuxi (Stock & Kleeton, 1963)
- Scambicornis*
petiti (Stock & Kleeton, 1963)
prehensilis (Sars, 1918)
tenuecaudus (Sars, 1918)
- Family Synaptiphilidae**
- Presynaptiphilus*
acrocinae Bocquet & Stock, 1960
- Synaptiphilus*
cantacuzenei Bocquet & Stock, 1957
luteus Canu & Cuénot, 1892
tridens (T. & A. Scott, 1893)
- Family Taeniacanthidae**
- Phagus*
muraenae (Brian, 1906)
- Taeniacanthus*
balistae (Claus, 1864)
gobii (Brian, 1906)
laqueus (Leigh-Sharpe, 1935)
lucipetus (Holmes, 1985)
onosi (T. Scott, 1902)
wilsoni (A. Scott, 1929)
zeugopteri (T. Scott, 1902)
- incertae sedis*
- Akessonnia*
occulta Bresciani & Lützen, 1962
- Axinophilus*
thyasirae Bresciani & Ockelmann, 1966
- Conchoheres*
malleolatus Sars, 1918
- Gonophysema*
gullmarense Bresciani & Lützen, 1960
- Jeannelia*
minor T. Scott, 1902
- Ophelicola*
drachi Laubier, 1978
- Pachos*
punctatum (Claus, 1863)
- Tarificola*
bulbosus López-González, Bresciani & Conradi, 1998
- Teredoika*
serpentina Stock, 1959
- Family Ventriculinidae**
- Endocheres*
obscurus Bocquet & Stock, 1956
- Myzotheridion*
sequenziae Laubier & Bouchet, 1976

- Family Xenocoelomidae**
- Aphanodomus*
terebellae (Levinsen, 1878)
 - Xenocoeloma*
alleni (Brumpt, 1897)
- Order Siphonostomatoida**
- Family Artotrogidae**
- Arctopontius*
expansus Sars, 1915
 - Artotrogus*
orbicularis Boeck, 1859
 - Bradyponтиus*
caudatus Sars, 1915
 - chelifer* Giesbrecht, 1895
 - dentatus* Hansen, 1923
 - magniceps* (Brady, 1880)
 - major* Sars, 1915
 - papillatus* (T. Scott, 1888)
 - siphonatus* Giesbrecht, 1895
 - unidens* Hansen, 1923
 - Cribropontius*
normani (Brady & Robertson, 1876)
 - Cryptopontius*
brevifurcatus (Giesbrecht, 1895)
 - capitalis* Giesbrecht, 1895
 - minor* Stock, 1965
 - tenuis* Giesbrecht, 1895
 - iherelli* Giesbrecht, 1895
 - DysPontius*
curticaudatus T. Scott, 1905
 - fringilla* (Giesbrecht, 1895)
 - passer* (Giesbrecht, 1897)
 - striatus* Thorell, 1859
 - Dystrogus*
gladiator Giesbrecht, 1899
 - Glannapontius*
maculatus Holmes, 1998
 - Metapontius*
latispinis Hansen, 1923
 - Myzopontius*
pungens Giesbrecht, 1895
 - Neopontius*
angularis T. Scott, 1898
 - Pteropontis*
cristatus Giesbrecht, 1895
 - Sestropontius*
bullifer Giesbrecht, 1899
- Family Asterocheridae**
- Acontiophorus*
antennatus Hansen, 1923
 - armatus* Brady, 1880
 - bracatus* Stock & Kleeton, 1963
 - ornatus* (Brady & Robertson, 1876)
 - scutatus* (Brady & Robertson, 1873)
 - Asterocheres*
bacescui (Marcus, 1965)
 - boecki* (Brady, 1880)
 - canui* Giesbrecht, 1897
 - complexus* Stock, 1960
 - corneliae* Schirl, 1973
 - dentatus* Giesbrecht, 1897
 - echinicola* (Norman, 1868)
 - ellisi* Hamond, 1968
 - genodon* Stock, 1966
 - kervillei* Canu, 1898
 - latus* (Brady, 1880)
 - lilljeborgi* Boeck, 1859
 - minutus* Giesbrecht, 1897
 - mucronipes* Stock, 1960
 - parvus* Giesbrecht, 1897
 - renaudi* Canu, 1891
 - simplex* Schirl, 1973
 - stimulans* (T. Scott, 1898)
 - siphonatus* (Thorell, 1859)
- stimulans* Giesbrecht, 1897
 - suberitis* Giesbrecht, 1899
 - uncinatus* (Kricagin, 1873)
 - ventricosum* (Brian, 1928)
 - violaceus* (Claus, 1889)
 - Collocheres*
breeei Stock, 1960
 - canui* Giesbrecht, 1897
 - elegans* Scott, A., 1896
 - gracilicauda* (Brady, 1880)
 - gracilipes* Stock, 1966
 - Dermatomyzon*
nigripes (Brady & Robertson, 1876)
 - Mesocheres*
anglicus Norman & Scott, 1905
 - Psilomyzon*
pauciseta Stock, 1965
 - Rhynchomyzon*
falco Giesbrecht, 1895
 - purpurocinctum* (T. Scott, 1893)
 - rubrovittatum* Sars, 1915
 - Scottocheres*
elongatus (T. & A. Scott, 1894)
 - laubieri* Stock, 1967
 - longifurca* Giesbrecht, 1897
 - Scottomyzon*
gibberum (T. & A. Scott, 1894)
 - Tuphacheres*
micropus Stock, 1965
- Family Brychiopontiidae**
- Brychiopontius*
falcatus Humes, 1974
- Family Caligidae**
- Caligodes*
laciniatus (Krøyer, 1863)
 - Caligopsis*
ponticus Markewisch, 1940
 - Caligus*
affinis Heller, 1866
 - alalongae* Krøyer, 1863
 - angustatus* Krøyer, 1863
 - belones* Krøyer, 1863
 - biaculeatus* Brian, 1914
 - bonito* Wilson, 1905
 - brevicaudatus* A. Scott, 1901
 - centrodonti* Baird, 1850
 - coryphaenae* Steenstrup & Lütken, 1861
 - curtus* Müller, 1785
 - deformis* Brian, 1924
 - diaphanus* von Nordmann, 1832
 - dicentrarchi* Cabral & Raibaut, 1985
 - dieuzeidei* Brian, 1932
 - elongatus* von Nordmann, 1832
 - guerini* Guiart, 1913
 - gurnadi* Krøyer, 1863
 - hyalinus* Chernyavsky, 1868
 - labracis* T. Scott, 1902
 - lacustris* Steenstrup & Lütken, 1861
 - leptochilus* Leuckart in Frey & Leuckart, 1847
 - lichiae* Brian, 1906
 - ligisticus* Brian, 1906
 - mauritanicus* Brian, 1924
 - minimus* Otto, 1821
 - pagelli* Delamare Deboutteville & Nunes-Ruivo, 1958
 - pageti* Russell, 1925
 - pelamydis* Krøyer, 1863
 - productus* Dana, 1852
 - robustus* Bassett-Smith, 1898
 - vexator* Heller, 1865
 - zei* Norman & Scott, 1906
 - Diphyllogaster*
thompsoni Brian, 1899
 - Lepeophtheirus*
crassus (Wilson & Bere, in Bere, 1936)
 - dissimilatus* Wilson, 1905

- europensis* Zeddam, Berrebi, Renaud, Raibaut & Gabrion, 1988
hippoglossi (Krøyer, 1837)
longipes Wilson, 1905
nordmanni (Milne Edwards, 1840)
pectoralis (Müller, 1776)
pollachius Bassett-Smith, 1896
rotundiventralis Bassett-Smith, 1898
salmonis (Krøyer, 1837)
sturionis (Krøyer, 1838)
thompsoni Baird, 1850
Pseudocaligus
apodus (Brian, 1924)
brevipedis (Bassett-Smith, 1896)
Scienophilus
tenuis van Beneden, 1852
- Family Calverocheridae**
Calveroheres
oblongus Stephensen, 1935
- Family Cancerillidae**
Cancerilla
tubulata Dalyell, 1851
Parartotrogus
arcticus (T. Scott, 1901)
richardi T. & A. Scott, 1893
- Family Cecropidae**
Cecrops
latreilli Leach, 1816
Luetkenia
asterodermi Claus, 1864
integra Richiardi, 1880
Orthagoriscola
muricatus (Krøyer, 1837)
Philorthagoriscus
serratus (Krøyer, 1863)
- Family Dicellesthiidae**
Anthosoma
crassum (Abildgaard, 1794)
Dicellestium
oblongum (Abildgaard, 1794)
- Family Dinopontiidae**
Dinopontius
acuticauda Stock, 1960
- Family Dirivultidae**
Aphotopontius
atlanteus Humes, 1996
temperatus Humes, 1997
Stygiopontius
pectinatus Humes, 1987
rimivagus Humes, 1997
- Family Entomolepidae**
Entomopsyllus
adriae (Eiselt, 1959)
- Family Eudactylinidae**
Eudactylinia
acanthii A. Scott, 1901
acuta van Beneden, 1853
aspera Heller, 1865
insolens T. & A. Scott, 1913
minuta T. Scott, 1904
rhinobati Raibaut & Essafi, 1979
similis T. Scott, 1902
turgipes Bere, 1936
vilelai Nunes-Ruivo, 1956
Eudactylinella
alba Wilson, 1932
Nemesis
lamna (A. Scott, 1929)
robusta (van Beneden, 1851)
Protodactylinia
pamelae Laubier, 1966
- Family Euryphoridae**
- Alebion*
carchariae Krøyer, 1863
Euryphorus
brachypterus (Giesbrecht, 1853)
- Family Hatschekiidae**
Congericola
pallidus van Beneden, 1854
Hatschekia
cadenati Nunes-Ruivo, 1954
cernae Goggio, 1905
cluthae (T. Scott, 1902)
gerro Leigh-Sharpe, 1936
hippoglossi (Cuvier, 1830)
ischnon Leigh-Sharpe, 1936
labracis (van Beneden, 1871)
mulli (van Beneden, 1851)
obesa Goggio, 1905
pagellibognaravei (Hesse, 1879)
pygmaea T. & A. Scott, 1913
sargi (Brian, 1902)
- Family Kroyeriidae**
Kroyeria
carchariaeglauci Hesse, 1878
lineata van Beneden, 1853
- Family Lernaeopodidae**
Advena
paradoxa (van Beneden, 1851)
Albionella
globosa (Leigh-Sharpe, 1918)
Atella
macrotrachelus (Brian, 1906)
pagelli (Krøyer, 1863)
Brachiella
thynni Cuvier, 1830
Charopinus
dalmanni (Retzius, 1829)
dubius Scott, T., 1900
Clavella
adunca Strøm, 1762
alata Brian, 1906
denticis Krøyer, 1863
stellata (Krøyer, 1838)
Clavellosa
emarginata (Krøyer, 1837)
sombri (Kurz, 1877)
Clavelistes
lampri (T. & A. Scott, 1913)
Clavelloides
rugosus (Krøyer, 1837)
Clavellosis
characis (Richiardi, 1880)
fallax (Heller, 1865)
pagri (Krøyer, 1863)
sargi (Kurz, 1877)
strumosa (Brian, 1906)
Eubachiella
mugilis Kabata *et al.*, 1971
Lernaeopoda
bidiscalis Kane, 1892
galei Krøyer, 1837
longibrachia Brian, 1912
Lernaeopodina
longimana (Olsson, 1869)
spinacis (Brian, 1908)
Naobranchia
cygniformis Hesse, 1863
Neobachiella
bispinosa (von Nordmann, 1832)
chevreuxii (van Beneden, 1891)
elegans (Richiardi, 1880)
exigua (Brian, 1906)
hostilis (Heller, 1865)
impudica (von Nordmann, 1832)
insidiosa (Heller, 1865)

- merluccii* (Bassett-Smith, 1896)
ramosa (Richiardi, 1880)
richiardii Ben Hassine & Raibaut, 1978
rostrata (Krøyer, 1837)
triglae (Claus, 1860)
Ommatokoita
elongata (Grant, 1827)
Pseudocharopinus
bicaudatus (Krøyer, 1837)
malleus (von Nordmann, 1832)
pteromylaei Raibaut & Essafi, 1979
Salmicola
edwardsi (Olsson, 1869)
gordoni Gurney, 1933
salmoneus (Linnaeus, 1758)
thymalli (Kessler, 1868)
Schistobrachia
ramosa (Krøyer, 1863)
Sparidicola
lithognathi (Kensley & Grindley, 1973)
Thysanote
ramosa (Richiardi, 1880)
Vanbenedenia
kroeyeri Malm, 1860
Family Lernanthropidae
Lernanthropus
gisleri van Beneden, 1832
kroyeri van Beneden, 1851
micropterygis Richiardi, 1885
mugilis Brian, 1898
scribae Krøyer, 1863
trachuri Brian, 1903
tylosuri Richiardi, 1880
vorax Richiardi, 1879
Family Megapontiidae
Hyalopontius
alatus Boxshall, 1979
cinctus Boxshall, 1979
enormis Boxshall, 1979
hulsemannae Boxshall, 1979
roei Boxshall, 1979
spinatus Boxshall, 1979
typicus Sars, 1909
Family Micropontiidae
Micropontius
ovoides Gooding, 1957
Family Nanaspidae
Allantogynus
delamarei Changeux, 1958
Nanaspis
ninae Bresciani & Lützen, 1962
Family Nicothoidae
Aspidoezia
normani Giard & Bonnier, 1889
Choniorhiza
enaulis Boxshall & Lincoln, 1983
Choniosphaera
maenadis (Bloch & Gallien, 1933)
Choniostoma
hanseni Giard & Bonnier, 1886
rotundatum Stock, 1958
Diexanthema
apoda Boxshall & Harrison, 1988
corrugatum Boxshall & Harrison, 1988
nudum Boxshall & Harrison, 1988
ritchiei Boxshall & Harrison, 1988
Hadrothoe
crenulata Boxshall & Defaye, 1995
Homoeoscelis
elongata Boxshall & Defaye, 1995
frigida Hansen, 1923
longipes Hansen, 1923
mediterranea Hansen, 1897
minuta Hansen, 1897
sedentaria (Bonnier, 1898)
Mysidion
abyssorum Hansen, 1897
commune Hansen, 1897
Nicorhiza
humesi Lincoln & Boxshall, 1983
rockallensis Lincoln & Boxshall, 1983
Nicothoe
astaci Audouin & Milne Edwards, 1826
Rhizorhina
aesthetes Boxshall & Harrison, 1988
ampeliscae Hansen, 1892
hystrix Boxshall & Harrison, 1988
leptostracae Gotto, 1984
tanaidaceae Gotto, 1984
Sphaeronella
aeginae Hansen, 1923
affinis Hansen, 1897
amphilochi Hansen, 1897
atyli Hansen, 1897
callisoma T. Scott, 1904
cluthae T. Scott, 1904
danica Hansen, 1897
decorata Hansen, 1897
devosae Stock, 1960
diadema Giard & Bonnier, 1895
dispar Hansen, 1897
dulichiae Hansen, 1897
ecaudata Stock, 1960
frontalis Hansen, 1897
giardii Hansen, 1897
gitanopsidis Hansen, 1897
insignis Hansen, 1897
intermedia Hansen, 1897
irregularis Hansen, 1897
leptocheiri Hansen, 1897
leuckartii Salensky, 1868
longipes Hansen, 1897
marginata Hansen, 1897
melphidippae Hansen, 1923
messinensis Hansen, 1897
metopae Hansen, 1897
microcephala Giard & Bonnier, 1893
minuta T. Scott, 1904
modesta Hansen, 1897
monothrix (Bowman & Kornicker, 1967)
munnae Hansen, 1897
nanonisci Hansen, 1923
norvegica Hansen, 1897
paradoxa Hansen, 1897
pleurogonii Hansen, 1923
pygmaea T. Scott, 1904
rotundata Hansen, 1923
valida T. Scott, 1905
vararensis T. Scott, 1905
vestita Hansen, 1897
Stenothenocheres
egregius Hansen, 1897
sarsi Hansen, 1897
Family Pandaridae
Demoleus
ferox (Krøyer, 1838)
heptapus (Otto, 1821)
Dinemoura
latifolia (Steenstrup & Lütken, 1861)
producta (Müller, 1785)
Echthrogaleus
coleoptratus (Guerin-Meneville, 1837)
Gangliopus
pyriformis Gerstaecker, 1854
Nessipus
orientalis Heller, 1865
Pandarus
ambiguus T. Scott, 1907
bicolor Leach, 1816

- borealis* (Steenstrup & Lütken, 1861)
cranchii Leach, 1819
- Perissopus*
dentatus Steenstrup & Lütken, 1861
- Phyllohyreus*
cornutus Milne Edwards, 1840
- Family Pennellidae**
- Cardiodes*
bellottii (Richiardi, 1882)
- Exopenna*
crimmeni Boxshall, 1986
- Haemobaphes*
ambiguus T. Scott, 1902
cycloptera (Fabricius, 1780)
- Lernaenicus*
encrasicoli (Turton, 1807)
gracilis (Heller, 1865)
sprattae (Sowerby, 1806)
vorax Richiardi, 1877
- Lernaeocera*
branchialis (Linnaeus, 1767)
lisci (Bassett-Smith, 1896)
minuta T. Scott, 1900
- Lernaeolophus*
sultanus von Nordmann, 1839
- Peniculus*
fistula Rudolphi, 1880
- Pennella*
balaenoptera Koren & Danielssen, 1877
filosa (Linnaeus, 1758)
- Peroderma*
cylindricum (Heller, 1865)
- Sarcotretes*
eristaliformis (Brian, 1908)
scopeli Junghsen, 1911
- Family Pontocellidae**
- Pontocella*
abyssicola (T. Scott, 1894)
- Family Pseudocycnidae**
- Pseudocycnus*
appendiculatus Heller, 1865
- Family Rataniidae**
- Ratania*
atlantica Farran, 1926
flava Giesbrecht, 1892
- Family Sphyriidae**
- Lophoura*
edwardsi Kolliker, 1853
simplex Boxshall, 2000
- Paeon*
versicolor Wilson, 1919
- Sphyriion*
lumpi (Krøyer, 1845)
- Tripaphylus*
musteli (van Beneden, 1851)
- Family Sponginticolidae**
- Spongintcola*
uncifer Topsent, 1928
- Family Spongicizontidae**
- Apodomyzon*
brevicorne Stock, 1970
longicorne Stock, 1970
- Spongicnizon*
lobatum Stock, 1967
petiti Stock & Kleeton, 1964
vermiformis Stock, 1967
- Family Trebiidae**
- Treibius*
caudatus Krøyer, 1938
- Order Mormonilloidea**
- Family Mormonillidae**
- Mormonilla*
- minor* Giesbrecht, 1891
phasma Giesbrecht, 1891
- Order Monstrilloida**
- Family Monstrillidae**
- Cymbasoma*
claparedei (Giesbrecht, 1892)
frondipes (Isaac, 1975)
longispinosum (Bourne, 1890)
malaquini (Caulery & Mesnil, 1914)
pallidum Isaac, 1974
reticulatum (Giesbrecht, 1892)
rigidum Thompson, 1888
rostratum (T. Scott, 1904)
similirostratum Isaac, 1974
striatum Isaac, 1974
tenuis Isaac, 1974
thompsoni (Giesbrecht, 1892)
tumorigrons Isaac, 1974
zelandicum (T. Scott, 1904)
- Haemocera*
danae Malaquin, 1901
- Monstrilla*
anglica Lubbock, 1857
conjunctiva Giesbrecht, 1902
filogranarum (Malaquin, 1901)
gracilicauda Giesbrecht, 1892
grandis Giesbrecht, 1891
helgolandica Claus, 1863
intermedia Krichagin, 1877
longicornis Thompson, 1890
longiremis Giesbrecht, 1892
minuta Isaac, 1974
obesa Isaac, 1974
roscovita (Malaquin, 1901)
serricornis Sars, 1921
- Monstrillopsis*
angustipes Isaac, 1974
dubia (T. Scott, 1904)
gracilis (Gurney, 1927)
sarsi Isaac, 1974
zernovi Dolgopol'skaya, 1948
- Order Platycopioidea**
- Family Platycopiidae**
- Platycopia*
perplexa Sars, 1911
pygmaea Sars, 1919
robusta Andronov, 1985
- Order Misophrioida**
- Family Misophriidae**
- Misophria*
pallida Boeck, 1864
- Archimisophria*
discoveryi Boxshall, 1983
- Arcticomisophria*
hispida Jaume & Boxshall, 1997
- Benthomisophria*
cornuta Hulsemann & Grice, 1964
pallidata Sars, 1909
- Fosshageniella*
glabra Jaume & Boxshall, 1997
- Misophriella*
tetraspina Boxshall, 1983
- Misophriopsis*
dichotoma Boxshall, 1983
- Family Speleophriidae**
- Boxshallia*
bulbantennulata Huys, 1988
- Dimisophria*
cavernicola Boxshall & Iliffe, 1987
- Expansophria*
dimorpha Boxshall & Iliffe, 1987
sarda Jaume & Boxshall, 1996
- Speleophria*

- gymnesica* Jaume & Boxshall, 1996
Speleophriopsis
balearicus Jaume & Boxshall, 1996
canariensis Jaume & Boxshall, 1996
- Family Palpophriidae**
Palpophria
aesthetra Boxshall & Iliffe, 1987
- Order Calanoida**
- Family Acartiidae**
- Acartia*
adriatica Steuer, 1910
bifilosa (Giesbrecht, 1881)
centrura Giesbrecht, 1889
clausi Giesbrecht, 1889
danae Giesbrecht, 1889
discaudata (Giesbrecht, 1881)
enzoi Crisafi, 1974
fossae Gurney, 1927
italica Steuer, 1910
longiremis (Lilljeborg, 1853)
margalefi Alcaraz, 1976
negligens Dana, 1849
plumosa T. Scott, 1894
teclae Bradford, 1976
tonsa Dana, 1849
- Paracartia*
dubia T. Scott, 1894
grani Sars, 1904
latisetosa (Kriczagin, 1873)
- Pteriacartia*
josephinae (Crisafi, 1974)
- Family Aetideidae**
- Aetideopsis*
armata (Boeck, 1872)
carinata Bradford, 1969
modesta (With, 1915)
multiserrata (Wolfenden, 1904)
nasuta (With, 1915)
rostrata Sars, 1903
- Aetideus*
acutus Farran, 1929
arcuatus (Vervoort, 1949)
armatus (Boeck, 1872)
giesbrechti Cleve, 1904
- Batheuchaeta*
lamellata Brodsky, 1950
- Bradyetes*
brevis Farran, 1936
inermis Farran, 1905
matthei Johannessen, 1976
- Bradyidius*
bradyi (Sars, 1884)
luluæ Grice, 1972
similis (Sars, 1902)
- Chiridiella*
abyssalis Brodsky, 1950
gibba Deevey, 1974
kuniae Deevey, 1974
macrodactyla Sars, 1907
ovata Deevey, 1974
subaequalis Grice & Hulsemann, 1965
- Chiridius*
gracilis Farran, 1908
molestus Tanaka, 1957
obtusifrons Sars, 1902
poppei Giesbrecht, 1892
- Chirundina*
streetsi Giesbrecht, 1895
- Comantenna*
brevicornis (Sars, 1902)
- Euchirella*
amoena Giesbrecht, 1888
bella Giesbrecht, 1888
bitumida With, 1915
- curticauda* Giesbrecht, 1888
maxima Wolfenden, 1905
messinensis (Claus, 1863)
pulchra (Lubbock, 1856)
rostrata (Claus, 1866)
truncata Esterly, 1911
- Gaetanus*
antarcticus Wolfenden, 1905
armiger Giesbrecht, 1888
brachyurus Sars, 1907
curvicornis Sars, 1905
kruppi Giesbrecht, 1903
latifrons Sars, 1905
miles Giesbrecht, 1888
minor Farran, 1905
pileatus Farran, 1903
- Gaidius*
affinis Sars, 1905
brevicaudatus (Sars, 1907)
brevispinus (Sars, 1900)
inermis (Sars, 1905)
minutus Sars, 1907
pungens Giesbrecht, 1895
robustus Sars, 1905
tenuispinus (Sars, 1900)
- Jaschnovia*
brevis (Farran, 1936)
- Paracomantenna*
minor (Farran, 1905)
- Pseudeuchaeta*
brevicauda Sars, 1905
- Pseudochirella*
notacantha (Sars, 1905)
obesa Sars, 1920
obtusa (Sars, 1905)
palliata (Sars, 1907)
pustulifera (Sars, 1905)
scopularis (Sars, 1905)
spectabilis (Sars, 1900)
batillipa Park, 1978
calcarata Sars, 1920
cryptospina (Sars, 1905)
divaricata (Sars, 1905)
dubia (Sars, 1905)
fallax Sars, 1920
lobata (Sars, 1907)
major (Sars, 1907)
- Pseudotharybis*
zetlandicus T. Scott, 1909
- Undeuchaeta*
incisa Esterly, 1911
major Giesbrecht, 1888
plumosa (Lubbock, 1856)
- Valdiviella*
insignis Farran, 1908
minor Wolfenden, 1911
oligartha Steuer, 1904
- Family Ariettellidae**
- Ariettellus*
aculeatus T. Scott, 1894
armatus Wolfenden, 1911
giesbrechti Sars, 1905
pavoninus Sars, 1905
plumifer Sars, 1905
setosus Giesbrecht, 1892
simplex Sars, 1905
- Crassarietellus*
huyasi Ohtsuka, Boxshall & Roe, 1994
- Metacalanus*
acutiperulum Ohtsuka, 1984
inaequicornis (Sars, 1902)
- Paramisophria*
ammophila Fosshagen, 1968
bathyalis Jaume, Cartes & Boxshall, in press
cluthae T. Scott, 1897

- intermedia* Jaume, Cartes & Boxshall, in press
lilbetana Campolmi, Zagami & Costanzo, in press
mediterranea Jaume, Cartes & Boxshall, in press
reducta Ohtsuka, Fosshagen & Iliffe, 1993
spooneri Krishnaswamy, 1959
Paraugaptilus
archimedi Gaudy, 1972
buchani Wolfenden, 1904
meridionalis Wolfenden, 1911
Sarsariellus
abyssalis (Sars, 1905)
- Family Augaptilidae**
- Augaptilina**
scopifera Sars, 1920
- Augaptilus**
anceps Farran, 1908
cornutus Wolfenden, 1911
glacialis Sars, 1900
longicaudatus (Claus, 1863)
megalurus Giesbrecht, 1889
spinifrons Sars, 1907
- Centraugaptilus**
cucullatus (Sars, 1905)
horridus (Farran, 1908)
rattrayi (T. Scott, 1894)
- Euaugaptilus**
affinis Sars, 1920
angustus (Sars, 1905)
atlanticus Roe, 1975
brodskyi Hulsemann, 1967
bullifer Giesbrecht, 1889
clavatus (Sars, 1907)
digitatus Sars, 1920
elongatus (Sars, 1905)
facilis (Farran, 1908)
farrani Sars, 1920
secundus Tanaka & Omori, 1974
filigerus (Claus, 1863)
gibbus (Wolfenden, 1904)
gracilis (Sars, 1905)
grandicornis Sars, 1920
hecticus Giesbrecht, 1889
humilis Farran, 1926
hyperboreus Brodsky, 1950
laticeps (Sars, 1905)
latifrons (Sars, 1907)
longicirrus (Sars, 1905)
longimanu (Sars, 1905)
longiseta Grice & Hulsemann, 1965
magnus (Wolfenden, 1904)
maxillaris Sars, 1920
mixtus (Sars, 1907)
niveus Tanaka, 1953
nodifrons (Sars, 1905)
oblongus (Sars, 1905)
palumbii (Giesbrecht, 1889)
paroblongus Matthews, 1972
penicillatus Sars, 1920
placitus (A. Scott, 1909)
rectus Grice & Hulsemann, 1967
rigidus (Sars, 1907)
roei Matthews, 1972
sarsi Grice & Hulsemann, 1965
similis (Farran, 1908)
squamatus (Giesbrecht, 1889)
tenuicaudis (Sars, 1905)
tenuispinus Sars, 1920
truncatus (Sars, 1905)
vicus Sars, 1920
- Haloptilus**
acutifrons (Giesbrecht, 1892)
angusticeps Sars, 1907
bulliceps Farran, 1926
chierchiai (Giesbrecht, 1889)
fertilis (Giesbrecht, 1892)
- sons* Farran, 1908
furcatus Sars, 1920
longiceps Tanaka, 1964
longicirrus Brodsky, 1950
longicornis (Claus, 1863)
mucronatus (Claus, 1863)
ornatus (Giesbrecht, 1892)
oxycephalus (Giesbrecht, 1889)
paralongicirrus Park, 1970
plumosus (Claus, 1863)
spiniceps (Giesbrecht, 1892)
tenuis Farran, 1908
validus Sars, 1920
- Heteroptilus**
acutilobus (Sars, 1905)
attenuatus (Sars, 1905)
- Pachyptilus**
abbreviatus (Sars, 1905)
eurygnathus Sars, 1920
lobatus Sars, 1920
- Pontoptilius**
mucronatus Sars, 1905
muticus Sars, 1905
ovalis Sars, 1907
pertenuis Sars, 1907
robustus Sars, 1905
- Pseudaugaptilus**
longiremis Sars, 1907
- Pseudhaloptilus**
longimanus (Wolfenden, 1906)
- Family Bathypontiidae**
- Bathypontia**
elegans Sars, 1920
elongata Sars, 1905
minor (Wolfenden, 1906)
sarsi Grice & Hulsemann, 1965
spinifera A. Scott, 1909
- Foxtonia**
barbatula Hulsemann & Grice, 1963
- Sognocalanus**
confertus Fosshagen, 1967
- Temorites**
brevis Sars, 1900
discoveryae Grice & Hulsemann, 1965
- Zenkevitchiella**
atlantica Grice & Hulsemann, 1965
- Family Calanidae**
- Calanoides**
carinatus (Krøyer, 1848)
- Calanus**
euxinus Hulsemann, 1991
simarchicus (Gunnerus, 1770)
glacialis Jaschnov, 1955
helgolandicus (Claus, 1863)
hyperboreus (Krøyer, 1838)
- Mesocalanus**
tenuicornis (Dana, 1849)
- Nannocalanus**
minor (Claus, 1863)
- Neocalanus**
cristatus (Krøyer, 1845)
gracilis (Dana, 1849)
robustior (Giesbrecht, 1888)
- Undinula**
vulgaris (Dana, 1849)
- Family Candaciidae**
- Candacia**
armata (Boeck, 1872)
bipinnata (Giesbrecht, 1889)
curta (Dana, 1849)
elongata (Boeck, 1872)
ethiopica (Dana, 1849)
giesbrechti Grice & Lawson, 1977
ketchumi Grice, 1961

- longimana* (Claus, 1863)
norvegica (Boeck, 1864)
pachydactyla (Dana, 1849)
paenelongimana Fleminger & Bowman, 1956
tenuimana (Giesbrecht, 1889)
varicans (Giesbrecht, 1892)
Paracandacia
bispinosa (Claus, 1863)
simplex (Giesbrecht, 1889)
- Family Centropagidae**
Centropages
aucklandicus Kramer, 1895
brachiatus (Dana, 1849)
bradyi Wheeler, 1901
calaninus (Dana, 1849)
caribbeanensis Park, 1970
chierchiae Giesbrecht, 1889
furcatus (Dana, 1849)
hamatus (Lilljeborg, 1853)
kroyeri Giesbrecht, 1892
ponticus Karavaev, 1894
typicus Krøyer, 1849
violaceus (Claus, 1863)
Isius
clavipes Boeck, 1864
- Family Clausocalanidae**
Clausocalanus
arcuicornis (Dana, 1849)
furcatus (Brady, 1883)
jobei Frost & Fleminger, 1968
lividus Frost & Fleminger, 1968
mastigophorus (Claus, 1863)
parapergens Frost & Fleminger, 1968
paululus Farran, 1926
pergens Farran, 1926
Ctenocalanus
vanus Giesbrecht, 1888
- Farrania**
lyra (Rose, 1937)
oblonga Sars, 1920
- Microcalanus**
pygmaeus (Sars, 1900)
- Pseudocalanus**
acuspes (Giesbrecht, 1881)
elongatus (Boeck, 1865)
major Sars, 1900
minutus (Krøyer, 1845)
- Spicipes**
nansenii Grice & Hulsemann, 1965
- Family Diaixidae**
Diaixis
durani Corral Estrada, 1972
helenae Andronov, 1979
hibernica (A. Scott, 1896)
pygmaea (T. Scott, 1899)
trunovi Andronov, 1979
- Family Discoidae**
Disco
atlanticus Gordeeva, 1974
erythraeus Gordeeva, 1974
fjordicus Fosshagen, 1967
hartmanni Schulz, 1993
inflatus Grice & Hulsemann, 1965
longus Grice & Hulsemann, 1965
marinus Gordeeva, 1974
mediterraneus Gordeeva, 1974
minutus Grice & Hulsemann, 1965
oceanicus Gordeeva, 1974
populosus Gordeeva, 1976
robustipes Gordeeva, 1974
vulgaris Gordeeva, 1974
- Paradisco**
gracilis Gordeeva, 1975
mediterraneus (Gordeeva, 1974)
- nudus* Schulz, 1993
Prodisco
secundus Gordeeva, 1975
- Family Eucalanidae**
Eucalanus
subtenuis Giesbrecht, 1888
- Rhincalanus**
cornutus (Dana, 1849)
nasutus Giesbrecht, 1888
- Subeucalanus**
crassus (Giesbrecht, 1888)
monachus (Giesbrecht, 1888)
muconatus (Giesbrecht, 1888)
pileatus (Giesbrecht, 1888)
subcrassus (Giesbrecht, 1888)
subtenuis (Giesbrecht, 1888)
- Family Euchaetidae**
Euchaeta
acuta Giesbrecht, 1892
concinna Dana, 1849
marina (Prestandrea, 1833)
media Giesbrecht, 1888
pubera Sars, 1907
spinosa Giesbrecht, 1892
- Pareuchaeta**
albibulla Park, 1995
barbata (Brady, 1883)
bisinuata (Sars, 1907)
bradyi (With, 1915)
glacialis (Hansen, 1887)
gracilis (Sars, 1905)
hanseni (With, 1915)
hebes (Giesbrecht, 1888)
incisa (Sars, 1905)
kurilensis Heptner, 1971
norvegica (Boeck, 1972)
parabbreviata Park, 1995
polaris Brodsky, 1950
pseudotonsa (Fontaine, 1967)
rubicunda (Farran, 1908)
sarsi (Farran, 1908)
scotti (Farran, 1908)
tumidula (Sars, 1905)
- Family Heterorhabdidae**
Disseta
palumboi Giesbrecht, 1889
- Hemirhabus**
grimaldii (Richard, 1893)
- Heterorhabdus**
abyssalis (Giesbrecht, 1889)
clausi (Giesbrecht, 1889)
compactus (Sars, 1900)
norvegicus (Boeck, 1872)
papilliger (Claus, 1863)
robustus Farran, 1908
spinifer Park, 1970
spinifrons (Claus, 1863)
tenuis Tanaka, 1964
vipera (Giesbrecht, 1889)
- Heterostylites**
longicornis (Giesbrecht, 1889)
major (F. Dahl, 1894)
- Mesorhabdus**
angustus Sars, 1907
brevicaudatus (Wolfenden, 1905)
gracilis Sars, 1907
- Microdisseta**
minuta (Grice & Hulsemann, 1965)
- Neorhabdus**
latus (Sars, 1905)
- Family Lucicutiidae**
Lucicutia
anomala Brodsky, 1950
bicornuta Wolfenden, 1905

- clausi* (Giesbrecht, 1889)
curta Farran, 1905
flavicornis (Claus, 1863)
formosa Hulsemann, 1966
gaussae Grice, 1963
geminæ Farran, 1926
intermedia Sars, 1905
longicornis (Giesbrecht, 1889)
longiserrata (Giesbrecht, 1889)
longispina Tanaka, 1963
lucida Farran, 1908
macrocera Sars, 1920
magna Wolfenden, 1903
maxima Steuer, 1904
ovalis (Giesbrecht, 1889)
parva Grice & Hulsemann, 1965
pellucida Hulsemann, 1966
pera A. Scott, 1909
sarsi Hulsemann, 1966
tenuicauda Sars, 1907
wolfendeni Sewell, 1932
- Family Mecynoceridae**
Mecynocera
clausi I.C. Thompson, 1888
- Family Megacalanidae**
Bathycalanus
bradyi (Wolfenden, 1905)
eximus Markhaseva, 1983
princeps (Brady, 1883)
richardi Sars, 1905
Bradykalanus
sarsi (Farran, 1939)
Megacalanus
princeps Wolfenden, 1904
- Family Mesaiokeratidae**
Mesaiokeras
heptneri Andronov, 1973
nansenii Matthews, 1961
- Family Metridinidae**
Gaussia
princeps (T. Scott, 1894)
Metridia
alata Roe, 1975
brevicauda Giesbrecht, 1889
curticauda Giesbrecht, 1889
discreta Farran, 1946
effusa Grice & Hulsemann, 1967
gurjanovae Brodsky, 1950
longa (Lubbock, 1854)
lucens Boeck, 1864
macrura Sars, 1905
princeps Giesbrecht, 1889
venusta Giesbrecht, 1889
Pleuromamma
abdominalis (Lubbock, 1856)
borealis (F. Dahl, 1893)
gracilis (Claus, 1863)
piseki Farran, 1929
quadrungulata (F. Dahl, 1893)
robusta (F. Dahl, 1893)
xiphias (Giesbrecht, 1889)
- Family Paracalanidae**
Acrocalanus
andersoni Bowman, 1958
gibber Giesbrecht, 1888
longicornis Giesbrecht, 1888
Calocalanus
adriaticus Shmeleva, 1965
africanus Shmeleva, 1979
alboranus Shmeleva, 1979
atlanticus Shmeleva, 1975
contractus Farran, 1926
elegans Shmeleva, 1965
elongatus Shmeleva, 1968
- fusiformis* Shmeleva, 1978
gresei Shmeleva, 1974
indicus Shmeleva, 1973
kristalli Shmeleva, 1968
latus Shmeleva, 1968
lomonosovi Shmeleva, 1975
longisetosus Shmeleva, 1965
longispinus Shmeleva, 1978
minor Shmeleva, 1975
nambiensis Andronov, 1973
neptunus Shmeleva, 1965
omaniensis Shmeleva, 1975
ovalis Shmeleva, 1965
paracontractus Shmeleva, 1974
parelongatus Shmeleva, 1979
pavo (Dana, 1849)
pavoninus Farran, 1936
plumatus Shmeleva, 1965
pseudocontractus Bernard, 1958
styliremis Giesbrecht, 1888
teniculus Andronov, 1973
vivesi Shmeleva, 1979
- Delius*
nudus (Sewell, 1929)
- Ischnocalanus*
equalicauda (Bernard, 1958)
gracilis (Tanaka, 1956)
plumulosus (Claus, 1863)
tenuis (Farran, 1926)
- Paracalanus*
aculeatus Giesbrecht, 1888
denudatus Sewell, 1929
indicus Wolfenden, 1905
nanus Sars, 1907
parvus (Claus, 1863)
- Family Parapontellidae**
Parapontella
brevicornis (Lubbock, 1857)
- Family Phaenidae**
Brachycalanus
atlanticus (Wolfenden, 1904)
Cephalophanes
frigidus Wolfenden, 1911
refulgens Sars, 1907
Cornucalanus
chelifer (Thompson, 1903)
indicus Sewell, 1929
simplex Wolfenden, 1905
Onchocalanus
affinis With, 1915
cristatus (Wolfenden, 1904)
hirtipes Sars, 1905
magnus (Wolfenden, 1906)
trigoniceps Sars, 1905
Phaenna
spinifera Claus, 1863
Talacalanus
greeni (Farran, 1905)
Xanthocalanus
agilis Giesbrecht, 1892
borealis Sars, 1900
claviger (T. Scott, 1909)
difficilis Grice & Hulsemann, 1965
echinatus Sars, 1907
fallax Sars, 1925
giesbrechti I.C. Thompson, 1903
incertus Sars, 1920
minor Giesbrecht, 1892
mixtus Sars, 1920
muticus Sars, 1905
obtusus Farran, 1905
pinguis Farran, 1905
profundus Sars, 1907
propinquus Sars, 1902
subagilis Wolfenden, 1904

<i>tenuiremis</i> T. Scott, 1909	
Family Phyllopodidae	
<i>Phyllopus</i>	
<i>aequalis</i> Sars, 1920	<i>obtusifrons</i> Sars, 1905
<i>bidentatus</i> Brady, 1883	<i>rotunda</i> Wolfenden, 1906
<i>helgae</i> Farran, 1908	<i>Amallothrix</i>
<i>impar</i> Farran, 1908	<i>arcuata</i> (Sars, 1920)
<i>muticus</i> Sars, 1907	<i>emarginata</i> (Farran, 1905)
Family Pontellidae	
<i>Anomalocera</i>	
<i>patersoni</i> Templeton, 1837	<i>falcifer</i> (Farran, 1926)
<i>Calanopia</i>	<i>farrani</i> Rose, 1942
<i>americana</i> F. Dahl, 1894	<i>gracilis</i> (Sars, 1905)
<i>elliptica</i> (Dana, 1849)	<i>paravalida</i> Brodsky, 1950
<i>media</i> Gurney, 1927	<i>pseudoarcuata</i> (Park, 1970)
<i>Labidocera</i>	<i>robusta</i> (T. Scott, 1894)
<i>acuta</i> (Dana, 1849)	<i>sarsi</i> Rose, 1942
<i>acutifrons</i> (Dana, 1849)	<i>Archescolecithrix</i>
<i>aestiva</i> Wheeler, 1901	<i>auropecten</i> (Giesbrecht, 1892)
<i>brunescens</i> (Czerniavski, 1868)	<i>Heteramalla</i>
<i>detruncata</i> (Dana, 1849)	<i>sarsi</i> Roe, 1975
<i>kroyeri</i> (Brady, 1883)	<i>Landrumius</i>
<i>madurae</i> A. Scott, 1909	<i>insignis</i> (Sars, 1920)
<i>nerii</i> (Krøyer, 1849)	<i>Lophothrix</i>
<i>orsinii</i> Giesbrecht, 1889	<i>frontalis</i> Giesbrecht, 1895
<i>pavo</i> Giesbrecht, 1889	<i>humilifrons</i> Sars, 1905
<i>wollastoni</i> (Lubbock, 1857)	<i>latipes</i> (T. Scott, 1894)
<i>Pontella</i>	<i>quadrispinosa</i> Wolfenden, 1911
<i>atlantica</i> (Milne Edwards, 1840)	<i>Mixtocalanus</i>
<i>lobiancoi</i> (Canu, 1888)	<i>alter</i> (Farran, 1929)
<i>mediterranea</i> (Claus, 1863)	<i>Racovitzanus</i>
<i>securifer</i> Brady, 1883	<i>antarcticus</i> Giesbrecht, 1902
<i>spinipes</i> Giesbrecht, 1889	<i>levis</i> Tanaka, 1961
<i>Pontellina</i>	<i>Scaphocalanus</i>
<i>elegans</i> Claus, 1892	<i>affinis</i> (Sars, 1905)
<i>plumata</i> (Dana, 1849)	<i>amplius</i> Park, 1970
<i>Pontellopsis</i>	<i>angulifrons</i> Sars, 1920
<i>brevis</i> (Giesbrecht, 1889)	<i>bogorovi</i> Brodsky, 1955
<i>regalis</i> (Dana, 1849)	<i>brevicornis</i> (Sars, 1900)
<i>villosa</i> Brady, 1883	<i>brevirostris</i> Park, 1970
Family Pseudocyclopidae	
<i>Pseudocyclops</i>	
<i>arguinensis</i> Andronov, 1986	<i>curtus</i> (Farran, 1926)
<i>crassiremis</i> Brady, 1872	<i>echinatus</i> (Farran, 1905)
<i>mirus</i> Andronov, 1986	<i>elongatus</i> A. Scott, 1909
<i>obtusatus</i> Brady & Robertson, 1873	<i>insignis</i> Brodsky, 1950
<i>pumilis</i> Andronov, 1986	<i>invalidus</i> Hure & Scotto di Carlo, 1968
<i>umbraticus</i> Giesbrecht, 1893	<i>longifurca</i> (Giesbrecht, 1888)
Family Pseudocyclopiidae	
<i>Pseudocyclopia</i>	
<i>caudata</i> T. Scott, 1894	<i>magnus</i> (T. Scott, 1894)
<i>crassicornis</i> T. Scott, 1892	<i>medius</i> (Sars, 1907)
<i>giesbrechti</i> Wolfenden, 1902	<i>pseudobrevirostris</i> Schulz, 1987
<i>insignis</i> Andronov, 1986	<i>similis</i> Hure & Scotto di Carlo, 1968
<i>minor</i> T. Scott, 1892	<i>subbrevicornis</i> (Wolfenden, 1911)
<i>Stygocyclopia</i>	
<i>balearica</i> Jaume & Boxshall, 1995	<i>subcurtus</i> Park, 1970
<i>Thompsonopia</i>	
<i>mediterranea</i> Jaume, Fosshagen & Iliffe, 1999	<i>Scolecithricella</i>
<i>stephoidea</i> (Thompson, 1895)	<i>abyssalis</i> (Giesbrecht, 1888)
Family Pseudodiaptomidae	
<i>Calanipedia</i>	<i>canariensis</i> Roe, 1975
<i>aquaedulcis</i> Krtschagin, 1873	<i>curticauda</i> A. Scott, 1909
Family Ridgewayiidae	
<i>Exumella</i>	<i>dentata</i> (Giesbrecht, 1892)
<i>mediterranea</i> Jaume & Boxshall, 1995	<i>lobata</i> Sars, 1920
<i>Ridgewayia</i>	<i>lobophora</i> Park, 1970
<i>minorcaensis</i> Razouls & Carola, 1996	<i>marquesa</i> Vervoort, 1965
Family Ryocalanidae	
<i>Ryocalanus</i>	<i>minor</i> (Brady, 1883)
<i>admirabilis</i> Andronov, 1992	<i>obtusifrons</i> (Sars, 1905)
Family Scolecitrichidae	
<i>Amalophora</i>	<i>orientalis</i> Mori, 1937
	<i>ovata</i> (Farran, 1905)
	<i>profunda</i> (Giesbrecht, 1892)
	<i>propinqua</i> Sars, 1920
	<i>spinata</i> Tanaka, 1962
	<i>unispinosa</i> Grice & Hulsemann, 1965
	<i>vittata</i> (Giesbrecht, 1892)
	<i>Scolecithrix</i>
	<i>bradyi</i> Giesbrecht, 1888
	<i>danae</i> (Lubbock, 1856)
	<i>fowleri</i> Farran, 1926
	<i>laminata</i> Farran, 1926
	<i>longipes</i> Giesbrecht, 1892
	<i>tenuiserrata</i> Giesbrecht, 1892
	<i>valens</i> Farran, 1926
	<i>Scopalatum</i>
	<i>dubia</i> (T. Scott, 1894)
	<i>farrani</i> Roe, 1975

gibbera Roe, 1975
Scottocalanus
corystes Owre & Foya, 1967
heleneae (Lubbock, 1856)
persecans (Giesbrecht, 1895)
securifrons (T. Scott, 1894)
thomasi A. Scott, 1909
thori With, 1915

Family Spinocalanidae

Damkaeria
falcifera Fosshagen, 1983
Mimocalanus
brodskyi Razouls, 1974
cultifer Farran, 1908
inflatus Davis, 1949
major Sars, 1920
nudus Farran, 1908
ovalis (Grice & Hulsemann, 1965)
Monacilla
tenera Sars, 1907
typica Sars, 1905
Spinocalanus
abruptus Grice & Hulsemann, 1965
abyssalis Giesbrecht, 1888
angusticeps Sars, 1920
brevicaudatus Brodsky, 1950
caudatus Sars, 1920
hirtus Sars, 1907
horridus Wolfenden, 1911
longicornis Sars, 1900
magnus Wolfenden, 1904
oligospinosus Park, 1970
polaris Brodsky, 1950
spinosus Farran, 1908
validus Sars, 1920
Teneriforma
meteorae Schulz, 1989
naso (Farran, 1936)
pentarichodes Schulz, 1989

Family Stephidae

Parastephos
pallidus Sars, 1902
Stephos
balearensis Carola & Razouls, 1996
canariensis Boxshall, Stock, Sanchez, 1990
cryptospinosus Zagami, Campolmi & Costanzo, 2000
fultoni T & A. Scott, 1898
gyrans (Giesbrecht, 1892)
lamellatus Sars, 1902
margalefi Riera, Vives & Gili, 1991
minor T. Scott, 1892
rustadi Stromgren, 1969
scotti Sars, 1902

Family Temoridae

Eurytemora
affinis (Poppe, 1880)
americana Williams, 1906
grimmi (Sars, 1897)
lacustris (Poppe, 1887)
raboti Richard, 1897
velox (Lilljeborg, 1853)
Temora
longicornis Müller, 1792)
stylifera (Dana, 1849)
turbinata (Dana, 1849)
Temoropia
mayumbaensis T. Scott, 1894
minor Deevey, 1972
setosa Schulz, 1986

Family Tharybidae

Neoscolecithrix
farrani Smirnov, 1935
koehleri Canu, 1896
watersae (Grice, 1972)

Pseudohaenna
typica Sars, 1902
Rythabis
atlantica Schulz, 1995
Tharybיס
angularis Schulz, 1995
crenata Schulz, 1995
groenlandicus (Tupitzky, 1982)
macrophthalma Sars, 1902
minor Schulz, 1981
Undinella
brevipes Farran, 1908
oblonga Sars, 1900
strini Grice, 1971

Harpacticoida

Compiled by Rony Huys

North-west European data on the primarily benthic harpacticoid copepods are derived from Huys, Gee, Moore & Hamond (1996) and an unpublished checklist prepared by these authors, from the monograph by Lang (1948) for Baltic and Arctic records and from several studies by Gerd Schriever for North Atlantic deepwater species. Mediterranean records postdating Lang (1948) have been compiled primarily from Bodin (1964) and Soyer (1971) for the Western Mediterranean and from Noodt (1955) and Por (1964) for the Eastern Mediterranean. Data on harpacticoids from the Black Sea basin were almost exclusively drawn from Apostolov & Marinov's (1988) *Fauna Bulgarica*. A research database maintained at the Natural History Museum was used to identify missing entries and we envisage the checklist to be over 95% complete. The family level classification is based on Huys, Gee, Moore & Hamond (1996) and includes recent additions of two anchialine cave-dwelling families by Huys (1996) and Huys & Iliffe (1998).

Order Harpacticoida**Family Adenopleurellidae**

Miroslavia
longicaudata Apostolov, 1980
Sarsocletodes
typicus (Sars, 1920)

Family Aegisthidae

Aegisthus
aculeatus Giesbrecht, 1891
mucronatus Giesbrecht, 1891
spinulosus Farran, 1905

Family Ambunguipedidae

Ambunguipes
rufocincta (Norman in Brady, 1880)

Family Ameiridae

Ameira
atlantica subsp. *atlantica* Noodt, 1958
atlantica subsp. *mediterranea* Kunz, 1975
divagans subsp. *divagans* Nicholls, 1939
divagans subsp. *pontica* Marinov, 1973
exigua T Scott, 1894
faroerensis Schriever, 1982
listensis Mielke, 1973
longipes Boeck, 1865
lusitanica Galhano, 1970
minuta Boeck, 1865
parascotti Chislenko, 1977
parvula subsp. *nana* Willey, 1935
parvula subsp. *parvula* (Claus, 1866)
parvula subsp. *tenuiseta* Willey, 1929
parvuloides Lang, 1965
pusilla T Scott, 1903

- reducta* Petkovski, 1954
scotti Sars, 1911
speciosa Monard, 1935
tenuicornis T Scott, 1902
usitata Klie, 1950
Ameiopsis
angulifera Sars, 1911
brevicornis Sars, 1907
longicornis Sars, 1907
minor (Sars, 1920)
mixta Sars, 1907
nobilis Sars, 1911
Ameiropsyllus
arianus (Monard, 1928)
monardi Bodin, 1979
Anoplosoma
sordidum Sars, 1911
stryx Por, 1964
Filexilia
attenuata (Thompson, 1893)
azorica Conroy-Dalton & Huys, 1997
brevipes (Kunz, 1954)
gravellicola (Guille & Soyer, 1966)
intermedia (Galhano, 1970)
longifurca (Bodin, 1964)
marinovi Conroy-Dalton & Huys, 1997
pestae (Petkovski, 1955)
Haifameira
archibenthicoa Por, 1964
Interleptomesochra
attenuata (A Scott, 1896)
elongata (Bozic, 1955)
eulitoralis (Noodt, 1952)
noodti Galhano, 1968
tenuicornis (Sars, 1911)
Leptomesochra
confluens Sars, 1911
infima Monard, 1928
macintoshii (T & A Scott, 1895)
nichollsii Noodt, 1955
theodoridis Soyer, 1966
Limameira
mediterranea Soyer, 1975
Malacopsyllus
fragilis Sars, 1911
Nitokra
affinis subsp. *affinis* Gurney, 1927
affinis subsp. *californica* Lang, 1965
affinis subsp. *rijekana* Petkovski, 1954
balnearia Por, 1964
baltica Arlt, 1983
cari Petkovski, 1954
divaricata Chappuis, 1923
elegans (T Scott, 1905)
elongata Marcus, 1968
fallaciosa subsp. *baltica* Lang, 1965
fallaciosa subsp. *fallaciosa* Klie, 1937
fragilis Sars, 1905
hamata Bodin, 1970
hibernica subsp. *bulgarica* Apostolov, 1976
hibernica subsp. *hibernica* (Brady, 1880)
lacustris subsp. *azorica* Kunz, 1983
lacustris subsp. *lacustris* (Shmankevich, 1875)
lacustris subsp. *sinoi* Marcus & Por, 1961
mediterranea subsp. *mediterranea* Brian, 1928
mediterranea subsp. *pontica* Apostolov, 1980
pontica Jakubisiak, 1938
pusilla Sars, 1911
reducta subsp. *fluviatilis* Galhano, 1968
reducta subsp. *reducta* Schäfer, 1936
spinipes Boeck, 1865
stygia (Apostolov, 1976)
typica subsp. *adriatica* Petkovski, 1954
typica subsp. *typica* Boeck, 1865
Parameiropolis
rapiens Becker, 1974
Parapseudoleptomesochra
polychaeta (Noodt, 1952)
Parevansula
elegans (Marinov, 1974)
mediterranea Guille & Soyer, 1966
vermiformis Moore, 1976
wellsi (Marinov, 1973)
Praeleptomesochra
africana (Kunz, 1951)
Proameira
arenicola (Lang, 1935)
dubia (Sars, 1920)
echinipes Soyer, 1975
hiddensoensis (Schäfer, 1936)
phaedra (Monard, 1935)
psammophila Wells, 1963
signata Por, 1964
simplex (Norman & T Scott, 1905)
Psammameira
hyalina Noodt, 1952
Pseudameira
antennulata Schriever, 1984
breviseta Klie, 1950
crassicornis Sars, 1911
furcata Sars, 1911
gracilis Sars, 1920
limicola Soyer, 1975
minutissima Monard, 1928
mixta subsp. *adriatica* Apostolov & Petkovski, 1980
mixta subsp. *mixta* Sars, 1920
perplexa Soyer, 1975
reducta Klie, 1950
reflexa (T Scott, 1894)
trisetosa Schriever, 1984
Pseudoleptomesochrella
halophila (Noodt, 1952)
Pseudosarsameira
exilis (T & A Scott, 1894)
Psyllocamptus
eridani Ceccherelli, 1988
minutus subsp. *minutus* Sars, 1911
monachus Chappuis, 1938
propinquus (T Scott, 1896)
Sarsameira
boecki (Lang, 1936)
elongata (Sars, 1909)
giraulti Monard, 1935
longifurcata Becker, 1974
longiremis (T Scott, 1894)
major (Sars, 1907)
parva (Boeck, 1873)
peresi Bodin, 1970
propinqua (T Scott, 1902)
sarsi Lang, 1948
Sicameira
gracilis (A Scott, 1896)
intermedia Marinov, 1973
leptoderma Klie, 1950
Stenocopia
longicaudata subsp. *longicaudata* (T Scott, 1892)
longicaudata subsp. *pontica* Griga, 1962
setosa Sars, 1907
spinosa (T Scott, 1892)
Family Ancorabolidae
Ancorabolus
mirabilis Norman, 1903
Arthropyllus
serratus subsp. *serratus* Sars, 1909
Ceratonotus
coineau Soyer, 1964
pectinatus subsp. *elaphus* Por, 1965
pectinatus subsp. *pectinatus* Sars, 1909
Dorsiceratus
octocornis Drzycimski, 1967

- Echinocletodes*
armatus (T Scott, 1903)
- Echinopsyllus*
normani Sars, 1909
- Laophontodes*
armatus Lang, 1936
bicornis A Scott, 1896
expansus Sars, 1908
gracilipes Lang, 1936
typicus (T Scott, 1894)
- Paralaophontodes*
echinatus (Willey, 1930)
- Family Arenopontiidae**
- Arenopontia*
- nesiae* Cottarelli, 1975
problematica Masry, 1970
stygia Noodt, 1955
subterranea Kunz, 1937
acantha subsp. *acantha* Chappuis, 1954
phreatica Cottarelli, Bruno & Venanzetti, 1995
 speluncae Cottarelli, Bruno & Venanzetti, 1995
- Pararenopontia*
breviarticulata (Mielke, 1975)
- Family Argestidae**
- Argestes*
mollis Sars, 1910
- Argestigens*
abyssalis Becker, 1979
glacialis Lang, 1936
- Corallicletodes*
boutierei Soyer, 1966
- Eurypletodes*
laticauda (Boeck, 1873)
rectangulatus Lang, 1936
serratus Sars, 1920
abyssi Lang, 1936
aculeatus Sars, 1920
arcticus Lang, 1936
echinatus Lang, 1936
ephippiger Por, 1964
irelandica Roe, 1959
latus (T Scott, 1892)
major Sars, 1909
minutus Sars, 1920
oblongus Sars, 1920
parasimilis Por, 1959
petiti Soyer, 1964
profundus Becker, 1979
quadrispinosa Schriever, 1986
similis (T Scott, 1895)
- Fultonia*
bougisi Soyer, 1964
gascognensis Bodin, 1968
hirsuta T Scott, 1902
- Hemicletodes*
typicus Lang, 1936
- Hypalocletodes*
aberrans (Marinov, 1973)
- Leptocletodes*
debilis Sars, 1920
- Mesocletodes*
abyssicola (T & A Scott, 1901)
ameliae Soyer, 1975
arenicola Noodt, 1952
bathybia Por, 1964
bodini Soyer, 1975
brevifurca Lang, 1936
carpinei Soyer, 1975
duosetosus Schriever, 1985
faroerensis Schriever, 1985
fladensis Wells, 1965
glaber Por, 1964
guillei Soyer, 1964
inermis Sars, 1920
irrasus (T & A Scott, 1894)
- katharinae* Soyer, 1964
kunzi Schriever, 1985
monensis (Thompson, 1893)
parabodini Schriever, 1983
quadrispinosa Schriever, 1985
robustus Por, 1965
sarsi Becker, 1979
soyeri Bodin, 1968
thieli Schriever, 1985
trisetosa Schriever, 1983
variabilis Schriever, 1983
- Neoargestes*
incertus Becker, 1979
variabilis Drzycimski, 1967
- Odiliacletodes*
gracilis Soyer, 1964
- Parargestes*
tenuis (Sars, 1921)
- Family Balaenophilidae**
- Balaenophilus*
unisetus Aurivillius, 1879
- Family Canthocamptidae**
- Bathycampus*
minutus (Wells, 1965)
- Boreolimella*
dubia (Wells, 1965)
nympha (Por, 1964)
- CletoCampus*
confluens (Schmeil, 1894)
retrogressus Shmankevich, 1875
- Hemimesochra*
clavularis Sars, 1920
- Heteropsyllus*
confluens Soyer, 1975
curticaudatus T Scott, 1894
exiguus (Sars, 1911)
major (Sars, 1920)
masculus Kunz, 1971
meridionalis Soyer, 1975
nanus (Sars, 1920)
rostratus subsp. *rostratus* (Sars, 1920)
- Itunella*
intermedia Apostolov, 1975
muelleri (Gagern, 1922)
tenuiremis (T Scott, 1893)
- Mesochra*
aestuarii subsp. *aestuarii* Gurney, 1921
aestuarii subsp. *baltica* Schäfer, 1936
anomala Klie, 1950
armoricana Monard, 1935
heldti Monard, 1935
inconspicua (T Scott, 1900)
lilljeborgii Boeck, 1865
lindbergi Petkovski, 1964
pestai Lang, 1948
pontica Marcus, 1965
pygmaea (Claus, 1863)
rapiens (Schmeil, 1894)
reducta Klie, 1950
rostrata Gurney, 1927
schmidti Mielke, 1974
timsae Gurney, 1927
xenopoda Monard, 1935
- Mesopsyllus*
atargatis Por, 1960
secundus (Wells, 1965)
- Nannomesochra*
aruensis (Brian, 1925)
zavodniki Petkovski & Apostolov, 1974
- Parepactophanes*
dubia Noodt, 1958
minuta Kunz, 1935
- Poria*
derketo (Por, 1964)
- Psammocampus*

- axi* Mielke, 1975
Pusillargillus
nixe (Por, 1964)
Family Canuellidae
Brianola
stebleri (Monard, 1926)
Canuella
furcigera Sars, 1903
perplexa T & A Scott, 1893
pontica Apostolov, 1971
Canuellina
insignis Gurney, 1927
Canuellopsis
mediterranea Soyer, 1966
swedmarki Por, 1964
typica Lang, 1936
Nathaniella
reichi (Por, 1964)
Scottolana
longipes (Thompson & A Scott, 1903)
bulbosa (Por, 1964)
Sunaristes
paguri Hesse, 1867
Family Cerviniidae
Cervinia
bradyi (Norman in Brady (1878, 1878))
pilosa Lang, 1948
synarthra Sars, 1910
Cerviniella
lagarderei Bodin, 1968
langi Bodin, 1968
talpa (Por, 1964)
Cerviniopsis
clavicornis Sars, 1903
intermedia Lang, 1936
langi Soyer, 1970
longicaudata Sars, 1903
stylicaudata Lang, 1936
Eucanuella
langi Por, 1964
reticulata Soyer, 1970
spinifera T Scott, 1901
Hemicervinia
stylifera (Thompson, 1893)
Pontostratiotes
abyssicola Brady, 1883
barnetti Dinet, 1978
cecilae Dinet, 1978
denticulatus Dinet, 1978
fontani Dinet, 1981
gladius Dinet, 1978
horrida Brotskaya, 1959
lubricus Dinet, 1978
minor Dinet, 1981
scotti Brotskaya, 1959
uxoris Dinet, 1978
vasconiensis Dinet, 1978
vitielloi Dinet, 1978
vivierae Dinet, 1978
Stratiopontotes
mediterraneus Soyer, 1970
Family Cletodidae
Acrenhydrosoma
perplexum (T Scott, 1899)
Cletodes
brucei T & A Scott, 1901
carthaginiensis Monard, 1935
endopodita (Schriever, 1984)
latirostris Drzycimski, 1967
limicola Brady, 1872
longicaudatus (Boeck, 1873)
longifurca Lang, 1948
pusillus Sars, 1920
reductus Moore, 1978
reyssi Soyer, 1964
setosus Marinov & Apostolov, 1985
smirnovi Bodin, 1970
tenuipes T Scott, 1897
Enhydrosoma
caeni Raibaut, 1965
curticauda Boeck, 1873
curvirostre (T Scott, 1894)
gariene Gurney, 1930
hopkinsi Lang, 1965
longicauda Marinov & Apostolov, 1983
longifurcatum Sars, 1909
micrurum Monard, 1928
migoti Monard, 1926
pontica Jakubisiak, 1938
propinquum (Brady & Robertson in Brady, 1880)
sarsi (T Scott, 1905)
sordidum Monard, 1926
wellsi Bodin, 1968
Enhydrosomella
setiensis Raibaut, 1965
staufferi Monard, 1935
Limnocletodes
behningi Borutzky, 1926
Nannopodella
denisi Monard, 1928
Paracrenhydrosoma
normani Gee, 1999
Pontocletodes
ponticus Apostolov, 1980
Strongylacron
buchholzi (Boeck, 1873)
Stylicletodes
longicaudatus (Brady & Robertson in Brady, 1880)
minutus Bodin, 1968
oligochaeta Bodin, 1968
reductus Wells, 1965
Family Cletopsyllidae
Isocletopsyllus
quartus (Soyer, 1966)
tertius (Por, 1964)
Family Clytemnestridae
Clytemnestra
rostrata (Brady, 1883)
scutellata Dana, 1848
Family Cristacoxidae
Noodtorthopsyllus
psammophilus (Noodt, 1955)
Family Cylindropsyllidae
Boreopontia
heipi Willem, 1981
Cylindropsyllus
govaerei Huys & Willem, 1993
ibericus Huys & Willem, 1993
kunzi Huys, 1988
laevis Brady, 1880
remanei Kunz, 1949
Cylinula
arganoi Cottarelli & Venanzetti, 1989
Evansula
incerta (T Scott, 1892)
pygmaea (T Scott, 1903)
Selenopsyllus
abyssalis (Becker, 1979)
profundus (Becker, 1979)
Stenocaris
baltica Arlt, 1983
gracilis Sars, 1909
klei Kunz, 1936
minor (T Scott, 1892)
minuta Nicholls, 1935
pontica Chappuis & Serban, 1953
pygmaea Noodt, 1955
Stenocaropsis

- pristina* (Wells, 1968)
similis Cottarelli & Venanzetti, 1989
valkanovi (Marinov, 1974)
- Willemia*
calceola Huys & Conroy-Dalton, 1993
- Family Darythompsoniidae**
- Darythompsonia*
neglecta Redeke, 1953
fairliensis (T Scott, 1899)
- Horsiella*
brevicornis Douwe, 1904
- Leptocaris*
biscayensis (Noodt, 1955)
canariensis Lang, 1965
igneus Cottarelli & Baldari, 1982
insularis (Noodt, 1958)
marinus (Por, 1964)
minimus (Jakobi, 1954)
minutus T Scott, 1899
trisetosus (Kunz, 1935)
- Family Diosaccidae**
- Amonardia*
arctica (T Scott, 1899)
normani (Brady, 1872)
pelophila Por, 1964
phyllopus (Sars, 1906)
similis (Claus, 1866)
- Amphiascoides*
breviarticulatus Kunz, 1983
brevifurca (Czerniavski, 1868)
debilis (Giesbrecht, 1881)
dispar (T & A Scott, 1894)
golikovi Chislenko, 1977
koltuni Chislenko, 1977
limicola (Brady, 1899)
littoralis (T Scott, 1903)
nanoides (Sars, 1911)
nanus (Sars, 1906)
neglecta (Norman & T Scott, 1905)
sterilis (Monard, 1926)
subdebilis (Willey, 1935)
- Amphiascopis*
cinctus (Claus, 1866)
thalestroides (Sars, 1911)
- Amphiascus*
amblyops Sars, 1911
angustipes Gurney, 1927
caudaespinosus Brian, 1927
gauthieri Monard, 1936
gracilis Lang, 1936
graciloides Klie, 1950
hirtus Gurney, 1927
humphriesi Roe, 1958
longarticulatus Marcus, 1974
minutus (Claus, 1863)
pacificus Sars, 1905
paracaudaespinosus Roe, 1958
parvus Sars, 1906
perplexus (Thompson & A Scott, 1903)
polapinvus Marcus, 1966
propinquus Sars, 1906
sinuatus Sars, 1906
tenellus Sars, 1906
tenuiremis (Brady & Robertson in Brady (1880),
1880)
ultimus Monard, 1928
varians (Norman & T Scott, 1905)
- Bulbamphiascus*
denticulatus (Thompson, 1893)
imus (Brady, 1872)
minutus Dinet, 1971
- Dactylopodamphiascopis*
latifolius (Sars, 1909)
- Diosaccopsis*
rubeus Brian, 1925
- Diosaccus*
tenuicornis (Claus, 1863)
truncatus Gurney, 1927
varicolor subsp. *biarticulatus* (Monard, 1924)
varicolor subsp. *pentasetosus* (Noodt, 1955)
varicolor subsp. *varicolor* (Farran, 1913)
- Eoschizopera*
sylensis (Mielke, 1973)
gligici (Petkovski, 1957)
- Haloschizopera*
bathyalis Schriever, 1984
bulbifera (Sars, 1911)
clotensis Moore & O'Reilly, 1993
conspicua Por, 1964
exigua (Sars, 1906)
latisetifera Marinov, 1973
lionensis Moore & O'Reilly, 1993
marmarae Noodt, 1955
mathoi (Monard, 1935)
minima Por, 1964
noodti Bodin, 1968
nuditerga Moore & O'Reilly, 1993
pauciseta Por, 1959
pygmaea (Norman & T Scott, 1905)
- Ialysus*
rufus Brian, 1927
- Metamphiascopis*
banyulensis (Monard, 1928)
hirsutus subsp. *hirsutus* (Thompson & A Scott, 1903)
- Paramphiascella*
bodini Marcotte, 1974
brucei (T & A Scott, 1901)
bulifer Guille & Soyer, 1966
couilli Marcotte, 1974
delamarei Guille & Soyer, 1966
faurei Bodin, 1968
fulvofasciata Rosenfield & Coull, 1974
hispidia (Brady, 1880)
hyperborea (T Scott, 1903)
intermedia (T Scott, 1897)
invaginata (Monard, 1926)
langi (Monard, 1936)
mediterranea Lang, 1948
roberti (Monard, 1935)
robinsoni (A Scott, 1902)
sirbonica Por, 1973
vararenensis (T Scott, 1903)
- Paramphiascopis*
giesbrechti (Sars, 1906)
longirostris (Claus, 1863)
pallidus (Sars, 1906)
paromolae Soyer, 1973
soyeri Lang, 1965
triarticulatus Moore, 1976
- Pararobertsonia*
abyssi (Boeck, 1873)
- Protopsammotopa*
norvegica Geddes, 1968
- Psammotopa*
chappuisi Noodt, 1955
phyllotosa (Noodt, 1952)
polyphilla Noodt, 1955
vulgaris Pennak, 1942
- Pseudamphiascopis*
attenuatus (Sars, 1906)
herdmani (A Scott, 1896)
- Pseudodiosaccopsis*
mesogaea Por, 1964
rufescens (Brian, 1925)
- Pseudodiosaccus*
propinquus (T & A Scott, 1893)
- Rhyncholagena*
lagenirostris (Sars, 1911)
levantina Por, 1964
pestai subsp. *pestai* (Monard, 1935)

- spinifer* (Farran, 1913)
Robertgurneya
dictydiophora (Monard, 1924)
ecaudata (Monard, 1936)
erythraeus (A Scott, 1902)
ilieveensis (Monard, 1935)
intermedia Bozic, 1955
oligochaeta Noodt, 1955
remanei Klie, 1950
rostrata (Gurney, 1927)
similis subsp. *bulbampliascoides* Noodt, 1955
similis subsp. *similis* (A Scott, 1896)
simulans (Norman & T Scott, 1905)
soyeri Apostolov, 1974
spinulosa (Sars, 1911)
Robertsonia
angolensis (Monard, 1934)
diademata Monard, 1926
knoxi subsp. *kieliensis* Becker, 1970
knoxi subsp. *knoxi* (Thompson & A Scott, 1903)
monardi (Klie, 1937)
propinquia (T Scott, 1894)
salsa Gurney, 1927
tenuis (Brady & Robertson in Brady (1880, 1880))
Schizopera
arconae Arlt, 1983
chaetosa Petkovski, 1954
meridionalis subsp. *listensis* Mielke, 1975
meridionalis subsp. *meridionalis* Petkovski, 1954
monardi Petkovski, 1955
ornata Noodt & Purasjoki, 1953
pratensis Noodt, 1958
baltica Lang, 1965
borutzkyi Monchenko, 1967
brusinae Petkovski, 1954
cicolanii Galassi & Pesce, 1988
clandestina (Klie, 1924)
compacta De Lint, 1922
jugurtha subsp. *stephanidesi* Pesta, 1938
kunzi Apostolov, 1967
langi Petkovski, 1954
minuta Noodt, 1955
nana Noodt, 1955
neglecta Akatova, 1935
parvula Noodt, 1955
petkovski Apostolov, 1971
pontica Chappuis & Serban, 1953
elatensis Kahan & Bar-El, 1982
Schizoperopsis
arenicola (Chappuis & Serban, 1953)
varnensis (Apostolov, 1967)
Stenelia
adriatica Marinov & Apostolov, 1981
arctica (T Scott, 1899)
bocqueti Soyer, 1972
caulerpae (Por, 1964)
coineauae Soyer, 1972
confluens Lang, 1948
cornuta Lang, 1936
elisabethae Por, 1960
giesbrechti (T & A Scott, 1896)
hanstromi Lang, 1948
incerta Por, 1964
intermedia Marinov & Apostolov, 1981
islandica Schriever, 1982
longicaudata Boeck, 1873
magnacaudata Monard, 1928
mastigochaeta Wells, 1965
minuta (A Scott, 1902)
noodti Schriever, 1982
normani subsp. *normani* (T Scott, 1905)
normani subsp. *polluta* Monard, 1928
palustris subsp. *bispinosa* Bodin, 1970
palustris subsp. *palustris* (Brady, 1869)
pygmaea (Brady, 1905)
- reflexa* (Brady & Robertson in Brady (1880, 1880))
saharae Marinov & Apostolov, 1985
tethysensis Monard, 1928
aemula (T Scott, 1893)
curviseta Lang, 1936
gibba Boeck, 1865
proxima Sars, 1906
hirsuta Thompson, 1893
Teissierella
massiliensis (Bodin, 1964)
pontica (Apostolov, 1968)
salamboi Monard, 1935
Typhlamphiascus
blanchardi (T & A Scott, 1895)
bouligandi Soyer, 1972
confusus (T Scott, 1902)
drachi Soyer, 1963
gracilis Por, 1963
lamellifer subsp. *lamellifer* (Sars, 1911)
lutincola Soyer, 1963
typhloides (Sars, 1911)
typhlops (Sars, 1906)
- Family Ectinosomatidae**
- Arenosetella*
duriensis Galhano, 1970
germanica subsp. *germanica* Kunz, 1937
incerta Chappuis, 1954
littoralis Bodin, 1979
longiseta Kunz, 1983
tenuissima (Klie, 1929)
- Bradya*
congenera Sars, 1920
furcata Sars, 1920
macrochaeta Sars, 1920
minutiseta Soyer, 1973
proxima T Scott, 1912
scotti Sars, 1920
simulans Sars, 1920
theodori Soyer, 1973
typica Boeck, 1873
atlantica Bodin, 1968
bodini Bozic, 1980
confluens Lang, 1936
dilatata Sars, 1904
pugiochaeta Arlt, 1983
- Bradyellopsis*
aruipinensis Steuer, 1941
briani Steuer, 1941
subniger Brian, 1925
tumidus Brian, 1925
- Ectinosoma*
breviarticulatum Lang, 1965
californicum Lang, 1965
compressum Sars, 1920
dentatum Steuer, 1940
litorale (Noodt, 1958)
mediterraneum Kunz, 1975
melaniceps Boeck, 1865
normani T & A Scott, 1896
obtusum Sars, 1920
paradentatum Bozic, 1965
pruovi Soyer, 1972
reductum subsp. *listensis* Mielke, 1975
reductum subsp. *reductum* Bozic, 1955
soyeri Apostolov, 1975
tenuipes T & A Scott, 1896
vervoorti Soyer, 1972
- Ectinosomella*
nitidula Sars, 1910
- Ectinosomoides*
pectinatus (Chappuis, 1954)
- Halectinosoma*
abrau (Krichagin, 1877)
abyssicola Bodin, 1968
angulifrons (Sars, 1919)

- argyllensis* Clément & Moore, 1995
armiferum (T & A Scott, 1896)
bodotriaensis Clément & Moore, 1999
brevirostre (Sars, 1904)
brittanicum Clément & Moore, 1999
brunneum (Brady, 1905)
canaliculatum (Por, 1964)
chislenki Clément & Moore, 1995
chrystalli (T Scott, 1894)
clavatum (Sars, 1920)
cooperatum Bodin, Bodiou & Soyer, 1971
crenulatum Clément & Moore, 1995
curticorne (Boeck, 1873)
denticulatum Clément & Moore, 1995
diops (Por, 1964)
distinctum (Sars, 1920)
elongatum (Sars, 1904)
erythrops (Brady, 1880)
gascognense Bodin, 1968
gothiceps (Giesbrecht, 1881)
gracile (T & A Scott, 1896)
herdmani (T & A Scott, 1896)
huysi Clément & Moore, 1999
inopinatum (Por, 1964)
itoi Clément & Moore, 1999
longicorne (T & A Scott, 1896)
mixtum (Sars, 1904)
monardi Soyer, 1973
neglectum (Sars, 1904)
oblongum (Kunz, 1949)
paraspinicaua Bodin, 1979
pilosum Clément & Moore, 1999
proximum (Sars, 1919)
pseudosarsi Clément & Moore, 1995
pterinum Moore, 1974
rouchi Soyer, 1973
similidistinctum Lang, 1965
smirnovi (Chislenko, 1967)
spinicauda (Wells, 1961)
tenerum (Sars, 1920)
tenuireme (T & A Scott, 1896)
travei Soyer, 1973
valeriae Soyer, 1973
Halophytophilus
fusiformis Brian, 1918
similis Lang, 1948
spinicornis Sars, 1920
Hastigerella
abbotti Lang, 1965
bodini Apostolov, 1974
bozici Soyer, 1974
chappuisi Soyer, 1974
leptoderma (Klie, 1929)
meridionalis (Chappuis, 1954)
monniotae (Guille & Soyer, 1966)
psammae (Noodt, 1955)
scheibeli Mielke, 1975
soyeri Bodin, 1976
unisetosa (Wells, 1965)
Klieosoma
spinosum (Hicks & Schriever, 1983)
trarticulatum (Klie, 1949)
Lineosoma
iscensis Wells, 1965
Microsetella
norvegica (Boeck, 1865)
rosea (Dana, 1947)
Noodtiella
arenosetelloides (Noodt, 1958)
gracile Mielke, 1975
lusitanica Wells, 1965
wellsi Apostolov, 1974
Peltobradya
bryozoophila Médioni & Soyer, 1967
Pseudectinosoma
- minor* Kunz, 1935
Pseudobradya
acuta Sars, 1904
ambigua Sars, 1920
attenuata Sars, 1920
banyulensis Soyer, 1974
beduina subsp. *beduina* Monard, 1935
beduina subsp. *faialensis* Kunz, 1983
breviceaudata Soyer, 1974
brevicornis (T Scott, 1893)
digitata Sars, 1920
distinctum Wells, 1968
elegans (T & A Scott, 1896)
exilis Sars, 1920
fusca (T & A Scott, 1896)
hirsuta (T & A Scott, 1896)
kusnezovi Chislenko, 1967
leptognatha Sars, 1920
major (Olofsson, 1917)
maxima Lang, 1935
minor (T & A Scott, 1896)
oligochaeta Chislenko, 1967
parvula Sars, 1920
pectinifera Lang, 1965
peresi Soyer, 1974
psammophila Bodiou, 1974
pulchella Sars, 1920
pygmaea Sars, 1920
rhea Bodin, 1979
robusta Sars, 1910
scabriuscula Sars, 1920
similis (T & A Scott, 1896)
soyeri Bodiou, 1974
spinulosa Soyer, 1974
tenella Sars, 1920
truncatiseta Soyer, 1974
Sigmatidium
difficile Giesbrecht, 1881
noodti Kunz, 1975
parvulum Mielke, 1974
rouchi Kunz, 1975
Tetanopsis
medius Perkins, 1956
smithi Perkins, 1956
- Family Euterpinidae**
Euterpina
acutifrons (Dana, 1847)
- Family Harpacticidae**
Harpacticus
alevtinae Chislenko, 1977
chelifer (OF Müller, 1776)
compsonyx Monard, 1926
depressus Boeck, 1865
flexulosus Ceccherelli, 1988
flexus Brady & Robertson, 1873
giesbrechti Klie, 1927
gracilis Claus, 1863
littoralis Sars, 1910
longiantennata Apostolov & Petkovski, 1980
macrodactylus Fischer, 1860
nicaensis Claus, 1866
obscurus T Scott, 1895
ponticus Marcus, 1967
septentrionalis Klie, 1939
tenellus Sars, 1920
uniaremis subsp. *uniaremis* (Kröyer in Gaimard (1842-1845?, 1842-1845?)
virescens Dana, 1848
- Perissocope**
adiastalus Wells, 1968
xenus (Monard, 1926)
- Tigriopus**
brevicornis (OF Müller, 1776)
fulvus (Fischer, 1860)
- Zaus**

- abbreviatus* Sars, 1904
aurelii Poppe, 1884
caeruleus Campbell, 1929
goodsiri Brady, 1880
spinatus subsp. *spinatus* Goodsir, 1845
- Family Huntemannidae**
- Huntemannia*
jadensis Poppe, 1884
micropus Monard, 1935
- Merlathuntemannia*
arctica Schriever, 1984
atlantica Schriever, 1983
crassa (Por, 1965)
dovpori Bodin, 1968
drzycinskii Soyer, 1970
iberica Becker, 1979
mediterranea Soyer, 1970
pseudomagniceps Schriever, 1983
smirnovi Bodin, 1968
spinosa (Klie, 1939)
triarticulata Schriever, 1984
- Monocletodes*
variana (T Scott, 1903)
- Nannopus*
palustris Brady, 1880
- Pontopolites*
typicus T Scott, 1894
- Pseudocletodes*
vararenensis T & A Scott, 1893
- Talpina*
bifida Schriever, 1984
curticauda (Becker, 1979)
- Family Laophontidae**
- Afrolaophonte*
pori Masry, 1970
- Appanola*
hirsuta (Thompson & A Scott, 1903)
- Archesola*
hamondi Huys & Lee, 2000
longiremis (T Scott, 1905)
typhlops (Sars, 1908)
typhlops subsp. *pontoica* (Por, 1959)
- Asellopsis*
bacesqui Por, 1959
duboscqui Monard, 1926
hispida Brady & Robertson, 1873
intermedia (T Scott, 1895)
penicillata Por, 1964
sarmatica Jakubisiak, 1938
- Bathylaophonte*
azorica Lee & Huys, 1999
faroensis (T Scott, 1902)
- Corbulaseta*
bulligera (Farran, 1913)
- Coullia*
platychelipusoides (Noodt, 1958)
- Echinolaoiphonte*
armiger subsp. *armiger* (Gurney, 1927)
brevispinosa (Sars, 1908)
horrida (Norman, 1876)
minuta Cottarelli & Forniz, 1991
veniliae Cottarelli, Forniz & Bascherini, 1992
- Esola*
bulbifera (Norman, 1911)
profunda Huys & Lee, 2000
- Harrietella*
simulans (T Scott, 1894)
- Hemilaophonte*
janinae Jakubisiak, 1932
- Heterolaophonte*
brevipes Roe, 1958
curvata subsp. *curvata* (Douwe, 1929)
curvata subsp. *micrarthros* Marcus & Por, 1960
denticulata Roe, 1958
furcata Noodt, 1958
- hamondi* Hicks, 1975
letovae Huys, 1990
littoralis (T & A Scott, 1893)
longisetigera Klie, 1950
mendax (Klie, 1939)
minuta (Boeck, 1873)
murmanica Letova, 1982
norvegica Drzycimski, 1968
phyacobates (Monard, 1935)
setigera (Kricagin, 1873)
stroemii subsp. *brevicaudata* (Monard, 1928)
stroemii subsp. *paraminuta* Noodt, 1955
stroemii subsp. *stroemii* (Baird, 1837)
uncinata (Czerniavski, 1868)
- Kleonychocampoides*
remanei Noodt, 1958
- Kleonychocampus*
discipes Noodt, 1958
kliei subsp. *adriaticus* (Petkovski, 1954)
kliei subsp. *confluens* Noodt, 1958
kliei subsp. *kliei* (Monard, 1935)
kliei subsp. *ponticus* Marcus, 1971
ponticus (Serban & Plesa, 1957)
- Laophonte*
adamsiae Raibaut, 1966
aldonae Chislenko, 1977
baltica Klie, 1929
borceai Jakubisiak, 1938
brevifurca Sars, 1920
commensalis Raibaut, 1961
cornuta Philippi, 1840
danversae Hamond, 1969
denticornis T Scott, 1894
depressa T Scott, 1894
dinocerata Monard, 1926
dominicalis Monard, 1935
drachi Médioni & Soyer, 1967
elongata subsp. *barbata* Lang, 1934
elongata subsp. *elongata* Boeck, 1873
elongata subsp. *triarticulata* Monard, 1928
farrani Roe, 1958
foxi Harding, 1956
inopinata T Scott, 1892
inornata A Scott, 1902
longicaudata Boeck, 1865
nordgaardi Sars, 1908
parvula Sars, 1908
parvuloides Monard, 1935
serrata (Claus, 1863)
setosa Boeck, 1865
sima Gurney, 1927
thoracica Boeck, 1865
trilobata Willey, 1929
- Laophontina*
acantha Noodt, 1955
dubia Norman & T Scott, 1905
noodti Kunz, 1983
posidoniae Fiers, 1986
- Lipomelum*
adriaticum (Petkovski, 1955)
- Loureirophonte*
cesareae (Por, 1964)
majacola Fiers, 1993
mediterranea Fiers, 1993
- Onychocampus*
mohammed (Blanchard & Richard, 1891)
- Paralaophonte*
brevirostris (Claus, 1863)
congenera (Sars, 1908)
dieuzeidei (Monard, 1936)
hyperborea (Sars, 1909)
innae Chislenko, 1977
karmensis (Sars, 1911)
macera (Sars, 1908)
maja Petkovski, 1964

- octavia* (Monard, 1935)
ormieresii Raibaut, 1968
perplexa (T Scott, 1899)
quaterspinata (Brian, 1917)
royi (Jakubisiak, 1932)
spitzbergensis Mielke, 1974
taurina (Monard, 1928)
tenera (Sars, 1920)
zimmeri (Douwe, 1929)
- Paronychocampus*
curticaudatus (Boeck, 1865)
nanus (Sars, 1908)
- Pilifera**
gracilis (T Scott, 1903)
- Platyelipus*
laophontoides Sars, 1908
littoralis Brady, 1880
- Platylaophonte*
delamarei Bodin, 1968
- Pontophonte*
grigae Lee & Huys, 1999
leuke (Por, 1959)
- Pseudolaophonte*
glemareci Bodin, 1977
proteus Klie, 1950
spinosa (Thompson, 1893)
- Pseudonychocampus*
abbreviatus (Sars, 1920)
carthyi Hamond, 1968
colomboi Ceccherelli, 1988
koreni (Boeck, 1873)
marinovi Apostolov & Petkovski, 1980
proximus (Sars, 1908)
- Quinquaophonte*
quinquespinosa (Sewell, 1924)
- Troglophonte*
spelaea (Chappuis, 1938)
- Family Laophontopsidae**
- Aculeopsis*
longisetosa Huys & Willems, 1989
- Laophontopsis*
borealis Huys & Willems, 1989
lamellifera (Claus, 1863)
monardi Huys & Willems, 1989
- Family Latiremidiae**
- Delamarella*
arenicola Chappuis, 1954
galateae Cottarelli, 1971
karamani Petkovski, 1957
- Family Leptastacidae**
- Archileptastacus*
aberrans (Chappuis, 1954)
- Arenocaris*
bifida Nicholls, 1935
reducta Huys, 1992
- Leptastacus*
corsicaensis Huys, 1992
kwintei Huys, 1992
laticaudatus Nicholls, 1935
macronyx (T Scott, 1892)
operculatus Masry, 1970, 1970
pygmaeus Huys, 1992
uncinatus Cottarelli & Venanzetti, 1989
- Meliorastacus*
ctenidis Huys & Todaro, 1997
- Minervella*
baccettii Cottarelli & Venanzetti, 1989
perplexa (Wells & Clark, 1965)
- Paraleptastacus*
brevicaudatus CB Wilson, 1932
espinulatus Nicholls, 1935
holsaticus Kunz, 1937
kliei (Gagern, 1923)
monensis Whybrew, 1986
- moorei* Whybrew, 1986
spinicauda (T & A Scott, 1895)
supralitoralis Mielke, 1975
triseta Noodt, 1954
- Psamathea*
brittanica Huys, Bodiou & Bodin, 1996
natarum Cottarelli & Venanzetti, 1989
- Psammastacus*
confuens Nicholls, 1935
- Schizothrix*
pontica (Griga, 1964)
rostrata (Nicholls, 1940)
- Family Leptopontiidae**
- Leptopontia*
curvicauda T Scott, 1902
dowpori Huys & Conroy-Dalton, 1996
flandrica Huys & Conroy-Dalton, 1996
mediterranea Huys & Conroy-Dalton, 1996
punctata Huys & Conroy-Dalton, 1996
- Syrticola*
flandricus Willems & Claeys, 1982
mediterraneus Willems, Claeys & Fiers, 1987
trispinosus (A Scott, 1896)
- Family Longipediidae**
- Longipedia*
brevispinosa Gurney, 1927
coronata Claus, 1863
ferox Kricagin, 1877
helgolandica Klie, 1949
minor T & A Scott, 1893
pirgos Apostolov, 1972
pontica Kricagin, 1877
rosea Sars, 1903
scotti Sars, 1903
weberi A Scott, 1909
- Family Louriniidae**
- Lourinia*
armata (Claus, 1866)
- Family Metidae**
- Laubieria*
corallicola Soyer, 1966
- Metis*
holothuriae (Edwards, 1891)
ignea subsp. *halmyricola* Marcus & Por, 1961
ignea subsp. *ignea* Philippi, 1843
- Family Miraciidae**
- Distioculus*
minor (T Scott, 1894)
- Macrosetella*
gracilis (Dana, 1847)
- Miracia*
efferrata Dana, 1849
- Oculosetella*
gracilis (Dana, 1849)
- Family Neobradyidae**
- Marseinia*
bozici (Bodin, 1968)
ibericus (Becker, 1974)
sarsi (Bodin, 1968)
similis Drzycimski, 1968
typica Drzycimski, 1968
- Neobradya*
pectinifera T Scott, 1892
- Family Normanellidae**
- Normanella*
dubia (Brady & Robertson in Brady (1880, 1880)
minuta (Boeck, 1873)
mucronata Sars, 1909
obscura Lee & Huys, 1999
paratemnifurca Lee & Huys, 1999
quarta Monard, 1935
reducta Noodt, 1955
sarsi Lee & Huys, 1999

- semitica* Monard, 1935
serrata Por, 1959
similis Lang, 1936
tenuifurca Sars, 1909
Sagamiella
aberrans Bodin, 1968
- Family Novocriniidae**
Atergopedia
vetusta Martinez Arbizu & Moura, 1998
- Family Orthopsyllidae**
Orthopsyllus
linearis subsp. *illgi* (Chappuis, 1958)
linearis subsp. *linearis* (Claus, 1866)
linearis subsp. *setosus* Boer, 1971
sarsi Klie, 1941
- Family Paramesochridae**
Apodopsyllus
africanus subsp. *listensis* Mielke, 1975
arenicolus (Chappuis, 1954)
littoralis (Nicholls, 1939)
lynceorum Cottarelli, 1973
melitae Kunz, 1992
spinipes (Nicholls, 1939)
- Diarthrodelta**
convexa Kunz, 1983
lancifera Kunz, 1983
orbiculata Klie, 1949
parorbiculata subsp. *parorbiculata* Wells, 1963
psammophila (Bocquet & Bozic, 1955)
secunda subsp. *secunda* Kunz, 1954
- Kliopsyllus**
atlanticus Kunz, 1983
coelebs (Monard, 1935)
constrictus subsp. *constrictus* (Nicholls, 1935)
constrictus subsp. *orotavae* (Noodt, 1958)
holsticus subsp. *holsticus* (Klie, 1929)
holsticus subsp. *listensis* Mielke, 1984
holsticus subsp. *longicaudatus* Galhano, 1970
laurenticus (Nicholls, 1939)
longifurcatus Scheibel, 1975
major (Nicholls, 1939)
masryi Bodin, 1979
miguelensis Kunz, 1983
paraholsticus Mielke, 1975
perharidiensis (Wells, 1963)
ponticus (Serban, 1959)
pygmaeus (Nicholls, 1939)
- Leptopsyllus**
abyssalis Becker, 1979
celticus Bodin & Jackson, 1987
elongatus Drzycimski, 1967
harveyi Wells, 1963
paratypicus Nicholls, 1939
reductus Lang, 1948
typicus T Scott, 1894
arcticus (Lang, 1936)
- Meiopsyllus**
marinae Cottarelli & Forniz, 1995
- Paramesocha**
acutata Klie, 1934
borealis Geddes, 1981
brevifurca subsp. *brevifurca* Galhano, 1970
brevifurca subsp. *mediterranea* Huys, 1987
dubia T Scott, 1892
helgolandica subsp. *helgolandica* Kunz, 1936
mielkei Huys, 1987
pterocaudata Kunz, 1936
similis Kunz, 1936
- Remanea**
arenicola Klie, 1929
- Scotopsyllus**
intermedius (T & A Scott, 1895)
minutus (Nicholls, 1939)
smirnovi Kunz, 1992
- herdmani* (Thompson & A Scott, 1899)
langi subsp. *continentalis* Kunz, 1992
minor (T & A Scott, 1895)
robertsoni (T & A Scott, 1895)
gigas (Wells, 1965)
- Family Parasteneliidae**
Parastenelia
anglica Norman & T Scott, 1905
hornelli Thompson & A Scott, 1903
ornatissima (Monard, 1935)
reducta Apostolov, 1975
spinosa subsp. *bulbosa* Wells, 1963
spinosa subsp. *spinosa* (Fischer, 1860)
- Family Parastenocarididae**
Parastenocaris
chappuisi Serban, 1960
phyllura Kiefer, 1938
vicesima subsp. *vicesima* Klie, 1934
- Family Peltidiidae**
Alteutha
aberrans Czerniavski, 1868
depressa (Baird, 1837)
interrupta (Goodsir, 1845)
oblonga (Goodsir, 1845)
roeae Hicks, 1982
sarsi Monard, 1924
typica Czerniavski, 1868
- Eupelte*
gracilis Claus, 1860
- Oniscidium*
incertum Haller, 1879
sculptum Haller, 1879
triarticulatum Haller, 1879
- Peltidium*
gracile (Claus, 1889)
purpureum Philippi, 1839
robustum (Claus, 1889)
sacesphorum Monard, 1928
- Zausoscidia*
folii Haller, 1879
- Family Phyllognathopodidae**
Phyllognathopus
viguieri (Maupas, 1892)
- Family Porcellidiidae**
Porcellidium
imbriatum Claus, 1863
lecanoides Claus, 1889
ovatum Haller, 1879
sarsi Bocquet, 1948
tenuicauda Claus, 1860
viride (Philippi, 1840)
- Family Psammopsyllidae**
Ichnusella
eione Cottarelli, 1971
pasquinii (Cottarelli, 1969)
- Psammopsyllus*
arenarius Enckell, 1965
delamarei (Chappuis, 1954)
mariae Cottarelli, Saporto & Puccetti, 1983
operculatus Nicholls, 1945
- Family Pseudotachidiidae**
Anapophysia
borealis (Klie, 1939)
- Archisenia*
sibirica (Sars, 1898)
- Bathypsammis*
longifurca (Bodin, 1968)
- Cylindronampus*
bispinosus Schriever, 1985
- elongatus* (Becker, 1979)
- Dactylopodella*
clypeata Sars, 1911
- flava* (Claus, 1866)

- rostrata* (T Scott, 1893)
vervoorti Moore, 1976
- Danielssenia**
quadriseta Gee, 1988
spitsbergensis Gee & Huys, 1994
typica Boeck, 1873
- Fladenia**
robusta (Sars, 1921)
- Idomene**
borealis Sars, 1911
coronata (T Scott, 1894)
forficata Philippi, 1843
kabylica Monard, 1936
parasimulans Médioni & Soyer, 1967
pectinata (T & A Scott, 1898)
purpureocincta (Norman & T Scott, 1905)
- Jonesiella**
fusiformis (Brady & Robertson in Brady (1880, 1880)
- Leptotachidina**
iberica Becker, 1974
- Micropsammis**
noodti Mielke, 1975
- Mucrosenia**
kendalli Gee & Huys, 1994
- Paradanielssenia**
biclavata Gee, 1988
christineae Gee & Huys, 1994
kathleenae Gee & Huys, 1994
kunzi Soyer, 1970
- Paranannopus**
abyssi (Sars, 1920)
bahusiene Por, 1964
caheti Soyer, 1964
denticulatus Schriever, 1985
hicksi Schriever, 1985
kunzi Schriever, 1985
philistinus Por, 1964
plumosus Schriever, 1983
reductus Becker, 1979
sarsi Lang, 1936
singulasetosus Schriever, 1985
triarticulatus Wells, 1965
trisetosus Schriever, 1985
uniarticulatus Schriever, 1985
variabilis Schriever, 1985
wellsi Soyer, 1976
- Peltisenia**
aberrans (Por, 1964)
- Psammis**
longisetosa Sars, 1910
- Pseudomesochra**
aberrans Bodin, 1968
abyssalis Becker & Schriever, 1979
beckeri Becker & Schriever, 1979
brucei (T & A Scott, 1901)
divaricata (Sars, 1906)
latifurca (Sars, 1911)
longifurcata T Scott, 1902
media (Sars, 1911)
perplexa Bodin, 1968
scheibeli Schriever, 1982
similis Lang, 1935
tatianae Drzycimski, 1968
- Pseudotachidius**
coronatus T Scott, 1898
ibericus Becker & Schriever, 1979
similis T Scott, 1903
vikingus Drzycimski, 1968
- Telopsammis**
secunda (Mielke, 1975)
- Family Rhizothricidae**
- Rhizothrix**
curvata (Brady & Robertson in Brady (1880), 1880)
gracilis (T Scott, 1903)
minuta (T Scott, 1903)
- pubescens* Por, 1959
reducta subsp. *noodti* Galhano, 1970
reducta subsp. *reducta* Noodt, 1952
scotti Lang, 1936
wilsoni Bodin, 1979
- Tryphoema**
bocqueti (Bozic, 1953)
lusitanica (Wells & Clark, 1965)
porca Monard, 1926
scilloniensis (Wells, 1968)
- Family Rotundiclipidae**
Rotundiclipus
canariensis Huys, 1988
- Family Superornatiremidiae**
- Intercrusia**
garciai Jaume, 1997
- Neoechinophora**
karaytugi Huys, 1996
xoni Jaume, 1997
- Superornatiremis**
mendai Jaume, 1997
- Family Tachidiidae**
Geeopsis
incisipes (Klie, 1913)
- Microarthridion**
berberum (Monard, 1935)
fallax Perkins, 1956
littorale (Poppe, 1881)
reductum (Monard, 1935)
- Tachidius**
discipes Giesbrecht, 1881
- Family Tegastidae**
- Parategastes**
haphe Leigh-Sharpe, 1936
sphaericus (Claus, 1863)
- Syngastes**
cornalinus Monard, 1924
macrognathus Monard, 1924
- Tegastes**
areolatus Monard, 1935
calcaratus Sars, 1910
clausi Sars, 1910
dalmatinus Petkovski, 1955
elenae Marcus, 1963
falcatus (Norman, 1868)
flavidus Sars, 1904
grandimanus Sars, 1904
knoepflii Médioni & Soyer, 1967
longimanus (Claus, 1863)
nanus Sars, 1904
neapolitanus (Claus, 1863)
porosus Petkovski, 1955
riedli Pesta, 1959
satyrus (Claus, 1860)
seurati Monard, 1936
- Family Tetragonicitidae**
- Aigondiceps**
bocki (Lang, 1948)
brevicauda (Huys, 1995)
kunzi (Marinov, 1974)
- Laophontella**
horrida subsp. *horrida* (Por, 1964)
- Paraschizopera**
menaiensis (Geddes, 1968)
- Phyllopodopsyllus**
berrieri Monard, 1936
bradyi (T Scott, 1892)
briani Petkovski, 1955
curtus Marcus, 1976
danielae Bodin, 1964
furciger Sars, 1907
hardingi (Roe, 1955)
hibernicus (Roe, 1955)
lasalmensis Marinov, 1973

- laticauda* Por, 1964
longicaudatus A Scott, 1909
longipalpus subsp. *longipalpus* (Chappuis, 1954)
medius Por, 1964
pauli Crisafi, 1960
thiebaudi subsp. *thiebaudi* Petkovski, 1955
Protagoniceps
hebraeus Por, 1964
Pteropsyllus
consimilis (T Scott, 1894)
Tetragoniceps
bergensis Por, 1965
bookhouti Coull, 1971
brevicauda T Scott, 1900
malleolatus Brady, 1880
scotti Sars, 1911
- Family Thalestridae**
Amenophia
peltata Boeck, 1865
pulchella Sars, 1906
Dactylopodopsis
dilatata Sars, 1911
Dactylopusia
brozkiae (Chislenko, 1967)
euryhalina Monard, 1935
glacialis subsp. *glacialis* Sars, 1909
longyearbyensis (Mielke, 1974)
micronyx Sars, 1905
neglecta Sars, 1905
pontica (Apostolov, 1968)
tisboides (Claus, 1863)
vulgaris subsp. *dissimilis* Brian, 1921
vulgaris subsp. *inornata* (Lang, 1965)
vulgaris subsp. *vulgaris* Sars, 1905
Dactylopusioides
macrolabris (Claus, 1866)
Diarthrodes
aegideus (Brian, 1927)
andrewi (T Scott, 1894)
assimilis (Sars, 1906)
drachi Bodiou, 1974
dubius (Brian, 1921)
fahrenbachi Bodin, 1968
feldmanni Bocquet, 1953
gravellicola Soyer, 1975
major (T & A Scott, 1895)
minutus (Claus, 1863)
nanus (T Scott, 1914)
nobilis (Baird, 1845)
ponticus subsp. *orientalis* Apostolov, 1975
ponticus subsp. *ponticus* (Kricagin, 1877)
pygmaeus (T & A Scott, 1895)
roscoffensis (Monard, 1935)
zavodniki Apostolov & Petkovski, 1980
Donsiella
anglica Hicks, 1988
limnoriae Stephensen, 1936
Eudactylopus
latipes (T Scott, 1893)
robustus (Claus, 1863)
spectabilis (Brian, 1923)
Paradactylopodia
bathybates (Monard, 1936)
brevicornis (Claus, 1866)
latipes (Boeck, 1865)
striata Kunz, 1983
Parathalestris
cambriensis Wells, 1964
clausi (Norman, 1868)
croni (Kröyer in Gaimard (1842-1845?), 1842-1845?)
dovi Marcus, 1966
harpactoides (Claus, 1863)
hibernica (Brady & Robertson, 1873)
intermedia Gurney, 1930
irelandica Roe, 1958
- jacksoni* (T Scott, 1899)
plumiseta Moore, 1976
Peltthestris
tripartita Monard, 1924
Phyllothalalestris
mysis (Claus, 1863)
Rhynchothalalestris
helgolandica (Claus, 1863)
Thalestris
brunnea Sars, 1905
curticornis Boeck, 1865
denti Brady, 1905
filifera Kricagin, 1877
frigida T Scott, 1899
gibba (Kröyer in Gaimard (1842-1845?), 1842-1845?)
longimana Claus, 1863
longipes Boeck, 1865
microphylla Claus, 1863
normani T Scott, 1903
northumbrica (Brady, 1905)
purpurea Sars, 1905
rhodymeniae (Brady, 1894)
rufoviolascens Claus, 1866
Tisemus
pulchellus Monard, 1928
- Family Thompsonulidae**
Thompsonula
hyaenae (Thompson, 1889)
- Family Tisidae**
Bathyidia
remota Farran, 1926
Cholidya
polypi Farran, 1914
Cholidyella
intermedia (Bresciani, 1970)
Idyanthe
dilatata (Sars, 1905)
pusilla (Sars, 1905)
Idyella
exigua Sars, 1905
kunzi Bodin, 1968
major Sars, 1920
pallidula Sars, 1905
Idyelopsis
typica Lang, 1944
Peresime
reducta (Becker & Schriever, 1979)
Sacodiscus
fasciatus (Norman, 1868)
humesi Stock, 1960
littoralis (Sars, 1904)
Scutellidium
arthuri Poppe, 1884
fucicolum (T Scott, 1912)
hippolytes subsp. *hippolytes* (Kröyer, 1863)
ligusticum (Brian, 1920)
longicauda subsp. *longicauda* (Philippi, 1840)
Tachidiella
minuta Sars, 1909
reducta Lee & Huys, 1999
Tachidiopsis
cyclopoides Sars, 1911
Tisbe
angusta (Sars, 1905)
aragoi Battaglia & Volkmann-Rocco, 1969
battagliai Volkmann-Rocco, 1972
bulbisetosa Volkmann-Rocco, 1972
clodiensis Battaglia & Fava, 1968
cluthae (T Scott, 1899)
compacta (Sars, 1920)
cucumariae Humes, 1957
dilatata Klie, 1949
dobzhanskii Volkmann-Rocco & Battaglia, 1972
elegantula (Sars, 1905)
elongata (A Scott, 1896)

ensifer Fischer, 1860
finmarchica (Sars, 1905)
furcata subsp. *furcata* (Baird, 1837)
gigantea Volkmann, 1979
gracilis (T Scott, 1895)
graciloides (Sars, 1920)
histriana Marcus & Por, 1961
holothuriae Humes, 1957
inflatiseta Volkmann, 1979
lagunaris Volkmann-Rocco, 1972
longicornis (T & A Scott, 1895)
longisetosa Gurney, 1927
marmorata Volkmann-Rocco, 1973
minor (T & A Scott, 1896)
palaeocrystica (Norman, 1878)
pentataenia Volkmann-Rocco, 1973
perplexa Volkmann, 1979
persimilis Volkmann-Rocco & Fava, 1969
pontina Volkmann-Rocco, 1969
pori Betouhim-El & Kahan, 1972
reductans Volkmann-Rocco, 1968
remanei Volkmann, 1975
reticulata Bocquet, 1951
similis Kricagin, 1873
tenella (Sars, 1911)
tenera (Sars, 1905)
variipes Marcus, 1974
wirkerissae Chislenko, 1967
Zosime
atlantica Bodin, 1968
bergensis Drzycimski, 1968
gisleni Lang, 1948
incrassata subsp. *bathybia* Bodin, 1968
incrassata subsp. *incrassata* Sars, 1910
major Sars, 1919
mediterranea Lang, 1948
paramajor Bodin, 1968
paratypica Becker & Schriever, 1979
rubra Thompson, 1888
typica Boeck, 1873
valida Sars, 1919

Tantulocarida

Compiled by Geoff Boxshall

The classification follows Huys (1991). These microscopic crustaceans, recognised in the 1980s, occur as ectoparasites on other host crustaceans, including the copepods, isopods, tanaids and amphipods. The data are drawn largely from Huys (1991), Boxshall & Vader (1993), Huys, Andersen and Kristensen (1992), and Huys, Boxshall and Casanova (1993).

References

- Boxshall, G.A. & Vader, W. 1993. A new genus of Tantulocarida (Crustacea) parasitic on an amphipod host from the North Sea. *J. nat. Hist.* 27:977-988.
 Huys, R., 1991. Tantulocarida (Crustacea: Maxillopoda): A new taxon from the temporary meiobenthos, Marine Ecology: *Pubblicazione della Stazione Zoologica di Napoli*, 12: 1-34.
 Huys, R., Andersen, P.F. and Kristensen, R.M., 1992. *Tantulacus hoegi* gen. et sp. nov. (Tantulocarida: Deuterterhidae) from the meiobenthos of the Faroe Bank, North Atlantic, *Sarsia*, 76: 287-297.
 Huys, R., Boxshall, G.A. and Casanova, J.-P., 1993. A redescription of *Cumoniscus* Bonnier, 1903 (Crustacea: Tantulocarida) from a cumacean host in the Bay of Biscay, *Systematic Parasitology*, 26: 75-79.

Subclass Tantulocarida

Family Deuterterhidae

- Tantulacus*
hoegi Huys, Andersen & Kristensen, 1992
Cumoniscus

kruppi Bonier, 1903
Boreotantulus
kunzi Huys & Boxshall, 1988
Amphitanthus
harpiniacheres Boxshall & Vader, 1993
Dicrotrichura
tricincta Huys, 1989
Family Basipodellidae
Basipodella
atlantica Boxshall & Lincoln, 1983
Stygotantulus
stocki Boxshall & Huys, 1989
Family Microdajidae
Microdajus
gaelicus Boxshall & Lincoln, 1987
langi Greve, 1965
pectinatus Boxshall, Huys & Lincoln, 1988
Family Doryphallophoridae
Doryphallophora
harrisoni (Boxshall & Lincoln, 1987)
Family Onceroxenidae
Onceroxenus
birdi Boxshall & Lincoln, 1987
curtus Boxshall & Lincoln, 1987

Cirripedia

Free-living

Compiled by A. J. Southward

Note here the parasitic groups Ascothoracida and Rhizocephala are not included, and indeed the latter are often considered to be separate from the true Cirripedia.

The taxonomic classification is based on Darwin (1851, 1854), Gruvel (1905), Pilsbry (1907, 1916), Tomlinson (1967) and Zevina (1981, 1982), as revised by Newman (1996). Other details of taxonomy and geographical limits are based on Darwin's volumes and on Bassindale (1964), Crisp *et al.* (1981), Marine Biological Association (1957), Nilsson-Cantell (1978), Newman and Ross (1976), Hayward and Ryland (1990), Relini (1980), Southward, (1976), Southward and Crisp (1963) and Young (1998a, 1998b, 1998c), with revisions from personal records of A.J. Southward and the late D.J. Crisp. Note that some geographical distributions in Nilsson-Cantell (1978) are incorrect, notably those for *Chthamalus* species and for *Semibalanus balanoides*. Species recently described, or not covered in detail or not included in the above compilations are described by Noll (1872), Southward (1995, 1999) and Turquier (1967).

References

- Bassindale, R., 1964. British Barnacles. *Synopses of the British Fauna*, no. 14.
 Crisp, D.J., Southward, A.J. & Southward, E.C. 1981. On the distribution of the intertidal barnacles *Chthamalus stellatus*, *Chthamalus montagui* and *Euraphia depressa*. *Journal of the Marine Biological Association of the United Kingdom* 61,359-380.
 Darwin, C., 1851. *A Monograph on the Sub-Class Cirripedia*, vol. 1. *The Lepadidae; or, pedunculated cirripedes*. London: Ray Society.
 Darwin, C., 1854. *A Monograph on the Sub-Class Cirripedia*, vol. 2. *The Balanidae, the Verrucidae, etc.* London: Ray Society.
 Gruvel 1905. Monographie des Cirripèdes ou Thecostracés. Paris, Masson et cie.
 Hayward, P.J. & Ryland, J.S. (eds), 1990. *The marine fauna of the British Isles and North-west Europe*, vol. 1. Oxford: Clarendon Press

- Marine Biological Association, 1957. *Plymouth Marine Fauna*, 2nd Edition. Plymouth: Marine Biological Association.
- Newman, W.A., 1997. Sous-Classe des Cirripèdes. Super-Ordres des Thoraciques et des Acrothoraciques. In *Traité de Zoologie*, Tome 7 (J. Forest, ed.), fasc. 2, pp. 453-450. Masson et cie, Paris
- Newman, W.A. & Ross, A. 1971. Antarctic Cirripedia. *Antarctic Research Series*, 14, American Geophysical Union, Washington, 257 pp.
- Newman, W.A. & Ross, A., 1976. Revision of the balanomorph barnacles: including a catalog of the species. *Memoirs of the San Diego Society of Natural History*, 9, 108pp.
- Nilsson-Cantell, C.A., 1978. Cirripedia Thoracica and Acrothoracica. *Marine Invertebrates of Scandinavia*, no. 5, 133 pp.
- Noll, F.C., 1872. Kochlorine hamata n., ein bohrender Cirripede. *Berichte der Senckenberg naturforschungs Gesellschaft*, 4, 50-58.
- Pilsbry, H.A., 1907. The barnacles (Cirripedia) contained in the collections of the U.S. National Museum. *Bulletin of the United States National Museum* 60, 1-122.
- Pilsbry, H.A., 1916. The sessile barnacles (Cirripedia) contained in the collections of the U.S. National Museum; including a monograph of the American species. *Bulletin of the United States National Museum*, 93, 1-366.
- Relini, G. 1980. 'Cirripedi Toracici'. Guida per il riconoscimento delle specie animali acque lagunari e costiere italiane. Consiglio Nazionale delle Ricerche, Genova, 112pp.
- Southward, A.J. 1976. On the taxonomic status and distribution of *Chthamalus stellatus* (Cirripedia) in the north-east Atlantic region: with a key to the common intertidal barnacles of Britain. *Journal of the Marine Biological Association of the United Kingdom*, 56, 1007-1028.
- Southward, A.J., 1995. Occurrence in the English Channel of a warm-water cirripede, *Solidobalanus fallax*. *Journal of the Marine Biological Association of the United Kingdom*, 75, 199-210.
- Southward, A.J. 1999. New observations on barnacles (Crustacea: Cirripedia) of the Azores region. *Arquipelago*, 16A: 11-27.
- Southward, A.J. & Crisp, D.J., 1963. Barnacles. *Catalogue of main marine fouling organisms*, 1, 46pp. Paris: O.E.C.D. Publications.
- Tomlinson, J.T. 1969. The burrowing barnacles (Cirripedia: Order Acrothoracica). *Bulletin of the United States National Museum*, 296, 1-162.
- Turquier, Y., 1967. Description d'un nouveau *Trypetesa* Norman (= *Alcippe* Hancock), Cirripede Acrothoracique des côtes françaises de la Manche. *Cahiers de Biologie Marine*, 8, 75-87.
- Young, P.S., 1998a. Cirripedia (Crustacea) from the "Campagne Biaçores" in the Azores region, including a generic revision of Verrucidae. *Zoosystema* 20(1):31-92
- Young, P.S., 1998b. Cirripedes (Crustacea) from the Mid-Atlantic Ridge collected by the submersible *Nautile*. *Cahiers de Biologie Marine*, 39, 109-119.
- Young, P.S., 1998c. The Cirripedia (Crustacea) collected by the "Fisherries Steamer Meteor" in the Eastern Atlantic. *Arquivos do Museu Nacional, Rio de Janeiro* 58, 1-54.
- Zevina, G.B. 1981. Barnacles of the suborder Lepadomorpha of the world ocean. I. Family Scalpellidae. *Fauna USSR* vol. 127, 406pp, Nauka, Leningrad. (in Russian)
- Zevina, G.B. 1982. Barnacles of the suborder Lepadomorpha of the world ocean. II. *Fauna USSR* vol. 133, 222pp, Nauka, Leningrad. (in Russian)
- Subclass Cirripedia**
- Superorder Acrothoracica**
- Order Pygophora**
- Suborder Pygophora**
- Family Lithoglyptidae**
Weltneria
exargilla Newman, 1974
- Kochlorine**
hamata Noll, 1872
- Order Apygophora**
- Suborder Apygophora**
- Family Trypetidae**
Trypetesa
lampas Hancock, 1849
= *Alcippe lampas*
nasserooides Turquier, 1967
- Superorder Thoracica**
- Order Pedunculata**
- Suborder Pedunculata**
- Family Heteralepadidae**
Heteralepas
microstoma Gravell, 1902
meteorensis Carriol, 1998
cornuta Darwin, 1851
Alepas
parasita Rang, 1829
Paralepas
minuta Philippi, 1836
- Family Anelasmatidae**
Anelasma
squalicola Darwin, 1851
- Suborder Lepadomorpha**
- Family Oxynaspidae**
Oxynaspis
celata Darwin, 1851
patens Aurivillius, 1894
- Family Poecilasmatidae**
Poecilasma
aurantia Darwin, 1851
crassa Gray, 1848
Glyptelasma
hamatum Calman, 1919
carinatum Hoek, 1883
rectum Pilsbry, 1907
Octolasmus
lowei Darwin, 1851
Dichelaspis
sessilis Hoek, 1883
thieli Young, 1998
darwini (Philippi, 1861)
= *Octolasmus darwini*
- Family Lepadidae**
Lepas
anatifera Linnaeus, 1758
anserifera Linnaeus, 1758
hilli Leach, 1818
pectinata Spengler, 1793
Dosima
fascicularis Ellis & Solander, 1786
Conchoderma
virgatum Spengler, 1789
auritum (Linnaeus, 1758)
- Suborder Scalpellomorpha**
- Family Calanticidae**
Scillaelepas
gemma (Aurivillius, 1894)
Gruvelialepas
pilsbryi (Gruvel, 1911)
Aurivillialepas
calyculus (Aurivillius, 1898)
falcata (Aurivillius, 1898)
Smilium
acutum (Hoek, 1883)
- Family Pollicipedidae**
Pollicipes
pollicipes (Gmelin, 1789)
= *Pollicipes cornucopia*

- Family Scalpellidae**
- Subfamily Scalpellinae**
- Scalpellum*
scalpellum (Linnaeus, 1758)
 - Ornatocalpellum*
stroemii M. Sars, 1859
 - Hamatoscalpellum*
hamatum G.O. Sars, 1879
 - Litoscalpellum*
meteoria Young, 1998
 - Neoscalpellum*
debile (Aurivillius, 1898)
- Subfamily Arcoscalpellinae**
- Arcoscalpellum*
michelottianum (Seguenza, 1876)
 - eponkos* Young, 1998
 - tritonis* (Hoek, 1883)
 - Planoscalpellum*
limpidus Zevina, 1976
 - Tarasovium*
cornutum (G.O. Sars, 1879)
 - Weltnerium*
nymphocola (Hoek, 1883)
 - pusillum* (Aurivillius, 1898)
 - Verum*
parazelandiae Young, 1998
 - striolatum* (G.O. Sars, 1877)
 - Catherinum*
recurvitergum (Gruvel, 1900)
 - Amigdoscalpellum*
hispidum (G.O. Sars, 1890)
 - mammilatum* (Aurivillius, 1898)
 - rigidum* (Aurivillius, 1898)
 - Trianguloscalpellum*
regium (Wyville-Thomson, 1877)
 - ovale* (Hoek, 1883)
 - darwini* (Hoek, 1883)
 - gigas* (Hoek, 1883)
 - Teloscalpellum*
anceps (Aurivillius, 1896)
 - atlanticum* (Gruvel, 1900)
 - gracile* (Gruvel, 1920)
 - incisum* (Aurivillius, 1898)
 - luteum* (Gruvel, 1900)
- Order Sessilia**
- Suborder Verrucomorpha**
- Family Verrucidae**
- Altiverruca*
costata (Aurivillius, 1898)
 - crenata* (Aurivillius, 1898)
 - erecta* (Gruvel, 1900)
 - gibbosa* (Hoek, 1883)
 - inermis* (Aurivillius, 1898)
 - carinata* (Gruvel, 1900)
 - obliqua* (Hoek, 1883)
 - Newmanverruca*
ferruginea Young, 1998
 - Metaverruca*
aequalis (Aurivillius, 1898)
 - recta* (Aurivillius, 1898)
 - cornuta* (Aurivillius, 1898)
 - sensibilis* Young, 1998
 - Verruca*
stroemia O.F. Müller, 1776
 - spengleri* Darwin, 1854
- Suborder Balanomorpha**
- Superfamily Pachylasmatoidea**
- Family Pachylasmatidae**
- Pachylasma*
giganteum Philippi, 1836
- Superfamily Chthamaloidea**
- Family Chthamalidae**
- Subfamily Euraphiinae**
- Euraphia*
depressa (Poli, 1795)
= *Chthamalus depressus*
- Subfamily Chthamalinae**
- Chthamalus*
stellatus (Poli, 1795)
montagui Southward, 1976
- Superfamily Coronuloidea**
- Family Chelonibiidae**
- Chelonibia*
carettæ Spengler, 1790
 - testudinaria* (Linnaeus, 1758)
 - patula* (Ranzani, 1820)
- Family Platylepadidae**
- Platylepas*
hexastylus (Fabricius, 1798)
 - Stomatolepas*
elegans (Costas, 1883)
- Family Coronulidae**
- Coronula*
diadema (Linnaeus, 1758)
 - Xenobalanus*
globicipitis Steenstrup, 1851
- Superfamily Tetractitoidea**
- Family Bathylasmatidae**
- Subfamily Bathylasmatinae**
- Bathylasma*
hirsutum (Hock, 1883)
- Subfamily Hexelasmatinae**
- Hexelasma*
americanum Pilsbry, 1916
- Family Tetractitidae**
- Subfamily Tetractitinæ**
- Tesseropora*
atlantica Newman & Ross, 1976
- Superfamily Balanoidea**
- Family Archaeobalanidae**
- Subfamily Archaeobalaninae**
- Chirona*
hameri (Ascanius, 1767)
= *Balanus hameri*
 - Solidobalanus*
fallax (Broch, 1927)
- Subfamily Acastinae**
- Acasta*
spongites (Poli, 1795)
 - Conopea*
calceola (Ellis, 1758)
= *Balanus calceolus*
- Subfamily Elminiinae**
- Elminius*
modestus Darwin, 1854
- Subfamily Semibalaninae**
- Semibalanus*
balanoides (Linnaeus, 1758)
= *Balanus balanoides*
- Family Pyrgomatidae**
- Subfamily Megatrematinae**
- Megatrema*
anglicum (Sowerby, 1823)
= *Pyrgoma anglicum*
- Family Balanidae**
- Subfamily Balaninae**
- Balanus*
balanus (Linnaeus, 1758)
= *Balanus porcatus*
 - crenatus* Bruguière, 1789
 - amphitrite* Darwin, 1854

= *Balanus amphitrite* subsp. *amphitrite*
eburneus Gould, 1841
improvisus Darwin, 1854
spongicola Brown, 1844
trigonus Darwin, 1854
perforatus Bruguère, 1789

Subfamily Megabalaninae

Megabalanus
tintinnabulum (Linnaeus, 1758)
= *Balanus tintinnabulum*
azoricus (Pilsbry, 1916)
= *Balanus tintinnabulum*
tulipiformis (Ellis, 1758)

Ascothoracida

Compiled by Geoff Boxshall

The classification follows Grygier (1987). These highly transformed thecostrakan crustaceans occur as parasites of echinoderms and corals. The data for European waters are drawn largely from the unpublished PhD thesis of Grygier (Grygier, 1983a), from Grygier (1983b), from Moyse (1983) and Stone & Moyse (1985).

References

- Grygier, M.J. (1983a) *Comparative morphology and ontogeny of the Ascothoracida, a step toward a phylogeny of the Maxillopoda*. University of California, San Diego, unpublished PhD thesis. 417pp.
- Grygier, M.J. (1983b) Revision of *Synagoga* (Crustacea: Maxillopoda: Ascothoracida). *J. nat. Hist.* 17: 213-239.
- Grygier, M.J. (1987) Classification of the Ascothoracida (Crustacea). *Proc. Biol. Soc. Wash.* 100: 452-458.
- Moyse, J. (1983) *Isidascus bassindalei* gen.nov., sp.nov. (Ascothoracida : Crustacea) from North-East Atlantic with a note on the origin of barnacles. *Journal of the Marine Biological Association of the United Kingdom*. 63: 161-180.
- Stone, C.J. & Moyse, J. (1985) *Bifurgaster*, a new genus of Ascothoracida (Crustacea: Maxillopoda) parasitic in deep water asteroids. *J. nat. Hist.* 19: 1269-1279.

Subclass Ascothoracida

Order Dendrogastrida

Family Ascothoracidae

Ascothorax
ophioctenis Djakonov, 1914
pilocaudatus Grygier, 1983

Family Dendrogastridae

Dendrogaster
astericola Knipovich, 1890
Bifurgaster
freyellae Stone & Moyse, 1985
Dendrogaster
muranensis Wagin, 1950
Ulophysema
oresundense Brattström, 1936

Order Laurida

Family Lauridae

Laura
gerardiae Lacaze-Duthiers, 1865

Family Petrarcidae

Petrarca
azorica Grygier, 1985

Family Synagogidae

Isidascus
bassindalei Moyse, 1983
Synagoga
mira Norman, 1888

Rhizocephala

Compiled by Geoff Boxshall

The classification follows Høeg & Lützen (1985) and was updated to incorporate the changes proposed by Høeg & Rybakov (1992). These highly transformed crustaceans occur as parasites on host crustaceans. The data for Northern European waters are drawn largely from the excellent account by Høeg & Lützen (1985) of the Rhizocephala of Scandinavia and for Mediterranean waters, from the classic monograph by Smith (1906) on the Fauna of the Gulf of Naples, and from Høeg & Rybakov (1992).

References

- Høeg, J.T. & Lützen, J. 1985. Crustacea Rhizocephala. *Marine Invertebrates of Scandinavia* No. 6: 1-92.
- Høeg, J.T. & Rybakov, A.V. 1992. Revision of the Rhizocephala Akentrogonida (Cirripedia), with a list of all the species and a key to the identification of families. *J. Crust. Biol.* 12: 600-609.
- Smith, G.W. 1906. Rhizocephala. *Fauna Flora Golf. Neapel* 29: 1-123.

Superorder Rhizocephala

Order Akentrogonida

Family Chthamalophilidae
Boschmaella
balani (Bocquet-Védrine, 1967)
Chthamalophilus
delagei Bocquet-Védrine, 1957

Family Clistosaccidae

Clistosaccus
paguri Lilljeborg, 1860
Sylon
hippolytes M. Sars, 1870

Family Duplorbidae

Cryptogaster
cumacei Bocquet-Védrine & Bourdon, 1984
Duplorbis
calathurae Smith, 1906

incertae sedis

Polysaccus
mediterraneus (Caroli, 1929)

Order Kentrogonida

Family Lernaeodiscidae

Lernaeodiscus
ingolfi Boschma, 1928
squamiferae Pérez, 1922
Triangulus
galatheae (Norman & Scott, 1906)
munidae Smith, 1906

Family Peitogastriidae

Cyphosaccus
norvegicus Boschma, 1962
Galatheascus
striatus Boschma, 1929

Parthenopea
subterranea Kossmann, 1874

Peltogaster
curvatus Kossmann, 1874
paguri Rathke, 1842

Peltogastrella
sulcata (Lilljeborg, 1859)
Tortugaster

boschmai (Brinkman, 1939)
Trachelosaccus
hymenodora (G.O. Sars, 1879)

Family Sacculinidae

Drepanorchis
neglecta (Fraisse, 1877)

Sacculina

atlantica Boschma, 1927
bourdoni Boschma, 1960
carcini Thompson, 1836
gerberi Bonnier, 1887
gibbsi (Hesse, 1867)
inflata Leuckart, 1859
lobata Boschma, 1965
triangularis Anderson, 1862

Stomatopoda

Compiled by Jacob van der Land

The ERMS list was derived from URMO which is based on Müller (1993) and corrected by P. Noël.

References

Müller, H.-G., 1993. *World Catalogue and Bibliography of the Recent Stomatopoda*. Müller Wissenschaft. Verlag, Wetzlar.
 [also on floppy disk]

Order Stomatopoda

Family Bathysquillidae

Bathysquilla
microps (Manning, 1961)

A

Family Coronidae

Coronida
bradyi (A. Milne Edwards, 1869)

A

Family Euryssquillidae

Euryssquilla
galathea Manning, 1977

A

Manningia
posteli Manning, 1977

A

Family Lysiosquillidae

Lysiosquilla
hoevenii (Herklotz, 1851)
monodi Manning, 1977

A

Lysiosquilloides
aulacorhynchus (Cadenat, 1957)

A

Family Nannosquillidae

Allosquilla
africana (Manning, 1970)

A M

Bigelowina
septemspinosa (Miers, 1881)

A

Nannosquilloides
occultus (Giesbrecht, 1910)

A M

Platysquilla
eusebia (Risso, 1816)

A M

Platysquilloides
lillyae (Manning, 1977)

A M

Family Parasquillidae

Parasquilla
ferussaci (Roux, 1828)

A M

Pseudosquillopsis
cerisii (Roux, 1828)

A M

Family Protosquillidae

Protosquilla
folini (A. Milne-Edwards, 1867)

A

Family Squillidae

Alima
hyalina Leach, 1817

A

Erugosquilla
massavensis (Kossmann, 1880)

M

Rissoïdés

africanus (Manning, 1974)
calypso (Manning, 1974)
desmaresti (Risso, 1816)

A

pallidus (Giesbrecht, 1910)

A M

Squilla

A M

mantis (Linnaeus, 1758)

Euphausiacea

Compiled by Jacob van der Land

The European list was based on Brinton *et al.* The list has not been checked by an expert on the European species.

References

Brinton E., M.D. Ohman, A.W. Townsend, M.D. Knith & A.L. Bridgeman. Euphausiids of the World Ocean. ETI CD-Rom.

Order Euphausiacea

Family Bentheuphausiidae

Bentheuphausia
amblyops G.O. Sars, 1885

A M

Family Euphausiidae

Euphausia
americana Hansen, 1911
brevis Hansen, 1905
gibboidea Ortmann, 1893
hanseni Zimmer, 1915
hemigibba Hansen, 1910
krohni (Brandt, 1851)
mutica Hansen, 1905
pseudogibba Ortmann, 1893

A M

A M

A

A M

A M

A

Meganyctiphanes

norvegica (M. Sars, 1857)

A M

Nematobrachion

boopis (Calman, 1905)
flexipes (Ortmann, 1893)
sexspinosum Hansen, 1911

A M

A M

A M

Nematozelis

atlantica Hansen, 1910
megalops G.O. Sars, 1883
micros G.O. Sars, 1883

A M

A M

tenella G.O. Sars, 1883
Nyctiphanes

A M

couchi (Bell, 1853)

A M

Stylocheiron

abbreviatum G.O. Sars, 1883
affine Hansen, 1910

A M

A M

carinatum G.O. Sars, 1883

A M

elongatum G.O. Sars, 1883
longicorne G.O. Sars, 1883

A M

A M

maximum Hansen, 1908
robustum Brinton, 1962

A M

A M

suhmi G.O. Sars, 1883
Thysanoessa

A M

A M

gregaria G.O. Sars, 1883

A M

inermis (Krøyer, 1846)
longicaudata (Krøyer, 1846)

A M

A M

parva Hansen, 1905

A M

raschii (M. Sars, 1864)

A M

Thysanopoda

acutifrons Holt & Tattersall, 1905
aequalis Hansen, 1905
cornuta Illig, 1905
cristata G.O. Sars, 1883
egregia Hansen, 1905
microphthalmia G.O. Sars, 1885
monacantha Ortmann, 1893
obtusifrons G.O. Sars, 1883
pectinata Ortmann, 1893
tricuspidata Milne-Edwards, 1837

A M

A M

A M

A M

A M

A M

A M

Decapoda

Compiled by Michael Türkay

The present list of European Decapoda has been adapted from a database which was compiled by myself before the project started. In 1999 a comprehensive list of European decapods was published by d'Udekem d'Acoz

and it was decided to make the present list as much compatible with that source as possible, because subsequent researchers will definitively base themselves on the published list containing also distribution and literature records. The present list thus also follows the classification used by d'Udekem d'Acoz (1999) on the family and generic level, however with some exceptions. The generic system of Pérez-Farfante & Kensley (1997) has been used for the Penaeoid shrimps rather than the one in d'Udeken d'Acoz (1999). The hippolytid genera *Eualus* and *Thoralus* are still separated, because if the mandible palpus character is removed as a primary one, the whole shrimp classification gets disputable. I do not think that such a discussion is possible at the present level of knowledge. In stenopodoid shrimps two families (viz. Spongicolidae and Stenopodidae) have been recognised instead of one, which is in accordance with Holthuis (1993). For the Callianassidae the classification of Sakai (1999b) has been used. The system proposed by d'Udekem d'Acoz (1999) for the polybiine genera of Portunidae has not been used, as I cannot see any progress in lumping all the genera together. Before such major changes are implemented a thorough revision must take place and the arguments be fully exposed and discussed. For the time being a certain "conservatism" seems to be more prudent.

On the species level all species known only from the Suez-Canal have been excluded in order to stick to a merely European framework. Also all species with doubtful identity have been excluded, as well as such that have been recorded only once a long time ago close to harbours and never have been found again. The following table gives a list of such species, also ones known from other sources, with the reason for their exclusion.

Metapenaeopsis vaillanti (Nobili, 1904) - Only Suez Canal; *Penaeus chinensis* (Osbeck, 1765) - Potential aquaculture species; *Penaeus monodon* Fabricius, 1798 - Potential aquaculture species; *Pasiphaea princeps* Smith, 1884 - Questionable records at abt. 25°N; *Plesionika holthuiisi* Crosnier & Forest, 1968 - Probably a misidentification; *Hippolyte orientalis* Heller, 1862 - Only Suez-Canal; *Gasconella parvula* De Saint Laurent, 1985 - *Nomen nudum*, invalid name; *Palaeomon pacificus* (Stimpson, 1860) - Only Suez Canal; *Jasus lalandii* (H. Milne-Edwards, 1837) - Single record, not established; *Homarus americanus* H. Milne-Edwards, 1837 - Probably escaped commercial specimens, not established; *Thenus orientalis* (Lund, 1793) - Single record from Rijeka port; *Paguristes perspicax* Nobili, 1906 - Only Suez Canal; *Galathea pygmaea* H. Milne-Edwards & Bouvier, 1894 - *incertae sedis*; *Munidopsis convexa* Noël, 1992 - *Nomen nudum*, invalid name; *Petrolisthes bosci* (Audouin, 1826) - Old record (1832) probable misidentification of *Pisidia*; *Dorippe frascone* (Herbst, 1795) - Only Suez-Canal; *Ebalia granulata* (Rüppell, 1830) - Only Suez-Canal; *Phlyra globosa* (Bosc, 1801) - Old record (1832) probabale misidentification of *Ilia*; *Schizophrys aspera* (H. Milne-Edwards, 1834) - Only Suez-Canal; *Liocarcinus subcorrugatus* (A. Milne-Edwards, 1861) - Doubtfully from Suez-Canal; *Portunus sanguinolentus* (Herbst, 1793) - Single record from Rijeka port; *Chaceon maritae* (Manning & Holthuis, 1981) - Only doubtful records; *Actaeodes hirsutissimus* (Rüppell, 1830) - Only Suez-Canal; *Actaea savignii* (H. Milne-Edwards, 1834) - Only

Suez-Canal; *Macromedaeus voeltzkowi* (Lenz, 1905) - Single record from English Channel (1913), not established; *Parapilumnus malardi* (De Man, 1913) - Single record from English Channel (1913), not established; *Parapilumnus truncatospinosus* (De Man, 1913) - Single record from English Channel (1913), not established; *Pilumnus longicornis* Hilgendorf, 1879 - Single record from English Channel (1913), not established; *Pilumnus savignyi* Heller, 1861 - Only Suez Canal; *Plagusia chabrus* (Linnaeus, 1758) - Single record from Marseille Port (1872); *Metopograpsus messor* (Forskal, 1775) - Only Suez Canal; *Macrobrachium depressus* (Rüppell, 1830) - Only Suez Canal; *Ocypode saratan* (Forskal, 1775) - Only Suez Canal; *Uca coarctata* (H. Milne-Edwards, 1852) - Single records from Adriatic and Black Seas; *Dotilla sulcata* (Forskål, 1775) - Only Suez Canal; *Serenella leachii* (Audouin, 1826) - Only Suez Canal; *Platymaia wyvillethomsoni* Miers, 1886 - Single record Adriatic Sea (1918); *Parthenope notialis* Manning & Holthuis, 1981 - Specimen from Cape Bojador (ca. 26°N) of unclear identity; because of new publications some species names had to be changed and new species had to be added: Galil (1999) recorded *Melicertus hathor* (Burkenroad, 1959) as a new Lessepsian migrant; and Sakai (1999b) described *Callianassa whitei* as a new species from the Adriatic Sea. Sakai (1999a) examined the classical collection of J. F. W. Herbst kept at the Berlin Museum and in consequence some names of well known European crabs had to be changed: *Pisa corallina* is now *P. hirticornis*, *Liocarcinus arcuatus* is now *L. navigator*, and *Xantho incisus* is now *X. hydrophilus*.

References

- Most of the references cited in this introduction can be found in d'Udekem d'Acoz (1999), the newer ones are cited here separately.
- Galil, B. S. (1999): *Melicertus hathor* Burkenroad, 1959 - a Red Sea penaeid prawn new to the Mediterranean. - *Crustaceana*, 72(9): 1226-1228.
- Sakai, K. (1999a): J. F. W. Herbst-Collection of decapod Crustacea of the Berlin Zoological Museum with remarks on certain species. - *Naturalists, Tokushima biol. Lab.*, 6: 1-45.
- Sakai, K. (1999b): Synopsis of the family Callianassidae, with keys to subfamilies, genera and species, and the description of new taxa (Crustacea: Decapoda: Thalassinidea). - *Zool. Verh.*, 326: 1-152.
- d'Udekem d'Acoz, C. (1999): Inventaire et distribution des crustacés décapodes de l'Atlantique nord-oriental, de la Méditerranée et des eaux continentales adjacentes au nord de 25°N. - *Patrimoines naturels* (M.N.H.N./S.P.N.), 40: I-X, 1-383; Paris.

Order Decapoda

Family Aristeidae

- Aristaeopsis edwardsiana* (Johnson, 1867)
- Aristeomorpha foliacea* (Risso, 1827)
- Aristeus antennatus* (Risso, 1816)
- Hepomadus tener* Smith, 1884
- Plesiopenaeus armatus* (Bate, 1881)

Family Benthesicymidae

- Benthogenmema intermedia* (Bate, 1888)
- Benthesicymus bartletti* Smith, 1882

- brasiliensis* Bate, 1881
iridescens Bate, 1881
laciniatus Rathbun, 1906
longipes Bouvier, 1906
Benthonectes
filipes Smith, 1885
Gennadas
brevirostris Bouvier, 1905
capensis Calman, 1925
elegans (Smith, 1882)
scutatus Bouvier, 1906
tinayrei Bouvier, 1906
valens (Smith, 1884)
- Family Solenoceridae**
Hymenopenaeus
aphoticus Burkenroad, 1936
chacei Crosnier & Forest, 1969
debilis Smith, 1882
Solenocera
africana Stebbing, 1917
crassicornis (H. Milne Edwards, 1837)
membranacea (Risso, 1816)
- Family Penaeidae**
Funchalia
danae Burkenroad, 1940
villosa (Bouvier, 1908)
woodwardi Johnson, 1867
Marsupenaeus
japonicus (Bate, 1888)
Melicertus
hathor (Burkenroad, 1959)
kerathurus (Forskål, 1775)
Metapenaeopsis
aegyptia Galil & Golani, 1990
mogiensis Rathbun, 1902 subsp. *consobrina* (Nobili, 1904)
Metapenaeus
monoceros (Fabricius, 1798)
stebbingi Nobili, 1904
Parapenaeus
longirostris (Lucas, 1846)
Pelagopenaeus
balboae (Faxon, 1893)
Penaeopsis
serrata Bate, 1881
Penaeus
semisulcatus De Haan, 1844
Trachysalambria
curvirostris (Stimpson, 1860)
- Family Sicyoniidae**
Sicyonia
carinata (Brünnich, 1768)
- Family Sergestidae**
Petalidium
obesum (Krøyer, 1855)
Sergestes
arachnipodus (Cocco, 1832)
arcticus Krøyer, 1855
armatus Krøyer, 1855
atlanticus H. Milne-Edwards, 1830
cornutus Krøyer, 1855
edwardsii Krøyer, 1855
nudus Illig, 1914
pectinatus Sund, 1920
sargassi Ortmann, 1893
vigilax Stimpson, 1860
- Sergia*
grandis (Sund, 1920)
japonica (Bate, 1881)
robusta (Smith, 1882)
splendens (Sund, 1920)
tenuiremis (Krøyer, 1855)
- Family Luciferidae**
Lucifer
- hanseni* Nobili, 1905
typus H. Milne-Edwards, 1837
- Family Stenopodidae**
Odontozona
addaia Pretus, 1990
edwardsi (Bouvier, 1908)
minoica Dounas & Koukouras, 1989
Richardina
fredericci Lo Bianco, 1903
spinicina A. Milne-Edwards, 1881
Stenopus
spinosus Risso, 1827
- Family Spongicolidae**
Spongicoloides
evolutus (Bouvier, 1905)
koehleri (Caullery, 1896)
profundus Hansen, 1908
- Family Pasiphaeidae**
Eupasiphae
gilesii (Wood-Mason, 1892)
serrata (Rathbun, 1902)
Glyphus
marsupialis Filhol, 1884
Leptocheila
aculeoaudata Paulson, 1875
pugnax De Man, 1916
Parapasiphae
compta Smith, 1884
cristata Smith, 1884
sulcatifrons Smith, 1884
Pasiphaea
hoplocerca Chace, 1940
liocerca Chace, 1940
multidentata Esmark, 1866
sivado (Risso, 1816)
tarda (Krøyer, 1845)
Psathyrocaris
fragilis Wood Mason, 1893
infirma Alcock & Anderson, 1894
- Family Alvinocarididae**
Chorocaris
chacei (Williams & Rona, 1986)
Mirocaris
fortunata (Martin & Christiansen, 1995)
Rimicaris
exoculata Williams & Rona, 1986
- Family Bresiliidae**
Bresilia
atlantica Calman, 1896
corsicana Forest & Cals, 1977
Kirnasia
nesini Burukovsky, 1988
siedlecki Burukovsky, 1988
- Family Ophelophoridae**
Acanthephyra
brevirostris Smith, 1885
curtirostris Wood Mason, 1891
eximia Smith, 1884
pelagica (Risso, 1816)
purpurea A. Milne-Edwards, 1881
stylorostratis (Bate, 1888)
tenuipes (Bate, 1888)
Ephyrina
benedicti Smith, 1885
bifida Stephensen, 1923
sigueirai Crosnier & Forest, 1973
Heterogenys
microphthalmia (Smith, 1885)
Hymenodora
acanthitelsonis Wasmer, 1972
glacialis (Buchholz, 1874)
gracilis Smith, 1886
Janicella

- spinicauda* (A. Milne-Edwards, 1883)
- Kemphyra**
- corallina* (A. Milne-Edwards, 1883)
- Meningodora**
- compsa* (Chace, 1940)
 - miccyla* (Chace, 1940)
 - mollis* Smith, 1882
 - vesca* (Smith, 1886)
- Notostomus**
- croshnieri* Macpherson, 1984
 - distirus* Chace, 1940
 - elegans* A. Milne-Edwards, 1881
 - robustus* Smith, 1884
- Oplophorus**
- spinosus* (Brullé, 1839)
- Systaspis**
- braueri* subsp. *braueri* (Balss, 1914)
 - cristata* Faxon, 1893
 - debilis* (A. Milne-Edwards, 1881)
 - pellucida* (Filhol, 1885)
- Family Nematocarcinidae**
- Nematocarcinus**
- ensifer* (Smith, 1882)
 - exilis* (Bate, 1888)
 - gracilipes* Filhol, 1884
- Nigmatellinus**
- acanthitelsonis* (Pequegnat, 1970)
- Family Pandalidae**
- Bitias**
- stocki* Fransen, 1990
- Chlorotocus**
- crassicornis* (Costa, 1871)
- Dichelopandalus**
- bonnieri* Caullery, 1896
- Heterocarpus**
- ensifer* subsp. *ensifer* A. Milne-Edwards, 1881
 - grimaldii* A. Milne-Edwards & Bouvier, 1900
 - laevigatus* Bate, 1888
- Pandalina**
- brevirostris* (Rathke, 1843)
 - profunda* Holthuis, 1946
- Pandalus**
- borealis* subsp. *borealis* Krøyer, 1838
 - latirostris* Rathbun, 1902
 - montagui* Leach, 1814
 - propinquus* Sars, 1870
- Physetocaris**
- microphthalmia* Chace, 1940
- Plesionika**
- acanthonotus* (Smith, 1882)
 - antigai* Zariquey-Alvarez, 1955
 - edwardsii* (Brandt, 1851)
 - ensis* (A. Milne-Edwards, 1881)
 - geniculatus* (A. Milne-Edwards, 1883)
 - giglioli* (Senna, 1902)
 - heterocarpus* (Costa, 1871)
 - martia* subsp. *martia* (A. Milne-Edwards, 1883)
 - narval* (Fabricius, 1787)
 - rossignoli* Crosnier & Forest, 1968
 - williamsi* Forest, 1974
- Stylopandalus**
- richardi* (Coutière, 1905)
- Family Ogyrididae**
- Ogyrides**
- njobergi* (Balss, 1921)
 - rariospina* Holthuis, 1951
- Family Hippolytidae**
- Bythocaris**
- akidopleura* Fransen, 1994
 - curvirostris* Kobjakova, 1957
 - elegans* Bryazgin, 1982
 - grumanti* Burukovsky, 1966
 - leucopsis* Sars, 1880
 - payeri* (Heller, 1875)
- simplicirostris* Sars, 1870
- Caridion**
- gordoni* (Bate, 1858)
 - steveni* Lebour, 1930
- Cryptocheles**
- pygmaea* Sars, 1870
- Eualus**
- belcheri* (Bell, 1855)
 - drachi* Noël, 1978
 - gaimardii* (H. Milne-Edwards, 1837)
 - lebourae* Holthuis, 1951
 - occultus* (Lebour, 1936)
 - pusiolus* (Krøyer, 1841)
- Hippolyte**
- coeruleescens* (Fabricius, 1775)
 - garciaraso* D'Udekem d'Acoz, 1996
 - holthuiisi* Zariquey-Alvarez, 1953
 - inermis* Leach, 1815
 - lagarderei* D'Udekem d'Acoz, 1995
 - leptocerus* (Heller, 1863)
 - leptometrae* Ledoyer, 1969
 - niezabitowskii* D'Udekem d'Acoz, 1996
 - prideauxiana* Leach, 1817
 - sapphica* D'Udekem d'Acoz, 1993
 - varians* Leach, 1814
- Latreutes**
- fucorum* (Fabricius, 1798)
- Lebbeus**
- groenlandicus* (Fabricius, 1775)
 - polaris* (Sabine, 1824)
- Leontocaris**
- lar* Kemp, 1906
- Ligur**
- ensiferus* (Risso, 1816)
- Lysmata**
- grabhami* (Gordon, 1935)
 - intermedia* (Kingsley, 1878)
 - nilita* Dohrn & Holthuis, 1950
 - olavo* Fransen, 1991
 - seticaudata* (Risso, 1816)
 - unicornis* Holthuis & Maurin, 1952
- Merhippolyte**
- ancistrotota* Crosnier & Forest, 1973
- Spirontocaris**
- lilljeborgii* (Danielssen, 1859)
 - phippsii* (Krøyer, 1841)
 - spinus* (Sowerby, 1805)
- Thor**
- amboinensis* (De Man, 1888)
- Thoralus**
- cranchii* (Leach, 1817)
 - sollaudi* (Zariquey-Cenarro, 1935)
- Trachycaris**
- restricta* (A. Milne-Edwards, 1878)
- Family Alpheidae**
- Alpheopsis**
- africana* Holthuis, 1952
- Alpheus**
- dentipes* Guérin-Méneville, 1832
 - edwardsii* (Audouin, 1826)
 - glaber* (Olivier, 1792)
 - lobidens* De Haan, 1849
 - macrocheles* (Hailstone, 1835)
 - migrans* Lewinsohn & Holthuis, 1978
 - platydactylus* Coutière, 1897
 - rapacida* De Man, 1908
 - talismani* Coutière, 1898
- Athanas**
- amazonae* Holthuis, 1951
 - grimaldii* Coutière, 1911
 - nitescens* (Leach, 1814)
- Automate**
- branchialis* Holthuis & Gottlieb, 1958
- Salmoneus**
- jarli* (Holthuis, 1951)

- sketi* Fransen, 1992
- Synalpheus**
- gambareloides* (Nardo, 1847)
 - tumidomanus* Paulson, 1875 subsp. *africanus* Crosnier & Forest, 1965
- Family Gnathophyllidae**
- Gnathophyllum*
 - americanum* Guérin-Méneville, 1855
 - elegans* (Risso, 1816)
- Family Processidae**
- Processa*
 - acutirostris* Nouvel & Holthuis, 1957
 - canaliculata* Leach, 1815
 - edulis* subsp. *edulis* (Risso, 1816)
 - edulis* (Risso, 1816) subsp. *arcassonensis* (Nouvel & Holthuis, 1957)
 - edulis* (Risso, 1816) subsp. *crassipes* (Nouvel & Holthuis, 1957)
 - elegantula* Nouvel & Holthuis, 1957
 - intermedia* Holthuis, 1951
 - macrodactyla* Holthuis, 1952
 - macrophthalma* Nouvel & Holthuis, 1957
 - modica* subsp. *modica* Williamson & Rochanaburanon, 1979
 - modica* Williamson & Rochanaburanon, 1979 subsp. *carolii* Williamson & Rochanaburanon, 1979
 - nouveli* subsp. *nouveli* Al-Adhub & Williamson, 1975
 - nouveli* Al-Adhub & Williamson, 1975 subsp. *holthuisi* Al-Adhub & Williamson, 1975
 - parva* Holthuis, 1951
 - pontica* (Sowinsky, 1882)
 - robusta* Nouvel & Holthuis, 1957
- Family Bathypalaemonellidae**
- Bathypalaemonella*
 - serratipalma* Pequegnat, 1970
- Family Palaemonidae**
- Balssia*
 - gasti* (Balss, 1921)
 - noeli* Bruce, 1998 - Brachycarpus*
 - biunguiculatus* (Lucas, 1846) - Leander*
 - tenuicornis* (Say, 1818) - Palaemon*
 - adspersus* Rathke, 1837
 - elegans* Rathke, 1837
 - longirostris* subsp. *longirostris* H. Milne-Edwards, 1837
 - longirostris* H. Milne-Edwards, 1837 subsp. *garciacaudi* Zariquey-Alvarez, 1968
 - serratus* (Pennant, 1777)
 - xiphias* Risso, 1816 - Palaemonella*
 - atlantica* Holthuis, 1951
 - rotumana* (Borradaile, 1898) - Palaemonetes*
 - antennarius* (H. Milne-Edwards, 1837)
 - mesogenitor* Sollaard, 1912
 - mesopotamicus* Pesta, 1913
 - turcorum* Holthuis, 1961
 - varians* (Leach, 1814)
 - zariqueyi* Sollaard, 1939 - Periclimenes*
 - aegylios* Grippo & d'Udekem d'Acoz, 1996
 - amethysteus* (Risso, 1827)
 - calmani* Tattersal, 1921
 - eleutheriou* Koukouras & Türlky, 1996
 - granulatus* Holthuis, 1950
 - kornii* (Lo Bianco, 1903)
 - sagittifer* (Norman, 1861)
 - scriptus* (Risso, 1822)
 - wirtzi* d'Udekem d'Acoz, 1996 - Pontonia*
 - domestica* Gibbes, 1850
- flavomaculata* Heller, 1864
- pinnophylax* (Otto, 1821)
- Tuleariocaris**
- neglecta* Chace, 1969
- Typton**
- gnathophylloides* Holthuis, 1951
 - spongicola* O.G. Costa, 1844
- Family Rhynchocinetidae**
- Cinetorhynchus*
 - rigens* (Gordon, 1936)
- Family Stylodactylidae**
- Styloceratulus*
 - serratus* A. Milne-Edwards, 1881
- Family Amphionidae**
- Amphionides*
 - reynaudi* (H. Milne-Edwards, 1833)
- Family Crangonidae**
- Aegaeon*
 - cataphractus* (Olivier, 1792)
 - lacazei* (Gourret, 1887) - Argis*
 - dentata* (Rathbun, 1902) - Crangon*
 - allmanni* Kinahan, 1857
 - crangon* (Linnaeus, 1758) - Metacrangon*
 - jacquetii* subsp. *jacquetii* (A. Milne-Edwards, 1881) - Parapontophilus*
 - abyssi* (Smith, 1884)
 - gracilis* (Smith, 1882)
 - talismanni* (Crosnier & Forest, 1973) - Philoceras*
 - bispinosus* (Hailstone, 1835) subsp. *neglectus* (Sars, 1883)
 - bispinosus* subsp. *bispinosus* (Hailstone, 1835)
 - echinulatus* (Sars, 1861)
 - fasciatus* (Risso, 1816)
 - monacanthus* (Holthuis, 1961)
 - sculptus* (Bell, 1847)
 - trispinosus* (Hailstone, 1835) - Pontophilus*
 - norvegicus* (Sars, 1861)
 - spinosis* (Leach, 1815) - Sabinea*
 - hystric* (A. Milne-Edwards, 1881)
 - sarsi* Smith, 1879
 - septemcarinata* (Sabine, 1824) - Sclerocrangon*
 - boreas* (Phipps, 1774)
 - ferox* (Sars, 1877)

Family Glyphocrangonidae

 - Glypocrangon*
 - atlantica* Chace, 1939
 - longirostris* (Smith, 1882)
 - sculpta* (Smith, 1882)

Family Enoplometopodidae

 - Enoplometopus*
 - antillensis* Lütken, 1865
 - callistus* Intés & Le Loeuff, 1970

Family Nephropidae

 - Homarus*
 - gammarus* (Linnaeus, 1758) - Nephrops*
 - norvegicus* (Linnaeus, 1758) - Nephropsis*
 - atlantica* Norman, 1882

Family Polychelidae

 - Polycheles*
 - crucifer* (Willemoes-Suhm, 1874)
 - granulatus* Faxon, 1893
 - typhlops* subsp. *typhlops* Heller, 1862
 - validus* (A. Milne-Edwards, 1880) - Stereomastis*

- grimaldii* (Bouvier, 1905)
nana (Smith, 1884)
sculpta subsp. *sculpta* (Smith, 1880)
Willemoesia
forceps A. Milne-Edwards, 1880
leptodactyla (Willemoes-Suhm, 1873)
- Family Palinuridae**
Palinurus
elephas (Fabricius, 1787)
mauritanicus Gruvel, 1911
Panulirus
echinatus Smith, 1869
ornatus (Fabricius, 1798)
regius De Brito Capello, 1864
- Family Scyllaridae**
Scyllarides
latus (Latreille, 1803)
Scyllarus
arcutus (Linnaeus, 1758)
caparti Holthuis, 1952
posteli Forest, 1963
pygmaeus (Bate, 1888)
- Family Axiidae**
Axius
stirynchus Leach, 1815
Calocarides
coronatus (Trybom, 1904)
Levantocaris
hornungae Galil & Clark, 1993
- Family Calocarididae**
Calastacus
laevis de Saint Laurent, 1972
Calocaris
macandreae Bell, 1846
- Family Laomediidae**
Jaxea
nocturna Nardo, 1847
- Family Callianassidae**
Callianassa
acanthura Caroli, 1946
candida (Olivier, 1792)
subterranea (Montagu, 1808)
truncata Giard & Bonnier, 1890
tyrrhena (Petagna, 1792)
whitei Sakai, 1999
Calliapagrops
charcoti de Saint Laurent, 1973
Calliax
lobata (de Gaillande & Lagardère, 1966)
Paraglypturus
punica (de Saint Laurent & Manning, 1982)
- Family Ctenochelidae**
Gourretia
denticulata (Lutze, 1937)
- Family Upogebiidae**
Gebiacantha
talismani (Bouvier, 1915)
Upogebia
deltaura (Leach, 1815)
mediterranea Noël, 1992
pusilla (Petagna, 1792)
stellata (Montagu, 1808)
tipica (Nardo, 1868)
- Family Parapaguridae**
Oncopagurus
bicristatus (A. Milne-Edwards, 1880)
Paragiopagurus
ruticheles (A. Milne-Edwards, 1891)
Parapagurus
abyssorum (Filhol, 1885)
alaminos Lemaire, 1986
nudus (A. Milne-Edwards, 1891)
- pilosimanus* Smith, 1879
Strobopagurus
gracilipes (A. Milne-Edwards, 1891)
Sympagurus
acinops Lemaire, 1898
- Family Diogenidae**
Calcinus
paradoxus Bouvier, 1922
tubularis (Linnaeus, 1767)
Clibanarius
aequabilis Dana, 1851
erythropus (Latreille, 1818)
Dardanus
arrosor (Herbst, 1796)
calidus (Risso, 1827)
Diogenes
pugilator (Roux, 1829)
Paguristes
eremita (Linnaeus, 1767)
marocanus A. Milne-Edwards & Bouvier, 1891
rubropictus A. Milne-Edwards & Bouvier, 1892
streaensis Pastore, 1984
syrtensis De Saint Laurent, 1971
- Family Paguridae**
Anapagurus
adriaticus Garcia-Gomez, 1994
alboranensis Garcia-Gomez, 1994
bicorniger A. Milne-Edwards & Bouvier, 1892
breviaculeatus Fenizia, 1937
chiroacanthus (Lilljeborg, 1856)
curvidactylus Chevreux & Bouvier, 1892
hyndmanni (Bell, 1845)
laevis (Bell, 1845)
longispina A. Milne-Edwards & Bouvier, 1900
petiti Dechancé & Forest, 1962
pusillus Henderson, 1888
smythi Ingle, 1993
Catapaguroides
iris Bouvier, 1922
megalops A. Milne-Edwards & Bouvier, 1892
microps A. Milne-Edwards & Bouvier, 1892
Cestopagurus
timidus (Roux, 1830)
Michelopagurus
atlanticus (Bouvier, 1922)
richardi (Bouvier, 1922)
Nematopagurus
longicornis A. Milne-Edwards & Bouvier, 1892
Pagurus
alatus Fabricius, 1775
anachoretus Risso, 1827
bernhardus (Linnaeus, 1758)
carneus (Pocock, 1889)
chevreuxi (Bouvier, 1896)
cuanensis Bell, 1845
excavatus (Herbst, 1791)
forbesii Bell, 1845
irregularis (A. Milne-Edwards & Bouvier, 1892)
prideaux Leach, 1815
pubescens Krøyer, 1838
pubescentulus (A. Milne-Edwards & Bouvier, 1892)
Spiropagurus
elegans Miers, 1881
- Family Lithodidae**
Lithodes
maja (Linnaeus, 1758)
Neolithodes
grimaldii (A. Milne-Edwards & Bouvier, 1894)
Paralithodes
camtschatcica (Tilesius, 1815)
Paralomis
bouvieri Hansen, 1908
microps Filhol, 1884
spectabilis Hansen, 1908

Family Chirostyliidae

- Eumunida*
bella de Saint Laurent & Macpherson, 1990
Gastropytchus
formosus (Filhol, 1884)
Uroptychus
bouvieri Caullery, 1896
concolor (A. Milne-Edwards & Bouvier, 1894)
maroccanus Türkay, 1976
rubrovittatus (A. Milne-Edwards, 1881)

Family Galatheidae

- Galathea*
agassizii A. Milne-Edwards, 1880
bolivari Zariquey-Alvarez, 1950
cenarroi Zariquey-Alvarez, 1968
dispersa Bate, 1859
faiali Nunes-Ruivo, 1961
intermedia subsp. *intermedia* Lilljeborg, 1851
intermedia Lilljeborg, 1851 subsp. *parroceli* Gourret, 1887
machadoi Barrois, 1888
nexa Embleton, 1834
rufipes A. Milne-Edwards & Bouvier, 1894
squamifera Leach, 1814
strigosa (Linnaeus, 1767)

Munida

- curvimanus* A. Milne-Edwards & Bouvier, 1894
intermedia A. Milne-Edwards & Bouvier, 1899
microphthalma A. Milne-Edwards, 1880
perarmata A. Milne-Edwards & Bouvier, 1894
rugosa (Fabricius, 1775)
rutllanti Zariquey-Alvarez, 1952
sanctipauli Henderson, 1885
sarsi Huus, 1935
speciosa von Martens, 1878
subcaeca Bouvier, 1922
tenuimana Sars, 1872
- Munidopsis*
abyssorum A. Milne-Edwards & Bouvier, 1897
acuta (A. Milne-Edwards, 1881)
acutispina Benedict, 1902
allae Khodkina & Duri_, 1989
antonii (Filhol, 1884)
bairdii (Smith, 1884)
bermudezi Chace, 1939
crassa Smith, 1885
curvirostra Whiteaves, 1874
livida (Perrier, 1886)
marionis (A. Milne-Edwards, 1882)
parfaii (A. Milne-Edwards & Bouvier, 1894)
polymorpha Koelbel, 1892
rostrata (A. Milne-Edwards, 1880)
serricornis (Lovén, 1853)
similis Smith, 1885
subsquamosa Henderson, 1885
sundi Sivertsen & Holthuis, 1956
thieli Türkay, 1975
vaillanti (A. Milne-Edwards, 1881)

Family Porcellanidae

- Pisidia*
blutelia (Risso, 1816)
longicornis (Linnaeus, 1767)
longimana (Risso, 1816)
- Porcellana*
platycheles (Pennant, 1777)

Family Albuneidae

- Albunea*
carabus (Linnaeus, 1758)
paretii Guérin-Méneville, 1853

Family Homolodromiidae

- Dicranodromia*
mahieuxii A. Milne-Edwards, 1883
pequegnati Guinot, 1995

Family Homolidae

- Homola*
barbata (Fabricius, 1793)
Homologenus
boucheti Guinot & Richer de Forges, 1995
Paromola
cuvieri (Risso, 1816)

Family Latreilliidae

- Latreillia*
elegans subsp. *elegans* Roux, 1830

Family Dromiidae

- Dromia*
marmorea Forest, 1974
nodosa A. Milne-Edwards & Bouvier, 1898
personata (Linnaeus, 1758)
spinirostris Miers, 1881

Family Cymonomidae

- Cymonomus*
granulatus (Norman, 1873)
normani Lankester, 1903

Family Raninidae

- Notopus*
dorsipes (Linnaeus, 1758)

Family Dorippidae

- Ethusa*
mascarone (Herbst, 1785)
microphthalmalma Smith, 1881
rosacea A. Milne-Edwards & Bouvier, 1897
- Ethusina*
alba (Filhol, 1884)
talismani A. Milne-Edwards & Bouvier, 1897
- Medorippa*
lanata (Linnaeus, 1767)

Family Calappidae

- Acanthocarpus*
brevispinis Monod, 1946
- Calappa*
granulata (Linnaeus, 1758)
pelii Herklots, 1851
rissiana Pastore, 1995
turkayana Pastore, 1995
- Cryptosoma*
cristatum Brullé, 1837

Family Matutidae

- Ashtoret*
lunarit (Forskål, 1775)

Family Leucosiidae

- Ebalia*
affinis Miers, 1881
cranchii Leach, 1817
deshayesi Lucas, 1846
edwardsii Costa, 1838
fragifera Miers, 1881
granulosa H. Milne-Edwards, 1837
nux A. Milne-Edwards, 1883
tuberculata Miers, 1881
tuberosa (Pennant, 1777)
tumefacta (Montagu, 1808)

- Ilia*
nucleus (Linnaeus, 1758)
spinosa Miers, 1881

- Ixa*
monodi Holthuis & Gottlieb, 1956

- Leucosia*
signata Paulson, 1875

- Meroeryptus*
boletifer A. Milne-Edwards & Bouvier, 1894
obsoletus A. Milne-Edwards & Bouvier, 1898

- Myra*
fugax (Fabricius, 1798)

Family Majidae

- Acanthonyx*
brevifrons A. Milne-Edwards, 1869

- lunulatus* (Risso, 1816)
- Achaeus**
- cranchii* Leach, 1817
 - gracilis* O.G. Costa, 1839
- Chionoecetes**
- opilio* (Fabricius, 1788)
- Dorhynchus**
- thomsoni* Thomson, 1873
- Ergasticus**
- clouei* A. Milne-Edwards, 1882
- Eury nome**
- aspera* (Pennant, 1777)
 - spinosa* Hailstone, 1835
- Herbstia**
- condyliata* (Fabricius, 1787)
 - rubra* A. Milne-Edwards, 1869
- Hyas**
- araneus* (Linnaeus, 1758)
 - coarctatus* subsp. *coarctatus* Leach, 1815
- Hyastenus**
- hilgendorfi* De Man, 1887
- Inachus**
- aguiarii* De Brito Capello, 1876
 - communissimus* Rizza, 1839
 - dorsettensis* (Pennant, 1777)
 - grallator* Manning & Holthuis, 1981
 - guentheri* (Miers, 1879)
 - leptochirus* Leach, 1817
 - nanus* Manning & Holthuis, 1981
 - parvirostris* (Risso, 1816)
 - phalangium* (Fabricius, 1775)
 - thoracicus* Roux, 1830
- Libinia**
- dubia* H. Milne-Edwards, 1834
- Lissa**
- chiragra* (Fabricius, 1775)
- Macropodia**
- czernjawskii* (Brandt, 1880)
 - deflexa* Forest, 1978
 - hesperiae* Manning & Holthuis, 1981
 - intermedia* Bouvier, 1940
 - linaresi* Forest & Zariquey-Alvarez, 1964
 - longipes* (A. Milne-Edwards & Bouvier, 1899)
 - longirostris* (Fabricius, 1775)
 - parva* Van Noort & Adema, 1985
 - rostrata* (Linnaeus, 1761)
 - tenuirostris* (Leach, 1814)
- Maja**
- brachydactyla* Balss, 1922
 - crispata* Risso, 1827
 - goltziana* d'Oliveira, 1888
 - squinado* (Herbst, 1788)
- Micippa**
- thalia* (Herbst, 1803)
- Micropisa**
- ovata* Stimpson, 1858
- Pisa**
- armata* (Latreille, 1803)
 - carinimana* Miers, 1879
 - hirticornis* (Herbst, 1804)
 - muscosa* (Linnaeus, 1758)
 - nodipes* (Leach, 1815)
 - tetraodon* (Pennant, 1777)
- Rochinia**
- carpenteri* (Thomson, 1873)
 - crassa* (A. Milne-Edwards, 1879)
 - rizzoana* (Roux, 1828)
- Stenorhynchus**
- lanceolatus* (Brullé, 1837)
- Family Parthenopidae**
- Heterocrypta*
- marionis* A. Milne-Edwards, 1881
- Parthenope*
- angulifrons* Latreille, 1825
 - expansa* (Miers, 1879)
- macrochelos* (Herbst, 1790)
- massena* (Roux, 1830)
- miersii* (A. Milne-Edwards & Bouvier, 1898)
- Family Corystidae**
- Corystes*
- cassivelalaunus* (Pennant, 1777)
- Family Atelecyclidae**
- Atelecyclus*
- rotundatus* (Olivi, 1792)
 - undecimdentatus* (Herbst, 1793)
- Family Cancridae**
- Cancer*
- bellianus* Johnson, 1861
 - pagurus* Linnaeus, 1758
- Family Pirimelidae**
- Pirimela*
- denticulata* (Montagu, 1808)
- Sirpus*
- ponticus* Vereshchaka, 1989
 - zariqueyi* Gordon, 1953
- Family Thiidae**
- Thia*
- scutellata* (Fabricius, 1793)
- Family Portunidae**
- Bathynectes*
- longipes* (Risso, 1816)
 - maravigna* (Prestandrea, 1839)
- Callinectes*
- danae* Smith, 1869
 - sapidus* Rathbun, 1896
- Carcinus*
- aestuarii* Nardo, 1847
 - maenas* (Linnaeus, 1758)
- Charybdis*
- helleri* (A. Milne-Edwards, 1867)
 - longicollis* Leene, 1938
- Laleonectes*
- vocans* (A. Milne-Edwards, 1878)
- Liocarcinus*
- bolivari* (Zariquey-Alvarez, 1948)
 - corrugatus* (Pennant, 1777)
 - depurator* (Linnaeus, 1758)
 - holstus* (Fabricius, 1798)
 - maculatus* (Risso, 1827)
 - marmoreus* (Leach, 1814)
 - macleayi* (Barnard, 1947)
 - navigator* subsp. *navigator* (Herbst, 1794)
 - navigator* (Herbst, 1794) subsp. *rondeletii* (Risso, 1816)
 - pusillus* (Leach, 1815)
 - vernalis* (Risso, 1827)
 - zariqueyi* (Gordon, 1968)
- Macropipus*
- rugosus* (Doflein, 1904)
 - tuberculatus* (Roux, 1830)
- Necora*
- puber* (Linnaeus, 1767)
- Polybius*
- henslowii* Leach, 1820
- Portumnus*
- latipes* (Pennant, 1777)
 - lysianassa* (Herbst, 1796)
- Portunus*
- hastatus* (Linnaeus, 1767)
 - inaequalis* (Miers, 1881)
 - pelagicus* (Linnaeus, 1758)
 - sayi* (Gibbes, 1850)
- Thalamita*
- gloriensis* Crosnier, 1962
 - poissonii* (Audouin, 1826)
- Xaiva*
- biguttata* (Risso, 1816)
- Family Geryonidae**

- Chaceon**
affinis (A. Milne-Edwards & Bouvier, 1894)
gordonae (Ingle, 1985)
inglei Manning & Holthuis, 1989
mediterraneus Manning & Holthuis, 1989
- Geryon**
longipes A. Milne-Edwards, 1882
trispinosus (Herbst, 1803)
- Zariquiey**
inflatus Manning & Holthuis, 1989
- Family Panopeidae**
Dyspanopeus
sayi (Smith, 1869)
Panopeus
africanus A. Milne-Edwards, 1867
- Rhithropanopeus**
harrisi (Gould, 1841)
- Family Trapeziidae**
Coralliope
parvula (A. Milne-Edwards, 1869)
- Domecia**
acanthophora (Desbonne & Schramm, 1867) subsp.
africana Guinot, 1964
- Family Xanthidae**
Atergatis
roseus (Rüppell, 1830)
Daira
perlata (Herbst, 1790)
Euryoxius
bouvieri (A. Milne-Edwards, 1869)
Glyptoxanthus
cavernosus (A. Milne-Edwards, 1878)
- Microcassiope**
minor (Dana, 1852)
- Monodaeus**
couchi (Couch, 1851)
guinotae Forest, 1976
rouxi (Capart, 1951)
- Nanocassiope**
melanodactyla (A. Milne-Edwards, 1867)
- Paractaea**
monodi Guinot, 1969
- Paragaleone**
longicrura (Nardo, 1868)
- Paraxanthias**
eriphoides (A. Milne-Edwards, 1867)
- Platypodiella**
picta (A. Milne-Edwards, 1869)
- Progeryon**
paucidens Bouvier, 1922
- Xantho**
granulicarpus Forest, 1953
hydrophilus (Herbst, 1790)
pilipes A. Milne-Edwards, 1867
poressa (Olivi, 1792)
sexdentatus (Miers, 1881)
- Family Eriphiidae**
Eriphia
verrucosa (Forskål, 1775)
- Globopilumnus**
africanus (A. Milne-Edwards, 1867)
- Sphaerozus**
nitidus Stimpson, 1858
- Family Pilumnidae**
Heteropanope
laevis (Dana, 1852)
- Pilumnopeus**
vauquelinii (Audouin, 1826)
- Pilumnus**
hirsutus Stimpson, 1858
hirtellus (Linnaeus, 1761)
inermis A. Milne-Edwards & Bouvier, 1894
spinifer H. Milne-Edwards, 1834
vilosissimus (Rafinesque, 1814)
- Family Pilumnoididae**
Pilumnoides
inglei Guinot & Macpherson, 1987
- Family Bythograeidae**
Segonzacia
mesatlantica (Williams, 1988)
- Family Carcinoplacidae**
Carcinoplax
barnardi Capart, 1951
- Family Euryplacidae**
Eucrate
crenata De Haan, 1835
- Machaerus**
atlanticus (Miers, 1881)
- Family Gonoplacidae**
Goneplax
rhombooides (Linnaeus, 1758)
- Family Grapsidae**
Brachynotus
atlanticus Forest, 1957
foresti Zariquiey-Alvarez, 1968
gemmellari (Rizza, 1839)
sedentatus (Risso, 1827)
- Eriocheir**
sinensis H. Milne-Edwards, 1853
- Euchirograpsus**
liguricus H. Milne-Edwards, 1853
- Grapsus**
adscencionis (Osbeck, 1765)
- Hemigrapsus**
penicillatus (De Haan, 1835)
- Pachygrapsus**
marmoratus (Fabricius, 1787)
maurus (Lucas, 1846)
transversus (Gibbes, 1850)
- Percnon**
gibbesi (H. Milne-Edwards, 1853)
- Plagusia**
depressa (Fabricius, 1775)
tuberculata Lamarck, 1818
- Planes**
cyaneus Dana, 1851
minutus (Linnaeus, 1758)
- Family Ocypodidae**
Macrophthalmus
graeffei A. Milne-Edwards, 1873
- Ocypode**
cursor (Linnaeus, 1758)
- Uca**
tangeri (Eyraud, 1835)
- Family Pinnotheridae**
Afropinnotheres
monodi Manning, 1993
- Asthenognathus**
atlanticus Monod, 1933
- Nepinnotheres**
pinnotheres (Linnaeus, 1758)
- Pinnotheres**
ascidicola Hesse, 1872
marioni Gourret, 1887
pectunculi Hesse, 1872
pisum (Linnaeus, 1767)
- Family Palicidae**
Palicus
caronii (Roux, 1830)
- Family Cryptochiridae**
Detocarcinus
balssi (Monod, 1956)

Thermosbaenacea

Compiled by Jacob van der Land

List derived from URMO.

Suborder Thermosbaenacea**Family Monodellidae**

- Tethysbaena*
aiakos Wagner, 1994
argentarii (Stella, 1951)
halophila (Karaman, 1953)
scabra (Pretus, 1991)
tarsiensis Wagner, 1994
Monodella
stygicola Ruffo, 1949

Family Halosbaenidae

- Halosbaena*
fortunata Bowman & Iliffe, 1986

Mysidacea

Compiled by Jacob van der Land and Torleiv Brattegard

List derived from URMO which is based on Müller (1993).

References

- Müller, H.-G., 1993. *World Catalogue and Bibliography of the Recent Mysidacea*. Müller Wissenschaft. Verlag, Wetzlar.
[also on floppy disk].

Order Mysidacea**Family Eucopiidae**

- Eucopia*
australis Dana, 1852
grimaldii Nouvel, 1942
hanseni Nouvel, 1942
major Hansen, 1910
sculpticauda Faxon, 1893
unguiculata (Willemoes-Suhm, 1875)
= *Chalaraspis unguiculata*

Family Gnathophausiidae

- Gnathophausia*
affinis G.O. Sars, 1883
gracilis Willemoes-Suhm, 1875
ingens (Dohrn, 1870)
= *G. bengalensis*
= *G. calcarata*
= *G. doryphora*
= *Lophogaster ingens*
zoea Willemoes-Suhm, 1875
Neognathophausia
gigas (Willemoes-Suhm, 1875)
= *Gnathophausia drepanephora*
= *Gnathophausia gigas*

Family Lophogastridae

- Lophogaster*
affinis Colosi, 1930
challengeri Fage, 1940
subglaber Hansen, 1927
typicus M. Sars, 1857
Paralophogaster
glaber Hansen, 1910

Family Mysidae

- Acanthomysis*
longicornis (Milne-Edwards, 1837)
= *Dasymysis longicornis*
= *Mysis longicornis*
= *Neomysis longicornis*
strauchi (Czerniavsky, 1882)
= *Metamysis strauchi*
= *Paramysis strauchi*
Amblyops
abbreviata (G.O. Sars, 1869)

= <i>Amblyopsis abbreviatum</i>	
<i>kempi</i> (Holt & Tattersall, 1905)	A
= <i>Pseudomma kempi</i>	
<i>spinifera</i> Nouvel & Lagardère, 1976	A
<i>tenuicauda</i> W. Tattersall, 1911	A
<i>trisetosa</i> Nouvel & Lagadère, 1976	A
<i>Amblyopsoides</i>	
<i>ohlini</i> (W. Tattersall, 1951)	A
= <i>Amblyops ohlini</i>	
<i>Anchialina</i>	
<i>agilis</i> (G.O. Sars, 1877)	A M
= <i>A. mediterranea</i>	
= <i>Anchialus agilis</i>	
<i>oculata</i> Hoenigman, 1960	M
<i>sanzoi</i> Coiffmann, 1937	M
<i>Anisomysis</i>	
<i>vasseuri</i> Ledoyer, 1974	M
<i>Arachnomysis</i>	
<i>leuckarti</i> Chun, 1887	A M
<i>Atlanteryrops</i>	
<i>crassipes</i> Nouvel & Lagardère, 1976	A
<i>Bathymysis</i>	
<i>helgae</i> W. Tattersall, 1907	A
<i>Boreomysis</i>	
<i>arctica</i> Krøyer, 1861	A M
= <i>Arctomyysis arctica</i>	
= <i>B. tregouhoffi</i>	
= <i>Mysis arctica</i>	
<i>atlantica</i> Nouvel, 1942	A
<i>bispinosa</i> O. Tattersall, 1955	A
<i>incisa</i> Nouvel, 1942	A
<i>inermis</i> Willemoes-Suhm, 1876	A
<i>longispina</i> Birstein & Tchindanova, 1958	A
<i>megalops</i> G.O. Sars, 1872	A M
<i>microps</i> G.O. Sars, 1883	A
<i>nobilis</i> G.O. Sars, 1885	A
<i>richardi</i> Nouvel, 1942	A
<i>scyphops</i> G.O. Sars, 1879	A
= <i>B. inermis</i> autt.	
<i>semicaeca</i> Hansen, 1905	A
<i>tridens</i> G.O. Sars, 1870	A
<i>vanhoeffeni</i> Zimmer, 1914	A
<i>Burrimysis</i>	
<i>palmeri</i> Jaume & Garcia, 1993	M
<i>Caesaromysis</i>	
<i>hispida</i> Ortmann, 1893	A M
<i>Calyptomma</i>	
<i>puritani</i> W. Tattersall, 1909	M
<i>Chunomysis</i>	
<i>diadema</i> Holt & Tattersall, 1905	A
<i>Dactylamblyops</i>	
<i>goniops</i> W. Tattersall, 1907	A
<i>sarsi</i> (Ohlin, 1901)	A
= <i>Amblyops sarsi</i>	
<i>thaumatops</i> W. Tattersall, 1907	A
<i>Dactyleryllops</i>	
<i>bidigitata</i> W. Tattersall, 1907	A
<i>dactylops</i> Holt & Tattersall, 1905	A
<i>dimorpha</i> Nouvel & Lagardère, 1976	A
<i>gracilura</i> W. Tattersall, 1907	A
<i>Diamysis</i>	
<i>bahirensis</i> (G.O. Sars, 1877)	A M
= <i>Mysis bahirensis</i>	
<i>pengoi</i> (Czerniavsky, 1882)	M
= <i>Potamomysis pengoi</i>	
<i>Erythrops</i>	
<i>abyssorum</i> G.O. Sars, 1869	A
<i>elegans</i> (G.O. Sars, 1863)	A M
= <i>E. pygmaea</i>	
= <i>Nematopus elegans</i>	
<i>erythrophthalma</i> (Goes, 1864)	A
= <i>E. goësi</i>	
<i>glacialis</i> G.O. Sars, 1885	A
<i>microps</i> (G.O. Sars, 1864)	A
= <i>E. microphthalmia</i>	

= <i>Nematopus microps</i>		= <i>Mysis hispida</i>	
<i>neapolitana</i> Colosi, 1929	A M	= <i>Nysidopsis hispida</i>	
<i>peterdohrni</i> Bacescu & Schiecke, 1974	M	<i>heterophila</i> Wittmann, 1986	A M
<i>serrata</i> (G.O. Sars, 1863)	A	<i>lingvura</i> (G.O. Sars, 1866)	A M
= <i>Nematopus serratus</i>		= <i>L. marioni</i>	
<i>Euchaetomera</i>		<i>mediterranea</i> (G.O. Sars, 1877)	A M
<i>glyphidophthalmica</i> Illig, 1906	A M	<i>megalops</i> Zimmer, 1915	A M
<i>intermedia</i> Nouvel, 1942	A M	<i>posidoniae</i> Wittmann, 1986	M
<i>richardi</i> Nouvel, 1945	M	<i>truncata</i> (Heller, 1863)	M
<i>tenuis</i> Sars, 1884	A M	= <i>L. sardica</i>	
= <i>Brutomyysis vogtii</i>		= <i>Mysis truncata</i>	
= <i>E. fowleri</i>		<i>Limnomyasis</i>	
<i>typica</i> Sars, 1884	A M	<i>benedeni</i> Czerniavsky, 1882	M
<i>Euchaetomeropsis</i>	A M	= <i>Mysidella bulgarica</i>	
<i>merolepis</i> (Illig, 1908)	A M	<i>Longithorax</i>	
= <i>Euchaetomera merolepis</i>		<i>alicei</i> Nouvel, 1942	A
<i>Gastrosaccus</i>		<i>fucus</i> Hansen, 1908	A
<i>lobatus</i> Nouvel, 1951	A M	<i>nouveli</i> O. Tattersall, 1955	A
<i>mediterraneus</i> Bacescu, 1970	M	<i>Mesopodopsis</i>	
<i>normani</i> G.O. Sars, 1877	A M	<i>aegyptica</i> Wittmann, 1992	M
<i>rosoffensis</i> Bacescu, 1970	A	<i>slabberi</i> (van Beneden, 1861)	A M
<i>sancus</i> (van Beneden, 1861)	A M	= <i>Mysis slabberi</i>	
= <i>Mysis sancta</i>		= <i>Podopsis cornuta</i>	
= <i>Pontomysis caucasica</i>		= <i>Podopsis goësi</i>	
= <i>Pontomysis widhalmi</i>		<i>Metamblyops</i>	
<i>spinifer</i> (Goës, 1864)	A M	<i>oculata</i> W. Tattersall, 1907	A
= <i>Acanthocaris livingstoneana</i>		<i>Meterythrops</i>	
= <i>G. spiniferus</i>		<i>picta</i> Holt & Tattersall, 1905	A M
= <i>Mysis spinifera</i>		<i>robusta</i> S.I. Smith, 1879	A
<i>Haplostylus</i>		<i>Michthyops</i>	
<i>bacescui</i> Hatzakis, 1977	M	<i>parva</i> (Vanhöffen, 1897)	A M
<i>magnilobatus</i> (Bacescu & Schiecke, 1974)	M	= <i>Pseudomma parvum</i>	
= <i>Gastrosaccus magnilobatus</i>		<i>theeli</i> (Ohlin, 1901)	A
<i>Hemimysis</i>		= <i>Pseudomma theeli</i>	
<i>abyssicola</i> G.O. Sars, 1869	A M	<i>Mysideis</i>	
<i>anomala</i> Sars, 1907	A M	<i>insignis</i> (G.O. Sars, 1864)	A
<i>lamornae</i> (Couch, 1856)	A M	= <i>Mysidopsis hibernica</i>	
= <i>H. pontica</i>		= <i>Mysis insignis</i>	
= <i>Mysis aurantia</i>		<i>parva</i> Zimmer, 1915	A M
= <i>Mysis lamornae</i>		<i>Mysidella</i>	
<i>margalefi</i> Alcaraz, Riera & Gili, 1986	M	<i>biscayensis</i> Lagardère & Nouvel, 1980	A M
<i>serrata</i> Bacescu, 1938	M	<i>typlops</i> G.O. Sars, 1872	A
<i>speluncola</i> Ledoyer, 1963	M	<i>typica</i> G.O. Sars, 1872	A M
<i>Heteroerythrops</i>	A	<i>Mysidetes</i>	
<i>purpura</i> O. Tattersall, 1955		<i>farrani</i> (Holt & Tattersall, 1905)	A M
<i>Heteromysis</i>		<i>Mysidopsis</i>	
<i>armoricana</i> Nouvel, 1940	A	<i>angusta</i> (G.O. Sars, 1864)	A M
<i>eideri</i> Bacescu, 1941	M	= <i>Mysis serraticauda</i>	
<i>lybiana</i> Bacescu, 1976	M	<i>didelphys</i> (Norman, 1863)	A M
<i>microps</i> (G.O. Sars, 1877)	A M	= <i>Mysis didelphys</i>	
= <i>Chiromysis microps</i>		<i>gibbosa</i> G.O. Sars, 1864	A M
<i>muelleri</i> Bacescu, 1976	M	<i>Mysis</i>	
<i>norvegica</i> G.O. Sars, 1883	A M	<i>litoralis</i> (Banner, 1948)	A
= <i>H. formosa</i> auctt.		= <i>Pugetomyysis litoralis</i>	
<i>Heteromysoides</i>		<i>mixta</i> Lilljeborg, 1852	A
<i>cotti</i> (Calman, 1932)	A	= <i>Mesomysis mixta</i>	
= <i>Heteromysis cotti</i>		= <i>Michtheimysis mixta</i>	
<i>Hypererythrops</i>		= <i>M. latitans</i>	
<i>richardi</i> Bacescu, 1941	M	<i>oculata</i> (Fabricius, 1780)	A
<i>seriventer</i> Holt & Tattersall, 1905	A	= <i>Megalophthalmus fabricianus</i>	
<i>zimmeri</i> Li, 1937	M	= <i>M. fabricii</i>	
<i>Imysis</i>		= <i>M. spinulosus</i>	
<i>atlantica</i> (Nouvel, 1942)	A	<i>polaris</i> Holmquist, 1959	A
= <i>Tenagomysis atlantica</i>		<i>relicta</i> Lovén, 1862	A
<i>Kainomatomysis</i>		<i>Neomysis</i>	
<i>foxi</i> W. Tattersall, 1927	M	<i>integer</i> (Leach, 1814)	A M
<i>Katamysis</i>		= <i>Mysis scoticus</i>	
<i>warpachowsky</i> G.O. Sars, 1877	M	= <i>Mysis vulgaris</i>	
<i>Katerythrops</i>	A	= <i>Praunus integer</i>	
<i>oceanae</i> Holt & Tattersall, 1905		<i>Paraleptomysis</i>	
<i>Leptomyysis</i>		<i>apiops</i> (G.O. Sars, 1877)	M
<i>buergii</i> Bacescu, 1966	M	= <i>Leptomyysis apiops</i>	
= <i>L. bürgei</i>		<i>banyulensis</i> (Bacescu, 1966)	M
<i>gracilis</i> (G.O. Sars, 1864)	A M	= <i>Leptomyysis apiops</i> subsp. <i>banyulensis</i>	
= <i>Mysis gracilis</i>		<i>Paramblyops</i>	

<i>bidigitata</i> W. Tattersall, 1911	A	= <i>Mysis kervillei</i>	
<i>rostrata</i> Holt & Tattersall, 1905	A M	= <i>Paramysis kervillei</i>	
<i>Paramysis</i>		<i>ornata</i> (G.O. Sars, 1864)	A
<i>agigensis</i> Bacescu, 1938	M	= <i>Mysis ornata</i>	
<i>arenosa</i> (G.O. Sars, 1877)	A M	= <i>Paramysis ornata</i>	
= <i>Mysis arenosa</i>		<i>parkeri</i> Norman, 1892	A
= <i>Schistomysis arenosa</i>		= <i>Mysis parkeri</i>	
<i>bacescoi</i> Labat, 1953	A	= <i>Paramysis parkeri</i>	
<i>baeri</i> Czerniavsky, 1882	M	= <i>Synmysis parkeri</i>	
<i>bakuensis</i> G.O. Sars, 1895	M	<i>spiritus</i> (Norman, 1860)	A
<i>festae</i> Colosi, 1920	M	= <i>Mysis spiritus</i>	
<i>helleri</i> (G.O. Sars, 1877)	M	= <i>Paramysis spiritus</i>	
= <i>Austromysis helleri</i>		= <i>Synmysis spiritus</i>	
= <i>Mysis helleri</i>		<i>Siriella</i>	
= <i>Schistomysis helleri</i>		<i>adriatica</i> Hoenigman, 1960	M
<i>intermedia</i> (Czerniavsky, 1882)	M	<i>armata</i> (Milne-Edwards, 1837)	A M
= <i>Heteromysis intermedia</i>		= <i>Cynthia flemingii</i>	
<i>kessleri</i> (Grimm, 1875)	M	= <i>Cynthilia frontalis</i>	
= <i>Mysis relicta</i> (Grim, 1875)		= <i>Mysis frontalis</i>	
<i>kosswigi</i> Bacescu, 1948	M	= <i>Mysis griffithsiae</i>	
<i>kroyeri</i> (Czerniavsky, 1882)	M	= <i>S. intermedia</i>	
= <i>Mesomysis kroyeri</i>		<i>castellabatensis</i> Ariani & Spagnuolo, 1975	M
<i>lacustris</i> (Czerniavsky, 1882)	A M	<i>clausi</i> G.O. Sars, 1877	A M
<i>nouveli</i> Labat, 1953	A	<i>jaltensis</i> Czerniavsky, 1868	A M
<i>pontica</i> (Bacescu, 1938)	M	= <i>S. gracilipes</i>	
<i>portzicensis</i> Nouvel, 1950	A	<i>norvegica</i> G.O. Sars, 1869	A M
<i>proconnesia</i> Colosi, 1922	M	<i>thompsonii</i> (H. Milne Edwards, 1837)	A M
<i>ullskyi</i> (Czerniavsky, 1882)	M	= <i>Cynthia inermis</i>	
<i>Parapseudomma</i>		= <i>Promysis galatheae</i>	
<i>calloplura</i> (Holt & Tattersall, 1905)	A M	= <i>S. edwardsii</i>	
= <i>Pseudomma calloplura</i>		<i>wolffi</i> O. Tattersall, 1961	A
<i>Parerythrops</i>		<i>Stiomysis</i>	
<i>bispinosa</i> Nouvel & Lagardère, 1976	A	<i>grandis</i> (Goes, 1863)	A
<i>lobiancoi</i> W. Tattersall, 1909	M	= <i>Mysideis grandis</i>	
<i>obesa</i> (G.O. Sars, 1864)	A M	= <i>Mysis grandis</i>	
= <i>Nematopus obesa</i>		<i>Teraterythrops</i>	
= <i>P. abyssicola</i>		<i>parva</i> (Zimmer, 1914)	A
<i>paucispinosa</i> Nouvel & Largardère, 1976	A	= <i>Katerynthia parva</i>	
<i>spectabilis</i> G.O. Sars, 1885	A	<i>Thalassomysis</i>	
<i>Praonus</i>	A	<i>tattersalli</i> Nouvel, 1942	A
<i>flexuosus</i> (Müller, 1776)		Family Petalophthalmidae	
= <i>Macromysis brevispinosus</i>		<i>Bacescomysis</i>	
= <i>Mysis flexuosa</i>		<i>abyssalis</i> (Lagardère, 1983)	A
= <i>Mysis spinulosus</i>		= <i>Hansenomysis abyssalis</i>	
<i>inermis</i> (Rathke, 1843)	A	<i>atlantica</i> (Lagardère, 1983)	A
= <i>Mysis cornuta</i>		= <i>Hansenomysis atlantica</i>	
= <i>Mysis inermis</i>		<i>Hansenomysis</i>	
= <i>Mysis truncatula</i>		<i>fallae</i> (Hansen, 1887)	A
<i>neglectus</i> (G.O. Sars, 1869)	A	= <i>Arctomysis fallae</i>	
= <i>Macromysis nigra</i>		<i>nouveli</i> Lagardère, 1983	A
= <i>Mysis neglecta</i>		<i>pseudofallae</i> Lagardère, 1983	A
= <i>Synmysis neglecta</i>		<i>Petalophthalmus</i>	
<i>Pseudomma</i>		<i>armiger</i> Willemoes-Suhm, 1875	A
<i>affine</i> G.O. Sars, 1870	A M		
<i>chattoni</i> Bacescu, 1941	M		
<i>frigidum</i> Hansen, 1908	A		
<i>jasi</i> Meland & Brattegard, 1995	A		
<i>kruppi</i> W. Tattersall, 1909	A M		
<i>nanum</i> Holt & Tattersall, 1906	A		
<i>roseum</i> G.O. Sars, 1870	A		
<i>truncatum</i> S.I. Smith, 1879	A		
<i>Pseudomysis</i>			
<i>abyssi</i> G.O. Sars, 1885	A		
<i>Pyroleptomytis</i>			
<i>perezi</i> (Bacescu, 1966)	M		
= <i>Leptomytis perezi</i>			
<i>rubra</i> (Wittmann, 1985)	M		
<i>Rhopalophthalmus</i>			
<i>flagellipes</i> Illig, 1906	A		
<i>mediterraneus</i> Nouvel, 1960	M		
<i>Schistomysis</i>			
<i>assimilis</i> (G.O. Sars, 1877)	M		
= <i>Mysis assimilis</i>			
= <i>Synmysis assimilis</i>			
<i>kervillei</i> (G.O. Sars, 1885)	A		

Amphipoda

Compiled by Denise Bellan-Santini and Mark J. Costello
The north-Atlantic lists were compiled by Mark J. Costello with assistance from Louise Collier from Costello *et al.* (1989), Brattegard (1997) and Vader *et al.* (1997). The Mediterranean and south Atlantic lists were compiled by Denise Bellan-Santini from Bellan-Santini *et al.* (1998), Marques and Bellan-Santini (1990, 1991), and Lopes *et al.* (1993). Additional species were then found in Dauvin (1999), and for Arctic seas in Palerud and Vader (1991) and Vader (1998).

Suborder, family, author and year were taken from Bellan-Santini *et al.* (1998) for species included in the 'Amphipoda of the Mediterranean' volumes, and from Vinogradov *et al.* (1996) for the Hyperiidea. The classification follows Bellan-Santini *et al.* (1998) and

Vinogradov *et al.* (1996). For other species in the list this information was obtained directly from the 'source'. Where this information was not available, it was obtained from Holmes *et al.* (1997), McCain (1968) and Stebbing (1906).

We thank Sandro Ruffo and J.C. Dauvin for their help in producing regional lists and comments on the amphipod list.

References

- Bellan-Santini, D., Karaman G.S., Ledoyer, M., Myers, A.A., Ruffo, S. and Vader, W. 1998. The Amphipoda of the Mediterranean, Part 4. *Mémoires de l'Institut Océanographique*, Monaco 13: XXVII- XLIV, 815-959.
- Bellan-Santini, D. and Dauvin J.C. 1997. Ampeliscidae (Amphipoda) from Iceland with a description of a new species (contribution to the BIOICE research programme). *Journal of Natural History* 31: 1157-1173.
- Brattegård, T. 1997. Order Amphipoda, Suborder Caprellidea (Phylum Crustacea) – caprellidean amphipods or skeleton shrimps. In: Brattegård, T., Holthe, T. (editors), *Distribution of marine, benthic macro-organisms in Norway. A tabulated catalogue*. Directorate for Nature Management, Trondheim, 213-214.
- Costello, M.J., Holmes, J.M.C., McGrath, D. and Myers, A.A. 1989. A review and catalogue of the Amphipoda (Crustacea) in Ireland. *Irish Fisheries Investigations*, Series B (Marine), No.33: 1-70.
- Dauvin J.C. 1999. Mise à jour de la liste des espèces d'Amphipodes (Crustacés Péracarides) présents en Manche. *Cahiers de Biologie Marine* 40: 165-183.
- Holmes, J.M.C., Costello, M.J. and Connor, D.W. 1997. Crustacea. In: C.M. Howson & B.E. Picton, *The Species Directory of the marine fauna and flora of the British Isles and surrounding seas*. Ulster Museum and The Marine Conservation Society, Belfast and Ross-on-Wye, 153-223.
- Lopes, M.F.R., Marques, J.C. and Bellan-Santini, D. 1993. The benthic amphipod fauna of the Azores (Portugal): an up-to-date annotated list of species, and some biogeographic considerations. *Crustaceana* 65: 204-217.
- Marques, J.C. and Bellan-Santini, D. 1993. Biodiversity in the ecosystem of the Portuguese continental shelf: distributional ecology and the role of benthic amphipods. *Marine Biology* 115: 555-564.
- McCain, J. 1968. *The Caprellidae (Crustacea: Amphipoda) of the Western North Atlantic*. Smithsonian Institution Press, Washington, D.C.
- Palerud R. and Vader W. 1991. *Marine Amphipoda Gammaridea in North-East Atlantic and Norwegian Arctic*. Tromsø 68: 97 pp.
- Stebbing, T.R.R. 1906. *Amphipoda I. Gammaridea*. Verlag von R. Friedlander und Sohn, Berlin.
- Vader W. 1998. Checklist of Amphipoda Gammaroidea in Barents Sea area. <http://www.imv.uit.no/ommuseet/enheter/zoo/wim/index.html>
- Vader, W., Brattegård, T., Buhl Mortensen, L. and Larsen, K. M. 1997. Order Amphipoda, Suborder Gammaridea (Phylum Crustacea) – gammaridean amphipods. In: Brattegård, T., Holthe, T. (editors). *Distribution of marine, benthic macro-organisms in Norway. A tabulated catalogue*. Directorate for Nature Management, Trondheim, 191-212.
- Vinogradov, M. E., Volkov, A.F. and Semenova T. N. 1996. *Hyperiid amphipods (Amphipoda, Hyperiidea) of the world oceans*. Science Publishers Inc., Lebanon USA, 632 pp.
- Order Amphipoda**
- Suborder Caprellidea**
- Family Caprellidae**
- Aeginella spinosa* Boeck, 1861
Aeginina longicornis (Kroyer, 1843)
- = *Aeginina echinata*
= *Aeginina laevia*
= *Aeginella spinosissima*
- Caprella**
- acanthifera* Leach, 1814
= *C. hystricula* (Kroyer)
andreae Mayer, 1890
ciliata Sars, 1883
daniilevskii Czerniavski, 1868
dilatata Kroyer, 1843
equilibra Say, 1818
= *C. aequilibra*
= *C. esmarckii*
= *C. laticornis* (Boeck)
erethizon Mayer, 1901
fretensis Stebbing, 1878
grandimana Mayer, 1882
hirnsuta Mayer, 1890
lilliput Krapp-Schickel & Ruffo, 1987
linearis Linnaeus, 1767
= *C. lobata* (Müller, Kroyer)
liparotensis Haller, 1879
= *C. dentata*
microtuberculata Sars, 1879
mitis Mayer, 1890
penantis (Leach, 1814)
= *C. acutifrons* Latreille, 1816
rapax Mayer, 1890
santosrosasi Sanchez-Moyano, Jimenez-Martin & Garcia-Gomez, 1995
septentrionalis Kroyer, 1838
= *C. loveni*
= *C. monocera* (Sars)
= *C. punctata* (Boeck)
telarpax Mayer, 1890
tuberculata Bate & Westwood, 1868
- Deutella**
- schieckeii* Cavedini, 1982
- Liropus**
- elongatus* Mayer, 1890
minimus Mayer, 1890
- Pariambus**
- typicus* (Kroyer, 1844)
= *Podalirius typicus* (Kroyer)
- Parvipalpus**
- capillaceus* (Chevreux, 1877)
linea Mayer, 1890
major Carausu, 1941
norvegicus (Stephensen, 1931)
- Pedoculina**
- bacescui* Carausu, 1940
garciagomezi Sanchez-Moyano, Carballo & Estacio, 1995
- Phthisica**
- marina* Slabber, 1749
= *Proto pedata* (Leach)
= *Proto ventricosa* (Müller)
- Proaeginina**
- norvegica* (Stephensen, 1931)
- Pseudolirius**
- kroyerii* (Haller, 1879)
- Pseudoprotella**
- phasma* (Montagu, 1804)
= *Protella phasma*
- Suborder Gammaridea**
- Family Acanthonotozomatidae**
- Acanthonotozoma**
- cristatum* (Ross, 1835)
inflatum (Kroyer, 1842)
magnum Just, 1978
monodontatum Kudzjaschov, 1965
rusanovae Bryazgin, 1974
serratum (Fabricius, 1780)
sinuatum Just, 1978

- Family Amathilopsidae**
- Amathilopsis*
 - affinis* Miers, 1881
 - atlantica* Chevreux, 1935
 - spinigera* Heller, 1875
- Family Ampeliscidae**
- Ampelisca*
 - abyssicola* Stebbing, 1888
 - aequicornis* Bruzelius, 1859
 - amblyops* Sars, 1895
 - anomala* Sars, 1883
 - anophthalma* Bellan-Santini & Kaim-Malka, 1977
 - antennata* Bellan-Santini & Kaim-Malka, 1977
 - armoricana* Bellan-Santini & Dauvin, 1981
 - brevicornis* (Costa, 1853)
 - = *A. bellianus*
 - = *A. laevigata* (Costa, 1853)
 - calypsonis* Bellan-Santini & Kaim-Malka, 1977
 - compacta* Norman, 1882
 - dalmatina* Karaman, 1975
 - declivitatis* Mills, 1967
 - diadema* (Costa, 1853)
 - = *A. assimilis* (Costa, 1853)
 - eschrichtii* Kroyer, 1842
 - gibba* Sars, 1882
 - heterodactyla* Schellenberg, 1925
 - intermedia* Bellan-Santini & Diviacco, 1990
 - islandica* Bellan-Santini & Dauvin, 1997
 - jaffensis* Bellan-Santini & Kaim-Malka, 1977
 - latifrons* Schellenberg, 1925
 - ledoyeri* Bellan-Santini & Kaim-Malka, 1977
 - lusitanica* Bellan-Santini & Marques, 1987
 - macrocephala* Liljeborg, 1852
 - = *A. latipes* Stephensen, 1925
 - massiliensis* Bellan-Santini & Kaim-Malka, 1977
 - melitae* Dauvin & Bellan-Santini, 1985
 - multispinosa* Bellan-Santini & Kaim-Malka, 1977
 - odontoplax* Sars, 1895
 - parabyblisoides* Dauvin & Bellan-Santini, 1996
 - planierensis* Bellan-Santini & Kaim-Malka, 1977
 - provincialis* Bellan-Santini & Kaim-Malka, 1977
 - pseudosarsi* Bellan-Santini & Kaim-Malka, 1977
 - pseudospinimana* Bellan-Santini & Kaim-Malka, 1977
 - pusilla* Sars, 1895
 - remora* Bellan-Santini & Dauvin, 1986
 - rubella* A. Costa, 1864
 - ruffoi* Bellan-Santini & Kaim-Malka, 1977
 - sarsi* Chevreux, 1888
 - serraticaudata* Chevreux, 1888
 - sorbei* Dauvin & Bellan-Santini, 1996
 - spinifer* Reid, 1951
 - spinimana* Chevreux, 1900
 - spinipes* Boeck, 1861
 - spooneri* Dauvin & Bellan-Santini, 1982
 - tenuicornis* Liljeborg, 1855
 - toulemoni* Dauvin & Bellan-Santini, 1982
 - truncata* Bellan-Santini & Kaim-Malka, 1977
 - typica* (Bate, 1856)
 - uncinata* Chevreux, 1887
 - unidentata* Schellenberg, 1936
 - verga* Reid, 1951
 - vervecei* Bellan-Santini & Kaim-Malka, 1977
- Byblis*
- abyssi* Sars, 1879
 - affinis* Sars, 1879
 - crassicornis* Metzger, 1875
 - erythrops* Sars, 1883
 - gaimardi* (Kroyer, 1846)
 - guernei* Chevreux, 1888
 - longicornis* Sars, 1891
 - minuticornis* Sars, 1879
 - = *B. serra*
- Haploops*
- abyssorum* Chevreux, 1908
 - dellavallei* Stebbing, 1893
- gascogni* Dauvin & Bellan-Santini, 1996
- laevis* Hoek, 1882
- nirae* Kaim-Malka, 1976
- proxima* Chevreux, 1919
- setosa* Boeck, 1871
- = *H. robusta* (Sars, 1891)
- = *H. sibirica* Gurjanova, 1929
- similis* Stephensen, 1925
- tenuis* Kanneworff, 1966
- tubicola* Liljeborg, 1855
- vallifera* Stephensen, 1925
- Family Amphilochidae**
- Amphilochoides*
 - boecki* Sars, 1892
 - = *A. odontonyx* (Sars, 1892)
 - serratipes* (Norman, 1869)
 - Amphilochopsis*
 - hamatus* Stephensen, 1925
 - Amphilochus*
 - borealis* Enequist, 1949
 - brunneus* Della Valle, 1893
 - manudens* Bate, 1862
 - neapolitanus* Della Valle, 1893
 - picadurus* Barnard, 1962
 - planierensis* Ledoyer, 1977
 - spencebati* (Stebbing, 1876)
 - = *Probolium spence-bati*
 - tenuimanus* Boeck, 1871
 - Gitana*
 - abyssicola* Sars, 1892
 - longicarpus* Ledoyer, 1977
 - rostrata* Boeck, 1871
 - sarsi* Boeck, 1871
 - Gitanopsis*
 - arctica* Sars, 1892
 - bispinosa* (Boeck, 1871)
 - inermis* Sars, 1882
 - Paramphilochoides*
 - intermedius* (Scott, 1896)
 - odontonyx* (Boeck, 1871)
 - = *Amphilochoides pusillus* (Boeck, 1871)
 - Peltocoxa*
 - brevirostris* (Scott & Scott, 1893)
 - = *Cyproidea brevirostris*
 - damnoniensis* (Stebbing, 1885)
 - gibbosa* (Schiecke, 1977)
 - marioni* Catta, 1875
 - mediterranea* Schiecke, 1977
 - Stegoplax*
 - longirostris* Sars, 1882
- Family Ampithoidae**
- Amphitholina*
 - cuniculus* (Stebbing, 1874)
 - Ampithoe*
 - fastidiosus* Mateus & Mateus, 1991
 - ferox* (Chevreux, 1902)
 - gammaroides* (Bate, 1856)
 - = *Pleonexes gammaroides*
 - helleri* Karaman, 1975
 - = *A. neglecta* Lincoln, 1976
 - pomboi* (Mateus & Alfonso, 1974)
 - ramondi* Audouin, 1826
 - = *A. vaillanti* Lucas
 - riedli* Krapp-Schickel, 1968
 - rubricata* (Montagu, 1808)
 - = *A. littorina* Bate & Westwood
 - = *Gammarus punctatus* Johnson
 - spuria* Krapp-Schickel, 1978
 - valida* Smith, 1873
 - Cymadusa*
 - crassicornis* (Costa, 1857)
 - filosa* Savigny, 1816
 - Sunamphitoe*
 - pelagica* (Milne-Edwards, 1830)

Family Anisogammaridae

Barrowgammarus
macginitiei Shoemaker, 1955

Family Aoridae

Aora
atlantidea Reid, 1951
gracilis (Bate, 1857)
spinicornis Afonso, 1976
Arctolembos
arcticus (Hansen, 1887)
Autoneo
angularis (Ledoyer, 1970)
borealis Myers, 1976
denticarpus (Myers & McGrath, 1978)
karamani Myers, 1976
longidigitana (Bonnier, 1896)
longipes (Liljeborg, 1852)
= *Lembos longipes* (Liljeborg, 1852)
megacheir Sars, 1879
rubromaculatus (Ledoyer, 1973)
spiniventris (Della Valle, 1893)
viduarum (Myers, 1974)

Bemlos

leptocheirus (Walker, 1909)

Lembos

denticarpus Myers & McGrath, 1978
hirsutipes Stebbing, 1895
websteri Bate, 1857
= *Autoneo websteri* Sars, 1894

Leptocheirus

bispinosus Norman, 1908
guttatus (Grube, 1864)
hirsutimanus (Bate, 1862)
longimanus Ledoyer, 1973
mariae Karaman, 1973
pectinatus (Norman, 1869)
pilosus Zaddach, 1844
tricristatus (Chevreux, 1887)

Microdeutopus

aligcola Della Valle, 1893
anomalus (Rathke, 1843)
= *Gammarus anomalus* Rathke, 1843
armatus Chevreux, 1887
bifidus Myers, 1977
chelifer (Bate, 1862)
damnoniensis (Bate, 1856)
gryllotalpa Costa, 1853
obtusatus Myers, 1973
similis Myers, 1977
sporadhi Myers, 1969
stationis Della Valle, 1893
versiculatus (Bate, 1856)
= *Coremapus versiculatus*

Tethylembos

viguieri (Chevreux, 1911)

Uncinotarsus

pellucidus L'Hardy & Truchot, 1964

Unciola

crassipes Hansen, 1887
crenatiopala (Bate, 1862)
incerta Bonnier, 1896
laticornis Hansen, 1887
leucopis (Kroyer, 1845)
petalocera (Sars, 1876)
planipes Norman, 1867
tenuipes Chevreux, 1919-20

Unciolella

lunata Chevreux, 1911

Family Argissidae

Argissa
hamatipes (Norman, 1869)
= *A. typica* (Boeck, 1871)
stebbingi Bonnier, 1896

Family Astyridae*Astyra*

abyssi Boeck, 1871

Family Biancolinidae

Biancolina
aligcola Della Valle, 1893

Family Bogidiellidae

Aurobogidiella
italica (Karaman, 1979)
Bogidiella
chappuisi Ruffo, 1952
dalmatina Karaman, 1953
helenae Mateus & Maciel, 1967
Marinobogidiella
tyrrhenica Schiecke, 1979

Family Carangoliopsidae

Carangoliopsis
spinulosa Ledoyer, 1970

Family Cheluridae

Chelura
terebrans Philippi, 1839

Family Colomastigidae

Colomastix
pusilla Grube, 1861
= *Exungia stilipes*

Family Corophiidae

Arctolembos
arcticus (Hansen, 1887)
= *Lembos arcticus* (Hansen, 1887)

Bathyphotis
tridentata Stephensen, 1944

Chevreuxius
grandimanus Bonnier, 1896

Corophium

acherusicum Costa, 1851
aculeatum Chevreux, 1908
acutum Chevreux, 1908
affine Bruzelius, 1859
annulatum Chevreux, 1908
arenarium Crawford, 1937
bonnellii (Milne-Edwards, 1830)
crassicornue Bruzelius, 1859
curvispinum Sars, 1895
insidiosum Crawford, 1937
lacustre Vanhoffen, 1911
minimum Schiecke, 1979
multisetosum Stock, 1952
orientale Schellenberg, 1928
rotundirostre Stephensen, 1915
runcicorne Della Valle, 1893
sextonae Crawford, 1937
volutator (Pallas, 1766)
= *C. grossipes* (Templeton, 1836)
= *C. longicornue* Latreille

Hansenella
longicornis Chevreux, 1909

Neohela

monstrosa (Boeck, 1861)
Siphonoecetes
dellavallei Stebbing, 1899
kroyeranus Bate, 1856
= *S. colletti* Boeck, 1862
neapolitanus Schiecke, 1979
pallidus Sars, 1882
sabatieri de Rouville, 1894
striatus Myers & McGrath, 1979

Family Cressidae

Cressa
abyssicola Sars, 1879
= *Danaia abyssicola* Sar, 1885
carinata Stephensen, 1931
cristata Myers, 1969
dubia (Bate, 1857)
mediterranea Ruffo, 1979

- minuta* Boeck, 1871
quinquedentata Stephensen, 1931
Cressina
monocuspis Stephensen, 1931
- Family Dexaminidae**
- Atylus*
carinatus (Fabricius, 1793)
falcatus Metzger, 1871
= *Neotropis falcatus*
= *Nototropis falcatus*
= *Paratylus unciatus* (Sars)
guttatus (Costa, 1851)
massiliensis Bellan-Santini, 1975
nordlandicus Boeck, 1871
smitti Goës, 1866
swammerdami (Milne-Edwards, 1830)
vedlomensis (Bate & Westwood, 1862)
- Dexamine*
spiniventris (Costa, 1853)
spinos (Montagu, 1813)
thea Boeck, 1861
- Guernea*
coailia (Norman, 1868)
- Tritaeta*
gibbosa (Bate, 1862)
- Family Epimeriidae**
- Epimeria*
cornigera (Fabricius, 1779)
= *Acanthonotus owenii* Bate & Westwood, 1863
= *Acanthonotus testudo* Bate, 1962
loricata Sars, 1879
parasitica (Sars, 1858)
tuberculata Sars, 1893
- Paramphithoe*
hystrix (Ross, 1835)
= *Acanthozone cuspidata* (Ross, 1835)
polyacantha (Murdoch, 1885)
- Family Eusiridae**
- Amphithopsis*
depressa Schiecke, 1976
longicaudata Boeck, 1861
- Apherusa*
alacris Krapp-Schickel, 1969
bispinosa (Bate, 1857)
= *Dexamine bispinosa*
chiereghinii Giordani- Soika, 1950
cirrus (Bate, 1862)
= *A. borealis* Sars, 1893
= *Pherusa bicuspis*
clevei Sars, 1904
corbeli Lagardere, 1968
glacialis (Hansen, 1887)
henneguyi Chevreux & Fage, 1925
jurinei (Milne-Edwards, 1830)
mediterranea Chevreux, 1911
megalops (Buchholz, 1874)
ovalipes Norman & Scott, 1906
ruffoi Krapp-Schickel, 1969
sarsi Shoemaker, 1930
= *A. megalops* (Sars, 1882)
tridentata (Bruzelius, 1859)
vexatrix Krapp-Schickel, 1979
- Bouvierella*
carcinophila Chevreux, 1900
- Calliopius*
laeviusculus (Kroyer, 1838)
= *C. rathkei*
- Cleippides*
bicuspis Stephensen, 1931
quadricuspis Heller, 1875
tricuspidis (Kroyer, 1846)
- Cleonardo*
appendiculatus (Sars, 1879)
microdactylus Stephensen, 1912
- spinicornis* Chevreux, 1908
Dautzenbergia
dentata (Chevreux, 1920)
grandimana (Chevreux, 1900)
megacheir (Walker, 1897)
- Eusirella*
elegans Chevreux, 1908
- Eusirogenes*
propinquus Scott, 1909
- Eusirooides*
dellavallei Chevreux, 1899
sarsi Chevreux, 1900
- Eusirus*
abyssi Stephensen, 1944
biscayensis Bonnier, 1896
cuspidatus Kroyer, 1845
holmi Hansen, 1887
leptocarpus Sars, 1893
longipes Boeck, 1861
minutus Sars, 1893
propinquus Sars, 1893
tjalfsiensis Stephensen, 1912
- Halirages*
elegans Norman, 1882
= *H. quadridentatus*
fulvocincta (Sars, 1858)
mixta Stephensen, 1931
nilssonii Ohlin, 1895
quadridentata Sars, 1876
- Haliragooides*
abyssi Gurjanova, 1951
inermis (Sars, 1882)
- Laothoes*
polylovi Gurjanova, 1945
meinerti Boeck, 1871
= *Laothoe meinerti* (Boeck, 1871)
- Leptamphopus*
massiliensis Ledoyer, 1977
sarsi Vanhoffen, 1897
= *L. longimanus* (Boeck, 1871)
- Oradarea*
longimana Walker, 1903
- Pontogeneia*
inermis (Kroyer, 1838)
- Rhachotropis*
aculeata (Lepechin, 1780)
arii Thurston, 1980
caeca Ledoyer, 1977
distincta (Holmes, 1908)
faeroensis (Stephensen, 1944)
gislii Thurston, 1980
glabra Ledoyer, 1977
grimaldii (Chevreux, 1888)
helleri (Boeck, 1871)
inermis Ledoyer, 1977
infata (Sars, 1882)
= *R. tumida* Sars, 1893 & 95
integricauda Carausu, 1948
kergueleni Stebbing, 1888
leucophthalma Sars, 1893
lomonosovi Gurjanova, 1934
macropus Sars, 1893
oculata (Hansen, 1887)
proxima Chevreux, 1911
rostrata Bonnier, 1896
thordisae Thurston, 1980
thorkelli Thurston, 1980
- Rozinante*
fragilis (Goes, 1866)
- Stenopleura*
atlantica Stebbing, 1888
- Family Gammaracanthidae**
- Gammaracanthus*
loricatus (Sabine, 1821 & 24)
- Family Gammarellidae**

- Gammarellus**
angulosus (Rathke, 1843)
 = *Amathilla angulosa* (Sars)
 = *Amathilla sabini* (Leach)
 = *G. carinatus* (Rathke, 1837)
homari (Fabricius, 1779)
- Family Gammaridae**
- Echinogammarus*
berilli (Catta, 1878)
dahli (Stock, 1968)
foxi (Schellenberg, 1928)
olivii (Milne-Edwards, 1830)
 = *Gammarus olivii*
pirloti (Sexton & Spooner, 1940)
 = *Chaetogammarus pirloti*
 = *Gammarus pirloti*
planicrurus (Reid, 1940)
 = *Pectenogammarus planicrurus* Reid, 1940
pungens (Milne-Edwards, 1840)
stocki Karaman, 1969
stoerensis (Reid, 1938)
 = *Chaetogammarus stoerensis*
 = *Gammarus stoerensis*
veneris (Heller, 1865)
- Gammarus*
aequicauda (Martyinov, 1931)
chevreuxi Sexton, 1913
crinicornis Stock, 1966
duebeni Liljeborg, 1852
 = *G. campylops*
 = *G. fluviatilis*
 = *G. wilkitzkii* Birula, 1897
finmarchicus Dahl, 1938
 = *Marinogammarus finmarchicus* (Dahl, 1938)
inaequicauda Stock, 1966
insensibilis Stock, 1966
locusta (Linnaeus, 1758)
 = *G. campylops* Leach
marinus Leach, 1815
 = *Echinogammarus marinus* (Leach, 1815)
 = *G. olivii* Milne-Edwards, 1830
 = *Marinogammarus marinus* (Leach, 1815)
obtusatus Dahl, 1938
 = *Eulimnogammarus obtusatus* (Dahl, 1938)
 = *Marinogammarus obtusatus* (Dahl, 1938)
oceanicus Segerstråle, 1947
pirloti (Sexton et Spooner, 1940)
salinus Spooner, 1947
setosus Dementieva, 1931
subtypicus Stock, 1966
tigrinus Sexton, 1939
wilkitzkii Birula, 1897
zaddachi Sexton, 1912
- Longigammarus*
bruni Karaman, 1969
- Lunulogammarus*
turcicus Krapp-Schickel, Ruffo & Schiecke, 1944
- Neogammarus*
adriaticus Karaman, 1973
festai Ruffo, 1937
nudus Stock, 1971
- Rhipidogammarus*
karamani Stock, 1971
rhipidiophorus (Catta, 1878)
- Weyprechtia*
heuglini (Buchholz, 1874)
pinguis (Kroyer, 1838)
- Family Haustoriidae**
- Haustorius*
algeriensis Mulot, 1968
arenarius (Slabber, 1767)
- Family Hyalidae**
- Hyale*
camptonyx (Heller, 1866)
- crassipes* (Heller, 1866)
grimaldii Chevreux, 1891
perieri (Lucas, 1849)
pontica Rathke, 1837
 = *H. lubboniana* Bate, 1875
prevosti (Milne-Edwards, 1830)
 = *H. nilssoni*
schmidti (Heller, 1866)
spinidactyla Chevreux, 1925
stebbingi Chevreux, 1888
 = *H. nilssoni*
Micropythia
carinata (Bate, 1862)

Parhyale

aqulina (Costa, 1857)
eburnea Krapp-Schickel, 1974
plumicornis (Heller, 1866)

Parhyalella
richardi (Chevreux, 1902)

Family Hyperiopsidae

Hyperiopsis
voringi Sars, 1885
 = *H. nilssoni*
Parargissa
nasuta Chevreux, 1908

Family Iphimedidiidae

Coboldus
nitior Krapp-Schickel, 1974

Iphimedia
brachygynatha Ruffo & Schiecke, 1979
carinata Heller, 1866
eblanae Heller, 1867
 = *I. carinata* Heller, 1866
 = *Panoploea eblanae*
gibbula Ruffo & Schiecke, 1979
jugoslavica Karman, 1975
minuta G.O. Sars, 1882
 = *Panoplea minuta* Stebbing, 1906, Chev. & Fage, 1925
nexa Myers & McGrath, 1987
obesa Rathke, 1843
perplexa Myers & Costello, 1987
quasimodus Ruffo & Schiecke, 1979
serratipes Ruffo & Schiecke, 1979
spatula Myers & McGrath, 1987
vicina Ruffo & Schiecke, 1979

Family Isaeidae

Cheiropotis
mediterraneus Myers, 1985

Gammaropsis
abyssalis Stephensen, 1944
crenulata Krapp-Schickel & Myers, 1979
dentata Chevreux, 1900
emancipata Krapp-Schickel & Myers, 1979
lobata (Chevreux, 1920)
maculata (Johnston, 1828)
 = *Eurystheus erythrophthalmus* (Liljeborg)
 = *G. erythrophalma* Liljeborg, 1855
melanops Sars, 1879
nitida (Stimpson, 1853)
 = *Podoceropsis excavata* Sars, 1894
 = *Podoceropsis nitida*
ostroumowi Sowinski, 1898
palmata (Stebbing & Robertson, 1891)
 = *Eurystheus palmatus*
 = *G. nana* Sars, 1894
pseudostroumowi Ledoyer, 1977
sophiae (Boeck, 1861)
 = *Gammarellus sophiae*
 = *Noenia undata* Bate & Westwood, 1863
 = *Podoceropsis sophiae* Boeck
togoensis (Schellenberg, 1925)
ulrici Krapp-Schickel & Myers, 1979

Isaea
elmhirsti Patience, 1909

- montagui* Milne-Edwards, 1830
Megamphopus
brevidactylus Myers, 1976
cornutus Norman, 1869
longicornis Chevreux, 1911
Microprotopterus
longimanus Chevreux, 1887
maculatus Norman, 1867
Photis
lamellifera Schellenberg, 1928
longicaudata (Bate & Westwood, 1862)
longipes (Della Valle, 1893)
pollex Walker, 1895
reinhardti Kroyer, 1842
= *P. pollex* Walker, 1895
tenuicornis Sars, 1882
- Family Ischyroceridae**
Bonnierella
abyssi (Chevreux, 1900)
abyssorum (Bonnier, 1896)
Cerapus
crassicornis (Bate, 1856)
Erichthonius
argenteus Krapp-Schickel, 1993
brasiliensis (Dana, 1855)
diformis Milne-Edwards, 1830
= *E. hunteri*
fasciatus (Stimpson, 1853)
megalops (Sars, 1879)
punctatus (Bate, 1857)
= *Cerapus abditus* Tempeleton
= *E. abditus* Bate
= *E. brasiliensis* (Dana, 1852)
rubricornis (Stimpson, 1853)
= *E. hunteri*
stephensi Myers & McGrath, 1984
Goesia
depressa (Goes, 1866)
= *Protomediea depressa*.
Ischyrocerus
anguipes Kroyer, 1838
= *I. minutus* Liljeborg
brevicornis (Sars, 1879)
commensalis Chevreux, 1900
enigmaticus Gurjanova, 1934
hanseni Stephensen, 1944
inexpectatus Ruffo, 1959
latipes Kroyer, 1842
= *I. assimilis*
megacheir (Boeck, 1871)
= *I. spitzbergensis*
megalops Sars, 1894
nanoides (Hansen, 1888)
stephensi Gurjanova, 1951
tenuicornis (Sars, 1885)
tuberculatus (Hoek, 1882)
Jassa
falcata (Montagu, 1808)
= *Cerapus falcata* Montagu
= *J. dentex* Czerniavski
= *J. pulchella*
= *Podocerus falcata*
herdmani (Walker, 1893)
marmorata (Holmes, 1903)
ocia (Bate, 1862)
pusilla (Sars, 1894)
= *J. dentex*
slatteryi Conlan, 1990
Microjassa
cumbrensis (Stebbing & Robertson, 1891)
= *Podoceropsis cumbrensis*
Parajassa
pelagica (Leach, 1814)
= *Janassa capillata* Sars, 1894
Protomediea
- fasciata* Kroyer, 1842
grandimana Brüggen, 1905
gurjanovae Bulycheva, 1951
- Family Lafystiidae**
Lafystius
sturionis Kroyer, 1842
= *L. sturionensis*
= *Laphystius sturionis* Kroyer, 1842
- Family Laphystiopsidae**
Iaphystiopsis
planifrons Sars, 1893
- Family Lepechinellidae**
Lepechinella
arctica Schellenberg, 1926
chrysotheras Stebbing, 1908
grimi Thurston, 1980
helgii Thurston, 1980
manco Barnard, 1973
skarphedini Thurston, 1980
Lepechinelloides
karti Thurston, 1980
- Family Leucothoidae**
Leucothoe
euryonyx Walker, 1901
incisa Robertson, 1892
lilljeborgi Boeck, 1861
= *L. imparicornis*
oboia Karaman, 1971
occulta Krapp-Schickel, 1975
pachycera Della Valle, 1893
procera Bate, 1857
richardii Lessona, 1865
rostrata Chevreux, 1908
serraticarpa Della Valle, 1893
spinicarpa (Abildgaard, 1789)
= *L. articulata*
spinulosa Chevreux, 1919-20
uschakovi Gurjanova, 1951
venetiarum Giordani- Soika, 1950
- Family Liljeborgiidae**
Idunella
aequicornis (Sars, 1876)
excavata (Schecke, 1973)
longirostris (Chevreux, 1920)
nana (Schecke, 1973)
pirata Krapp-Schickel, 1975
Liljeborgia
dellavallei Stebbing, 1906
fissicornis (Sars, 1858)
inermis Chevreux, 1919-20
kinahani (Bate, 1862)
macronyx (Sars, 1894)
pallida (Bate, 1857)
= *L. brevicornis* (Bruzelius)
psaltrica Krapp-Schickel, 1975
Listriella
dentipalma Dauvin & Gentil, 1983
mollis Myers & McGrath, 1983
picta Norman, 1889
spinifera Dauvin & Gentil, 1983
Sextonia
longirostris Chevreux, 1920
- Family Lysianassidae**
? *Gronella*
lobata (Chevreux, 1907)
Abyssorhomene
abyssorum (Stebbing, 1888)
chevreuxi (Stebbing, 1906)
Acidostoma
laticorne Sars, 1879
nodiferum Stephensen, 1923
= *A. sarsi*
obesum (Bate & Westwood, 1861)

- = *A. neglectum* (Bate & Westwood, 1861)
- Amaryllis*
haswelli Stebbing, 1888
= *Bathymaryllis haswelli* (Stebbing, 1888)
pulchella Bonnier, 1896
- Ambasia*
atlantica (Milne-Edwards, 1830)
= *A. danielsseni* (Milne-Edwards, 1830)
- Ambasiella*
murmanica (Brüggen, 1905)
= *Ambasia murmanica* (Brüggen, 1905)
- Anonyx*
affinis Ohlin, 1895
bispinosus Steele, 1967
compactus Gurjanova, 1962
debruyni Hoek, 1882
laticoxae Gurjanova, 1962
lilljeborgii Boeck, 1870
nugax (Phipps, 1774)
ochoticus Gurjanova, 1962
sarsi Steele & Brunel, 1968
stappersi Steele, 1989
- Aristias*
commensalis Bonnier, 1896
falcatus Stephensen, 1923
megalops Sars, 1895
microps Sars, 1895
neglectus Hansen, 1887
= *A. audouinianus*
topsenti Chevreux, 1900
tumidus (Kroyer, 1846)
- Aroui*
setosus Chevreux, 1911
- Boeckosinus*
abyssi (Oldwig, 1959)
= *Onisimus abyssi*
affinis (Hansen, 1887)
= *Onisimus affinis*
brevicaudatus (Hansen, 1887)
= *Onisimus brevicaudatus*
caricus (Hansen, 1887)
= *Onisimus caricus*
crassini (Gorbunov, 1946)
derjugini Gurjanova, 1929
= *Onisimus derjugini*
dubius Schellenberg, 1935
= *Onisimus dubius*
edwardsi (Kroyer, 1846)
= *Onisimus edwardsi*
krassini Gorbunov, 1946
= *B. crassini*
leucopis Sars, 1879
= *Onisimus leucopis*
normani (Sars, 1895)
= *Onisimus normani*
plautus Kroyer, 1845
= *Onisimus plautus*
sextonae Chevreux, 1925
= *Onisimus sextonae*
simus Gurjanova, 1962
= *Onisimus simus*
- Caeconyx*
caeculus Sars, 1891
= *Tmetonyx caeculus* Sars, 1891
- Centromedon*
calcaratus Sars, 1879
productus (Goes, 1866)
pumilus (Lilljeborg, 1865)
typhlops (Sars, 1879)
- Coximedon*
latimanus Sars, 1966
- Crybelocephalus*
megalurus Tattersall, 1906
- Cyclocaris*
guileimi Chevreux, 1899
- Cyphocaris*
anonyx Boeck, 1871
bouvieri Chevreux, 1916
polaris Gurjanova, 1951
richardi Chevreux, 1905
- Danaella*
mimonectes Stephensen, 1925
- Ensaya*
carpinei Bellan-Santini, 1974
- Euonyx*
biscayensis Chevreux, 1908
chelatus Norman, 1867
talismani Chevreux, 1919
- Eurythenes*
gryllus Lichtenstein, 1822
obesus (Chevreux, 1905)
- Gronella*
groenlandica (Hansen, 1888)
= *Orchomenella groenlandica*
- Hippomedon*
ambiguus Ruffo, 1946
bidentatus Chevreux, 1903
denticulatus (Bate, 1857)
= *Annoyx denticulatus* Bate
= *H. holbollii* Kroyer, 1846
frigidus Stephensen, 1923
gorbunovi Gurjanova, 1929
holbollii (Kroyer, 1846)
longimanus Stebbing, 1888
massiliensis Bellan-Santini, 1965
nasutus Stephensen, 1923
oculatus Chevreux & Fage, 1925
propinquus Sars, 1890
= *H. squamosus*
propinquus subsp. *petschoricus* Bryazgin, 1974
reticulatus Stephensen, 1923
robustus Sars, 1895
rylovi Gurjanova, 1933
serratipes Stephensen, 1923
striolatus Stephensen, 1923
- Hirondella*
triloculata Chevreux, 1889
- Hirondellea*
abyssalis (Stephensen, 1923)
- Ichnopus*
spinicornis Boeck, 1861
taurus Costa, 1853
- Katius*
obesus Chevreux, 1905
- Kerguelenia*
borealis Sars, 1891
reducta Ledoyer, 1977
- Koroga*
megalops Holmes, 1908
- Lepidepecrella*
cymba Goes, 1866
- Lepidepecreum*
clypeatum Chevreux, 1888
crypticum Ruffo & Schiecke, 1977
longicornis (Bate & Westwood, 1862)
= *L. carinatum* Bate & Westwood, 1868
= *L. longicorne*
serratum Stephensen, 1925
subclypeatum Ruffo & Schiecke, 1977
typhlops Bonnier, 1896
umbo (Goes, 1866)
- Lysianassa*
caesarea Ruffo, 1987
= *L. longicornis* Lucas
= *L. spinicornis* Costa
= *Lysianax certinus* Walker
ceratina (Walker, 1889)
costae (Milne-Edwards, 1830)
= *L. plumosa* Boeck, 1871
insperata (Lincoln, 1979)

- longicornis* (Lucas, 1849)
pilicornis (Heller, 1866)
plumosa Boeck, 1871
= *L. septentrionalis* Della Valle
Lysianella
dellavallei Stebbing, 1906
petalocera Sars, 1882
Martensia
martensi (Goes, 1866)
Menigrates
maslovi Bryazgin, 1974
obtusifrons (Boeck, 1861)
Menigratopsis
svennilssoni Dahl, 1945
Metacyphocaris
helgae Tattersall, 1906
Metambasia
faeroensis Stephensen, 1923
Nannonyx
goesi (Boeck, 1871)
propinquus Chevreux, 1911
spinimanus Walker, 1895
Normanion
abyssi Chevreux, 1903
amblyops Sars, 1895
chevreuxi Diviacco & Vader, 1988
quadrimanus (Bate et Westwood, 1868)
ruffoi Diviacco & Vader, 1988
sarsi Stebbing, 1906
= *N. quadrimanus* (Bate & Westwood, 1868)
Onesimoides
mediterraneus Bellan-Santini, 1974
Onisimus
birulai (Gurjanova, 1929)
glacialis (Sars, 1900)
litoralis (Kroyer, 1845)
= *Alibrotus litoralis* (Kroyer, 1845)
nanseni (Sars, 1900)
turgidus (Sars, 1879)
Opisa
eschrichtii (Kroyer, 1842)
Orchomene
abyssorum (Stebbing, 1865)
amblyops Sars, 1890
commensalis Chevreux et Fage, 1925
crenata Chevreux, 1925
crispatus (Goes, 1866)
faeroensis Stephensen, 1923
gerulicoris Shulenberger & Barnard, 1976
= *Orchomenella gerulicoris*
glabrus (Lagardere, 1968)
= *Orchomenella glabrus*
grimaldii Chevreux, 1890
humilis (Costa, 1853)
= *O. batei* Costa, 1853
laevipes Stephensen, 1923
lepidulus Gurjanova, 1962
= *Orchomenella lepidulus*
lobata (Chevreux, 1907)
macroserratus Shoemaker, 1930
massiliensis Ledoyer, 1977
minutus (Kroyer, 1846)
nanus (Kroyer, 1846)
= *Orchomenella nanus*
oystomus Stephensen, 1923
pectinatus Sars, 1882
pinguis (Boeck, 1861)
serratus (Boeck, 1861)
sibirjakovi Gurjanova, 1951
= *Orchomenella sibirjakovi*
similis (Chevreux, 1912)
thorii Stephensen, 1923
triangulus (Stephensen, 1925)
= *Orchomenella triangulus*
tschernyschevi Bruggen, 1909
- Orchomenella*
ciliata Sars, 1883
= *O. nanus*
minuta (Kroyer, 1846)
nana (Kroyer, 1846)
= *O. nanus*
obtusa (Sars, 1891)
= *Orchomene obtusus* Sars, 1895
pinguis (Boeck, 1861)
Pachychelium
davidis Stephensen, 1925
Paracallisoma
alberti Chevreux, 1903
Paracentromedon
crenatum Chevreux, 1900
Parachevreuxiella
lobata Andres, 1987
Paracyphocaris
praedator Chevreux, 1905
Paralibrotus
setosus Stephensen, 1923
Paralicella
caperesca Shulenberger & Barnard, 1976
tenuipes Chevreux, 1908
Paratryphosites
abyssi (Goes, 1866)
Pardia
punctata (Costa, 1851)
Paronesimus
barentsi Stebbing, 1894
= *Onisimus barentsi*
Perrierella
audouiniana (Bate, 1857)
Podopriion
bolivari Chevreux, 1891
Podopriionella
fissicaudata Ledoyer, 1977
norvegica Sars, 1895
Prachynella
mediterranea Ruffo, 1975
Rhinolabia
parthenopeia Ruffo, 1971
Schisturella
pulchra (Hansen, 1887)
Scopelocheiropsis
abyssalis Schellenberg, 1926
Scopelocheirus
abyssi Oldewig, 1959
crenatus Bate, 1857
= *Callisoma crenata* Bate, 1855
hopei (Costa, 1851)
polymedus Bellan-Santini, 1985
Socarnes
bidenticulatus (Bate, 1835)
erythrophthalmus Robertson, 1892
jilicornis (Heller, 1866)
= *S. crenulatus* (Chevreux, 1911)
= *Socarnopsis jilicornis* (Heller, 1866)
vahlii (Kroyer, 1838)
Sophrosyne
hispana Chevreux, 1888
robertsoni Stebbing & Robertson, 1891
Stephonyx
biscayensis (Chevreux, 1908)
talismanni (Chevreux, 1919-20)
Thoriella
islandica Stephensen, 1915
Tmetonyx
acutus (Sars, 1891)
albidus (Sars, 1891)
cicada (Fabricius, 1780)
= *Anonyx gulosus* Kroyer
= *Hoplornx gulosus*
gulosa (Kroyer, 1845)
leucophthalmus (Sars, 1891)

- morbiensis* Oleröd, 1987
nardonis (Heller, 1866)
rotundatus Chevreux, 1926
similis (Sars, 1891)
stephensi Chevreux, 1935
Trischizostoma
nicaeense (Costa, 1853)
= *Guerina nicaeensis* Chevreux, 1905
raschi Esmark & Boeck, 1861
Tryphosella
abyssalis (Stephensen, 1925)
angulata (Sars, 1895)
barentsi Gurjanova, 1929
biloba Stephensen, 1925
caecula (Sars, 1891)
= *Caeconyx caecula* Sars, 1891
compressa (Sars, 1895)
dilatata (Chevreux, 1903)
= *Orchomene dilatatus*
gracilipes Stephensen, 1925
horingi (Boeck, 1871)
= *Tryphosa horingi* (Boeck, 1871)
insignioidea Stephensen, 1925
insignis Bonnier, 1895
laevis (Bonnier, 1896)
longichela Stephensen, 1925
longidactyla Ruffo, 1985
minima Chevreux, 1911
nanoides (Lilljeborg, 1865)
orchomenoides Stephensen, 1925
oxystoma Stephensen, 1925
propinqua (Chevreux, 1926)
pusilla Sars, 1879
rotundata Stephensen, 1925
rusanovi Goes, 1866
= *Centromedon productus*
= *Centromedon rusanovi*
sarsi Bonnier, 1893
= *Orchomene nana*
= *Orchomene nanus*
= *Tryphosa nana*
= *T. gradimana*
scheideri (Stephensen, 1921)
simillima Ruffo, 1985
spitzbergensis (Chevreux, 1926)
= *Tmetonyx stephensi*
triangula Stephensen, 1925
trionyx Stephensen, 1925
tuberculimana Lagardere, 1968
Tryphosites
allenii Sexton, 1911
longipes (Bate & Westwood, 1861)
Uristes
abyssalis Stephensen, 1925
abyssi (Norman, 1900)
umbonatus (Sars, 1882)
Family Megaluropidae
Megaluropus
agilis Hoeck, 1889
massiliensis Ledoyer, 1976
monasteriensis Ledoyer, 1976
Family Melitidae
Abludomelita
aculeata (Chevreux, 1911)
gladiosa (Bate, 1862)
obtusata (Montagu, 1813)
= *Melita obtusata*
Allomelita
pellucida (Sars, 1882)
= *Melita pellucida* Sars, 1883
Ceradocus
orchestriipes Costa, 1853
semiserratus (Bate, 1862)
= *Maera semiserratus*
torelli (Goes, 1866)
Cheirocratus
assimilis (Lilljeborg, 1852)
intermedius Sars, 1894
monodontus Karaman, 1977
robustus Sars, 1894
sundevalli (Rathke, 1843)
= *Protomedia whitei* Bate
Degochirocratus
spani Karaman, 1989
Elasmopus
affinis Della Valle, 1893
brasiliensis (Dana, 1855)
pecteniferus (Bate, 1862)
perditus Reid, 1951
pocillimanus (Bate, 1862)
rapax Costa, 1853
vachoni Mateus & Mateus, 1966
Eriopisa
elongata (Bruzelius, 1859)
Eriopisella
pusilla Chevreux, 1920
ruffoi Marti & Villora-Moreno, 1996
Gammarella
fucicola (Leach, 1814)
= *Amphithoe fucicola* & *Pherusa fucicola*
= *G. brevicaudata* (Milne-Edward)
garciai Marti & Villora-Moreno, 1995
Maera
ariadne Krapp, Marti & Ruffo, 1996
aurora Krapp, Marti & Ruffo, 1996
grossimana (Montagu, 1808)
hamigera Haswell, 1880
hirondellei Chevreux, 1900
inaequipes (Costa, 1857)
knudseni Reid, 1951
loveni (Bruzelius, 1859)
othonis (Milne-Edwards, 1830)
= *Gammarus longimanus* Leach
= *Megamoera longimanus*
pachytelson Karaman & Ruffo, 1971
prionochira Bruggen, 1907
revelata Krapp, Marti & Ruffo, 1996
schieckei Karaman & Ruffo, 1971
schmidti Stephensen, 1915
sodalis Karaman & Ruffo, 1971
tenera Sars, 1885
Maerella
tenuimana (Bate, 1862)
= *Maera tenuimana*
Maeropsis
perrieri Chevreux, 1919
Melita
abyssorum Stephensen, 1944
bulla Karaman, 1978
coroninii Heller, 1866
dentata (Kroyer, 1842)
formosa Murdoch, 1866
gladiosa Bate, 1862
hergensis Reid, 1939
pallida Sars, 1879
palmata (Montagu, 1804)
= *Gammarus palmata*
quadrispinosa Vosseler, 1889
reidi Hamond, 1965
richardi Chevreux, 1900
valesi Karaman, 1955
virgula Krapp-Schickel, Ruffo & Schiecke, 1994
Psammogammarus
caecus Karaman, 1955
= *Chaetogammarus caecus*
gracilis Ruffo & Schiecke, 1976
Pseudoniphargus
adriaticus Karaman, 1955
africanus Chevreux, 1901
Family Melphidippidae

- Melphidippa*
borealis Boeck, 1871
goesi Stebbing, 1899
= *M. spinosa*
macrura Sars, 1894
Melphidippella
macra (Norman, 1869)
- Family Odiidae**
Odius
carinatus (Bate, 1862)
- Family Oedicerotidae**
Acanthostepheia
behrungiensis (Lockington, 1877)
incarinata Gurjanova, 1929
malmgreni (Goes, 1866)
- Aceroides*
goesi Just, 1980
latipes (Sars, 1883)
- Arrhinopsis*
longicornis Stappers, 1911
- Arrhis*
mediterraneus Ledoyer, 1983
phyllonyx (Sars, 1858)
= *Aceros phyllonyx* (Sars, 1858)
phyllonyx subsp. *arcticus* Bryazgin, 1974
- Bathymedon*
acutifrons Bonnier, 1896
= *Synchelidium acutifrons*?
banyulsensis Ledoyer, 1983
longimanus (Boeck, 1871)
monoculodiformis Ledoyer, 1983
obtusifrons (Hansen, 1883)
saussurei (Boeck, 1871)
- Halicreion*
aequicornis (Norman, 1869)
= *H. longicaudatus*
- Monoculodes*
acutipes Ledoyer, 1983
borealis Boeck, 1871
carinatus (Bate, 1857)
crassirostris Hansen, 1888
gibbosus Chevreux, 1888
griseus Della Valle, 1893
kroyeri Boeck, 1871
latimanus (Goes, 1866)
latissimanus Stephensen, 1931
longirostris (Goes, 1866)
norvegicus (Boeck, 1861)
packardi Boeck, 1871
pallidus Sars, 1892
rostratus Stephensen, 1931
schneideri Sars, 1895
simplex Hansen, 1888
subnudus Norman, 1889
= *M. falcatus* (G.O. Sars)
tenuirostratus Boeck, 1871
tesselatus Schneider, 1883
tuberculatus Boeck, 1871
- Monoculopsis*
longicornis (Boeck, 1871)
- Oedicerina*
ingolfi Stephensen, 1931
- Oedicerooides*
pilosa Ledoyer, 1983
proxima Bonnier, 1896
- Oediceropsis*
brevicornis (Lilljeborg, 1865)
proxima Chevreux, 1908
- Oediceros*
borealis (Boeck, 1871)
moigni Lagardere, 1968
saginatus Kroyer, 1842
- Paroediceros*
curvirostris (Hansen, 1888)
intermedius Stebbing, 1906
- lynceus* (Sars, 1858)
macrocheir (Sars, 1879)
propinquus (Goes, 1866)
- Perioculodes*
aequimanus (Korssman, 1880)
longimanus subsp. *angustipes* Ledoyer, 1983
longimanus subsp. *longimanus* (Bate & Westwood, 1868)
- Pontocrates*
altamarinus (Bate & Westwood, 1862)
- arcticus* Sars, 1895
= *P. norvegicus*
arenarius (Bate, 1858)
= *P. norvegicus*
- Synchelidium*
haplocheles (Grube, 1864)
= *Pseudocolodes brevicarpum*
= *S. brevicarpum*
intermedium Sars, 1892
= *S. haplocheles* Sars
longidigitatum Ruffo, 1947
maculatum Stebbing, 1906
tenuimanum Norman, 1895
- Westwoodilla*
acutifrons (Sars, 1892)
= *W. caecula* s. l.
brevicalcar Goes, 1866
caecula (Bate, 1857)
= *Halimedon parvimanus*
= *W. megalops* G.O. Sars, 1882
megalops (Sars, 1882)
= *W. caecula* s. l.
rectirostris (Della Valle, 1893)
- Family Pardaliscidae**
Arculia
trago subsp. *mediterranea* Karaman, 1986
- Halicre*
abyssi Boeck, 1871
aculeata Chevreux, 1912
walkeri (Ledoyer, 1973)
- Halicoedes*
anomalus Walker, 1893
tertia Stephensen, 1931
- Nicippe*
tumida Bruzelius, 1859
- Pardalisca*
abyssi Boeck, 1871
brachydactyla Bellan-Santini, 1985
cuspidata Kroyer, 1842
mediterranea Bellan-Santini, 1984
tenuipes Sars, 1893
- Pardalisella*
boecki (Malm, 1870)
- Pardaliscoides*
stebbingi Ledoyer, 1970
tenellus Stebbing, 1888
- Princaxelia*
stephensi Dahl, 1959
- Family Phlantidae**
Pereionotus
testudo (Montagu, 1808)
- Family Phoxocephalidae**
Eusyrophoxus
tenuipes Stephensen, 1925
- Harpinia*
abyssi Sars, 1879
agna Karaman, 1987
ala Karaman, 1987
amundseni Gurjanova, 1946
antennaria Meinert, 1890
= *H. neglecta* Sars
bidentata Stephensen, 1925
clivicola Watling, 1961
crenulata (Boeck, 1871)

- crenuloides* Stephensen, 1925
curtipes Stephensen, 1925
dellavallei Chevreux, 1910
excavata Chevreux, 1900
laevis Sars, 1891
latipes Norman, 1900
mucronata Sars, 1879
pectinata Sars, 1891
plumosa (Kroyer, 1842)
propinqua Sars, 1891
serrata Sars, 1878
truncata Sars, 1891
zavodniki Karaman, 1987
- Harpiniopsis*
similis Stephensen, 1925
- Joubinella*
ciliata Chevreux, 1908
- Leptophoxus*
falcatus (Sars, 1882)
- Metaphoxus*
fultoni (Scott, 1890)
= *Parametaphoxus fultoni*
= *Phoxocephalus fultoni*
gruneri Karaman, 1986
pectinatus (Walker, 1896)
= *Phoxocephalus pectinatus*
= *Phoxocephalus simplex* (Bate)
simplex (Bate, 1857)
- Paraphoxus*
oculatus (Sars, 1879)
- Phoxocephalus*
aquosus Karaman, 1985
holbotti (Kroyer, 1842)
- Pseudharpinia*
brevirostris Chevreux, 1920
excavata (Chevreux, 1887)
- Family Pleustidae**
- Cleonardopsis*
carinata Barnard, 1915
- Mesopleustes*
abyssorum Stebbing, 1888
- Neopleustes*
pulchellus (Kroyer, 1846)
= *Paramphithoe pulchella* R.M. Bruzelius, 1859
= *Pleusymtes pulchella* (Sars, 1876)
- Parapleustes*
assimilis (Sars, 1882)
= *Paramphithoe assimilis* (Sars, 1882)
bicuspis (Kroyer, 1838)
= *Neopleustes bicuspis*
= *Neopleustes m.*
= *Paramphithoe bicupis* (Sars, 1893)
= *Paramphithoe monocuspis* (Sars, 1893)
gracilis (Buchholz, 1874)
= *Paramphithoe brevicornis* (Sars, 1882)
monocuspis (Sars, 1893)
= *Neopleustes monocuspis*
- Pleustes*
cataphractus (Stimpson, 1853)
panopla (Kroyer, 1838)
= *P. panoplus* (Kroyer, 1838)
= *P. tuberculata*
tuberculata Bate, 1858
- Pleustomesus*
medius (Goes, 1866)
- Pleusymtes*
glaber (Boeck, 1861)
= *Parapleustes gaber* (Boeck, 1861)
glabroides (Dunbar, 1954)
mediterraneus (Ledoyer, 1986)
megacheir (Walker, 1897)
= *Parapleustes megacheir*
pulchella (Sars, 1876)
- Stenopleustes*
eldingi Gurjanova, 1930
- latipes* (Sars, 1858)
= *Parapleustes latipes*
malmgreni (Boeck, 1871)
nodifer (Sars, 1882)
- Family Podoceridae**
- Dulichia*
falcata (Bate, 1857)
spinossissima Kroyer, 1845
tuberculata Boeck, 1870
= *D. curticauda* Boeck, 1870
- Dulichiopsis*
abyssi Stephensen, 1944
bispinis (Gurjanova, 1930)
macera (Sars, 1879)
nordlandica (Boeck, 1870)
= *Dulichia nordlandica* Boeck, 1870
- Dyopedos*
hirticornis (Sars, 1876)
knipowitschi (Gurjanova, 1933)
monacantha (Metzger, 1875)
normani (Sars, 1895)
orrectus (Bate, 1857)
= *Dulichia porrectus* Bate, 1857
spinosus (Stephensen, 1944)
- Laetmatophilus*
ledoyeri Ruffo, 1986
purus Stebbing, 1888
tuberculatus Bruzelius, 1859
= *L. armatus* (Norman, 1869)
- Paradulichia*
typica Boeck, 1870
- Parunciola*
seurati Chevreux, 1911
- Podocerus*
chelonophilus (Chevreux & Guerne, 1888)
cristatus Thomson, 1879
schiekei Ruffo, 1987
variegatus Leach, 1814
- Xenodice*
frauenfeldti Boeck, 1871
- Family Pontoporeidae**
- Bathyporeia*
elegans Watkin, 1938
gracilis Sars, 1891
guilliamsoniana (Bate, 1857)
= *B. norvegica*
leucophthalma Bellan-Santini, 1973
lindstromi Stebbing, 1906
megalops Chevreux, 1911
nana Toulmond, 1966
pelagica (Bate, 1856)
phaiophthalma Bellan-Santini, 1973
pilosa Lindström, 1855
pseudopelagica Bellan-Santini & Vader, 1988
sardoa Bellan-Santini & Vader, 1988
sarsi Watkin, 1938
= *B. pilosa* of Russian authors
= *B. robertsoni*
sophiae Bellan-Santini & Vader, 1988
sunnivae Bellan-Santini & Vader, 1988
tenuipes Meinert, 1877
- Pontoporeia*
affinis Lindström, 1855
= *Monoporeia affinis*
femorata Kroyer, 1842
- Priscillina*
armata Boeck, 1861
- Family Sebidae**
- Seba*
aloe Karaman, 1971
armata (Chevreux, 1889)
saunderii Stebbing, 1875
- Family Stegocephalidae**
- Andaniella*

- pectinata* Sars, 1882
Andaniexis
abyssi (Boeck, 1871)
 = *Andania abyssi* (Boeck, 1871)
lupus Berge & Vader, 1997
mimonectes Ruffo, 1975
Andaniopsis
nordlandica (Boeck, 1871)
Andaniotes
islandica (Thompson, 1882)
 = *A. corpulentus*
Euandania
gigantea (Stebbing, 1888)
Phippsia
romeri Schellenberg, 1925
gibbosa (Sars, 1882)
 = *Aspidopleurus gibbosus* Sars, 1966
Phippsiella
abyssicola Oldewig, 1959
minima Stephensen, 1925
pseudophippsia Bellan-Santini, 1985
similis (Sars, 1891)
 = *Stegocephalus similis* (Sars, 1895)
Stegocephalina
ingolfi Stephensen, 1925
Stegocephaloides
auratus (Sars, 1882)
christianiensis Boeck, 1871
wagini (Gurjanova, 1936)
Stegocephalopsis
ampulla (Phipps, 1774)
Stegocephalus
inflatus Kroyer, 1842
Family Stenothoidae
Hardametopa
carinata (Hansen, 1888)
nasuta (Boeck, 1871)
Mesometopa
neglecta (Hansen, 1888)
Metopa
abyssalis Stephensen, 1931
aequicornis Sars, 1879
affinis Boeck, 1871
alderi (Bate, 1857)
 = *M. spectabilis*
boeckii Sars, 1892
borealis Sars, 1882
bruzelii (Goes, 1866)
clypeata (Kroyer, 1842)
glacialis (Kroyer, 1842)
 = *M. cariana*
groenlandica Hansen, 1888
invalida Sars, 1892
latimana Hansen, 1888
leptocarpa Sars, 1882
longicornis Boeck, 1871
norvegica (Liljeborg, 1851)
 = *M. pollexiana*
palmata Sars, 1892
propinqua Sars, 1892
pusilla Sars, 1892
robusta Sars, 1892
sinuata Sars, 1892
solsbergi Schneider, 1884
spitzbergensis Bruggen, 1907
tenuimana Sars, 1895
wiesei Gurjanova, 1933
Metopella
longimana (Boeck, 1871)
neglecta (Hansen, 1888)
Microstenothe
ascidiae Pirlot, 1933
Parametopa
kervillei Chevreux, 1901
Proboloides
calcarata (Sars, 1882)
 = *Probolium calcaratum* Sars
grandimimum Bonnier, 1896
gregaria (Sars, 1882)
 = *Probolium gregarium* Sars
 = *P. gregarius* (Sars, 1892)
Stenothe
aequicornis Stephensen, 1931
antennulariae Della Valle, 1893
ascidiae (Pirlot, 1933)
bosphorana Sowinsky, 1898
brevicornis Sars, 1882
cavimana Chevreux, 1908
contieri Chevreux, 1908
crassicornis Walker, 1897
dolifusi Chevreux, 1887
edwardi Krapp Schickel, 1975
 = *S. cattai* Reid, 1951
elachista Krapp Schickel, 1975
elachistoides Myers & McGrath, 1980
gallensis Walker, 1904
macrophthalma Stephensen, 1931
mandragora Krapp-Schickel, 1996
marina (Bate, 1856)
megacheir (Boeck, 1871)
microps Sars, 1892
monoculoides (Montagu, 1815)
 = *Montagua monoculoides*
 = *S. brevicornis* Sars, 1882
pieropan Krapp-Schickel, 1996
richardi Chevreux, 1895
setosa Norman, 1900
tenella Sars, 1882
tergestina Nebeski, 1881
valida Dana, 1852
Stenula
nordmanni (Stephensen, 1931)
rubrovittata Sars, 1883
 = *Amphithopsis latipes* Boeck
 = *Stenothoides latipes*
Family Synopiidae
Austrosyrrhoe
fimbriatus (Stebbing & Robertson, 1891)
septentrionalis Stephensen, 1931
Bruzelia
tuberculata Sars, 1882
typica Boeck, 1871
Ileraustroe
ilergetes subsp. *ilergetes* (Barnard, 1964)
Pseudotiron
bouvieri Chevreux, 1895
Stephobruzelia
dentata (Stephensen, 1931)
Syrrhoe
affinis Chevreux, 1908
angulipes Ledoyer, 1977
crenulata Goës, 1866
Syrrhoites
barnardi Karaman, 1986
capricornia Bellan-Santini, 1984
cornuta Bellan-Santini, 1984
pusilla Enequist, 1949
serrata (Sars, 1879)
walkeri (Bonnier, 1896)
Tiron
spiniferus (Stimpson, 1853)
 = *T. acanthurus*
 = *T. spiniferum*
Family Talitridae
Arcitalitrus
dorrieni (Hunt, 1925)
Macarorchestia
remyi (Schellenberg, 1950)
Orchestia
aestuarensis Wildish, 1987

- cavimana* Heller, 1865
gammarellus (Pallas, 1766)
 = *O. littorea*
guernei Chevreux, 1889
kossugi Ruffo, 1949
mateusi Afonso, 1977
mediterranea Costa, 1853
 = *O. laevis*
micropalma Armanieu and Salvat, 1963
montagui Audouin, 1826
roffensis Wildish, 1969
 = *O. remyi*
 = *O. roffensis*
spinifera Mateus, 1962
stephensi Cecchini, 1928
- Platorchestia**
platensis Kroyer, 1845
 = *Orchestia platensis* Kroyer, 1845
- Talitrus**
saltator (Montagu, 1808)
 = *T. locusta* Sars, 1890
- Talorchestia**
brito Stebbing, 1891
deshayesi (Audouin, 1826)
 = *Orchestia deshayesi* Heller, 1866
 = *Orchoidea deshayesi*
pelecaniformis Bellan-Santini & Ruffo, 1986
ugolini Bellan-Santini & Ruffo, 1991
- Family Urothoidae**
- Urothoe*
brevicornis Bate, 1862
corsica Bellan-Santini, 1965
elegans (Bate, 1857)
 = *U. abbreviata* Sars, 1879
 = *U. norvegica* Boeck, 1861
grimaldii Chevreux, 1895
hesperiae Conradi, Lopez-Gonzalez & Bellan-Santini, 1995
intermedia Bellan-Santini & Ruffo, 1986
marina (Bate, 1857)
poseidonis Reibisch, 1905
 = *U. grimaldi* var. *inermis*
 = *U. inermis* Chevreux & Fage, 1925
poucheti Chevreux, 1888
pulchella (Costa, 1853)
- Family Valettiidae**
- Valettietta*
gracilis Lincoln & Thurston, 1983
punctata Bellan-Santini, 1985
- Valettiopsis*
macrodactyla Chevreux, 1909
- Suborder Hyperiidea**
- Family Brachyscelidae**
- Brachyscelus*
crusculum Bate, 1861
 = *B. mediterranea* G1142
- Family Cystisomatidae**
- Cystisoma*
fabricii Stebbing, 1888
latipes (Stephensen, 1918)
pellucida (Willemoes-Suhm, 1874)
 = *C. parkinsoni* Stebbing
spinosum (Fabricius, 1775)
- Family Dairellidae**
- Dairella*
latissima Bovallius, 1887
- Family Hyperiidae**
- Hyperia*
galba (Montagu, 1815)
 = *H. latreille* Milne-Edwards
medusarum (Muller, 1776)
spinigera Bovallius, 1889
- Hyperioidea*
- longipes* Chevreux, 1900
Hyperoche
medusarum (Kroyer, 1838)
Themisto
abyssorum Boeck, 1870
 = *Parathemisto oblivia* (Kroyer)
gaudichaudi (Guerin, 1825)
 = *Euthemisto bispinosa* Boeck, 1876
 = *Euthemisto compressa* (Goes, 1865)
 = *P. gracilipes* Norman, 1869
- Family Lanceolidae**
- Lanceola*
borealis Bate & Westwood, 1868
loveni Bovallius, 1885
 = *L. aestiva* Stebbing, 1888
pacifica Stebbing, 1888
sayana Bovallius, 1885
serrata Bovallius, 1885
- Family Mimonectidae**
- Mimonectes*
gaussi (Woltereck, 1904)
 = *M. fowleri* (Stebbing, 1904)
 = *Parascina fowleri*
loveni Bovallius, 1885
- Family Phronimidae**
- Phronima*
sedentaria (Forskal, 1775)
 = *P. atlantica* Guerin-Meneville, 1836
- Family Phrosinidae**
- Phrosina*
semilunata Risso, 1882
- Primno*
evansi Shearer, 1986
- Family Platyscelidae**
- Platyscelus*
ovooides (Risso, 1816)
- Family Pronoidae**
- Paralycea*
gracilis Claus, 1879
- Family Scinidae**
- Acanthoscina*
acanthodes (Stebbing, 1895)
- Scina*
borealis (Sars, 1882)
crassicornis (Fabricius, 1775)
marginata (Bovallius, 1885)
 = *S. lepisma* Chun, 1889
oedicarpus Stebbing, 1895
rattrayi Stebbing, 1895
spinosa Vosseler, 1901
 = *S. uncipes* Stebbing, 1904
submarginata Tattersall, 1906
tullbergi (Bovallius, 1885)
- Family Tryphanidae**
- Tryphana*
malmi Boeck, 1870
- Family Vibiliidae**
- Vibilia*
armata Bovallius, 1887
borealis Bate & Westwood, 1868
propinquua Stebbing, 1888
pyripes Bovallius, 1887
- Suborder Ingolfiellidea**
- Family Ingolfiellidae**
- Ingolfiella*
ischitana Schiecke, 1973
- Cumacea**
- Compiled by Les Watling

The list follows the classification of Bacescu (1988, 1992), for the most part, with modifications from Watling (1991a, 1991b) for the Leuconidae and Nannastacidae, respectively. Besides the distributional data in Bacescu (1988, 1992), lists from Brattegard & Holthe (1997) and Gerken & Watling (1999) were used. Most of the literature describing the European Atlantic species is listed in Gerken & Watling (1999).

The list includes species from the shallow seas, shelves, and upper slopes to about 600 m depth from Novaya Zemlya in the Arctic, Greenland in the northwest, south to western Africa and east to the Black Sea. Although the European Seas are among the best to be studied for cumaceans, there are still many new species to be described, especially from the outer shelves and upper slope habitats. Future lists will record many of these new species as well as those from the abyssal plains.

Dr T. Brattegard is thanked for his careful checking of an earlier version of this list.

References

- Bacescu, M. 1988. Cumacea I (Fam. Archaeocumatidae, Lampropidae, Bodotriidae, Leuconidae). *Crustaceorum Catalogus* Pars 7:1-173.
 Bacescu, M. 1992. Cumacea II (Fam. Nannastacidae, Diastylidae, Pseudocumatidae, Gynodiastylidae and Ceratocumatidae). *Crustaceorum Catalogus* Pars 8: 175-468.
 Brattegard, T. & Holthe, T. (eds). 1997. Distribution of marine, benthic macro-organisms in Norway. A tabulated catalogue. Preliminary edition. - Research Report for DN 1997-1. Directorate for Nature Management. 409 pp. ISBN 82-7072-251-0.
 Gerken, S. & Watling L. 1999. Cumacea (Crustacea) of the Faeroe Island region. *Fróskaparit* 47: 199-227.
 Watling, L. 1991a. Revision of the Cumacean Family Leuconidae. *Journal of Crustacean Biology* 11:569-582.
 Watling, L. 1991b. Revision and rediagnosis of some Nannastacidae (Crustacea, Cumacea). *Proceedings of the Biological Society of Washington* 104:751-757.

Order Cumacea

Family Bodotriidae

Subfamily Bodotriinae

<i>Bodotria</i>			
<i>armoricana</i> LeLoeuff & Intes, 1977			
<i>gibba</i> (Sars, 1878)	M		
<i>intermedia</i> LeLoeuff & Intes, 1977	M		
<i>pulchella</i> (Sars, 1878)	M		
<i>scorpioides</i> (Montagu, 1804)	M		
<i>Cyclaspis</i>	A M		
<i>longicaudata</i> Sars, 1865			
<i>Cyclaspoides</i>	A		
<i>sarsi</i> Bonnier, 1896			
<i>Eocuma</i>			
<i>calmani</i> Fage, 1928			
<i>dolfusi</i> Calman, 1907			
<i>ferox</i> (Fischer, 1872)	M		
<i>Iphinoe</i>			
<i>acutirostris</i> Ledoyer, 1965	M		
<i>adriatica</i> Bacescu, 1988	M		
<i>armata</i> Ledoyer, 1965	M		
<i>douniae</i> Ledoyer, 1965	M		
<i>elisae</i> Bacescu, 1950	M		
<i>hupferi</i> Zimmer, 1916			
<i>inermis</i> Sars, 1878	M		
<i>maculata</i> Ledoyer, 1965	M		
<i>rhodaniensis</i> Ledoyer, 1965	M		
<i>serrata</i> Norman, 1867	M		
<i>tenella</i> Sars, 1878	M		
<i>trispinosa</i> (Goodsir, 1843)	M		
<i>Subfamily Vaunthompsoniinae</i>			
<i>Bathycuma</i>			
<i>elongatum</i> Hansen, 1895		A	
<i>Cumopsis</i>			
<i>fagei</i> Bacescu, 1956			
<i>goodsir</i> (VanBeneden, 1861)		M	
<i>longipes</i> (Dohrn, 1869)		M	
<i>Vaunthompsonia</i>			
<i>cristata</i> Bate, 1858		M	
Family Ceratocumatidae			
<i>Ceratocuma</i>			
<i>horridum</i> Calman, 1905		A	
<i>Cimmerius</i>			
<i>reticulatus</i> Jones, 1973		A	
Family Diastylidae			
<i>Brachydiastylis</i>			
<i>resima</i> (Krøyer, 1846)		A	
<i>Diastylis</i>			
<i>boecki</i> Zimmer, 1930			
<i>bradyi</i> Norman, 1879			
<i>charcoti</i> Reyss, 1974		M	
<i>cornuta</i> (Boeck, 1864)		A	
<i>dolfusi</i> Fage, 1928			
<i>echinata</i> Bate, 1865		A	
<i>edwardsii</i> (Krøyer, 1841)		A	
<i>glabra</i> Zimmer, 1900		A	
<i>goodsiri</i> (Bell, 1855)		A	
<i>jonesi</i> Reyss, 1972		M	
<i>laevia</i> Norman, 1869		M	
<i>lepechini</i> Zimmer, 1926		A	
<i>lucifera</i> (Krøyer, 1837)		A	
<i>neapolitana</i> Sars, 1879		M	
<i>oxyrhyncha</i> Zimmer, 1926		A	
<i>polaris</i> Sars, 1871		A	
<i>rathkei</i> (Krøyer, 1841)		A	
<i>rostrata</i> Goodsir, 1885			
<i>rugosa</i> Sars, 1865		M	
<i>scorpioides</i> (Lepechin, 1780)		A	
<i>spinulosa</i> Heller, 1875		A	
<i>tumida</i> (Liljeborg, 1855)			
<i>Diastyloides</i>			
<i>bacescoi</i> Fage, 1940		M	
<i>biplicatus</i> (Sars, 1865)		M	
<i>carpinei</i> Bacescu, 1969		M	
<i>scaber</i> Hansen, 1920		A	
<i>serratus</i> (Sars, 1865)		M	
<i>Diastylopsis</i>			
<i>bosphorica</i> Bacescu, 1982		M	
<i>Ekleptostylis</i>			
<i>walkeri</i> (Calman, 1907)		M	
<i>Ekonodiastylis</i>			
<i>nimia</i> (Hansen, 1920)		A	
<i>robusta</i> Gerken, Watling & Klitgaard, 2000			
<i>Leptostylis</i>			
<i>ampullacea</i> (Liljeborg, 1855)		A	
<i>longicaudata</i> Bonnier, 1896			
<i>longimana</i> (Sars, 1865)		A	
<i>macrura</i> Sars, 1870		A M	
<i>producta</i> Norman, 1879			
<i>villosa</i> Sars, 1869		A	
<i>Makrokylindrus</i>			
<i>insignis</i> Sars, 1871		M	
<i>josephinae</i> (Sars, 1871)			
<i>longicaudatus</i> (Bonnier, 1896)		A	
<i>longipes</i> (Sars, 1871)		M	
<i>serricauda</i> (T. Scott, 1912)			
<i>tubulicauda</i> (Calman, 1905)		A	
<i>aegaeus</i> Reyss, 1974		M	
<i>dubius</i> (Bonnier, 1896)			
<i>spiniventris</i> Hansen, 1920			
<i>Vemakylindrus</i>			
<i>charcoti</i> (Reyss, 1974)		M	
<i>doryphora</i> (Fage, 1940)		M	
Family Lampropidae			
<i>Chalarostylis</i>			

<i>elegans</i> Norman, 1879			
<i>Hemilamprops</i>			
<i>assimilis</i> Sars, 1883	A		
<i>cristatus</i> (Sars, 1870)	A		
<i>normani</i> Bonnier, 1896			
<i>pellucidus</i> Zimmer, 1908			
<i>roseus</i> (Norman, 1863)			
<i>uniplicatus</i> (Sars, 1872)	A		
<i>Lamprops</i>			
<i>fasciatus</i> Sars, 1863	A		
<i>fuscatus</i> Sars, 1865	A		
<i>profundus</i> Reyss, 1978			
<i>Mesolamprops</i>			
<i>denticulatus</i> Ledoyer, 1983	M		
<i>Paralamprops</i>			
<i>orbicularis</i> (Calman, 1905)	A		
<i>Platysympus</i>			
<i>tricarinatus</i> Hansen, 1920			
<i>typicus</i> (Sars, 1870)	A M		
Family Leuconidae			
<i>Eudorella</i>			
<i>arctica</i> Hansen, 1920	A		
<i>emarginata</i> (Krøyer, 1846)	A		
<i>gottliebi</i> Bacescu, 1961	M		
<i>gracilis</i> Sars, 1871			
<i>groenlandica</i> Zimmer, 1926	A		
<i>hirsuta</i> (Sars, 1869)			
<i>hispida</i> Sars, 1871	A		
<i>intermedia</i> Hansen, 1920	A		
<i>nana</i> Sars, 1879	M		
<i>parvula</i> Hansen, 1920	A		
<i>pusilla</i> Sars, 1871	A		
<i>spitzbergensis</i> Zimmer, 1926	A		
<i>truncatula</i> (Bate, 1856)	A M		
<i>Eudorellopsis</i>			
<i>deformis</i> (Krøyer, 1846)	A		
<i>Leucon</i>			
<i>pallidus</i> Sars, 1865	A		
<i>macrorhinus</i> Fage, 1951	M		
<i>norrevangi</i> Watling & Gerken, 1999			
<i>tener</i> Hansen, 1920	A		
<i>longirostris</i> Sars, 1871	A M		
<i>tenuirostris</i> Sars, 1887	A		
<i>acutirostris</i> Sars, 1865	A		
<i>affinis</i> Fage, 1951	M		
<i>sulvus</i> Sars, 1865	A		
<i>mediterraneus</i> Sars, 1878	M		
<i>nasica</i> (Krøyer, 1841)	A		
<i>nasicoides</i> (Krøyer, 1841)	A		
<i>nathorsii</i> Ohlin, 1901	A		
<i>robustus</i> Hansen, 1920	A		
<i>serratus</i> Norman, 1879	A		
<i>siphonatus</i> Calman, 1905	A M		
<i>spinulosus</i> Hansen, 1920	A		
Family Nannastacidae			
<i>Campylaspis</i>			
<i>affinis</i> Sars, 1870			
<i>africana</i> Bacescu & Muradian, 1972	A		
<i>alba</i> Hansen, 1920			
<i>bonetti</i> Bacescu & Muradian, 1972			
<i>caperata</i> Jones, 1974	A		
<i>costata</i> (Sars, 1865)	A		
<i>glabra</i> Sars, 1878	A M		
<i>globosa</i> Hansen, 1920	A		
<i>horrida</i> Sars, 1870	A		
<i>horridoidea</i> Stephensen, 1915	M		
<i>intermedia</i> Hansen, 1920	A		
<i>jonesi</i> Bacescu & Muradian, 1972			
<i>laticarpa</i> Hansen, 1920			
<i>legendrei</i> Fage, 1951	M		
<i>macrophthalma</i> Sars, 1878	M		
<i>mauritanica</i> Bacescu & Muradian, 1972			
<i>nitens</i> Bonnier, 1896			
<i>rostrata</i> Calman, 1905	M		
<i>rubicunda</i> (Liljeborg, 1855)	A		
		<i>serratipes</i> Hansen, 1920	A
		<i>squamifera</i> Fage, 1929	A M
		<i>sulcata</i> Sars, 1870	M
		<i>undata</i> Sars, 1864	
		<i>verrucosa</i> Sars, 1866	M
		<i>vitrea</i> Calman, 1906	A M
		<i>Cumella</i>	
		<i>africana</i> Bacescu, 1972	
		<i>angustata</i> Jones, 1984	A
		<i>carinata</i> (Hansen, 1887)	A
		<i>decipiens</i> Jones, 1984	A
		<i>limicola</i> Sars, 1879	M
		<i>pygmaea</i> Sars, 1865	M
		<i>tarda</i> Hansen, 1920	
		<i>Cumellopsis</i>	
		<i>helgae</i> Calman, 1905	A
		<i>puritani</i> Calman, 1906	A M
		<i>Nannastacus</i>	
		<i>brevicaudatus</i> Calman, 1905	M
		<i>euxinicu</i> Bacescu, 1951	M
		<i>turcicus</i> Bacescu, 1982	M
		<i>unguiculatus</i> (Bate, 1859)	M
		<i>Procampylaspis</i>	
		<i>arguini</i> Bacescu & Muradian, 1972	
		<i>armata</i> Bonnier, 1896	A M
		<i>bacescoi</i> Reyss & Soyer, 1966	M
		<i>bituberculata</i> Hansen, 1920	
		<i>bonnieri</i> Calman, 1906	M
		<i>lutensis</i> Jones, 1984	A
		<i>macronyx</i> Hansen, 1920	
		<i>mediterranea</i> Ledoyer, 1988	M
		<i>thalassae</i> Bacescu & Muradian, 1972	A
		<i>Scherocumella</i>	
		<i>longirostris</i> (Sars, 1878)	M
		<i>Schizotrema</i>	
		<i>atlanticum</i> Bacescu & Muradian, 1972	
		<i>Styloptocuma</i>	
		<i>gracillimum</i> (Calman, 1905)	A
		Family Pseudocumatidae	
		<i>Fontainella</i>	
		<i>mediterranea</i> Bacescu & Muradian, 1978	M
		<i>Petalosarsia</i>	
		<i>declivis</i> (Sars, 1865)	A
		<i>Pseudocuma</i>	
		<i>ciliatum</i> Sars, 1879	M
		<i>gilsoni</i> Bacescu, 1950	
		<i>longicorne</i> (Bate, 1858)	M
		<i>simile</i> Sars, 1900	M
		<i>graciloides</i> Sars, 1894	M
		<i>laeve</i> Sars, 1914	M
		<i>Pterocuma</i>	
		<i>pectinatum</i> (Sowinsky, 1893)	M
		<i>rostratum</i> (Sars, 1894)	M
		<i>sowinskyi</i> (Sars, 1894)	M
		<i>Schizorhamphus</i>	
		<i>eudorelloides</i> (Sars, 1894)	M
		<i>scabriusculus</i> (Sars, 1868)	M
		<i>Strauchia</i>	
		<i>taurica</i> Czerniavsky, 1868	M

Tanaidacea

Compiled by Graham J. Bird

Relatively few comprehensive faunal lists of Tanaidacea exist and the compilation has necessarily drawn on many individual papers on taxonomy or benthic ecology, although Sieg (1983b) has been invaluable. Major published sources for the NE Atlantic included Hansen (1913), Holdich & Jones (1983a), Holdich & Bird (1985), and two series by Bacescu (eg. 1981a, 1982a, 1985) and Bird & Holdich (eg. 1988, 1989a, 1989c). Mediterranean and Black Sea records were more

disparate and the list may be less comprehensive than that for the Atlantic. In this context, an unpublished thesis that included many 'new' species from the Bay of Naples was *not* used (Schiecke 1973).

Geographic coverage conformed to that in Figure 1 with Hansen's species records for West Greenland and some by Monod and others from Mauritania being excluded, even though the chance of their being found in the study area is high. Deep-sea taxa are especially prevalent in the list but recourse to unpublished material available to the compiler was kept to the minimum (Holdich & Bird 1989).

Throughout, the listing of old and obsolete names (many of them basionyms) has been minimised but reference to the comprehensive and excellent bibliography (Sieg 1983b) will enable those interested to follow nomenclatural changes. Linked to this, the reference attached to each record is generally a more familiar, or later, publication than the original except where this is unique or particularly definitive.

The classification of the two suborders, Apseudomorpha and Tanaidomorpha, is problematical and contentious but, for the purpose of the ERMS list, the former are classified following Gutu (1996) and the latter by Sieg (1986b). An alternative classification would place the apseudomorphan genera *Carpoapseudes* Lang, 1968 and *Leviapseudes* Sieg, 1980 in the family Whiteleggidae Gutu, 1972 rather than as a subfamily within the Apseudidae Leach, 1814. Also, within the tanaidomorphan family Anarthruridae Lang, 1971 are subsumed the better known Leptognathiidae Sieg, 1976 and Agathotanaidae Lang, 1971, but not the typhlotanaids which stand as a separate family. It is probable that this complex will receive more attention in the next few years, with many species of *Leptognathia* Sars, 1882 and *Typhlotanais* Sars, 1882 being transferred to new or existing genera.

The compiler is grateful for useful and detailed revisionary comments from Dr. Modest Gutu, of the Muzeul de Istorie Natural "Grigore Antipa", Bucarest.

References

- Bacescu, M. 1939. Les premiers Tanaidacea recueillis dans les eaux Roumaines de la Mer Noire. *Ann. Sci. Univ. Iassy* 25 (2): 205-208.
- Bacescu, M. 1949. Données sur la faune carcinologique de la Mer Noire le long de la côte Bulgare. *Trav. St. Biol. Marine Varna* 14: 1-24.
- Bacescu, M. 1960. Citeva animale necunoscute inca in Marea Neagra si descriera unor Malacostracei noi (*Elaphognathia monodi* n.sp. si *Pontotanais borceai* n.g. n.sp.) provenind din apele pontice prebosforice. *Studii Cerc. Biol. anim* 12 (2): 107-124.
- Bacescu, M. 1961. Contribution à la connaissance des Tanaidacés de la Méditerranée orientale -- I. Les Apseudidae et Kalliapseudidae des côtes d'Israël. *Bulletin of the Research Council of Israel* 10B: 137-170.
- Bacescu, M. 1977. *Heterotanais longidactylus* n.sp. and *Synapseudes mediterraneus* n.sp., Tanaidacea new for the eastern Mediterranean fauna. *Revue Roumaine de Biologie - Série de Biologie Animale* 22 (1): 119-125.
- Bacescu, M. 1978. *Atlantapseudes nigrichela* n.g., n.sp., Tanaidacea nouveau capturé par le navire "Thalassa" dans les eaux Portugaises. *Cahiers de Biologie Marine* 19: 317-322.
- Bacescu, M. 1980a. *Anuropoda francispori*, genre nouveau et espèce nouvelle de Monokonophora (Crustacea, Tanaidacea) des eaux de la Méditerranée levantine. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 22: 381-384.
- Bacescu, M. 1980b. Contribution à la connaissance des Monokonophora (Crustacea, Tanaidacea) de la Mer Méditerranée; description de deux espèces nouvelles, *Apseudes sicilianus* sp.n et *A. miserai* sp.n. *Revue Roumaine de Biologie - Série de Biologie Animale* 25: 83-91.
- Bacescu, M. 1980c. Contribution to the knowledge of some Kalliapseudidae (Crustacea, Tanaidacea) from the NW of the Indian Ocean. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 22: 359-379.
- Bacescu, M. 1981a. Nouvelle contribution à la connaissance de la faune d'Apseudoidea Leach 1914 (Crustacea) des eaux profondes du N.E. Atlantique. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 23: 33-71.
- Bacescu, M. 1981b. *Carpoapseudes kudinovae* n.sp. in the French waters of the Atlantic. *Revue Roumaine de Biologie - Série de Biologie Animale* 26 (1): 9-14.
- Bacescu, M. 1982a. *Carpoapseudes laubieri* sp.n. et *C. curticarpus* sp.n. de l'Atlantique du N.E. (Bassin W.-Européen) et quelques détails nouveaux sur la valabilité du genre. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 24: 55-68.
- Bacescu, 1982b. Sur la position systématique de *Leviapseudes hansenii* (Lang, 1968) emend. Bacescu. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 24: 69-77.
- Bacescu, M. 1984. Considérations sur deux sous-espèces nouvelles d'*Apeudes* - *A. spinosus albidas* ssp.n. et *A. robustus erythraicus* ssp.n. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 25: 91-98.
- Bacescu, M. 1985. Apseudoidea (Crustacea, Tanaidacea). In *Peuplements profonds du Golfe de Gascogne*, eds. L. Laubier & C. Monniot, IFREMER: 435-440.
- Bacescu, M. & Carausu, A. 1947. *Apseudopsis ostroumovi* n.sp. dans la Mer Noire (morphologie, affinités phylogénétiques, écologie). *Bull. sect. Sc. Ac. Roum.* 29 (6): 366-383.
- Bacescu, M. & Gutu, M. 1971. Contribution à la connaissance du genre *Apseudes* de la Méditerranée. *Fageapsuedes* n.g. et *Tuberapseudes* n.sg. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 11: 59-70.
- Bamber, R.N. 1990. A new species of *Zeuxo* (Crustacea: Tanaidacea) from the French Atlantic coast. *Journal of Natural History* 24: 1587-1596.
- Bird, G.J. & Holdich, D.M. 1984. New deep-sea tanaids (Crustacea, Tanaidacea) from the North-east Atlantic. *Zoologica Scripta* 13 (4): 285-315.
- Bird, G.J. & Holdich, D.M. 1985. A remarkable tubicolous tanaid (Crustacea: Tanaidacea) from the Rockall Trough. *Journal of the Marine Biological Association of the United Kingdom* 65: 563-572.
- Bird, G.J. & Holdich, D.M. 1988. Deep-sea Tanaidacea (Crustacea) of the North-east Atlantic: the tribe Agathotanini. *Journal of Natural History* 22: 1591-1621.
- Bird, G.J. & Holdich, D.M. 1989a. Deep-sea Tanaidacea (Crustacea) of the North-east Atlantic: the genus *Paranarthrura* Hansen. *Journal of Natural History* 23: 137-187.
- Bird, G.J. & Holdich, D.M. 1989b. Recolonisation of artificial sediments in the deep Bay of Biscay by tanaidaceans (Crustacea: Peracarida), with a description of a new species of *Pseudotanais*. *Journal of the Marine Biological Association of the United Kingdom* 69: 307-317.
- Bird, G.J. & Holdich, D.M. 1989c. Deep-sea Tanaidacea (Crustacea) of the North-east Atlantic: the subfamily Pseudotanainae (Pseudotanaidae) and the family Nototanidae. *Zoological Journal of the Linnean Society* 97: 233-298.
- Carpine, C. 1970. Ecologie de l'étage bathyal dans la Méditerranée occidentale. *Mémoires de l'Institut océanographique*, Monaco 2, 146pp.
- Dojiri & Sieg, J. 1997. The Tanaidacea. In *Taxonomic Atlas of the Benthic Fauna of the Santa Maria Basin and Western Santa Barbara Channel. Volume 11. The Crustacea Part 2 - The Isopoda, Cumacea and Tanaidacea*. Santa Barbara Museum of Natural History: 181-268.

- Dollfus, A. 1897. Note préliminaire sur les Tanaidæ recueillis aux Açores pendant les Campagnes de l'Hirondelle (1877-1888). *Bull. Soc. Zool. Fr.* 22: 207-215.
- Dollfus, A. 1898. Campagnes de la Melita. Tanaidæ recoltés par M. Ed. Chevreux dans l'Atlantique et dans la Méditerranée. *Mém. Soc. Zool. Fr.* 11: 35-47.
- Gardiner, L.F. 1975. The systematics, postmarsupial development and ecology of the deep-sea Family Neotanaidæ (Crustacea: Tanaidacea). *Smithsonian Contributions to Zoology* 170, 265pp.
- Greve, L. 1964. The records of *Leptognathia dentifera* G.O. Sars (Tanaidacea). *Sarsia* 15: 71.
- Gutu, M. 1986. Description of *Apseudes olimpiae* n.sp. and of *Tanabnormia cornicauda* n.g., n.sp. (Crustacea, Tanaidacea). *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 28: 37-48.
- Gutu, M. 1989. La description de *Calozodion dollfusi*, espèce nouvelle de tanaïdacs (Crustacea) provenant des eaux européennes de l'Atlantique. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 30: 129-133.
- Gutu, M. 1996. The synoptic table and key to superspecific taxa of recent Apseudomorpha (Crustacea, Tanaidacea). *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 36: 135-1456.
- Gutu, M. & Illiffe, T.M. 1998. *Sphyrapoides bicornis*, new genus and new species from a submarine escarpment in the Bahamas, and some remarks on Sphyrapinae. *Travaux du Muséum d'Histoire naturelle "Grigore Antipa"* 40: 211-225.
- Hansen, H.J. 1909. Revidered fortægnelse over Danmarks marine Arter af Isopoda, Tanaidacea, Cumacea, Mysidacea of Euphausiacea. *Vidensk. Meddr. Naturh. Foren.* 1: 197-263 [226-233].
- Hansen, H.J. 1913. Crustacea Malacostraca II, IV, the Order Tanaidacea. *Danish Ingolf Expedition* 3 (3): 1-145.
- Holdich, D.M. & Bird, G.J. 1985. A preliminary report on 'dikonophoran' tanaids (Crustacea). In *Peuplements profonds du Golfe de Gascogne*, eds. L. Laubier & C. Monniot, IFREMER: 441-447.
- Holdich, D.M. & Bird, G.J. 1986. Tanaidacea (Crustacea) from sublittoral waters off West Scotland, including the description of two new genera. *Journal of Natural History* 20: 79-100.
- Holdich, D.M. & Bird, G.J. 1989. The systematics, distribution and habitat preferences of sublittoral and bathybenthic Tanaidacea in the North-east Atlantic. Unpublished report to Natural Environment Research Council, GR3/4983 A.
- Holdich, D.M. & Jones, J.A. 1983a. *Tanaids*. Synopsis of the British Fauna 27, Cambridge University Press, 98pp.
- Holdich, D.M. & Jones, J.A. 1983b. The distribution and ecology of British shallow-water tanaid crustaceans (Peracarida, Tanaidacea). *Journal of Natural History* 17: 157-183.
- Johnson, S.B. & Attramadal, Y.G. 1982a. Reproductive behaviour and larval development of *Tanaidæ cavolinii* (Crustacea, Tanaidacea). *Marine Biology* 71: 11-16.
- Johnson, S.B. & Attramadal, Y.G. 1982b. A functional-morphological model of *Tanaidæ cavolinii* Milne-Edwards (Crustacea, Tanaidacea) adapted to a tubicolous life-strategy. *Sarsia* 67 (1): 29-42.
- Jones, J.A. & Holdich, D.M. 1983. A new species of the tanaid genus *Tanaissus* Norman & Scott from British waters. *Crustaceana* 45 (2): 214-219.
- Kudinova-Pasternak, R.K. 1982. Deep-sea Tanaidacea (Crustacea, Malacostraca) from the Mediterranean Sea. *Trudy Instituta okeanologii. Akademija nauk SSSR* 117: 151-162.
- Kudinova-Pasternak, R.K. 1983. The abyssal Tanaidacea (Crustacea) of the Iberian and West-European hollows of the Atlantic Ocean. *Zool. Zhur.* 62 (8): 1170-1176.
- Lang, K. 1968. Deep-sea Tanaidacea. *Galathea Report* 9: 23-209.
- Lang, K. 1973. Taxonomische und phylogenetische Untersuchungen über die Tanaidaceen (Crustacea). 8. *Zoologica Scripta* 2: 197-229.
- Larwood, H.J. 1954. Crustacea Tanaidacea and Isopoda from the Suez Canal. *Annals Mag. Nat. Hist. London*, Series 12, 7: 561-577.
- Monod, T. 1923. Notes Isopodologiques. I. Sur un *Heterotanais*. *Bulletin de la Société Zoologique de France* 48: 506-511.
- Norman, A.M. & Stebbing, T.R.R. 1886. On the Crustacea Isopoda of the 'Lightning', 'Porcupine' and 'Valorous' expeditions. *Transactions of the Zoological Society of London* 12: 77-141.
- Richardson, H. 1911. Les Crustaces Isopodes du Travailleur et du Talisman; formes nouvelles. *Bull. Mus. Natn. Hist. Nat.*, Paris 7: 1-17.
- Riggio, S. 1973. On the presence of the genus *Synapseudes* Miller, 1940 (Crustacea Peracarida Anisopoda), in the Mediterranean, with a preliminary description of *Synapseudes shiinoi* n.sp. *Memorie di Biologia Marina e di Oceanografia* N.S. 3 (1): 11-19.
- Riggio, S. 1976. Dati tassonomici sugli Anisopoda (Crustacea Peracarida) del Porto di Palermo. *Archo Oceanogr. Limnol.* 18, suppl. 3: 501-506.
- Riggio, S. 1977. *Synapseudes shiinoi* Riggio, 1973, a species of Tanaidacea found in the Mediterranean. *Crustaceana* 33 (2): 153-162.
- Sars, G.O. 1886. Nye bidrag til Kundskaben om Middelhavets Invertebratfauna, III: Middelhavets Saxispoder (Isopoda Chelifera). *Arch. Math. Naturv.* 11: 263-368.
- Sars, G.O. 1896. Isopoda. *An Account of the Crustacea of Norway* 2: 1-270.
- Schiecke, U. 1973. *Ein Beitrag zur Kenntnis der Systematik, Biologie und Autokologie Mariner Peracarida (Amphipoda, Isopoda, Tanaidacea) des Golfs von Neapel*. Unpublished thesis, Universität Kiel, 408 pp.
- Sieg, J. 1980. Taxonomische Monographie der Tanaidæ Dana, 1849 (Crustacea, Tanaidacea). *Abh. Senckenb. Naturforsch. Ges.* 537: 1-267.
- Sieg, J. 1983a. Erstnachweis von *Exspina typica* Lang für den Atlantischen Ozean (Tanaidacea). *Crustaceana* 36 (2): 189-190.
- Sieg, J. 1983b. *Tanaidacea. Crustaceorum Catalogus, Pars 6*, Dr. W. Junk Publishers, The Hague, 552pp.
- Sieg, 1986a. *Crustacea Tanaidacea of the Antarctic and the Subantarctic. I. On material collected at Tierra del Fuego, Isla de los Estados, and the West Coast of the Antarctic Peninsula*. Antarctic Research Series Vol.45, Biology of the Antarctic Seas XVIII, 180 pp.
- Sieg, 1986b. *Tanaidacea (Crustacea) von der Antarktis uns Subantarktis. II. Tanaidacea gesammelt von Dr. J.W. Wagele während der deutschen Antarktis Expedition, 1983*. Mitteilungen aus dem Zoologischen Museum der Universität Kiel II (4): 1-80.
- Sieg, J. & Bird, G.J. 1989. Remarks on the genus *Mesotanais* Dollfus, 1897 (Crustacea, Tanaidacea). Redescription of the type species and description of *M. elongatus* sp.nov. *Bull. Mus. Natn. Hist. Nat.*, Paris 11, sect A. 1: 165-182.
- Sieg, J. & Dojiri, M. 1989. Remarks on *Araphura* Bird & Holdich (Crustacea, Tanaidacea) and allied genera, including descriptions of three new species. *Zoologica Scripta* 18 (1): 115-137.
- Smith, G. 1906. High and low dimorphism. With an account of certain tanaidæ of the Bay of Naples. *Mitt. Zool. Stn Neapel* 17: 312-340.
- Stephensen, K. 1915. Isopoda, Tanaidacea, Cumacea, Amphipoda (excl. Hyperiidae). In *Report on the Danish Oceanographical Expeditions 1908-10 to the Mediterranean and adjacent seas* Vol.2, Biology: 26-29.
- Tattersall, W.M. 1905. The marine fauna of the coast of Ireland. Pt.V. Isopoda. *Fisheries Ireland. Scientific Investigations*, 1904, II: 1-90.
- Tattersall, W.M. 1925. New Tanaidacea and Isopoda from the West Coast of Africa. *Bull. Soc. Sc. Nat. Maroc.* 5 (3): 77-85.
- Willemoes-Suhm, R. 1879. On some Atlantic Crustacea from the Challenger Expedition. *Transactions of the Linnean Society of London*, Series 2, Zoology 1: 23-59.

Order Tanaidacea		
Suborder Apseudomorpha		
Family Apseudidae		
Subfamily Apseudinae		
<i>Apseudes</i>		
<i>acutifrons</i> G.O. Sars, 1882	M	
= <i>A. ostroumovi</i> (Bacescu & Carausu, 1947)		
= <i>Apseudopsis ostroumovi</i>		
<i>africanus</i> Tattersall, 1925	M	
<i>africanus</i> subsp. <i>orientalis</i> Bacescu, 1961	M	
<i>caecus</i> Willemoes-Suhm, 1879	A	
<i>elisae</i> Bacescu, 1961	M	
<i>graciloides</i> Stephensen, 1915	A M	
<i>graziela</i> Bacescu, 1981	A	
<i>grossimanus</i> Norman & Stebbing, 1886	A M	
<i>hastifrons</i> Norman & Stebbing, 1886		
= <i>A. ostroumovi</i> (Bacescu & Carausu, 1947)		
= <i>Apseudopsis ostroumovi</i>		
<i>holthuisi</i> Bacescu, 1962	M	
<i>intermedius</i> Hansen, 1985	A M	
<i>latreillii</i> (Milne-Edwards, 1828)	A M	
= <i>A. latreillei</i>		
= <i>A. latreillii</i> subsp. <i>mediterraneus</i> Bacescu, 1961		
<i>miserai</i> Bacescu, 1980	M	
<i>robustus</i> G.O. Sars, 1882	M	
<i>robustus</i> subsp. <i>israeliticus</i> Bacescu, 1961	M	
<i>rotundifrons</i> Bacescu, 1981	A	
<i>setiferus</i> Bacescu, 1981	A	
<i>sicilianus</i> Bacescu, 1980	M	
<i>spinosus</i> (M. Sars, 1858)	A	
= <i>A. koehleri</i> Bonnier, 1896		
= <i>A. spinosus</i> subsp. <i>albidus</i> Bacescu, 1984		
= <i>A. spinosus</i> subsp. <i>koehleri</i> Bacescu, 1981		
<i>talpa</i> (Montagu, 1808)	A	
= <i>A. hibernicus</i> Walker, 1898		
<i>tenuis</i> Hansen, 1913	A	
<i>tenuimanus</i> G.O. Sars, 1882	M	
<i>tenuimanus</i> subsp. <i>gottliebi</i> Bacescu, 1961	M	
<i>uncidigitatus</i> Norman & Stebbing, 1886	M	
<i>vicinus</i> Hansen, 1913	A	
<i>Atlantapseudes</i>		
<i>nigrichela</i> Bacescu, 1978		
<i>Fageapseudes</i>		
<i>retusifrons</i> (Richardson, 1912)	A M	
= <i>Apseudes obtusifrons</i> Norman & Stebbing, 1886		
= <i>Apseudes retusifrons</i>		
<i>Tuberapseudes</i>		
<i>echinatus</i> (G.O. Sars, 1882)	M	
= <i>Apseudes echinatus</i>		
= <i>Apseudes lunaris</i> Norman & Stebbing, 1886		
Subfamily Leviapseudinae		
<i>Carpapseudes</i>		
<i>curticarpus</i> Bacescu, 1982	A	
<i>kudinovae</i> Bacescu, 1981	A	
<i>laubieri</i> Bacescu, 1982	A	
<i>serratispinosus</i> Lang, 1968	A	
<i>simplicirostris</i> (Norman & Stebbing, 1886)	A	
= <i>Apseudes simplicirostris</i>		
<i>Leviapseudes</i>		
<i>gracilis</i> (Norman & Stebbing, 1886)	A	
= <i>Apseudes gracilis</i>		
= <i>Leiopus gracilis</i>		
<i>gracillimus</i> (Hansen, 1913)	A	
= <i>Leiopus gracillimus</i>		
<i>hanseni</i> (Lang, 1968)	A	
= <i>Leiopus hansenii</i>		
<i>leptodactylus</i> (Beddard, 1886)	A	
= <i>Apseudes leptodactylus</i>		
= <i>Leiopus leptodactylus</i>		
<i>longissimus</i> Bacescu, 1982	A	
<i>segonzaci</i> Bacescu, 1981	A	
<i>zenkevitchioides</i> Bacescu, 1981	A	
Family Kalliapseudidae		
Subfamily Kalliapseudinae		
<i>Cristapseudes</i>		
<i>omercooperi</i> (Larwood, 1954)		M
= <i>Kalliapseudes omer-cooperi</i>		
Family Metapseudidae		
Subfamily Metapseudinae		
<i>Calozodion</i>		
<i>dollfusi</i> Gutu, 1989		A
Subfamily Synapseudinae		
<i>Synapseudes</i>		
<i>mediterraneus</i> Bacescu, 1977		M
Family Parapseudidae		
<i>Parapseudes</i>		
<i>latifrons</i>		
<i>Anuropoda</i>		
<i>francispora</i> Bacescu, 1980		M
<i>Parapseudes</i>		
<i>latifrons</i> (Grube, 1864)		A M
Family Sphyrapidae		
Subfamily Pseudosphyrapinae		
<i>Francapseudes</i>		
<i>debilis</i> Bacescu, 1981		A
<i>uniarticulatus</i> Bacescu, 1981		A
<i>Pseudosphyrapus</i>		
<i>anomalus</i> (G.O. Sars, 1869)		A
= <i>Sphyrapus anomalus</i>		
<i>centobi</i> Bacescu, 1981		A
<i>serratus</i> (G.O. Sars, 1882)		A
= <i>Sphyrapus serratus</i>		
<i>Sphyrapus</i>		
<i>malleolus</i> Norman & Stebbing, 1886		A M
Subfamily Spbyrapinae		
<i>Sphyrapoides</i>		
<i>tudes</i> Norman & Stebbing, 1886		A
Suborder Neotanaidomorpha		
Family Neotanaidae		
<i>Neotanais</i>		
<i>affinis</i> Wolff, 1956		A
<i>americanus</i> Beddard, 1886		A
= <i>Alao tanais serratispinosus</i>		
= <i>N. serratispinosus</i> (Norman & Stebbing, 1886)		
<i>hadalis</i> Wolff, 1956		A
<i>hastiger</i> (Norman & Stebbing, 1886)		A
= <i>Alao tanais hastiger</i>		
= <i>N. Edwardsi</i> Dollfus, 1898		
<i>laevipinosus</i> (Norman & Stebbing, 1886)		A
= <i>Alao tanais laevipinosus</i>		
= <i>N. Edwardsi</i> Dollfus, 1898		A
<i>micromopher</i> Gardiner, 1975		A
<i>peculiaris</i> Lang, 1968		A
<i>sandersi</i> Gardiner, 1975		A
Suborder Tanaidomorpha		
Family Tanaidae		
Subfamily Pancolinae		
Tribe Pancolini		
<i>Hexapleomera</i>		
<i>robusta</i> (Moore, 1894)		M
= <i>Tanais robustus</i>		
= <i>Tanais testudinicola</i> Dollfus, 1898		
Subfamily Sinelobinae		
<i>Parasinelobus</i>		
<i>chevreuxi</i> (Dollfus, 1898)		A
= <i>Tanais chevreuxi</i>		
Subfamily Tanainae		
<i>Tanais</i>		
<i>dulongii</i> (Audouin, 1826)		A M
= <i>T. cavolini</i>		
<i>grimaldii</i> Dollfus, 1897		A M
Subfamily Pancolinae		
Tribe Anatanaini		

<i>Zeuxo</i>		<i>Arhaphuroides</i>	
<i>holdichi</i> Bamber, 1990	A	<i>brevispina</i> (Bird & Holdich, 1984)	A
Family Leptocheiliidae		= <i>Araphura brevispina</i>	
<i>Heterotanais</i>		<i>Araphura</i>	
<i>oerstedi</i> (Kroyer, 1842)	A M	<i>filiformis</i> (Lilljeborg, 1864)	A M
= <i>H. gurneyi</i> Norman, 1906		= <i>Leptognathia filiformis</i>	
= <i>H. oerstedii</i>		<i>Collettea</i>	
<i>Leptochelia</i>		<i>cylindrata</i> (G.O. Sars, 1882)	A M
<i>savignyi</i> (Kroyer, 1842)	A M	= <i>Strongylura cylindrata</i>	
= <i>L. algicola</i> Harger, 1878		<i>minima</i> (Hansen, 1913)	A
= <i>L. corsica</i> Dollfus, 1898		= <i>Strongylura minima</i>	
= <i>L. dubia</i> Kroyer, 1842		<i>rotundicauda</i> Kudinova-Pasternak, 1983	A
= <i>L. Edwardsii</i> (Kroyer, 1842)		<i>Filitanais</i>	
= <i>L. neapolitana</i> G.O. Sars, 1882		<i>rebainsi</i> Kudinova-Pasternak, 1975	
<i>Mesotanais</i>		<i>Haplocope</i>	
<i>dubius</i> Dollfus, 1897	A	<i>abyssorum</i> Dollfus, 1897	A
<i>elongatus</i> Sieg & Bird, 1989	A	<i>angusta</i> G.O. Sars, 1882	A
Pseudoleptochelia		<i>linearis</i> Hansen, 1913	A
<i>anomala</i> (G.O. Sars, 1882)	M	<i>Leptognathiella</i>	
= <i>Heterotanais algiricus</i> Dollfus, 1898		<i>abyssi</i> (Hansen, 1913)	A
= <i>Heterotanais anomalus</i>		= <i>Leptognathia abyssi</i>	
<i>longydactylus</i> (Bacescu, 1977)	M	<i>clivicola</i> Bird & Holdich, 1984	A
= <i>Heterotanais longidactylus</i>		<i>spinicauda</i> Bird & Holdich, 1984	A
<i>magna</i> (Smith, 1906)	M	<i>subaequalis</i> (Hansen, 1913)	A
= <i>Heterotanais magnus</i>		= <i>Leptognathia subaequalis</i>	
<i>mercantilis</i> (Smith, 1906)	M	<i>Leptognathoides</i>	
= <i>Leptochelia mercantilis</i>		<i>polita</i> (Hansen, 1913)	A
<i>merginellae</i> (Smith, 1906)	M	= <i>Leptognathia polita</i>	
= <i>Leptochelia marginellae</i>		<i>potens</i> Bird & Holdich, 1984	A
<i>provincialis</i> (Dollfus, 1898)		<i>Leptognathiopsis</i>	
= <i>Heterotanais provincialis</i>		<i>antennuata</i> Holdich & Bird, 1986	A
Family Paratanaidae		<i>Macrinella</i>	
<i>Paratanais</i>		<i>clavipes</i> (Hansen, 1913)	
<i>atlanticus</i> Dollfus, 1897	A	<i>Robustochelia</i>	
Family Anarthuridae		<i>longa</i> Kudinova-Pasternak, 1983	A
Subfamily Anarthurinae		<i>Stenotanais</i>	
Tribe Agathotanaini		<i>crassiseta</i> Bird & Holdich, 1984	A
<i>Agathotanais</i>		<i>hamicauda</i> Bird & Holdich, 1984	A
<i>ingolfi</i> Hansen, 1913	A	<i>Strongylurella</i>	
<i>Metagathotanais</i>		<i>indivisa</i> Hansen, 1913	
<i>insulcatus</i> Bird & Holdich, 1988	A	<i>Subulella</i>	
<i>Paragathotanais</i>		<i>scotti</i> Holdich & Bird, 1986	A
<i>gracilis</i> Bird & Holdich, 1988	A	<i>Tanabnormia</i>	
<i>nanus</i> Bird & Holdich, 1988	A	<i>cornicauda</i> Gutu, 1986	A
<i>robustus</i> Bird & Holdich, 1988	A	<i>Tanaella</i>	
= <i>P. typicus</i> Lang, 1971		<i>ochracea</i> Hansen, 1913	A
<i>Paranarthrura</i>		= <i>Leptognathia ochracea</i>	
<i>borealis</i> Bird & Holdich, 1989	A	<i>unguicillata</i> Norman & Stebbing, 1886	A M
<i>crassa</i> Bird & Holdich, 1989	A	= <i>Leptognathia unguicillata</i>	
<i>insignis</i> Hansen, 1913	A	Subfamily Leptognathiinae	
<i>intermedia</i> Kudinova-Pasternak, 1982	A M	<i>Exspina</i>	
= <i>P. similis</i> Lang, 1970		<i>typica</i> Lang, 1968	A
<i>lusitanus</i> Bird & Holdich, 1989	A	<i>Leptognathia</i>	
<i>subtilis</i> Hansen, 1913	A M	<i>acanthifera</i> Hansen, 1913	A
<i>tridens</i> Bird & Holdich, 1989	A	<i>alba</i> Hansen, 1913	A
Tribe Anarthurini		<i>arctophylax</i> (Norman & Stebbing, 1886)	A
<i>Anarthrura</i>		= <i>Cryptocoope arctophylax</i>	
<i>simplex</i> G.O. Sars, 1882	A M	<i>armata</i> Hansen, 1913	A
<i>Leptognathia</i>		= <i>L. Amdrupii</i> Hansen, 1913	
<i>latiremis</i> Hansen, 1913	A	<i>brachiata</i> Hansen, 1913	A
Subfamily Akanthophoreinae		<i>breviremis</i> (Lilljeborg, 1864)	A
<i>Akanthophoreus</i>		<i>crassa</i> Hansen, 1913	A
<i>gracilis</i> (Kroyer, 1842)	A	<i>hastata</i> Hansen, 1913	A
= <i>Leptognathia gracilis</i>		<i>laticaudata</i>	
= <i>Leptognathia hansenii</i> Vanhoffen, 1907		= <i>Tanaopsis laticaudata</i>	
= <i>Leptognathia sarsi</i> Hansen, 1909		<i>longiremis</i>	
<i>inermis</i> (Hansen, 1913)	A	<i>manca</i> G.O. Sars, 1882	A M
<i>longiremis</i> (Lilljeborg, 1864)	A M	<i>paramanca</i> Lang, 1958	A
<i>multiserratus</i> (Hansen, 1913)	A	<i>rigida</i> (Bate & Westwood, 1868)	A
<i>Araphura</i>		<i>tenella</i> Hansen, 1913	A
<i>brevimanus</i> (Lilljeborg, 1864)	A M	<i>uncinata</i> Hansen, 1913	A
= <i>Leptognathia brevimanus</i>		<i>ventralis</i> Hansen, 1913	A
= <i>Leptognathia brevimanus</i>		<i>voeringi</i> (G.O. Sars, 1877)	A
		= <i>Cryptocoope Voringii</i>	
		= <i>L. Voringii</i>	

<i>zezinae</i> Kudinova-Pasternak, 1973	A	<i>jonesi</i> Sieg, 1977	A
<i>Mirandotanais</i>	A	<i>lilljeborgi</i> G.O. Sars, 1882	A
<i>vorax</i> Kussakin & Tsareva, 1974	A	<i>longispinus</i> Bird & Holdich, 1989	A
<i>Nematotanais</i>	A	<i>macrocheles</i> G.O. Sars, 1882	A M
<i>mirabilis</i> Bird & Holdich, 1985	A	<i>mediterraneus</i> G.O. Sars, 1882	A M
<i>Pseudoparatanais</i>	A M	= <i>P. mediterraneus</i>	
<i>batei</i> (G.O. Sars, 1882)	A M	<i>scalpellum</i> Bird & Holdich, 1989	A
= <i>Paratanais batei</i>	A M	<i>spatula</i> Bird & Holdich, 1989	A
<i>Tanaopsis</i>	A M	<i>spicatus</i> Bird & Holdich, 1989	A
<i>graciloides</i> (Lilljeborg, 1864)	A M	<i>vulsella</i> Bird & Holdich, 1989	A
= <i>T. laticaudata</i>	A M		
Family Typhlotanaidae		Family Nototanaidae	
<i>Meromonakantha</i>	A	<i>Bathytanaissus</i>	
<i>macrocephala</i> (Hansen, 1913)	A	<i>spinulosus</i> Bird & Holdich, 1989	A
= <i>Typhlotanais macrocephalus</i>	A	<i>Tanaissus</i>	
<i>Peraeospinosus</i>	A	<i>elongatus</i> Jones & Holdich, 1983	A
<i>mixtus</i> (Hansen, 1913)	A	<i>lilljeborgi</i> (Stebbing, 1891)	A
= <i>Typhlotanais mixtus</i>	A	= <i>Leptognathia crassimana</i> Dollfus, 1898	
<i>Typhlotanais</i>	A M		
<i>aequiremis</i> (Lilljeborg, 1864)	A		
<i>assimilis</i> G.O. Sars, 1882	A		
<i>brevicornis</i> (Lilljeborg, 1864)	A		
<i>cornutus</i> G.O. Sars, 1879	A		
<i>eximus</i> Hansen, 1913	A		
<i>finmarchicus</i> G.O. Sars, 1882	A		
<i>gracilipes</i> Hansen, 1913	A		
<i>grandis</i> Hansen, 1913	A		
<i>inaequipes</i> Hansen, 1913	A		
<i>inermis</i> Hansen, 1913	A		
<i>irregularis</i> Hansen, 1913	A		
<i>kerguelensis</i> Beddard, 1886	A		
<i>longimanus</i> Dollfus, 1897	A		
<i>messinensis</i> G.O. Sars, 1882	A M		
<i>microcheles</i> G.O. Sars, 1882	A		
<i>mucronatus</i> Hansen, 1913	A		
<i>penicillatus</i> G.O. Sars, 1882	A		
<i>plebejus</i> Hansen, 1913	A		
<i>proctagon</i> Tattersall, 1904	A		
<i>profundus</i> Hansen, 1913	A		
<i>pulcher</i> Hansen, 1913	A		
<i>richardi</i> Dollfus, 1897	A		
<i>solidus</i> Hansen, 1913	A		
<i>spinicauda</i> Hansen, 1913	A		
<i>spinipes</i> Kudinova-Pasternak, 1982	M		
<i>spiniventris</i> Dollfus, 1897	A		
<i>tenuicornis</i> G.O. Sars, 1882	A		
<i>tenuimanus</i> (Lilljeborg, 1864)	A		
= <i>T. tenuimanus</i>	A		
<i>trispinosus</i> Hansen, 1913	A		
<i>variabilis</i> Hansen, 1913	A		
Family Pseudotanaidae		Isopoda	
Subfamily Cryptocopinae		Compiled by Jacob van der Land	
<i>Cryptocope</i>		The ERMS-list of Isopoda is mainly based on the "World List of Marine and Freshwater Crustacea Isopoda", compiled by Brian Kensley and Marilyn Schotte, Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian Institution Washington, D.C. 20560. It has been compiled from a variety of sources, including numerous individual papers and monographic works, and the collections and catalogues of the National Museum of Natural History, Smithsonian Institution. The following experts co-operated: Dr Thomas E. Bowman, National Museum of Natural History, Smithsonian Institution, (Asellidae in part, Cymothoidae); Dr Niel Bruce, Queensland Museum, South Brisbane, Australia (Aegidae, Cirolanidae, Sphaeromatidae); Dr Richard Brusca, Columbia University Biosphere 2 Center, Oracle, Arizona U.S.A. (Cirolanidae); Dr Hans-Georg Mueller, Germany (Anthuridea, Asellota); Dr Gary C. B. Poore, Museum of Victoria, Melbourne, Australia (Anthuridea, Valvifera); Dr Fahmida Rafi, Canadian Museum of Nature, Ottawa, Canada (Valvifera); Dr Ernest Williams, and Dr Lucy Bunkley Williams, University of Puerto Rico, Mayaguez, Puerto Rico (Cymothoidae). The main list of isopods has not been checked in detail by an expert on the European species. The isopod fauna of deeper waters and parts of the Mediterranean has been least studied (Waegele, personal communication).	
<i>abbreviata</i> (G.O. Sars, 1868)	A M	The Internet address is: http://www.nmnh.si.edu/gopher-menus/WorldListofMarineandFreshwaterCrustacealsopoda.html	
<i>longa</i> Kudinova-Pasternak, 1982	M		
<i>vijazi</i> Kudinova-Pasternak, 1982	M		
<i>Cryptocopoides</i>	A		
<i>arcticus</i> (Hansen, 1886)	A	References	
= <i>Cryptocope arctica</i>	A		
Subfamily Pseudotanainae			
<i>Mystrocentrus</i>	A	Bate, C. S., and J. O. Westwood. 1868. <i>A history of the British sessile-eyed Crustacea</i> . vi + 536 pages. London: John van Voorst.	
<i>serratus</i> Bird & Holdich, 1989	A	Bonnier, J. 1900. Contribution à l'étude des épicarides. Les Bopyridae. <i>Travaux de l'Institut Zoologique de Lille et du Laboratoire de Zoologie Maritime de Wimereux</i> 8: 1-478.	
<i>Parapseudotanais</i>	A	Bourdon, R. 1968. Les Bopyridae des mers européennes. <i>Mémoires du Muséum National d'Histoire Naturelle de Paris</i> , Nouvelle Série (A) 50 (2): 77-424.	
<i>abyssalis</i> Bird & Holdich, 1989	A	Bourdon, R. 1981. Bopyriens nouveaux pour la faune européenne de l'Atlantique (Isopoda Epicaridea). <i>Bulletin du Muséum National d'Histoire Naturelle</i> , Paris (4) 3 (Section A) (no. 2): 615-634.	
<i>Pseudotanais</i>	A	Carayon, J. 1943. Sur les épicarides du Bassin d'Arcachon (2e note). <i>Bulletin de la Société Zoologique de France</i> 68: 43-48.	
<i>longipes</i> Hansen, 1913	A		
<i>similis</i> Sieg, 1977	A		
<i>affinis</i> Hansen, 1887	A		
<i>borceai</i> Bacescu, 1960	A		
<i>colonus</i> Bird & Holdich, 1989	A		
<i>falcicula</i> Bird & Holdich, 1989	A		
= <i>P. abyssi</i> Hansen, 1913	A		
<i>denticulatus</i> Bird & Holdich, 1989	A		
<i>forcipatus</i> (Lilljeborg, 1864)	A		

Caroli, E. 1930. Notizia di tre specie nuove ed una poco nota di Bopiridi addominali, parassiti di Caridei del golfo di Napoli. (<i>Contributo alla conoscenza del genere Phrixus Rathke</i>). <i>Bulletino della Società Naturale di Napoli</i> 41: 258-269.	<i>caprii</i> Wägele, 1981 <i>mjoelniri</i> Negoescu & Svarsson, 1997	M
Codreanu, R., and M. Codreanu. 1963. Sur plusieurs bopyriens parasites branchiaux des anomoures de la Mer Noire, de la Méditerranée et du Viet-Nam. <i>Rapports et Procès verbaux de réunions de la Commission internationale pour l'Exploration scientifique de la mer Méditerranée</i> 17: 283-285.	<i>Malacanthura</i> <i>cantabrica</i> Kensley, 1982 <i>truncata</i> (Hansen, 1916)	A
	<i>Monodanthura</i> <i>maroccana</i> Wägele & Platvoet, 1982	A
	<i>Pilosanthura</i> <i>fresii</i> (Wägele, 1980)	AM
	<i>Quantanthura</i> <i>tyri</i> Negoescu & Svarsson, 1997	A
Giard, A., and J. Bonnier. 1887. Contributions à l'étude des bopyriens. <i>Travaux de l'Institut Zoologique de Lille et du Laboratoire Marine de Wimereux</i> 5: 1-272.	Family Hyssuridae	
Markham, J. C. 1985. A review of the bopyrid isopods infesting caridean shrimps in the northwestern Atlantic Ocean, with special reference to those collected during the Hourglass cruises in the Gulf of Mexico. <i>Memoirs of the Hourglass Cruises</i> 7 (3): 1-156.	<i>Eisothistos</i> <i>macrurus</i> Wägele, 1979 <i>pumilus</i> Wägele, 1979	M
Nobili, G. 1906. Nuovi Bopiridi. <i>Atti della Real Accademia della Scienze di Torino</i> 41: 1-18.	<i>Hyssura</i> <i>ligurica</i> Wägele, 1981 <i>producta</i> Norman & Stebbing, 1886 <i>profunda</i> Barnard, 1925	M
Norman, A. M. 1886. <i>Museum Normanianum, or a catalogue of the Invertebrata of Europe, and the Arctic and North Atlantic Oceans, which are contained in the collection of the Rev. Canon A. M. Norman, M.A.D.C.L., F.L.S. III-Crustacea</i> . Houghton-Le-Spring: Morton. 26 pp.	<i>Kupellonura</i> <i>flexibilis</i> (Pasternak, 1982) <i>mediterranea</i> Barnard, 1925 <i>serritelson</i> Wägele, 1981	M
Popov, V. K. 1927. [Rhizocephala and Bopyridae of the Bay of Sevastopol]. <i>Trudy Sevastopol'skoy Biologicheskoy Stantsii Akademii Nauk SSSR</i> 1 (1-26).	<i>Neohyssura</i> <i>spinicauda</i> (Walker, 1901)	M
Richardson, H. 1905. A monograph on the isopods of North America. <i>Bulletin of the United States National Museum</i> 54: i-iii, I- 727.	<i>Stellanthura</i> <i>cryptobia</i> Wägele, 1979	M
Sars, G. O. 1898. <i>An account of the Crustacea of Norway with short descriptions and figures of all the species</i> . Volume II. <i>Isopoda</i> . Part XI, XII, pp. 185-232. <i>Oniscidae (concluded), Bopyridae, Dajidae</i> . Bergen: Bergen Museum.	Family Paranthuridae	
Tattersall, W. M. 1905. The marine fauna of the coast of Ireland. Part 5. Isopoda. <i>Scientific Investigations for 1904, Fisheries Branch, Ireland</i> 2: 1-90.	<i>Apanthuroides</i> <i>mediterranea</i> (Negoescu, 1981) <i>spathulicauda</i> (Wägele, 1981)	M
Trilles, J.-P., 1999. Ordre des isopodes. Sous-ordre des Épicarides (Epicaridea Latreille, 1825). In Forest, J., ed., <i>Traité de zoologie: Anatomie, systématique, biologie</i> . Tome VII, Fascicule IIIA. <i>Mémoires de l'Institut Océanographique de Monaco</i> 19: 279-352.	<i>Bullowanthura</i> <i>aquitanica</i> Kensley, 1982	A
Order Isopoda	<i>Calathura</i> <i>brachiata</i> (Stimpson, 1854)	A
Suborder Anthuridea	<i>Curassanthura</i> <i>canariense</i> Wägele, 1985	A
Family Antheluridae	<i>Leptanthura</i> <i>affinis</i> (Bonnier, 1896) <i>agulhasensis</i> Kensley, 1975 <i>apalpata</i> Wägele, 1981 <i>chardyi</i> Negoescu, 1992 <i>laevigata</i> (Stimpson, 1855) <i>muelleri</i> Negoescu, 1980 <i>sculpta</i> Pasternak, 1982 <i>tenuis</i> (Sars, 1873) <i>thori</i> Barnard, 1925 <i>victori</i> Negoescu, 1985	A
<i>Ananthura</i> <i>abyssorum</i> (Norman & Stebbing, 1886)	<i>Paranthura</i> <i>costana</i> Bate & Westwood, 1868 <i>nigropunctata</i> (Lucas, 1846)	M
<i>Anthelura</i> <i>elongata</i> Norman & Stebbing, 1886		
<i>ovalis</i> (Barnard, 1925)		M
= <i>Ananthura ovalis</i>		
<i>sulcataicauda</i> (Barnard, 1925)		
= <i>Ananthura sulcataicauda</i>		
<i>truncata</i> (Hansen, 1916)		
Family Anthuridae		
<i>Alloanthura</i> <i>larwoodi</i> Wägele, 1981		M
<i>Amakusanthura</i> <i>iberica</i> Reboreda & Wägele, 1992		A
<i>Anthura</i> <i>gracilis</i> (Montagu, 1808)		AM
<i>Apanthura</i> <i>corsica</i> Amar, 1953		M
<i>libyana</i> Negoescu, 1980		M
<i>tyrrhenica</i> Wägele, 1980		M
<i>Apanthuroides</i> <i>mediterranea</i> (Negoescu, 1981)		M
<i>spatulicauda</i> Wägele, 1981		M
<i>Cyathura</i> <i>carinata</i> (Krøyer, 1847)		A
<i>Haliophasma</i> <i>alaticauda</i> Amar, 1966		M
Suborder Asellota		
Family Acanthaspidiidae		
<i>Acanthaspidia</i> <i>decorata</i> (Hansen, 1895) <i>laevis</i> Chardy, 1978 <i>typhlops</i> (G.O. Sars, 1879)		A
<i>Ianthopsis</i> <i>pulchra</i> (Hansen, 1916)		A
Family Dendrotiidae		
<i>Acanthomunna</i> <i>hystrix</i> (Hansen, 1916)		A
<i>Dendromunna</i> <i>compsa</i> Lincoln & Boxshall, 1983		A
<i>Dendron</i> <i>elegans</i> Lincoln & Boxshall, 1983		A
<i>paradoxum</i> Hansen, 1916		A
<i>setosum</i> Lincoln & Boxshall, 1983		A
<i>spinosum</i> Sars, 1872		A
<i>Munella</i> <i>dantei</i> Bonnier, 1896		A
Family Desmosomatidae		
<i>Balbidocolon</i>		

<i>atlanticum</i> Hessler, 1970		
<i>polare</i> Malyutina & Kussakin, 1996	A	
<i>Chelator</i>	M	
<i>chelatus</i> (Stephensen, 1915)		
<i>stellae</i> Malyutina & Kussakin, 1996	A	
<i>striatus</i> (Menzies, 1962)		
<i>verecundus</i> Hessler, 1970		
<i>vulgaris</i> Hessler, 1970		
<i>Cryodesma</i>	A	
<i>agnari</i> Svavarsson, 1988		
<i>cryoabyssale</i> Svavarsson, 1987	A	
<i>Desmosoma</i>	M	
<i>affine</i> Fresi & Schiecke, 1969		
<i>atypicum</i> Schiecke & Fresi, 1969	M	
<i>elegans</i> Fresi & Schiecke, 1969	M	
<i>elongatum</i> Bonnier, 1896	A	
<i>lineare</i> G.O. Sars, 1864	A	
<i>puritanum</i> Fresi & Schiecke, 1969	M	
<i>serratum</i> Fresi & Schiecke, 1969	M	
<i>strombergi</i> Svavarsson, 1988	A	
<i>thoracicum</i> Fresi & Schiecke, 1969	M	
<i>tyrrhenicum</i> Fresi & Schiecke, 1969	M	
<i>Desmosomella</i>		
<i>armata</i> (G.O. Sars, 1864)		
<i>Disparella</i>		
<i>pachythrix</i> Hessler, 1970		
<i>valida</i> Hessler, 1970		
<i>Echinopleura</i>	A	
<i>aculeata</i> (G.O. Sars, 1864)		
<i>Eugerda</i>	A	
<i>arctica</i> Svavarsson, 1988		
<i>dubia</i> Malyutina & Kussakin, 1996	A	
<i>fragilis</i> (Kussakin, 1965)	A	
<i>fulcimandibulata</i> Hessler, 1970		
<i>gurjanovae</i> Malyutina & Kussakin, 1996	A	
<i>imbricata</i> Hessler, 1970		
<i>mandibulata</i> Malyutina & Kussakin, 1996	A	
<i>pannosa</i> Hessler, 1970		
<i>reticulata</i> (Gurjanova, 1946)	A	
<i>seiflura</i> Hessler, 1970		
<i>tenuimana</i> (G.O. Sars, 1868)		
<i>tertaria</i> Hessler, 1970		
<i>Eugerdella</i>	A	
<i>armata</i> (G.O. Sars, 1864)		
<i>coarctata</i> (G.O. Sars, 1899)	A	
<i>cornuta</i> Hessler, 1970		
<i>hessleri</i> Just, 1980	A	
<i>ischnomesoides</i> Hessler, 1970	A	
<i>lateralis</i> (G.O. Sars, 1899)	A	
<i>longimana</i> (Vanhöffen, 1914)	A	
<i>polita</i> (Hansen, 1916)	A	
<i>pugillator</i> Hessler, 1970		
<i>zenkewitschi</i> (Gurjanova, 1946)	A	
<i>Mirabilicoxa</i>	M	
<i>acuminata</i> Hessler, 1970		
<i>curticolalis</i> Pasternak, 1982		
<i>exopodata</i> Hessler, 1970		
<i>fletcheri</i> Paul & George, 1975	A	
<i>minuta</i> Hessler, 1970		
<i>palpata</i> Hessler, 1970		
<i>plana</i> Hessler, 1970		
<i>Momedossa</i>		
<i>profunda</i> Hessler, 1970		
<i>Oecidiobranchus</i>	A	
<i>glacialis</i> Malyutina & Kussakin, 1996		
<i>nansenii</i> Just, 1980	A	
<i>plebejum</i> (Hansen, 1916)	A	
<i>polare</i> (Gurjanova, 1946)	A	
<i>Prochelator</i>	A	
<i>abyssalis</i> Hessler, 1970		
<i>hampsoni</i> Hessler, 1970		
<i>incomitatus</i> Hessler, 1970		
<i>lateralis</i> (G.O. Sars, 1899)		
<i>litus</i> Hessler, 1970		
<i>uncatus</i> Hessler, 1970		
<i>Pseudogerda</i>		
<i>filipes</i> (Hult, 1936)		
<i>intermedia</i> (Hult, 1936)		
<i>Tau mastosoma</i>		
<i>platycarpus</i> Hessler, 1970		
<i>Torwolia</i>		
<i>creper</i> Hessler, 1970		
<i>subchelatus</i> Hessler, 1970		
<i>Whoia</i>		
<i>angusta</i> (G.O. Sars, 1899)		A
<i>dumbshafensis</i> Svavarsson, 1988		A
<i>variabilis</i> Hessler, 1970		
Family Eurycopidae		
<i>Acanthocope</i>		
<i>annulatus</i> Menzies, 1962		A
<i>armata</i> Chardy, 1972		
<i>carinata</i> Chardy, 1972		
<i>Amuleta</i>		
<i>abyssorum</i> (Richardson, 1911)		A
<i>Baeonectes</i>		
<i>muticus</i> (Sars, 1864)		A
<i>Bathyopsurus</i>		
<i>abyssiculus</i> (Beddard, 1885)		A
<i>Bellibos</i>		
<i>buzwilsoni</i> Haugsness & Hessler, 1979		A
<i>dageti</i> (Chardy, 1975)		A
<i>monicae</i> (Chardy, 1975)		A
<i>Belonectes</i>		
<i>parvus</i> (Bonnier, 1896)		A
<i>Disconectes</i>		
<i>coxalis</i> Kussakin, 1983		A
<i>furcatus</i> (G.O. Sars, 1870)		A
<i>latirostris</i> (Sars, 1882)		A
<i>phalangium</i> (Sars, 1864)		A
<i>picardi</i> (Amar, 1957)		A
<i>Eurycope</i>		
<i>brevirostris</i> Hansen, 1916		A
<i>canariensis</i> Wilson, 1983		A
<i>complanata</i> Bonnier, 1896		A
<i>cornuta</i> Sars, 1864		A
<i>dahli</i> Svavarsson, 1987		A
<i>errabunda</i> Wilson, 1983		A
<i>hessleri</i> Wilson, 1983		A
<i>iphthima</i> Wilson, 1981		A
<i>laticuneata</i> Pasternak, 1982		M
<i>longiflagrata</i> Wilson, 1983		A
<i>nobili</i> Richardson, 1911		A
<i>producta</i> G.O. Sars, 1868		A
<i>pygmaea</i> Sars, 1870		A
<i>ratmanovi</i> Gurjanova, 1946		A
<i>septentrionalis</i> Malyutina & Kussakin, 1996		A
<i>vasinae</i> Malyutina & Kussakin, 1996		A
<i>Lipomera</i>		
<i>lamellata</i> Tattersall, 1905		A
<i>Munneurycope</i>		
<i>glacialis</i> Malyutina & Kussakin, 1996		A
<i>incisa</i> (Gurjanova, 1946)		A
<i>murrayi</i> (Walker, 1903)		
<i>nodifrons</i> (Hansen, 1916)		
<i>Munnopsurus</i>		
<i>arcticus</i> Richardson, 1912		A
<i>atlanticus</i> (Bonnier, 1896)		A
<i>giganteus</i> subsp. <i>giganteus</i> (Sars, 1877)		A
<i>murrayi</i> (Walker, 1903)		A
<i>Storthygura</i>		
<i>atlantica</i> (Beddard, 1885)		A
<i>Syneurycope</i>		
<i>dageti</i> Chardy, 1975		A
<i>Tythocope</i>		
<i>beddardi</i> (Bonnier, 1896)		A
<i>laktionovi</i> (Gurjanova, 1946)		A
<i>megalura</i> (Sars, 1872)		A
<i>neupokoevi</i> (Gurjanova, 1946)		A
<i>pygmaea</i> (Sars, 1879)		A
Family Gnathostenetroididae		

Caecostenetroides			
<i>ischitanum</i> Fresi & Schiecke, 1968	M	<i>hessleri</i> Kussakin, 1988	
<i>ruderalis</i> Stock & Vonk, 1990	A	<i>norvegicus</i> Svavarsson, 1984	A
Gnathostenetroides			
<i>laodicense</i> Amar, 1957	M	Family Janirellidae	
Family Haplomunnidae		<i>Dactylostylis</i>	
<i>Munella</i>		<i>acutispinis</i> Richardson, 1911	A
<i>dantei</i> Bonnier, 1896	A	<i>Janirella</i>	
Family Haploniscidae		<i>abyssicola</i> Richardson, 1911	A
<i>Antennuloniscus</i>		<i>bocqueti</i> Chardy, 1974	
<i>dilatatus</i> Chardy, 1974	A	<i>bonnieri</i> Stephensen, 1915	M
<i>diversus</i> Lincoln, 1985	A	<i>glabra</i> Richardson, 1911	
<i>simplex</i> Lincoln, 1985	A	<i>hessleri</i> Chardy, 1975	A
<i>Haploniscus</i>		<i>laevis</i> Hansen, 1916	
<i>aduncus</i> Lincoln, 1985	A	<i>laubieri</i> Chardy, 1974	
<i>ampliatus</i> Lincoln, 1985	A	<i>nansenii</i> Bonnier, 1896	A
<i>angustus</i> Lincoln, 1985	A	<i>priseri</i> Chardy, 1972	A
<i>bicuspis</i> subsp. <i>bicuspis</i> (G.O. Sars, 1877)	A	<i>spongicola</i> Hansen, 1916	A
<i>bicuspis</i> subsp. <i>tepidus</i> Wolff, 1962	A	Family Janiridae	
<i>borealis</i> Lincoln, 1985	A	<i>Caecijaera</i>	
<i>charcoti</i> Chardy, 1975	A	<i>borealis</i> Kussakin, 1962	A
<i>foresti</i> Chardy, 1974		<i>mirabilis</i> Kussakin, 1962	
<i>furcatus</i> Chardy, 1974		<i>Carpias</i>	
<i>hamatus</i> Lincoln, 1985	A	<i>galloprovincialis</i> (Amar, 1950)	A
<i>ingolfi</i> Wolff, 1962	A	<i>stebbingi</i> (Monod, 1933)	M
<i>monodi</i> Chardy, 1974		<i>Ianiropsis</i>	
<i>myriamae</i> Chardy, 1974		<i>breviremis</i> (Sars, 1883)	
<i>obtusifrons</i> Chardy, 1974		<i>Jaera</i>	
<i>reyssi</i> Chardy, 1974		<i>albifrons</i> subsp. <i>albifrons</i> Leach, 1814	A
Hydroniscus		<i>bocqueti</i> Veuille & Kocatas, 1979	M
<i>abyssi</i> Hansen, 1916		<i>forsmani</i> Bocquet, 1950	
<i>vandeli</i> Chardy, 1974		<i>hopeana</i> Costa, 1853	M
Family Ilyarachnidae		<i>ischiosetosa</i> Forsman, 1949	A
<i>Echinozone</i>		<i>istri</i> Veuille, 1979	M
<i>coronata</i> (Sars, 1870)		<i>italica</i> Kesselyak, 1938	
<i>Ilyarachna</i>		<i>nordica</i> Lemercier, 1958	M
<i>arctica</i> (Hansen, 1916)	A	<i>nordmanni</i> (Rathke, 1837)	M
<i>bergendahli</i> Ohlin, 1901	A	<i>nordmanni</i> subsp. <i>guernei</i> Dollfus, 1889	A
<i>calva</i> Pasternak, 1982	M	<i>nordmanni</i> subsp. <i>illyrica</i> Veuille, 1979	M
<i>clypeata</i> (Sars, 1870)	A	<i>nordmanni</i> subsp. <i>massiliensis</i> Lemercier, 1960	M
<i>derjugini</i> Gurjanova, 1946	A	<i>nordmanni</i> subsp. <i>occidentalis</i> Veuille, 1979	M
<i>frami</i> Just, 1980	A	<i>petiti</i> Schulz, 1953	M
<i>hirticeps</i> Sars, 1870	A	<i>praehirsuta</i> Forsman, 1949	A
<i>longicornis</i> (G.O. Sars, 1864)	A	<i>sarsi</i> Valkanov, 1936	M
<i>mediorientalis</i> Chardy, 1974	M	<i>schellenbergi</i> Kesselyak, 1938	
<i>polita</i> Bonnier, 1896	A	<i>sorrentina</i> Verhoeff, 1943	M
<i>torleivi</i> Svavarsson, 1988	A	<i>syei</i> Bocquet, 1950	
<i>Pseudorachna</i>		<i>Janira</i>	
<i>hirsuta</i> G.O. Sars, 1864	A	<i>alta</i> (Stimpson, 1853)	A
Family Incertae sedis		<i>denticulata</i> Gourret, 1891	M
<i>Tole</i>		<i>maculosa</i> Leach, 1814	A
<i>laciniosa</i> (Sars, 1872)	A	<i>Janiralata</i>	
<i>speciosa</i> (Bovallius, 1881)	A	<i>holmesi</i> (Richardson, 1905)	
<i>Urias</i>		<i>tricornis</i> (Krøyer, 1847)	
<i>spinosus</i> Richardson, 1911	A	<i>Microjaera</i>	
Family Ischnomesidae		<i>anisopoda</i> Bocquet & Levi, 1955	
<i>Haploemesus</i>		<i>Microjanira</i>	
<i>angustus</i> Hansen, 1916	A	<i>dentifrons</i> Schiecke & Fresi, 1970	M
<i>bicuspis</i> (G.O. Sars, 1877)		Family Joeropsidae	
<i>biscayensis</i> Chardy, 1975	A	<i>Joeropsis</i>	
<i>quadrispinosus</i> (Sars, 1879)	A	<i>antarctica</i> Menzies & Schultz, 1968	A
<i>tropicalis</i> Menzies, 1962	M	<i>brevicornis</i> subsp. <i>brevicornis</i> Koehler, 1885	A
<i>Heteromesus</i>		<i>brevicornis</i> subsp. <i>littoralis</i> Amar, 1949	M
<i>dentatus</i> Hansen, 1916	A	<i>dollfusi</i> Norman, 1899	M
<i>frigidus</i> Hansen, 1916	A	<i>legrandi</i> Juchault, 1962	M
<i>greeni</i> (Tattersall, 1905)	A	Family Katianiridae	
<i>schmidti</i> Hansen, 1916	A	<i>Katianira</i>	
<i>similis</i> Richardson, 1911	A	<i>bilobata</i> Gurjanova, 1930	A
<i>spinosus</i> (Beddard, 1886)	A	<i>chelifera</i> Hansen, 1916	A
<i>wolffi</i> Chardy, 1974		Family Macrostyliidae	
<i>Ischnomesus</i>		<i>Macrostyliis</i>	
<i>bispinosus</i> Sars, 1868	A	<i>elongata</i> Hansen, 1916	A
<i>chardyi</i> Kussakin, 1988		<i>longipes</i> Hansen, 1916	A
<i>gracilis</i> Chardy, 1974		<i>longiremis</i> (Meinert, 1890)	A
		<i>polaris</i> Malyutina & Kussakin, 1996	A
		<i>spinifera</i> G.O. Sars, 1864	A

<i>subinermis</i> Hansen, 1916	A	<i>Pleurogonium</i>	
Family Microparasellidae		<i>inerme</i> G.O. Sars, 1882	A
<i>Angleria</i>		<i>intermedium</i> Hansen, 1916	A
<i>phreaticola</i> Chappuis & Delamare, 1952	A	<i>pulchrum</i> Hansen, 1916	A
<i>Microcharon</i>		<i>rubicundum</i> (G.O. Sars, 1864)	A
<i>boutini</i> Boulanouar, Yacoubi, Messouli & Coineau, 1995	A	<i>spinosissimum</i> (G.O. Sars, 1866)	A
		<i>variable</i> Schiecke & Modigh-Tota, 1976	M
Family Munnidae		Family Pseudomesidae	
<i>Munna</i>		<i>Pseudomesus</i>	
<i>acanthifera</i> Hansen, 1916	A	<i>bispinosus</i> Chardy, 1974	M
<i>armoricana</i> Carton, 1962	A	<i>brevicornis</i> Hansen, 1916	A
<i>coeca</i> Gurjanova, 1930	A		
<i>fabricii</i> Krøyer, 1846	A	Family Stenetriidae	
<i>groenlandica</i> Hansen, 1916	A	<i>Hansenium</i>	
<i>hanseni</i> Stappers, 1911	A	<i>caicoensis</i> (Kensley & Heard, 1991)	M
<i>kroyeri</i> Goodsir, 1842	A	<i>Stenetrium</i>	
<i>limicola</i> Sars, 1866	A	<i>mediterraneum</i> Hansen, 1905	M
<i>minuta</i> Hansen, 1916		<i>Tristeniun</i>	
<i>palmata</i> Lilljeborg, 1851	A	<i>longicorne</i> (Lucas, 1849)	M
<i>pellucida</i> Gurjanova, 1930	A		
<i>roemeri</i> Gurjanova, 1930	A	Family Thambematidae	
<i>similis</i> Fresi & Mazzella, 1971	M	<i>Thambema</i>	
<i>spitzbergensis</i> Gurjanova, 1930	A	<i>amicorum</i> Stebbing, 1912	A
<i>wolffi</i> Fresi & Mazzella, 1974	M	<i>statum</i> Harrison, 1987	A
<i>Uromunna</i>	M	<i>golanachum</i> Harrison, 1987	A
<i>petiti</i> (Amar, 1948)		<i>tanum</i> Harrison, 1987	A
Family Munnopsidae			
<i>Lipomera</i>		Suborder Flabellifera	
<i>lamellata</i> Tattersall, 1905	A	Family Aegidae	
<i>Munnopsis</i>		<i>Aega</i>	
<i>beddardi</i> (Tattersall, 1905)	A	<i>arctica</i> Lutken, 1859	A
<i>typica</i> M. Sars, 1861	A	<i>bicarinata</i> Leach, 1818	A
<i>Paramunnopsis</i>		<i>crenulata</i> Lutken, 1859	A
<i>justi</i> Svavarsson, 1988	A	<i>deshaysiana</i> (H. Milne Edwards, 1840)	M
<i>oceanica</i> (Tattersall, 1905)		<i>emarginata</i> Leach, 1818	
Family Nannoniscidae		<i>gracilipes</i> Hansen, 1895	A
<i>Austroniscus</i>		<i>hirsuta</i> Schiödte & Meinert, 1879	A
<i>coronatus</i> Schiecke & Modigh-Tota, 1976	M	<i>incisa</i> Schiödte & Meinert, 1879	M
<i>groenlandicus</i> (Hansen, 1916)	A	<i>loveni</i> Bovallius, 1886	A
<i>nordi</i> Svavarsson, 1982	A	<i>megalops</i> Norman & Stebbing, 1886	A
<i>Hebefustis</i>		<i>microphthalmma</i> Dana, 1853	A
<i>alleni</i> Siebenaller & Hessler, 1977	A	<i>monophthalma</i> Johnston, 1834	A
<i>Nannoniscoides</i>		<i>nordensioides</i> Bovallius, 1885	A
<i>angulatus</i> Hansen, 1916	A	<i>psora</i> (Linnaeus, 1758)	A
<i>Nannonicus</i>		<i>rosacea</i> (Risso, 1816)	M
<i>aequiremus</i> Hansen, 1916	A	<i>stroemii</i> Lutken, 1859	A
<i>affinis</i> Hansen, 1916	A	<i>tridens</i> Leach, 1815	A
<i>arcticus</i> Hansen, 1916	A	<i>ventrosa</i> M. Sars, 1859	A
<i>arctoabyssalis</i> Just, 1980	A	<i>webi</i> (Guérin-Ménéville, 1836)	A
<i>bicuspid</i> G.O. Sars, 1877	A		
<i>laticeps</i> Hansen, 1916	A	<i>Altitropus</i>	
<i>oblongus</i> G.O. Sars, 1870	A	<i>foveolatus</i> Schiödte & Meinert, 1879	
<i>plebejus</i> Hansen, 1916	A	<i>Rocinela</i>	
<i>profundus</i> Svavarsson, 1982	A	<i>danmoniensis</i> Leach, 1818	A
<i>reticulatus</i> Hansen, 1916	A	<i>dumerili</i> (Lucas, 1849)	M
<i>simplex</i> Hansen, 1916	A		
<i>spinicornis</i> Hansen, 1916	A	<i>Syscenus</i>	
<i>teres</i> Siebenaller & Hessler, 1981	A	<i>atlanticus</i> Kononenko, 1988	A
<i>Panetela</i>		<i>lilljeborgii</i> (Bovallius, 1885)	A
<i>compacta</i> Malyutina & Kussakin, 1996	A		
<i>Rapaniscus</i>		<i>Xenuraega</i>	
<i>crassipes</i> (Hansen, 1916)	A	<i>pilosera</i> Tattersall, 1909	A
Family Paramunnidae			
<i>Paramunna</i>		Family Cirolanidae	
<i>bilobata</i> G.O. Sars, 1866		<i>Cirolana</i>	
<i>brevisipes</i> (Bate & Westwood, 1868)		<i>cranchii</i> Leach, 1818	A
<i>typica</i> Tattersall, 1905	A	<i>ferruginosa</i> Risso, 1826	M
Family Pleurocopidae		<i>perlata</i> Barnard, 1936	A
<i>Pleurocope</i>			
<i>dasyura</i> Walker, 1901	M	<i>Conilera</i>	
Family Pleurogoniidae		<i>cylindracea</i> (Montagu, 1804)	A
<i>Neasellus</i>			
<i>oculatum</i>		<i>Eurydice</i>	
		<i>affinis</i> Hansen, 1916	A
		<i>caeca</i> Hansen, 1916	A
		<i>clymeneia</i> Monod, 1926	A
		<i>czerniavskyi</i> Bacescu, 1948	M
		<i>dolfusi</i> Monod, 1930	A
		<i>grimaldii</i> Dollfus, 1888	A
		<i>inermis</i> Hansen, 1890	A
		<i>longispina</i> Jones, 1969	M
		<i>pontica</i> (Czerniavsky, 1868)	M

<i>pulchra</i> Leach, 1815	A	<i>laticauda</i> Milne Edwards, 1840	
<i>racovitzai</i> Bacescu, 1949	M	<i>Lironeca</i>	
<i>rotundicauda</i> Norman, 1906	A	<i>galinae</i> Kononenko, 1989	A
<i>truncata</i> (Norman, 1868)	A	<i>ricinoides</i> Risso, 1816	M
<i>valkanovi</i> Bacescu, 1949	M	<i>Livonectus</i>	
<i>Faucheria</i>		<i>pomatoi</i> Gaillat Airoldi, 1940	M
<i>faucherii</i> (Dollfus & Vire, 1900)	A	<i>Mothocya</i>	
<i>Metacirolana</i>		<i>contracta</i> Costa, 1851	M
<i>hanseni</i> (Bonnier, 1896)	A	<i>Nerocila</i>	
<i>Natalolana</i>		<i>cuspisata</i> Costa, 1851	M
<i>borealis</i> (Lilljeborg, 1851)	A	Family incertae sedis	
<i>caeca</i> (Dollfus, 1903)	M	<i>Aega</i>	
<i>gallica</i> (Hansen, 1905)	A	<i>affinis</i> Milne Edwards, 1840	
<i>imicola</i> (Dollfus, 1903)	A	Family Limnoriidae	
<i>neglecta</i> (Hansen, 1890)	M	<i>Limnoria</i>	
<i>Politolana</i>		<i>borealis</i> Kussakin, 1963	A
<i>micropthalmia</i> (Hoek, 1882)	A	<i>carinata</i> Menzies & Becker, 1957	M
<i>Saharolana</i>		<i>lignorum</i> (Rathke, 1799)	A
<i>seurati</i> Monod, 1930	M	<i>quadripunctata</i> Holthuis, 1949	A
Family Cymothoidae		<i>tuberculata</i> Sowinsky, 1884	M
<i>Ceratothoa</i>			
<i>capri</i> (Trilles, 1964)	A	Family Serolidae	
<i>collaris</i> Schiödte & Meinert, 1883	M	<i>Serolidae i.s.</i>	
<i>gobii</i> Schiödte & Meinert, 1883	M	<i>acutangula</i> Grube, 1871	
<i>italica</i> Schiödte & Meinert, 1883	M	<i>rugosa</i> Kussakin, 1982	
<i>oestroides</i> (Risso, 1816)	M	Family Sphaeromatidae	
<i>parallela</i> (Otto, 1828)	M	<i>Anoplocopea</i>	
<i>potassoniensis</i> (Penso, 1939)	A	<i>hanseni</i> Racovitzta, 1908	M
<i>steindachneri</i> Koelbel, 1879		<i>ischiana</i> Verhoeff, 1943	
<i>trigonocephala</i> (Leach, 1818)		<i>lusitanica</i> Nolting; Reboreda & Wägele, 1998	A
<i>Colombia</i>		Bathyporeaa	
<i>atlantica</i> Schiödte & Meinert, 1881	A	<i>typhlops</i> Tattersall, 1905	A
<i>frontalis</i> Milne Edwards, 1840	M	<i>Campylocpea</i>	
<i>hedenborgi</i> Bovallius, 1887	M	<i>hirsuta</i> (Montagu, 1804)	A
<i>physodes</i> (Linnaeus, 1758)	M	<i>Ciliacea</i>	
<i>rissoniana</i> (Leach, 1818)		<i>latreillei</i> Leach, 1818	
<i>Cymothoa</i>		<i>Cymodoce</i>	
<i>carryensis</i> Gourret, 1892	M	<i>emarginata</i> Leach, 1818	A
<i>curta</i> Schiödte & Meinert, 1884		<i>erythraea</i> subsp. <i>euxinica</i> Bacescu, 1958	M
<i>dufresni</i> Leach, 1818		<i>hanseni</i> Dumay, 1972	M
<i>gibbosa</i> Gourret, 1892	M	<i>pilosa</i> Milne Edwards, 1840	M
<i>globosa</i> Schiödte & Meinert, 1884	A	<i>rubopunctata</i> (Grube, 1864)	M
<i>ichtiola</i> (Brünnich, 1764)		<i>spinosa</i> (Risso, 1816)	M
<i>nigropunctata</i> Risso, 1816	M	<i>tattersalli</i> Torelli, 1929	M
<i>oestrum</i> (Linnaeus, 1758)		<i>truncata</i> Leach, 1814	A
<i>paradoxa</i> (Fabricius, 1787)		<i>tuberculata</i> Costa in Hope, 1851	M
<i>scopulorum</i> (Linnaeus, 1758)	A	<i>Dynamene</i>	
<i>slusarskii</i> Rokicki, 1986		<i>bicolor</i> (Rathke, 1837)	M
<i>Elthusa</i>		<i>bidentatus</i> (Adams, 1800)	A
<i>atlantnroi</i> (Kononenko, 1988)	A	<i>bifida</i> Torelli, 1930	M
<i>Emetha</i>		<i>edwardsi</i> (Lucas, 1849)	M
<i>adriatica</i> Bovallius, 1885	M	<i>magnitorata</i> Holdich, 1968	M
<i>audouini</i> (H. Milne Edwards, 1840)	M	<i>torelliae</i> Holdich, 1968	M
<i>Glossobius</i>		<i>tubicauda</i> Holdich, 1968	M
<i>albinae</i> Kononenko, 1986	A	<i>Dynamenella</i>	
<i>auritus</i> Bovallius, 1885		<i>savignii</i> (H. Milne Edwards, 1840)	M
<i>Idusa</i>		<i>Exosphaeroma</i>	
<i>dieuzeidei</i> Dollfus, 1950	M	<i>gigas</i> (Leach, 1818)	
<i>Mothocya</i>		<i>Ischyromene</i>	
<i>belonae</i> Bruce, 1986	M	<i>bicarinata</i> Harrison, 1981	M
<i>epimerica</i> Costa, 1851	M	<i>lacazei</i> Racovitzta, 1908	M
<i>taurica</i> (Czerniavsky, 1868)	M	<i>Lekanesphaera</i>	
<i>Nerocila</i>		<i>bocqueti</i> (Daguerre de Hureaux, Hoestlandt & Lejuez, 1961)	A
<i>aculeata</i> Milne Edwards, 1840		<i>ephippium</i> (Costa, 1882)	M
<i>acuminata</i> Schiödte & Meinert, 1881	M	<i>glabella</i> Jacobs, 1987	A
<i>bivittata</i> (Risso, 1816)		<i>hoestlandti</i> (Daguerre de Hureaux, Elkaim & Lejuez, 1965)	A
<i>blainvilli</i> Leach, 1818		<i>hookeri</i> (Leach, 1814)	A
<i>depressa</i> Milne Edwards, 1840	M	<i>levii</i> (Argano & Ponticelli, 1981)	A
<i>orbignyi</i> (Guérin-Ménéville, 1832)	M	<i>marginata</i> (H. Milne Edwards, 1840)	M
<i>swainsoni</i> Leach, 1818	M	<i>monodi</i> (Arcangeli, 1934)	M
<i>Olcincira</i>		<i>panousei</i> (Daguerre de Hureaux, Elkaim & Lejuez, 1964)	A
<i>lamarcii</i> Leach, 1818		<i>rugicauda</i> (Leach, 1814)	A
<i>Ourozeuktes</i>			
<i>owenii</i> Milne Edwards, 1840			
<i>Ceratothoa</i>			

<i>sardoa</i> (Arcangeli, 1934)	M	<i>Arcturinella</i>	
<i>teissieri</i> (Bocquet & Lejeuz, 1967)	A	<i>banyulensis</i> Poisson & Maury, 1931	M
<i>terceirae</i> Jacobs, 1987	A	<i>Arcturopsis</i>	
<i>weilli</i> (Elkaim, 1967)	M	<i>giardi</i> (Bonnier, 1896)	M
Monolistra		<i>Arcturus</i>	
<i>bericum</i> subsp. <i>bericum</i> (Fabiani, 1901)	M	<i>scabrosus</i> Norman, 1904	A
<i>boldorii</i> subsp. <i>boldorii</i> Brian, 1931	M	<i>Astacilla</i>	
<i>lavalensis</i> Stoch, 1984	M	<i>arietina</i> G.O. Sars, 1883	A
Sphaeroma		<i>axeli</i> Castello, 1997	A
<i>boryi</i> Guérin-Ménéville, 1832	M	<i>bocagei</i> Nobre, 1903	A
<i>curtum</i> (Leach, 1818)	A	<i>bonnieri</i> Stephens, 1915	M
<i>dumerilii</i> Leach, 1818		<i>gorgonophila</i> Monod, 1925	A
<i>emarginatum</i> Grube, 1864	M	<i>granulata</i> (G.O. Sars, 1877)	A
<i>laevigatum</i> Philippi		<i>intermedia</i> (Goodsir, 1841)	A
<i>podicipitis</i> Monod, 1931	A	<i>longicornis</i> (Sowerby, 1806)	A
<i>prideauxianum</i> Leach, 1818	A	<i>mediterranea</i> Koehler, 1911	M
<i>serratum</i> (Fabricius, 1787)		<i>pusilla</i> (G.O. Sars, 1873)	A
<i>venustissimum</i> Monod, 1931	A	<i>Pleuroprion</i>	
<i>weilli</i> Elkaim, 1967	A	<i>frigidum</i> Hansen, 1916	A
Cymodoce		<i>hystrix</i> (G.O. Sars, 1877)	A
<i>bifida</i> Leach, 1818		<i>Spectarcturus</i>	
Suborder Gnathiidea		<i>multispinatus</i> Schultz, 1981	A
Family Gnathiidae		<i>Arcturus</i>	
<i>Bathynathia</i>		<i>serrulatus</i>	
<i>bathybia</i> (Beddard, 1886)	A	Family Chaetiliidae	
<i>monodi</i> Cals, 1974	A	<i>Chiridotea</i>	
<i>oedipus</i> (Stebbing, 1913)	A	<i>caeca</i> (Say, 1818)	A
<i>Caecognathia</i>		<i>tufsi</i> (Stimpson, 1853)	A
<i>abyssorum</i> (G.O. Sars, 1872)	A	<i>Parachiridotea</i>	
<i>bicolor</i> (Hansen, 1916)	A	<i>panousei</i> Daguerre de Hureaux & Elkaim, 1972	A
<i>caeca</i> (Richardson, 1911)		<i>Proidotea</i>	
<i>elongata</i> (Krøyer, 1849)	A	<i>haughi</i> Racovitz & Sevastos, 1910	M
<i>hirsuta</i> (G.O. Sars, 1877)	A	<i>Saduria</i>	
<i>robusta</i> (G.O. Sars, 1879)	A	<i>entomon</i> (Linnaeus, 1758)	A
<i>schistifrons</i> (Stebbing, 1912)	A	<i>megalura</i> (G.O. Sars, 1879)	A
<i>stygia</i> (G.O. Sars, 1877)	A	<i>sabini</i> (Krøyer, 1849)	A
<i>Elaphognathia</i>		<i>Saduriella</i>	
<i>bacescoi</i> (Kussakin, 1969)	M	<i>losadai</i> Holthuis, 1964	A
<i>Gnathia</i>		Family Holognathidae	
<i>albescens</i> Hansen, 1916	A	<i>Cleantis</i>	
<i>arctica</i> Gurjanova, 1929	A	<i>prismatica</i> (Risso, 1826)	M
<i>dentata</i> (G.O. Sars, 1872)	A	Family Idoteidae	
<i>fallax</i> Monod, 1926	A	<i>Idotea</i>	
<i>illepidus</i> (Wagner, 1869)	M	<i>balthica</i> (Pallas, 1772)	A
<i>inopinata</i> Monod, 1925	M	<i>chelipes</i> (Pallas, 1766)	A M
<i>maxillaris</i> (Montagu, 1804)	A	<i>emarginata</i> (Fabricius, 1793)	A
<i>oxyuraea</i> (Lilljeborg, 1855)	A	<i>granulosa</i> Rathke, 1843	A
<i>panousei</i> Daguerre de Hureaux, 1971	A	<i>hectica</i> (Pallas, 1772)	A
<i>phallonajopsis</i> Monod, 1925	M	<i>linearis</i> (Linnaeus, 1766)	A M
<i>serrulatifrons</i> Monod, 1926		<i>metallica</i> Bosc, 1802	A M
<i>teissieri</i> Cals, 1972	A	<i>neglecta</i> G.O. Sars, 1897	A
<i>vorax</i> (Lucas, 1849)	M	<i>ostroumovi</i> Sowinsky, 1895	
<i>Paragnathia</i>		<i>pelagica</i> Leach, 1815	A
<i>formica</i> (Hesse, 1864)	A	<i>Paridotea</i>	
Suborder Microcerberidea		<i>ungulata</i> (Pallas, 1772)	A
Family Microcerberidae		<i>Synidotea</i>	
<i>Coxicerberus</i>		<i>nodulosa</i> (Krøyer, 1846)	A
<i>adriaticus</i> Karaman, 1955	M	<i>Synisoma</i>	
<i>arenicola</i> Chappuis & Delamare-Deboutteville, 1952	M	<i>acuminatum</i> (Leach, 1815)	A M
<i>remani</i> Chappuis, Delamare-Deboutteville & Paulian,		<i>appendiculata</i> (Risso, 1816)	M
1956	M	<i>bellonae</i> Daguerre de Hureaux, 1968	A
<i>ruffoi</i> Chappuis, 1953	M	<i>capito</i> (Rathke, 1837)	M
<i>Microcerberus</i>		<i>carinata</i> (Lucas, 1849)	M
<i>remyi</i> Chappuis, 1953	M	<i>lancifer</i> (Miers, 1881)	A M
Suborder Valfifera		<i>mediterranea</i> Rezig, 1989	M
Family Arcturidae		<i>nadejda</i> Rezig, 1989	M
<i>Arcturella</i>		<i>spinosa</i> Amar, 1957	M
<i>carlosteroi</i> Reboreda, Wägele & Garmendia, 1994	A	<i>teissieri</i> Prunus & Pantousier, 1976	M
<i>cornuta</i> Koehler, 1911	A	<i>Idotea</i>	
<i>danmoniensis</i> (Stebbing, 1874)	A	<i>brevicornis</i> Rathke, 1843	A
<i>dilatata</i> (Sars, 1883)	A	<i>carinata</i> Lucas, 1849	M
<i>dolfusi</i> Monod, 1925	A	<i>whymperi</i> Miers, 1881	
<i>poorei</i> Castello, 1997	A		

Suborder Epicaridea**Family Bopyridae***Acobelione**reverberii* (Restivo, 1970)*Anisarthrus**pelseneeri* Giard, 1907*Asymmetrione**foresti* (Bourdon, 1968)*Athelges**bilobus* G. O. Sars 1898*cladophorus* Gerstaecker, 1862*guitarra* Gaid & Bonnier, 1890*intermedius* Hesse, 1876*paguri* (Rathke, 1843)*pelagosae* Babi_, 1912*prideauxii* Giard & Bonnier, 1890*tenuicaudis* G. O. Sars, 1898*Bopyrina**ocellata* (Czerniavsky, 1869)*Bopyrissa**diogeni* (Popov, 1927)*frassei* (Carayon, 1943)*Bopyroides**cluthae* (Scott, 1902)*hippolytes* (Krøyer, 1838)*Bopyrus**squillarum* Latreille, 1802*Cancricepon**elegans* (Giard & Bonnier, 1886)*pilula* (Giard & Bonnier, 1886)*Diplophryxus**alveolatus* Bourdon, 1981*Eophrixus**enchophilus* Caroli, 1930*laevimanus* Caroli, 1930*lysmaiae* Caroli, 1930*Epipenaeon**ingens* Nobili, 1906*Epiphrixus**adriaticus* Nierstrasz & Brender à Brandis, 1932*Ergyne**cervicornis* Risso, 1816*Grapoicepon**edwardsi* Bonnier, 1900*Gyge**branchialis* Comalia & Panceri, 1861*Hemiarthrus**abdominalis* (Krøyer, 1840-1841)*Ione**thoracica* (Montagu, 1808)*vicina* Bonnier, 1900*Paragigantione**papillosa* Barnard, 1920*Paraphrixus**adriaticus* Nierstrasz & Brender à Brandis, 1931*Parathelges**cardonae* R. & M. Codreanu, 1968*carolii* R. Codreanu, 1968*racovitzai* R. Codreanu, 1940*Pleurocrypta**galateae* Hesse, 1865*longibranchiata* (Bate & Westwood, 1868)*microbranchiata* G. O. Sars, 1898*piriformis* Bourdon, 1968*porcellanae* Hesse, 1876*strigosa* Codreanu, Codreanu, & Pike, 1968*Pleurocryptella**superba* Bourdon, 1981*Pliophrixus**philonika* Caroli, 1930*Progebiophilus**euxinicus* (Popov, 1929)*Pseudione**affinis* (G. O. Sars, 1882)*borealis* Caspers, 1939*callianassae* Kossmann, 1881*confusa* (Norman, 1886)*crenulata* G. O. Sars, 1898*dohrnii* Bonnier, 1900*hyndmanni* (Bate & Westwood, 1868)*Scyracepion**tuberculosa* Tattersall, 1905*Urobopyrus**processae* Richardson, 1904

CHILOPODA

Compiled by Alessandro Minelli

Centipedes (Chilopoda) and millipedes (Diplopoda) are terrestrial arthropods, most abundant in forest environments. A few species, however, have a more or less obligate relationship to the shore and are usually found under jetsam. Centipedes are predators, whereas most millipedes feed on litter, dead wood and other decaying plant matter. The sources of the present list were Brölemann (1932), Eason (1964), Foddai *et al.* (1995), Strasser & Minelli (1984).

References

- Brolemann H.W. (1932) Chilopodes. In: *Faune de France*, Paris, 25.
 Eason E.H. (1964) *Centipedes of the British Isles*. F. Warne & Co. Ltd. London.
 Foddai D., Minelli A., Scheller U. & Zapparoli M. (1995) Chilopoda, Diplopoda, Pauropoda, Symphyla, in Minelli A., Ruffo S. e La Posta S. (a cura di) *Checklist delle specie della fauna italiana*, 32: 1-35. Calderini, Bologna.
 Strasser C. & Minelli A. (1984) Elenco dei Diplopodi d'Italia. *Lavori Soc. ven. Sc. nat.*, 9:193-212.

Class Chilopoda

Order Geophilomorpha

Family Dignathodontidae

Henia bicarinata (Meinert, 1870)

Family Schendylidae

Hydropschendyla submarina (Grube, 1872)

Family Geophilidae

Tuoba poseidonis (Verhoeff, 1901)

Geophilus fucorum Brölemann, 1900

algarum Brölemann, 1909

Family Linotaeniidae

Strigamia maritima (Leach, 1817)

DIPLOPODA

(see Chilopoda above)

Class Diplopoda

Order Julida

Family Nemasomatidae

Thalassiosobates litoralis (Silvestri)

Family Julidae

Dolichoilulus tongiorgii (Strasser, 1973)

INSECTA

Compiled by Anastasios Legakis and Declan Murray

The list of the marine insects is definitely not complete. There are two reasons for this situation. One is that references to marine insects are scattered in various entomological publications some of which are old and not readily available. Second, the definition of a marine insect is not commonly agreed. Some authors consider them as insects found in marine habitats while others include those that are sporadically associated with the marine environment. The definition proposed by Chang & Frank (1993) considers as marine species an insect which spends at least one of its developmental stages habitually in a marine habitat. Even in this case there are some who consider the supralittoral zone as terrestrial (Bigot *et al.*, 1984). In this list, the supralittoral zone is excluded as a very large number of species would have to be included. The list was compiled using several sources (Riedl, 1970; Chang & Frank, 1993; Hayward & Ryland, 1995).

The majority of juvenile Chironomidae inhabit freshwater but, uncommonly among insects, larvae of some are true halobionts and have been recorded from coasts all over the world (Pinder 1995). The following Chironomidae are known from European coastal waters (Ashe and Cranston, 1990).

References

- Ashe, P. & Cranston, P.S., 1990, Chironomidae. In Soos, A. & Papp, L. (Eds) *Catalogue of Palaearctic Diptera: 2 Psychodidae - Chironomidae*. 113-499. Akadémiai Kiadó, Budapest
 Bigot L., Picard J., Roman M.L. 1984.- Signification des peuplements d'invertébrés des plages et dunes du delta du Rhône; délimitation des domaines marin et terrestre. *C.R. Acad. Sci. Paris* (III)2980(1): 5-7.
 Chang L. & Frank J.H. 1993.- Marine insects and their reproduction. *Oceanogr. Mar. Biol. Ann. Rev.* 31: 479-506.
 Hayward P.J. & Ryland J.S. 1995.- *Handbook of the Marine Fauna of North-West Europe*. Oxford Univ. Press, pp. 462-476.
 Pinder, L.C.V., 1995, The habitats of chironomid larvae. In (Eds) Armitage, P.D., Cranston, P.S., & Pinder, L.C.V., *The Chironomidae - Biology and ecology of non-biting midges*. 107-135. Chapman and Hall, London
 Riedl R. 1970.- *Fauna und Flora der Adria*. Verlag Paul Parey, pp. 371-380.

Class Insecta

Order Archaeognatha

Family Machilidae

Petrobius maritimus (Leach, 1809)
brevistylis Carpenter

Order Coleoptera

Family Carabidae

Aepopsis robinii (Laboulbene, 1894)
Aepus marinus (Stroem, 1788)
Eurynebria complanata (L., 1767)
Broscus cephalotes (L., 1758)

Family Staphylinidae

Micralymma marinum (Stroem, 1785)
Diglotta

submarina (Fairmaire & Laboulbene, 1856)

Bledius

spectabilis (Kraatz, 1857)

Order Collembola

Family Entomobryidae

Pseudosinella

halophila (Bagnall)

Family Hypogastruridae

Hypogastrura

viatica (Tullberg, 1872)

Family Isotomidae

Axelsonia

littoralis (Moniez, 1890)

Family Neanuridae

Anurida

maritima (Guerin, 1838)

Anuridella

marina (Willem)

Family Onychiuridae

Onychiurus

debilis (Moniez)

thalassophila (Bagnall)

Order Dermaptera

Family Carcinophoridae

Anisolabis

maritima (Bonelli, 1832)

Order Diptera

Family Chironomidae

Clunio

adriaticus Schiner, 1856

balticus Heimbach, 1978

marinus Haliday, 1855

ponticus Michailova, 1980

Halocladius

braunsi (Goetghebuer, 1942)

fucicola (Edwards, 1926)

mediterraneus Hirvenoja, 1973

millenarius (Santos Abreu, 1918)

variabilis (Staeger, 1839)

varians (Staeger, 1839)

Telmatogeton

japonicus Tokunaga, 1933

pectinata (Deby, 1889)

Thalassomyia

frauensfeldi Schiner, 1856

Thalassosmittia

atlantica (Stora, 1936)

thalassophila (Bequaert & Goetghebuer, 1913)

Family Ephydriidae

Ephydria

macellaria Egger, 1862

Order Hemiptera

Family Saldidae

Aepophilus

bonnairei (Signoret)

PHORONIDA

Compiled by C. C. Emig, C. Roldán & J. M. Viéitez

The Phoronida is an exclusively marine group of lophophorate animals: infaunal, suspension-feeders, with a vermiform body enclosed in a slender, chitinous tube in which it moves freely and it is anchored by the ampulla, the end-bulb of the body. The tube is embedded in hard or soft substrata.

Phoronids are found in all oceans and seas and are not uncommon in favourable situations. In some habitats they are very abundant, reaching several tens of thousand individuals per square metre. Phoronids occur at depths ranging from the intertidal zone to about 400 m depth, but mainly between 0 to 70 m. Almost all species occurring in European waters have wide geographical ranges and most are probably cosmopolitan.

Several authors regard Phoronida as constituting a separate phylum; but others, including myself (Emig 1997), consider them as a class within the phylum Lophophorata, which also includes the Bryozoa and Brachiopoda. Recently, Cohen (2000) included the Phoronida in a subphylum named Phoroniformea within the Brachiopoda. The Phoronida are particularly distinguished from the other two lophophorate groups on the basis of its cylindrical body form living in a tube which can be compared to the shell of the Brachiopoda and to the exo-skeleton in the Bryozoa.

Currently only two genera, *Phoronis*, Wright, and *Phoronopsis*, Gilchrist, are recognised, together with respectively seven and three well-defined species (Emig 1982). The name *Phoronis* Wright, 1856 from the genus name, is one of the numerous epithets of the Egyptian goddess Isis. The genus *Phoronopsis* has an epidermal invagination at the base of the lophophore.

The characteristic larva of the Phoronida, named actinotroch or *Actinotrocha* Müller, 1846, has been described near Helgoland (Germany). He considered it as an adult form, and named it *Actinotrocha branchiata*. The transformation of the actinotroch into an adult phoronid was described for the first time by Kowalevsky (1867) who, following the metamorphosis, realised that the *Actinotrocha* was a larval stage of Wright's *Phoronis* species.

Separate names for larval and adult forms are still used in taxonomy. Despite the priority of the larval name *Actinotrocha*, the International Commission of Zoological Nomenclature accepted also as valid the name *Phoronis*. Consequently, the actinotroch keeps a separate "generic" name considered as a technical term under *Actinotrocha* which is sometimes still different from the adult species name.

From data of recent ecological surveys in Europe, mainly in the south of the Iberian Peninsula, the Chafarinas Islands and Canary Islands (Emig *et al.* 1999, 2000), the number of phoronid species occurring in the European waters increased to 9 of 10 species known in the world. The species not recorded in Europe, *Phoronis ijimai*, is presently known from Pacific and N. W. Atlantic waters. The Iberian Peninsula and the surrounding islands represent a privileged area for the Phoronida because all 9 species has been recorded along their coasts.

For more detailed information on Phoronida see the website at <http://www.com.univ-mrs.fr/DIMAR/Phoro/>.

References

- Bailey-Brock J. H. & C. C. Emig, 2000. Hawaiian Phoronida (Lophophorata) and their distribution in the Pacific region. *Pacific Science* 54 (2), 119-126.
- Cohen B. L., 2000. Monophyly of brachiopods and phoronids: reconciliation of molecular evidence with Linnaean classification (the subphylum Phoroniformea nov.). *Proceedings of the Royal Society, London, Series B* 267, 225-231.
- Emig C. C. & A. N. Golikov, 1990. On Phoronids of the Far Eastern Seas of the USSR and their distribution in the Pacific Ocean (in Russian). *Zool. Zh.* 69, 22-30.
- Emig C. C. & C. Roldán, 1992. The occurrence in Australia of three species of Phoronida (Lophophorata) and their distribution in the Pacific area. *Rec. S. Aust. Mus.* 26 (1), 1-8.
- Emig C. C., 1982. The biology of Phoronida. *Advances in Marine Biology* 19, 1-89.
- Emig C. C., 1984. On the origin of the Lophophorates. *Z. zool. System. Evolut.-forsch.* 22 (2), 91-94.
- Emig C. C., 1985. Phylogenetic systematics in Phoronida (Lophophorata). *Z. zool. System. Evolut.-forsch.* 23 (3), 184-193.
- Emig C. C., 1997. Les Lophophorates constituent-ils un embranchemen? *Bull. Soc. zool. Fr.* 122 (3), 279-288.
- Emig C. C., García Carrascosa A. M., Roldán C. & J. M. Viéitez, 1999. The occurrence in the Chafarinas Islands (S.E. Alboran Sea, western Mediterranean) of four species of Phoronida (Lophophorata) and their distribution in the north-eastern Atlantic and Mediterranean areas. *Cahiers de Biologie Marine* 40, 129-133.
- Emig C. C., Roldán C. & J. M. Viéitez, 2000. Foronídeos del litoral luso-español. *Boletín de la Real Sociedad Española de Historia Natural (Sección Biología)*, 96 (1-2), in press.
- Viéitez J. M., C. C. Emig, C. Rodríguez Babío & A. M. García Carrascosa, 1987. Foronídeos de las costas de la Península Ibérica e Islas Baleares: sistemática, ecología, distribución y estados larvarios. *Bol. Inst. esp. Oceanogr.* 4 (2), 63-78.

PHYLUM PHORONIDA

- Phoronis*
ovalis Wright, 1856
hippocrepia Wright, 1856
australis Haswell, 1883
muelleri Selys-Lonchamps, 1903
psammophila Cori, 1889
pallida Silén, 1952
- Phoronopsis*
albomaculata Gilchrist, 1907
harmeri Pixell, 1912
californica Hilton, 1930

BRYOZOA (ECTOPROCTA)

Compiled by Peter J. Hayward

This catalogue of European Bryozoa incorporates the checklists prepared by Hayward (1997) and Brattegård (1995, also checked by Hayward), the species lists in Hayward and Ryland (1998, 1999), Zabala and Maluquer (1988), and relevant primary sources published after 1985. All faunistic and systematic papers relating to shelf and coastal European Bryozoa listed in the Zoological Record from 1985 to date have been consulted, but only those including new species or new European records are referenced here. Synonyms of most species included are to be found in the sources cited above; an unsourced entry indicates either a recently described species or unpublished records held by the compiler. The ERMS list was checked by J. Harmelin.

Taxonomic and faunistic research on European Bryozoa has been continuous through the last three decades, but geographical coverage and frequency of investigation have been extremely variable. The British Isles, Ireland and the Atlantic coasts of north-west Europe have attracted most research effort and the faunas of these regions are reasonably well known, although completely new species continue to be discovered even in such well studied areas as the English Channel (Reverter-Gil and Fernandez-Pulpeiro, 1996; Hayward and Hansen, 1999). The Faeroe Islands have been the subject of two surveys (Kramp, 1934; Hayward, 1994), the latter resulting in 15 new species and 13 new records for the temperate northeast Atlantic. The faunas of Iceland and west Greenland are similarly poorly known but are currently under review. The synoptic key of Zabala and Maluquer (1988) was founded on numerous sources relating to the western Mediterranean and Adriatic faunas, on the few available accounts of Aegean Bryozoa and the even sparser records for the southern coasts of the Mediterranean. Mediterranean Bryozoa are still only incompletely known, at best, and it is certain that many more undescribed species will be discovered as research continues.

References

- Alvarez, J. A. 1987. Notas sobre la fauna briozoologica marina Iberica. II. Especies de la Costa Vasca: estudio zoogeográfico. *KOBIE (Serie Ciencias Naturales)* Bilbao, 16: 215 - 222
- Alvarez, J. A. 1990. Una colección de briozos procedentes de la costa de Alicante. *Boletín Inst. Esp. Oceanogr.*, 6: 21 - 40
- Alvarez, J. A. 1992. Sobre algunas especies de la familia Lichenoporidae Smitt, 1866 (Bryozoa, Cyclostomida) en la región Atlántico-Mediterránea. Parte I: género *Disporella* Gray, 1848. *Cahiers de Biología Marine*, 33: 201 - 243
- Alvarez, J. A. 1993. *Fenestrulina barrosoi* sp. nov. (Bryozoa: Cheilostomida) with a review of the genus *Fenestrulina* on the Iberian Peninsula. *Journal of the Marine Biological Association of the U. K.*, 73: 831 - 835
- Alvarez, J. A. 1994. La famille des Lichenoporidae (Bryozoa, Cyclostomida) dans les provinces Atlantico-Méditerranéenne et boréale Partie III: étude des collections du Muséum National d'Histoire Naturelle de Paris. *Cahiers de Biología Marine*, 35: 491 - 509
- Alvarez, J. A. 1995. New data on the family Lichenoporidae Smitt (Bryozoa: Cyclostomida) from the Mediterranean region. *Journal of Natural History*, 29: 1067 - 1079.
- Aristegui, J. 1984. *Briozosos Quilostomados (Ectoprocta, Cheilostomata) de Canarias: estudio sistemático, faunístico y biogeográfico*. Tesis doctoral, Universidad de La Laguna. 524 pp.

- Baldazzi, A., Barbieri, M. and Gristina, M. 1991. Morphology and life strategies of *Aetea* (Bryozoa: Cheilostomata) living on some western Mediterranean *Posidonia oceanica* meadows. *Bulletin de la Société des Sciences Naturelles de l'Ouest de la France, Mémoire HS 1*: 1 - 12.
- Brattegård, T. 1995. Bryozoa. Pp. 278 - 295 in Brattegård, T. and Holthe, T. (eds), *Distribution of marine, benthic macro-organisms in Norway*. Research Report for DN 1997-1. Directorate for Nature Management: Trondheim.
- Fernandez-Pulpeiro, E., Reverter-Gil, O. and Ramil, F. 1990. Inventario de los briozos de Galicia. *Thalassas*, 8: 57 - 67.
- Harmelin, J.-G. 1997. Espèces affines microsympatriques chez *Puellina* (Bryozoa, Cheilostomata) et description d'espèces nouvelles. *Zoologica Scripta*, 17: 25 - 38.
- Harmelin, J.-G. and Aristegui, J. 1988. New Cribrilinidae (Bryozoa, Cheilostomata) from the upper bathyal of the Atlanto-Mediterranean region. *Journal of Natural History*, 22: 507 - 535.
- Harmelin, J.-G. and d'Hondt, J.-L. 1992a. Bryozoaires des parages de Gibraltar (campagne océanographique BALGIM, 1984) 1 - Cheilostomes. *Bulletin du Muséum National d'Histoire Naturelle*, Paris, série 4, 14: 23 - 67.
- Harmelin, J.-G. and d'Hondt, J.-L. 1992b. Bryozoaires des parages de Gibraltar (campagne océanographique Balgim, 1984) 2 - Ctenostomes et Cyclostomes. *Bulletin du Muséum National d'Histoire Naturelle*, Paris, série 4, 14: 605 - 621.
- Harmelin, J.-G., Boronat, J., Moisette, P. and Rosso, A. 1989. *Distansescharella seguenzai* Cipolla, 1921 (Bryozoa, Cheilostomata), nouvelles données morphologiques et écologiques tirées de spécimens fossiles (Miocène, Pliocène) et actuels de Méditerranée. *Geobios*, 22: 484 - 501.
- Hayward, P. J. 1994. New species and new records of cheilostomatous Bryozoa from the Faroe Islands, collected by BIOFAR. *Sarsia*, 79: 181 - 206.
- Hayward, P. J. 1997. Bryozoa. Pp. 273 - 282, in Howson, C. M. and Picton, B. E. (eds), *The species directory of the marine fauna and flora of the British Isles and surrounding seas*. Belfast and Ross-on-Wye: Ulster Museum and the Marine Conservation Society.
- Hayward, P. J. and Hansen, K. B. 1999. Three newly recognised cheilostomate bryozoans from the British sea area. *Journal of the Marine Biological Association of the U.K.* 79: 917 - 921.
- Hayward, P. J. and Ryland, J. S. 1998. Cheilostomatous Bryozoa. Part 1, Aeteoidea - Cribriolinaidea. *Synopsis of the British Fauna* (new series), 10 (second edition): 1 - 366.
- Hayward, P. J. and Ryland, J. S. 1999. Cheilostomatous Bryozoa. Part 2, Hippothooidea - Celleporoidea. *Synopsis of the British Fauna* (new series), 14 (second edition): 1 - 416.
- Hondt, J.-L. d' 1984. Un nouvel immigrant dans le Bassin d'Arcachon, *Watersipora aterrima* (Ortmann, 1890) (Bryozaire Cheilostome). *Congrès national des Sociétés savantes, Dijon, 1984, sciences*, fasc. II: 237 - 245.
- Hondt, J.-L. d' 1988. Bryozoa from the coast of Israel. *Boll. Zool.* 3: 191 - 203.
- Kramp, P. L. 1934. Bryozoa. *Zoology of the Faroes*, 58: 1 - 39.
- Reverter-Gil, O. 1994. *Briozos de la Ría de Ferrol*. Tesis doctoral, Universidad de Santiago de Compostela.
- Reverter-Gil, O. and Fernandez-Pulpeiro, E. 1996. Cribrilinidae from the Ria de Ferrol (NW Spain). *Journal of Natural History*, 30: 1247 - 1260.
- Reverter-Gil, O. and Fernandez-Pulpeiro, E. 1997. Two new species of *Schizomavella* (Bryozoa, Cheilostomatida). *Cahiers de Biologie Marine*, 38: 1 - 6.
- Reverter-Gil, O., d'Hondt, J.-L. and Fernandez-Pulpeiro, E. 1995. Mise à jour de l'inventaire des bryozoaires de Roscoff publié par Echalier et Prentant (1951). *Cahiers de Biologie Marine*, 36: 123 - 131.
- Rosso, A. 1996. Popolamenti e Tanatocenosi a briozoi di fondi mobili circalitorali del Golfo di Noto (Sicilia, Italia). *Naturalista sicil. S. IV*, 30: 189 - 225.
- Rosso, A. 1999. Recent and fossil species of *Characodoma* Maplestone, 1900 (Bryozoa) from the Mediterranean, with description of two new species. *Journal of Natural History*, 33: 415 - 437.
- Zabala, M. and Maluquer, P. 1988. Illustrated keys for the classification of Mediterranean Bryozoa. *Treballs del Museu de Zoologia, Barcelona*, 4: 1 - 294.
- Zabala, M., Maluquer, P. and Harmelin, J.-G. 1993. Epibiotic bryozoans on deep-water scleractinian corals from the Catalonia slope (western Mediterranean, Spain, France). *Scientia Marina*, 57: 65 - 78.
- ## PHYLUM BRYOZOA
- ### Class Gymnolaemata
- #### Order Cheilostomatida
- ##### Family Adeonidae
- Adeonella*
calveti (Canu & Bassler, 1930)
polystomella (Reuss, 1847)
- ##### *Adeonellopsis*
- distoma* (Busk, 1858)
multiporosa Aristegui, 1985
- ##### *Anarthropora*
- monodon* (Busk, 1860)
- ##### *Repadileonella*
- insidiosa* (Jullien, 1903)
violacea (Johnston, 1847)
- ##### Family Aeteidae
- Aetea*
anguina (Linnaeus, 1758)
lepadiformis Waters, 1906
ligulata Busk, 1852
longicollis (Jullien, 1903)
sica (Couch, 1844)
truncata (Landsborough, 1852)
- ##### Family Ascidiidae
- Ascobia*
pandora Jullien, 1882
- Fedora*
edwardsi Jullien, 1882
- ##### Family Beanidae
- Beania*
hirtissima (Heller, 1967)
hirtissima subsp. *cylindrica* (Hincks, 1886)
magellanica (Busk, 1852)
mirabilis Johnston, 1840
robusta (Hincks, 1881)
thula Hayward, 1994
- ##### Family Bitectiporidae
- Codonellina*
lacunata Hayward & Hansen, 1999
- Hippomonavella*
borealis (Waters, 1900)
- Hippoporina*
harmsworthi (Waters, 1900)
lineolifera (Hincks, 1886)
pertusa (Esper, 1796)
reticulatopunctata (Hincks, 1877)
- Metroperiella*
lepralioides (Calvet, 1903)
- Pentapora*
fascialis (Pallas, 1766)
= *Pentapora foliacea* (see Hayward & Ryland 1999)
- ottomulleriana* (Möll, 1803)
- Pseudofistularia*
anderssoni Kluge, 1946
hincksi Kluge, 1946
minima Hayward, 1994
sinuosa (Andersson, 1902)
solida (Stimpson, 1854)
virgula Hayward, 1994
- Schizomavella*
arrogata (Waters, 1879)
auriculata (Hassall, 1842)
auriculata subsp. *asymetrica* (Calvet, 1927)

- auriculata* subsp. *hirsuta* (Calvet, 1927)
auriculata subsp. *inordinata* Canu & Bassler, 1930
auriculata subsp. *leontiniensis* (Waters, 1878)
auriculata subsp. *lineata* (Nordgaard, 1896)
cristata (Hincks, 1879)
cuspidata (Hincks, 1880)
discoidea (Busk, 1859)
fischeri (Jullien, 1882)
gautieri Reverter Gil & Fernandez Pulpeiro, 1997
hastata (Hincks, 1862)
hondti Reverter Gil & Fernandez Pulpeiro, 1996
linearis (Hassall, 1841)
linearis subsp. *profunda* Harmelin & d'Hondt, 1992
mamillata (Hincks, 1880)
marsupifera (Busk, 1884)
monoecensis (Calvet, 1927)
neptuni (Jullien, 1882)
ochracea (Hincks, 1880)
porifera (Smitt, 1868)
rudis (Manzoni, 1869)
sarniensis Hayward & Thorpe, 1995
teresae Reverter Gil & Fernandez Pulpeiro, 1996
triangularis Reverter Gil & Fernandez Pulpeiro, 1997
- Family Bryocryptellidae**
- Bryocryptella*
koehleri (Calvet, 1896)
torquata (Jullien, 1903)
- Cystisella*
saccata (Busk, 1856)
- Marguettia*
pulchra Jullien, 1903
- Palmicellaria*
cf. aviculifera (Canu & Bassler, 1928)
- Palmiskenea*
aquila Hayward, 1994
faroeensis Hayward, 1994
lorea (Alder, 1864)
plana (Hincks, 1888)
skenei (Ellis & Solander, 1786)
- Porella*
acutirostris Smitt, 1868
alba Nordgaard, 1906
belli Dawson, 1859
concinna subsp. *tubulata* Calvet, 1927
compressa (J. Sowerby, 1805)
concinna (Busk, 1854)
cymosa Hayward, 1994
groenlandica (Norman, 1894)
laevis (Fleming, 1828)
minuta (Norman, 1868)
patula (M. Sars, 1851)
peristomata (Nordgaard, 1905)
princeps Norman, 1903
proboscidea Hincks, 1888
smitti Kluge, 1907
struma (Norman, 1868)
tubulata (Busk, 1861)
turgidula Hayward, 1994
- Rhamphostomella*
argentea (Hincks, 1881)
bilaminata (Hincks, 1877)
costata Lorenz, 1886
hincksi Nordgaard, 1906
ovata (Smitt, 1868)
plicata (Smitt, 1868)
radiatula (Hincks, 1877)
scabra (O. Fabricius, 1780)
spinigera Lorenz, 1886
- Family Bugulidae**
- Bicellariella*
ciliata (Linnaeus, 1758)
- Bicellarina*
alderi (Busk, 1859)
- Bugula*
aperta (Hincks, 1886)
- avicularia* (Linnaeus, 1758)
calathus Norman, 1864
calathus subsp. *calathus* Ryland, 1962
calathus subsp. *minor* Ryland, 1962
flabellata (J V Thompson, in Gray, 1848)
fulva Ryland, 1960
gautieri Ryland, 1961
germanae Calvet, 1902
gracilis (Busk, 1852)
harmsworthi Waters, 1900
neritina (Linnaeus, 1758)
plumosa (Pallas, 1766)
prenanti Castric-Fey, 1971
purpuroincta Norman, 1868
simplex Hincks, 1886
spicata (Hincks, 1880)
stolonifera Ryland, 1960
tricuspis Kluge, 1915
turbinata Alder, 1857
- Bugulella*
elegans Hayward, 1978
fragilis Verrill, 1879
- Corynoporella*
tenuis Hincks, 1888
- Dendrobeania*
decorata (Verrill, 1879)
= *Bugula elongata* Nordgaard
elongata (Nordgaard, 1906)
fessa Kluge, 1955
fruticosa (Packard, 1863)
murrayana (Bean, in Johnston, 1847)
pseudomurrayana Kluge, 1955
quadrivalvata (Loven, 1834)
sessilis (d'Hondt, 1974)
- Kinetoskias*
arborescens Danielssen, 1868
smitti Danielssen, 1868
- Nordgaardia*
cornucopoides d'Hondt, 1983
pusilla (Nordgaard, 1907)
- Uschakovia*
gorbunovi Kluge, 1946
- Family Calescharidae**
- Coronellina*
jagei (Gautier, 1962)
- Family Calloporidae**
- Alderina*
imbellis (Hincks, 1860)
- Ammatophora*
nodulosa (Hincks, 1877)
- Amphilestrum*
auritum (Hincks, 1877)
flemingii (Busk, 1854)
lyrulatum (Calvet, 1907)
septentrionalis (Kluge, 1906)
solidum (Packard, 1860)
- Aplousina*
capriensis (Waters, 1898)
filum (Jullien, 1903)
- Callopora*
craticula (Alder, 1856)
depressa Cook, 1968
discreta (Hincks, 1862)
dumerilii (Audouin, 1826)
dumerilii subsp. *pouilleti* (Alder, 1857)
lineata (Linnaeus, 1767)
minuta Harmelin, 1973
rylandi Bobin & Prenant, 1965
whiteavesi Norman, 1903
- Cauloramphus*
cymbiformis (Hincks, 1877)
spiniferum (Johnston, 1832)
- Clavodesia*
biradiculata Harmelin & d'Hondt, 1992
- Copidozoum*

- exiguum* (Barroso, 1920)
macilenta (Jullien, 1882)
planum (Hincks, 1880)
tenuirostre (Hincks, 1880)
- Crassimarginatella**
crassimarginata (Hincks, 1880)
maderensis (Waters, 1898)
solidula (Hincks, 1860)
tensa (Norman, 1903)
- Doryporella**
spathulifera (Smitt, 1868)
reticulata Ryland, 1963
- Ellisina**
antarctica Hastings, 1945
cf. *antarctica*
gautieri Fernandez Pulpeiro & Reverter Gil, 1993
- Megapora**
ringens (Busk, 1856)
- Parellisina**
curvirostris (Hincks, 1862)
- Ramphonotus**
minax (Busk, 1860)
gorbunovi Kluge, 1946
- Retevirgula**
akdenizae Chimenz et al., 1997
- Tegella**
arctica (d'Orbigny, 1851)
armifera (Hincks, 1880)
kildinensis Kluge, 1955
norvegica Nordgaard, 1918
spitzbergensis (Bidenkap, 1897)
unicornis (Fleming, 1828)
- Family Candidae**
- Caberea**
boryi (Audouin, 1826)
ellisi (Fleming, 1814)
- Notoplites**
damicornis Hayward & Ryland, 1978
evocata (Jullien, 1882)
= *Notoplites normani*
harmeri Ryland, 1963
jeffreysii (Norman, 1868)
sibirica (Kluge, 1929)
smitti (Norman, 1868)
- Scrupocellaria**
aegeensis Harmelin, 1969
arctica (Busk, 1855)
bertholleti (Audouin, 1826)
delilii (Audouin, 1826)
elongata (Smitt, 1868)
incurvata Waters, 1896
inermis Norman, 1867
intermedia Norman, 1893
jolloisi (Audouin, 1826)
- Scupocellaria**
macrorhyncha Gautier, 1962
- Scrupocellaria**
maderensis Busk, 1860
reptans (Linnaeus, 1767)
scabra (van Beneden, 1848)
scruposa Busk, 1852
scruposa (Linnaeus, 1758)
spatulata d'Orbigny, 1851
- Tricellaria**
gracilis (van Beneden, 1848)
inopinata d'Hondt et al., 1985
peachii (Busk, 1851)
ternata (Ellis & Solander, 1786)
- Family Cellariidae**
- Cellaria**
fistulosa (Linnaeus, 1758)
normani Hastings, 1946
salicornioides Lamouroux, 1816
sinuosa (Hassall, 1840)
- Euginoma**
- reticulata* d'Hondt, 1981
vermiformis Jullien, 1882
- Family Celleporidae**
- Buffonellaria**
divergens (Smitt, 1873)
- Buskea**
billardi (Calvet, 1906)
dichotoma (Hincks, 1862)
nitida (Heller, 1867)
quinuncialis (Norman, 1867)
- Cellepora**
pumicosa Waters, 1877
pumicosa (Pallas, 1766)
- Celleporaria**
sardonica (Waters, 1879)
- Celleporina**
caminata (Waters, 1879)
canariensis Aristegui, 1989
decipliens Hayward, 1976
hassallii (Johnston, 1847)
lucida (Hincks, 1880)
pygmaea (Norman, 1868)
surcularis (Packard, 1863)
tubulosa (Hincks, 1880)
ventricosa (Lorenz, 1886)
- Lagenipora**
lepraloides (Norman, 1868)
- Omalosecosa**
ramulosa (Linnaeus, 1767)
- Palmicellaria**
elegans Alder, 1864
- Rhamphostomellina**
posidoniae Hayward, 1975
- Turbicellepora**
avicularis (Hincks, 1860)
boreale Hayward & Bille Hansen, 1999
camera Hayward, 1978
canaliculata (Busk, 1886)
cantabra (Barroso, 1919)
coronopus (Wood, 1844)
coronopusoidea Calvet, 1931
crenulata Hayward, 1978
magnicostata (Barroso, 1919)
nodulosa Lorenz, 1886
smitti (Kluge, 1962)
torquata Hayward, 1978
tubigera (Busk, 1859)
- Family Chaperiidae**
- Chaperiopsis**
annulus (Manzoni, 1870)
- Larnacicus**
corniger (Busk, 1859)
- Family Cheiloporinidae**
- Cheilopora**
sincera (Smitt, 1868)
- Cheiloporina**
circumcincta Neviani, 1896
- Family Chlidoniidae**
- Chlidonia**
pyriformis (Bertoloni, 1810)
- Crepis**
longipes Jullien, 1882
- Family Chorizoporidae**
- Chorizopora**
brongniartii (Audouin, 1826)
- Family Cleidochasmidae**
- Characodoma**
mamilatum (Seguenza, 1880)
- Cleidochasma**
oranense (Waters, 1918)
porcellanum (Busk, 1860)
strangulatum (Calvet, 1906)
- Cleidochasmidra**

- canakkalense* Unsal & d'Hondt, 1979
- Family Cribrilinidae**
- Collarina*
 - balzaci* (Audouin, 1826)
 - Cribrilina*
 - alcicornis* Jullien, 1882
 - annulata* (O. Fabricius, 1780)
 - cryptoecium* Norman, 1903
 - punctata* (Hassall, 1841)
 - spiculifera* Kluge, 1955
 - watersi* Andersson, 1902
 - Distansescharella*
 - seguenzai* Cipolla, 1921
 - Figularia*
 - figularis* (Johnston, 1847)
 - Gephyrotes*
 - nitidopunctata* (Smitt, 1868)
 - Membraniporella*
 - nitida* (Johnston, 1838)
 - Puellina*
 - arrecta* Bishop & Househam, 1987
 - bathyalis* Harmelin & Aristegui, 1988
 - bifida* (d'Hondt, 1970)
 - cassidainsis* (Harmelin, 1984)
 - corbula* Bishop & Househam, 1987
 - denticulata* Harmelin & Aristegui, 1988
 - directa* Bishop & Househam, 1987
 - gattyae* (Landsborough, 1852)
 - hincksi* (Friedl, 1917)
 - innominata* (Couch, 1844)
 - minima* (Harmelin, 1984)
 - modica* Bishop & Househam, 1987
 - orientalis* subsp. *azorensis* Harmelin, 1988
 - orientalis* subsp. *lusitanica* (Harmelin, 1988)
 - orientalis* subsp. *orientalis* Harmelin & Aristegui, 1987
 - pseudoradiata* subsp. *canariensis* Harmelin & Aristegui, 1988
 - pseudoradiata* subsp. *pseudoradiata* Harmelin & Aristegui, 1988
 - parva* Reverter Gil & Fernandez Pulpeiro, 1996
 - pedunculata* Gautier, 1956
 - picardi* Harmelin, 1987
 - praecox* Bishop & Househam, 1987
 - pseudoradiata* Harmelin & Aristegui, 1988
 - radiata* (Moll, 1803)
 - setiformis* Harmelin & Aristegui, 1988
 - setiformis* subsp. *romana* Harmelin & Aristegui, 1988
 - setiformis* subsp. *setiformis* Harmelin & Aristegui, 1988
 - setosa* (Waters, 1899)
 - venusta* Canu & Bassler, 1925
- Family Cryptosulidae**
- Cryptosula*
 - pallasiana* (Moll, 1803)
 - Hameria*
 - scutelata* (Busk, 1855)
- Family Electridae**
- Conopeum*
 - reticulum* (Linnaeus, 1767)
 - seurati* (Canu, 1928)
- Family Cupuladriidae**
- Cupuladria*
 - biporosa* (Canu & Bassler, 1923)
 - canariensis* (Busk, 1859)
 - doma* (d'Orbigny, 1851)
 - multispinata* (Canu & Bassler, 1923)
- Family Cyclicoporidae**
- Lepraliooides*
 - nordlandica* (Nordgaard, 1905)
- Family Electridae**
- Aspidelectra*
 - melolontha* (Landsborough, 1852)
- Electra**
- crustulenta* (Pallas, 1766)
 - monostachys* (Busk, 1854)
 - pilosa* (Linnaeus, 1767)
 - posidoniae* Gautier, 1957
 - verticillata* (Ellis & Solander, 1786)
- Pyripora**
- catenularia* (Fleming, 1828)
- Family Epistomiidae**
- Epistomia*
 - bursaria* (Linnaeus, 1758)
 - Synnotum*
 - aegytiacum* (Audouin, 1826)
- Family Eucrateidae**
- Eucrata*
 - loricata* (Linnaeus, 1758)
- Family Flustridae**
- Carbasea*
 - carbasea* (Ellis & Solander, 1786)
 - Chartella*
 - barleei* (Busk, 1860)
 - membranaceotruncata* (Smitt, 1868)
 - papyracea* (Ellis & Solander, 1786)
 - tenella* (Hincks, 1887)
 - Flustra*
 - foliacea* (Linnaeus, 1758)
 - Gregarinidra*
 - gregaria* (Heller, 1867)
 - Hincksina*
 - flustroides* subsp. *crassispinata* Calvet
 - longispinosa* Harmelin & d'Hondt, 1992
 - sceletos* (Busk, 1858)
 - Hincksinoflustra*
 - octodon* (Busk, 1852)
 - Sarsiflustra*
 - abyssicola* (Sars, 1872)
 - Securiflustra*
 - securifrons* (Pallas, 1766)
 - Terminoflustra*
 - calvetti* (Guerin-Ganivet, 1911)
- Family Gigantoporidae**
- Coscinioipsis*
 - ambita* Hayward, 1974
 - Cylindroporella*
 - tubulosa* (Norman, 1868)
- Family Haplopomidae**
- Haplopoma*
 - bimucronatum* (Moll, 1803)
 - bimucronatum* subsp. *occiduum* (Waters, 1877)
 - graniferum* (Johnston, 1847)
 - graniferum* subsp. *carinatum* (Calvet, 1902)
 - impressum* (Audouin, 1826)
 - planum* Ryland, 1963
 - sciaphilum* Silén & Harmelin, 1976
- Family Heliodomidae**
- Setosellina*
 - capriensis* (Waters, 1926)
 - constricta* Harmer, 1926
 - roulei* Calvet, 1906
- Family Hincksiporidae**
- Hincksipora*
 - spinulifera* (Hincks, 1889)
- Family Hippaliosinidae**
- Hippaliosina*
 - depressa* (Busk, 1854)
- Family Hippopodinidae**
- Hippomenella*
 - mucronelliformis* (Waters, 1899)
 - Hippopodina*
 - feegeensis* (Busk, 1884)
- Family Hippoporididae**
- Hagiosynodos*

- latus* (Busk, 1856)
kirchenpaueri (Heller, 1867)
kirchenpaueri subsp. *tregoubovii* Gautier, 1962

Hippoporellidae

obesa (Waters, 1900)
Hippoporidra
lusitania Taylor & Cook, 1981
picardi Gautier, 1962

Family Hippothoidae

Celleporella
angusta Alvarez, 1991
gallica Cesar-Aldariz *et al.*, 1999
hyalina (Linnaeus, 1767)

Haplota
clavata (Hincks, 1857)

Hippothoa
divaricata Lamouroux, 1821
flagellum Manzoni, 1870

Plesiothoa
gigerium (Ryland & Gordon, 1977)

Family Lacernidae

Cribellopora
trichotoma (Waters, 1918)

Family Lanceoporidae

Calyptotheca
obscura Harmelin *et al.*, 1989
rugosa Hayward, 1974
triarmata Hayward, 1974

Family Lepraliellidae

Buchneria
sayalensis (Waters, 1888)
Lepraliella
contigua (Smitt, 1868)
hippopus (Smitt, 1867)

Family Margarettidae

Margareta
cereoides (Ellis & Solander, 1786)

Family Membraniporidae

Membranipora
arborescens (Canu & Bassler, 1928)
membranacea (Linnaeus, 1767)
tenuis Desor, 1848

Family Microporellidae

Calloporina
decorata (Reuss, 1848)
Diporula
verrucosa (Peach, 1868)

Fenestrulina

asturiasensis Alvarez, 1992
barrosoi Alvarez, 1993
joannae (Calvet, 1902)
malusii (Audouin, 1826)

Microporella
appendiculata (Heller, 1867)
arctica Norman, 1903
ciliata (Pallas, 1766)
coronata (Audouin, 1826)
orientalis Harmer, 1957
umbracula (Audouin, 1826)

Family Microporidae

Calpensia
nobilis (Esper, 1796)

Micropora
coriacea (Johnston, 1847)
normani Levinson, 1909

Mollia
circumcincta (Heller, 1867)
patellaria (Moll, 1816)

Monoporella
nodulifera (Hincks, 1881)

Rosseliana
rosselii (Audouin, 1826)

Steraechmella

buski Lagajai, 1952

Family Myriaporidae

Myriapora
coarctata (M. Sars, 1863)
subgracile (d'Orbigny, 1852)
truncata (Pallas, 1766)

Family Onychocellidae

Onychocella
angulosa (Reuss, 1847)
marioni (Jullien, 1881)
vibraculifera Neviani, 1895

Family Calloporidae

Rectonychoella
disjuncta Canu & Bassler, 1930

Family Onychocellidae

Smittipora
solida (Nordgaard, 1907)

Family Pasytheidae

Gemmellipora
eburnea (Smitt, 1873)

Family Phidoloporidae

Hippellozoon
mediterraneum (Waters, 1894)

Jaculina
blanchardi Jullien, 1903
parallelata (Waters, 1895)
tessellata Hayward, 1979

Phidolopora
elongata (Smitt, 1868)

Reteporella
aporosa (Waters, 1894)
atlantica (Busk, 1884)
beaniana (King, 1846)
complanata (Waters, 1894)
couchii (Hincks, 1878)
elegans Harmelin, 1976
feuerbornii Hass, 1948
grimaldii (Jullien, 1903)
harmeri Hass, 1948
incognita Hayward & Ryland, 1996
jermanensis (Waters, 1909)
mediterranea Hass, 1948
septentrionalis Harmer, 1933
sparteli (Calvet, 1906)
sudbourniensis Gautier, 1962
watersi (Nordgaard, 1907)

Reteporellina
delicatula Hayward, 1974
hyperborea Hayward, 1994

Rhynchozoon
bispinosum (Johnston, 1847)
globosum Harmer, 1957
neapolitanum Gautier, 1962
pseudodigitatum Zabala & Maluquer, 1988
quadrispinatum Zabala & Maluquer, 1988

Schizoretepora
imperati (Busk, 1884)
longisetae (Canu & Bassler, 1928)
solanderia (Risso, 1826)

Schizotheca
divisa (Norman, 1864)
fissa (Busk, 1856)
serratimargo Hincks, 1886
tubigera (Jullien, 1903)

Stephanollona
armata (Hincks, 1862)

Family Phoceanidae

Phocea
columnaris Jullien, 1903

Sertulipora
guttata Harmelin & d'Hondt, 1992

Family Romancheinidae

Arctonula

- arctica* (M. Sars, 1851)
- Escharella**
- abyssicola* (Norman, 1869)
 - acuta* Zabala *et al.*, 1993
 - connectens* (Ridley, 1881)
 - hexaspinosa* Aristegui, 1986
 - immersa* (Fleming, 1828)
 - indivisa* Levinse, 1916
 - klugei* Hayward, 1979
 - labiata* (Boeck MS, in Smitt, 1868)
 - labiosa* (Busk, 1856)
 - laqueata* (Norman, 1864)
 - levinsemi* Hayward, 1994
 - longicollis* (Jullien, 1882)
 - octodentata* (Hincks, 1880)
 - praealta* (Calvet, 1906)
 - rylandi* Geraci, 1994
 - variolosa* (Johnston, 1838)
 - ventricosa* (Hassall, 1842)
- Escharoides**
- bidenkapi* (Kluge, 1946)
 - coccinea* (Abildgaard, 1806)
 - jacksoni* (Waters, 1900)
 - mamillata* (Wood, 1844)
 - megarostris* (Canu & Bassler, 1928)
- Exechonella**
- antillea* (Osburn, 1927)
- Hemicyclopora**
- discrepans* (Jullien, 1903)
 - emucronata* (Smitt, 1872)
 - labrata* Hayward, 1994
 - microstoma* (Norman, 1864)
 - multispinata* (Busk, 1861)
 - polita* (Norman, 1864)
- Neolagenipora**
- collaris* (Norman, 1867)
 - eximia* (Hincks, 1860)
 - rugosa* Hayward, 1994
- Family Savignyellidae**
- Savignyella*
 - lafontii* (Audouin, 1826)
- Family Schizoporellidae**
- Arthropoma*
 - cecili* (Audouin, 1826)
 - Escharina*
 - alderi* (Busk, 1856)
 - boreale* Hayward, 1994
 - dutertrei* (Audouin, 1826)
 - dutertrei* subsp. *haywardi* Zabala *et al.*, 1993
 - dutertrei* subsp. *protecta* Zabala *et al.*, 1993
 - hyndmanni* (Johnston, 1847)
 - johnstoni* (Quelch, 1884)
 - porosa* (Smitt, 1973)
 - vulgaris* (Moll, 1803)
 - Phaeostachys*
 - spinifera* (Johnston, 1847)
 - Schizobrachiella*
 - sanguinea* (Norman, 1868)
 - Schizoporella*
 - cornualis* Hayward & Ryland, 1995
 - crustacea* (Smitt, 1868)
 - dunkeri* (Reuss, 1848)
 - = *Schizoporella longirostris* Hincks, 1886
 - elmwoodiae* (Waters, 1900)
 - errata* (Waters, 1878)
 - hesperia* Hayward & Ryland, 1995
 - hexagona* Nordgaard, 1905
 - limbata* Lorenz, 1886
 - magnifica* Hincks, 1886
 - magniporata* Nordgaard, 1906
 - mutabilis* Calvet, 1927
 - neptuni* (Jullien, 1882)
 - patula* Hayward & Ryland, 1995
 - smitti* Kluge, 1962
 - tetragona* (Reuss, 1848)
- unicornis* (Johnston, in Wood, 1844)
- Family Scrupariidae**
- Scruparia*
 - ambigua* (d'Orbigny, 1841)
 - chelata* (Linnaeus, 1758)
- Family Setosellidae**
- Setosella*
 - cavernicola* Harmelin, 1977
 - solini* (Jullien, 1882)
 - vulnerata* (Busk, 1860)
- Family Smittinidae**
- Parasmittina*
 - jeffreysi* (Norman, 1903)
 - parsevalii* (Audouin, 1826)
 - raigii* (Audouin, 1826)
 - trispinosa* (Johnston, 1838)
 - tropica* (Waters, 1909)
 - Smittina*
 - affinis* (Hincks, 1862)
 - bella* (Busk, 1860)
 - cervicornis* (Pallas, 1766)
 - cheilostoma* (Manzoni, 1870)
 - colleti* (Jullien, 1903)
 - crystallina* (Norman, 1867)
 - inerna* (Calvet, 1906)
 - jordii* Reverter Gil & Fernandez Pulpeiro, 1999
 - jeffreysi* Norman, 1903
 - landsborovii* (Johnston, 1847)
 - majuscula* (Smitt, 1868)
 - malleolus* (Hincks, 1884)
 - minuscula* (Smitt, 1868)
 - remotorostrata* Canu & Bassler, 1928
 - rigida* Lorenz, 1886
 - smitti* (Kirchenpauer, 1874)
 - tubulifera* (Heller, 1867)
- Smittoidea**
- amplissima* Hayward, 1979
 - exilis* Hayward, 1994
 - glaciata* Waters, 1900
 - marmorea* (Hincks, 1877)
 - microoechia* Hayward, 1994
 - ophidiana* (Waters, 1879)
 - propinqua* (Smitt, 1868)
 - reticulata* (MacGillivray, 1842)
- Family Stomachetosellidae**
- Cigclisula*
 - turrita* (Smitt, 1873)
 - Stomachetosella*
 - cruenta* (Busk, 1854)
 - normani* Hayward, 1994
 - sinuosa* (Busk, 1860)
- Family Tessaradomidae**
- Tessaradoma*
 - boreale* (Busk, 1860)
- Family Tcuchoporidae**
- Phylactella*
 - labrosa* (Busk, 1854)
 - Teuchopora*
 - edwardsi* (Jullien, 1882)
- Family Trypostegidae**
- Trypostega*
 - claviculata* (Hincks, 1884)
 - venusta* (Norman, 1864)
- Family Umbonulidae**
- Hippopleurifera*
 - pulchra* (Manzoni, 1870)
 - Posterula*
 - sarsi* (Smitt, 1868)
 - Ragionula*
 - rosacea* (Busk, 1856)
 - Umbonula*
 - littoralis* Hastings, 1944
 - ovicellata* Hastings, 1944

- Family Watersiporidae**
Watersipora
aterrima (Ortmann, 1890)
complanata (Norman, 1864)
cucullata (Busk, 1854)
subovoidea (d'Orbigny, 1852)
subtorquata (d'Orbigny, 1852)
- Order Ctenostomatida**
- Family Aethozoontidae**
Aethozoon
pellucidum Hayward, 1978
- Family Alcyoniidae**
Alcyonium
albidum Alder, 1857
candidum Ryland, 1963
cellariooides Calvet, 1900
diaphanum (Hudson, 1778)
disciforme Smitt, 1872
duplex Prouho, 1892
gelatinosum (Linnaeus, 1761)
hirsutum (Fleming, 1828)
= several species
mamillatum Alder, 1857
mytili Dalyell, 1848
parasiticum (Fleming, 1828)
variegatum (Prouho, 1892)
- Family Arachnididae**
Arachnidium
clavatum Hincks, 1877
fibrosum Hincks, 1880
hippothooides Hincks, 1859
- Arachnoidea**
annosciae d'Hondt & Geraci, 1976
thalassae (d'Hondt, 1978)
- Tanganella*
muelleri Kraepelin, 1887
- Family Benedeniporidae**
Benedenipora
catenata Fergens, 1889
delicatula d'Hondt & Geraci, 1975
- Family Buskiidae**
Buschia
nitens Alder, 1856
socialis Hincks, 1887
- Family Clavoporidae**
Ascorhiza
mawatarii d'Hondt, 1983
- Clavopora*
hystricis (Busk, 1874)
- Metalcyonium*
gautieri d'Hondt, 1976
- Family Claviporidae**
Pseudalcyonidium
bobinæ d'Hondt, 1976
- Family Flustrellidae**
Flustrellidra
corniculata (Smitt, 1872)
hispida (O. Fabricius, 1780)
- Family Hypophorellidae**
Hypophorella
expansa Ehlers, 1876
- Family Immergentiidae**
Immergentia
suecica Silén, 1946
- Family Lobiancoporidae**
Lobiancopora
hyalina Pergens, 1889
- Family Mimosellidae**
Bantariella
verticillata (Heller, 1867)
- Mimosella*
- crosslandi* d'Hondt, 1983
gracilis Hincks, 1851
verticillata (Heller, 1867)
- Family Monohryzoontidae**
Monohryzozon
ambulans Remane, 1936
- Family Nolellidae**
Anguinella
palmata van Beneden, 1845
- Nolella*
dilatata (Hincks, 1860)
- Family Nollellidae**
gigantea (Busk, 1856)
- Family Nolellidae**
limicola (Franzén, 1960)
pusilla (Hincks, 1880)
stipata (Gosse, 1855)
- Panolicella*
nutans Jebram, 1985
- Family Penetrantiidae**
Penetrantia
brevis Silén, 1946
concharum Silén, 1946
- Family Pherusellidae**
Pherusella
tubulosa (Ellis & Solander, 1786)
- Family Spathiporidae**
Spathipora
sertum (Fischer, 1866)
- Family Terebriporidae**
Immergentia
orbigniana Fischer, 1866
- Spathipora*
comma (Soule, 1950)
- Family Triticellidae**
Farrella
repens (Farre, 1837)
- Triticella*
flava Dalyell, 1848
= *Triticella korenii* G.O. Sars
pedicellata (Alder, 1857)
- Triticellopsis*
tissieri Gautier, 1962
- Family Vesiculariidae**
Amathia
distrans Busk, 1886
guernseii Chimonides, 1987
intermedis Chimonides, 1987
lendigera (Linnaeus, 1758)
pruvoti Calvet, 1911
semiconvoluta (Lamouroux, 1824)
tortuosa Tenison-Woods, 1880
vidovici (Heller, 1867)
- Bowerbankia*
citrina (Hincks, 1877)
gracilis Leidy, 1855
gracillima (Hincks, 1877)
imbricata (Adams, 1798)
pustulosa (Ellis & Solander, 1786)
- Vesicularia*
spinosa (Linnaeus, 1967)
- Zoobotryon*
verticillatum (delle Chiaje, 1822)
- Family Victorellidae**
Bulbella
abscondita Braem, 1951
- Victorella*
muelleri (Kraepelin, 1887)
pavida Saville Kent, 1870
- Family Walkeriidae**
Walkeria

- atlantica* Busk, 1886
tuberosa Heller, 1867
uva (Linnaeus, 1758)
- Class Stenolaemata**
- Order Cyclostomatida**
- Family Annectocymidae**
- Annectocyma*
arcuata (Hartnelin, 1976)
indistincta Canu & Bassler, 1929
major (Johnston, 1847)
tubulosa (Busk, 1875)
- Entalophoroecia*
balgimae Harmelin & d'Hondt, 1992
clavata (Busk, 1859)
elegans (Norman, 1909)
gracilis Hartnelin, 1976
robusta Harmelin, 1976
deflexa (Couch, 1842)
producta (M. Sars, 1869)
- Family Crisiidae**
- Bicrisia*
abyssicola Kluge, 1862
gibraltarensis Harmelin, 1990
- Crisia*
aculeata Hassall, 1841
calyptostoma Hayward & Ryland, 1978
conferta Busk, 1875
constans Kluge, 1946
cuneata Maplestone, 1905
denticulata (Lamarck, 1816)
denticulata var. *arctica* M. Sars, 1863
eburnea (Linnaeus, 1758)
elongata (Milne Edwards, 1838)
fistulosa Heller, 1867
klugei Ryland, 1967
occidentalis (Trask, 1857)
oranensis Waters, 1916
pyrula Harmelin, 1990
ramosa Harmer, 1891
sigmoidea Waters, 1916
tenella Calvet, 1906
- Crisidia*
cornuta (Linnaeus, 1758)
- Crisiella*
producta (Smitt, 1865)
- Filicrisia*
geniculata (Milne Edwards, 1838)
smitti (Kluge, 1946)
- Family Diastoporidae**
- Cardioecia*
watersi (O'Donoghue & de Waterville, 1939)
- Diplosolen*
obelia (Johnston, 1838)
intricarius (Smitt, 1872)
- Eurystrota*
compacta (Norman, 1866)
occulta (Hartnelin, 1976)
- Liripora*
amphorae (Hartnelin, 1974)
violacea (Hartnelin, 1976)
- Plagioecia*
dorsalis (Waters, 1879)
inoedificata (Julien, 1882)
patina (Lamarck, 1816)
platydiscus Harmelin
sarniensis (Norman, 1864)
- Family Horneridae**
- Hornera*
frondiculata Lamouroux, 1821
lichenoides (Linnaeus, 1758)
- Family Lichenoporidae**
- Coronopora*
truncata (Fleming, 1928)
- Defrancia*
lucernaria M. Sars, 1851
- Disporella*
alboranensis Alvarez, 1992
borgi Alvarez, 1995
boutani Alvarez, 1995
harmeri (Neviani, 1939)
hispida (Fleming, 1828)
piramidata Alvarez, 1992
robusta Alvarez, 1992
smitti (Calvet, 1906)
spinulosa Jullien, 1888
zurigae Alvarez, 1992
- Lichenopora*
aztiensis Alvarez, 1994
distincta Alvarez, 1993
flosculus Hincks, 1862
mediterranea de Blainville
radiata (Audouin, 1826)
verrucaria (O. Fabricius, 1780)
- Family Mecynoeciidae**
- Mecynoecia*
delicatula (Busk, 1875)
- Family Oncousoeciidae**
- Oncousoecia*
canadensis (Osborn, 1933)
diastoporides (Norman, 1869)
dilatans (Johnston, 1847)
polygonalis (Kluge, 1915)
- Stomatopora*
semirecta Jullien, in Jullien & Calvet, 1903
- Stomatoporina*
incurvata (Hincks, 1859)
- Family Stegohorneridae**
- Stegohornera*
arctica (Kluge, 1946)
violacea (M. Sars, 1863)
- Family Terviidae**
- Tervia*
irregularis (Meneghini, 1844)
- Family Tubuliporidae**
- Idmidronea*
atlantica (Forbes, in Johnston, 1847)
bidenkapi (Kluge, 1955)
coerulea Harmelin, 1976
flexuosa (Pourtales, 1867)
- Platonea*
stoechas Harmelin, 1976
- Tubulipora*
aperta (Harmer, 1898)
flabellaris (O. Fabricius, 1780)
hemiphragmata Harmelin, 1976
liliacea (Pallas, 1766)
lobifera Hastings, 1963
notomale (Busk, 1875)
penicillata (O. Fabricius, 1780)
phalangea Couch, 1844
plumosa Thomson, in Hartner, 1898
ziczac Harmelin, 1976

BRACHIOPODA

Compiled by C. M. Howson

The phylum Brachiopoda is a group of lophophorate organisms which have in the past frequently been classed with the phoronids and bryozoans in the phylum Lophophorata. The brachiopods are well represented in the fossil record, and have a long evolutionary history; there are many thousand fossil species known, but only about 300 living species. Their systematics have thus primarily been the province of paleontologists, and the resulting classification, reflecting the complex phylogeny of the group, is based on the large amount of descriptive work available on both fossil and recent species. The classification used in this checklist follows Ryland (1990) who provides good illustrations and descriptions of many of the British species.

This project has not been able to compile a list of living marine brachiopods for Europe. However, the list produced for Britain, Ireland and adjacent seas by Howson (1997) is considered a valuable starting point. This is the provided here. Indeed, it includes all the species recorded for Norwegian waters by Thomsen and Brattegard (1996).

The taxonomy of recent brachiopods and the species known from the British Isles have changed little since Brunton & Curry (1979), and the checklist is therefore based on this publication.

References

- Brunton, C.H.C. & Curry, G.B. 1979. *British Brachiopods*. Synopses of the British Fauna (N.S.), No. 17. 64 pp. Linnean Society, London and E.B.S.A. Academic Press, London.
- Curry, G.B. 1982. Ecology and population structure of the recent brachipod *Terebratulina* from Scotland. *Palaeontology* 25(2), 227 - 246.
- Curry, G.B. 1983. Ecology of the recent deep water rhynchonellid brachiopod *Cryptopora* from the Rockall Trough. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 44, 93 - 102.
- Lee & Brunton, C.H.C. 1986. *Neocrania* n. gen., and a revision of Cretaceous - recent brachiopod genera in the family Craniidae. *Bulletin of the British Museum of Natural History (Geol.)* 40(4), 141 - 160.
- Ryland, J.S. 1990. Lophophorate phyla. In: *The marine fauna of the British Isles and North West Europe*. P.J. Hayward & J. S. Ryland, eds, 2: 794-838, Clarendon Press, Oxford.
- Thomsen, E. & Brattegard T. 1997. Brachiopoda. In: Brattegard T. and Holthue T., *Distribution of marine, benthic macro-organisms in Norway*. Research Report for DN 1997-1, Directorate for Nature Management, 296-297 pp.

SUBPHYLUM CRANIIFORMEA

Class Craniata

Order Craniida

- Family Craniidae**
Neocrania anomala (O. F. Müller 1776)

SUBPHYLUM RHYNCHONELLIFORMEA

Class Rhynchonellata

Order Rhynchonellida

- Family Cryptoporidae**
Cryptopora gnomon Jeffreys 1869

- Family Hemithirididae**
Hemithiris psittacea (Gmelin 1790)

Family Frieleiidae

- Hispanirhynchia cornea* (Davidson 1887)

Order Terebratulida

- Family Terebratulidae**
Gryphus vitreus (Born 1778)

Family Cancellothyrididae

- Terebratulina retusa* (Linnaeus 1758)
septentrionalis (Couthouy 1838)

Family Megathyrididae

- Argyrotheca cistellula* (Searles-Wood 1841)
Argyrotheca cuneata (Risso 1826)

Family Gwynioidea

- Gwynia capsula* (Jeffreys 1859)
Megathiris detruncata (Gmelin 1790)

Family Platidiidae

- Platidia anomioides* (Scacchi & Philippi 1844)
davidsoni (Deslongchamp 1885)

Family Kraussinidae

- Megerlia truncata* (Linnaeus 1767)

Family Dallinidae

- Dallina septigera* (Lovén 1846)
Fallax dalliniiformis Atkins 1960
Glaciarcula spitzbergensis (Davidson 1852)

Family Macandreviidae

- Macandrevia cranium* (O. F. Müller 1776)

CHAETOGNATHA

Compiled by Jacob van der Land and Helga Kapp

The European list was derived from URMO, which is mainly based on Bieri (1991).

References

Bieri, R., 1991. Systematics of the Chaetognatha. In: Q. Bone, et al. *The Biology of Chaetognaths*. Oxford Univ. Press: p.122-136.

PHYLUM CHAETOGNATHA

Family Eukrohniidae

<i>Eukrohnia</i>	
<i>bathypelagica</i> Alvarino, 1962	A M
<i>fowleri</i> Ritter-Zahony, 1909	A M
<i>hamata</i> (Möbius, 1875)	A M
<i>Heterokrohnia</i>	
<i>curvichaeta</i> Casanova, 1986	A
<i>davidii</i> David, 1985	A
<i>discovery</i> Casanova, 1994	
<i>furnestinae</i> Casanova & Chidgey, 1987	A
<i>heterodonta</i> Casanova, 1986	A
<i>mirabilis</i> Ritter-Zahony, 1911	A M
<i>murina</i> Casanova, 1986	A
<i>palpifera</i> (Casanova, 1986)	M
<i>rubra</i> (Casanova, 1986)	A
<i>sorbei</i> (Casanova, 1993)	

Family Krohnittellidae

<i>Krohnittella</i>	
<i>boureei</i> Germain & Joubin, 1912	

Family Krohnittidae

<i>Krohnitta</i>	
<i>pacifica</i> (Aida, 1897)	
<i>subtilis</i> (Grassi, 1881)	

Family Pterosagittidae

<i>Pterosagitta</i>	
<i>draco</i> (Krohn, 1853)	A M
= <i>Spadella draco</i> Krohn, 1853	

Family Sagittidae

<i>Caecosagitta</i>	
<i>macrocephala</i> (Fowler, 1904)	A M
= <i>Sagitta macrocephala</i> Fowler, 1904	
<i>Ferosagitta</i>	
<i>hispida</i> (Conant, 1895)	A
= <i>Sagitta hispida</i> Conant, 1895	
<i>paulilata</i> kassatkina, 1971	
= <i>Sagitta tokioi</i> Alvarino, 1967 ??	
<i>Flaccisagitta</i>	
<i>enflata</i> (Grassi, 1881)	A M
= <i>Sagitta enflata</i> Grassi, 1881	
<i>hexaptera</i> (d'Orbigny, 1836)	A M
= <i>Sagitta hexaptera</i> d'Orbigny, 1836	
<i>Mesosagattia</i>	
<i>decipiens</i> (Fowler, 1905)	A M
= <i>Sagitta decipiens</i> Fowler, 1905	
<i>minima</i> (Grassi, 1881)	A M
= <i>Sagitta minima</i> Grassi, 1881	
<i>neodecipiens</i> (Tokioka, 1959)	
<i>sibogae</i> (Fowler, 1906)	A M
<i>Parasagitta</i>	
<i>elegans</i> (Verrill, 1873)	A
= <i>Sagitta elegans</i> Verrill, 1873	
<i>friderici</i> (Ritter-Zahony, 1911)	A
= <i>Sagitta friderici</i> Ritter-Zahony, 1911	
<i>megalophthalma</i> (Dallot & Ducret, 1969)	
<i>setosa</i> (Müller, 1847)	A
= <i>Sagitta batava</i> (Biersteker & van der Spoel, 1966)	
<i>Pseudosagitta</i>	
<i>lyra</i> (Krohn, 1853)	A M
= <i>Sagitta lyra</i> Krohn, 1853	
<i>maxima</i> (Conant, 1896)	A M
= <i>Sagitta maxima</i> Conant, 1896	
<i>Sagitta</i>	

<i>abyssicola</i> Chidgey, 1989	
<i>bipectinata</i> Quoy & Gaimard, 1828	A M
<i>zetesios</i> (Fowler, 1905)	
<i>Serratosagitta</i>	
<i>serratodentata</i> (Krohn, 1853)	A M
= <i>Sagitta serratodentata</i> Krohn, 1853	
<i>tasmanica</i> (Thompson, 1947)	A M
= <i>Sagitta tasmanica</i> Thompson, 1947	
<i>Solidosagitta</i>	
<i>planctonis</i> (Steinhaus, 1896)	A M
= <i>Sagitta planctonis</i> Steinhaus, 1896	
Family Spadellidae	
<i>Spadella</i>	
<i>birostrata</i> Casanova, 1987	
<i>equidentata</i> Casanova, 1987	
<i>ledoyerii</i> Casanova, 1986	
<i>cephaloptera</i> (Busch, 1851)	A M

HEMICORDATA

Compiled by Jacob van der Land.

The European list was derived from URMO with names provided by J. Hallan. No expert has checked the accuracy of the ERMS list.

PHYLUM HEMICORDATA

Class Enteropneusta

Family Harrimaniidae

<i>Harrimania</i>	
<i>kupfferi</i> (von Willemoes-Suhm, 1871)	A
<i>Saccoglossus</i>	
<i>horsti</i> Brambell & Goodhart, 1941	A
<i>kowalevskii</i> (Agassiz, 1873)	A
<i>mereschkowskii</i> (Wagner, 1885)	A M
<i>pygmaeus</i> Hinrichs & Jacobi, 1938	A
<i>ruber</i> Tattersall, 1905	A
<i>Stereobalanus</i>	
<i>canadensis</i> (Spengel, 1893)	A
<i>Protoglossus</i>	
<i>koehleri</i> (Caulery & Mesnil, 1900)	A

Family Ptychoderidae

<i>Balanoglossus</i>	
<i>clavigerus</i> delle Chiaje, 1829	A M
<i>Glossobalanus</i>	
<i>elongatus</i> Spengel, 1904	M
<i>marginatus</i> Meek, 1922	A
<i>minutus</i> Kowalevsky, 1866	A M
<i>sarniensis</i> (Koehler, 1886)	A

Family Spengelidae

<i>Glandiceps</i>	
<i>talaboti</i> (Marion, 1876)	M

Family incertae sedis

<i>Planctosphaera</i>	
<i>pelagica</i> (Spengel, 1932)	A

Class Pterobranchia

Family Rhabdopleuridae

<i>Rhabdopleura</i>	
<i>compacta</i> Hincks, 1880	A
<i>normani</i> Allmann, 1869	A M

ECHINODERMATA

Compiled by Hans G. Hansson

The list of European Echinodermata was compiled by Hans G. Hansson. It used the Echinoderm part of the compilers NEAT lists (available via Internet at address <http://www.tmbi.gu.se/libdb/taxon/taxa.html>) as a starting point. NEAT (North East Atlantic Taxa) are preliminary check-lists, covering Scandinavian marine animal taxa. The chapter by F.J. Fell on Echinodermata in Parker (1982) was used as the systematic framework. The asteroids are mainly listed from the recent monograph on Atlantic species by Clark & Downey (1992). Mortensen (1927) is used as a basal source for the other groups. Additions from several other sources have been made, and the electronic part of the Zoological Record (covering recent decades) has been utilised. However, examination of the reports from the large expeditions (except the Norwegian North Atlantic and the Ingolf) and other literature since WWII, may add to the species list.

The compiler is generally interested in marine fauna, but is not a specialist on any Echinoderm taxon, so there are certainly several omissions and lapses in this draft list. The assistance of experts in groups of Echinodermata was thus invaluable. Dr Sabine Stöhr, Stockholm, assisted with ophiuroid names, Dr Claude Massin, Brussels, assisted with the holothurians and added 9 omitted species of this taxon to the list. Dr Andrey Gebruk, Moscow, together with his friends Alexander Mironov and Alexei Smirnov kindly helped with several corrections and additions to the holothurians, the crinoids, the echinoids and the genus *Caulaster* among the asteroids. Prof. Dusan Zavodnik, Rovinj, Croatia, provided several corrections, new distribution notes (particularly for the Adriatic Sea area), information about synonyms, and added a few omitted species. Dr Mateo Garrido, Gran Canaria, provided additional records from the Canaries.

Authors of most genus and species names in the list below have been found, and most often also the years of description, but with several exceptions. Also, some of the years of authority in this list may be wrong, because different sources often have different opinions regarding which year a certain species is described (e.g. gen. *Poliometra* Clark; the original publication: 1923, Neave: 1924, McKenzie in Howson & Picton 1997: 1913; in this case the year from the original publication must be correct). Thus, further study of the primary literature may improve the accuracy of this list.

References

- Belyaev G.M. & A.N. Mironov, 1996. Starfishes Porcellanaster and *Caulaster* (Porcellanasteridae) from the Atlantic and the Antarctic. *Zoologichesky zhurnal*, 75 (6): 886 - 899 (In Russian, English summary).
- Brattegard, T & T. Holthe (Eds.). 1997. *Distribution of marine, benthic macro-organisms in Norway. A tabulated catalogue*. Preliminary edition. Research Report for DN, Nr. 1997-1. Trondheim. 409 pp. (Echinoderm chapters by T. Høisæter, T Brattegard & J.-A. Snell, pp. 301-315).
- Clark, A.H. 1923. *Crinoidea*. The Danish Ingolf Expedition. Vol. IV. Luno, Copenhagen. 58 pp.
- Clark, A.M. 1970. Echinodermata Crinoidea. *Marine Invertebrates of Scandinavia*. 3. Universitetsforlaget, Oslo. 55 pp.
- Clark, A.M. & M.E. Downey 1992. *Starfishes of the Atlantic*. Chapman & Hall, London. 794 pp.

- D'yakonov, A.M. 1967. *Ophiuroids of the USSR Seas*. Israel Program for Scientific Translations, Jerusalem 123 pp.
- D'yakonov, A.M. 1968. *Sea Stars (Asteroids) of the USSR Seas*. Israel Program for Scientific Translations, Jerusalem 183 pp.
- D'yakonov, A.M. 1969. Echinodermata, Vol. I. Sea-Urchins (Echinoidea) No. I. *Fauna of Russia and adjacent Countries*. Israel Program for Scientific Translations, Jerusalem 265 pp.
- Fell, H.B. 1960. Synoptic Keys to the Genera of Ophiuroidea. *Zoology Publications from Victoria University of Wellington* No. 26. 34 pp + literature index.
- Fiege, D. & Y. Liao 1996. *Penipidida ludwigi* (Marenzeller, 1893) (Holothurioidea: Elpidiidae) rediscovered from the Eastern Mediterranean Sea. In: Uiblein, F., Ott, J., Stachowitsch, M. (Eds) Deep-sea and extreme shallow-water habitats: affinities and adaptations. *Biosystematics and Ecology Series*, 11, 61-66.
- Gebruk, A.V. 1989. Revision of the family Pelagothuriidae (Holothurioidea, Elasipodida) with a review of swimming holothurians. Part I. Revision of Pelagothuriidae. *Zool. Zhurnal* 68 (12): 57-66. (In Russian, English summary).
- Gebruk, A.V., 1990. *Deep-sea holothurians of the family Elpidiidae*. Moscow, Nauka, 160 p. [In Russian].
- Gulliksen, B., Palerud, R., Brattegård, T. & J.-A. Snell, 1999. Distribution of marine benthic macro-organisms at Svalbard (including Bear Island) and Jan Mayen. Research report for DN 1999 - 4. 148 pp. Trondheim, Norway.
- Hansen, B. 1975. Systematics and biology of the deep-sea holothurians. *Galathea Report*, 13, 1-262.
- Heding, S.G. 1935. *Holothurioidea I. The Danish Ingolf Expedition*. Vol. IV. Hagerup, Copenhagen. 84 pp + 8 plates.
- Heding, S.G. 1935 Echinoderms. *Meddelelser om Grønland*. Bd. 104, Nr. 13. 68 pp + 1 plate.
- Heding, S.G. 1936 Echinoderms. *Meddelelser om Grønland*. Bd. 108, Nr. 1. 34 pp.
- Heding, S.G. 1942. *Holothurioidea II. The Danish Ingolf Expedition*. Vol. IV. Hagerup, Copenhagen. 39 pp + 2 plates.
- Heding, S.G. & A. Panning 1954. *Phyllophoridae*. Spolia Zooligica Musei Hauniensis XIII. 209 pp.
- Howson, C.M. & B.E. Picton 1997 (Eds.). *The Species Directory of the Fauna and Flora of the British Isles and Surrounding Seas*. The Ulster Museum, Belfast & The Marine Conservation Society, Ross-on-Wye. 508 pp. (Echinoderm chapter by J.D. McKenzie, pp. 287-295).
- Høisæter, T. 1990. An annotated check-list of the echinoderms of the Norwegian coast and adjacent waters. *Sarsia* 75 : 83-106
- Kier P.M. & M. H. Lawson, 1978. Index of Living and Fossil Echinoids 1924-1970. *Smithson. Contrib. Paleobiol.* 34 : vi+182 pp. (not yet seen)
- Madsen, F.J. 1947. The Echinoderms collected by the "Skagerak" expedition in the eastern Atlantic 1946. I. Asteroidea, Ophiuroidea, Echinoidea, and Holothuroidea. *Göteborgs Kungl. Vetenskaps- och Vitterhets-Samhälles Handlingar*, Sjätte följdjen. Ser. B, Band 5. N:o 7. 16 pp.
- Madsen, F.J. 1987. The *Henricia sanguinolenta* complex (Echinodermata, Asteroidea) of the Norwegian Sea and adjacent waters. A re-evaluation, with notes on related species. *Steenstrupia* 13 (5), pp. 201-268.
- Madsen, F.J. & B. Hansen 1994. Echinodermata. Holothurioidea. *Marine Invertebrates of Scandinavia*. 9. Scandinavian University Press. 143 pp.
- Miller, J.E and Pawson, D.L. 1990. Swimming sea cucumbers (Echinoderata:Holothuroidea): a survey, with analysis of swimming behaviour in four bathyal species. *Smithsonian Contributions to Marine Sciences*, 35, 18p.
- Mironov A.N., 1974. Pourtalesiid sea urchins of the Antarctic and Subantarctic (Echinoidea: Pourtalesiidae). *Transactions of the P.P. Shirshov Institute of oceanology*, 98: 240 - 252. (In Russian, English summary).
- Mironov A.N., 1985. Circumcontinental zonation in distribution of Atlantic echinoids. *Transactions of the P.P. Shirshov Institute of oceanology*, 120, 70-95. (In Russian, English summary).
- Mironov A.N. 1995. Holasteroid echinoids. 2. *Pourtalezia*. *Zoologichesky zhurnal*, 74 (12), 59-75. (In Russian, English summary).
- Mironov A.N. 1997. Holasteroid echinoids. 4. *Echinosigra*. *Zoologichesky zhurnal*, 76, (2): 173-186. (In Russian, English summary).
- Mortensen, T. 1903. *Echinoidea I. The Danish Ingolf Expedition*. Vol. IVa. Luno, Copenhagen. 183 pp + 21 plates.
- Mortensen, T. 1907. *Echinoidea II. The Danish Ingolf Expedition*. Vol. IVa. Luno, Copenhagen. 200 pp + 19 plates.
- Mortensen, T. 1927. *Handbook of the Echinoderms of the British Isles*. Clarendon Press, Oxford. 471 pp.
- Mortensen, T. 1933. *Ophiuroidea. The Danish Ingolf Expedition*. Vol. IV. Luno, Copenhagen. 121 pp + 3 plates.
- Neave, S.A. 1939-50. *Nomenclator Zoologicus*. Vol. I-5. The Zoological Society of London. + M.A. Edwards & A.T. Hopwood (Eds.) 1966 Vol. 6 + M.A. Edwards & H.G. Vevers (Eds.) Vol. 7.
- Parker, S.P. (Ed.) 1982. *Synopsis and Classification of Living Organisms*. 2 Vols. McGraw-Hill, New York. 1166 + 1232 pp.
- Paterson, G.L.J. 1985. The deep-sea Ophiuroidea of the north Atlantic Ocean. *Bulletin of the British Museum (Natural History). Zoology*, 49 (1). 162 pp. (not seen)
- Pereira, Mariano Olivéiro Rego 1997. Checklist of the littoral echinoderms of the Azores. *Açoreana* 8 (3): 331-337.
- Picton, B.E. 1993. *A Field Guide to the Shallow-water Echinoderms of the British Isles*. Immel Publishing. 96 pp.
- Sherborn, C.D. 1902. *Index Animalium sive index nominum quae ab a.d. MDCCCLVIII generibus et speciebus animalium imposita sunt. Sectio prima a kalendis Ianuariis, MDCCCLVIII usque ad finem Decembris, MDCCC. C. J. Clay & Sons, Cambridge Univ. Press, London. 1195 pp.*
- Sherborn, C.D. 1922-32. *Index Animalium sive index nominum quae ab a.d. MDCCCLVIII generibus et speciebus animalium imposita sunt. Sectio secunda a kalendis Ianuariis, MDCCCI usque ad finem Decembris, MDCCCL*. The Trustees of the British Museum, London. 7056 + 1098 pp.
- Tortesone, E. 1965. Echinodermata. *Fauna d'Italia*. Vol. 6. Edizioni Calderini, Bologna. 422 pp.

PHYLUM ECHINODERMATA

Class Crinoidea

Order Millericrinida

Family Hyocrinidae

Gephyrocrinus grimaldii Koehler & Bather, 1902

Order Cyrtocrinida

Family Holopodidae

Cyathidium foresti Cherbonnier & Guille, 1972
= *C. meteorensis* Fecheter, 1973

Order Bourgueticrinida

Family Bourgueticrinidae

Conocrinus cabioshi Roux, 1976
cherbonnieri Roux, 1976

Democrinus

parfaiti Perrier, 1883
= *Rhizocrinus magnus* Gislén, 1947
= *Rhizocrinus rawsoni* Koehler, 1909, non Pourtalès, 1874
conifer (A.H. Clark, 1909)
= *Bythocrinus intermedius* A.H. Clark, 1915
= *Rhizocrinus conifer* A.H. Clark, 1909
= *Rhizocrinus robustus* A.H. Clark, 1909
rawsonii (Pourtalès, 1874)
= *Rhizocrinus rawsonii* Pourtalès, 1874
Rhizocrinus

- Iosotensis* M. Sars, 1868
 = *R. verrilli* A.H. Clark, 1908
- Family Bathycrinidae**
- Bathycrinus*
aldrichianus Wyville Thomson, 1877
 = *B. serratus* A.H. Clark, 1908
carpenterii (Danielssen & Koren, 1877)
 = *Illycrinus carpenterii* Danielssen & Koren, 1877
gracilis Wyville Thomson, 1877
- Monachocrinus*
perrieri (Koehler & Vaney, 1910)
 = *Bathycrinus perrieri* Koehler & Vaney, 1910
 = *M. coelus* H.L. Clark, 1923
recuperatus (Perrier, 1885)
 = *Illycrinus recuperatus* Perrier, 1885
sexradiatus A.H. Clark, 1919
- Family Phryncrinidae**
- Zeuctocrinus*
gisleni A.M. Clark, 1973
- Family Porphyrocrinidae**
- Porphyrocrinus*
incrassatus (Gislen, 1933)
thalassae Roux, 1977
- Order Isocrinida**
- Family Isocrinidae**
- Diplocrinus*
(Annocrinus) wyvillethomsoni (Jeffreys, 18???)
 = *Pentacrinus wyville-thomsoni* Jeffreys, 18???
- Order Comatulida**
- Family Pentametrocrinidae**
- Pentametrocrinus*
atlanticus (Perrier, 1883)
 = *Eudicocrinus atlanticus* Perrier, 1883
- Thaumatacrinus*
jungensi A.H. Clark, 1923
- Family Antedonidae**
- Antedon*
bifida (Pennant, 1777)
 = ? *Antedon duebeni* Böhlse, 1866
 = *A. gorgia* de Freminville, 1811
 = *A. maroccana* A.H. Clark, 1910
 = *Asterias bifida* Pennant, 1777
mediterranea (de Lamarck, 1816)
 = *A. adriatica* A.H. Clark, 1911
 = *Comatula annulata* Risso, 1826
 = *Comatula mediterranea* de Lamarck, 1816
petasus (Düben & Koren, 1846)
 = *Alecto petasus* Düben & Koren, 1846
- Hathrometra*
sarsi (Düben & Koren, 1846)
 = *Antedon tenella* Carpenter non (Retzius)
 = *H. tenella* (Carpenter, 1888) non (Retzius, 1783)
- Heliometra*
glacialis (Owen, 1833 ex Leach MS)
 = *Alecto eschrichtii* J. Müller, 1841
 = *Alecto glacialis* Owen, 1833 ex Leach MS
 = *Antedon quadrata* P.H. Carpenter, 1884
- Leptometra*
celtica (M'Andrew & Barrett, 1858)
 = *Comatula celtica* M'Andrew & Barrett, 1858
 = *Comatula woodwardi* Barrett, 1857, non Forbes, 1852
 = *L. phalangium* (Carpenter, 1881) non (J. Müller, 1841)
phalangium (J. Müller, 1841)
 = *Alecto phalangium* J. Müller, 1841
- Meteorometra*
monticola A.M. Clark, 1980
- Orthometra*
hibernica (A.H. Clark, 1913)
 = *Trichometra hibernica* A.H. Clark, 1913
- Poliometra*
prolixa (Sladen, 1881)
 = *Antedon prolixa* Sladen, 1881
 = *Hathrometra prolixa* (Sladen)
- Trichometra*
cubensis (Pourtalès, 1869)
 = *Antedon cubensis* Pourtalès, 1869
 = *T. aspera* A.H. Clark, 1908
delicata A.H. Clark, 1911
- Thaumatometra*
septentrionalis A.H. Clark, 1923
- Family Atelecrinidae**
- Atelecrinus*
balanooides P.H. Carpenter, 1881
 = *A. helgae* A.H. Clark, 1913
- Family Comasteridae**
- Neocomatella*
europaea A.H. Clark, 19??
- Family Thalassometridae**
- Koehlermetra*
flava (Koehler)
 = *Antedon flava* Koehler
orrecta (P.H. Carpenter, 18???)
 = *Antedon porrecta* P.H. Carpenter, 18???
- Thalassometra*
lusitanica (P.H. Carpenter, 1884) A.M. Clark, 1980
 = *Antedon lusitanica* P.H. Carpenter, 1884
 = *Stiremetra lusitanica* (P.H. Carpenter, 1884)
omissa (Koehler)
 = *Antedon omissa* Koehler
- Class Asteroidea**
- Order Paxillosida**
- Family Astropectinidae**
- Astropecten*
aranciacus (Linnaeus, 1758)
 = *Asterias aranciaca* Linnaeus, 1758
 = *Asterias aurantiaca* Tiedemann, 1816
 = *A. antarcticus* Studer, 1876
 = *A. crenaster* Dujardin & Hupé, 1862
 = *A. gruveli* Koehler, 1912
 = *A. meridionalis* Studer, 1876
 = *A. perarmatus* Perrier, 1869
bispinosus (Otto, 1823)
 = *Asterias bispinosa* Otto, 1823
 = *A. myosurus* Perrier, 1869
hermatophilus Sladen, 1883
ibericus Perrier, 1894
irregularis subsp. *irregularis* (Pennant, 1777)
 = *Asterias irregularis* Pennant, 1777
 = *A. acicularis* Norman, 1865
 = *A. echinulatus* J. Müller & Troschel, 1844
 = *A. helgolandicus* Greeff, 1871
 = *A. muelleri* J. Müller & Troschel, 1844
irregularis subsp. *pentacanthus* (Delle Chiaje, 1827)
 = *Asterias pentacantha* Delle Chiaje, 1827
- jonstoni* (Delle Chiaje, 1827)
 = *Asterias jonstoni* Delle Chiaje, 1827
 = *A. aster* De Filippi, 1859
 = *A. squamatus* J. Müller & Troschel, 1844
platyacanthus (Philippi, 1837)
 = *Asterias platyacantha* Philippi, 1837
spinulosus (Philippi, 1837)
 = *Asterias spinulosa* Philippi, 1837
- Bathybiaster*
vexillifer (W. Thomson, 1873)
 = *Archaster robustus* Verrill, 1884
 = *Archaster vexillifer* W. Thomson, 1873
 = *Astropecten pallidus* Danielssen & Koren, 1877
 = *Illyaster mirabilis* Danielssen & Koren, 1884
 = *Phoxaster pumilus* Sladen, 1889
- Dytaster*
grandis subsp. *grandis* (Verrill, 1884)
 = *Archaster grandis* Verrill, 1884
 = *D. agassizi* Perrier, 1894
 = *D. biserialis* Sladen, 1889
 = *D. exilis* var. *carinata* Sladen, 1889
 = *D. madreporeif* Sladen, 1889

- = *D. rigidus* Perrier, 1894
intermedius Perrier, 1891
mollis (Perrier, 1885)
= *Crenaster mollis* Perrier, 1885
semispinosus (Perrier, 1894)
= *Crenaster semispinosus* Perrier, 1894
= *D. parvulus* Koehler, 1909
- Leptychaster**
arcticus (M. Sars, 1851)
= *Astropecten arcticus* M. Sars, 1851
= *Astropecten lutkeni* Barrett, 1857
- Lonchotaster**
tartareus Sladen, 1889
- Persephonaster**
leptactis H.L. Clark, 1941
patagiatus (Sladen, 1889)
= *P. spinulosus* H.L. Clark, 1941
= *Psilaster patagiatus* Sladen, 1889
sphenoplax (Bell, 1892)
= *Astropecten sphenoplax* Bell, 1892
= *Psilasteropsis humilis* Koehler, 1907
- Plutonaster**
agassizi subsp. *notatus* Sladen, 1889
= ?*Gonipecten edwardsi* Perrier, in Milne-Edwards, 1882
= ?*Gonipecten inermis* Perrier, 1885 (*nomen nudum*)
= *P. abbreviatus* Sladen, 1889
= *P. granulosus* Perrier, 1891
= *P. notatus* Sladen, 1889
bifrons (Wyville Thomson, 1873)
= *Archaster bifrons* Wyville Thomson, 1873
= *P. proteus* H.L. Clark, 1923
- Psilaster**
andromeda subsp. *andromeda* (J. Müller & Troschel, 1842)
= *Astropecten andromeda* J. Müller & Troschel, 1842
= *Astropecten cristii* Düben & Koren, in Düben, 1845
cassiope Sladen, 1889
= ?*Archaster pulcher* Perrier, 1881
= *P. andromeda* var. *cassiope* Grieg, 1907
= *P. squameus* H.L. Clark, 1941
- Tethyaaster**
subinermis (Philippi, 1837)
= *Asterias subinermis* Philippi, 1837
- Family Luidiidae**
Luidia
atlantidea Madsen, 1950
ciliaris (Philippi, 1837)
= *Asterias imperati* Delle Chiaje, 1841
= *L. fragilissima* Forbes, 1839
sarsi subsp. *sarsi* Düben & Koren, in Düben, 1845
= *Astrella simplex* Perrier, in Milne-Edwards, 1882
= *L. africana* Sladen, 1889
- Family Ctenodiscidae**
Ctenodiscus
crispatus (Retzius, 1805)
= *Asterias arancia* Dewhurst, 1834
= *Asterias crispa* Retzius, 1805
= *Asterias polaris* Sabine, 1824
= *C. corniculatus* Perrier, 1875
= *C. pygmaeus* J. Müller & Troschel, 1842
- Family Radiasteridae**
Radiaster
tizardi (Sladen, 1882)
= *Mimaster tizardi* Sladen, 1882
- Family Porcellanasteridae**
Caulaster
eremicus (Sladen, 1889)
= *Albatrotrossaster richardi* Koehler, 1909
= *Porcellanaster eremicus* Sladen, 1889
pedunculatus Perrier, 1882
- Hyphalaster**
inermis Sladen, 1883
= *H. antonii* Perrier, 1885
- = *H. fortis* Koehler, 1907
= *H. gracilis* Koehler, 1907
= *H. parfaiti* Perrier, 1885
- Porcellanaster**
ceruleus Wyville Thomson, 1877
= *Albatrossia semimarginalis* Ludwig, 1905
= *Caulaster sladeni* Perrier, 1885
= *P. granulosus* Perrier, 1885
= *P. inermis* Perrier, 1885
= *P. irregularis* Lieberkind, 1932
- Stylocaster**
armatus Sladen, 1883
= *Machairaster pictus* Perrier, 1884
= *S. edwardsi* Perrier, 1885
= *S. spinosus* Perrier, 1885
chuni Ludwig, 1907
elongatus Koehler, 1907
horridus Sladen, 1883
= *Chunaster scapanophorus* Ludwig, 1907
- Thoracaster**
cylindratus Sladen, 1883
= *Pseudaster cordifer* Perrier, 1885
= *T. magnus* Ludwig, 1907
- Order Notomyotida**
- Family Benthopectinidae**
Benthopecten
simplex subsp. *simplex* (Perrier, 1881)
= *Archaster simplex* Perrier, 1881
= *B. spinosus* Grieg, 1918 (p.p.), non Verrill, 1889
= *Pararchaster armatus* Sladen, 1889
spinossissimus (Sladen, 1889)
= *Pararchaster spinosissimus* Sladen, 1889
- Cheiraster**
sepitus (Verrill, 1885)
= *Archaster sepitus* Verrill, 1885
= *Luidiaster dubius* H.L. Clark, 1941
= *Pectinaster dispar* Verrill, 1915
= *Pectinaster echinulatus* Ludwig, 1910 (p.p.)
= *Pontaster perplexus* Perrier, 1894
= *Pontaster venustus* Sladen, 1889
- Pectinaster**
filholi Perrier, 1885
= *Pontaster forcipatus* Sladen, 1889
= *Pontaster pristinus* Sladen, 1889
- Pontaster**
tenuispinus (Düben & Koren, 1846)
= *Astropecten tenuispinus* Düben & Koren, 1846
= *Crenaster marionis* Perrier, 1885
= *P. hebitus* Sladen, 1889
= *P. limbatus* Sladen, 1889
- Order Valvatida**
- Family Chaetasteridae**
Chaetaster
longipes (Retzius, 1805)
= *Asterias longipes* Retzius, 1805
= *Asterias subulata* de Lamarck, 1816
= *Nepanthia tessellata* Gray, 1840
- Family Odontasteridae**
Hoplasaster
spinosa Perrier, in Milne-Edwards, 1882
= *Pentagonaster lepidus* Sladen, 1889
- Odontaster**
mediterraneus (von Marenzeller, 1893)
= *Gnathaster mediterraneus* von Marenzeller, 1893
- Family Goniasteridae**
Ceramaster
granularis subsp. *granularis* (Retzius, 1783)
= *Asterias granularis* Retzius, 1783
= *Asterias tessellatus* de Lamarck, 1816
= *Astrogonium boreale* Barrett, 1857
= *Pentagonaster eximius* Verrill, 1894
= *Pentagonaster simplex* Verrill, 1895
grenadensis subsp. *grenadensis* (Perrier, 1881)

- = *Astrogonium greeni* Bell, 1889
 = *Pentagonaster affinis* Perrier, 1884
 = *Pentagonaster balteatus* Sladen, 1891
 = *Pentagonaster crassus* Perrier, 1885
 = *Pentagonaster deplasi* Perrier, 1885
 = *Pentagonaster gosselini* Perrier, 1884
 = *Pentagonaster grenadensis* Perrier, 1881
 = *Pentagonaster haesitans* Perrier, 1885
 = *Pentagonaster hystericis* von Marenzeller, 1893
 = *Pentagonaster kerghoensi* Koehler, 1895
- Evoplosoma**
scorpio Downey, 1981
- Goniaster**
tessellatus (de Lamarck, 1816)
 = *Asterias granularis* Gmelin, 1791 (p.p.)
 = *Asterias tessellata* de Lamarck, 1816
 = *Astrogonium dubium* Perrier, 1869
 = *G. africanus* Verrill, 1871
 = *G. americanus* Verrill, 1871
 = *G. cuspidatus* Gray, 1840
 = *G. sebae* Gray, 1840
- Hippasteria**
phrygiana subsp. *phrygiana* (Parelius, 1768)
 = *Asterias equestris* de Lamarck, 1816, non Retzius, 1805
 = *Asterias johnstoni* Gray, in Johnston, 1836
 = *Asterias phrygianus* Parelius, 1768
 = *Astrogonium aculeatum* Barrett, 1857
 = *Goniaster abbensis* Gray, 1843
 = *H. (Nehippasteria) insignis* Dons, 1938
 = *H. cornuta* Gray, 1840
 = *H. europaea* Gray, 1840
 = *H. plana* Gray, 1840
 = *H. sebae* Gray, 1840
- insignis* Dons, 1937
- Mediaster**
bairdi subsp. *bairdi* (Verrill, 1882)
 = *Archaster bairdi* Verrill, 1882
 = *M. stellatus* Perrier, 1891
- Nymphaster**
arenatus (Perrier, 1881)
 = *Dorigona jacqueti* Perrier, 1894
 = *Dorigona prehensilis* Perrier, 1885
 = *N. albidus* Sladen, 1889
 = *N. basilicus* Sladen, 1889
 = *N. protentus* Sladen, 1889
 = *Pentagonaster arenatus* Perrier, 1881
 = *Pentagonaster subspinosus* Perrier, 1881
 = *Pentagonaster ternalis* Perrier, 1881
- Paragonaster**
subtilis (Perrier, 1881)
 = *Archaster formosus* Verrill, 1884
 = *Goniopecten subtilis* Perrier, 1881
 = *P. cylindratus* Sladen, 1889
 = *P. strictus* Perrier, 1894
 = *Pentagonaster elongatus* Perrier, 1885
- Peltaster**
placenta (J. Müller & Troschel, 1842)
 = *Goniaster nidarosiensis* Storm, 1881
 = *Goniodiscus acutus* Heller, 1863
 = *Goniodiscus placenta* J. Müller & Troschel, 1842
 = *Goniodiscus placentaformis* Heller, 1863
 = *P. hebes* Verrill, 1899
 = *Pentagonaster minor* Koehler, 1895
 = *Pentagonaster mirabilis* Perrier, 1875
 = *Pentagonaster planus* Verrill, 1895
 = *Pentagonaster vincenti* Perrier, 1885
- Plinthaster**
dentatus (Perrier, 1884)
 = *Pentagonaster concinnus* Sladen, 1891
 = *Pentagonaster dentatus* Perrier, 1884
 = *Pentagonaster grandis* Perrier, 1885
 = *Pentagonaster perrieri* Sladen, 1889
 = *Tosia (Plinthaster) compita* Verrill, 1899
 = *Tosia (Plinthaster) nitida* Verrill, 1899
- Pseudarchaster**
gracilis subsp. *gracilis* (Sladen, 1889)
- = *Aphroditaster gracilis* Sladen, 1889
 = *Astrogonium aequabile* Koehler, 1907
 = *Astrogonium aphrodite* Perrier, 1894
 = *Astrogonium eminens* Koehler, 1907
 = *Astrogonium marginatum* Koehler, 1909
 = *Astrogonium necator* Perrier, 1894
 = *P. concinnus* Verrill, 1894
 = *P. granuliferus* Verrill, 1899
 = *P. ordinatus* Verrill, 1899
 = *Parellii* (Düben & Koren, 1846)
 = *Astrogonium annectens* Perrier, 1894
 = *Astrogonium fallax* Perrier, 1885
 = *Astrogonium hystrix* Perrier, 1894
 = *Astrogonium longobrachiale* Koehler, 1907
 = *Astropecten parellii* Düben & Koren, 1846
 = *P. intermedius* var. *insignis* Verrill, 1895
 = *P. intermedius* Sladen, 1889
 = *P. tessellatus* var. *arcticus* Sluiter, 1895
- Sphaeriodiscus**
bourgeti (Perrier, 1885)
 = *Stephanaster bourgeti* Perrier, 1885
- Family Ophidiasteridae**
- Hazelia**
attenuata Gray, 1840
 = *Asterias coriacea* Grube, 1840
 = *Asterias laevigata* subsp. *varietas* de Lamarck, 1816
 = *Asterias variolata* Risso, 1826, non Retzius, 1805
 = *Ophidiaster (Hazelia) attenuatus* Gray, 1840
 = *Ophidiaster lessonae* Gasco, 1876
- Narcissia**
canariensis (d'Orbigny, 1839)
 = *Asterias canariensis* d'Orbigny, 1839
 = *Fromia narcissae* Perrier, 1885
 = *Fromia narcissiae* Perrier, 1894
 = *N. teneriffae* Gray, 1840
 = *trigonaria* Sladen, 1889
- Ophidiaster**
ophidianus (de Lamarck, 1816)
 = *Asterias ophidiana* de Lamarck, 1816
 = *O. aurantius* Gray, 1840
 = *O. canariensis* Greeff, 1872
 = *reissyi* Sibuet, 1977
- Family Poraniidae**
- Chondraster**
grandis (Verrill, 1878)
 = *?Marginaster fimbriatus* Sladen, 1889
 = *Porania grandis* Verrill, 1878
- Marginaster**
caprensis (Gasco, 1876)
 = *Asteropsis caprensis* Gasco, 1876
- Porania**
(Porania) pulvillus subsp. *pulvillus* (O.F. Müller, 1776)
 = *Asteropsis ctenacantha* J. Müller & Troschel, 1842
 = *Goniaster templetoni* Thomson, in Forbes, 1839
 = *P. gibbosa* Gray, 1840 (*nomen nudum*)
 = *(Pseudoporania) stormi* (Dons, 1935)
 = *Pseudoporania stormi* Dons
- Poraniomorpha**
(Poraniomorpha) bidens Mortensen, 1932
(Poraniomorpha) hispida subsp. *hispida* (M. Sars, 1872)
 = *?Rhegaster murrayi* Sladen, 1883
 = *Asterina borealis* Verrill, 1878
 = *Goniaster hispidus* M. Sars, 1872
 = *Porania spinulosa* Verrill, 1880
(Poraniomorpha) hispida subsp. *rosea* Danielssen & Koren, 1881
 = *Lasiaster villosus* Sladen, 1889
 = *P. rosea* Danielssen & Koren, 1881
(Poraniomorpha) tumida (Stuxberg, 1878)
 = *Rhegaster tumidus* (Stuxberg) auctt.
 = *Solaster tumida* Stuxberg, 1878
 = *Sphaeriaster bjoerklykki* (Dons, 1938)
(Culcitopsis) borealis (Süssbach & Breckner, 1911)

- = *?Marginaster pentagonus* Perrier, in Milne-Edwards, 1882
= *Culcita borealis* Süßbach & Breckner, 1911
= *Sphaeraster berthae* Dons, 1938
= *Sphaeraster berthae* Dons, 1939
Tylaster
willei Danielssen & Koren, 1881
- Family Sphaerasteridae**
Podosphaeraster
thalassae Cherbonnier, 1970
- Family Asterinidae**
Allopatiria
ocellifera (Gray, 1847)
= *Palmipes lobiancoi* Ludwig, 1897
= *Parasterina africana* Engel & Croes, 1960
= *Patiria ocellifera* Gray, 1847
= *Patiria pulla* Koehler & Vaney, 1906
= *Patiria rosea* Koehler & Vaney, 1906
Anseropoda
lobiancoi (Ludwig, 1897)
= *Palmipes lobiancoi* Ludwig, 1897
placenta (Pennant, 1777)
= *Asterias cartilaginea* Fleming, 1828
= *Asterias membranacea* Retzius, 1783
= *Asterias placenta* Pennant, 1777
= *Asteriscus palmipes* J. Müller & Troschel, 1842
Asterina
gibbosa (Pennant, 1777)
= *Asterias gibbosa* Pennant, 1777
= *Asterias verruculata* Retzius, 1805
= *A. minuta* Nardo, 1834
= *Asteriscus arrecifiensis* Greeff, 1872
pancerli (Gasco, 1870)
= *Asteriscus pancerii* Gasco, 1870
phylactica Emson & Crump, 1979
wega Perrier
Tremaster
mirabilis Verrill, 1880
= *T. laevis* H.L. Clark, 1941
- Order Spinulosida**
- Family Echinasteridae**
Echinaster
(*Echinaster*) *sepositus* subsp. *sepositus* (Retzius, 1783)
= *Asterias sagena* Retzius, 1805
= *Asterias seposita* Retzius, 1783
= *Rhopia mediterranea* Gray, 1840
Henricia
caudani (Koehler, 1895)
= *Cribella caudani* Koehler, 1895
cylindrella (Sladen, 1883)
= *Cribella abyssalis* Perrier, 1894
= *Cribella oculata* var. *cylindrella* Sladen, 1883
= *H. abyssicola* Perrier, 1885 & Farran, 1913 & Mortensen, 1927
eschrichti (J. Müller & Troschel, 1842)
= *Echinaster eschrichti* J. Müller & Troschel, 1842
= *Magdalenaaster arcticus* Koehler, 1907
hedingi Madsen, 1987
lisa A.H. Clark, 1949
lisa subsp. *ingolfi* Madsen, 1987
oculata (Pennant, 1777)
= *Asterias oculata* Pennant, 1777
= *H. sanguinolenta* auctt., non (O.F. Müller) (p.p.)
perforata (O.F. Müller, 1776)
= *Asterias perforata* O.F. Müller, 1776
= *Cribella oculata* var. *murina* Perrier, 1875
= *Cribella sanguinolenta* f. *scabrior* Michailovskij, 1902
= *Echinaster scrobiculatus* Danielssen & Koren, 1883
= *H. sanguinolenta* auctt., non (O.F. Müller)
= *H. crossispika* Heding, in Jägerskiöld, 1971 (*nomen nudum*)
= *H. fisheri* subsp. *echinata* Heding, in Jägerskiöld, 1971 (*nomen nudum*)
- = *H. knipowitschi* D'yakonov, 1950
= *H. sanguinolenta* Rasmussen, 1965 non (O.F. Müller)
pertusa (O.F. Müller, 1776)
= *?Cribella biscayensis* Koehler, 1895
= *Asterias pertusa* O.F. Müller, 1776
= *Cribella sanguinolenta* var. *abyssicola* Norman, 1869
= *Echinaster oculatus* f. *major* Düben & Koren, 1846
sanguinolenta (O.F. Müller, 1776)
= *Asterias sanguinolenta* O.F. Müller, 1776
= *Cribella pectinata* Verrill, 1894
= *Cribella sanguinolenta* var. *curta* Norman, 1869
= *Cribella sanguinolenta* f. *laevior* Michailovskij, 1902
= *Echinaster oculatus* f. *minor* Düben & Koren, 1846
= *Echinaster sarsi* J. Müller & Troschel, 1844
= *H. perforata* Rasmussen, 1965 non (O.F. Müller)
= *H. sanguinolenta* subsp. *miliaris* Verrill, 1914
skorikovi D'yakonov, 1950
solida D'yakonov, 1950
spongiosa (O. Fabricius, 1780)
- Order Velatida**
- Family Solasteridae**
Crossaster
papposus (Linnaeus, 1767)
= *Asterias affinis* Brandt, 1835
= *Asterias helianthoides* Pennant, 1777
= *Asterias papposa* Linnaeus, 1767
squamatus (Döderlein, 1900)
= *Solaster affinis* Danielssen & Koren, 1877 non (Brandt, 1835)
= *Solaster papposus* var. *squamata* Döderlein
= *Solaster squamatus* Döderlein
Lophaster
furcifer (Düben & Koren, 1846)
= *Solaster furcifer* Düben & Koren, 1846
Solaster
earlli Verrill, 1879
endeca (Linnaeus, 1771)
= *Asterias alboverrucosa* Brandt, 1835
= *Asterias aspera* O.F. Müller, 1776
= *Asterias endeca* Linnaeus, 1771
= *Asterias rumphii* Parelius, 1768
= *S. endeca* subsp. *decemradiata* Sladen, 1889
= *S. galaxides* Verrill, 1909
= *S. intermedius* Sluiter, 1895
glacialis Danielssen & Koren, 1881
= *S. echinatus* Storm, 1888
syrizensis Verrill, 1894
- Family Korethrasteridae**
Korethraster
hispidus Wyville Thomson, 1873
= *K. setosus* Perrier, in Milne-Edwards, 1882
- Family Pterasteridae**
Diploptaster
multipes (M. Sars, 1866)
= *Pteraster multipes* M. Sars, 1866
Hymenaster
giboryi Perrier, 1894
pellucidus Thomson, 1873
= *H. giganteus* Sladen, 1891
= *H. membranaceus* Thomson, 1877
= *H. perissonotus* Fisher, 1910
regalis Verrill, 1895
= *H. kieri* Downey, 1979
reticulatus Sibuet, 1976
rex Perrier, 1885
= *H. gennaeus* H.L. Clark, 1923
roseus Koehler, 1907
temuropinus Sibuet, 1976
Pteraster
acicula (Downey, 1970)
= *Marsipaster acicula* Downey, 1970
alveolatus Perrier, 1894
hystrix Harvey, 1989
militaris (O.F. Müller, 1776)

- = *Asterias militaris* O.F. Müller, 1776
 = *P. aporus* Ludwig, 1886
obscurus (Perrier, 1891)
 = *Hexaster obscurus* Perrier, 1891
 = *P. (Temnaster) hexactis* Verrill, 1894
personatus Sladen, 1891
 = ?*Pteraster sordidus* Perrier, 1894
 = *P. reductus* Koehler, 1907
pulvillus (M. Sars, 1861)
- Family Myxasteridae**
- Myxaster*
perrieri Koehler, 1896
sol Perrier, 1885
- Pythonaster*
atlanticus A.H. Clark, 1948
- Order Brisingida**
- Family Brisingidae**
- Brisinga*
endecacnemos Asbjørnsen, 1856
hirsuta Perrier, 1894
- Brisingella*
coronata (G.O. Sars, 1871)
 = *Brisinga coronata* G.O. Sars, 1871
 = *Brisingia mediterranea* Perrier, 1885
- Novodinia*
homonyma Downey, 1986
 = *Brisinga elegans* Perrier, 1885, non Verrill, 1884
 = *Odinia elegans* Perrier, 1894
pandina (Sladen, 1889) Downey, 1986
 = *Odinia pandina* Sladen, 1889
semicoronata (Perrier, 1885) Downey, 1986
 = *Brisinga robusta* Perrier, 1885
 = *Brisinga semicoronata* Perrier, 1885
- Family Freyellidae**
- Colpaster*
edwardsi (Perrier, 1882) Downey, 1986
 = *Brisinga edwardsi* Perrier, 1882
scutigerula Sladen, 1889
- Freyastera*
benthophila (Sladen, 1889) Downey, 1986
 = *Freyella benthophila* Sladen, 1889
sexradiata (Perrier, 1885) Downey, 1986
 = *Freyella sexradiata* Perrier, 1885
tuberculata (Sladen, 1889) Downey, 1986
 = *Freyella tuberculata* Sladen, 1889
- Freyella*
elegans (Verrill, 1884)
 = *Brisinga elegans* Verrill, 1884
 = *F. aspera* Verrill, 1895
 = *F. bracteata* Sladen, 1889
 = *F. laubieri* Cherbonnier & Sibuet, 1973
 = *F. spinosa* var. *abyssicola* Perrier, 1894
 = *F. spinosa* Perrier, 1885
recta Koehler, 1907
- Order Forcipulatida**
- Family Asteriidae**
- Asterias*
rubens Linnaeus, 1758
 = *A. clathrata* Pennant, 1777
 = *A. holsatica* Retzius, 1805
 = *A. minuta* Linnaeus, 1761
 = *A. rubens* subsp. *violacea* O.F. Müller, 1776
 = *A. spinosa* Say, 1825
 = *A. vulgaris* Packard, 1863
- Coscinasterias*
tenuispina (de Lamarck, 1816)
 = *Asterias atlantica* Verrill, 1868
 = *Asterias savarensis* Delle Chiaje, 1827
 = *Asterias tenuispina* de Lamarck, 1816
 = *Lytaster inaequalis* Perrier, 1894
- Icastarias*
panopla (Stuxberg, 1878)
 = *Asterias panopla* Stuxberg
- Leptasterias*
canuti Heding, 1936
clavispina Heding, 1936
(Leptasterias) danica (Levinsen, 1887)
degerboelli Heding, 1935
floccosa (Levinsen, 18???)
floccosa var. *crassa* Heding, 1935
(Endogenasterias) groenlandica (Steenstrup, 1857)
 = *Asteracanthion groenlandica* Steenstrup, 1857
 = *Asterias cribaria* Stimpson, 1862
(Leptasterias) hyperborea (Danielssen & Koren, 1883)
 = *Asterias hyperborea* Danielssen & Koren, 1883
 = *Asterias normani* Danielssen & Koren, 1884
hyperborea var. *intermedia* Heding, 1935
(Leptasterias) islandica (Levinsen, 1887)
 = *Asterias mülleri* var. *islandica* Levinsen, 1887
(Leptasterias) muelleri subsp. *muelleri* (M. Sars, 1846)
 = *Asteracanthion mülleri* M. Sars, 1846
(Leptasterias) muelleri subsp. *celtica* Clark & Downey, 1992
 = *Asterias hispida* Pennant, 1777
(Hexasterias) polaris (J. Müller & Troschel, 1842)
 = *Asteracanthion polaris* J. Müller & Troschel, 1842
 = *Asterias acervata* Stimpson, 1862
 = *Asterias borealis* Perrier, 1875
- Marthasterias*
glacialis (Linnaeus, 1758)
 = *Asteracanthion africanus* J. Müller, & Troschel, 1842
 = *Asterias angulosa* O.F. Müller, 1788
 = *Asterias echinophora* Delle Chiaje, 1827
 = *Asterias glacialis* Linnaeus, 1758
 = *Asterias madeirensis* Stimpson, 1862
 = *Asterias rarispina* Perrier, 1875
 = *Asterias spinosa* Pennant, 1777
 = *M. foliacea* Jullien, 1878
 = *Sellonia webbianae* d'Orbigny, 1839
- Sclerasterias*
guernei Perrier, 1891
neglecta (Perrier, 1891)
 = *Asterias edmundi* Ludwig, 1897
 = *Stolasterias neglecta* Perrier, 1891
richardi (Perrier, 1882)
 = *Asterias richardi* Perrier, 1882
- Stephanasterias*
albula (Stimpson, 1853)
 = *Asteracanthion albulus* Stimpson, 1853
 = *Asteracanthion problema* Steenstrup, 1855
 = *Asterias gracilis* Perrier, 1881
- Stichasterella*
ambigua (Farran, 1913)
 = *Stichaster roseus* var. *ambiguus* Farran, 1913
rosea (O.F. Müller, 1776)
 = *Asterias rosea* O.F. Müller, 1776
 = *Chaetaster hermanni* J. Müller & Troschel, 1842
 = *Stichaster arcticus* Danielssen & Koren, 1883
- Urasterias*
linckii (J. Müller & Troschel, 1842)
 = *Asteracanthion linckii* J. Müller & Troschel, 1842
 = *Asteracanthion stellionura* E. Perrier, 1869
 = *Asterias enopla* Verrill, 1895
 = *Asterias gunneri* Danielssen & Koren, 1884
- Family Pedicellasteridae**
- Hydrasterias*
sexradiata (Perrier, in Milne-Edwards, 1882)
 = *Pedicellaster sexradiatus* Perrier, in Milne-Edwards, 1882
- Pedicellaster*
typicus M. Sars, 1861
 = *Archaster bifrons* W. Thomson, 1873
 = *Asteracanthion palaeocrystallus* Duncan & Sladen, 1877
- Family Zoroasteridae**
- Zoroaster*

- fulgens* Thomson, 1873
- Family Labidiasteridae**
- Coronaster*
- volsellatus* (Sladen, 1889)
 - = *Asterias (Stolasterias) volsellatus* Sladen, 1889
 - = *C. antonii* Perrier, 1894
- Family Neomorphasteridae**
- Neomorphaster*
- margaritaceus* (Perrier, *in* Milne-Edwards, 1882)
 - A.M. Clark & Downey, 1992
 - = *Calycaster monoecus* Perrier, 1891
 - = *N. eustichus* Sladen, 1889
 - = *N. parfaitei* Koehler, 1896
 - = *Pedicellaster margaritaceus* Perrier, *in* Milne-Edwards, 1882
 - = *Stichaster talismani* Perrier, 1885
- Class Ophiuroidea**
- Order Phrynomphiurida**
- Family Ophiomyxidae**
- Astrogeron*
- supinus* (Lyman, 1883)
 - = *Ophiogeron supinus* Lyman, 1883
- Ophiogeron*
- edentulus* Lyman, 1878
- Ophilomyxa*
- pentagona* (de Lamarck, 1816)
 - = *Ophiura pentagona* de Lamarck, 1816
 - serpentaria* Lyman, 1883
- Ophiophrixus*
- spinosis* (Storm, 1881)
 - = *Ophiobrysa hystricis* Lyman, 1883
 - = *Ophiobryssa hystricis* (Lyman, 1883)
- Ophioscolex*
- glacialis* J. Müller & Troschel, 1842
 - purpureus* Döbken & Koren, 1846
 - = ?*Ophioscolex quadrispinus* Verrill, 1885
- Order Euryalina**
- Family Asteronychidae**
- Asteronyx*
- loveni* J. Müller & Troschel, 1842
 - = *A. cooperi* Bell, 1909
 - = *A. dispar* Lütken & Mortensen, 1899
 - = *A. locardi* Koehler, 1896
 - = *Ophiuropsis lymani* Studer, 1884
- Astroadia*
- tenuispina* (Verrill, 1884)
 - = *A. bispinosa* Koehler, 1922
- Family Gorgonocephalidae**
- Astrospartus*
- mediterraneus* (Risso, 1826)
 - = *Euryale arborescens* L. Agassiz, 1839
 - = *Euryale mediterraneus* Risso, 1826
 - = *Gorgonocephalus verrucosus* Grube, 1840
- Gorgonocephalus*
- arcticus* Leach, 1819
 - = *Astrophyton agassizi* Stimpson, 1853
 - caput-medusae* (Linnaeus, 1758)
 - = *Asterias caput-medusae* Linnaeus, 1758
 - = *Astrophyton linckii* J. Müller & Troschel, 1842
 - = *Astrophyton scutatum* Fleming, 1828
 - eucnemis* (J. Müller & Troschel, 1842) Lyman, 1882
 - = *Astrophyton eucnemis* J. Müller & Troschel, 1842
 - = *Astrophyton malmsgreni* Danielssen & Koren, 1877
 - = *G. caryi* Lyman, 1860
 - = *G. japonicus* Döderlein, 1907
 - lamarckii* (J. Müller & Troschel, 1842) Lyman, 1882
 - = *Astrophyton lamarckii* J. Müller & Troschel, 1842
- Family Asteroschematidae**
- Asterschema*
- ornatum* Koehler, 19??
- Ophiocreas*
- oedipus* Lyman, 1879
- Order Ophiurida**
- Family Ophiotrichidae**
- Ophiothrix*
- fragilis* (Abildgaard, *in* O.F. Müller, 1789)
 - = *Asteria cuvieri* Delle Chiaje, 1828
 - = *Asterias echinata* Delle Chiaje, 1822-29
 - = *Asterias fragilis* Abildgaard, *in* O.F. Müller, 1789
 - = *Asterias pentaphylla* Pennant, 1777
 - = *Asterias rubra* Delle Chiaje, 18??
 - = *Ophiocoma minuta* Forbes, 1839
 - = *Ophiocoma rosula* Forbes, 1839
 - = *O. alopecurus* J. Müller & Troschel, 1842
 - = *O. ferussacii* J. Müller & Troschel, 1842
 - = *O. lusitanica* Ljungman, 1872
 - = *O. rammelsbergii* J. Müller & Troschel, 1842
 - = *O. rubra* Ljungman, 1872
 - = *Ophiura quinquemaculata* Delle Chiaje, 1828
 - indigna* Koehler, 1906
 - luetkeni* Wyville Thomson, 1873
 - maculata* Ljungman, 1872
 - = *O. inducta* Koehler, 1906
- Family Ophiactidae**
- Histampica*
- duplicata* (Lyman, 1875)
- Ophiactis*
- abyssicola* (M. Sars, 1861) Ljungman, 1866
 - = ?*Ophiactis cuspidata* Lyman, 1882
 - = *Amphiura abyssicola* M. Sars, 1861
 - = *O. corallicola* Koehler, 1896
 - = *O. echinata* Koehler, 1898
 - = *O. poa* Lyman, 1882
 - balli* (W. Thompson, 1840)
 - = *Ophiocoma balli* W Thompson, 1840
 - = *Ophiocoma goodsir* Forbes, 1840 (Nov.)
 - canotia* Lyman, 1882
 - lymani* Ljungman, 1872
 - nidarosiensis* Mortensen, 1920
 - profundi* Lütken & Mortensen, 1899
 - savignyi* (J. Müller & Troschel, 1842)
 - = *O. brocki* de Loriol, 1893
 - = *O. krebsii* Lütken, 1857
 - = *Ophiolepis savignyi* J. Müller & Troschel, 1842
 - virens* (M. Sars, 1857)
 - = *Amphiura virens* M. Sars, 1857
- Ophiopholis*
- aculeata* (Linnaeus, 1767)
 - = *Ophiocoma bellis* Forbes, 1839
 - = *Ophiolepis scolopendrica* J. Müller & Troschel, 1842
 - = *Polypholis echinata* Duncan, 18??
- Ophiopus*
- arcticus* Ljungman, 1867
 - = *Ophiaregma abyssorum* G.O. Sars, 1872
- Family Amphiuridae**
- Amphichilus*
- daleus* (Lyman, 1882) Matsumoto, 1917
 - = *Amphiura dalea* Lyman, 1882
- Amphioplus*
- verrilli* (Lyman, 1879)
 - = *Silix pulvinus* Cherbonnier & Sibuet, 1972
- Amphipholis*
- instructa* Koehler, 1906
 - muranica* D'yakonov, 1929
 - squamata* (Delle Chiaje, 1829)
 - = *A. lineata* Ljungman, 1872
 - = *A. tenera* Lütken, 18??
 - = *Amphiura elegans* (Leach, 1815)
 - = *Asterias squamata* Delle Chiaje, 1829
 - = *Axiognathus squamata* (Delle Chiaje)
 - torelli* Ljungman, 1872
- Amphiura*
- abyssorum* Norman, *in* Jeffreys, 1876
 - = *A. digna* Koehler, 1906
 - bellis* Lyman, 1879
 - bellis* var. *tritonis* Hoyle, 1884

- borealis* (G.O. Sars, 1871) Ljungman, 1872
 = *Ophiopeltis borealis* G.O. Sars, 1871
(Acrocnida) brachiata (Montagu, 1804)
 = *Amphiocnida brachiata* (Montagu)
 = *A. neapolitana* M. Sars, 1857
 = *Asterias brachiata* Montagu, 1804
cherbonnieri Guille, 1972
chiajei Forbes, 1843
delamarei Cherbonnier, 1958?
denticulata Koehler, 189
filiformis (O.F. Müller, 1776)
 = *Asterias filiformis* O.F. Müller, 1776
fragilis Verrill, 1885
 = *A. denticulata* Koehler, 1898
grandisquama Lyman, 1869
 = *A. iris* Lyman, 1879
 = *A. josephinae* Ljungman, 1872
 = *A. longispina* Koehler
griegi Mortensen, 1920
incana Lyman, 1879
medierranea Lyman, 1882
otteri Ljungman, 1872
 = *A. grandis* Koehler, 1896
palmeri Lyman, 1882
richardi Koehler, 1906
sarsi Ljungman, 1872
(Ophiopeltis) securigera (Düben & Koren, 1846)
 = *Ophiopeltis securigera* Düben & Koren, 1846
(Acrocnida) semi squamata (Koebler)
senegalensis Madsen, 1970
stepanovi D'yakonov, 1954
 = *A. stepanovi* Chernyavskii, 1861 (*nomen nudum*)
 = *A. stepanovi* Lyman, 1869 (*nomen nudum*)
sundevalli (J. Müller & Troschel, 1842)
 = *Ophiolepis sundevalli* J. Müller & Troschel, 1842
- Monophiura**
apicula (Cherbonnier, 1957)
 = *Amphiura apicula* Cherbonnier, 1957
- Paramphiura**
punctata (Forbes, 1841)
 = *Ophiocoma punctata* Forbes, 1841
- Family Amphilepididae**
Amphilepis
norvegica (Ljungman, 1865)
 = *Amphiura norvegica* Ljungman, 1865
ingolfiana Mortensen, 1933
- Order Laemophiurina**
- Family Ophiacanthidae**
Ophiacantha
abyssicola G.O. Sars, 1871
 = *O. millespina* Verrill, 1879
angolensis Koehler, 1923
anomala G.O. Sars, 1871
 = ?*Ophiacantha nodosa* Lyman, 18??
aristata Koehler
bidentata (Retzius, 1805)
 = *Asterias bidentata* Retzius, 1805
 = *O. fraterna* Verrill, 1885
 = *O. groenlandica* J. Müller & Troschel, 1844
 = *O. hibernica* Farran, 1913
 = *O. spinulosa* J. Müller & Troschel, 1842
brevispina Koehler, 1898?
composita Koehler, 1907
crassidens Verrill
cuspidata Lyman, 1882
(Ophiotreta) danae Mortensen, 1933
 = *O. valenciennesi* Lieberkind, 1929, *non* Lyman
decipiens Koehler, 1906
densa Farran, 1913?
enopla subsp. *veterna* (Koehler, 1907)
hirsuta Lyman, 1875
lineata Koehler, 1909
mesembria H.L. Clark, 1915
nodosa Lyman, 1882
- notata* Koehler, 1906
opercularis (Koehler, 1909)
 = *Ophiolimna opercularis* Koehler, 1909
parcita Koehler, 1906
setosa (Retzius, 1805)
 = *Ophiura setosa* Retzius, 1805
smitti Ljungman, 1872
simulans Koehler, 1896
 = *O. composita* Koehler, 1906, *non* Koehler, 1904
 = *O. composita* Mortensen, 1927
spectabilis G.O. Sars, 1871
spinosella Mortensen, 1933
valenciennesi Lyman, 1879
- Ophiambix**
devaneyi Paterson, 1985
meteoris Bartsch, 1983
- Ophiochondrus**
armatus (Koehler, 1909)
 = *Ophioplus armatus* Koehler, 1909
- Ophiocomina**
nigra (Abildgaard, *in* O.F. Müller, 1789)
 = *Asterias nigra* Abildgaard, *in* O.F. Müller, 1789
 = *Ophiacantha danae* Mortensen, 1933
 = *Ophiocoma granulata* Forbes, 1839
 = *Ophiocoma nigra* (Abildgaard)
 = *Ophiocoma nilssonii* J. Müller & Troschel, 1842
 = *Ophiocoma raschi* G.O. Sars, 18??
- Ophiolebes**
bacata Koehler, 1909
reecta (Koehler, 1896)
 = *Ophioscolex reectus* Koehler, 1896
- Ophiomitra**
hamula Mortensen, 1933
- Ophiopsila**
annulosa (M. Sars, 1859)
 = *Ophianoplus annulosus* M. Sars, 1859
aranea Forbes, 1845
 = *Ophianoplus marmoreus* M. Sars, 18??
aff. guineensis (Koehler, 1914)
- Ophiocamax**
dominans Koehler, 19??
- Ophiomitrella**
clavigera (Ljungman, 1865)
 = ?*Ophiomitrella cordifera* Koehler, 1909
 = *Ophiactis clavigera* Ljungman, 1865
 = *Ophiolebes acanellae* Verrill, 1885
 = *Ophiolebes claviger* (Ljungman, 1865)
globulifera (Koehler, 1896)
- Ophioplinthaca**
carduus (Lyman, 1882)
chelys (Lyman, 1882)
- Ophiothamnus**
affinis Ljungman, 1872
 = *Ophioleda minima* Koehler, 1906
 = *Ophioplinthaca occlusa* Koehler, 1907
 = *O. minima* Matsumoto, 1917
chariis H.L. Clark, 1915
 = *O. gracilis* (Verrill, 1885), *non* (Studer, 1822)
- Ophiolimna**
bairdi (Lyman, 1883) Verrill, 1899
 = *Ophiacantha bairdi* Lyman, 1883
- Ophiomedea**
duplicata Koehler, 1906
- Ophioprium**
rosea (Lyman, 1879) Baker & Devaney, 1981
 = *Ophiacantha rosea* Lyman, 1879
 = *Ophiacantha truncata* Koehler, 1930
- Ophiotoma**
bartletti (Lyman, 18??)
coriacea Lyman, 1883
- Ophiotrema**
alberti Koehler, 1896
- Ophiomyces**
fructosus Lyman, 1869
grandis Lyman, 1878

Order Chilophiurina**Family Ophiuridae***Amphiophiura*

- bullata* (Wyville Thomson, 1873) Matsumoto, 1915
- = *Ophioglypha abdita* Koehler, 1909
- = *Ophioglypha bullata* Wyville Thomson, 1873
- bullata* subsp. *convexa* (Lyman, 1878)
- = *Ophioglypha convexa* Lyman, 1878
- bullata* subsp. *vittata* Litvinova, 1971
- = *A. vittata* Litvinova, 1971
- saurura* (Verrill, 1894)
- = *Ophioglypha aspera* Koehler, 1898
- = *Ophioglypha saurura* Verrill, 1894

Anthophiura

- ingolfi* Fasmer, 1930

Aspidophiura

- minuta* (Lyman, 1878)
- = *Ophioglypha minuta* Lyman, 1878

Homophiura

- abyssorum* (Lyman, 1883)
- abyssorum* var. *africana* (Madsen, 1947)
- = *Homalophiura abyssorum* var. *africana* Madsen, 1947
- tessellata* (Verrill, 1894)
- = *Homalophiura tessellata* (Verrill, 1894)
- = *Ophioglypha tessellata* Verrill, 1894

Ophiocten

- abyssicolum* von Marenzeller, 18??
- affinis* (Lütken, 1858) Sumida, Tyler, Gage & Nørrevang, 1998
- = *Ophiusa affinis* Lütken, 1858
- centobi* Paterson, Tyler & Gage, 1982
- gracilis* (G.O. Sars, 1871)
- = *O. sericeum* auct., non (Forbes, 1852) (p.p.)
- = *Ophioglypha gracilis* G.O. Sars, 1871
- = *Ophioglypha signata* Verrill, 1882
- = *Ophiusa signata* (Verrill, 1882)
- hastatum* Lyman, 1878
- = *Ophiocten longispinum* Koehler, 1898
- = *O. latens* Koehler, 1906
- = *O. pacificum* Lütken & Mortensen, 1899
- scutatum* Koehler, 1898
- sericeum* (Forbes, 1852)
- = *O. kroeyeri* Lütken, 1855
- = *Ophiusa sericea* Forbes, 1852

Ophioctenella

- acies* Tyler, Paterson, Sibuet, Guille, Murton & Segonzac, 1995

Ophiolepis

- paucispina*?

Ophiomisidium

- pulchellum* (Wyville Thomson, 18??)
- = *Ophiomusium pulchellum* Wyville Thomson, 18??

Ophiomusium

- africanum* Koehler, 1909
- lymani* Wyville Thomson, 1873

Ophiodyskis

- mirabilis* Koehler, 1901

Ophiolepta

- aurantiaca* (Verrill, 1882) Verrill, 1882
- = *Ophioglypha aurantiaca* Verrill, 1882
- borealis* Danielssen & Koren, 1877
- = *Luetkenia arctica* Duncan, 1878

Ophiosphalma

- armigerum* (Lyman, 1878) H.L. Clark, 1941
- = *Ophiomusium armigerum* Lyman, 1878
- planum* (Lyman, 1878) H.L. Clark, 1941
- = *Ophiomusium planum* Lyman, 1878

Ophiotjalfa

- vivipara* Mortensen, 1913

Ophiotypa

- simplex* Koehler, 1897

Ophiozonella

- molesta* (Koehler, 1906)

- = *Ophiozona molesta* Koehler, 1906

- sincera* (Koehler, 1906)

= *Ophiozona sincera* Koehler, 1906

tfalfiana (Mortensen, 1913) Matsumoto, 1915

= *Ophiozona tfalfiana* Mortensen, 1913

Ophiura

- albida* Forbes, 1839

africana (Koehler)

= *Ophiocten africana* Koehler

(*Dictenophiura*) *carnea* Lütken, 1858 ex M. Sars MS

clemens (Koehler, 1906)

= *Ophioglypha clemens* Koehler, 1906

flagellata (Lyman, 1878)

= *Gymnophiura caerulescens* Lütken & Mortensen, 1899

= *Ophioglypha flagellata* Lyman, 1878

grubei Heller, 1863

imprudens (Koehler, 1906)

= *Ophioglypha imprudens* Koehler, 1906

inermis (Lyman, 1882)

= *Ophioglypha inermis* Lyman, 1882

(*Ophiuroglypha*) *irrorata* (Lyman, 1878)

= *Ophioglypha grandis* Verrill, 1894

= *Ophioglypha involuta* Koehler, 1899

= *Ophioglypha irrorata* Lyman, 1878

= *Ophioglypha orbiculata* Lyman, 1882

= *Ophioglypha tumulosa* Lütken & Mortensen, 1899

(*Ophiuroglypha*) *irrorata* subsp. *concreta* (Koehler, 1901)

= *Ophioglypha concreta* Koehler, 1901

(*Ophiuroglypha*) *irrorata* subsp. *polyacantha*

Mortensen, 1933

= *Homalophiura multispina* Cherbonnier & Sibuet, 1972

ljungmani (Lyman, 1878)

= *Ophioglypha lepida* Lyman, 1882

= *Ophioglypha ljungmani* Lyman, 1878

= *Ophioglypha thouleti* Koehler, 1896

madseni (Belyaev, 1972)

= *Homalophiura madseni* (Belyaev, 1972)

(*Ophiuroglypha*) *mundata* Koehler, 1906

niida Mortensen, 1933

ophiura (Linnaeus, 1758)

= *Asterias ophiura* Linnaeus, 1758

= *O. ciliaris* (Linnaeus, 1766)

= *O. ciliata* (Retzius, 1783) non (O.F. Müller, 1776)

= *O. lacertosa* (Pennant, 1777)

= *O. texturata* de Lamarck, 1816

robusta (Ayres, 1851)

= *Ophiolepis robusta* Ayres, 1851

= *O. fasciculata* Forbes, 1854

= *O. squamosa* Lütken, 1859

sarsi Lütken, 1858

violaine (Cherbonnier & Sibuet, 1972)

= *Homalophiura violaine* Cherbonnier & Sibuet, 1972

Ophiurolepis

inornata (Lyman, 1878) Paterson, 1985

= *Homalophiura inornata* (Lyman, 1878)

= *Ophioglypha indivisa* Lütken & Mortensen, 1899

= *Ophioglypha inornata* Lyman, 1878

Stegophiura

nodosa (Lütken, 1858) Matsumoto, 1917

= *Ophiura nodosa* Lütken, 1858

stuvitzii (Lütken, 1858) Matsumoto, 1917

= *Ophiura stuvitzii* Lütken, 1858

Uriopha

ios Paterson, 1980

Family Ophioleucidae*Ophiermus*

adspersus Lyman, 1883

vallincola Lyman, 1878

= *O. abyssalis* Koehler, 1896

Ophiocrates

intervallus Madsen, 1947

secundus Koehler, 1906

Ophiopyren

striatum Mortensen, 1933

Family Ophionereididae*Ophiochiton*

- solutus* Koehler, 1906
ternispinus Lyman, 1883
 = *O. grandis* Verrill, 1884
 = *O. tenuispinus* Hoyle, 1884
- Ophionereis*
reticulata (Say, 1825)
 = *Ophiura reticulata* Say, 1825
- Family Ophiodermatidae**
- Bathypectinura*
heros (Lyman, 1879)
 = *B. elata* (Koehler, 1906)
 = *Pectinura elata* Koehler, 1906
 = *Pectinura heros* Lyman, 1879
- Cryptopelta*
brevispina (Ludwig, 1879)
 = *Ophioconis brevispina* Ludwig, 1879
- Ophiarachnella*
semicincta (Studer, 18???)
 = *Pectinura semicincta* Studer, 18???
- Ophioconis*
forbesi (Heller, 1863)
 = *Pectinura forbesi* Heller, 1863
vivipara Mortensen, 19???
- Ophiiderma*
longicauda (Retzius, 1805)
 = *Asterias longicauda* Retzius, 1805
 = *O. lacertosum* Lamarck, 1801
 = *Ophiura laevis* Lyman, 1865
- Pectinura*
vestita Forbes, 1843
- Class Echinoidea**
- Order Cidaroida**
- Family Cidaridae**
- Cidaris*
cidaris (Linnaeus, 1758)
 = *C. borealis* Düben & Koren, 1846
 = *C. papillata* Leske, 1778
 = *Cidarites hystrix* de Lamarck, 1816
 = *Dorocidaris abyssicola* A. Agassiz, 1869?
 = *Echinus cidaris* Linnaeus, 1758
tribuloides de Lamarck, 1816
- Poriocidaris*
purpurata (Wyville Thomson, 1875)
 = *Porocidaris gracilis* Sladen, 1891
 = *Porocidaris purpurata* Wyville Thomson, 1875
- Stereocidaris*
ingolfiana Mortensen, 1903
- Stylocidaris*
affinis (Philippi, 1845)
 = *Cidaris affinis* Philippi, 1845
 = *Cidaris stokesii* L. Agassiz & Desor, 1846-47
- Order Echinothuroidea**
- Family Echinothuridae**
- Calveriosoma*
fenestratum (Wyville Thomson, 1872)
 = *Araeosoma violaceum* Mortensen, 1903
 = *Calveria fenestrata* Wyville Thomson, 1872
- hystrix* (Wyville Thomson, 1872)
 = *Asthenosoma reynoldsi* A. Agassiz, 1880
 = *Calveria hystrix* Wyville Thomson, 1872
- Hygrosoma*
petersii (A. Agassiz, 1880)
 = *Phormosoma luculentum* Koehler, 1896, non A. Agassiz, 18???
 = *Phormosoma petersii* A. Agassiz, 1880
 = *Phormosoma uranus* Koehler, 1898, non Wyville Thomson, 1877
- uranus* (Wyville Thomson, 1877)
 = *Phormosoma uranus* Wyville Thomson, 1877
- Phormosoma*
placenta Wyville Thomson, 1874
 = *P. sigsbei* A. Agassiz, 1883
- Sperosoma*
- grimaldii* Koehler, 1897
Tromikosoma
koehleri Mortensen, 1903
- Order Diadematoida**
- Family Aspidodiadematidae**
- Aspidodiadema*
jacobyi A. Agassiz, 1883
- Plesiodiadema*
antillarum (A. Agassiz, 1883)
 = *Aspidodiadema antillarum* A. Agassiz, 1883
- Family Diadematidae**
- Centrostephanus*
longispinus (Philippi, 1845)
 = *Cidaris (Diadema) longispina* Philippi, 1845
- Diadema*
antillarum Philippi, 1845
 = *Cidaris (Diadema) antillarum* Philippi, 1845
- Order Pedinoida**
- Family Pedinidae**
- Caenopedia*
cubensis A. Agassiz, 1869
- Order Salenoida**
- Family Saleniidae**
- Salenia*
pattersoni A. Agassiz
- Salenocidaris*
profundi subsp. *profundi* (Duncan, 1877)
 = ?*Salenia hastigera* A. Agassiz
 = *Salenia hastigera* subsp. *acuminata* Mortensen, 1935
 = *Salenia profundi* subsp. *occlusa* Mortensen, 1934
 = *Salenia profundi* Duncan, 1877
profundi subsp. *brevispina* Markov, 1988
varispina A. Agassiz, 1869
- Order Arbacoida**
- Family Arbacidae**
- Arbacia*
lixula (Linnaeus, 1758)
 = *Cidaris pustulosa* Leske, 1778
 = *Echinus aequituberculatus* de Blainville, 1825
 = *Echinus lixula* Linnaeus, 1758
 = *Echinus neopolitanus* Delle Chiaje, 1825
- Arbaciella*
elegans Mortensen, 1910
- Coelopleurus*
floridanus A. Agassiz, 1872
- Habrocidaris*
scutata (A. Agassiz, 1880)
- Podocidaris*
sculpia A. Agassiz
- Order Temnopleuroidea**
- Family Temnopleuridae**
- Genocidaris*
maculata A. Agassiz, 1869
 = *Arbacina pallaryi* Gauthier, 1897
 = *Temnechinus maculatus* A. Agassiz, 1869
- Hypsiechinus*
coronatus Mortensen, 1903
- Trigonocidaris*
albida A. Agassiz
- Family Toxopneustidae**
- Sphaerechinus*
granularis (de Lamarck, 1816)
 = *Echinus (Toxopneustes) brevispinosus* Risso, 1826
 = *Echinus granularis* de Lamarck, 1816
 = *S. roseus* Russo, 1893
- Order Echinoida**
- Family Echinidae**
- Echinus*
acutus de Lamarck, 1816
 = ?*Echinus depressus* G.O. Sars, 1871

- = *Echinus microstoma* Wyville Thomson, 1875
 = *Echinus norvegicus* Dübén & Koren, 1846
 = *Echinus rarispinus* G.O. Sars, 1872
 = *E. flemingii* Forbes, 1841
 = *E. sardicus* Delle Chiaje, 1841
affinis Mortensen, 1903
alexandri Danielssen & Koren, 1883
elegans Dübén & Koren, 1846
 = *Echinus wallisi* A. Agassiz, 1883
esculentus Linnaeus, 1758
 = *E. aurantiacus* de Blainville, 1825
 = *E. globiformis* de Lamarck, 1801
 = *E. pseudomelo* de Blainville, 1825
 = *E. quinqueangulatus* de Blainville, 1825
 = *E. quinqueangulosus* de Blainville, 1834
 = *E. schwartzii* Nilsson & Holst, 1817
 = *E. sphaera* O.F. Müller, 1776
 = *E. subangulosus* de Blainville, 1826
 = *E. violaceus* de Blainville, 1825
esculentus var. *glacialis* D'yakonov, 1923
melo Lamarck, 1816
tenuispinus Norman, 1868
 = *E. esculentus* var. *tenuispinus* Norman, 1868
- Paracentrotus**
lividus (de Lamarck, 1816)
 = *Echinus lividus* de Lamarck, 1816
 = *Echinus purpureus* Risso, 1826
 = *Echinus saxatilis* Tiedemann, 1816
 = *Echinus vulgaris* de Blainville, 1825
- Psammechinus**
microtuberculatus (de Blainville, 1825) Heller, 1868
 = *Echinus microtuberculatus* de Blainville, 1825
miliaris (P.L.S. Müller, 1771)
 = *Echinus miliaris* P.L.S. Müller, 1766
 = *Parechinus miliaris* Mortensen, 1903
- Family Strongylocentrotidae**
Strongylocentrotus
droebachiensis (O.F. Müller, 1776)
 = *Echinus droebachiensis* O.F. Müller, 1776
 = *Echinus granularis* Say, 1826
 = *Echinus neglectus* de Lamarck, 1816
 = *S. chlorocentrotus* Brandt, 1835
 = *Toxopneustes pictus* Norman, 18??
pallidus (G.O. Sars, 1871)
 = *?Strongylocentrotus granularis* (Say, 1827)
 = *S. echinoides* A. Agassiz & H.L. Clark, 1907
 = *S. pictus* (Norman, 1868)
 = *S. sachalinicus* Clark, 1912
- Order Clypeasteroida**
- Family Fibulariidae**
Echinocyamus
grandiporus Mortensen, 1907
macrostomus Mortensen, 1907
pusillus (O.F. Müller, 1776)
 = *E. angulosus* Leske, 1778
 = *E. parthenopaeus* Costa, 1869
 = *E. speciosus* Costa, 1869
 = *Echinus minutus* Gmelin, 1790
 = *Fibularia equina* Aradas, 1850
 = *Fibularia tarentina* de Lamarck, 1816
 = *Spatagus pusillus* O.F. Müller, 1776
- Order Cassiduloida**
- Family Neolampadidae**
Neolampas
rostellata A. Agassiz, 1869
- Order Pourtalesioida**
- Family Pourtalesiidae**
Echinosigra
phiale (Wyville Thomson, 1873) Mortensen, 1907
 = *E. paradoxa* (Mortensen, 1905) Mortensen, 1907
 = *Pourtalezia paradoxa* Mortensen, 1905
 = *Pourtalezia phiale* Wyville Thomson, 1873
- Pourtalezia*
- aff. *alcocki* Koehler, 1914
 = *P. miranda* auctt., non A. Agassiz, 1869
jeffreysi Wyville Thomson, 1873
wandeli Mortensen, 1905
- Family Urechinidae**
Urechinus
naresianus A. Agassiz, 1881
- Family Plexechinidae**
Plexechinus
hirsutus Mortensen, 1905
- Family Calymnidae**
Calymne
relicta Wyville Thomson, 1877?
- Order Spatangoida**
- Family Hemiasteridae**
Hemiaster
expergitus Lovén, 1874
 = *H. gibbosus* A. Agassiz, 1881
 = *H. mentzii* A. Agassiz, 1883
 = *H. zonatus* A. Agassiz, 1881
- Family Spatangidae**
Spatangus
inermis Mortensen, 1913
purpureus (O.F. Müller, 1776)
 = *Spatagus purpureus* O.F. Müller, 1776
 = *S. meridionalis* Risso, 1825
 = *S. reginae* Gray, 1855
 = *S. spinosissimus* Desor, 1847
raschi Lovén, 1869
 = *S. meridionalis* Norman, 18??, non Risso, 1825
- Family Schizasteridae**
Brisaster
fragilis (Dübén & Koren, 1846)
 = *Brissus fragilis* Dübén & Koren, 1846
 = *Schizaster fragilis* (Dübén & Koren)
Schizaster
canaliferus (de Lamarck, 1816)
 = *Spatangus canaliferus* de Lamarck, 1816
- Family Brissidae**
Brissopsis
atlantica Mortensen, 1907
atlantica var. *mediterranea* Mortensen, 1913
lyrifera (Forbes, 1841)
 = *B. parma* Val., 18??
 = *Brissus lyrifer* Forbes, 1841
 = *Brissus pulvinatus* Philippi, 1845
 = *Schizaster incertus* Aradas, 1850
- Brissus*
unicolor (Leske, 1778)
 = *B. brissus* auctt.
 = *B. scillae* L. Agassiz, 1836
 = *Spatangus brissus* subsp. *unicolor* Leske, 1778
 = *Spatangus columbaris* de Lamarck, 1816
- Plagiobrissus*
affinis?
costai (Gasco, 1876)
 = *Metalia costae* Gasco, 1876
 = *Rhabdobraissus perrieri* Bernard
- Family Loveniidae**
Echinocardium
cordatum (Pennant, 1777)
 = *Amphidetus kürzii* Girard, 1852
 = *E. sebae* Gray, 18??
 = *Echinus cordatus* Pennant, 1777
 = *Spatangus arcarius* de Lamarck, 1816
fenaui Péquignat, 1963
flavescens (O.F. Müller, 1776)
 = *Amphidetus roseus* Forbes, 1841
 = *Spatagus flavescens* O.F. Müller, 1776
 = *Spatangus ovatus* Leske, 1778
mediterraneum (Forbes, 1844)
 = *Amphidetus mediterraneus* Forbes, 1844

- = *E. gibbosus* L. Agassiz & Desor, 1847
mortenseni Thiéry, 1909
= ?*Echinocardium laevigaster* A. Agassiz, 18??
= ?*Echinocardium pennatifidum* Koehler, 1898, non Norman, 1868
= *E. intermedium* Mortensen, 1907
pennatifidum Norman, 1868
= *E. gibbosus* Barrett, 1857 non L. Agassiz & Desor, 1847
- Family Aeropsidae**
Aceste
bellidifera Wyville Thomson, 1877
Aeropsis
rostrata (Wyville Thomson, 1877) Mortensen, 1907
= *Aerope rostrata* Wyville Thomson, 1877
- Family Asterostomatidae**
Palaeotropus
josephinae Lovén, 18??
= *Palaeobrissus hilgardi* A. Agassiz
= *P. hirondellei* Koehler
Peripatagus
cinctus Koehler
- Class Holothuroidea**
- Order Dendrochirotida**
- Family Psolididae**
Psolidium
complanatum Cherbonnier, 1969
Psolus
fabricii (Düben & Koren, 1846)
= *Cuvieria fabricii* Düben & Koren, 1846
hypsinotus Heding, 1942
nummularis R. Perrier, 1899
operculatus (Pourtalès, 1868)
= *Cuvieria operculata* Pourtalès, 1868
phantapus (Strassenfels, 1765) Jaeger, 1833
= *Holothuria phantapus* Strassenfels, 1765
= *P. granulatus* Ayres, 1851
= *P. laevigatus* Ayres, 1851
= *P. regalis* Verrill, 1866
pourtalesii Théel, 1886
squamatus (O.F. Müller, 1776) Lütken, 1857
= *Holothuria squamata* O.F. Müller, 1776
= *P. valvatus* Östergren, 1904
tesselatus Koehler, 1896
- Family Phyllophoridae**
Ekmania
barthii (Troschel, 1846)
= *Orcula barthii* Troschel, 1846
= *Orcula luminosa* Lampert, 1885
= *Thyonidium pellucidum* Düben & Koren, 1846
- Euthyonidiella**
atlantica (Ludwig & Heding, 1935)
= *Pseudocucumis atlantica* Ludwig & Heding, 1935
- Neocucumis**
atlantica (Ludwig & Heding, 1935)
marioni (von Marenzeller, 1878)
= *Cucumaria marioni* von Marenzeller, 1878
= *Pseudocucumis marioni* Ludwig, 1874
- Neopentadactyla**
mixta (Östergren, 1898) Deichmann, 1944
= *Pseudocucumis cuenoti* Koehler & Vaney, 1905
= *Pseudocucumis mixta* Östergren, 1898
- Phyllophorus**
drachi Cherbonnier & Guille, 1968
granulatus (Grube, 1840)
= *Psolus granulatus* Grube, 1840
(*Phyllophorus*) *pedinaequalis* Cherbonnier, 1969
(*Phyllophorus*) *urna* Grube, 1840
= *Holothuria penicillus* Delle Chiaje, 1828
- Thyonidium**
hyalinum (Forbes, 1841)
= *Holothuria pellucida* Fleming, 1828, non Vahl, 1806
drummondii (Thompson, 1840)
- = *Cucumaria communis* Forbes, 1841
= *Cucumaria nobilis* Ludwig, 1874
= *Cucumaria perspicua* Ludwig, 1874
= *Holothuria drummondii* Thompson, 1840
= *Thyone portlocki* Forbes, 1841
= *T. düberi* Norman, 1869
- Family Cucumiidae**
Abyssocucumis
abyssorum (Théel, 1886)
= *Cucumaria abyssorum* auctt., non Théel, 1886
= *Cucumaria abyssorum* Théel, 1886
= *Cucumaria shuteri* Ohshima, 1915
= *Staurocucumis ingolfi* Clark & Deichmann, 1936
- Aslia*
lefevrei (Barrois, 1882)
= *Cucumaria lefevrei* Barrois, 1882
= *Cucumaria normani* Allen & Pace, 1904
= *Holothuria montagui* Fleming, 1828 (p.p.)
- Cucumaria*
arcuata (Hérouard, 1921)
= *Psolidium arcuatum* Hérouard, 1921
compressa (R. Perrier, 1898)
= *Ocnus compressus* R. Perrier, 1898
frondosa (Gunnerus, 1767)
= *C. assimilis* Düben & Koren, 1846
= *Holothuria frondosa* Gunnerus, 1767
= *Holothuria fucicola* Forbes & Goodsir, 1839
= *Holothuria grandis* Forbes & Goodsir, 1839
= *Holothuria pentactes* Linnaeus, 1767
paraglacialis Heding, 1942
- Havelockia*
inermis (Heller, 1868)
= *Thyone elegans* Norman, 1869
= *Thyone inermis* Heller, 1868
= *Uroxia aurantiaca* Costa, 1869
- Neocnus*
incubans Cherbonnier, 1972
- Ocnus*
glacialis (Ljungman, 1879)
= *Cucumaria glacialis* Ljungman, 1879
grubei (von Marenzeller, 1874)
= *Cucumaria dicquemarii* M. Sars, 1857
= *Cucumaria grubei* von Marenzeller, 1874
= *Pseudocnus grubei* Panning, 1949
koellikeri (Semper, 1868)
= *Cucumaria köllikeri* Semper, 1868
= *Pseudocnus dubiosus* (Semper) Panning, 1962
= *Pseudocnus köllikeri* Panning, 1949
lacteus (Forbes & Goodsir, 1839)
= *Cucumaria mosterensis* Grieg, 1889
= *Holothuria lactea* Forbes & Goodsir, 1839
petiti (Cherbonnier, 1958)
= *Ludwigia petiti* Cherbonnier, 1958
planci (Brandt, 1835)
= ?*Holothuria brunneus* (W. Thompson, 1840, ex Forbes MS), non Chamisso & Eysenhardt, 1821
= ?*Ocnus brunneus* Forbes & Goodsir, in Forbes, 1841
= *Cladodactyla plonci* Brandt, 1835
syracusana (Grube, 1840) Panning, 1949
= *Cladodactyla syracusana* Grube, 1840
= *Cucumaria hyndmanni* W. Thompson, 1840
= *Cucumaria syracusana* auctt.
= *Pseudocnus syracusana* auctt.
- Paracucumaria*
mauritanica (Hérouard, 1929)
= *Cucumaria mauritanica* Hérouard, 1929
thalassae Cherbonnier, 1969
- Pawsonia*
saxicola (Brady & Robertson, 1871)
= *Colochirus lacazei* Hérouard
= *Cucumaria saxicola* Brady & Robertson, 1871
= *Holothuria montagui* Fleming, 1828 (p.p.)
aff. *saxicola* Garrido (email check-list of Canarian echinoderms)
- Pentamera*

- calcigera* (Stimpson, 1851)
 = *Cucumaria koreni* Lütken, 1857
 = *Pentacta calcigera* Stimpson, 1851
- Pseudothyone*
furnesti Cherbonnier, 1969
raphanus (Düben & Koren, 1846)
 = *Thyone poucheti* Barrois, 1882
 = *Thyone raphanus* Düben & Koren, 1846
sculponea Cherbonnier, 1958
serrifera (Östergren, 1898) Jensenius Madsen & Hansen, 1994
 = *Thyone serrifera* Östergren, 1898
- Stereoderma*
incerta Cherbonnier, 1969
kirchsbergii (Heller, 1868)
 = *Cucumaria kirchsbergii* Heller, 1868
 = *Ocnus?* *kirchsbergii* von Marenzeller, 1874
- Thyone*
cherbonnieri Reys, 1960
fusus (O.F. Müller, 1776)
 = ?*Thyone flexus* Hodge, 1867
 = *Holothuria fusus* O.F. Müller, 1776
 = *Holothuria papillosa* Abildgaard, 1789
 = *T. subvillosa* Hérouard, 1890
fusus subsp. *mediterranea* Madsen, 1941
gadeana R. Perrier, 1898
 = *T. wahrbergii* Madsen, 1941
roscovia Hérouard, 1890
- Leptopentacta*
elongata (Düben & Koren, 1846)
 = ?*Cucumaria cucumis* (Risso, 18??)
 = *Cucumaria elongata* Düben & Koren, 1846
 = *Cucumaria pentactes* Forbes, 1841 non (Linnaeus, 1767)
tergestina (M. Sars, 1857)
 = *Cucumaria incurvata* Perrier, 1902
 = *Cucumaria tergestina* M. Sars, 1857
- Order Dactylochirotida**
- Family Ypsilothuriidae**
- Echinocucumis*
hispida (Barrett, 1857)
 = *Cucumaria hispida* (Barrett, 1857)
 = *E. typica* M. Sars, 1859
 = *Eupyrgus hispidus* Barrett, 1857
- Ypsilothuria*
bitentaculata subsp. *attenuata* E. Perrier, 1886
talismani subsp. *talismani* E. Perrier, 1886
 = ?*Echinocucumis hispida* var. *abyssalis* Koehler, 1896
- Order Aspidochirotida**
- Family Stichopodidae**
- Eostichopus*
regalis (Cuvier, 1817)
 = *Holothuria columnae* Cuvier, 1817
 = *Holothuria regalis* Cuvier, 1817
 = *Holothuria triquetra* Delle Chiaje, 1841
 = *Stichopus regalis* (Cuvier, 1817)
- Parastichopus*
tremulus (Gunnerus, 1767)
 = ?*Stichopus griegi* Östergren, 1896
 = *Holothuria ecalcarea* M. Sars, 1859
 = *Holothuria elegans* O.F. Müller, 1781
 = *Holothuria tremula* Gunnerus, 1767
 = *Stichopus richardi* Hérouard, 1896
 = *Stichopus tremulus* (Gunnerus)
- Family Synallactidae**
- Bathyplotes*
natans (M. Sars, 1868) Östergren, 1896
 = *B. assimilis* Koehler & Vaney, 1905
 = *B. heterostylides* Heding, 1942
 = *B. ostergreni* Ohshima, 1915
 = *B. papillosus* Koehler & Vaney, 1905
 = *B. patagiatus* Fisher, 1907
 = *Holothuria natans* M. Sars, 1868
- = *Stichopus moseleyi* Théel, 1886
pourtalesii (Théel, 1886)
 = *B. bipartitus* Hérouard, 1923
 = *Stichopus pourtalesii* Théel, 1886
tizardi (Théel, 1882)
 = *B. fallax* Östergren, 1896
 = *B. reptans* R. Perrier, 1902
 = *Stichopus tizardi* Théel, 1882
- Benthothuria*
junebris R. Perrier, 1898
 = *Benthothuria valdiviae* Heding, 1940
- Gephyrothuria*
alcocki Koehler & Vaney, 1905
 = *G. europeensis* Hérouard, 1923
- Mesothuria*
bifurcata Hérouard, 1906
candelabrum Hérouard, 1923
cathedralis Heding, 1940
connectens (R. Perrier, 1898)
 = *Zygothuria connectens* R. Perrier, 1898
expectans R. Perrier, 1898
intestinalis (Ascanius, 1805) Östergren, 1896
 = *Fistularia mollis* M. Sars, 1835
 = *Holothuria intestinalis* Ascanius, 1805
 = *Thyonidium scabrum* M. Sars, 1868
lactea (Théel, 1886)
 = ?*Holothuria aspera* Bell, 1892
 = *Holothuria lactea* Théel, 1886
 = *Zygothuria lactea* (Théel, 1886)
maroccana Perrier, 1902
murrayi var. *grandipes* Hérouard
roulei (Koehler, 1896)
 = *Holothuria roulei* Koehler, 1896
rugosa Hérouard, 1912
triradiata Heding, 1942
verrilli (Théel, 1886)
 = *Holothuria verrilli* Théel, 1886
 = *M. oktaknemus* Sluiter, 1901
- Meseres*
acaudum Heding, 1935
ingolfi (Heding, 1940)
 = *Plicastichopus ingolfi* Heding, 1942
- Paelopaites*
atlantica Hérouard, 1902
grisea R. Perrier, 1902
- Paroriza*
pallens (Koehler, 1896)
 = *Stichopus pallens* Koehler, 1896
prahoi Hérouard, 1902
- Pseudostichopus*
atlanticus R. Perrier, 1898
depressus Hérouard, 1902
globigerinae Hérouard, 1902
lapidus Hérouard
marenzelleri Hérouard, 1902
occultatus von Marenzeller, 1893
vilosus Théel, 1886
- Scotothuria*
herringi Hansen, 1978
 = ?*Dendrothuria similis* Koehler & Vaney, 1905
- Synallactes*
crucifera R. Perrier, 1898
- Family Holothuriidae**
- Gastroturia*
limbata R. Perrier, 1899
- Holothuria*
(Roweothuria) arguinensis Koehler & Vaney, 1906
dakarensis Panning, 19??
forskali Delle Chiaje, 1823
 = *H. catanensis* Grube, 1840
 = *H. nigra* Peach, 1845
 = *Stichopus selenkae* Barrois, 1882
helleri von Marenzeller, 1878
 = *H. affinis* Heller, 1868, non Brandt, 1835
(Thymiosicya) impatiens (Forskål, 1775)

- = *Fistularia impatiens* Forskål, 1775
 = *H. botellus* Selenka, 1867
 = *H. impatiens* subsp. *bicolor* Clark, 1938
 = *H. impatiens* subsp. *concolor* Clark, 1921
 = *H. impatiens* subsp. *lutea* Clark, 1921
 = *H. impatiens* subsp. *pulchra* Clark, 1921
 = *H. truncata* Lampert, 1885
(Paneyothuria) lentiginosa von Marenzeller, 1892
mammata Grube, 1840
(Halodeima) mexicana Ludwig, 1875
(Platyperona) parvula (Selenka, 18???)
polii Delle Chiaje, 1823
sanctori Delle Chiaje, 1823
= *H. sarcimen* Selenka, 1867
(Holothuria) stellata Delle Chiaje, 1823
(Holothuria) tubulosa Gmelin, 1788
= *H. cavelinii* Delle Chiaje, 1841
= *H. columnae* Delle Chiaje, 1823, non Cuvier, 1817
= *H. maxima* Delle Chiaje, 1823
= *H. pentagnaiae* Delle Chiaje, 1823
- Order Elasipodida**
- Family Elpidiidae**
Achlyonice
myriamae Gebruk, 1997
Amperima
furcata (Hérouard, 1899)
= *Kolga furcata* Hérouard, 1899
rosea (R. Perrier, 1896)
= *Periamma rosea* R. Perrier, 1896
Ellipinion
delagei (Hérouard, 1896)
= *Scotoplanes delagei* Hérouard, 1896
Elpidia
glacialis Théel, 1876
echinata (R. Perrier, 1896)
= *Tutela echinata* R. Perrier, 1896
Irpa
abyssicola Danielssen & Koren, 1878
Kolga
hyalina Danielssen & Koren, 1879
= *Elpidia nana* Théel, 1879
Peniagone
azorica von Marenzeller, 1893
diaphana (Théel, 1882)
= *Scotoanassa translucida* Hérouard, 1899
gracilis (Ludwig, 1894)
islandica Deichmann, 1930
porcella R. Perrier, 1900
purpurea (Théel, 1882)
= *P. ferruginea* Grieg, 1921
= *P. vexillum* R. Perrier, 1900
wyvillii Théel, 1882
Penilpidea
ludwigi (von Marenzeller, 1893)
= *Kolga ludwigi* von Marenzeller, 1893
Psychroplanes
obsoleta (Hérouard, 1899)
= *Kolga foliacea* Hérouard, 1912
= *Kolga obsoleta* Hérouard, 1899
= *Peniagone foliacea* (Hérouard, 1912)
= *Peniagone obsoleta* (Hérouard, 1899)
- Family Deimatidae**
Deima
validum subsp. *validum* Théel, 1879
= *D. atlanticum* Hérouard, 1898
= *D. blakei* Théel, 1886
Oneirophanta
mutabilis Théel, 1879
= *O. alternata* R. Perrier, 1900
- Family Laetmogonidae**
Benthogone
rosea Koehler, 1896
= *B. quadrilineata* Heding, 1940
Laetmogone
- = *Violacea* Théel, 1879
= *Cryodora spongiosa* Théel, 1879
= *L. bronniarti* E. Perrier, 1886
= *L. journainii* Petit, 1885
= *L. wyvillethomsoni* Théel, 1879
Pannychia
glutinosa Hérouard, 1902
- Family Psychropotidae**
Benthodytes
janthina von Marenzeller, 1893
= *B. lingua* R. Perrier, 1896
typica Théel, 1882
= *B. glutinosa* R. Perrier, 1902
Psychropotes
depressa (Théel, 1882)
= *Euphronides auricularia* R. Perrier, 1902
= *Euphronides cornuta* Verrill, 1884
= *Euphronides talismani* R. Perrier, 1896
= *Euphronides violacea* Perrier, 1902
grimaldii Hérouard, 1896
longicauda Théel, 1882
= *Nectothuria translucida* Belyaev & Vinogradov, 1969
= *P. buglossa* R. Perrier, 1896
= *P. furcata* R. Perrier, 1896
semperviana Théel, 1882
= *Euphronides anchora* Hérouard, 1912
= *Euphronides kerhervei* (Hérouard, 1896)
= *P. kerhervei* Hérouard, 1896
- Family Pelagothuridae**
Enypniastes
eximia Théel, 1882
= *Pelagothuria bouvieri* Hérouard, 1906
= *Plankothuria diaphana* Gilchrist, 1920
- Order Apodida**
- Family Synaptidae**
Epitomapta
roseola (Verrill, 1874)
= *Leptosynapta roseola* Verrill, 1874
Euapta
lappa (J. Müller, 1850)
= *Synapta lappa* J. Müller, 1850
Labidoplax
buskii (M'Intosh, 1866)
= *Synapta buskii* M'Intosh, 1866
= *Synapta tenera* Norman, 1865 (*nomen nudum*)
digitata (Montagu, 1815)
= *Holothuria digitata* Montagu, 1815
= *L. (Oestergrenia) adriatica* Heding, 1931
digitata var. *profundicola* Kemp, 1905
media Östergren, 1905
mortensi Heding, 1931
= *L. (Oestergrenia) mortensi* Heding, 1931
similimedia Gage, 1985
southwardorum Gage, 1985
thomsoni (Heraphath, 1865)
= *Synapta digitata* var. *thomsonii* Heraphath, 1865
Leptosynapta
bergensis (Östergren, 1905)
= *Synapta bergensis* Östergren, 1905
decaria (Östergren, 1905)
= *Synapta decaria* Östergren, 1905
gallienii (Heraphath, 1865)
= *Synapta gallienii* Heraphath, 1865
= *Synapta sarniensis* Heraphath, 1865
inhaerens (O.F. Müller, 1776)
= *Holothuria inhaerens* O.F. Müller, 1776
= *Synapta duvernaea* de Quatrefages, 18??
makrankyra (Ludwig, 1898)
= *Synapta makrankyra* Ludwig, 1898
minuta (Becher, 1906)
= *Synapta minuta* Becher, 1906
Protankyra
abyssicola (Théel, 1886)
= *Synapta abyssicola* Théel, 1886

- Rhabdomolgus*
ruber Keferstein, 1863
- Synapta*
reciproca (Forskal, 1775)
 = *Synapta reciproca* Forskål, 1775
- Family Chiridotidae**
- Achiridota*
profunda Heding, 1935
- Chiridota*
abyssicola von Marenzeller, 1893
laevis (O. Fabricius, 1780)
 = *C. groenlandica* Heding, 1935
 = *C. spirourna* Heding, 1935
 = *Holothuria laevis* O. Fabricius, 1780
 = *Holothuria pellucida* Vahl, 1806
- Trochodota*
furcipraedita Salvini Plawen, 1972
venusta (Semon, 1887)
 = *Chiridota venusta* Semon, 1887
- Family Myriotrochidae**
- Acanthotrochus*
mirabilis Danielssen & Koren, 1879
- Myriotrochus*
bathybius H.L. Clark, 1920
clarki Gage & Bilett, 1986
eurycyclus Heding, 1935
geminiradiatus Salvini-Plawen, 1977
giganteus Clark, 1920
rinkii Steenstrup, 1851
theeli Östergren, 1905
vitreus (M. Sars, 1872)
 = *Chiridota brevis* Huxley, 1852
 = *Chiridota laevis* C.G.J. Petersen, 1898
 = *Oligotrochus vitreus* M. Sars, 1872
- Parvotrochus*
belyaevi Gage & Bilett, 1986
- Siniotrochus*
myriodontus Gage & Bilett, 1986
- Trochoderma*
elegans Théel, 1877
- Order Molpadiida**
- Family Molpadiidae**
- Cherbonniera*
utriculus Sibuet, 1974
- Molpadia*
blakei (Théel, 1886)
 = *Paratrichostoma spiniferum* Heding, 1932
 = *Trochostoma blakei* var. *excentrica* Hérouard
 = *Trochostoma blakei* Théel, 1886
 = *Trochostoma grossularia* Hérouard
borealis M. Sars, 1859
 = *Ankyroderma affine* Danielssen & Koren, 1882
 = *Ankyroderma jeffreysii* Danielssen & Koren, 1878
 = *Haplodactyla arctica* von Marenzeller, 1877
 = *M. (Paramolpadia) diploa* Heding, 1935
 = *M. oolitica* Mortensen, 1914 non Pourtales, 1851
 = *Trochostoma arctica* Danielssen & Koren, 1880 non
 (von Marenzeller, 1877)
 = *Trochostoma boreale* (M. Sars)
 = *Trochostoma Thomsonii* Danielssen & Koren, 1879
loricata (R. Perrier, 1898)
 = *Ankyroderma loricatum* Perrier, 1898
maroccana (Perrier, 1898)
 = *Ankyroderma maroccana* Perrier, 1898
musculus Risso, 1826
 = *Ankyroderma danielsseni* Théel, 1886
 = *Ankyroderma hispanicum* Petit
 = *Ankyroderma perrieri* Petit
 = *Eumolpadia asaphes* Heding, 1935
 = *Haplodactyla mediterranea* Grube, 1840
 = *Trochostoma violaceum* Théel, 1886
oolitica (Pourtales, 1851)
- Family Eupyrgidae**
- Eupyrgus*

TUNICATA

Compiled by Claude Monniot

The list has been prepared from a personal database of the Ascidiacea and Sorberacea of the world, held in Muséum National d'Histoire Naturelle, Paris. The original database includes synonyms as well as subsequent usage of the names in the literature and the geographical distribution of species. The ERMS list represents the state-of-the-art of Ascidiacean systematics and nomenclature, and this may occasionally deviate from common usage; in cases where a commonly accepted name is no longer valid, it appears in the ERMS list as a synonym. The author thanks David Connor for comparing an earlier draft of the European list with the British List and for pointing out a number of inconsistencies and omissions. P. Lozouet and J. Marmayou checked and edited the list.

Superclass Tunicata

Class Ascidiacea

Order Pleobranchia

Family Agneziidae

- Adagnesia*
- charcoti* Monniot C. & Monniot F., 1973
- rimosa* Monniot C. & Monniot F., 1974

Agnezia

- atlantica* (Monniot C. & Monniot F., 1973)
- celtica* (Monniot C. & Monniot F., 1974)

Proagnesia

- depressa* (Millar, 1955)

Family Ascidiidae

Ascidia

- callosa* Stimpson, 1852
- celtica* Monniot, C., 1969
- conchilega* Müller, 1776
- correi* Monniot C., 1970
- dijmphniana* Templeton, 1834
- iberica* Monniot C. & Monniot F., 1988
- interrupta* Heller, 1878
- involuta* Heller, 1875
- malaca* (Traustedt, 1883)
- mediterranea* Péres, 1959
- mentula* Müller, 1776
- molguloides* Monniot, C., 1975
- muricata* Heller, 1874
- obliqua* Alder, 1863
- prunum* Müller, 1776
- salvatoris* (Traustedt, 1885)
- tritonis* Herdman, 1883
- virginea* Müller, 1776

Ascidia

- aspersa* (Müller, 1776)
- scabra* (Müller, 1776)

Phallusia

- arabica* Savigny, 1816
- colleta* Monniot C. & Monniot F., 1970
- fumigata* Grube, 1864
- ingeria* Traustedt, 1883
- mammillata* (Cuvier, 1815)
- nigra* Savigny, 1816

Psammascidia

- teissieri* Monniot F., 1962

Family Cionidae

Araneum

- sigma* Monniot C. & Monniot F., 1973

Ciona

- edwardsi* (Roule, 1886)
- fascicularis* Hancock, 1870
- gelatinosa* Bonnevie, 1896

- imperfecta* Monniot C. & Monniot F., 1977

intestinalis (Linnaeus, 1758)

Diazona

- violacea* Savigny, 1816

Pseudodiazona

- abyssa* Monniot C. & Monniot F., 1974

Rhopalaea

- hartmeyeri* (Salfi, 1927)

- neapolitana* (Philippi, 1843)

- orientalis*? Péres, 1958

Tylobranchion

- nordgaardi* (Hartmeyer, 1922)

Family Corellidae

Abyssascidia

- millari* Monniot F., 1971

Chelyosoma

- macleayanum* Broderip & Sowerby, 1830

Corella

- borealis* Traustedt, 1886

- halii* Kott, 1951

- parallelogramma* (Müller, 1876)

Corellopsis

- pedunculata* Hartmeyer, 1903

Corynascidia

- hartmeyeri* Monniot C. & Monniot F., 1994

- suhmi* Herdman, 1882

- translucida* (Monniot, C., 1969)

Rhodosoma

- callensis* (Lacaze-Dutbiers, 1865)

- turicum* (Savigny, 1816)

Xenobranchion

- tenue* Monniot F. & Monniot C., 1976

Family Octacnemidae

Dicopia

- antirrhinum* Monniot, C., 1972

Octacnemus

- ingolfi* Madsen, 1947

Situla

- lanosa* Monniot C. & Monniot F., 1973

Family Perophoridae

Ecteinascidia

- herdmani* (Lahille, 1887)

- styloides* (Traustedt, 1882)

- turbanata* Herdman, 1880

Perophora

- bermudensis* Berrill, 1932

- japonica* Oka, 1927

- listeri* Wiegman, 1835

- multiclavata* (Sluiter, 1904)

- viridis* Verrill, 1871

Order Aplousobranchia

Family Clavelinidae

Archidistoma

- aggregatum* Garstang, 1891

- productum* (Milne-Edwards, 1841)

Clavelina

- dellavallei* (Zirpolo, 1825)

- borealis* Savigny, 1816

- lepadiformis* Müller, 1776

- nana* Lahille, 1890

- oblonga* Herdman, 1880

- phlegraea* Salfi, 1929

- sabbadini* Brunetti, 1987

Pycnoclavella

- aurilucens* Garstang, 1891

- taureanensis* Brunetti, 1991

Family Didemnidae

Didemnum

- albidum* (Verrill, 1871)

- amourouxii* Lafargue, 1976

- coccineum* (Drasche, 1883)

- commune* (Della Valle, 1877)

- coriaceum* (Drasche, 1883)

- drachi* Lafargue, 1975

- fulgens* Milne-Edwards, 1841
granulosum (Drasche, 1883)
lahillei Hartmeyer, 1909
maculosum (Milne-Edwards, 1841)
= *Didemnum asperum* (Milne-Edwards, 1841)
perspicuum Giard, 1873
peyrefitense Brément, 1913
polare (Hartmeyer, 1903)
protectum (Daumézon, 1908)
pseudofulgens Médioni, 1970
romssae Marks, 1996
- Diplosoma**
carnosum Drasche, 1883
laflagueae Vazquez, 1993
listerianum (Milne-Edwards, 1841)
ponticum Romanov, 1989
singulare Lafargue, 1968
spongiforme Giard, 1872
- Leptoclinides**
faeroensis Bjerkan, 1905
- Lissoclinum**
aureum Verrill, 1871
perforatum (Giard, 1872)
rubrum Monniot F., 1975
weigelei Lafargue, 1968
- Polysyncraton**
amethysteum (Van Name, 1902)
asterix Monniot F., 1975
bilobatum Lafargue, 1968
canetense (Brément, 1913)
haranti Lafargue, 1975
lacazei (Giard, 1872)
- Trididemnum**
cereum (Giard, 1872)
delessertiae Lafargue, 1968
inarmatum (Drasche, 1883)
pedunculatum Lafargue *et al.*, 1993
tenerum (Verrill, 1871)
translucidum (Lafargue, 1968)
= *Didemnopsis translucidum*
- Family Polycitoridae**
- Archascidia**
neapolitana Julin, 1904
- Cystodites**
dellechiaiei Della Valle, 1877
durus Drasche, 1883
guinensis Michaelsen, 1914
- Distaplia**
bermudensis Van Name, 1902
clavata (Sars, 1851)
corolla Monniot F., 1975
livida (Sars, 1851)
magnilarva (Della Valle, 1881)
= *Distaplia stelligera* (Giard, 1899)
rosea Della Valle, 1881
- Eudistoma**
angolanum (Michaelsen, 1915)
banyulense (Brément, 1912)
clarum (Van Name, 1902)
costai (Della Valle, 1877)
magnum Médioni, 1968
mucosum (Drasche, 1883)
paesslerioides (Michaelsen, 1914)
planum Pérès, 1948
plumbeum (Della Valle, 1877)
posidonarium (Daumézon, 1908)
roseum Vazquez & Ramos-Espala, 1993
rubrum (Savigny, 1816)
tridentatum (Heiden, 1894)
vitreum (Sars, 1851)
- Polycitor**
adriaticus (Drasche, 1883)
crystallinus (Renier, 1804)
searli Kott, 1952
- Protoholozoa**
- pigra* Monniot F., 1975
- Rhombifera**
caerulea Pérès, 1956
- Sycozoa**
melopepona Monniot F., 1975
- Family Polyclinidae**
- Aplidiopsis**
atlanticus Monniot F., 1975
vitreum (Lahille, 1887)
- Aplidium**
aegeaensis Koukouras *et al.*, 1995
albicans (Milne-Edwards, 1841)
appendiculatum (Michaelsen, 1923)
argus (Milne-Edwards, 1841)
asperum Drasche, 1883
bermudae (Van Name, 1902)
brementi (Harant, 1925)
caeruleum Lahille, 1890
conicum (Olivier, 1792)
densem (Giard, 1872)
elegans (Giard, 1872)
= *Sidnum elegans*
enigmaticum Monniot C. & Monniot F., 1973
fuscum (Drasche, 1883)
gibbosum (Savigny, 1816)
glabrum (Verrill, 1871)
griseum Lahille, 1890
haouarianum (Pérès, 1956)
hyalinum (Pérès, 1956)
inversum (Pérès, 1959)
lobatum Savigny, 1816
mediterraneum (Hartmeyer, 1909)
mutabile (Sars, 1851)
nema Monniot F. & Monniot C., 1976
nordmani (Milne-Edwards, 1841)
ocellatum Monniot C. & Monniot F., 1987
pallidum (Verrill, 1871)
polytrema (Monniot C. & Monniot F., 1983)
proliferum (Milne-Edwards, 1841)
pseudolobatum (Pérès, 1956)
punctum (Giard, 1873)
radicosum (Monniot C. & Monniot F., 1979)
sagresensis Ramos-Espala, Turon & Vazquez, 1993
schaudinni Hartmeyer, 1903
spitzbergense Hartmeyer, 1903
tabarquensis Ramos-Espala, 1991
translucidum (Ritter, 1901)
tridentatum (Daumézon, 1909)
turbinatum (Savigny, 1816)
= *Sydnium turbinatum*
urgorii Vazquez, 1994
- Aplidum**
punctum (Giard, 1872)
- Polyclinella**
azemai Harant, 1930
- Polyclinum**
aurantium Milne-Edwards, 1841
saturnium Savigny, 1816
sibiricum Redikorzev, 1907
- Pseudodistoma**
arnbacki Pérès, 1959
crucigaster Gaill, 1972
cyrnusense Pérès, 1952
obscurum Pérès, 1959
- Ritterella**
glareosa Monniot F., 1975
- Sidneioides**
ivicense Pérès, 1957
- Synoicum**
beauchampi (Harant, 1927)
blochmanni (Heiden, 1894)
daicum Monniot C. & Monniot F., 1977
diaphanum Sluiter, 1927
incrassatum (Sars, 1851)
lacazei (Pérès, 1957)

- pomum* (Sars, 1851)
pulmonaria (Ellis & Solander, 1786)
turgens Phipps, 1774
vibei Lützen, 1959
- Order Stolidobranchia**
- Family Molgulidae**
- Anomopera*
ingolfiana Hartmeyer, 1923
- Eugyra*
adriatica Drasche, 1884
arenosa (Alder & Hancock, 1848)
pedunculata Traustedt, 1886
- Eugyrioides*
borealis (Monniot C. & Monniot F., 1977)
glutinans (Müller, 1842)
islandica (Millar, 1974)
- Gamaster*
dakarensis Pizon, 1896
- Molgula*
amesophleba (Codreanu & Mack-Fira, 1956)
appendiculata Heller, 1877
bleizi (Lacaze-Duthiers, 1877)
caminae Monniot C. & Monniot F., 1988
celtica Monniot, C., 1970
citrina Alder & Hancock, 1848
complanata Alder & Hancock, 1870
crustosa Monniot C. & Monniot F., 1988
dextrocarpa Monniot C. & Monniot F., 1974
echinosiphonica Lacaze-Duthiers, 1877
euprocta Drasche, 1884
griffithsi (Mac Leay, 1825)
helleri Drasche, 1884
herdmani Bjerkan, 1905
hirta Monniot F., 1965
impura Heller, 1877
kiaeri Hartmeyer, 1901
kolaensis Årnäs, 1928
manhattensis (De Kay, 1843)
mira (Årnäs, 1931)
occidentalis Traustedt, 1882
occulta Kupffer, 1875
oculata Forbes, 1848
plana Monniot, C., 1971
platybranchia Monniot, 1970
pumila Monniot F. & Monniot C., 1976
retortiformis Verrill, 1871
rheophila (Pérès, 1956)
romeri Hartmeyer, 1903
roulei Monniot C., 1970
satyrus Monniot C. & Monniot F., 1993
simplex Alder & Hancock, 1870
siphonalis Kiae, 1896
siphonata Alder, 1850
socialis Alder, 1863
solenata (Lacaze-Duthiers, 1877)
tethys Monniot F. & Monniot C., 1974
tubifera Orsted, 1844
vara Monniot C. & Monniot F., 1979
- Molguloides*
crenatum Monniot C. & Monniot F., 1974
- Pareugyrioides*
charadyi Monniot C. & Monniot F., 1977
- Protomolgula*
bythia Monniot F., 1971
triangularis Millar, 1982
- Family Pyuridae**
- Bathypyura*
celata Monniot C. & Monniot F., 1973
- Boltenia*
echinata (Linnaeus, 1767)
ovifera (Linnaeus, 1767)
pilosa (Millar, 1955)
- Bolteniopsis*
prenanti Harant, 1927
- sessilis* Monniot C. & Monniot F., 1970
- Cratostigma*
gravellophila (Pérès, 1955)
intermedia Vazquez & Ramos-Espal, 1993
regularis Monniot, C., 1963
simplex Millar, 1982
singularis (Van Name, 1912)
vestigialis Turon, 1988
- Culeolus*
suhmi Herdman, 1881
- Dextrogaster*
suecica Monniot F., 1962
- Halocynthia*
papillosa (Linnaeus, 1767)
pyriformis (Rathke, 1806)
spinosa Sluiter, 1905
- Herdmania*
momus (Savigny, 1816)
- Heterostigma*
fagei Monniot C. & Monniot F., 1961
gonochorica Monniot F., 1965
mediterranea Pérès, 1958
melitensis Monniot F. & Monniot C., 1976
reptans Monniot C. & Monniot F., 1963
separ Årnäs, 1924
- Microcosmus*
claudicans (Savigny, 1816)
exasperatus Heller, 1878.
glacialis (Sars, 1859)
hennius (Monniot C. & Monniot F., 1974)
nudistigma Monniot C., 1962
polymorphus Heller, 1877
pupa (Savigny, 1816)
sabatieri Roule, 1885
savignyi Monniot, 1962
vulgaris Heller, 1877
- Minipera*
papillosa Monniot C. & Monniot F., 1974
pedunculata Monniot C. & Monniot F., 1974
- Pyura*
dura Heller, 1877
gangelion (Savigny, 1816)
microcosmus (Savigny, 1816)
squamulosa (Alder, 1863)
tessellata (Forbes, 1848)
- Family Styelidae**
- Alloeocarpa*
loculosa Monniot, C., 1975
- Bathyconus*
mirabilis Herdman, 1882
- Bathystyeloides*
dubius Monniot C. & Monniot F., 1974
enderbyanus (Michaelsen, 1904)
laubieri Monniot C. & Monniot F., 1974
- Botryllus*
aureus Sars, 1851
leachi Savigny, 1816
 $=$ *Botrylloides leachi*
schlosseri (Pallas, 1766)
- Cnemidocarpa*
annectens (Hartmeyer, 1921)
bathyphila Millar, 1955
bythia (Herdman, 1881)
cirrata Årnäs, 1922
cipeata (Årnäs, 1921)
devia Årnäs, 1931
digonas Monniot C. & Monniot F., 1968
finmarkiensis (Kiae, 1893)
hartmeyeri Michaelsen, 1918
margaritifera Hartmeyer, 1918
mollis (Stimpson, 1852)
mollispina Årnäs, 1922
mortensenii Hartmeyer, 1912
platybranchia Millar, 1955
rhizopus (Redikorzev, 1907)

<i>sabulosa</i> Glémarec & Monniot C., 1966	<i>thalassae</i> Monniot, C., 1969	
<i>squamata</i> Lützen, 1970		
<i>Dendrodoa</i>		
<i>aggregata</i> Müller, 1776		
<i>carnea</i> (Rathke, 1806)		
<i>grossularia</i> (Van Beneden, 1846)		
<i>lineata</i> (Traustedt, 1880)		
<i>minuta</i> (Bonnevie, 1896)		
<i>pulchella</i> (Rathke, 1806)		
<i>uniplicata</i> (Bonnevie, 1896)		
<i>Dicarpa</i>		
<i>atlantica</i> Millar, 1964		
<i>Distomus</i>		
<i>fuscus</i> (Delle Chiaje, 1841)		
<i>hupferi</i> (Michaelsen, 1904)		
<i>variolosus</i> Gaertner, 1774		
<i>Eusynstyela</i>		
<i>beuzitti</i> (Monniot C., 1970)		
<i>hartmeyeri</i> Michaelsen, 1904		
<i>Kukenthalia</i>		
<i>borealis</i> (Gottschaldt, 1894)		
<i>Pelonia</i>		
<i>corrugata</i> Goodsir & Forbes, 1841		
<i>Polyandrocarpa</i>		
<i>zorritensis</i> (Van Name, 1931)		
<i>Polycarpa</i>		
<i>arnbackae</i> Monniot F., 1964		
<i>biscayensis</i> Monniot C. & Monniot F., 1974		
<i>caudata</i> Monniot C. & Monniot F., 1975		
<i>comata</i> (Alder, 1863)		
<i>discoidea</i> Heller, 1877		
<i>errans</i> Hartmeyer, 1909-1911		
<i>fibrosa</i> (Stimpson, 1852)		
<i>gracilis</i> Heller, 1877		
<i>itera</i> Monniot C. & Monniot F., 1977		
<i>kornogi</i> Glémarec & Monniot C., 1966		
<i>mamillaris</i> (Gaertner, 1774)		
<i>offa</i> Monniot C. & Monniot F., 1988		
<i>pentarhiza</i> Monniot F., 1965		
<i>poraria</i> (Savigny, 1816)		
<i>porculus</i> Monniot C. & Monniot F., 1979		
<i>pseudoalbatrossi</i> Monniot C. & Monniot F., 1968		
<i>pusilla</i> Herdman, 1884		
<i>rockallensis</i> Millar, 1982		
<i>scuba</i> Monniot C., 1971		
<i>tenera</i> Lacaze-Duthiers & Delage, 1892		
<i>violacea</i> (Alder, 1863)		
<i>Protostyela</i>		
<i>heterobranchia</i> Millar, 1954		
<i>longicauda</i> Monniot, Vazquez, White, 1995		
<i>Psammostyela</i>		
<i>delamarei</i> Weinstein, 1961		
<i>Seriocarpa</i>		
<i>rhizoides</i> Dielh, 1969		
<i>Stolonica</i>		
<i>socialis</i> Hartmeyer, 1903		
<i>Styela</i>		
<i>adriatica</i> Monniot F. & Monniot C., 1976		
<i>atlantica</i> (Van Name, 1912)		
<i>bathybia</i> Bonnevie, 1896		
<i>calva</i> Monniot C., Monniot F. & Millar, 1976		
<i>canopus</i> (Savigny, 1816)		
= <i>Styela partita</i> (Stimpson, 1852)		
<i>chaini</i> Monniot C. & Monniot F., 1970		
<i>charcoti</i> Monniot C. & Monniot F., 1973		
<i>clava</i> Herdman, 1881		
<i>coriacea</i> (Alder & Hancock, 1848)		
<i>crinita</i> Monniot C. & Monniot F., 1973		
<i>gelatinosa</i> Traustedt, 1886		
<i>gibbsii</i> Stimpson, 1864		
<i>loculosa</i> Monniot C. & Monniot F., 1969		
<i>plicata</i> (Lesueur, 1823)		
<i>rustica</i> Linnaeus, 1767		
<i>sigma</i> (Van Name, 1912)		
<i>similis</i> Monniot, C., 1970		
	<i>Tibitin</i>	
	<i>probatus</i> Monniot C. & Monniot F., 1987	
	Class Sorberacea	
	<i>Gasterascidia</i>	
	<i>lyra</i> Monniot C. & Monniot F., 1973	
	<i>sandersi</i> Monniot C. & Monniot F., 1968	
	<i>Hexadactylus</i>	
	<i>arcticus</i> (Hartmeyer, 1923)	
	<i>eunuchus</i> (Monniot F. & Monniot C., 1976)	
	<i>ledanoisi</i> (Monniot F. & Monniot C., 1990)	
	<i>Sorbera</i>	
	<i>unigonas</i> Monniot C. & Monniot F., 1974	
	Thaliacea	
	Compiled by Jacob van der Land and R. W. M. van Soest	
	The European list was derived from URMO, which is based on Godeaux (1998) and van Soest (1981).	
	References	
	Godeaux, J., 1998. The relationships and systematics of the Thaliacea, with keys for identification. In: Q. Bone (ed.), <i>The Biology of Pelagic Tunicates</i> : 273-294. Oxford Univ. Press. Oxford.	
	Soest, R.W.M. van, 1981. A monograph of the order Pyrosomatida (Tunicata, Thaliacea). <i>J. Plankton Research</i> 3(4) 1981: 603-631, illustr.	
	Class Thaliacea	
	Family Doliolidae	
	<i>Anchinia</i>	
	<i>savigniania</i> Rathke, 1851	M
	= <i>Doliopsis rubescens</i> Vogt	
	<i>Dolioletta</i>	
	<i>gegenbauri</i> (Uljanin, 1884)	A M
	<i>Doliolina</i>	
	<i>intermedium</i> Neumann	A M
	<i>krohni</i> Herdmann, 1888)	M
	<i>mulleri</i> (Krohn, 1852)	A M
	<i>Dolioloides</i>	
	<i>rarum</i> Gegenbaur	A M
	<i>Doliolum</i>	
	<i>denticulatum</i> Quoy & Gaimard, 1834	A M
	<i>nationalis</i> Borgert, 1894	A M
	Family Pyrosomatidae	
	<i>Pyrosoma</i>	
	<i>aherniosum</i> Seeliger, 1895	A
	<i>atlanticum</i> Péron, 1804	A M
	= <i>Pyrosoma benthica</i> Monniot & Monniot, 1966	
	<i>Pyrostremma</i>	
	<i>agassizi</i> (Ritter & Byxbe, 1905)	A
	<i>spinosum</i> (Herdman, 1888)	A M
	= <i>Pyrosoma sedentarium</i>	
	= <i>Pyrosoma spinosum</i>	
	Family Salpidae	
	<i>Brooksia</i>	
	<i>rostrata</i> (Traustedt, 1893)	
	<i>Cyclosalpa</i>	
	<i>affinis</i> (Chamisso, 1819)	A
	<i>floridana</i> (Apstein, 1894)	A
	<i>foxtoni</i> van Soest, 1974	A
	<i>pinnata</i> (Forskål, 1775)	A M
	<i>polae</i> Sigl, 1912	A
	<i>Helicosalpa</i>	
	<i>virgula</i> (Vogt, 1854)	A
	<i>Ihlea</i>	
	<i>punctata</i> (Forskål, 1775)	A M
	<i>Pegea</i>	
	<i>bicaudata</i> (Quoy & Gaimard, 1826)	A
	<i>confoederata</i> (Forskål, 1775)	A M
	= <i>Pegea socia</i>	

<i>Ritteriella</i>			
<i>amboinensis</i> (Apstein, 1904)	A	<i>stygius</i> Garstang, 1937	A
<i>retracta</i> (Ritter, 1906)	A M	<i>Folia</i>	
<i>Salpa</i>		<i>gracilis</i> Lohmann, 1892	A M
<i>aspera</i> Chamisso, 1819	A	<i>mediterranea</i> (Lohmann, 1899)	M
<i>fusiformis</i> Cuvier, 1804	A M	<i>Megalocercus</i>	
<i>maxima</i> Forskål, 1775	A	<i>abyssorum</i> Chun, 1887	A M
<i>younsi</i> van Soest, 1973	A	<i>Mesochordaeus</i>	
<i>Soestia</i>		<i>bahamasi</i> Fenaux & Youngbluth, 1990	A
<i>zonaria</i> (Pallas, 1774)	A M	<i>Mesoikopleura</i>	
= <i>Iasis zonaria</i>		<i>enterospira</i> Fenaux, 1993	A
<i>Thetys</i>		<i>gyroceanis</i> Fenaux, 1993	M
<i>vagina</i> Tilesius, 1802	A M	<i>haranti</i> (Vernières, 1934)	M
= <i>Tetys costata</i>		<i>youngbluthi</i> Fenaux, 1993	A
= <i>Tetys tilesii</i>		<i>Oikopleura</i>	
<i>Thalia</i>		<i>albicans</i> (Leuckart, 1853)	A M
<i>cicar</i> van Soest, 1973	A M	<i>caudaornata</i> (Fenaux & Youngbluth, 1991)	A
<i>democratica</i> (Forskål, 1775)	A M	<i>cophocerca</i> (Gegenbaur, 1855)	A M
<i>orientalis</i> Tokioka, 1937	A	<i>dioica</i> Fol, 1872	A M
<i>Traustedtia</i>		<i>fusiformis</i> Fol, 1872	A M
<i>multotentaculata</i> (Quoy & Gaimard, 1834)	A	<i>gaussica</i> Lohmann, 1905	A
<i>Weelia</i>		<i>gracilis</i> Lohmann, 1896	A M
<i>cylindrica</i> (Cuvier, 1804)	A	<i>inflata</i> (Fenaux & Youngbluth, 1991)	A
= <i>Salpa cylindrica</i>		<i>intermedia</i> Lohmann, 1896	A M
		<i>labradoriensis</i> Lohmann, 1892	A
		<i>longicauda</i> (Vogt, 1854)	A M
		<i>parva</i> Lohmann, 1896	A M
		<i>rufescens</i> Fol, 1872	A M
		<i>vanhoeffeni</i> Lohman, 1896	A
		<i>villafraancae</i> Fenaux, 1992	M
		<i>Pelagopleura</i>	
		<i>gracilis</i> (Lohmann, 1914)	A
		<i>oppressa</i> (Lohmann, 1914)	A
		<i>verticalis</i> (Lohmann, 1914)	A M
		<i>Stegosoma</i>	
		<i>magnum</i> (Langerhans, 1880)	A M

Appendicularia

Compiled by Jacob van der Land

The European list was derived from URMO, which is based on a recent publication by Fenaux (1998). The list has not been checked by an expert on the European species.

References

Fenaux, R., 1998. The classification of Appendicularia. In: Q. Bone (ed.), *The Biology of Pelagic Tunicates*: 295–306. Oxford Univ. Press, Oxford.

Class Appendicularia

Family Fritillariidae

<i>Appendicularia</i>		
<i>sicula</i> Fol, 1874	A	
<i>tregouboffi</i> Fenaux, 1960	M	
<i>Fritillaria</i>		
<i>aberrans</i> Lohmann, 1896	A M	
<i>abjornceni</i> Lohmann, 1909	A M	
<i>aequatorialis</i> Lohmann, 1896	A M	
<i>borealis</i> Lohman, 1896	A M	
<i>charybdas</i> Lohmann in Lohmann & Buckmann, 1926	M	
<i>drygalskii</i> Lohmann in Bückmann, 1923	A M	
<i>fagei</i> Fenaux, 1961	M	
<i>formica</i>	A M	
<i>fraudax</i> Lohmann, 1896	A M	
<i>gracilis</i> Lohmann, 1896	A M	
<i>haplostoma</i> Fol, 1872	A M	
<i>heleneae</i> Bückmann, 1924	A	
<i>megachile</i> Fol, 1872	A M	
<i>messanensis</i> Lohmann in Bückmann, 1924	M	
<i>pellucida</i> Busch, 1851	A M	
<i>tenella</i> Lohmann, 1896	A M	
<i>urticans</i> Fol, 1872	M	
<i>venusta</i> Lohman, 1896	A M	
<i>Tectillaria</i>		
<i>fertilis</i> (Lohmann, 1896)	A M	

Family Kowalewskiidae

<i>Kowalevskia</i>		
<i>oceanica</i> Lohman, 1899	A M	
<i>tenuis</i> Fol, 1872	A M	

Family Oikopleuridae

<i>Althoffia</i>		
<i>tumida</i> Lohmann, 1892	A M	
<i>Balchochordae</i>		
<i>charon</i> Chun, 1900	A M	

CEPHALOCHORDATA

Compiled by Jacob van der Land

The European list was derived from URMO, which is based on Poss & Boschung (1996). The list has not been checked by an expert on the European species.

References

- Poss, S.G. & H.T. Boschung, 1996. Lancelets (Cephalochordata: Branchiostomatidae): How many species are valid? In: C Gans et al. (eds), The lancelets (Cephalochordata): A new look at some old beasts. - Israel J. Zool. 42 (suppl.): S13-S66.

SUBPHYLUM CEPHALOCHORDATA

Family Branchiostomidae

<i>Epigonichthys lucayanum</i> (Andrews, 1893)	A
<i>Branchiostoma lanceolatum</i> (Pallas, 1774)	A M

PISCES

Compiled by Nicolas Bailly, William N. Eschmeyer, Rainer Froese, Jean-Claude Quéro, Jacob van der Land, Mark J. Costello, Dusan Zavodnik, Riccardo Serrão Santos and Filipe Mora Porteiro.

Three lists of fish in European seas are provided: Osteichthyes (bony fish); Chondrichthyes (cartilagenous fish such as sharks); and Agnatha (lampreys and hagfish). The lists, based on Hureau (1995-96), an updated version of Whitehead *et al.* (1984-86), were drafted by Jacob van der Land and Mark J. Costello, with assistance from Louise Collier, with additions from the Adriatic and Baltic provided by Dusan Zavodnik, and for the Azores by Riccardo Serrão Santos and Filipe Mora Porteiro. Filipe Mora Porteiro also carefully corrected the list for errors. We also thank Giambattista Bello for his helpful suggestions concerning the Chondrichthyes, and Bruno Chanet, Martine Desoutter, Bo Fernholm, Arnaud Filleul, Jean-Claude Hureau, Yuri Sazonov, Bernard Séret for their helpful comments. Jean-Claude Quéro provided his own updated list survey for the North-East Atlantic north to the 36° parallel.

The orders and families follow Eschmeyer (1998) strictly. The species names and authorities also follow Eschmeyer (1998) for the most part. The FishBase (FB) web site is a useful source of further information on the species (<http://www.fishbase.org>). Nicolas Bailly checked the list with the procedure "Check Names" within FB 99 on CD-ROM (Froese and Pauly 1999), then compared with Jean-Claude Quéro's list, and the MNHN fish collection database GICIM (<http://www.mnhn.fr/base/gicim.html>), then checked again with FB 2000 current version in early June 2000. The reader is directed to FB and CoF for the many synonyms of the species. Future lists could include synonyms still in common use, and in addition, species which have recently invaded the European fauna. For example, the International Commission for the Scientific Exploration of the Mediterranean Sea (CIESM) lists 29 fish species for the Mediterranean that are additional to the present list (<http://www.ciesm.org/atlas>). Most of these species have entered from the Indian Ocean via the Suez Canal.

All the subgenera of *Raja* have been upgraded to genus level according to McEachran and Dunn (1998). All the undescribed species, i.e. those noted as "aff." or "sp." have been omitted in the list, but are noted below. In most cases, the letter "i" at the end of species epithets referring to a persons name follow Eschmeyer (1998). We have tried to use the original spelling, even when not only the ending, but also when the whole name of a person was involved (e.g. *Parablennius ruxii* for Roux). In some cases, the names can be a noun in apposition, and do not need an ending "i" at all (e.g., *Arctozenus risso*, *Galeocerdo cuvier*, *Lophotus lacepede*, *Notacanthus bonaparte*, *Squalus blainville*, etc). If the final "i" or a more used spelling should be desirable, the emendation must be submitted to the International Commission of Zoological Nomenclature according to the prevailing use rule.

There remain some issues which should be solved by future publications (the names chosen here are cited first) before a final endorsement:

- presence of species: *Alepocephalus* aff. *australis* (not in the list, new species?), *Antennarius radiosus* (accidental), *Epinephelus nigritus* (accidental), *Monolene microstoma* (not in the list), *Mobula tarapacana*, *Polymetme corythaeola* (misidentification for *P. thaeocoryla*?), *Sphoeroides spengleri* (not in the list, misidentification for *S. marmoratus*?), *Sphyraena barracuda* (not in the list), *Squalus megalops* (new species?);
- validity of genus: *Apghanius*/Lebias, *Macropodus*, *Dalatias*/*Scymnorhinus*, *Diceratias*/*Phrynicthys*;
- validity of species: *Bathyraja* sp. (not in the list, to be described), *Centrophorus uyato*, *Cepola rubescens* (here considered as a junior synonym of *C. macrophthalmus*); *Chromogobius* sp. (not in the list, to be described), *Gaidropsarus* sp. (not in the list, to be described), *Hippocampus ramulosus/guttulatus*, *Leptoderma* sp. (not in the list, to be described), *Neoraja* sp. (not in the list, to be described), *Psednos* sp. (not in the list, to be described);
- genus assignation: *Arctozenus*/*Notolepis* *risso*, *Callionymus*/*Calliurichthys* *filamentosus*, *Histiobranchus*/*Synaphobranchus* *bathybius*, *Platytroctes*/*Platytroctegen* *mirus*, *Scophthalmus maximus*/*Psetta maxima*, *Solea*/*Synapturichthys klenii*, *Synchiropus*/*Foetorepus* *phaeton*, *Trachinus*/*Echiiithys* *vipera*;
- subspecies elevated to species level (only for practical layout reason, should be correctly considered as subspecies): *Apletodon bacescui* (subsp. *A. dentatus*), *Diplecogaster euxinica*, *D. pectoralis* (subsp. *D. bimaculata*), *Eumicrotremus egginsi* (subsp. *E. spinosus*), *Lepadogaster purpurea* (subsp. *L. lepadogaster*), *Molva macrophthalmus* (subsp. *M. dypterygia*), *Trachyscorpia echinata* (subsp. *T. cristulata*), *Tylosurus imperialis* (subsp. *T. acus*);
- senior synonym: *Lethenteron camtschaticum/japonicum*;
- homonym: subgenus *Chalinura*, *Raja africana*;
- spelling: *Careproctus reinhardtii/reinhardti*, *Canthidermis maculata/maculatus*, *Dipturus nidarosiensis/nigarosiensis/nidroensis*, *Kyphosus sectator/sectatrix*, *Leucobrotula adipata/adipatus*, *Microichthys sanzilansanzei*, *Parablennius ruxii/rouxi*, *Pleuronectes platessa/plateussus*, *Salarias basiliscus/basiliscus*.

N. Bailly was responsible for the final decisions on the list of fishes, and should be contacted for comments and corrections (bailly@mnhn.fr).

References

- Eschmeyer, W.N. (ed.) 1998. *Catalog of Fishes*. California Academy of Sciences, 3 vols: 1-2905, with Database CD-ROM.
- Froese, R. and D. Pauly, Editors. 1999. *FishBase 99: concepts, structure et sources de données*. ICLARM, Manilla, Philippines. 324 p.
- Hureau, J.-C. (ed.) 1995-1996. *Fishes of the North-Eastern Atlantic and the Mediterranean*. CD-ROM. ETI, Amsterdam; Springer-Verlag, Berlin. 1995: MacIntosh version; 1996: PC version.
- McEachran, J.D. and Dunn K.A. 1998. *Phylogenetic analysis of skates, a morphologically conservative clade of elasmobranchs* (Chondrichthyes: Rajidae). *Copeia* 1998(2), 271-290.

Santos, R.S., Porteiro, F.M. and Barreiros J.P. 1997. *Marine fishes of the Azores: annotated checklist and bibliography*. Universidade dos Açores, Horta (Açores), Portugal.

Whitehead, P.J.P., Bauchot, M.-L., Hureau J.-C., Nielsen J., and Tortonese E. (eds) 1984-1986. *Fishes of the North-Eastern Atlantic and the Mediterranean*. 3 Vols, 1473 pp, UNESCO, Paris.

Class Agnatha

Order Myxiniformes

Family Myxinidae

Myxine glutinosa Linnaeus, 1758
ios Fernholm, 1981

Order Petromyzontiformes

Family Petromyzontidae

Lampetra fluviatilis (Linnaeus, 1758)
Lethenteron camtschaticum (Tilesius, 1811)
Petromyzon marinus Linnaeus, 1758

Class Chondrichthyes

Order Hexanchiformes

Family Hexanchidae

Heptanchias perlo (Bonnaterre, 1788)
Hexanchus griseus (Bonnaterre, 1788)
nakamurai Teng, 1962

Family Chlamydoselachidae

Chlamydoselachus anguineus Garman, 1884

Order Orectolobiformes

Family Rhincodontidae

Rhincodon typus Smith, 1828

Family Ginglymostomatidae

Ginglymostoma cirratum (Bonnaterre, 1788)

Order Lamniformes

Family Odontaspidae

Carcharias taurus (Rafinesque, 1810)
Odontaspis ferox (Risso, 1810)
noronhai (Maul, 1955)

Family Mitsukurinidae

Mitsukurina owstoni Jordan, 1898

Family Lamnidae

Carcharodon carcharias (Linnaeus, 1758)
Isurus oxyrinchus Rafinesque, 1810
paucus Guitart Manday, 1966

Family Lamnidae

Cetorhinus maximus (Gunnerus, 1765)

Family Alopiidae

Alopias superciliosus (Lowe, 1841)
vulpinus (Bonnaterre, 1788)

Order Carcarhiniformes

Family Scyliorhinidae

- Apristurus***
aphyodes Nakaya & Stehmann, 1998
laurussonii (Saemundsson, 1922)
manis (Springer, 1979)
micros (Gilchrist, 1922)
nasutus de Buen, 1959
- Galeus***
atlanticus Vaillant, 1888
melastomus Rafinesque, 1810
murinus (Collett, 1904)
- Scyliorhinus***
canicula (Linnaeus, 1758)
stellaris (Linnaeus, 1758)
- Family Pseudotriakidae**
Pseudotriakis
microdon Capello, 1867
- Family Leptochariidae**
Leptocharias
smithii (Müller & Henle, 1839)
- Family Triakidae**
Galeorhinus
galeus (Linnaeus, 1758)
Mustelus
asterias Cloquet, 1821
mustelus (Linnaeus, 1758)
punctulatus Risso, 1827
- Family Hemigaleidae**
Paragaleus
pectoralis (Garman, 1906)
- Family Carcharhinidae**
Carcharhinus
altimus (Springer, 1950)
brevipinna (Müller & Henle, 1839)
falciformis (Müller & Henle (ex Bibron), 1839)
galapagensis (Snodgrass & Heller, 1905)
isodon (Müller & Henle (ex Valenciennes), 1839)
leucas (Müller & Henle (ex Valenciennes), 1839)
limbatus (Müller & Henle (ex Valenciennes), 1839)
longimanus (Poey, 1861)
melanopterus (Quoy & Gaimard, 1824)
obscurus (Lesueur, 1818)
plumbeus (Nardo, 1827)
signatus (Poey, 1868)
Galeocerdo
cuvier (Péron & Lesueur in Lesueur, 1822)
- Negaprion*
brevirostris (Poey, 1868)
- Prionace*
glauca (Linnaeus, 1758)
- Rhizoprionodon*
acutus (Rüppell, 1837)
- Family Sphyrnidae**
Sphyraena
lewini (Griffith & Smith, 1834)
mokarran (Rüppell, 1837)
tudes (Valenciennes, 1822)
zygaena (Linnaeus, 1758)
- Order Squaliformes**
- Family Dalatiidae**
Centroscyllium
fabricii (Reinhardt, 1825)
- Etmopterus*
princeps Collett, 1904
pusillus (Lowe, 1839)
spinax (Linnaeus, 1758)
- Centroscymnus*
coelolepis Bocage & Capello, 1864
crepidater (Bocage & Capello, 1864)
cryptacanthus Regan, 1906
- Scymnodalatias*
garricki Kukuyev & Konovalenko, 1988
- Scymnodon*
obscurus (Vaillant, 1888)
- ringens* Bocage & Capello, 1864
- Somniosus***
microcephalus (Bloch & Schneider, 1801)
rostratus (Risso, 1827)
- Oxynotus***
centrina (Linnaeus, 1758)
paradoxus Frade, 1929
- Dalatias***
licha (Bonnaterre, 1788)
- Squaliolus***
laticaudus Smith & Radcliffe in Smith, 1912
- Family Centrophoridae**
Centrophorus
granulosus (Bloch & Schneider, 1801)
lusitanicus Bocage & Capello, 1864
squamosus (Bonnaterre, 1788)
uyato (Rafinesque, 1810)
- Deania***
calcea (Lowe, 1839)
hystricosa (Garman, 1906)
profundorum (Smith & Radcliffe in Smith, 1912)
- Family Squalidae**
Squalus
acanthias Linnaeus, 1758
blainville (Risso, 1827)
megalops (Macleay, 1881)
- Family Echinorhinidae**
Echinorhinus
brucus (Bonnaterre, 1788)
- Order Squatiniformes**
- Family Squatinidae**
Squatina
aculeata, 1829
oculata Bonaparte, 1840
squatina (Linnaeus, 1758)
- Order Pristiformes**
- Family Pristidae**
Pristis
pectinata Latham, 1794
pristis (Linnaeus, 1758)
- Order Torpediniformes**
- Family Torpedinidae**
Torpedo (*Tetronarce*)
nobiliana Bonaparte, 1835
- Torpedo* (*Torpedo*)
marmorata Risso, 1810
torpedo (Linnaeus, 1758)
- Order Rajiformes**
- Family Rhinobatidae**
Rhinobatos
cemiculus Geoffroy Saint-Hilaire, 1817
rhinobatos (Linnaeus, 1758)
- Family Rajidae**
Amblyraja
hyperborea (Collett, 1879)
jensenii (Bigelow & Schroeder, 1950)
radiata (Donovan, 1808)
- Bathyraja*
pallida (Forster, 1967)
richardsoni (Garrick, 1961)
spinicauda (Jensen, 1914)
- Dipturus*
batis (Linnaeus, 1758)
linteus (Fries, 1838)
nidarosiensis (Storm, 1881)
oxyrinchus (Linnaeus, 1758)
- Leucoraja*
circularis (Couch, 1838)
fullonica (Linnaeus, 1758)
melitensis (Clark, 1926)
naevus (Müller & Henle, 1841)

- Malacoraja**
kreffti (Stehmann, 1977)
spinacidermis (Barnard, 1923)
- Neoraja**
caerulea (Stehmann, 1976)
- Raja**
africana, 1977
asterias Delaroche, 1809
brachyura Lafont, 1873
clavata Linnaeus, 1758
maderensis Lowe, 1838
microcellata Montagu, 1818
miraletus Linnaeus, 1758
montagui Fowler, 1910
polystigma Regan, 1923
radula Delaroche, 1809
rondeleti Bougis, 1959
undulata Lacepède, 1802
- Rajella**
bathyphila (Holt & Byrne, 1908)
bigelowi (Stehmann, 1978)
fallae (Litken, 1887)
kukujevi (Dolganov, 1985)
- Rostroraja**
alba (Lacepède, 1803)
- Family Dasyatidae**
Dasyatis
centroura (Mitchill, 1815)
pastinaca (Linnaeus, 1758)
tortonesei Capapé, 1975
violacea (Bonaparte, 1832)
- Himantura**
uarnak (Forsskål, 1775)
- Taeniura**
grabata (Geoffroy Saint-Hilaire, 1817)
- Family Gymnuridae**
Gymnura
altavela (Linnaeus, 1758)
- Family Myliobatidae**
Myliobatis
aquila (Linnaeus, 1758)
- Pteromylaeus**
bovinus (Geoffroy Saint-Hilaire, 1817)
- Rhinoptera**
marginata (Geoffroy Saint-Hilaire, 1817)
- Manta**
birostris (Walbaum, 1792)
- Mobula**
mobular (Bonnaterre, 1788)
tarapacana (Philippi, 1893)
- Order Chimaeriformes**
- Family Chimaeridae**
Chimaera
monstrosa Linnaeus, 1758
- Hydrolagus**
affinis (Capello, 1868)
mirabilis (Collett, 1904)
pallidus Hardy & Stehmann, 1990
- Family Rhinochimaeridae**
Harriotta
haeckeli Karrer, 1972
raleighana Goode & Bean, 1895
- Rhinochimaera**
atlantica Holt & Byrne, 1909
- Class Osteichthyes**
- Order Acipenseriformes**
- Family Acipenseridae**
Acipenser
gueldenstaedtii Brandt & Ratzburg, 1833
naccarii Bonaparte, 1836
- nudiventris** Lovetzký, 1828
stellatus Pallas, 1771
sturio Linnaeus, 1758
- Huso**
huso (Linnaeus, 1758)
- Order Elopiformes**
- Family Megalopidae**
Megalops
atlanticus Valenciennes in Cuvier & Valenciennes, 1847
- Order Notacanthiformes**
- Family Halosauridae**
Aldrovandia
affinis (Günther, 1877)
phalacra (Vaillant, 1888)
- Halosauropsis**
macrochir (Günther, 1878)
- Halosaurus**
johsonianus Vaillant, 1888
overii Johnson, 1864
- Family Notacanthidae**
Notacanthus
bonapartei Risso, 1840
chemnitzi Bloch, 1788
- Polyacanthonotus**
challengeri (Vaillant, 1888)
rissoanus (De Filippi & Verany, 1857)
- Order Anguilliformes**
- Family Anguillidae**
Anguilla
anguilla (Linnaeus, 1758)
- Family Heterenchelyidae**
Panturichthys
fowleri (Ben-Tuvia, 1953)
mauritanicus Pellegrin, 1913
- Family Chlopsidae**
Chlopsis
bicolor Rafinesque, 1810
- Family Muraenidae**
Anarchias
euryurus (Lea, 1913)
Enchelycore
anatina (Lowe, 1838)
- Gymnothorax**
maderensis (Johnson, 1862)
polygonius Poey, 1875
unicolor (Delaroche, 1809)
vicinus (Castelnau, 1855)
- Muraena**
helena Linnaeus, 1758
- Family Synaphobranchidae**
Dysomma
brevirostre (Facciolà, 1887)
- Ilyophis**
arx Robins & Martin in Robins & Robins, 1976
blachei Saldanha & Merrett, 1982
brunneus Gilbert, 1891
- Haptenchelys**
texis Robins & Martin in Robins & Robins, 1976
- Histiobranchus**
bathybius (Günther, 1877)
- Synaphobranchus**
kaupii Johnson, 1862
- Simenchenchelys**
parasitica Gill in Goode & Bean, 1879
- Family Ophichthidae**
Apterichtus
anguiformis (Peters, 1877)
caecus (Linnaeus, 1758)
- Dalophis**
imberbis (Delaroche, 1809)

- Echelus**
myrus (Linnaeus, 1758)
- Ophichthus**
ophis (Linnaeus, 1758)
rufus (Rafinesque, 1810)
- Ophisurus**
serpens (Linnaeus, 1758)
- Pisodonophis**
semicinctus (Richardson, 1848)
- Family Congridae**
- Bathymuroconger**
viginus (Vaillant, 1888)
- Conger**
conger (Linnaeus, 1758)
- Gnathophis**
codoniphorus Maul, 1972
mystax (Delaroche, 1809)
- Pseudophichthys**
splendens (Lea, 1913)
- Ariosoma**
baleanicum (Delaroche, 1809)
- Paraconger**
macrops (Günther, 1870)
- Heteroconger**
longissimus Günther, 1870
- Family Muraenesocidae**
- Cynoponticus**
ferox Costa, 1846
- Muraenesox**
cinereus (Forsskål, 1775)
- Family Derichthyidae**
- Derichthys**
serpentinus Gill, 1884
- Nessorhamphus**
ingolfianus (Schmidt, 1912)
- Family Nemichthysidae**
- Avocettina**
infans (Günther, 1878)
- Nemichthys**
curvirostris (Strömmann, 1896)
scolopaceus Richardson, 1848
- Family Serrivomeridae**
- Serrivomer**
beani Gill & Ryder, 1883
lanceolatoides (Schmidt, 1916)
- Family Nettastomatidae**
- Facciolella**
oxyrhyncha (Bellotti, 1883)
- Nettastoma**
melanurum Rafinesque, 1810
- Saurenchelys**
cancrivora Peters, 1864
- Venefica**
proboscidea (Vaillant, 1888)
- Order Saccopharyngiformes**
- Family Cyematidae**
- Cyema**
atrum Günther, 1878
- Family Saccopharyngidae**
- Saccopharynx**
ampullaceus (Harwood, 1827)
harrisoni Beebe, 1932
hjorti Bertin, 1938
paucovertebratus Nielsen & Bertelsen, 1985
ramosus Nielsen & Bertelsen, 1985
thalassa Nielsen & Bertelsen, 1985
trilobatus Nielsen & Bertelsen, 1985
- Family Eurypharyngidae**
- Eurypharynx**
pelecanoides Vaillant, 1882
- Family Monognathidae**
Monognathus
- bertini** Bertelsen & Nielsen, 1987
boehlkei Bertelsen & Nielsen, 1987
herringi Bertelsen & Nielsen, 1987
jesperseni Bertin, 1936
nigeli Bertelsen & Nielsen, 1987
- Order Clupeiformes**
- Family Clupeidae**
- Alosa**
alosa (Linnaeus, 1758)
caspia (Eichwald, 1838)
fallax (Lacepède, 1803)
pontica (Eichwald, 1838)
- Clupea**
harengus Linnaeus, 1758
- Clupeonella**
cultiriventris (Nordmann, 1840)
- Dussumieria**
acuta Valenciennes in Cuvier & Valenciennes, 1847
- Herklotsichthys**
punctatus (Rüppell, 1837)
- Sardina**
pilchardus (Walbaum, 1792)
- Sardinella**
aurita Valenciennes in Cuvier & Valenciennes, 1847
maderensis (Lowe, 1838)
- Spratelloides**
delicatulus (Bennett, 1832)
- Sprattus**
sprattus (Linnaeus, 1758)
- Family Engraulidae**
- Engraulis**
encrasiculus (Linnaeus, 1758)
- Order Osmeriformes**
- Family Argentinidae**
- Argentina**
silus (Ascanius, 1775)
sphyraena Linnaeus, 1758
- Glossanodon**
leioglossus (Valenciennes in Cuvier & Valenciennes, 1848)
- Family Microstomatidae**
- Microstoma**
microstoma (Risso, 1810)
- Nansenia**
groenlandica (Reinhardt, 1840)
iberica Matallanas, 1985
longicauda Kawaguchi & Butler, 1984
oblita (Facciolà, 1887)
tenera Kawaguchi & Butler, 1984
- Family Bathylagidae**
- Bathylagichthys**
greyae (Cohen, 1958)
- Bathylagus**
euryops Goode & Bean, 1896
- Dolicholagus**
longirostris (Maul, 1948)
- Melanolagus**
bericoides (Borodin, 1929)
- Family Opisthoproctidae**
- Bathylychnops**
exilis Cohen, 1958
- Dolichopteryx**
longipes (Vaillant, 1888)
- Opisthoproctus**
grimaldii Zugmayer, 1911
soleatus Vaillant, 1888
- Rhynchohyalus**
natalensis (Gilchrist & Von Bonde, 1924)
- Family Alepocephalidae**
- Alepocephalus**
agassizii Goode & Bean, 1883
bairdii Goode & Bean, 1879

- productus* Gill, 1883
rostratus Risso, 1820
Asquamiceps
velaris Zugmayer, 1911
Bajacalifornia
calcarea (Weber, 1913)
megalops (Lütken, 1898)
Bathylaco
nigricans Goode & Bean, 1896
Bathypirion
danae Marshall, 1966
Bathytroctes
macrogaster Sazonov, 1999
macrolepis Günther, 1887
michaelsarsi Koefoed, 1927
microlepis Günther, 1878
Conocara
fioneni Sazonov & Ivanov, 1979
macropterum (Vaillant, 1888)
microlepis (Lloyd, 1909)
murrayi (Koefoed, 1927)
salmoneum (Gill & Townsend, 1897)
werneri Nybelin, 1947
Einara
edentula (Alcock, 1892)
macrolepis (Koefoed, 1927)
Herwigia
krefftii (Nielsen & Larsen, 1970)
Leptoderma
macrops Vaillant, 1886
Mirognathus
normani Parr, 1951
Narctes
erimelas Alcock, 1890
stomias (Gilbert, 1890)
Photostylus
pycnopterus Beebe, 1933
Rinocetes
nasutus (Koefoed, 1927)
Rouleina
attrita (Vaillant, 1888)
maderensis Maul, 1948
Talismania
antillarum (Goode & Bean, 1896)
homoptera (Vaillant, 1888)
mekistonema Sulak, 1975
Xenodermichthys
copei (Gill, 1884)
- Family Platytroctidae**
- Barbantus**
curvifrons (Roule & Angel, 1931)
- Holtbyrnia**
anomala Krefft, 1980
macrops Maul, 1957
- Maulisia**
argipalla Matsui & Rosenblatt, 1979
mauli Parr, 1960
microlepis Sazonov & Golovan, 1976
- Mentodus**
rostratus (Günther, 1878)
- Normichthys**
operosus Parr, 1951
- Platytroctes**
apus Günther, 1878
mirus (Lloyd, 1909)
- Sagamichthys**
schnakenbecki (Krefft, 1953)
- Searsia**
koefoedi Parr, 1937
- Family Osmeridae**
- Mallotus**
villosus (Müller ex Olafsen, 1776)
- Osmerus**
eperlanus (Linnaeus, 1758)
mordax (Mitchill, 1814)
- Family Leptochilichthyidae**
Leptochilichthys
agassizii Garman, 1899
pinguis (Vaillant, 1886)
- Order Salmoniformes**
- Family Salmonidae**
- Coregonus**
albus (Linnaeus, 1758)
autumnalis (Pallas, 1776)
lavaretus (Linnaeus, 1758)
sardinella Valenciennes *in* Cuvier & Valenciennes, 1848
- Stenodus**
leucichthys (Güldenstädt, 1772)
- Oncorhynchus**
gorbuscha (Walbaum, 1792)
keta (Walbaum, 1792)
kisutch (Walbaum, 1792)
mykiss (Walbaum, 1792)
- Salmo**
salar Linnaeus, 1758
trutta Linnaeus, 1758
- Salvelinus**
alpinus (Linnaeus, 1758)
- Order Stomiiformes**
- Family Gonostomatidae**
- Bonapartia**
pedaliota Goode & Bean, 1896
- Cyclothona**
acclinidens Garman, 1899
alba Brauer, 1906
braueri Jespersen & Tåning, 1926
livida Brauer, 1902
microdon (Günther, 1878)
obscura Brauer, 1902
pallida Brauer, 1902
pseudopallida Mukhacheva, 1964
pygmaea Jespersen & Tåning, 1926
- Diplophos**
maderensis (Johnson, 1890)
taenia Günther, 1873
- Gonostoma**
atlanticum Norman, 1930
bathophilum (Vaillant *in* Filhol, 1884)
denudatum Rafinesque, 1810
elongatum Günther, 1878
- Margrethia**
obusirostra Jespersen & Tåning, 1919
- Family Sternopychidae**
- Argyripnus**
atlanticus Maul, 1952
- Maurolicus**
amethystinopunctatus Cocco, 1838
mulleri (Gmelin, 1789)
- Valenciennellus**
tripunctulatus (Esmark, 1871)
- Argyropelecus**
aculeatus Valenciennes *in* Cuvier & Valenciennes, 1850
affinis Garman, 1899
gigas Norman, 1930
hemigymnus Cocco, 1829
olfersi (Cuvier, 1829)
- Polyipnus**
polli Schultz, 1961
- Sternopyx**
diaphana Hermann, 1781
pseudoscura Baird, 1971
- Family Phosichthyidae**
- Ichthyococcus**
ovatus (Cocco, 1838)
- Pollichthys**
mauli (Poll, 1953)

- Polymetme*
corythaeola (Alcock, 1898)
thaecocyla Parin & Borodulina, 1990
- Vinciguerria*
attenuata (Cocco, 1838)
nimbaria (Jordan & Williams *in* Jordan & Starks, 1895)
poweriae (Cocco, 1838)
- Family Stomiidae**
- Chauliodus*
danae Regan & Trewavas, 1929
sloani Bloch & Schneider, 1801
- Stomias*
boa (Risso, 1810)
brevibarbus Ege, 1918
longibarbus (Brauer, 1902)
- Astronesthes*
atlanticus Parin & Borodulina, 1996
cyclophotus Regan & Trewavas, 1929
geminifer Goode & Bean, 1896
indicus Brauer, 1902
leucopogon Regan & Trewavas, 1929
macropogon Goodyear & Gibbs, 1970
micropogon Goodyear & Gibbs, 1970
neopogon Regan & Trewavas, 1929
niger Richardson, 1845
- Borostomias*
antarcticus (Lönnberg, 1905)
elucens (Brauer, 1906)
mononema (Regan & Trewavas, 1929)
- Neonesthes*
capensis (Gilchrist & Von Bonde, 1924)
- Rhadinesthes*
decimus (Zugmayer, 1911)
- Bathophilus*
brevis Regan & Trewavas, 1930
digitatus (Welsh, 1923)
nigerrimus Giglioli, 1882
pawnee Parr, 1927
vaillanti (Zugmayer, 1911)
- Chirostomias*
pliopterus Regan & Trewavas, 1930
- Echiostoma*
barbatum Lowe, 1843
- Eustomias*
braueri Zugmayer, 1911
filifer (Gilchrist, 1906)
furcifer Regan & Trewavas, 1930
longibarba Parr, 1927
macronema Regan & Trewavas, 1930
macrurus Regan & Trewavas, 1930
obscurus Vaillant *in* Filhol, 1884
schmidti Regan & Trewavas, 1930
simplex Regan & Trewavas, 1930
tetranema Zugmayer, 1913
- Flagellostomias*
boureei (Zugmayer, 1913)
- Grammatostomias*
circularis Morrow, 1959
flagellibarba Holt & Bryne, 1910
- Leptostomias*
gladiator (Zugmayer, 1911)
haplocaulus Regan & Trewavas, 1930
longibarba Regan & Trewavas, 1930
- Melanostomias*
bartonbeani Parr, 1927
biseriatus Regan & Trewavas, 1930
macrophotus Regan & Trewavas, 1930
melanopogon Regan & Trewavas, 1930
melanops Brauer, 1902
tentaculatus (Regan & Trewavas, 1930)
valdiviae Brauer, 1902
- Pachystomias*
microdon (Günther, 1878)
- Photonectes*
- braueri* (Zugmayer, 1913)
dinema Regan & Trewavas, 1930
margarita (Goode & Bean, 1896)
parvimanus Regan & Trewavas, 1930
- Trigonolampa*
miriceps Regan & Trewavas, 1930
- Aristostomias*
grimaldii Zugmayer, 1913
lunifer Regan & Trewavas, 1930
tittmanni Welsh, 1923
- Malacosteus*
niger Ayres, 1848
- Photostomias*
guernei Collett, 1889
Idiacanthus
fasciola Peters, 1877
- Order Ateleopodiformes**
- Family Atelcopidae**
- Guentherus*
altivelis Osório, 1917
- Ijimaia*
loppei Roule, 1922
- Order Aulopiformes**
- Family Aulopidae**
- Aulopus*
filamentosus (Bloch, 1792)
- Family Chlorophthalmidae**
- Chlorophthalmus*
agassizi Bonaparte, 1840
- Family Ipnopidae**
- Bathymicrops*
regis Hjort & Koefoed, 1912
- Bathypterois*
dubius Vaillant, 1888
grallator (Goode & Bean, 1886)
longipes Günther, 1878
phenax Parr, 1928
- Bathytyphlops*
sewelli (Norman, 1939)
- Family Scopelarchidae**
- Benthalbella*
infans Zugmayer, 1911
- Rosenblattichthys*
hubbsi Johnson, 1974
- Scopelarchus*
analis (Brauer, 1902)
michaelsarsi Koefoed, 1955
- Family Notosudidae**
- Ahliasaurus*
berryi Bertelsen, Kreft & Marshall, 1976
- Scopelosaurus*
argenteus (Maul, 1954)
lepidus (Kreft & Maul, 1955)
smithii Bean, 1925
- Family Synodontidae**
- Synodus*
saurus (Linnaeus, 1758)
synodus (Linnaeus, 1758)
- Saurida*
undosquamis (Richardson, 1848)
- Bathysaurus*
ferox Günther, 1878
mollis Günther, 1878
- Family Paralepididae**
- Arctozenus*
risso (Bonaparte, 1840)
- Lestidiops*
affinis (Ege, 1930)
jayakari (Boulenger, 1889)
sphyrenoides (Risso, 1820)
- Macroparalepis*
affinis Ege, 1933

- brevis* Ege, 1933
nigra (Maul, 1965)
Magnisudis
atlantica (Krøyer, 1868)
Paralepis
brevirostris (Parr, 1928)
coregonoides Risso, 1820
Sudis
hyalina Rafinesque, 1810
Uncisudis
longirostra Maul, 1956
quadrimaculata (Post, 1969)
- Family Evermannellidae**
Coccarella
atlantica (Parr, 1928)
Evermannella
balbo (Risso, 1820)
indica Brauer, 1906
Odontostomops
normalops (Parr, 1928)
- Family Omosudidae**
Omosudis
lowei Günther, 1887
- Family Alepisauridae**
Alepisaurus
brevirostris Gibbs, 1960
ferox Lowe, 1833
- Family Anophteridae**
Anophterus
pharao Zugmayer, 1911
- Order Myctophiformes**
- Family Neoscopelidae**
Neoscopelus
macrolepidotus Johnson, 1863
microchir Matsubara, 1943
Scopelengys
tristis Alcock, 1890
- Family Myctophidae**
Benthosema
glaciale (Reinhardt, 1838)
suborbitalis (Gilbert, 1913)
Bolinichthys
indicus (Nafpaktitis & Nafpaktitis, 1969)
supralateralis (Parr, 1928)
Centrobranchus
nigroocellatus (Günther, 1873)
Ceratoscopelus
maderensis (Lowe, 1839)
warmingii (Lütken, 1892)
Diaphus
adenomus Gilbert, 1905
bertelseni Nafpaktitis, 1966
brachycephalus Tåning, 1928
dumerilii (Bleeker, 1856)
effulgens (Goode & Bean, 1896)
holti Tåning, 1918
lucidus (Goode & Bean, 1896)
metopoclampus (Cocco, 1829)
mollis Tåning, 1928
rafinesquii (Cocco, 1838)
subtilis Nafpaktitis, 1968
termophilus Tåning, 1928
Diogenichthys
atlanticus (Tåning, 1928)
Electrona
risso (Cocco, 1829)
Gonichthys
cocco (Cocco, 1829)
Hygophum
benoiti (Cocco, 1838)
hygomii (Lütken, 1892)
reinhardti (Lütken, 1892)
taanungi Becker, 1965
- Lampadena*
anomala Parr, 1928
chavesi Collett, 1905
speculigera Goode & Bean, 1896
urophaos Paxton, 1963
Lampanyctus
alatus Goode & Bean, 1896
Nannobrachium
atrum (Tåning, 1928)
Lampanyctus
crocodilus (Risso, 1810)
Nannobrachium
cuprarium Tåning, 1928
Lampanyctus
festivus Tåning, 1928
intricarius Tåning, 1928
Nannobrachium
lineatum Tåning, 1928
Lampanyctus
macdonaldi (Goode & Bean, 1896)
photonus Parr, 1928
pusillus (Johnson, 1890)
Lepidophanes
gaussi (Brauer, 1906)
guentheri (Goode & Bean, 1896)
Lobianchia
dofleini (Zugmayer, 1911)
gemellarii (Cocco, 1838)
Loweina
interrupta (Tåning, 1928)
rara (Lütken, 1892)
Myctophum
nitidulum Garman, 1899
punctatum Rafinesque, 1810
selenops Tåning, 1928
Notolychnus
valdiviae (Brauer, 1904)
Notoscopelus (*Notoscopelus*)
caudispinosus (Johnson, 1863)
elongatus (Costa, 1844)
kroeyeri (Malm, 1861)
resplendens (Richardson, 1845)
Notoscopelus (*Pareiophus*)
bolini Nafpaktitis, 1975
Protomyctophum (*Hierops*)
arcticum (Lütken, 1892)
Symbolophorus
veranyi (Moreau, 1888)
Taaningichthys
bathophilus (Tåning, 1928)
minimus (Tåning, 1928)
paurollynchus Davy, 1972
- Order Lampriformes**
- Family Lampridae**
Lampris
guttatus (Brünnich, 1788)
- Family Lophotidae**
Lophotus
lacepede Giorna, 1809
- Family Radiicephalidae**
Radiicephalus
elongatus Osório, 1917
- Family Trachipteridae**
Trachipterus
arcticus (Brünnich, 1788)
trachypterus (Gmelin, 1789)
- Zu*
cristatus (Bonelli, 1819)
- Family Regalecidae**
Regalecus
glesne Ascanius, 1772
- Family Stylephoridae**
Stylephorus

- chordatus* Shaw, 1791
- Order Polymixiiformes**
- Family Polymixiidae**
- Polymixia nobilis* Lowe, 1838
- Order Gadiformes**
- Family Bregmacerotidae**
- Bregmaceros atlanticus* Goode & Bean, 1886
- Family Macrouridae**
- Bathygadus favosus* Goode & Bean, 1886
melanobranchus Vaillant, 1888
- Gadomus arcuatus* (Goode & Bean, 1886)
dispar (Vaillant, 1888)
longifilis (Goode & Bean, 1885)
- Trachyrincus murrayi* Günther, 1887
scabrus (Rafinesque, 1810)
- Caelorinchus caelorhincus* (Risso, 1810)
caudani (Koehler, 1896)
labiatus (Koehler, 1896)
- Cetonus*
globiceps (Vaillant in Filhol, 1884)
- Coryphaenoides* (*Chalinura*)
brevibarbis (Goode & Bean, 1896)
leptolepis Günther, 1877
mediterraneus (Giglioli, 1893)
profundiculus (Nybelin, 1957)
- Coryphaenoides* (*Coryphaenoides*)
guentheri (Vaillant, 1888)
rudis Günther, 1878
rupestris Gunnerus, 1765
thelestomus Maul, 1951
zaniophorus (Vaillant, 1888)
- Coryphaenoides* (*Lionurus*)
carapinus Goode & Bean, 1883
- Coryphaenoides* (*Nematorurus*)
armatus (Hector, 1875)
- Echinomacrurus mollis* Roule, 1916
- Hymenocephalus* (*Hymenocephalus*)
italicus Giglioli in Giglioli & Issel, 1884
- Hymenocephalus* (*Hymenogadus*)
gracilis Gilbert & Hubbs, 1920
- Macrourus berglax* Lacepède, 1801
- Malacocephalus* (*Malacocephalus*)
laevis (Lowe, 1843)
- Nezumia aequalis* (Günther, 1878)
longebarbata (Roule & Angel, 1933)
sclerorhynchus (Valenciennes, 1838)
- Odontomacrurus murrayi* Norman, 1939
- Pseudonezumia flagellicauda* (Koefoed, 1927)
- Sphagomacrurus grenadae* (Parr, 1946)
hirundo (Collett, 1896)
- Trachonurus villosum* (Günther, 1877)
- Family Moridae**
- Antimora rostrata* (Günther, 1878)
- Eretmophorus kleinenberghi* Giglioli, 1889
- Gadella maraldi* (Risso, 1810)
- Halargyreus johnsonii* Günther, 1862
- Laemonema latifrons* Holt & Byrne, 1908
robustum Johnson, 1862
yarrellii (Lowe, 1838)
- Lepidion**
- eques* (Günther, 1887)
guentheri (Giglioli, 1880)
lepidion (Risso, 1810)
schmidti Svetovidov, 1936
- Mora moro* (Risso, 1810)
- Physiculus dalwigki* Kaup, 1858
- Rhynchogadus hepaticus* (Facciolà, 1884)
- Svetovidovia lucullus* (Jensen in Koefod, 1953)
- Family Melanonidae**
- Melanonus zugmayeri* Norman, 1930
- Family Gadidae**
- Arctogadus glacialis* (Peters, 1872)
- Boreogadus saida* (Lepechin, 1774)
- Eleginus navaga* (Pallas, 1811)
- Gadiculus argenteus* Guichenot, 1850
- Gadus morhua* Linnaeus, 1758
- Melanogrammus aeglefinus* (Linnaeus, 1758)
- Merlangius merlangus* (Linnaeus, 1758)
- Micromesistius poutassou* (Risso, 1827)
- Pollachius pollachius* (Linnaeus, 1758)
virens (Linnaeus, 1758)
- Raniceps raninus* (Linnaeus, 1758)
- Theragra finnmarchica* Koefoed, 1956
- Trisopterus esmarkii* (Nilsson, 1855)
luscus (Linnaeus, 1758)
minutus (Linnaeus, 1758)
- Family Lotidae**
- Brosme brosme* (Ascanius, 1772)
- Ciliata**
- mustela* (Linnaeus, 1758)
septentrionalis (Collett, 1875)
- Enchelyopus cimbrius* (Linnaeus, 1766)
- Gaidropsarus argentatus* (Reinhardt, 1838)
biscayensis (Collett, 1890)
ensis (Reinhardt, 1838)
granti (Regan, 1903)
guttatus (Collett, 1890)
macrophthalmus (Günther, 1867)
mediterraneus (Linnaeus, 1758)
vulgaris (Cloquet, 1824)
- Molva dypterygia* (Pennant, 1784)
macrophthalmus (Rafinesque, 1810)
molva (Linnaeus, 1758)
- Family Phycidae**
- Phycis blennoides* (Brünnich, 1768)
phycis (Linnaeus, 1766)
- Irophycis*

- chuss* (Walbaum, 1792)
tenuis (Mitchill, 1814)
- Family Merlucciidae**
Macruronus (*Cynogadus*)
brachycolus (Holt & Byrne, 1906)
Merluccius
merluccius (Linnaeus, 1758)
senegalensis Cadenat, 1950
- Order Ophidiiformes**
- Family Ophidiidae**
Brotulotaenia
brevicauda Cohen, 1974
crassa Parr, 1934
nigra Parr, 1933
- Ophidion*
barbatum Linnaeus, 1758
rochei Müller, 1845
- Parophidion*
vassali (Risso, 1810)
- Barathrites*
iris Zugmayer, 1911
- Bassozetus*
taenia (Günther, 1887)
- Bathyonus*
laticeps (Günther, 1878)
- Benthocometes*
robustus (Goode & Bean, 1886)
- Holcomycteronus*
squamosus (Roule, 1916)
- Lamprogrammus*
niger Alcock, 1891
shcherbachevi Cohen & Rohr, 1993
- Monomitopus*
metriostoma (Vaillant, 1888)
- Spectrunculus*
grandis (Günther, 1877)
- Family Carapidae**
Carapus
acus (Brünnich, 1768)
- Echiodon*
dentatus (Cuvier, 1829)
drummondii Thompson, 1837
- Family Bythitidae**
Bellottia
apoda Giglioli, 1883
- Bythites*
islandicus Nielsen & Cohen, 1973
- Cataetyx*
alleni (Byrne, 1906)
laticeps Koefoed, 1927
- Grammonus*
ater (Risso, 1810)
- Thalassobathia*
pelagica Cohen, 1963
- Melodichthys*
hadrocephalus Nielsen & Cohen, 1986
- Family Aphyonidae**
Aphyonus
gelatinosus Günther, 1878
- Barathronus*
multidens Nielsen, 1984
parfauti (Vaillant, 1888)
- Meteoria*
erythrops Nielsen, 1969
- Nybelinella*
erikssoni (Nybelin, 1957)
- Sciadonus*
cryptophthalmus (Zugmayer, 1911)
galathaea (Nielsen, 1969)
pedicularis Garman, 1899
- Family Parabrotulidae**
Leucobrotula
adipata Koefoed, 1952
- Parabrotula*
plagiophthalma Zugmayer, 1911
- Order Batrachoidiformes**
- Family Batrachoididae**
Halobatrachus
didactylus (Bloch & Schneider, 1801)
- Order Lophiiformes**
- Family Lophiidae**
Lophius
budegassa Spinola, 1807
piscatorius Linnaeus, 1758
- Family Melanocetidae**
Melanocetus
johsonii Günther, 1864
- Family Antennariidae**
Antennarius
nummifer (Cuvier, 1817)
radiosus Garman, 1896
senegalensis Cadenat, 1959
- Histrio*
histrio (Linnaeus, 1758)
- Family Chaunacidae**
Chaunax
pictus Lowe, 1846
suttkusi Caruso, 1989
- Family Ogcocephalidae**
Dibranchus
atlanticus Peters, 1876
- Family Caulophrynidiae**
Caulophryne
jordani Goode & Bean, 1896
polynema Regan, 1930
- Family Melanocetidae**
Melanocetus
murrayi Günther, 1887
- Family Himantolophidae**
Himantolophus
albinares Maul, 1961
compressus (Osório, 1912)
groenlandicus Reinhardt, 1837
mauli Bertelsen & Krefft, 1988
- Family Diceratiidae**
Diceratias
wedli (Pietschmann, 1926)
- Family Oneirodidae**
Chaenophryne
draco Beebe, 1932
longiceps Regan, 1925
- Ctenochirichthys*
longimanus Regan & Trewavas, 1932
- Dolopichthys*
allector Garman, 1899
danae Regan, 1926
jubatus Regan & Trewavas, 1932
longicornis Parr, 1927
- Leptacanthichthys*
gracilispinis (Regan, 1925)
- Lophodolos*
acanthognathus Regan, 1925
- Microlophichthys*
microlophus (Regan, 1925)
- Oneirodes*
anisacanthus (Regan, 1925)
carlsbergi (Regan & Trewavas, 1932)
clarkei Swinney & Pietsch, 1988
eschrichtii Lütken, 1871
macronema (Regan & Trewavas, 1932)
macrosteus Pietsch, 1974
myrionemus Pietsch, 1974
- Pentherichthys*
venustus (Regan & Trewavas, 1932)

- Phyllorhinichthys**
micractis Pietsch, 1969
- Family Thaumatichthyidae**
- Lasiognathus*
bbeebei Regan & Trewavas, 1932
saccostoma Regan, 1925
- Family Centrophrynidæ**
- Centrophryne*
spinulosa Regan & Trewavas, 1932
- Family Ceratiidae**
- Ceratias*
holboelli Krøyer, 1845
- Cryptopsaras*
couesii Gill, 1883
- Family Gigantactinidae**
- Gigantactis*
vanhoefeni Brauer, 1902
- Family Linophrynidæ**
- Haplophryne*
mollis (Brauer, 1902)
- Linophryne*
brevibarbata Beebe, 1932
coronata Part, 1927
lucifer Collett, 1886
maderensis Maul, 1961
polypogon Regan, 1925
sexfiliis Bertelsen, 1973
- Order Gobiesociformes**
- Family Gobiesocidae**
- Apletodon*
bacescui (Murgoci, 1940)
dentatus (Facciola, 1887)
incognitus Hofrichter & Patzner, 1997
pellegrini (Chabanaud, 1925)
- Diplecogaster*
bimaculata (Bonnaterre, 1788)
euxinica Murgoci, 1964
pectoralis Briggs, 1955
- Gouania*
willdenowi (Risso, 1810)
- Lepadogaster*
candollei Risso, 1810
lepadogaster (Bonnaterre, 1788)
purpurea (Bonnaterre, 1788)
zebrina Lowe, 1839
- Opeatogenys*
gracilis (Canestrini, 1864)
- Order Atheriniformes**
- Family Atherinidae**
- Atherina* (*Atherina*)
hepsetus Linnaeus, 1758
- Atherina* (*Hepsetia*)
boyeri Risso, 1810
presbyter Cuvier, 1829
- Atherinomorus*
lacunosus (Forster in Bloch & Schneider, 1801)
- Order Cyprinodontiformes**
- Family Cyprinodontidae**
- Aphanius*
dispar (Rüppell, 1829)
fasciatus (Valenciennes, 1821)
iberus (Valenciennes, 1846)
- Order Beloniformes**
- Family Scomberesocidae**
- Nanichthys*
simulans Hubbs & Wisner, 1980
- Scomberesox*
saurus (Walbaum, 1792)
- Family Belonidae**
- Belone*
belone (Linnaeus, 1761)
svetovidovi Collette & Parin, 1970
- Platybelone*
argalus (Lesueur, 1821)
- Tylosurus*
acus (Lacepède, 1803)
choram (Rüppell, 1837)
imperialis (Rafinesque, 1810)
- Family Hemiramphidae**
- Hemiramphus*
far (Forsskål, 1775)
- Hyporhamphus*
picarti (Valenciennes in Cuvier & Valenciennes, 1847)
- Family Exocoetidae**
- Cheilopogon*
exsiliens (Linnaeus, 1771)
heterurus (Rafinesque, 1810)
pinnatibarbus (Bennett, 1831)
- Exocoetus*
obtusirostris Günther, 1866
volitans Linnaeus, 1758
- Hirundichthys*
rondeletii (Valenciennes in Cuvier & Valenciennes, 1847)
speculiger (Valenciennes in Cuvier & Valenciennes, 1847)
- Parexocoetus*
mento (Valenciennes in Cuvier & Valenciennes, 1847)
- Order Stephanoberyciformes**
- Family Stephanoberycidae**
- Acanthochaenus*
luetkenii Gill, 1844
- Family Melamphaidæ**
- Melamphaes*
longivelis Parr, 1933
microps (Günther, 1878)
simus Ebeling, 1962
suborbitalis (Gill, 1883)
typhlops (Lowe, 1843)
- Poromitra*
capito Goode & Bean, 1883
crassiceps (Günther, 1878)
megalops (Lütken, 1877)
- Scopeloberyx*
opisthoterus (Parr, 1933)
robustus (Günther, 1887)
rubriventer (Koefoed, 1953)
- Scopelogadus*
beanii (Günther, 1887)
mizolepis (Günther, 1878)
- Order Beryciformes**
- Family Trachichthyidae**
- Gephyroberyx*
darwinii (Johnson, 1866)
- Hoplostethus*
atlanticus Collett, 1889
cadenati Quéro, 1974
mediterraneus Cuvier in Cuvier & Valenciennes, 1829
- Family Diretmidae**
- Diretmichthys*
parini (Post & Quéro, 1981)
- Diretmoides*
pauciradiatus (Woods in Woods & Sonoda, 1973)
- Diretmus*
argenteus Johnson, 1864
- Family Anoplogasteridae**
- Anoplogaster*
cornuta (Valenciennes in Cuvier & Valenciennes, 1833)
- Family Berycidae**
- Beryx*

- decadactylus* Cuvier in Cuvier & Valenciennes, 1829
splendens Lowe, 1834
- Family Holocentridae**
Sargocentron
hastatum (Cuvier in Cuvier & Valenciennes, 1829)
rubrum (Forsskål, 1775)
- Order Cetomimiformes**
- Family Mirapinnidae**
Eutaeniophorus
festivus (Bertelsen & Marshall, 1956)
Mirapinna
esau Bertelsen & Marshall, 1956
Parataeniophorus
gulosus Bertelsen & Marshall, 1956
- Family Megalomycteridae**
Ataxolepis
apus Myers & Freihoffer, 1966
- Family Rondeletiidae**
Rondeletia
loricata Abe & Hotta, 1963
- Family Cetomimidae**
Cetomimus
hempeli Maul, 1969
Cetostoma
regani Zugmayer, 1914
Ditropichthys
storeri (Goode & Bean, 1895)
- Order Zeiformes**
- Family Macrurocyttidae**
Zenion
hololepis (Goode & Bean, 1896)
- Family Zeidae**
Cytopsis
rosea (Lowe, 1843)
Zenopsis
conchifer (Lowe, 1852)
Zeus
faber Linnaeus, 1758
- Family Oreosomatidae**
Neocytthus
helgae (Holt & Byrne, 1908)
Allocyttus
verrucosus (Gilchrist, 1906)
Pseudocytthus
maculatus Gilchrist, 1906
- Family Grammicolepididae**
Grammicolepis
brachiusculus Poey, 1873
- Family Caproidae**
Antigonia
capros Lowe, 1843
Capros
aper (Linnaeus, 1758)
- Order Gasterosteiformes**
- Family Gasterosteidae**
Gasterosteus
aculeatus Linnaeus, 1758
Pungitius
platygaster (Kessler, 1859)
pungitius (Linnaeus, 1758)
Spinachia
spinachia (Linnaeus, 1758)
- Order Syngnathiformes**
- Family Centriscidae**
Macrorhamphosus
scolopax (Linnaeus, 1758)
- Order Syngnathiiformes**
- Family Syngnathidae**
Entelurus
- aequoreus* (Linnaeus, 1758)
Minyichthys
sentus Dawson, 1982
Nerophis
lumbriciformis (Jenyns, 1835)
maculatus Rafinesque, 1810
ophidion (Linnaeus, 1758)
Syngnathus
abaster Risso, 1827
acus Linnaeus, 1758
phlegon Risso, 1827
rostellatus Nilsson, 1855
schmidtii Popov, 1927
taenianotus Canestrini, 1871
tenuirostris Rathke, 1837
typhle Linnaeus, 1758
variegatus Pallas, 1814
Hippocampus
hippocampus (Linnaeus, 1758)
ramulosus Leach in Shaw & Nodder, 1814
- Order Scorpaeniformes**
- Family Sebastidae**
Helicolenus
dactylopterus (Delaroche, 1809)
Sebastes
fasciatus Storer, 1856
marinus (Linnaeus, 1758)
mentella (Travin, 1951)
viviparus Krøyer, 1845
Trachyscorpia
cristulata (Goode & Bean, 1896)
echinata (Koehler, 1896)
- Family Setarchidae**
Setarches
guentheri Johnson, 1862
- Family Scorpaenidae**
Pontinus
kuhlii (Bowdich, 1825)
Scorpaena
azorica Eschmeyer, 1969
elongata Cadenat, 1943
laevis Troschel, 1866
lopezi Cadenat, 1943
madurensis Valenciennes, 1833
notata Rafinesque, 1810
porcus Linnaeus, 1758
scrofa Linnaeus, 1758
Scorpaenodes
arenai Torchio, 1962
- Family Dactylopteridae**
Dactylopterus
volitans (Linnaeus, 1758)
- Family Scorpaenidae**
Sebastapistes
mucialis (Günther, 1874)
- Family Triglidae**
Chelidonichthys
cuculus (Linnaeus, 1758)
gurnardus (Linnaeus, 1758)
lastoviza (Bonnaterre, 1788)
lucerna (Linnaeus, 1758)
obscurus (Walbaum, 1792)
Lepidotrigla
cavillone (Lacepède, 1801)
dieuzeidei Blanc & Hureau ex Andouin, 1973
Trigla
lyra Linnaeus, 1758
- Family Peristediidae**
Peristedion
cataphractum (Linnaeus, 1758)
- Family Platyccephalidae**
Platycephalus

- indicus* (Linnaeus, 1758)
- Family Cottidae**
- Artediellus*
 - atlanticus* Jordan & Evermann, 1898
 - scaber* Knipowitsch, 1907
 - uncinatus* (Reinhardt, 1835)
 - Cottus*
 - gobio* Linnaeus, 1758
 - poecilopus* Heckel, 1837
 - Gymnophanthus*
 - tricuspidis* (Reinhardt, 1830)
 - Icelus*
 - bicornis* (Reinhardt, 1840)
 - spatula* Gilbert & Burke, 1912
 - Micrenophrys*
 - lilljeborgii* (Collett, 1875)
 - Myoxocephalus*
 - scorpioides* (Fabricius, 1780)
 - scorpius* (Linnaeus, 1758)
 - Taurulus*
 - bubalis* (Euphrasen, 1786)
 - Triglops*
 - murrayi* Günther, 1888
 - nybelini* Jensen, 1944
 - pingelii* Reinhardt, 1837
 - Triglopsis*
 - quadricornis* (Linnaeus, 1758)
- Family Psychrolutidae**
- Cottunculus*
 - microps* Collett, 1875
 - sadko* Essipov, 1937
 - thomsonii* (Günther, 1882)
 - Ebinaria*
 - costaeccanariae* (Cervigon, 1961)
 - Psychrolutes*
 - subspinosus* (Jensen, 1902)
- Family Agonidae**
- Agonus*
 - cataphractus* (Linnaeus, 1758)
 - Leptagonus*
 - decagonus* (Bloch & Schneider, 1801)
 - Ulcina*
 - olrika* (Lütken, 1876)
- Family Cyclopteridae**
- Cyclopteropsis*
 - meclipini* (Fowler, 1914)
 - Cyclopterus*
 - lumpus* Linnaeus, 1758
 - Eumicrotremus*
 - derjugini* Popov, 1926
 - eggvinii* Koefoed, 1956
 - spinulosus* (Müller ex Fabricius, 1776)
 - Careproctus*
 - aciculipunctatus* Andriashev & Chernova, 1997
 - merretti* Andriashev & Chernova, 1988
 - micropus* (Günther, 1887)
 - reinhardti* (Krøyer, 1862)
 - Liparis*
 - fabricii* Krøyer, 1847
 - liparis* (Linnaeus, 1766)
 - montagui* (Donovan, 1804)
 - Paraliparis*
 - abyssorum* Andriashev & Chernova, 1997
 - bathybius* (Collett, 1879)
 - bipolaris* Andriashev, 1997
 - challengeri* Andriashev, 1993
 - edwardsi* (Vaillant, 1888)
 - hystrix* Merrett, 1983
 - leptoichirus* (Tortonese, 1959)
 - membranaceus* Günther, 1887
 - murieli* Matallanas, 1984
 - Pseudnos*
 - christinae* Andriashev, 1992
 - Rhodichthys*
- regina* Collett, 1879
- Order Perciformes**
- Family Percichthyidae**
- Howella*
 - brodiei* Ogilby, 1899
 - sherborni* (Norman, 1930)
- Family Moronidae**
- Dicentrarchus*
 - labrax* (Linnaeus, 1758)
 - punctatus* (Bloch, 1792)
- Family Polyprionidae**
- Polyprion*
 - americanus* (Bloch & Schneider, 1801)
- Family Serranidae**
- Serranus*
 - atrauda* Günther, 1874
 - cabrilla* (Linnaeus, 1758)
 - hepatus* (Linnaeus, 1758)
 - scriba* (Linnaeus, 1758)
 - Anthias*
 - anthias* (Linnaeus, 1758)
 - Epinephelus*
 - aeneus* (Geoffroy Saint-Hilaire, 1817)
 - caninus* (Valenciennes, 1843)
 - costae* (Steindachner, 1878)
 - haifensis* Ben-Tuvia, 1953
 - marginatus* (Lowe, 1834)
 - nigeritus* (Holbrook, 1855)
 - Mycteroperca*
 - fusca* (Lowe, 1838)
 - rubra* (Bloch, 1793)
- Family Callanthiidae**
- Callanthias*
 - ruber* (Rafinesque, 1810)
- Family Terapontidae**
- Pelates*
 - quadrilineatus* (Bloch, 1790)
 - Terapon*
 - puta* (Cuvier, 1829)
- Family Priacanthidae**
- Cookeolus*
 - japonicus* (Cuvier in Cuvier & Valenciennes, 1829)
 - Heteropriacanthus*
 - cruentatus* (Lacepedé, 1801)
 - Priacanthus*
 - arenatus* Cuvier in Cuvier & Valenciennes, 1829
- Family Apogonidae**
- Apogon* (*Apogon*)
 - imberbis* (Linnaeus, 1758)
 - Apogon* (*Nectamia*)
 - taeniatus* Cuvier ex Ehrenberg) in Cuvier & Valenciennes, 1828
- Family Epigonidae**
- Epigonus*
 - constanciae* (Giglioli, 1880)
 - denticulatus* Dieuzeide, 1950
 - telescopus* (Risso, 1810)
 - Microichthys*
 - coccoi* Rüppell, 1852
 - sanzoi* Spartà, 1950
- Family Pomatomidae**
- Pomatomus*
 - saltator* (Linnaeus, 1766)
- Family Rachycentridae**
- Rachycentron*
 - canadum* (Linnaeus, 1766)
- Family Echeneidae**
- Echeneis*
 - naucrates* Linnaeus, 1758
 - Phtheirichthys*
 - lineatus* (Menzies, 1791)

- Remora**
australis (Bennett, 1840)
brachyptera (Lowe, 1839)
osteochir (Cuvier, 1829)
remora (Linnaeus, 1758)
- Remorina**
albescens (Temminck & Schlegel, 1845)
- Family Carangidae**
- Alectis*
alexandrinus (Geoffroy Saint-Hilaire, 1817)
- Alepes*
djedaba (Forsskål, 1775)
- Campogramma*
glaycos (Lacepède, 1801)
- Caranx*
crysos (Mitchill, 1815)
hippos (Linnaeus, 1766)
latus Agassiz in Spix & Agassiz, 1831
lugubris Poey, 1860
rhonchus Geoffroy Saint-Hilaire, 1817
- Decapterus*
macarellus (Cuvier in Cuvier & Valenciennes, 1833)
punctatus (Cuvier, 1829)
- Elagatis*
bipinnulata (Quoy & Gaimard, 1825)
- Lichia*
amia (Linnaeus, 1758)
- Naucrates*
ductor (Linnaeus, 1758)
- Pseudocaranx*
dentex (Bloch & Schneider, 1801)
- Selar*
boops (Cuvier in Cuvier & Valenciennes, 1833)
- Selene*
dorsalis (Gill, 1863)
- Seriola*
carpenteri Mather, 1971
dumerili (Risso, 1810)
fasciata (Bloch, 1793)
rivoliana Valenciennes in Cuvier & Valenciennes, 1833
- Trachinotus*
ovatus (Linnaeus, 1758)
- Trachurus*
mediterraneus (Steindachner, 1868)
picturatus (Bowdich, 1825)
trachurus (Linnaeus, 1758)
trecae Cadenat, 1950
- Family Coryphaenidae**
- Coryphaena*
equiselis Linnaeus, 1758
hippurus Linnaeus, 1758
- Family Leiognathidae**
- Leiognathus*
kunzingeri (Steindachner, 1898)
- Family Bramidae**
- Brama*
brama (Bonnaterre, 1788)
- Pteraclis*
carolinus Valenciennes in Cuvier & Valenciennes, 1833
- Pterycombus*
brama Fries, 1837
- Taractes*
asper Lowe, 1843
rubescens (Jordan & Evermann, 1887)
- Taractichthys*
longipinnis (Lowe, 1843)
- Family Caristiidae**
- Caristius*
macropus (Bellotti, 1903)
maderensis Maul, 1949
opalescens (Zugmayer, 1911)
- Family Emmelichthyidae**
- Erythrocles*
monodi Poll & Cadenat, 1954
- Family Lobotidae**
- Lobotes*
surinamensis (Bloch, 1790)
- Family Haemulidae**
- Brachydeuterus*
auritus (Valenciennes in Cuvier & Valenciennes, 1832)
- Parapristipoma*
humile (Bowdich, 1825)
octolineatum (Valenciennes in Cuvier & Valenciennes, 1833)
- Plectrohinchus*
mediterraneus (Guichenot, 1850)
- Pomadasys*
incisus (Bowdich, 1825)
stridens (Forsskål, 1775)
- Family Sparidae**
- Boops*
boops (Linnaeus, 1758)
- Crenidens*
crenides (Forsskål, 1775)
- Dentex* (*Cheimerius*)
canariensis Steindachner, 1881
gibbosus (Rafinesque, 1810)
- Dentex* (*Dentex*)
dentex (Linnaeus, 1758)
- Dentex* (*Polysteganus*)
macrophthalmus (Bloch, 1791)
maroccanus Valenciennes in Cuvier & Valenciennes, 1830
- Diplodus*
annularis (Linnaeus, 1758)
bellottii (Steindachner, 1882)
cervinus (Lowe, 1838)
puntazzo (Cetti, 1777)
sargus (Linnaeus, 1758)
vulgaris (Geoffroy Saint-Hilaire, 1817)
- Lithognathus*
mormyrus (Linnaeus, 1758)
- Oblada*
melanura (Linnaeus, 1758)
- Pagellus*
acarne (Risso, 1827)
bellottii Steindachner, 1882
bogaraveo (Brünlich, 1768)
erythrinus (Linnaeus, 1758)
- Pagrus*
auriga Valenciennes, 1843
caeruleostictus (Valenciennes in Cuvier & Valenciennes, 1830)
pagrus (Linnaeus, 1758)
- Sarpa*
salpa (Linnaeus, 1758)
- Sparus*
auratus Linnaeus, 1758
- Spondylisoma*
canthus (Linnaeus, 1758)
- Family Centracanthidae**
- Centracanthus*
cirrus Rafinesque, 1810
- Spicara*
maena (Linnaeus, 1758)
melanurus (Valenciennes in Cuvier & Valenciennes, 1830)
smaris (Linnaeus, 1758)
- Family Sciaenidae**
- Argyrosomus*
regius (Asso, 1801)
- Pseudotolithus*
senegalensis (Valenciennes, 1833)
typus Bleeker, 1863

- Sciaena**
umbra Linnaeus, 1758
- Umbrina**
canariensis Valenciennes, 1843
cirrosa (Linnaeus, 1758)
ronchus Valenciennes, 1843
- Family Polynemidae**
Galeoides
decadactylus (Bloch, 1795)
- Family Mullidae**
Mullus
barbatus Linnaeus, 1758
surmuletus Linnaeus, 1758
- Pseudupeneus**
pragensis (Cuvier in Cuvier & Valenciennes, 1829)
- Upeneus**
asymmetricus Lachner, 1954
moluccensis (Bleeker, 1855)
- Family Kyphosidae**
Kyphosus
incisor (Cuvier in Cuvier & Valenciennes, 1831)
sextator (Linnaeus, 1766)
- Family Chaetodontidae**
Chaetodon
hoefti Steindachner, 1881
robustus Günther, 1860
- Prognathodes**
marcellae (Poll, 1950)
- Family Cepolidae**
Cepala
macrophthalma (Linnaeus, 1758)
- Family Mugilidae**
Chelon
labrosus (Risso, 1827)
- Liza*
aurata (Risso, 1810)
carinata (Valenciennes ex Ehrenberg) in Cuvier & Valenciennes, 1836
ramada (Risso, 1810)
saliens (Risso, 1810)
- Mugil*
capurrii (Perugia, 1892)
cephalus Linnaeus, 1758
- Oedalechilus*
lobeo (Cuvier, 1829)
- Family Pomacentridae**
Abudefduf
luridus (Cuvier in Cuvier & Valenciennes, 1830)
- Chromis*
chromis (Linnaeus, 1758)
limbata (Valenciennes in Cuvier & Valenciennes, 1833)
- Family Labridae**
Acantholabrus
palloni (Risso, 1810)
- Bodianus*
scrfa (Valenciennes in Cuvier & Valenciennes, 1839)
speciosus (Bowdich, 1825)
- Centrolabrus*
caeruleus Azevedo, 1999
exoletus (Linnaeus, 1758)
trutta (Lowe, 1834)
- Coris*
julis (Linnaeus, 1758)
- Ctenolabrus*
rupestris (Linnaeus, 1758)
- Labrus*
bergylta Ascanius, 1767
bimaculatus Linnaeus, 1758
merula Linnaeus, 1758
viridis Linnaeus, 1758
- Lappanella*
fasciata (Cocco, 1833)
- Sympodus (Crenilabrus)**
bailloni (Valenciennes in Cuvier & Valenciennes, 1839)
cinereus (Bonnaterre, 1788)
doderleini Jordan, 1890
mediterraneus (Linnaeus, 1758)
melanocercus (Risso, 1810)
melops (Linnaeus, 1758)
ocellatus Forsskål, 1775
roissali (Risso, 1810)
tinca (Linnaeus, 1758)
- Sympodus (Sympodus)**
rostratus (Bloch, 1791)
- Thalassoma**
pavo (Linnaeus, 1758)
- Xyrichtys**
novacula (Linnaeus, 1758)
- Family Scaridae**
Sparisoma (Euscarus)
cretense (Linnaeus, 1758)
- Family Zoarcidae**
Gymnelus
retrodorsalis Le Danois, 1913
viridis (Fabricius, 1780)
- Lycenchelys*
alba (Vaillant, 1888)
kolthoffi Jensen, 1904
muraena (Collett, 1878)
platyrhina (Jensen, 1902)
sarsi (Collett, 1871)
- Lycodes*
adolphi Nielsen & Fosså, 1993
esmarkii Collett, 1875
eudipleurostictus Jensen, 1902
frigidus Collett, 1879
jugoricus Knipowitsch, 1906
lavalei Vladkov & Tremblay, 1936
luetkenii Collett, 1880
microcephalus Jensen, 1902
pallidus Collett, 1879
polaris (Sabine, 1824)
reticulatus Reinhardt, 1835
rossi Malmgren, 1865
sagittarius McAllister, 1976
seminudus Reinhardt, 1837
squamiventer Jensen, 1904
terraenovae Collett, 1896
vahlii Reinhardt, 1831
- Lycodonus*
flagellicauda (Jensen, 1902)
- Melanostigma*
atlanticum Koefoed, 1952
- Pachycara*
bulbiceps (Garman, 1899)
crassiceps (Roule, 1916)
- Zoarces*
viviparus (Linnaeus, 1758)
- Family Stichaeidae**
Anisarchus
medius (Reinhardt, 1838)
- Chirolophis*
ascanii (Walbaum, 1792)
- Leptoclinus*
maculatus (Fries, 1837)
- Lumpenus*
fabricii Reinhardt, 1836
lumpenaeformis (Walbaum, 1792)
- Family Pholidae**
Pholis
gunnellus (Linnaeus, 1758)
- Family Anarhichadidae**
Anarhichas
denticulatus Krøyer, 1845
lupus Linnaeus, 1758

- minor* Olafsen, 1772
- Family Chiasmodontidae**
- Chiasmodon*
- bolangeri* Osório, 1909
 - niger* Johnson, 1864
- Kali*
- macrodon* (Norman, 1929)
 - macrura* (Parr, 1933)
- Pseudoscopelus*
- alipinnis* Parr, 1933
- Family Ammodytidae**
- Ammodytes*
- marinus* Raitt, 1934
 - tobianus* Linnaeus, 1758
- Gymnammodytes*
- cicerelus* (Rafinesque, 1810)
 - semisquamatus* (Jourdain, 1879)
- Hyperoplus*
- immaculatus* (Corbin, 1950)
 - lanceolatus* (Le Sauvage, 1824)
- Family Trachinidae**
- Trachinus*
- araneus* Cuvier ex Risso in Cuvier & Valenciennes, 1829
 - draco* Linnaeus, 1758
 - radiatus* Cuvier, 1829
 - vipera* Cuvier, 1829
- Family Uranoscopidae**
- Uranoscopus*
- scaber* Linnaeus, 1758
- Family Tripterygiidae**
- Tripterygion*
- delaisi* Cadenat & Blache, 1970
 - melanurus* Guichenot, 1850
 - tripteronotus* (Risso, 1810)
- Family Clinidae**
- Clinitrachus*
- argentatus* (Risso, 1810)
- Family Blenniidae**
- Aidablennius*
- sphyra* (Valenciennes in Cuvier & Valenciennes, 1836)
- Blennius*
- ocellaris* Linnaeus, 1758
- Coryphoblennius*
- galerita* (Linnaeus, 1758)
- Hyleurochilus*
- bananensis* (Poll, 1959)
- Lipophrys*
- adriaticus* (Steindachner & Kolombatovic, 1883)
 - canevae* (Vinciguerra, 1880)
 - dalmatinus* (Steindachner & Kolombatovic, 1883)
 - nigriceps* (Vinciguerra, 1883)
 - pholis* (Linnaeus, 1758)
- Ophioblennius*
- atlanticus* (Valenciennes in Cuvier & Valenciennes, 1836)
- Parablennius*
- gattorugine* (Linnaeus, 1758)
 - incognitus* (Bath, 1968)
 - parvicornis* (Valenciennes in Cuvier & Valenciennes, 1836)
 - pilicornis* (Cuvier, 1829)
 - ruxii* (Cocco, 1833)
 - ruber* (Valenciennes in Cuvier & Valenciennes, 1836)
 - sanguinolentus* (Pallas, 1814)
 - tentacularis* (Brünnich, 1768)
 - zonimiri* (Kolombatovic, 1892)
- Paralipophrys*
- trigloides* (Valenciennes in Cuvier & Valenciennes, 1836)
- Salaria*
- basilisca* (Valenciennes, 1836)
- pavo* (Risso, 1810)
- Scartella*
- cristata* (Linnaeus, 1758)
- Family Callionymidae**
- Callionymus*
- fasciatus* Valenciennes in Cuvier & Valenciennes, 1837
 - filamentosus* Valenciennes in Cuvier & Valenciennes, 1837
 - lyra* Linnaeus, 1758
 - maculatus* Rafinesque, 1810
 - pusillus* Delaroche, 1809
 - reticulatus* Valenciennes in Cuvier & Valenciennes, 1837
 - risso* Lesueur, 1814
- Protogrammus*
- sousai* (Maul, 1972)
- Synchiropus*
- phaeton* (Günther, 1861)
- Family Draconettidae**
- Centrodraco*
- acanthopoma* (Regan, 1904)
- Family Gobiidae**
- Aphia*
- minuta* (Risso, 1810)
- Benthophiloides*
- brauneri* Beling & Iljin, 1927
- Benthophilus*
- stellatus* (Sauvage, 1874)
- Buenia*
- affinis* Iljin, 1930
 - jeffreysii* (Günther, 1867)
- Caspiosoma*
- caspium* (Kessler, 1877)
- Chromogobius*
- quadrivittatus* (Steindachner, 1863)
 - zebratus* (Kolombatovic, 1891)
- Corycrogobius*
- liechtensteini* (Kolombatovic, 1891)
- Corygalops*
- ochetica* (Norman, 1927)
- Crystallogobius*
- linearis* (Düben, 1845)
- Deltentosteus*
- collonianus* (Risso, 1820)
 - quadrimaculatus* (Valenciennes in Cuvier & Valenciennes, 1837)
- Didogobius*
- bentuvii* Miller, 1966
- Gammogobius*
- steinizzi* Bath, 1971
- Gobius*
- ater* Bellotti, 1888
 - auratus* Risso, 1810
 - buccichi* Steindachner, 1870
 - cobitis* Pallas, 1814
 - couchi* Miller & El-Tawil, 1974
 - cruentatus* Gmelin, 1789
 - fallax* Sarato, 1889
 - gasteveni* Miller, 1974
 - geniporus* Valenciennes in Cuvier & Valenciennes, 1837
 - niger* Linnaeus, 1758
 - paganellus* Linnaeus, 1758
 - roulei* de Buen, 1928
 - strictus* Fage, 1907
 - vittatus* Vinciguerra, 1883
 - xanthocephalus* Heymer & Zander, 1992
- Gobiusculus*
- flavescens* (Fabricius, 1779)
- Knipowitschia*
- caucasica* (Berg ex Kavraisii, 1916)
 - longecaudata* (Kessler, 1877)
 - panizzae* (Verga, 1841)

- Lebetus***
guillei (Le Danois, 1913)
scorpioides (Collett, 1874)
- Lesueurigobius***
friesii (Malm, 1874)
heterofasciatus Maul, 1971
sanzi (de Buen, 1918)
suerii (Risso, 1810)
- Mauligobius***
maderensis (Valenciennes in Cuvier & Valenciennes, 1837)
- Mesogobius***
batrachocephalus (Pallas, 1814)
- Millerigobius***
macrocephalus (Kolombatovic, 1891)
- Neogobius***
cephalargoides Pinchuk, 1976
eurycephalus (Kessler, 1874)
fluvialis (Pallas, 1814)
gymnotrachelus (Kessler, 1857)
kessleri (Günther, 1861)
melanostomus (Pallas, 1814)
platyrostris (Pallas, 1814)
raian (Nordmann, 1840)
syrman (Nordmann, 1840)
- Odondebuenia***
balearica (Pellegrin & Fage, 1907)
- Oxyurichthys***
papuensis (Valenciennes, 1837)
- Pomatoschistus***
bathi Miller, 1982
canestrinii (Ninni, 1883)
knerii (Steindachner, 1861)
lozanoi (de Buen, 1923)
marmoratus (Risso, 1810)
micros (Krøyer, 1838)
minutus (Pallas ex Gronovius, 1770)
norvegicus (Collett, 1902)
pictus (Malm, 1865)
quagga (Heckel, 1837)
tortonesei Miller, 1969
- Proterorhinus***
marmoratus (Pallas, 1814)
- Pseudaphya***
ferreri (de Buen & Fage, 1908)
- Silhouettea***
aegyptia (Chabanaud, 1933)
- Speleogobius***
trigloides Zander & Jelinek, 1976
- Thorogobius***
ephippiatus (Lowe, 1839)
macrolepis (Kolombatovic, 1891)
- Vanneaugobius***
canariensis Van Tassell, Miller & Brito, 1988
pruvioti (Fage, 1907)
- Zebrus***
zebrus (Risso, 1827)
- Zosterisessor***
ophiocephalus (Pallas, 1814)
- Family Siganidae**
Siganus
luridus (Rüppell ex Ehrenberg, 1829)
rivulatus Forsskål, 1775
- Family Sillaginidae**
Sillago
sihama (Forsskål, 1775)
- Family Luvaridae**
Luvarus
imperialis Rafinesque, 1810
- Family Acanthuridae**
Acanthurus
monroviae Steindachner, 1876
- Family Scomrolabracidae**
Scomrolabrax
- heterolepis*** Roule, 1921
- Family Sphyraenidae**
Sphyraena
chrysotaenia Klunzinger, 1884
sphyraena (Linnaeus, 1758)
viridensis Cuvier in Cuvier & Valenciennes, 1829
- Family Gempylidae**
Diplospinus
multistriatus Maul, 1948
- Gempylus***
serpens Cuvier, 1829
- Lepidocybium***
flavobrunneum (Smith, 1843)
- Nealotus***
tripes Johnson, 1865
- Nesiarchus***
nasutus Johnson, 1862
- Promethichthys***
prometheus (Cuvier in Cuvier & Valenciennes, 1832)
- Ruvettus***
pretiosus Cocco, 1833
- Family Trichiuridae**
Aphanopus
carbo Lowe, 1839
- Benthodesmus***
elongatus (Clarke, 1879)
simonyi (Steindachner, 1891)
- Lepidopus***
caudatus (Euphrasen, 1788)
- Trichiurus***
lepturus Linnaeus, 1758
- Family Xiphidae**
Xiphias
gladius Linnaeus, 1758
- Family Istiophoridae**
Istiophorus
albicans (Latreille, 1804)
- Makaira***
nigricans Lacepède, 1802
- Tetrapturus***
albidus Poey, 1860
belone Rafinesque, 1810
georgii Lowe, 1841
pfluegeri Robins & de Silva, 1963
- Family Scombridae**
Acanthocybium
solandri (Cuvier in Cuvier & Valenciennes, 1832)
- Auxis***
rochei (Risso, 1810)
thazard (Lacepède, 1800)
- Euthynnus***
alleletteratus (Rafinesque, 1810)
- Katsuwonus***
pelamis (Linnaeus, 1758)
- Orcynopsis***
unicolor (Geoffroy Saint-Hilaire, 1817)
- Rastrelliger***
kanagurta (Cuvier, 1816)
- Sarda***
sarda (Bloch, 1793)
- Scomber***
japonicus Houttuyn, 1782
scombrus Linnaeus, 1758
- Scomberomorus***
commerson (Lacepède, 1800)
tritor (Cuvier in Cuvier & Valenciennes, 1832)
- Thunnus***
alalunga (Bonnaterre, 1788)
albacares (Bonnaterre, 1788)
obesus (Lowe, 1839)
thynnus (Linnaeus, 1758)
- Family Centrolophidae**
Centrolophus

- niger* (Gmelin, 1789)
Hyperoglyphe
perciformis (Mitchill, 1818)
Schedophilus
medusophagus Cocco, 1829
ovalis (Cuvier in Cuvier & Valenciennes, 1833)
- Family Nomeidae**
Cubiceps
gracilis (Lowe, 1843)
Nomeus
gronovii (Gmelin, 1789)
Psenes
maculatus Lütken, 1880
pellucidus Lütken, 1880
- Family Stromateidae**
Pampus
argenteus (Euphrasen, 1788)
Stromateus
fiatola Linnaeus, 1758
- Family Tetragonuridae**
Tetragonurus
atlanticus Lowe, 1839
cuvieri Risso, 1810
- Order Pleuronectiformes**
- Family Citharidae**
Citharus
linguatula (Linnaeus, 1758)
- Family Scophthalmidae**
Lepidorhombus
boscii (Risso, 1810)
whiffagonis (Walbaum, 1792)
Phrynorhombus
norvegicus (Günther, 1862)
regius (Bonnaterre, 1788)
Scophthalmus
maximus (Linnaeus, 1758)
rhombus (Linnaeus, 1758)
Zeugopterus
punctatus (Bloch, 1787)
- Family Bothidae**
Arnoglossus
imperialis (Rafinesque, 1810)
kessleri Schmidt, 1915
laterna (Walbaum, 1792)
rueppelii (Cocco, 1844)
thori Kyle, 1913
Bothus
podas (Delaroche, 1809)
- Family Pleuronectidae**
Glyptocephalus
cynoglossus (Linnaeus, 1758)
Hippoglossoides
platessoides (Fabricius, 1780)
Hippoglossus
hippoglossus (Linnaeus, 1758)
Limanda
limanda (Linnaeus, 1758)
Microstomus
kitt (Walbaum, 1792)
Platichthys
flesus (Linnaeus, 1758)
Pleuronectes
glacialis Pallas, 1776
platessa Linnaeus, 1758
Reinhardtius
hippoglossoides (Walbaum, 1792)
- Family Soleidae**
Bathysolea
profundicola (Vaillant, 1888)
Buglossidium
luteum (Risso, 1810)
Dicologlossa
- cuneata* (Moreau ex de la Pylaie, 1881)
Microchirus (*Microchirus*)
boscanion (Chabanaud, 1926)
ocellatus (Linnaeus, 1758)
variegatus (Donovan, 1808)
Microchirus (*Zevia*)
azevia (Capello, 1867)
hexopthalmus (Bennett, 1831)
Monochirus
hispidus Rafinesque, 1814
Pegasa
lascaris (Risso, 1810)
nasuta (Pallas, 1814)
Solea
aegyptiaca Chabanaud, 1927
senegalensis Kaup, 1858
solea (Linnaeus, 1758)
Synaptura
kleinii (Risso, 1827)
lusitanica Capello, 1868
- Family Cynoglossidae**
Cynoglossus
browni Chabanaud, 1949
sinusarabici (Chabanaud, 1931)
- Syphurus*
insularis Munroe, Brito & Hernandez, 2000
ligulatus (Cocco, 1844)
nigrescens Rafinesque, 1810
reticulatus Munroe, 1990
- Order Tetraodontiformes**
- Family Balistidae**
Balistes
carolinensis Gmelin, 1789
punctatus Gmelin, 1789
vetula Linnaeus, 1758
- Canthidermis*
maculata (Bloch, 1786)
sufflamen (Mitchill, 1815)
- Family Monacanthidae**
Aluterus
monoceros (Linnaeus, 1758)
- Stephanolepis*
diaspros Fraser-Brunner, 1940
hispidus (Linnaeus, 1766)
- Family Ostraciidae**
Acanthostracion
notacanthus (Bleeker, 1863)
- Family Tetraodontidae**
Canthigaster
rostrata (Bloch, 1786)
sanctaehelenae (Günther, 1870)
- Ephippion*
guttifer (Bennett, 1831)
- Lagocephalus*
lagocephalus (Linnaeus, 1758)
spadiceus (Richardson, 1845)
- Sphoeroides*
marmoratus (Lowe, 1838)
pachygaster (Müller & Troschel, 1848)
- Family Diodontidae**
Chiromycterus
aringa (Linnaeus, 1758)
- Diodon*
eydouxii Brisout de Barneville, 1846
holocanthus Linnaeus, 1758
hystrix Linnaeus, 1758
- Family Molidae**
Masturus
lanceolatus (Liénard, 1840)
- Mola*
mola (Linnaeus, 1758)
- Ranzania*
laevis (Pennant, 1776)

TETRAPODA

Compiled by Jacob van der Land

The European list of Reptilia, Aves and Mammalia was derived from URMO, and checked by M. Ramos and J. Templado. The Mammalia was based on Jeffersen *et al.* (1996).

References

Jeffersen, T.A., S. Leatherwood & M.A. Webber, 1996. *Marine Mammals of the World*. (ETI CD-ROM).

Order Testudines

Family Cheloniidae

<i>Caretta</i>		A M
<i>carettina</i> (Linnaeus, 1758)		
<i>Chelonia</i>		A M
<i>mydas</i> (Linnaeus, 1758)		
<i>Eretmochelys</i>		A M
<i>imbricata</i> (Linnaeus, 1766)		
<i>Lepidochelys</i>		A
<i>kempii</i> (Garman, 1880)		

Family Dermochelyidae

<i>Dermochelys</i>		A M
<i>coriacea</i> (Vandelli, 1761)		

Order Anseriformes

Family Anatidae

<i>Somateria</i>		A
<i>mollissima</i> (Linnaeus, 1758)		
<i>spectabilis</i> (Linnaeus, 1758)		
<i>Clangula</i>		A
<i>hyemalis</i> (Linnaeus, 1758)		
<i>Melanitta</i>		A
<i>nigra</i> (Linnaeus, 1758)		
<i>fusca</i> (Linnaeus, 1758)		

Order Procellariiformes

Family Oceanitidae

<i>Hydrobates</i>		A M
<i>pelagicus</i> (Linnaeus, 1758)		
<i>Oceanites</i>		A M
<i>oceanicus</i> (Kuhl, 1820)		
<i>Oceanodroma</i>		A
<i>castro</i> (Harcourt, 1851)		
<i>leucorhoa</i> (Vieillot, 1818)		A M

Family Procellariidae

<i>Bulweria</i>		A
<i>bulwerii</i> (Jardine & Selby, 1828)		
<i>Calonectris</i>		A
<i>diomedea</i> (Scopoli, 1769)		
<i>Fulmarus</i>		A
<i>glacialis</i> (Linnaeus, 1761)		
<i>Pterodroma</i>		A
<i>arminjoniana</i> (Giglioli & Salvadori, 1869)		
<i>feae</i> Mathews, 1934		
<i>madeirensis</i> Mathews, 1934		
= <i>P. mollis</i> subsp. <i>madeira</i>		
<i>Puffinus</i>		A
<i>assimilis</i> (Gould, 1838)		
<i>gravis</i> (O'Reilly, 1818)		
<i>griseus</i> (Gmelin, 1789)		A M
<i>puffinus</i> (Brünnich, 1764)		A
<i>yelkouan</i> (Acaerbi, 1827)		A M
= <i>P. puffinus</i> subsp. <i>yelkouan</i>		

Family Gaviidae

<i>Gavia</i>		M
<i>adamsii</i> (Gray, 1859)		
<i>arctica</i> (Linnaeus, 1758)		M
<i>immer</i> (Brünnich, 1764)		
<i>stellata</i> (Pontoppidan, 1763)		M

Order Pelecaniformes

Family Pelecanidae

<i>Pelecanus</i>		
<i>crispus</i> Bruch, 1832		
<i>onocrotalus</i> Linnaeus, 1758		
Family Phaethontidae		
<i>Phaethon</i>		A M
<i>aethereus</i> Linnaeus, 1758		
Family Phalacrocoracidae		
<i>Phalacrocorax</i>		
<i>aristotelis</i> (Linnaeus, 1761)		A M
<i>carbo</i> (Linnaeus, 1758)		A M
Family Sulidae		
<i>Sula</i>		A
<i>bassana</i> (Linnaeus, 1758)		
Order Charadriformes		
Family Alcidae		
<i>Alca</i>		A
<i>torda</i> Linnaeus, 1758		
<i>Alle</i>		A
<i>alle</i> (Linnaeus, 1758)		
<i>Cephus</i>		A
<i>grylle</i> (Linnaeus, 1758)		
<i>Fratercula</i>		A
<i>arctica</i> (Linnaeus, 1758)		
<i>Pinguinus</i>		A
<i>impennis</i> (Linnaeus, 1758)		
<i>Uria</i>		A
<i>aalge</i> (Pontoppidan, 1763)		
<i>lomvia</i> (Linnaeus, 1758)		A
Family Laridae		
<i>Chlidonias</i>		A
<i>hybridus</i> Pallas, 1811		
<i>leucopterus</i> (Temminck, 1815)		
<i>niger</i> (Linnaeus, 1758)		
<i>Larus</i>		A
<i>argentatus</i> Pontoppidan, 1763		
<i>audouinii</i> Payraudeau, 1826		A M
<i>cachinnans</i> Pallas, 1811		
<i>canus</i> Linnaeus, 1758		
<i>fuscus</i> Linnaeus, 1758		A
<i>genei</i> Brême, 1839		A M
<i>glaucocephalus</i> Meyer, 1822		A
<i>hyperboreus</i> Gunnerus, 1767		A
<i>marinus</i> Linnaeus, 1758		A
<i>melaenophanus</i> Temminck, 1820		M
<i>minutus</i> Pallas, 1776		
<i>ridibundus</i> Linnaeus, 1766		
<i>sabini</i> Sabine, 1819		A
= <i>Xema sabini</i>		
<i>tridactyla</i> (Linnaeus, 1758)		
<i>Pagophila</i>		A
<i>eburnea</i> (Phipps, 1774)		
<i>Rhodostethia</i>		
<i>rosea</i> (Macgillivray, 1924)		A
= <i>Larus rossii</i>		
<i>Rissa</i>		
<i>tridactyla</i> (Linnaeus, 1758)		
= <i>Larus tridactyla</i>		
Sterna		
<i>albifrons</i> Pallas, 1764		A
<i>bengalensis</i> Lesson		M
<i>caspia</i> Pallas, 1770		A M
<i>dougallii</i> Montagu, 1813		
<i>fuscata</i> Linnaeus, 1766		
<i>hirundo</i> Linnaeus, 1758		
<i>maxima</i> Boddaert, 1791		A
<i>nilotica</i> Gmelin, 1789		A M
<i>paradisea</i> Pontoppidan, 1763		
<i>sandvicensis</i> Latham, 1787		A M
Family Phalaropodidae		
<i>Phalaropus</i>		
<i>fulicarius</i> (Linnaeus, 1758)		A
<i>lobatus</i> (Linnaeus, 1758)		A
Family Stercorariidae		

Catharacta			
<i>skua</i> Brünnich, 1764	A	<i>mirus</i> True, 1913	A M
= <i>Stercorarius skua</i>		<i>Ziphius</i>	
Stercorarius		<i>cavirostris</i> Cuvier, 1823	A M
<i>longicaudus</i> Vieillot, 1819	A	Family Balaenidae	
<i>parasiticus</i> (Linnaeus, 1758)	A	<i>Balaena</i>	
<i>pomarinus</i> (Temminck, 1815)	A	<i>mysticetus</i> Linnaeus, 1758	
Order Podicipediformes		Family Balaenopteridae	
Family Podicipedidae		<i>Balaenoptera</i>	
<i>Podicipes</i>		<i>acutorostrata</i> Lacépède, 1804	A M
<i>cristatus</i> (Linnaeus, 1758)	A	<i>borealis</i> Lesson, 1828	A M
<i>griseogenus</i> Boddaert, 1783		<i>edeni</i> Anderson, 1871	A M
<i>auritus</i> (Linnaeus, 1758)	A	<i>musculus</i> (Linnaeus, 1758)	A M
<i>nigricollis</i> Brehm, 1831		<i>physalus</i> (Linnaeus, 1758)	A M
Order Cetacea		<i>Megaptera</i>	
Family Delphinidae		<i>novaeangliae</i> (Borowski, 1781)	A M
<i>Delphinus</i>		Family Eschrichtiidae	
<i>capensis</i> Gray, 1828	A M	<i>Eschrichtius</i>	
<i>delphis</i> Linnaeus, 1758	A	<i>robustus</i> (Lilljeborg, 1861)	A
<i>Feresa</i>		Order Carnivora	
<i>attenuata</i> Gray, 1875	A M	Family Mustelidae	
<i>Globicephala</i>		<i>Lutra</i>	
<i>macrorhynchus</i> Gray, 1846	A M	<i>lutra</i> (Linnaeus, 1758)	A
<i>melas</i> (Traill, 1809)	A M	Family Ursidae	
<i>Grampus</i>		<i>Ursus</i>	
<i>griseus</i> (Cuvier, 1812)	A M	<i>maritimus</i> Phipps, 1774	A
<i>Lagenodelphis</i>		Family Odobenidae	
<i>hosei</i> Fraser, 1956	A M	<i>Odobenus</i>	
<i>Lagenorhynchus</i>		<i>rosmarus</i> (Linnaeus, 1758)	A
<i>acutus</i> (Gray, 1828)	A	Family Phocidae	
<i>albirostris</i> (Gray, 1846)	A M	<i>Cystophora</i>	
<i>Orcinus</i>		<i>cristata</i> (Erxleben, 1777)	A
<i>orca</i> (Linnaeus, 1758)	A M	<i>Erignathus</i>	
<i>Peponocephala</i>		<i>barbatus</i> (Erxleben, 1777)	A
<i>electra</i> (Gray, 1846)	A	<i>Halichoerus</i>	
<i>Pseudorca</i>		<i>grypus</i> (Fabricius, 1791)	A
<i>crassidens</i> (Owen, 1846)	A M	<i>Monachus</i>	
<i>Stenella</i>		<i>monachus</i> (Hermann, 1779)	M
<i>attenuata</i> (Gray, 1846)	A M	<i>Phoca</i>	
<i>clymene</i> (Gray, 1850)	A	<i>groenlandica</i> (Erxleben, 1777)	A
<i>coeruleoalba</i> (Meyen, 1833)	A M	<i>hispida</i> (Schreber, 1775)	A
<i>frontalis</i> (Cuvier, 1829)	A	<i>vitulina</i> (Linnaeus, 1758)	
<i>longirostris</i> (Gray, 1828)	A M		
<i>Steno</i>			
<i>bredanensis</i> (Lesson, 1828)	A M		
<i>Tursiops</i>			
<i>truncatus</i> (Montagu, 1821)	A M		
Family Kogiidae			
<i>Kogia</i>			
<i>breviceps</i> (de Blainville, 1838)	A M		
<i>simus</i> Owen, 1866			
Family Monodontidae			
<i>Delphinapterus</i>			
<i>leucas</i> (Pallas, 1776)			
<i>Monodon</i>			
<i>monoceros</i> Linnaeus, 1758	A		
Family Phocoenidae			
<i>Phocoena</i>			
<i>phocoena</i> (Linnaeus, 1758)	A		
Family Physeteridae			
<i>Physeter</i>			
<i>macrocephalus</i> Linnaeus, 1758	A M		
= <i>P. catodon</i>			
Family Ziphiidae			
<i>Hyperoodon</i>			
<i>ampullatus</i> (Forster, 1770)	A		
<i>Mesoplodon</i>			
<i>bidens</i> (Sowerby, 1804)	A		
<i>densirostris</i> (de Blainville, 1817)	A M		
<i>europeaeus</i> Gervais, 1855	A		
<i>grayi</i> von Haast, 1876	A M		
<i>hectori</i> (Gray, 1871)	A M		

BIBLIOGRAPHY OF IDENTIFICATION GUIDES TO MARINE SPECIES IN EUROPE

Compiled by Philippe Bouchet and Julien Marmayou

The bibliography benefited from input by Nadia Ameziane, Guy Boucher, Sylvie Crasquin, Christer Erseus, Jean-Loup d'Hondt, Jean-Claude Hureau, Jean-Lou Justine, Claude Lévi, Françoise and Claude Monniot, Pierre Noël, Fredrik Pleijel, Bruno de Reviers, and Daniel Robineau.

Key references in bold.

GENERAL

- Bianchi, C. N., Dore, G. & Morri, C. 1995 *Guida del subacqueo naturalista. Mediterraneo e Tropici*. Sassari: Archivio Fotografico Sardo-Nuoro. 284 pp.
- Campbell, A. C. & Nicholls, J. 1979 *Guide de la faune et de la flore littorale des mers d'Europe*. Neuchâtel, Paris: Delachaux & Niestlé. 322 pp.
- Campbell, A. C. 1982 *The Hamlyn guide to the flora and fauna of the Mediterranean sea*. London: Hamlyn. 320 pp.
- Campbell, A. C. 1984 *Guía de la flora y fauna del Mar Mediterráneo*. Barcelona: Omega. 333 pp.
- Campbell, A. C. 1994 *Hamlyn Guide to Seashores and Shallow Seas of Britain and Europe*. 320 pp.
- Castric, A., Chassé, C. & Michel, C. 1984 *Flore et faune fixées sous-marines de Bretagne*. Brest, Concarneau: Laboratoire Maritime, Concarneau/Université de Bretagne Occidentale, Laboratoire d'Océanographie Biologique/ADMS Laboratoire Maritime
- Departamento de Biología. Facultad de Biología. University of La Laguna, 1982 *Estudio del Benthos Marino del Archipiélago Canario*: Gobierno de Canarias. Consejería de Agricultura y pesca. 807 pp.
- Erwin, D. & Picton, B. 1987 *Guide to inshore marine life*. London: Immel Publishing. 120 pp.
- Erwin, D. G., Picton, B. E., Connor, D. W., Howson, C. M., Gilleece, P. & Bogues, M. J. 1990 *Inshore Marine Life of Northern Ireland*. Belfast: HMSO. 148 pp.
- Fechter, R. 1992 *Fauna y flora de las costas plantas y animales de las costas europeas (Mediterraneo, Atlántico, Mar del Norte y Báltico)*. Barcelona: Naturart. 287 pp.
- Fiala-Médioni, A., Pétron, C. & Rives, C. 1987 *Guía submarina del Mediterráneo*. Madrid: Mundi-Prensa. 156 pp.
- Fischer, W., Bauchot, M.-L. & Schneider, M. (ed.) 1987 *Fiches FAO d'identification des espèces pour les besoins de la pêche. Méditerranée et Mer Noire, Zone de Pêche 37, revision 1. Volume I. Végétaux et Invertébrés*. Rome: FAO/CEE. xvi+760 pp.
- Fish, J. D. & Fish, S. 1996 *A student's guide to the seashore*. Cambridge: Cambridge University Press. 564 pp.
- García-Raso, J. E., Luque, A. A., Templado, J., Salas, C., Hergueta, E., Moreno, D. & Calvo, M. 1992 *Fauna y flora marinas del Parque natural de Cabo de Gata-Níjar*. Madrid. 288 pp.
- Hass, H. 1978 *Guida del subacqueo. II Mediterraneo*. Milano: Garzanti. 295 pp.
- Luther, W. & Fiedler, K. 1976 *Field Guide to the Mediterranean Sea Shore*. London: Collins. 272 pp.
- Melone, N., Picchetti, G. & Paccagnella, W. 1977 *Vita del mare. Guida Illustrata della fauna e flora del Mediterraneo*, vol. I. Milano: Primaris
- Melone, N. & Picchetti, G. 1980 *Vita del mare. Guida Illustrata della fauna e flora del Mediterraneo*, vol. II. Milano: Primaris. 172 pp.
- Moreno, I., Roca, I. & Salamanca, M. 1991 *Guía de la flora y fauna del mar balear*. Mallorca: Govern Balear. 279 pp.
- Riedl, R. 1970 *Fauna und Flora der Adria*. Hamburg: Paul Parey. 702 pp.
- Riedl, R. 1983 *Fauna und Flora des Mittelmeeres*. Hamburg, Berlin: Verlag Paul Parey. 836 pp.
- Riedl, R. 1986 *Fauna y flora del mar Mediterraneo*. Barcelona: Omega. 858 pp.
- Riedl, R. 1991 *Fauna e flora del Mediterraneo*. Padova: F. Muzzio. 777 pp.
- Schick, K. L. 1998 *Atlas submarino de la Costa del Sol*. Marbella (Málaga): Edición Schick. 71 pp.
- Simeonidis, D. 1997 *Guide sous-marin de l'Atlantique, côtes et îles de Dunkerque à Biarritz*. Lausanne, Paris: Delachaux & Niestlé. 160 pp.
- Simeonidis D. 1995 *Guide de la faune et de la flore sous-marines du bassin méditerranéen*. Lausanne: Delachaux & Niestlé. 160 pp.
- Tréguoboff, G. & Rose, M. 1957 *Manuel de planctonologie méditerranéenne*. Paris: CNRS. 587 pp.
- Weinberg, S. 1997 *Découvrir l'Atlantique, la Manche et la mer du Nord*. Nathan Nature. Paris: Nathan. 383 pp.

FUNGI

- Johnson, T. W. & Sparrow, F. K. 1961 *Fungi in oceans and estuaries*. Bibliotheca mycologica. 26. New York: Weinheim J. Kramer. xxii+668 pp.
- Kohlmeyer, J. & Kohlmeyer, E. 1979 *Marine mycology: the higher Fungi*. New York: Academic Press. xiv+690 pp.
- Kohlmeyer, J. & Volkmann-Kohlmeyer, B. 1991 *Illustrated key to filamentous higher marine fungi*. Botanica marina 38: 165-174.

PLANTS

General and other algal taxa

- Aasen, P. A. 1980 *Illustrert algeflora*. Oslo: Cappelen forlag. 64 pp.
- Ardré, F. 1970 Contribution à l'étude des algues du Portugal. I. La Flore. *Portugaliae Acta Biologica*, ser. B 10: 1-423.
- Bellinger, E. G. 1992 *A key to common algae. Freshwater, estuarine and some coastal species*. London: The Institution of Water and Environment Management
- Berthold, G. 1884 *Die Cryptonemaceen des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 12. Leipzig: W. Engelmann. 27 pp.
- Bourgeois, P. & Desprez, M. 1984 *Le littoral: 123 espèces rassemblées et dessinées*. Flore algale de Normandie. xvi+28 pp.
- Cabioch, J., Floc'h, J.-Y., Le Toquin, A., Boudouresque, C.-F., Meinesz, A. & Verlaque, M. (ed.) 1992 *Guide des Algues des Mers d'Europe. Manche, Atlantique, Méditerranée*. Lausanne: Delachaux & Niestlé. 231 pp.
- Cardinal, A. 1964 *Etudes sur les Ectocarpacées de la Manche*. Beihefte zur Nova Hedwigia. 15. 87 pp.
- Carrillo, J. A. & Sansón, M. 1999 *Algas Hongos y Fanerógamas Marinas de las Islas Canarias*: University of La Laguna. 254 pp.
- Chrétiennot-Dinet, M.-J. 1990 *Atlas du phytoplancton marin. III. Chlorarachniophycées, Chlorophycées, Clrysophycées, Cryptophycées, Euglenophycées, Eustigmatophycées, Prasinophycées, Prymnésiophycées, Rhodophycées et Tribophycées*. Paris: C.N.R.S. 261 pp.
- Christensen, T. 1987 *Tribophyceae (Xanthophyceae). Seaweeds of the British Isles*. 4. London: British Museum (Natural History). 36 pp.
- Coppejans, E. 1983 *Iconographie d'algues Méditerranéennes Chlorophyta, Phaeophyta, Rhodophyta*. Bibliotheca Phycologica. 63. xxvii+151 pp.
- Coppejans, E. 1995 Flore algologique des côtes du Nord de la France et de la Belgique. *Scripta Botanica Belgica* 9: 1-454.

- Feldmann, J. 1940 La végétation benthique de la Méditerranée. *Mémoires de la Société de Biogéographie* 7: 181-195.
- Feldmann, J. & Hamel, G. 1936 Floridées de France. VII. *Revue algologique* 9(1): 85-140.
- Gayral, P. 1966. *Les Algues des côtes françaises (Manche et Atlantique). Notions fondamentales sur l'écologie, la biologie et la systématique des algues marines*. Paris: Doin. 632 pp.
- Gayral, P. & Casson, J. 1986 *Connaître et reconnaître les algues marines*. Rennes: Ouest-France. 220 pp.
- Gruet, Y. & Le Neustiec, R. 1989 Algues des côtes rocheuses de Loire-Atlantique et de Vendée: richesses naturelles du littoral. *Bulletin de la Société de Sciences Naturelles de l'Ouest de la France*: 1-90.
- Hamel, G. 1924 Floridées de France. II. *Revue algologique* 1(4): 427-457.
- Hamel, G. 1925 Floridées de France. III. *Revue algologique* 2(1): 39-67.
- Hamel, G. 1925 Floridées de France. IV. *Revue algologique* 2(3-4): 280-309, pls 9-10.
- Hamel, G. 1928 Floridées de France. V. *Revue algologique* 3(1-4): 99-158.
- Hamel, G. 1930 Floridées de France. VI. *Revue algologique* 5(1): 61-109.
- Hamel, G. & Lemoine, M. 1953 Corallinacées de France et d'Afrique du Nord. *Archives du Muséum national d'Histoire naturelle* sér. 7 1: 15-136.
- Hoffmann, C. 1933 *Die Vegetation der Nord- und Ostsee. Tierwelt der Nord- und Ostsee*. 1c. Leipzig: Akademische Verlagsgesellschaft. 32 pp.
- Jaasund, E. 1965 Aspects of the marine algal vegetation of North Norway. *Botanica Gothoburgensis* 4: 1-174.
- Jones, W. E. A key to the Genera of the British Seaweeds. Menai Bridge
- Kiselev, I. A. 1950 *Dinoflagellata of the seas and freshwaters of the USSR*. Opredelitel po faune SSSR. 33: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 280 pp.
- Kornmann, P. & Sahling, P.-H. 1994 Meeresalgen von Helgoland. Zweite Ergänzung. *Helgol. Meeresunters* 48: 365-406.
- Kremer, B. 1975 *Meeresalgen. Ein Bestimmungsbuch für häufigere Arten des Atlantikküsten, der Nord- und Ostsee*. Ziemsen: Die Neue Brehm-Bücherei. 152 pp.
- Levring, T. 1937 Zur Kenntnis der Algenflora der norwegischen Westküste. *Lunds Universitets Årsskrifter. N. F. Avd. 2* 31(8): 1-148.
- Maggs, C. A. & Howson, C. M. 1985 *A photographic guide to some common subtidal seaweeds of the British Isles*. Ross-on-Wye: Marine Conservation Society
- Newton, L. 1931 *A handbook of the British seaweeds*. The trustees of the British Museum (Natural History). London. xiii+478 pp.
- Reichenow, E. 1930 *Parasitische Peridinea (einschliesslich Ellobiopsidae)*. Tierwelt der Nord- und Ostsee. 2d3. Leipzig: Akademische Verlagsgesellschaft. Pp. 85-100.
- Reichenow, E. 1931 *Parasitische Flagellata (ausschliesslich Peridinea)*. Tierwelt der Nord- und Ostsee. 2e1. Leipzig: Akademische Verlagsgesellschaft. 18 pp.
- Ribier, J. & Godineau, J.-C. 1984 *Les algues: connaissance, utilisation, culture. Avec des clés simples de détermination des algues du littoral de la Manche et de l'Atlantique*. Paris: Maison rustique, Flammarion. 281 pp.
- Ricard, M. 1987 *Atlas du phytoplancton marin. II. Diatomophycées*. Paris: CNRS 297 pp.
- Rueness, J. 1977 *Norsk algeflora*. Oslo: University Press. 266 pp.
- Skyes, J. B. 1981 *An illustrated guide to the diatoms of British Coastal plankton*. AIDGAP Guides. 5. Shrewbury: Field Studies Council. Pp. 425-468.
- Sournia, A. 1986 *Atlas du phytoplancton marin. I. Cyanophycées, Dictyochophycées, Dinophycées et Raphidophycées*. Paris: CNRS 219 pp.
- Sundene, O. 1953 The algal vegetation of Oslofjord. *Skrifter. Det Norske Videnskaps-akademiet i Oslo. I. Matematisk-Naturvitenskapelig Klasse* 1953(2): 1-244.
- Tomas, C. R. (ed.) 1996 *Identifying marine diatoms and dinoflagellates*. San Diego: Academic Press. xiii+598 pp.
- Tomas, C. R. (ed.) 1997 *Identifying marine phytoplankton*. San Diego: Academic Press. xv+858 pp.
- Tomas, C. R., Thronsdson, J. & Heimdal, B. 1993 *Marine phytoplankton: a guide to naked flagellates and coccolithophorids*. San Diego: Academic Press. xiii+263 pp.
- ### Seaweeds
- #### Bangiophyceae
- Barthold, G. 1882 *Die Bangiaceen des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 8. Leipzig: W. Engelmann. 28 pp.
- Hamel, G. 1924 *Floridées de France. I. Bangiales*. Paris. 15 pp.
- #### Chlorophyta
- Bliding, C. 1963 A critical survey of European taxa in Ulvales. I. Capsosiphon, Percusaria, Blidingia, Enteromorpha. *Bot. Not.* 8: 3-160.
- Bliding, C. 1969 A critical survey of European taxa in Ulvales. II. Ulva, Ulvaria, Monostroma, Kormannia. *Bot. Not.* 121: 535-629.
- Burrows, E. M. 1991 *Chlorophyta. Seaweeds of the British Isles. 2*. London: British Museum (Natural History). 238 pp.
- Hamel, G. 1930 Chlorophycées des côtes françaises. I. *Revue algologique* 5(1): 1-54.
- Hamel, G. 1931 Chlorophycées des côtes françaises. II. *Revue algologique* 5(3-4): 383-430.
- Hamel, G. 1931 Chlorophycées des côtes françaises. III. *Revue algologique* 6(1): 9-73.
- Hoek van den, C. 1963 *Revision of the European species of Cladophora*. Leiden: E.J. Brill éd. 248 pp.
- Söderström, J. 1963 *Studies in Cladophora*. Bot. Gotoburg. I: Act. Univer. Gotoburg. 147 pp.
- #### Phaeophyta
- Fletcher, R. L. 1987 *Fucophyceae (Phaeophyceae) Part 1. Seaweeds of the British Isles. 3*. London: British Museum (Natural History). 360 pp.
- Hamel, G. 1931 *Phéophycées de France. Part 1*. Paris: chez l'auteur. 80 pp.
- Hamel, G. 1935 *Phéophycées de France. Part 2*. Paris: chez l'auteur. Pp. 81-176.
- Hamel, G. 1937 *Phéophycées de France. Part 3*. Paris: chez l'auteur. Pp. 177-240.
- Hamel, G. 1938 *Phéophycées de France. Part 4*. Paris: chez l'auteur. Pp. 241-336.
- Hamel, G. 1939 *Phéophycées de France. Part 5*. Paris: chez l'auteur. xlvi+pp. 337-431.
- Hiscock, S. 1979 *A field key to the British brown seaweeds (Phaeophyta)*. AIDGAP Guides. 5. Shrewbury: Field Studies Council. 44 pp.
- Valiante, R. 1883 *Le Cystoceriae del Golfo di Napoli*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 7. Leipzig: W. Engelmann. 30 pp.
- #### Rhodophyta
- Bird, C. J. & McLachlan, J. L. 1992 Seaweed flora of the Maritimes I. Rhodophyta - the red algae. Bristol: Biopress
- Dixon, P. S. & Irvine, L. M. 1977 Rhodophyta. Part 1. Introduction, Nemaliales, Gigartinales. Seaweeds of the British Isles. 1. London: British Museum (Natural History). 252 pp.
- Falkenberg, P. 1901 *Die Rhodomelaceen des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 26. Berlin: R. Friedländer & Sohn. 754 pp.
- Hiscock, S. 1986 A field key to the British red seaweeds (Rhodophyta). AIDGAP guides. 13. London: Field Studies Council. 95 pp.
- Irvine, L. M. 1983 Rhodophyta. Part 2A. Cryptonemiales (sensu stricto), Palmariales, Rhodymeniales. Seaweeds

- of the British Isles. 1. London: British Museum (Natural History). 115 pp.
- Irvine, L. M. & Chamberlain, Y. M. 1994 Rhodophyta. Part 2B Corallinales, Hildenbrandiales. Seaweeds of the British Isles. 1. London: British Museum (Natural History). 276 pp.
- Maggs, C. A. & Hommersand, M. H. 1993 Rhodophyta. Part 3A: Ceramiales. Seaweeds of the British Isles. 1. London: HMSO. 464 pp.
- Solms-Laubach, G. 1881 Die corallinenalgen des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 4. Leipzig: W. Engelmann. 64 pp.
- ### PROTISTA
- Brandt, K. 1885 *Die koloniebildenden Radiolarien (Sphaerozoen) des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 13. Berlin: R. Friedländer & Sohn. 276 pp.
- Carey, P. G. 1992 *Marine interstitial ciliates. An illustrated key*. Chapman and Hall Identification guides. 2. London, New York: Chapman and Hall. xiii+351 pp.
- Foissner, W. 1993 Colpodea (Ciliophora). *Protozoenfauna* 4(1): 1-798.
- Jorgensen, E. 1927 *Ciliata: Tintinnidae*. Tierwelt der Nord- und Ostsee. 2 cl. 33 pp.
- Kahl, A. 1930 *Urtiere oder Protozoa. I. Wimpertiere oder Ciliata (Infusoria)*. Eine Bearbeitung der freilebenden und ectocommensalen Infusorien der Erde, unter Ausschluss der marin Tintinnidae. 1. Allgemeiner Teil un Protostomata IV. Die Tierwelt Deutschlands. 18. Jena: G. Fischer. 180 pp.
- Kahl, A. 1931 *Urtiere oder Protozoa. I. Wimpertiere oder Ciliata (Infusoria)*. Eine Bearbeitung der freilebenden und ectocommensalen Infusorien der Erde, unter Ausschluss der marin Tintinnidae. 2. Holotrichia ausser den inn 1 Teil behandelten Protostomata. Die Tierwelt Deutschlands. 21. Jena: G. Fischer. Pp. 181-398.
- Kahl, A. 1932 *Urtiere oder Protozoa. I. Wimpertiere oder Ciliata (Infusoria)*. Eine Bearbeitung der freilebenden und ectocommensalen Infusorien der Erde, unter Ausschluss der marin Tintinnidae. 3. Spirotricha. Die Tierwelt Deutschlands. 25. Jena: G. Fischer. Pp. 399-650.
- Kahl, A. 1933 *Ciliata libera et ectocommensalia*. Tierwelt der Nord- und Ostsee. 2 c3. Leipzig: Akademische verlagsgesellschaft. Pp. 29-146.
- Kahl, A. 1934 *Suctorria*. Tierwelt der Nord- und Ostsee. 2 c5. Leipzig: Akademische verlagsgesellschaft. Pp. 184-226.
- Kahl, A. 1934 *Ciliata ectocommensalia et parasitica*. Tierwelt der Nord- und Ostsee. 2 c4. Leipzig: Akademische verlagsgesellschaft. Pp. 147-183.
- Kahl, A. 1935 *Urtiere oder Protozoa. I. Wimpertiere oder Ciliata (Infusoria)*. Die Tierwelt Deutschlands. 30. Jena: G. Fischer. Pp. 651-886.
- Marshall, S. M. 1969 *Protozoa, order: Tintinnida*. Fiches d'Identification du Zooplanton. 117-127. 70 pp.
- Petrushevskaya, M. G. 1981 [Radiolaria of the order Nassellaria of the world's oceans] [in Russian]. Opredelitel po faune SSSR. 128. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 405 pp.
- Reichenow, E. 1931 *Sporozoa*. Tierwelt der Nord- und Ostsee. 2gl. Leipzig: Akademische verlagsgesellschaft. 48 pp.
- Reichenow, E. 1931 *Cnidosporidia*. Tierwelt der Nord- und Ostsee. 2g2. Leipzig: Akademische verlagsgesellschaft. Pp. 49-88.
- Schewiakoff, W. 1926 *Die Acantharia des Golfs von Neapel*. Fauna e flora del Golfo di Napoli. 37. Berlin: R. Friedländer & Sohn. 755 pp.
- Zuelzer, M. 1931 *Spirochaeta*. Tierwelt der Nord- und Ostsee. 2f. Leipzig: Akademische verlagsgesellschaft. 22 pp.

Foraminifera

- A major group of the meiobenthos and a minor group of the plankton (1161 species in the ERMS area). Coverage patchy all over.
- Be, A. W. H. 1967 *Foraminifera. Families : Globigerinidae and Globorotaliidae*. Fiches d'Identification du Zooplancton. 108. 9 pp.
- Christiansen, B. O. 1958 The Foraminifer Fauna in the Dröbak Sound in the Oslo Fjord (Norway). *Nytt Magasin for Zoologi* 6: 5-91.
- Cimernan, F. & Langer, M. R. 1991 *Mediterranean Foraminifera*. Ljubljana: Slovenska Akademija Znanosti in Umetnosti. 118 pp.
- Colom, G. 1974 Foraminiferos ibéricos. Introducción al estudio de las especies bentónicas recientes. *Investigación Pesquera* 38(1): 1-245.
- Murray, J. W. 1979 *British nearshore foraminiferids*. Synopses of the British Fauna (New Series). 16. London, New-York: Academic Press. vii+68 pp.
- Rhumbler, L. 1928 *Amoebozoa et Reticulosa (Foraminifera)*. Tierwelt der Nord- und Ostsee. 2a1. Leipzig: Akademische verlagsgesellschaft. 26 pp.

FAUNA

GENERAL

- Andrews, M. J., Aston, K. F. A., Rickard, D. G. & Steel, J. E. C. 1982 The macrofauna of the Thames Estuary. *London Naturalist* 61: 30-61.
- Barnes, R. S. K. 1994 *The brackish-water fauna of Northwestern Europe*. Cambridge: Cambridge University Press. 287 pp.
- Clément, J.-L. 1978 *Invertébrés du littoral Sud Armoricain*. Nantes: Cercle Naturaliste Étudiant Nantais, Faculté des Sciences. 92 pp.
- Crothers, J. 1994 *A key to the major groups of the British marine invertebrates*. AIDGAP Guides. Shrewsbury: Field Studies Council. 194 pp.
- Dahl, F. 1925 *Säugetiere, Vögel, Kriechtiere, Lurche, Fische, Weichtiere nebst Bestimmungsschlüssel für die Klassen und Ordnungen der Tiere und Für Vogelnester XXXV*. Die Tierwelt Deutschlands. 1. Jena: G. Fischer. 207 pp.
- Eno, N. C., Clark, R. A. & Sanderson, W. G. (ed.) 1997 *Non-native marine species in British waters: a review and directory*. Peterborough: Joint Nature Conservation Committee. 152 pp.
- Gosálvez López, M., Fernández Rubio, F. & Martínez Martínez, J. 1992 *Guía de la Fauna submarina del litoral mediterráneo continental español*. Madrid: Pirámide. 253 pp.
- Hannemann, H. J., Klausnitzer, B. & Senglau, K. 1992 *Wirbellose (ohne Insekten)*. Exkursionsfauna von Deutschland. 1. Berlin: Volk und wissen verlag GmbH. 637 pp.
- Hayward, P. J. 1988 *Animals on Seaweed*. Naturalists' Hanbooks. 9. Richmond: Richmond Publ. Co. 108 pp.
- Hayward, P. J. & Ryland, J. S. 1990 *The marine fauna of the British Isles and north-west Europe. Volume 1. Introduction and protozoans to arthropods*. Oxford: Clarendon Press. xvi+627 pp.
- Hayward, P. J. & Ryland, J. S. 1990 *The marine fauna of the British Isles and north-west Europe. Volume 2. Molluscs to chordates*. Oxford: Clarendon Press. xvi + pp. 628-996.
- Hayward, P. J. 1994 *Animals of sandy shores*. Naturalists' Handbooks. 21. Slough: Richmond Publ. Co. 104. pp.
- Hayward, P. J. & Ryland, J. S. 1995 *Handbook of the marine fauna of north-west Europe*. Oxford: Oxford University Press. x+800 pp.
- López Alegret, P. L. 1983 *La fauna submarina del mar Mediterráneo*. Barcelona: Omega. 229 pp.
- Luther, W. & Fiedler, K. 1986 *Guida della fauna marina costiera del Mediterraneo*. Padova: Franco Muzzio. 244 pp.

- Luther, W. & Fiedler, K. 1987 *Guide de la faune sous-marine des côtes méditerranéennes*. Neuchâtel, Paris: Delachaux & Niestlé. 270 pp.
- Micalef, H. & Evans, F. 1968 *The marine fauna of Malta*. Malta University Press. vi + 26 pp.
- Perez Sanchez, J. M. & Moreno Batet, E. 1990 *Invertebrados Marinos de Canarias*. Las Palmas de Gran Canaria: Ediciones del Abildo Insular de Gran Canaria. 335 pp.
- Sims, R. W. 1980 *Marine and brackish water animals*. Animal Identification. A reference guide. I. London: British Museum (Natural History). 111 pp.
- Smith, D. L. & Johnson, K. B. 1996 *Guide to marine coastal plankton and marine invertebrate larvae*. Dubuque: Kendall/Hunt Pub. Co. xv+221 pp.
- Somerfield, P. J. & Warwick, R. M. 1996 *Meiofauna in marine pollution monitoring. A laboratory manual*. Directorate of fisheries research. 71 pp.
- Todd, C. D., Laverack, M. S. & Boxshall, G. 1996 *Coastal marine zooplankton. A practical manual for students*. Cambridge: Cambridge University Press. 106 pp.
- Turquier, Y. & Loir, M. 1992 *Connaitre et reconnaître la faune du littoral*. Rennes: Ouest-France. 332 pp.
- Vodianitzkii, V. A. (ed.) 1972 [*Identification key to the fauna of Black and Azov seas. Volume 3. Free living invertebrates. Arthropods (besides Crustacea), Mollusca, Echinodermata, Chaetognatha, Chordata*]. Kiev: Naukova Dumka.
- Wirtz, P. 1994 *Unterwasserführer Madeira, Kanaren, Azoren: Fische*. Underwater guide Madeira, Canary Islands, Azores: fish. Stuttgart: S. Nagschmid. 259 pp.
- Wirtz, P. 1995 *Unterwasserführer Madeira, Kanaren, Azoren: Niedere Tiere*. Underwater guide Madeira, Canary Islands, Azores: invertebrates. Stuttgart: S. Nagschmid.

PORIFERA

- Ackers, R. G., Moss, D., Picton, B. E. & Stone, S. M. K. 1985 *Sponges of the British Isles ("Sponges IV")*. A colour guide and working document. Ross-on-Wye: Marine Conservation Society. 202 pp.
- Ackers, R. G., Moss, D., Picton, B. E. & Stone, S. M. K. 1992 *Sponges of the British Isles ("Sponge V")*. A colour guide and working document. Ross-on-Wye: Marine Conservation Society. 175 pp.
- Alander, H. 1942 *Sponges from the Swedish west coast and adjacent waters*. Dissertation. Göteborg: Stuves Boktryckeri. 95 pp.
- Arndt, W. 1934 *Porifera*. Tierwelt der Nord- und Ostsee. 3a. Leipzig: Akademische verlagsgesellschaft. 140 pp.
- Arndt, W., Broch, H., Krumbach, T., Pax, F. & Leiberkind, J. 1928 *Porifera, Schwämme, Spongien. Hydrozoen. Scyphozoa. Anthozoa. Ctenophora, Kammquallen oder Rippenquallen. Echinodermata, Stachelhäuter oder Echinodermen*. Die Tierwelt Deutschlands. 4. Jena: G. Fischer. 332 pp.
- Bibiloni, M. A. 1981 Estudio sistemático del orden Poecilosclerida de la costa Brava (Girona). *Boletín del Instituto Español de Oceanografía* 4(324): 103-153.
- Boury-Esnault, N., Pansini, M. & Uriz, M. J. 1994 Spongaires bathyaux de la Mer d'Alboran et du Golfe Ibero-Moroccain. *Mémoires du Muséum national d'Histoire naturelle* 160: 1-174.
- Bowerbank, J. S. 1864 *A monograph of the British Spongiidae*. 1. London: The Ray Society. 290 pp.
- Bowerbank, J. S. 1866 *A Monograph of the British Spongiidae*. 2. London: The Ray Society. 388 pp.
- Bowerbank, J. S. 1874 *A monograph of the British Spongiidae*. 3. London: The Ray Society. 362 pp.
- Bowerbank, J. S. 1882 *A monograph of the British Spongiidae*. 4. London: The Ray Society. 250 pp.
- Carballo, J. L., Sánchez Moyano, J. E. & García Gómez, J. C. 1994 Esponjas del Estrecho de Gibraltar. I. Esponjas cárneas. *Graellsia* 50: 35-56.
- Carballo, J. L. & García Gómez, J. C. 1994 Esponjas del Estrecho de Gibraltar y áreas próximas, con nuevas aportaciones para la fauna ibérica. *Cahiers de Biologie Marine* 35: 193-211.
- Koltun, V. M. 1959 *Cornacuspongida of the northern and Far Eastern seas of the USSR*. Opredelitel po faune SSSR. 67. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 236 pp.
- Koltun, V. M. 1966 *Tetraxonida (Spongia) of the northern and Far Eastern seas of the USSR*. Opredelitel po faune SSSR. 90. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 112 pp.
- Koltun, V. M. 1967 *Hyalospongiae of the northern and Far Eastern seas of the USSR*. Opredelitel po faune SSSR. 94. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 125 pp.
- Lopes, M. T. 1995 *Colheita, preparação e identificação de espongiários*. Lisbon: Museu Nacional de História Natural, Museu e Laboratorio Zoológico e Antropológico (Museu Bocage). 35 pp.
- Martínez Inglés, A. M. 1993 *Guía ilustrada de las esponjas del litoral alicantino*. Alicante: "Juan Gil-Albert". 172 pp.
- Pulitzer-Finali, G. 1983 A collection of Mediterranean Demospongiae (Porifera) with, in appendix, a list of the demospongiae hitherto recorded from the Mediterranean sea. *Annali del Museo Civico di Storia Naturale di Genova* 84: 445-621.
- Rezvoi, P. D. 1936 *Presnovodnye gubki*. Fauna SSSR (novaia seriiia). 3. Moskva: Akademiiia nauk SSSR, Zoologicheskii institut. 124 pp.
- Rubio, M. 1982 Estudio sistemático de las esponjas del orden Halichondrida del litoral de Blanes (Girona) y Alicante. *Boletín del Instituto Español de Oceanografía* 4(321): 60-73.
- Sarà, M. 1972 Guida ai poriferi della fauna italiana. *Quadrerni Civica Stazione Idrobiologica Milano* 3-4: 53-94.
- Soest van, R. W. M., Picton, B. & Morrow, M. 1999 *Sponges of the North East Atlantic*. Amsterdam: ETI. CD-ROM.
- Uriz, M. J. 1978 Estudio sistemático de las esponjas Astrophorida de los fondos de pesca de arrastre entre Tossa y Calella (Cataluña). *Boletín del Instituto Español de Oceanografía* 4(320): 7-57.
- Uriz, M. J. 1986 Clave de identificación de las esponjas más frecuentes de la Península Ibérica. *Miscelánea Zoologica* 10: 7-22.
- Vosmaer, G. C. J. 1933-1935 *The sponges of the Bay of Naples*. 1-3. The Hague: Martinus Nijhoff. 875 pp.

CNIDARIA

General

- Arndt, W., Broch, H., Krumbach, T., Pax, F. & Leiberkind, J. 1928 Porifera, Schwämme, Spongien. Hydrozoen. Scyphozoa. Anthozoa. Ctenophora, Kammquallen oder Rippenquallen. Echinodermata, Stachelhäuter oder Echinodermen. Die Tierwelt Deutschlands. 4. Jena: G. Fischer. 332 pp.
- Carlgren, O. 1945 Polypdyr (Coelenterata). 3. Koraldyr. Danmarks Fauna. 51. Kobenhavn: GEC, Gad. 168 pp.
- George, J. D. & Warren, L. 1986 British coelenterates - a photographic identification guide. Ross-on-Wye: Marine Conservation Society. 8 pp.
- Gili i Sardà, J.-M. 1982 Fauna de cnidaris de les illes Medes. Treballs de la Institució Catalana de Història Natural. 10. Barcelona: Institució Catalana de Història Natural. 175 pp.
- Kramp, P. L. 1935 Polypdyr (Coelenterata). I. Ferskvandspolypper og goplepolypper. Danmarks Fauna. 41. Kobenhavn: GEC, Gad. 207 pp.
- Kramp, P. L. 1937 Polypdyr (Coelenterata). II. Goppler. Danmarks Fauna. 43. Kobenhavn: GEC, Gad. 233 pp.
- Kramp, P. L. 1939 Medusae, Siphonophora and Ctenophora. Zoology Iceland. 2(5b). Copenhagen, Reykjavik: Levin & Munksgaard. 37 pp.
- Morri, C. 1981 Idrozoi lagunari. Guide per il riconoscimento delle specie animali delle acque lagunari e costiere italiane. 6: Consiglio Nazionale delle Ricerche. 105 pp.
- Nobre, A. 1931 Contribuições para o estudo dos coelenterados de Portugal. Fauna marinha de Portugal. 3. Porto: Imprensa Portuguesa. 82 pp.

- Rossi, L. 1971 Guida a Cnidari e Ctenofori della fauna italiana. *Quaderni della Civica Stazione Idrobiologica di Milano* 2: 101 pp.
- Russell, F. S. 1953 The medusae of the British Isles. Anthomedusae, Leptomedusae, Limnomedusae, Trachymedusae and Narcomedusae. Cambridge: Cambridge University Press. 530 pp.
- Russell, F. S. 1970 The medusae of the British Isles. II. Pelagic Scyphozoa, with a supplement to the first volume on hydromedusae. Cambridge: Cambridge University Press. 238 pp.
- Stephenson, T. A. 1935 The British sea anemones. II. Cape Town: Ray Society. Pp. 96-111.
- ### *Anthozoa*
- Andres, A. 1884 *Le Attinie*. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 9. Leipzig: W. Engelmann. 459 pp.
- Broch, H. 1912 Die Alcyonarien des Trondhjemsfjordes I. Alcyonacea. *Det Kongelige Norske Videnskabers Selskabs Skrifter* 1911(7): 1-48.
- Broch, H. 1912 Die Alcyonarien des Trondhjemsfjordes II. Gorgonacea. *Det Kongelige Norske Videnskabers Selskabs Skrifter* 1912(2): 1-48.
- Broch, H. 1913 Die Alcyonarien des Trondhjemsfjordes III. Pennatulacea. *Det Kongelige Norske Videnskabers Selskabs Skrifter* 1912(16): 1-59.
- Carpine, C. & Grasshoff, M. 1975 Les gorgonaires de la Méditerranée. *Bulletin de l'Institut Océanographique de Monaco* 71: 1-140.
- Doumenc, D., Chintiroglou, C. & Koukouras, A. 1985 Actiniens de la Mer Egée: méthodes d'identification, zoogéographie. *Bulletin du Muséum national d'Histoire naturelle, Paris, 4ème série, Section A*, 7(3): 497-529.
- Grasshoff, M. 1992 Die Flachwasser-Gorgonarien von Europa und West Afrika (Cnidaria, Anthozoa). *Courier Forsch.-Inst Senckenberg* 149: 10-135.
- Koch von, G. 1887 *Die Gorgoniden des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 15. Berlin: R. Friedländer & Sohn. 99 pp.
- Kramp, P. L. 1939 *Octocorallia*. Zoology Iceland. 2(7). Copenhagen, Reykjavik: Levin & Munksgaard. 13 pp.
- Kükenthal, W. 1915 Pennatularia. *Das Tierreich* 43: i-xv, 1-132.
- Manuel, R. L. 1983 *The Anthozoa of the British Isles - a colour guide*. Ross-on-Wye: Underwater Conservation Society. 79 pp.
- Manuel, R. L. 1988 *British anthozoans (Coelenterata: Octocorallia and Hexacorallia). Keys and notes for the identification of the species. Synopses of the British Fauna (New Series)*, 18. Leiden: E.J. Brill. 241 pp.
- Pax, F. 1934 *Anthozoa*. Tierwelt der Nord- und Ostsee. 3e le2. Leipzig: Akademische verlagsgesellschaft. 317 pp.
- Pax, F. & Müller, I. 1962 *Fauna anthozoa Jadrana = Die Anthozoenfauna der Adria*. Fauna e flora adriatica = Fauna i flora jadrana. 3. Split: Institut za oceanografiju i ribarstvo. 343 pp.
- Weinberg, S. 1976 Revision of the common Octocorallia of the mediterranean circalittoral. I. Gorgonacea. *Beaufortia* 24(313): 63-103.
- Weinberg, S. 1977 Revision of the common Octocorallia of the mediterranean circalittoral. II. Alcyonacea. *Beaufortia* 25(326): 131-165.
- Weinberg, S. 1978 Revision of the common Octocorallia of the mediterranean circalittoral. III. Stolonifera. *Beaufortia* 27(338): 139-175.
- Weinberg, S. 1978 Octocorallarios mediterraneos. *Inmersión y Ciencia* 12: 37-47.
- Williams, G. C. 1995 Living genera of sea pens (Coelenterata: Pennatulacea): illustrated key and synopses. *Zoological Journal of the Linnean Society* 113: 93-140.
- Zibrowius, H. 1980 Les scléractiniaires de la Méditerranée et de l'Atlantique nord-oriental. *Mémoires de l'Institut Océanographique de Monaco* 11: 1-227.
- ### *Scyphozoa*
- Krumbach, T. 1930 *Scyphozoa*. Tierwelt der Nord- und Ostsee. 3d. Leipzig: Akademische verlagsgesellschaft. 88 pp.
- Naumov, D. V. 1961 *Scyphomedusae of the seas of the USSR*. Opredelitel po faune SSSR. 75. Leningrad: izdavaemye Zoolicheskim institutom Akademii Nauk SSSR. 98 pp.
- ### *Hydrozoa*
- Brinckmann-Voss, A. 1970 *Anthomedusae / Athecatae (Hydrozoa, Cnidaria) of the Mediterranean. Part I Capitata*. Fauna und flora del Golfo di Napoli. 39. Napoli: Stazione Zoologica di Napoli. 96 pp.
- Broch, H. 1928 *Hydrozoa*. Tierwelt der Nord- und Ostsee. 3b,c. Leipzig: Akademische verlagsgesellschaft. 100 pp.
- Cornelius, P. F. S. 1975 A revision of the species of Lafoeidae and Haleciidae (Coelenterata: Hydroida) recorded from Britain and nearby seas. *Bulletin of the British Museum (Natural History), Zoological series* 28: 373-426.
- Cornelius, P. F. S. 1995 *North-west European thecate hydroids and their medusae. Part 1 Introduction, Laodiceidae to Haleciidae*. Synopses of the British fauna (New Series). 50. Field Studies Council. 357 pp.
- Cornelius, P. F. S. 1995 *North-west European thecate hydroids and their medusae. Part 2 Sertulariidae to Campanulariidae*. Synopses of the British fauna (New Series). 50. Field Studies Council. 386 pp.
- García Corrales, P., Aguirre Incharibe, A. & Gonzalez Moya, D. 1978 Contribución al conocimiento de los hidrozoos de las costas españolas. Parte 1: Halecidos, Campanularídos y Plumularíados. *Boletín del Instituto Español de Oceanografía* 253: 5-73.
- García Corrales, P., Aguirre Incharibe, A. & Gonzalez Moya, D. 1979 Contribución al conocimiento de los hidrozoos de las costas españolas. Parte 2: Lafoeidae, Campanulinidae y Syntheciidae. *Boletín del Instituto Español de Oceanografía* 273: 5-39.
- García Corrales, P., Aguirre Incharibe, A. & Gonzalez Moya, D. 1980 Contribución al conocimiento de los hidrozoos de las costas españolas. Parte 3: Sertulariidae. *Boletín del Instituto Español de Oceanografía* 296: 5-67.
- Kirkpatrick, P. A. & Pugh, P. R. 1984 *Siphonophores and velellids. Keys and notes for the identification of the species*. Synopses of the British Fauna (New Series). 29. Leiden: Backhuys Publishers. vii+154 pp.
- Naumov, D. V. 1960 *Hydroids and hydromedusae of marine, brackish, and freshwater basins of the USSR*. Opredelitel po faune SSSR. 70. Leningrad: izdavaemye Zoolicheskim institutom Akademii Nauk SSSR. 627 pp.
- Ramil, F. & Vervoort, W. 1992 Report on the Hydroids collected by the "BALGIM" expedition in and around the straits of Gibraltar. *Zoologische Verhandelingen* 277: 3-262.
- Rees, W. J. R., M. 1941 Hydroids of the Swedish west coast. *Acta Regiae Societatis Scientiarum et Litterarum Gothoburgensis Zoologica* 3: 1-24.
- Stepanyants, S. D. 1967 *Siphonophora of the seas of the USSR and northern part of the Pacific Ocean*. Opredelitel po faune SSSR. 96. Leningrad: izdavaemye Zoolicheskim institutom Akademii Nauk SSSR. 216 pp.
- Svoboda, A. & Cornelius, P. F. S. 1991 The European and Mediterranean species of Aglaophenia (Cnidaria, Hydrozoa). *Zoologische verhandelingen* (274): 1-72.
- Totton, A. K. 1965 *A synopsis of the Siphonophora*. London: British Museum (Natural History). 230 pp.
- Vervoort, W. 1946 *Hydrozoa (CJ), A. Hydropolypen*. Fauna van Nederland. 14. Leiden: A. W. Sijthoff. 336 pp.
- Zibrowius, H. W. & Cairns, S. D. 1992 Revision of the north east Atlantic and Mediterranean Styelasteridae (Cnidaria: Hydrozoa). *Mémoires du Muséum national d'Histoire naturelle, Paris. Série A, Zoologie* 153: 1-136.
- ### *CTENOPHORA*
- Arndt, W., Broch, H., Krumbach, T., Pax, F. & Leiberkind, J. 1928 Porifera, Schwämme, Spongiens. Hydrozoen.

- Scyphozoa. Anthozoa. Ctenophora, Kammquallen oder Rippenquallen. Echinodermata, Stachelhäuter oder Echinodermen. Die Tierwelt Deutschlands. 4. Jena: G. Fischer. 332 pp.
- Chun, C. 1880 Die Ctenophoren des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 1. Leipzig: W. Engelmann. 313 pp.
- Greve, W. 1975 Ctenophora. Fiches d'Identification du Zooplanton. 146. 6 pp.
- Krumbach, T. 1926 Ctenophora. Tierwelt der Nord- und Ostsee. 3f. Leipzig: Akademische verlagsgesellschaft. 50 pp.
- Rossi, L. 1971 Guida a Cnidari e Ctenofori della fauna italiana. Quaderni della Civica Stazione Idrobiologica di Milano 2: 1-101.

PLATYHELMINTHES

- Ax, P. 1956 Les Turbellariés des étangs côtiers du littoral méditerranéen de la France méridionale. *Vie et Milieu Suppl.* 5: 1-152.
- Ax, P. 1956 Monographie der Otoplanidae (Turbellaria), Morphologie und Systematik. *Abhandlungen der mathematisch-naturwissenschaftlichen Klasse* 13: 499-796.
- Ax, P. 1971 Zur Systematik und Phylogenie der Trigonostominae (Turbellaria, Neorhabdocoela). *Mikrofauna Meeresboden* 4: 1-84.
- Ball, I. R. & Reynoldson, T. B. 1981 *British Planarians, Platyhelminthes, Tricladida. Synopses of the British Fauna (New Series)*. 19. Cambridge, New-York: Cambridge University Press. 141 pp.
- Brinkmann, A. 1957 *Fish trematodes from Norwegian waters IIa. The Norwegian species of the orders Aspidogastrea and Digenea (Gasterostomata)*. Naturvitenskapelig rekke. 4: Universitetet i Bergen Årbok. 29 pp.
- Cannon, L. R. G. 1986 *Turbellaria of the world*. Brisbane: Queensland Museum. viii+136 pp.
- Chiriac, E. & Udrescu, M. 1973 *Trematoda*. Fauna Republicii socialiste România. 2(4). Bucuresti: Academiei Republicii socialiste România. 496 pp.
- Dawes, B. 1947 *The Trematoda of British Fishes*. London: Ray Society. 364 pp.
- Dorjcs, J. 1968 Die Acoela (Turbellaria) der deutschen Nordseeküste und ein neues System der Ordnung. *Zeitschrift für Zoologische Systematik und Evolutionforschung* 6: 56-452.
- Ehlers, U. 1974 Interstitial Typhloplanoida (Turbellaria) aus dem Litoral der Nordseeinsel Sylt. *Mikrofauna Meeresboden* 49: 425-527.
- Evdonin, L. A. 1977 *Kalyptorhynchia in the fauna of the USSR and adjoining countries. Turbellaria, I (1)*. Fauna SSSR (novaia seriia). 115. Leningrad. 400 pp.
- Faibel, A. 1974 Macrostomida (Turbellaria) von einem Sandstrand der Nordseeinsel Sylt. *Mikrofauna Meeresboden* 45: 339-370.
- Faibel, A. 1976 Interstitial Acoela (Turbellaria) aus dem Litoral der Nordfriesischen Inseln Sylt und Amrum (Nordsee). *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 73: 17-56.
- Faibel, A. 1983 The Polycladida, Turbellaria. Proposal and establishment of a new system. Part. I. The Acotylea. *Mitt. hamb. zool. Mus. Inst.* 80: 17-121.
- Faibel, A. 1984 The Polycladida, Turbellaria. Proposal and establishment of a new system. Part II. The Cotylea. *Mitt. hamb. zool. Mus. Inst.* 81: 189-259.
- Joyeux, C. & Baer, J. G. 1936 *Cestodes*. Faune de France. 30. Paris: Paul Lechevalier. 613 pp.
- Karling, T. G. 1958 Zur Kenntnis der Gattung *Coelogynopora Steinböck* (Turbellaria, Proseriata). *Ark. Zool.* 11: 559-567.
- Karling, T. G. 1974 Turbellarian fauna of the Baltic Proper. Identification, ecology and biogeography. *Fauna Fennica* 27: 1-101.
- Khalil, L. F., Jones, A. & Bray, R. A. 1994 *Keys to the cestodes parasites of vertebrates*. Wallingford: CAB International. xiii+751 pp.

- Khotenovskii, I. A. 1985 *Suborder Octomacrinae Khotenovskij. Monogeneans*. Fauna SSSR (novaia seriia). 132. Leningrad: Akademija nauk SSSR, Zoologicheskij institut. 263 pp.
- Lang, A. 1884 *Die Polycladen (seeplanarien) des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 11. Leipzig: W. Engelmann. 688 pp.
- Meixner, J. s.d. *Turbellaria (Strudelwürmer)*. Tierwelt der Nord- und Ostsee. 4b. Leipzig: Akademische verlagsgesellschaft. 146 pp.
- Norena-Janssen, C. & Faibel, A. 1992 Revision of the subfamily Mesostominae, Rhabdocoela, Platyhelminthes. *Mitt. hamb. zool. Mus. Inst.* 89: 7-47.
- Prudhoe, S. 1982 *British Polyclad Turbellarians. Synopses of the British Fauna (New Series)*. 26. Cambridge, New-York: Cambridge University Press. 77 pp.
- Prudhoe, S. 1985 *A monograph on polyclad Turbellaria*. Oxford: Oxford University Press. 259 pp.
- Roman-Chiriac, E. 1960 *Clasa Monogenoidea*. Fauna Republicii Populară Române. 2(1). Bucuresti: Academiei Republicii Populară Române. 151 pp.
- Schilke, K. 1970 *Kalyptorhynchia (Turbellaria) aus dem Eulitoral der deutschen Nordseeküste. Helgoländer wissenschaftliche Meeresuntersuchungen* 21: 143-265.
- Schmidt, G. D. 1985 *Handbook of tapeworm identification*. Boca Raton: CRC Press. 675 pp.
- Sluys, R. 1989 *A Monograph of the marine tricladids*. Rotterdam: Balkema. xii+463 pp.
- Sopott, B. 1972 Systematik und Ökologie von Proseriaten (Turbellaria) der deutschen Nordseeküste. *Mikrofauna Meeresboden* 13: 1-72.
- Sprehn, C. 1933 *Trematoda*. Tierwelt der Nord- und Ostsee. 4cl. Leipzig: Akademische verlagsgesellschaft. 60 pp.
- Sprehn, C. E. W. 1934 *Cestoidea*. Tierwelt der Nord- und Ostsee. 4c2. Leipzig: Akademische verlagsgesellschaft. Pp. 61-91.
- Wilhelmi, J. 1909 *Tricladida*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 32. Berlin: R. Friedländer & Sohn. 405 pp.

ROTIFERA

- Berzins, B. 1952 Contributions to the knowledge of the marine Rotatoria of Norway. *Universitetet i Bergen. Årbok. Naturvitenskapelig rekke* 1951(6): 1-11.
- Braioni, M. G. & Gelmini, D. 1983 *Rotiferi Monogononti (Rotatoria: Monogononta)*: Consiglio Nazionale delle Ricerche. 180 pp.
- Eriksen, G. G. 1968 Marine rotifers found in Norway, with description of two new species and one little known species. *Sarsia* 33: 23-34.
- Kutikova, L. A. 1970 *Rotifers of the USSR fauna (Rotatoria. Subclass Eurotatoria)*. Opredelitel po faune SSSR. 104. Leningrad: izdavaemye Zoologicheskij institutom Akademii Nauk SSSR. 744 pp.
- Remane, A. 1929 *Rotatoria*. Tierwelt der Nord- und Ostsee. 7e. 156 pp.
- Remane, A. 1949 Die psammobionten Rotatorien der Nord- und Ostsee. *Kieler Meeresf.* 6: 59-67.
- Rudescu, L. 1960 *Rotatoria, Trochelminthes*, fasc. 2. Fauna Republicii Populară Române. 2. s.l.: Editura Academiei Republicii Populară Române. 1192 pp.
- Thane-Fenchel, A. 1968 A simple key to the genera of marine and brackish-water Rotifers. *Ophelia* 5: 299-311.

NEMERTINI

- Brunberg, L. 1964 On the nemertean fauna of Danish waters. *Ophelia* 1: 77-111.
- Bürger, O. 1895 *Die Nemertinen des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 22. Berlin: R. Friedländer & Sohn. 743 pp.
- Friedrich, H. 1936 *Nemertini*. Tierwelt der Nord- und Ostsee. 4d. Leipzig: Akademische verlagsgesellschaft. 69 pp.

- Gibson, R. 1994 *British nemerteans. Synopses of the British Fauna (New Series)*. 24. Cambridge, New-York: Field Studies Council. vii+224 pp.
- Norenburg, J. 1988 Remarks on marine interstitial nemertines and key to the species. *Hydrobiologia* 156: 87-92.

GASTROTRICHA

- Balsamo, M. 1983 *Gastrotrichi (Gastrotricha)*. Consiglio Nazionale delle Ricerche. 92 pp.
- d'Hondt, J.-L. 1974 Clés tabulaires de détermination des genres marins de Gastrotriches. *Bulletin de la Société Zoologique de France* 99(4): 645-665.
- Remane, A. 1927 *Gastrotricha*. Tierwelt der Nord- und Ostsee. 7d1. Leipzig: Akademische verlagsgesellschaft. 56 pp.
- Rudescu, L. 1967 *Gastrotricha*. Fauna Republicii Socialiste România. 2(3). Bucuresti: Academia Republicii Socialiste România. 295 pp.
- Voigt, M. *Gastrotricha*. Die Tierwelt Mitteleuropas. I (4a). Leipzig: Quelle & Meyer. 74 pp.

CEPHALORHYNCHA

Kinorhyncha

- Higgins, R. P. 1985 The genus *Echinoderes* (*Kinorhyncha: Cyclorrhagida*) from the English Channel. *Journal of the Marine Biological Association of the United Kingdom* 65: 785-800.
- Remane, A. 1928 *Kinorhyncha*. Tierwelt der Nord- und Ostsee. 7d2. Leipzig: Akademische verlagsgesellschaft. Pp. 57-84.

Priapulida (see under *Sipunculida*)

Nematomorpha

- Heinze, K. 1941 *Würmer oder Vermes. III. Saitenwürmer oder Gordioidea (Nematomorpha)*. Die Tierwelt Deutschlands. 39. Jena: G. Fischer. 78 pp.
- Stekhoven, J. H. S. 1934 *Nematomorpha*. Tierwelt der Nord- und Ostsee. 6f. Leipzig: Akademische verlagsgesellschaft. 10 pp.

NEMATODA

- Platt, H. M. & Warwick, R. M. 1983 *Free-living marine nematodes. Part I: British enoplids. Pictorial key to world genera and notes for the identification of British species*. *Synopses of the British Fauna (New Series)*. 28. London, New-York: Cambridge University Press. vii+307 pp.
- Platt, H. M. & Warwick, R. M. 1988 *Free-living marine nematodes. Part II: British Chromadorids*. *Synopses of the British Fauna (New Series)*. 38. Cambridge, New York: Cambridge University Press. 522 pp.
- Schuurmans Stekhoven, J. H. 1935 *Nematoda parasitica*. Tierwelt der Nord- und Ostsee. 5c. Leipzig: Akademische verlagsgesellschaft. 50 pp.
- Schuurmans Stekhoven, J. H. 1935 *Nematoda errantia*. Tierwelt der Nord- und Ostsee. 5b. Leipzig: Akademische verlagsgesellschaft. 173 pp.
- Schuurmans Stekhoven, J. H. (not dated). *Nematoda parasitica. Ergänzungen und Berichtigungen*. Tierwelt der Nord- und Ostsee. 5c1. Leipzig: Akademische verlagsgesellschaft. Pp. 52-59.
- Tarjan, A. C. 1980 *An illustrated guide to the marine nematodes*. Institute of food and agricultural sciences. University of Florida. 198 pp.
- Warwick, R. M., Platt, H. M. & Somerfield, P. J. 1998 *Free-living marine nematodes: Part III monhysterids*. *Synopses of the British Fauna (New Series)*. 53. Leiden: Backhuys Publishers. 310 pp.

ACANTHOCEPHALA

- Amin, O. M. 1987 Key to the families and subfamilies of Acanthocephala, with the erection of a new class (Polyacanthocephala) and a new order

- (Polyacanthocephala). *Journal of Parasitology* 73(6): 1216-1219.

- Wulker, G. & Schuurmans Stekhoven, J. H. 1933 *Acanthocephala*. Tierwelt der Nord- und Ostsee. 6e. Leipzig: Akademische verlagsgesellschaft. 64 pp.

ECHIURA (see under *sipunculida*)

SIPUNCULIDA, ECHIURA and PRIAPULIDA

- Baltzer, F. 1917 *Monographie der Echiuriden des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 34. Berlin: R. Friedländer & Sohn. 234 pp.

- Cuénod, L. 1922 *Sipunculians, Echiuriens, Priapuliens*. Faune de France. 4. Paris: P. Lechevalier. 29 pp.

- Fischer, W. *Echiuridae, Sipunculidae, Priapulidae*. Tierwelt der Nord- und Ostsee. 6d. Leipzig: Akademische verlagsgesellschaft. 55 pp.

- Gibbs, P. E. 1977 *British sipunculans. Keys and notes for the identification of the species*. *Synopses of the British Fauna (New Series)*. 12. London: Academic Press. 35 pp.

- Murina, G. V. 1977 *Sipunculids of the Arctic and boreal waters of Eurasia*. Opredeliteli po faune SSSR. 111. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 284 pp.

- Murina, G. V. & Zadovnik, D. 1985 Sipuncula of the Adriatic Sea. *Thalassia Jugoslavica* 21/22(1/2): 23-73.

- Saiz Salinas, J. I. 1993 *Sipuncula. Fauna Iberica*. 4. Madrid: Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Científicas. 200 pp.

- Sáiz Salinas, J. I. 1986 *Los gusanos sipunculidos de los fondos litorales y circalitorales de las costas de la Península Ibérica, Islas Baleares, Canarias y mares adyacentes*. Monografías del Instituto Español de Oceanografía: Instituto Español de Oceanografía. 84 pp.

- Stephen, A. C. 1960 *British echiurids (Echiuroidea), sipunculids (Sipunculoidea) and priapulids (Priapuloidea)*. With keys and notes for the identification of the species. *Synopses of the British Fauna*. 12. London. 27 pp.

- Stephen, A. C. & S. J. Edmonds 1972 *The phyla Sipuncula and Echiura*. London: British Museum (Natural History). 528 pp.

- Ude, H., Johansson, L. & Ten Broecke, A. 1929 *Würmer oder Vermes. Oligochaeta. Hirudinea (Hegel)*. *Sipunculoidea und Echiuroidea*. Die Tierwelt Deutschlands. 15. Jena: G. Fischer. 168 pp.

- Wesenberg-Lund, E. 1939 *Pølseorme (Gephyrea). Sipunculider, Priapulider og Echiurider*. Danmarks Fauna. 45. Kobenhaven: GEC Gad. 60 pp.

ENTOPOCTA

- Cori, C. J. 1930 *Kamptozoa (Entoprocta)*. Tierwelt der Nord- und Ostsee. 4a. Leipzig: Akademische verlagsgesellschaft. 68 pp.

- Nielsen, C. 1989 *Entoprocts. Keys and notes for the identification of the species*. *Synopses of the British Fauna (New Series)*. 41. Leiden, New-York: E.J. Brill. 131 pp.

MOLLUSCA

General

- Barash, A. & Danin, Z. 1992 *Fauna Palaestina. Mollusca I. Annotated list of Mediterranean Molluscs of Israel and Sinai*. Fauna Palaestina. Jerusalem: The Israel Academy of Sciences and Humanities. 405 pp.

- Beedham, G. E. 1972 *Identification of the British Mollusca*. Hulton Group Keys. Amersham: Hulton Educational Publications Ltd. 239 pp.

- Boer de, T. W. & Bruyne de, R. H. 1991 *Schelpen van de Friese Waddeneilanden: overzicht van alle mariene autochtone weekdieren (Mollusca) en aangespoele schelpen*. Fryske akademy. Ljouwert: W. Backhuys. 292 pp.
- Bruyne de, R. H. 1991 *Schelpen van de Nederlandse kust*. Utrecht: Stichting Uitgeverij KNNV. 165 pp.
- Cossignani, T., Cossignani, V., Di Nisio, A. & Passamonti, M. 1992 *Atlante delle conchiglie del medio Adriatico. Atlas of shells from the central Adriatic sea*. Ancona: L'Informatore Piceno. 118 pp.
- Delamotte, M. 1994 *Shells from the Greek seas*. Annales Musei Goulandris. Greece: Goulandris NHM. 313 pp.
- Demetropoulos, A. 1969 Marine molluscs of Cyprus. *Fisheries Bulletin* 2-3.
- Hidalgo, J. G. 1917 *Fauna Malacológica de España, Portugal y las Baleares. Moluscos testaceos marinos*. Trabajos del Museo Nacional de Ciencias Naturales, Serie Zoológica. 30. Madrid. 752 pp.
- Muus, B. J. 1959 *Skallus, Søtander, Blæsprutter*. Danmarks Fauna. 65. København: G.E.C. Gads Forlag. 239 pp.
- Nobre, A. 1931 *Moluscos marinhos de Portugal*. Fauna Marinha de Portugal. 5. Pôrto: Instituto de Zoologia da Universidade do Pôrto. 466 pp.
- Nobre, A. 1936 *Moluscos marinhos de Portugal*. Vol. 2. Fauna Marinha de Portugal. 5. Barcelos: Companhia Editoria do Minho. 378 pp.
- Poppe, G. T. & Goto, Y. 1991 *European Seashells. Vol. I Polyplacophora, Caudofoveata, Soleogastera, Gastropoda*. Hemmen: Verlag Christa. 352 pp.
- Poppe, G. T. & Goto, Y. 1993 *European Seashells. Vol. II Scaphopoda, Bivalvia, Cephalopoda*. Hemmen: Verlag Christa. 221 pp.
- Rolan, M., Otero Schmitt, J. & Rolan Alvarez, E. 1990 *Moluscos de la ría de Vigo 2. Poliplacoforos, Bivalvos, Escafopodos, Cefalopodos*. 2. Santiago de Compostela: Servicio de publicaciones e intercambio científico. 276 pp.
- Starobogatov, Y. I. & Naumov, A. D. (ed.) 1987 *Molluscs of the White Sea*. Opredelitel po faune SSSR. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR.
- Tornaritis, G. 1987 *Mediterranean sea shells - Cyprus*. Nicosia: G. Tornaritis. 190 pp.
- Warén, A. 1989 New and little known Mollusca from Iceland. *Sarsia* 74: 1-28.
- Warén, A. 1991 New and little known Mollusca from Iceland and Scandinavia. *Sarsia* 76: 53-124.
- Warén, A. 1993 New and little known Mollusca from Iceland and Scandinavia. Part 2. *Sarsia* 78: 159-201.
- Warén, A. 1996 New and little known Mollusca from Iceland and Scandinavia. Part 3. *Sarsia* 81: 197-246.
- Zhadin, V. I. 1952 *Molluscs of fresh and brackish waters of the USSR*. Opredelitel po faune SSSR. 46. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 376 pp.
- Polyplacophora, Scaphopoda, Aplacophora**
- Belle van, R. A. 1983 *De Europese Polyplacophora*. Antwerpen: Belgische Vereinigung voor Conchyliologie. 251 pp.
- Benthem Jutting van, T. 1926 *Scaphopoda*. Tierwelt der Nord- und Ostsee. 9c2. Leipzig: Akademische verlagsgesellschaft. Pp. 67-80.
- Jones, A. M. & Baxter, J. M. 1987 *Molluscs: Caudofoveata, Soleogastera, Polyplacophora and Scaphopoda*. Synopses of the British Fauna (New Series). 37. London: E.J. Brill/Dr.W. Backhuys. 123 pp.
- Nierstrasz, H. F. & Hoffmann, H. 1929 *Mollusca Aculifera Aplacophora und Aculifera Placophora*. Tierwelt der Nord- und Ostsee. 9a. Leipzig: Akademische verlagsgesellschaft. 64 pp.
- Salvini-Plawen van, L. 1975 *Mollusca Caudofoveata*. Marine invertebrates of Scandinavia. 4. Oslo: Scandinavian University Books. 55 pp.
- Steiner, G. 1997 Escalópodos de las costas españolas. *Iberus* 15(1): 95-111.
- Yakovleva, A. M. 1952 *Shell-Bearing Mollusks (Loricata) of the Seas of the U.S.S.R.* Opredelitel po faune SSSR. 45. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 107 pp.
- Gastropoda**
- Ankel, W. E. 1926 *Prosobranchia*. Tierwelt der Nord- und Ostsee. 9b1. Akademische verlagsgesellschaft. 240 pp.
- Barletta, G. 1980 *Gasteropodi nudi (Pleurobranchomorpha, Sacoglossa, Aplysiomorpha e Nudibranchia)*. Guida per il riconoscimento delle specie animale delle acque lagunari e costiere italiane. 3. 128 pp.
- Benthem Jutting van, T. 1933 *Mollusca (I) A Gastropoda Prosobranchia et Pulmonata*. Fauna van Nederland. 7. Leiden: A.W. Sijthoff. 387 pp.
- Benthem Jutting van, T. & Engel, H. 1936 *Mollusca (I) B Gastropoda Opisthobranchia; Amphineura et Scaphopoda*. Fauna van Nederland. 8. Leiden: A. W. Sijthoff. 106 pp.
- Bobdanov, L. P. 1990 *Molluski podsemeistva Oenopotaiae (Gastropoda, Pectinibranchia, Turridae) morei SSSR*. Fauna SSSR (novaia seria). 142. Leningrad: Akademii nauk SSSR, Zoologicheskii institut. 221 pp.
- Bondesen, P. 1975 Danske havsnegle. *Natur og Museum* 16(3-4): 1-31.
- Bouchet, P. & Warén, A. 1980 Revision of the north-east Atlantic bathyal and abyssal Turridae (Mollusca, Gastropoda). *Journal of Molluscan Studies Suppl.* 8: 1-120.
- Bouchet, P. & Warén, A. 1985 Revision of the north-east Atlantic bathyal and abyssal Neogastropoda excluding Turridae (Mollusca, Gastropoda). *Bulletino Malacologico Suppl.* 1: 123-296.
- Bouchet, P. & Warén, A. 1986 Revision of the north-east Atlantic bathyal and abyssal Aclididae, Eulimidae, Epitonidae (Mollusca, Gastropoda). *Bulletino Malacologico Suppl.* 2: 300-576.
- Bouchet, P. & Warén, A. 1993 Revision of the north-east Atlantic bathyal and abyssal Mesogastropoda. *Bulletino Malacologico Suppl.* 3: 577-840.
- Brown, G. H. & Picton, B. E. 1985 *Nudibranchs of the British Isles - a colour guide*. Nottingham: R. Earll.
- Cachia, C., Mifsud, C. & Sammut, P. M. 1991 *The Marine shelled Mollusca of the Maltese Islands, Part 1: Archaeogastropoda*. Marsa, Malta: Grima Print. & Publ. Ind. xii+113 pp.
- Cachia, C., Mifsud, C. & Sammut, P. M. 1996 *The marine Mollusca of the Maltese islands, Part 2: Neotaenioglossa*. Leiden: Backhuys. 228 pp.
- Cattaneo-Vietti, R., Chemello, R. & Giannuzzi-Savelli, R. (ed.) 1990 *Atlas of Mediterranean nudibranchs. Atlante dei Nudibranchi del Mediterraneo*. Roma: La Conchiglia. 264 pp.
- Dales, R. P. 1957 *Heteropoda*. Families: Atlantidae, Carinariidae and Pterotracheidae. *Fiches d'identification du plancton* 66: 1-4.
- Fretter, V. & Graham, A. 1976 The prosobranch molluscs of Britain and Denmark. Part 1: Pleurotomariacea, Fissurellacea and Patellacea. *Journal of Molluscan Studies Suppl.* 1: 1-37.
- Fretter, V. & Graham, A. 1977 The prosobranch molluscs of Britain and Denmark. Part 2: Trochacea. *Journal of Molluscan Studies Suppl.* 3: 39-100.
- Fretter, V. & Graham, A. 1978 The prosobranch molluscs of Britain and Denmark. Part 4: Marine Rissacea. *Journal of Molluscan Studies Suppl.* 6: 153-241.
- Fretter, V. & Graham, A. 1978 The prosobranch molluscs of Britain and Denmark. Part 3: Neritacea, Viviparacea, Valvatacea, terrestrial and freshwater Littorinacea and Rissacea. *Journal of Molluscan Studies Suppl.* 5: 101-153.
- Fretter, V. & Graham, A. 1980 The prosobranch molluscs of Britain and Denmark. Part 5: Marine Littorinacea. *Journal of Molluscan Studies Suppl.* 7: 243-284.
- Fretter, V. & Graham, A. 1981 The prosobranch molluscs of Britain and Denmark. Part 6: Cerithiacea, Strombacea, Hippionicacea, Calyptraeacea, Lamellariacea, Cypraeacea,

- Naticacea, Tonnacea, Heteropoda. *Journal of Molluscan Studies Suppl.* 9: 285-363.
- Fretter, V. & Graham, A. 1982 The prosobranch molluscs of Britain and Denmark. Part 7: 'Heterogastropoda' (Cerithiopsacea, Triforacea, Epitonacea, Eulimacea). *Journal of Molluscan Studies Suppl.* 11: 363-434.
- Fretter, V. & Graham, A. 1985 The prosobranch molluscs of Britain and Denmark. Part 8: Neogastropoda. *Journal of Molluscan Studies Suppl.* 15: 435-556.
- Fretter, V., Graham, A. & Andrews, E. B. 1986 The prosobranch molluscs of Britain and Denmark. Part 9: Pyramidellacea. *Journal of Molluscan Studies Suppl.* 16: 557-649.
- Galkin, Y. I. 1955 *Gastropod molluscs (am. Trochidae) of Far Eastern and northern seas of the USSR*. Opredeliteli po faune SSSR. 57. Zoologicheski Institut, Akademii Nauk SSSR. 132 pp.
- Giannuzzi-Savelli, R., Pusateri, F., Palmeri, A. & Ebreo, C. 1994 *Atlante delle conchiglie marine del Mediterraneo. Atlas of the Mediterranean sea shells*. 1. Roma: La Conchiglia. 125 pp.
- Giannuzzi-Savelli, R., Pusateri, F., Palmeri, A. & Ebreo, C. 1996 *Atlante delle conchiglie marine del Mediterraneo. Atlas of the Mediterranean sea shells*. 2. Roma: La Conchiglia. 258 pp.
- Giannuzzi-Savelli, R., Pusateri, F., Palmeri, A. & Ebreo, C. 1999 *Atlante delle conchiglie marine del Mediterraneo. Atlas of the Mediterranean sea shells*. 3. Roma: Evolver. 126 pp.
- Giusti, F. & Pezzoli, E. 1980 *Gasteropodi 2 (Gastropoda: Prosobranchia: Hydrobioidea, Pyrguloidea)*. Consiglio Nazionale delle Ricerche. 67 pp.
- Golikov, A. N. 1963 *Briukhonogie molliuski roda Neptunea Bolten*. Fauna SSSR (novaia serii). 85. Leningrad: Akademii nauk SSSR, Zoolicheskii institut. 217 pp.
- Golikov, A. N. & Kusakin, O. G. 1978 *Gastropod molluscs of the intertidal zone of the USSR*. Opredeliteli po faune SSSR. 116. Zoologicheski Institut, Akademii Nauk SSSR. 256 pp.
- Graham, A. 1988 *Molluscs: prosobranch and pyramidellid gastropods. Keys and notes for the identification of the species. Synopses of the British Fauna (New Series)*. 2. Leiden, New-York: E. J. Brill / Dr W. Backhuys. viii+662 pp.
- Grossu, A. V. 1956 *Gastropoda Prosobranchia si Opisthobranchia*. Fauna Republicii Populare Romîne. 3(2). Bucuresti: Editura Academiei Republicii populare Romîne. 220 pp.
- Hoffmann, H. 1926 I. *Opisthobranchia. II. Pteropoda*. Tierwelt der Nord- und Ostsee. 9cl. Leipzig: Akademische verlagsgesellschaft. 66 pp.
- Hubendick, B. & Warén, A. 1976 *Framgående Snäckor från Svenska westkusten*. Göteborg: Rundqvists Boktryckeri. 73 pp.
- Just, H. & Edmunds, M. 1985 North Atlantic nudibranchs (Mollusca) seen by Henning Lemche. *Ophelia suppl.* 2: 1-170.
- Llera, E. M., Ortea, J. A. & Vizcaino, A. 1983 *Moluscos. I. Archaeogastropoda (Prosobranchia)*. Fauna Marina de Asturias. 1. Gijon. 75 pp.
- Macan, T. T. 1977 *A key to the British fresh- and brackish-water gastropods*. Ambleside: Freshwater Biological Association. 46 pp.
- Morton, J. E. 1957 Opisthobranchia. Order: Gymnosomata, family: Clionidae. *Fiches d'identification du plancton* 80: 1-4.
- Morton, J. E. 1957 Opisthobranchia. Order: Gymnosomata. *Fiches d'identification du plancton* 79: 1-4.
- Nordsieck, F. 1968 *Die europäischen Meeres-Gehäuseschnecken (Prosobranchia) Vom Eismeer bis Kapverden und Mittelmeer*. Stuttgart: Gustav Fischer Verlag. 273 pp.
- Nordsieck, F. 1972 *Die europäischen Meeres-Gehäuseschnecken (Opisthobranchia mit Pyramellidae)*; *Rissoacea) Vom Eismeer bis Kapverden, Mittelmeer und Schwarzes Meer*. Stuttgart: Gustav Fischer Verlag. 327 pp.
- Nordsieck, F. 1977 *The Turridae of the European seas*. Roma: La Piramide / La Conchiglia. 131 pp.
- Nordsieck, F. & Garcia-Talavera, F. 1979 *Moluscos marinos de Canarias y Madeira (Gastropoda)*. Madrid: Selecciones Graficas. 208 pp.
- Ortea, J., Valdés, A. & Garcia-Gomez, J. C. 1996 Revision de las especies atlanticas de la familia Chromodorididae (Mollusca: Nudibranchia) del grupo cromatico azul. *Avicennia suppl.* 1: 1-165.
- Peñas, A., Templado, J. & Martinez, J. L. 1996 Contribucion al conocimiento de los Pyramidelloidea (Gastropoda: Heterostropha) del Mediterraneo español. *Iberus* 14(1): 1-82.
- Picton, B. E. & Morrow, C. C. 1994 *A field guide to the nudibranchs of the British Isles*. London: Immel publishing. 143 pp.
- Pruvot-Fol, A. 1954 *Mollusques opisthobranches*. Faune de France. 58. Paris: P. Lechevalier. 460 pp.
- Rolan, E. 1983 *Moluscos de la ría de Vigo I. Gasteropodos*. 1. Santiago de Compostela: Servicio de publicaciones e intercambio científico. 383 pp.
- Rolan, E. 1987 *Moluscos gasteropodos de Galicia*. Santiago de Compostela: Universidad de Santiago de Compostela. 105 pp.
- Schmekel, L. & Portmann, A. 1982 *Opisthobranchia des Mittelmeeres. Nudibranchia und Sacoglossa*. Fauna e flora del golfo di Napoli. 40. Berlin, New-York: Springer-Verlag. x+410 pp.
- Smith, S. M. 1974 *Key to the British Gastropoda*. Royal Scottish Museum Information Series. Natural History. 2. Edinburgh: Royal Scottish Museum. viii+44 pp.
- Sneli, J.-A. 1975 *Mollusca Prosobranchia, Forgiellesnegler, nordiske marine arter*. Oslo: Scandinavian University Books. Universitetsforlaget. 55 pp.
- Thompson, T. E. 1988 *Molluscs: benthic opisthobranchs (Mollusca: Gastropoda)*. Synopses of the British Fauna (New Series). 8. London: Linnean Society of London/Estuarine & Brackish Water Sciences Association. 356 pp.
- Torelli, A. 1982 *Gasteropodi conchigliati*. Guida per il riconoscimento delle specie animali delle acque lagunari e costiere italiane. 8. Roma. 235 pp.
- Warén, A. 1992 New and little known "Skeneimorph" gastropods from the Mediterranean sea and the adjacent Atlantic ocean. *Bollettino Malacologico* 27(10-12): 149-248.
- Wirtz, P. 1998 Opisthobranch molluscs from the Azores. *Vita Marina* 45(1-2): 1-16.
- Wirtz, P. 1999 Opisthobranch molluscs from the Madeira archipelago. *Vita Marina* 46(1-2): 1-18.
- ### Cephalopoda
- Clarke, M. R. 1986 *Handbook for the identification of cephalopod beaks*. Oxford: Oxford University Press. 273 pp.
- Guerra, S. A. 1992 *Mollusca, Cephalopoda. Fauna Iberica*. 1. Madrid: Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Cientificas. 327 pp.
- Jaeckel, S. G. A. 1958 *Cephalopoden*. Tierwelt der Nord- und Ostsee. 9b3. Leipzig: Akademische verlagsgesellschaft. Pp. 479-723.
- Jatta, G. 1896 *I Cefalopodi viventi nel Golfo di Napoli (sistematica)*. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 23. Berlin: R. Friedländer & Sohn. 268 pp.
- Lacourt, A. W. & Huwae, P. H. M. 1981 *De Inktvissen (Cephalopoda) van de Nederlandse kust*. Wetenschappelijke mededelingen KNIN. 145. Hoogwoud: Koninklijke Nederlandse Natuurhistorische Vereniging. 32 pp.
- Naeff, A. 1972 *Cephalopoda. Part I. Vol. I. Fasc. I. Fauna and flora of the Bay of Naples*. 35. Jerusalem: Israel Program for Scientific Translations. 292 pp.

- Naef, A. 1972 *Cephalopoda (systematics). Part 1. Vol. 1.*
Fasc. 2. Fauna and flora of the Bay of Naples. 35.
Jerusalem: Israel Program for Scientific Translations. Pp.
293-917.
- Sweeney, M. J., et al. 1992 "Larval" and juvenile cephalopods:
a manual for their identification. Smithsonian Contribution
to Zoology. 513. Washington D.C.: Smithsonian Institution
Press. viii+282 pp.

Bivalvia

- Bentham Junting van, T. 1943 *Mollusca (I) C Lamellibranchia.*
Fauna van Nederland. 12. Leiden: A. W. Sijthoff. 477 pp.
- Bondesen, P. 1984 Danske havmuslinger. *Natur og Museum*
23(2): 1-31.
- Christensen, J. M., Larsen, S. & Nystöm, O. 1978 *Muslinger.*
Oslo: J.W. Cappelens Forlag. 125 pp.
- Gomez Rodriguez, R. & Pérez Sanchez, J. M. 1997 *Moluscos*
bivalvos de Canarias. Las Palmas de Gran Canarias:
Calbido Insular de Gran Canaria. 425 pp.
- Grossu, A. V. 1962 *Bivalvia (Scoici).* Fauna Republicii
Populare Române. 3(3). Bucuresti: Editura Academiei
Republicii populare Române. 426 pp.
- Haas, F. 1926 *Lamellibranchia.* Tierwelt der Nord- und Ostsee.
9dl. Leipzig: Akademische Verlagsgesellschaft. 96 pp.
- Jensen, A. D. & Spärck, R. 1934 *Blöddyr. II.*
Salvandsmuslinger. Danmarks Fauna. 40. København:
GEC, Gad. 208 pp.
- List, T. 1902 *Die Mytiliden des Golfes von Neapel und der*
Angrenzenden Meeres-Abschnitte. Fauna und flora des
Golfes von Neapel und der Angrenzenden Meeres-
Abschnitte. 27. Berlin: R. Friedländer & Sohn. 312 pp.
- Nordsieck, F. 1969 *Die europäischen Meeres-*
Gehäuseschnecken (Bivalvia) Vom Eismeer bis Kapverden,
Mittelmeer und Schwarzes Meer. Stuttgart: Gustav Fischer
Verlag. 256 pp.
- Piechocki, A. & Dyduch-Falniowska, A. 1993 *Mieczaki Malze*
(*Mollusca, Bivalvia of Poland.*) RCNP, Poland. 200 pp.
- Tebble, N. 1976 *British bivalve seashells. A handbook for*
identification. Edinburgh: Royal Scottish Museum. 213
pp.
- Zenetas, A. 1996 *The marine Bivalvia (Mollusca) of Greece.*
Fauna Graeciae. 7. Athens. 319 pp.

ANNELIDA

General

- Gerlach, S. A. 1967 *Annelida.* Die Tierwelt Mitteleuropas. 1-7a.
Leipzig: Quelle & Meyer. 8 pp.
- Remane, A. 1932 *Archannelida.* Tierwelt der Nord- und
Ostsee. 6a. 36 pp.

Polychaeta

- Bianchi, C. N. 1981 *Polichetti Serpuloidei.* Guida per il
riconoscimento delle specie animale delle acque lagunari e
costiere italiane. 5. Consiglio Nazionale delle Ricerche. 187
pp.
- Böggemann, M. 1997 Polychaeten aus der Deutschen Bucht:
Taxonomische Bearbeitung und Dokumentation der vom
Forschungsinstitut Senckenberg hauptsächlich in der
Deutschen Bucht gesammelten Polychaeten. *Courier*
Forschungsinstitut Senckenberg 202(Special issue): 1-315.
- Campoy, A. 1982 *Fauna de annelidos poliquetos de la*
Peninsula Iberica. Fauna de España. 7. 781 pp.
- Chambers, S. J. 1985 *Polychaetes from Scottish waters. Part 2*
Families Aphroditidae, Sigalionidae and Polyodontidae.
Edinburgh: Royal Scottish Museum Studies. 38 pp.
- Chambers, S. J. & Garwood, P. 1992 *Polychaetes from Scottish*
waters. Part 3 Family Nereidae. Edinburgh: National
Museums of Scotland. 65 pp.
- Chambers, S. J. & Muir, A. I. 1997 *Polychaetes: British*
Chrysopetaloidea, Pisinoidea and Aphroditoidae. Keys
and notes for the identification of the species. Synopses of
the British Fauna (New Series). 54. Shrewsbury: Field
Studies Council. 202 pp.

- Eisig, H. 1887 *Monographie der Capitelliden des Golfes von*
Neapel und der Angrenzenden Meeres-Abschnitte. Fauna
und flora des Golfes von Neapel und der Angrenzenden
Meeres-Abschnitte. 16. Berlin: R. Friedländer & Sohn. 906
pp.

- Eisig, H. 1906 *Ichthyotomus sanguinarius eine auf aalen*
schmarotzende annelide. Fauna und flora des Golfes von
Neapel und der Angrenzenden Meeres-Abschnitte. 28.
Berlin: R. Friedländer & Sohn. 300 pp.

- Fauchald, K. 1977 The polychaete worms. Definitions and keys
to the orders, families and genera. *Natural History Museum*
of Los Angeles County. Science Series 28: 1-188.

- Fauvel, P. 1923 *Polychètes errantes.* Faune de France. 5. Paris:
P. Lechevalier. 488 pp.

- Fauvel, P. 1927 *Polychètes sédentaires, addenda aux errantes,*
Archianélidés, Myzostomaires. Faune de France. 16. Paris:
P. Lechevalier. 494 pp.

- Fraipont, J. 1887 *Le genre Polygordius.* Fauna und flora des
Golfes von Neapel und der Angrenzenden Meeres-
Abschnitte. 14. Berlin: R. Friedländer & Sohn. 125 pp.

- Friedrich, H. 1938 *Polychaeta.* Tierwelt der Nord- und Ostsee.
6b. Leipzig: Akademische Verlagsgesellschaft. 201 pp.

- George, J. D. & Hartmann-Schröder, G. 1985 *Polychaetes:*
British Amphionimida, Spintherida and Eunicida. Keys
and notes for the identification of the species. Synopses of
the British Fauna (New Series). 32. London: E.J.
Brill/Dr. W. Backhuys. 221 pp.

- Hartley, J. P. 1981 The family Paraonidae, (Polychaeta) in
British waters, a new species and new records with a key to
species. *Journal of the Marine Biological Association of the*
United Kingdom 61(1): 133-150.

- Hartmann-Schröder, G. 1996 *Annelida, borstenwürmer,*
Polychatea. Die Tierwelt Deutschlands. 58. Jena: G.
Fischer. 648 pp.

- Hartmann-Schröder, G. 1988 *Key to the polychaete annelids*
from the North Sea and Baltic approaches. Edinburgh:
Heriot-Watt University, Institute of Offshore Engineering.
63 pp.

- Holthe, T. 1986 *Polychaeta - Terebellomorpha.* Marine
invertebrates of Scandinavia. 7. Trondheim: Norwegian
University Press. 194 pp.

- Holthe, T. 1992 Identification of Annelida Polychaeta from
northern European and adjacent arctic waters. *Gunneria* 66:
1-30.

- Katzmann, W., Laubier, L. & Ramos, J. 1974 Pilargidae
(Annelides Polychètes errantes) de Méditerranée. *Bulletin*
de l'Institut océanographique de Monaco 71(1428): 1-40.

- Katzmann, W. & Laubier, L. 1975 Paraonidae (Polychètes
sédentaires) de l'Adriatique. *Annalen des Naturhistorischen*
Museums in Wien 79: 568-588.

- Kirkegaard, J. B. 1992 *Havborsteorme 1.* Danmarks Fauna. 83.
416 pp.

- Kirkegaard, J. B. 1996 *Havborsteorme 2.* Danmarks Fauna. 86.
451 pp.

- Knight-Jones, P. & Knight-Jones, E. W. 1977 Taxonomy and
ecology of British Spirorbidae (Polychaeta). *Journal of the*
Marine Biological Association of the United Kingdom 57:
453-500.

- Lardicci, C. 1989 Censimento dei Polycheti dei mari Italiani
Spionidae Grube, 1850. *Atti della Società Toscana di*
Scienze Naturale Memorie, Serie B 96: 121-152.

- Laubier, L. & Ramos, J. 1973 Paraonidae (Polychètes
sédentaires) de Méditerranée. *Bulletin du Muséum national*
d'Histoire naturelle, Paris, 3ème série, Zoologie 113: 1097-
1148.

- Marinov, T. 1977 *Mnogochetinesti cherven (Polychaeta).* Fauna
na Bulgariia. 6. Sofia: Bulgarskata akademiya na naukite.
258 pp.

- O'Connor, B. D. S. 1987 The Glyceridae (Polychaeta) of the
North Atlantic and Mediterranean, with descriptions of two
new species. *Journal of Natural History* 21: 167-189.

- Pieranton, U. 1908 *Protodrilus.* Fauna und flora des Golfes von
Neapel und der Angrenzenden Meeres-Abschnitte. 31.
Berlin: R. Friedländer & Sohn. 226 pp.

- Plate, S. & Husemann, E. 1994 Identification guide to the planktonic polychaete larvae around the island of Helgoland (German Bight). *Helgolander Meeresuntersuchungen* 48(1): 1-58.
- Pleijel, F. & Dales, R. P. 1991 *Polychaetes: British Phyllodocoideans, Typhloscolecoides and Tomopteroideans*. Synopses of the British Fauna (New Series), 45. Oegstgeest: Universal Book Service / Backhuys Publishers. 202 pp.
- Pleijel, F. 1993 *Polychaeta Phyllodocidae*. Marine invertebrates of Scandinavia. 8. Scandinavian University Press. 159 pp.
- Rainer, S. F. 1991 The genus *Nephthys* (Polychaeta: Phyllodocida) of Northern Europe: a review of species, including the description of *N. pulchra* sp.n. and a key to the Nephtyidae. *Helgoländer Meeresuntersuchungen* 45: 65-96.
- Støp-Bowitz, C. 1941 Les glycériens de Norvège. *Nytt Magasin for Naturvitenskapene* 82: 181-250.
- Støp-Bowitz, C. 1946 Les ophéliens norvégiens. *Nytt Magasin for Naturvitenskapene* 85: 21-61.
- Støp-Bowitz, C. 1946 Les scalibregmiens de Norvège. *Nytt Magasin for Naturvitenskapene* 85: 63-87.
- Støp-Bowitz, C. 1948 Les flabelligériens norvégiens. *Bergens Museums Årbok. Naturvitenskapelig Rekke* 1946-1947(2): 1-59.
- Støp-Bowitz, C. 1948 Sur les polychètes arctiques des familles des glycériens, des ophéliens, des scalibregmiens et des flabelligériens. *Tromsø Museums Årshefter* 66(2): 1-58.
- Tebble, N. & Chambers, S. 1982 *Polychaetes from Scottish waters. A guide to identification. Part I. Family Polynoidae*. Edinburgh: Royal Scottish Museum Studies. 73 pp.
- Ushakov, P. V. 1972 *Mnogoschetinkovye chervi podotriada Phyllodociformia poliarnogo basseina i severo-zapadnoi chasti Tikhogo okeana: Semeistva Phyllodocidae, Alciopidae, Tomopteridae Typhloscolecidae i Lacydoniidae*. Fauna SSSR (novaia seriiia). 102. Leningrad: Akademiiia nauk SSSR, Zoologicheskii institut. 270 pp.
- Ushakov, P. V. 1974 *Polychaetes of the suborder Phyllodociformia of the polar basin and the northwestern part of the Pacific. Families Phyllodocidae, Alciopidae, Tomopteridae Typhloscolecidae and Lacydoniidae*. Fauna SSSR (novaia seriiia). 102. Jerusalem: Israël Program for Scientific Translations. 259 pp.
- Ushakov, P. V. 1982 *Mnogoschetinkovye chervi podotriada Aphroditiformia severnogo Ledovitogo okeana i severo-zapadnoi chasti Tikhogo okeana: Semeistva Aphroditidae i Polynoidae*. Fauna SSSR (novaia seriiia). 126. Leningrad: Leningradskoe otdelenie. 272 pp.
- Westhead, W. 1990 *Polychaetes: interstitial families. Keys and notes for the identification of the species. Synopses of the British Fauna (New Series)*, 44. Oegstgeest: Universal Book Service / Backhuys Publishers. VII + 152 pp.
- ### Oligochaeta
- Brinkhurst, R. O. 1982 *British and other marine and estuarine oligochaetes. Keys and notes for the identification of the species*. Synopses of the British Fauna (New Series). 21. Cambridge: Cambridge University Press. 127 pp.
- Chekanovskaya, O. V. 1962 *Aquatic oligochaetes in the USSR fauna*. Opredelitel po faune SSSR. 78. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 411 pp.
- Michaelsen, W. 1927 *Clitellata: Oligochaeta*. Tierwelt der Nord- und Ostsee. 6cl. Leipzig: Akademische verlagsgesellschaft. 44 pp.
- Tynen, M. J. & Nurminen, M. 1969 A key to the European littoral Enchytraeidae (Oligochaeta). *Ann. Zool. Fennici* 6: 150-155.
- Ude, H., Johanesson, L. & Ten Broecke, A. 1929 *Würmer oder Vermes. Oligochaeta. Hirudinea (Hegel). Sipunculoidea und Echiuroidea*. Die Tierwelt Deutschlands. 15. Jena: G. Fischer. 168 pp.
- ### Hirudinea
- Herter, K. 1935 *Hirudinea*. Tierwelt der Nord- und Ostsee. 6c2. Leipzig: Akademische Verlagsgesellschaft. Pp. 45-106.
- Lukin, E. I. 1976 *Leeches of fresh and brackish waters. Leeches*, 1. Fauna SSSR (novaia seriiia). 109. Leningrad: Akademiiia nauk SSSR, Zoologicheskii institut. 484 pp.
- Ude, H., Johanesson, L. & Ten Broecke, A. 1929 *Würmer oder Vermes. Oligochaeta. Hirudinea (Hegel). Sipunculoidea und Echiuroidea*. Die Tierwelt Deutschlands. 15. Jena: G. Fischer. 168 pp.
- ### Pogonophora
- Ivanov, A. V. 1960 *Pogonofory*. Fauna SSSR (novaia seriiia). 75. Moskva: Akademiiia nauk SSSR, Zoologicheskii institut. 271 pp.
- Ivanov, A. V. 1963 *Pogonophora*. London: Academic Press. 479 pp.
- ### TARDIGRADA
- Cuénot, L. 1932 *Tardigrades*. Faune de France. 24. Paris: P. Lechevalier. 96 pp.
- Marcus, E. 1928 *Spinnentiere oder Arachnoidea. IV. Bärtierchen (Tardigrada) VIII*. Die Tierwelt Deutschlands. 12. 230 pp.
- Maucci, W. 1986 *Tardigrada*. Fauna d'Italia. 24. Bologna: Calderini. 388 pp.
- Morgan, C. I. & King, P. E. 1976 *British tardigrades. Synopses of the British Fauna*. 9. London, New-York: Academic Press. 133 pp.
- Rahm, G. 1928 *Tardigrada*. Tierwelt der Nord- und Ostsee. 1lb1. Leipzig: Akademische verlagsgesellschaft. 25 pp.
- Ramazzotti, G. 1972 Il phylum Tardigrada. *Memorie dell'Istituto di Idrobiologia* 19: 1-732.
- Thulin, G. 1911 Beiträge zur Kenntnis der Tardigraden fauna Schwedens. *Arkiv for Zoologi* 7(16): 1-60.
- ### CHELICERATA
- #### General
- André, M. 1946 *Halacariens marins*. Faune de France. 46. Paris: P. Lechevalier. 152 pp.
- Green, J. & MacQuitty, M. 1987 *Halacarid Mites (Arachnida: Acari): Keys and Notes for the Identification of the Species. Synopses of the British Fauna (New Series)*, 36. Leiden: E. J. Brill/W. Backhuys. 178 pp.
- Kästner, A., Schenkel, E., Loman, J. C. C. & Sprehn, C. 1928 *Spinnentiere oder Arachnoidea. Opiliones (Webernechte, Kanker). Pseudoscorpionida (Afterskorpione). Pantopoda oder Asselspinnen (Meerspinnen). Pentastomida, Linguatulida (Zungenwürmer)*. Die Tierwelt Deutschlands. 8. Jena: G. Fischer. 94 pp.
- Legg, G. & Jones, R. E. 1988 *Pseudoscorpions*. Synopses of the British Fauna (New Series), 40. Leiden: E. J. Brill/W. Backhuys. 159 pp.
- Sokolov, I. I. 1952 *Vodianye kleshchi. Chast II. Halacarae*. Fauna SSSR (novaia seriiia). 53. Moskva: Akademiiia nauk SSSR, Zoologicheskii institut. 200 pp.
- Viets, K. 1936 *Spinnentiere oder Arachnoidea. VII. Wassermilben I (Hydrachnellae und Halacaridae)*. Die Tierwelt Deutschlands. 31/32. Jena: G. Fischer. 574 pp.
- #### Pycnogonida
- Bouvier, E.-L. 1923 *Pycnogonides*. Faune de France. 7. Paris: P. Lechevalier. 69 pp.
- Dohrn, A. 1881 *Die Pantopoden des Golfs von Neapel*. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 3. Leipzig: W. Engelmann. 252 pp.
- Helfer, H. 19?? *Pantopoda*. Tierwelt der Nord- und Ostsee. 1la3. Leipzig: Akademische verlagsgesellschaft. 5 pp.
- Kästner, A., Schenkel, E., Loman, J. C. C. & Sprehn, C. 1928 *Spinnentiere oder Arachnoidea. Opiliones (Webernechte, Kanker). Pseudoscorpionida (Afterskorpione). Pantopoda oder Asselspinnen (Meerspinnen). Pentastomida*,

- Lingualulida (Zungenwürmer).* Die Tierwelt Deutschlands. 8. Jena: G. Fischer. 94 pp.
- King, P. E. 1974 *British seaspiders.* Synopses of the British Fauna (New Series). 5. London, New-York: Academic Press. 68 pp.
- King, P. E. 1986 *Sea spiders: A revised key to the adults of littoral Pycnogonida in the British Isles.* AIDGAP Guides. 6. Shrewsbury: Field Studies Council. Pp. 493-516.
- Meisenheimer, J. 1927 *Pantopoda.* Tierwelt der Nord- und Ostsee. 11a1, a2. Leipzig: Akademische verlagsgesellschaft. 15 pp.
- Stephensen, K. 1933 *Havedderkopper (Pycnogonida) og rankefodder (Cirripedia).* Danmarks Fauna. 38. Kobenhaven: GEC Gad. 158 pp.

CRUSTACEA

General and other miscellaneous groups

- Anadon, R. 1975 Aportación al conocimiento de la fauna bentónica de la ría de Vigo (N.W. de España). I. Pycnogónidos y crustáceos de Panjón. *Investigación Pesquera* 39(1): 199-218.
- Cottarelli, V. & Mura, G. 1983 *Anostraci, notostraci, concostraci (Crustacea: Anostraca, Notostraca, Conchostraca): Consiglio Nazionale delle Ricerche.* 73 pp.
- Demetropoulos, A. & Neocleous, D. 1969 The Fishes and Crustaceans of Cyprus. *Fisheries Bulletin* 1.
- Enckell, P. H. 1980 *Kräftdjur. Feltfauna.* Lund: Signum. 685 pp.
- Lomakina, N. B. 1978 *Euphausiacea of the world's oceans.* Opredeliteli po faune SSSR. 118. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 224 pp.
- Manuilova, E. F. 1964 *Cladocera in the USSR fauna.* Opredeliteli po faune SSSR. 88. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 327 pp.
- Margaritora, F. G. 1985 *Cladocera.* Fauna d'Italia. 23. Bologna: Calderini. 400 pp.
- Mauchline, J. 1984 *Euphausiid, stomatopod and leptostracan crustaceans. Keys and notes for the identification of the species.* Synopses of the British Fauna (New Series). 30. London: E.J. Brill/Dr.W. Backhuys. 91 pp.
- Negrea, S. 1983 *Cladocera.* Fauna Republicii Socialiste România. 4(12). Bucuresti: Editura Academiei Republicii Socialiste România. 399 pp.
- Rammer, W. 1930 *Phyllopoda.* Tierwelt der Nord- und Ostsee. 10a. Leipzig: Akademische verlagsgesellschaft. 32 pp.
- Thiele, J. 1926 *Leptostraca.* Tierwelt der Nord- und Ostsee. 10g1. Leipzig: Akademische verlagsgesellschaft. 8 pp.
- Zimmer, C. 1932 *Euphausiacea.* Tierwelt der Nord- und Ostsee. 10g2. Leipzig: Akademische verlagsgesellschaft. Pp. 9-28.

Cladocera (Branchiopoda)

- Alonso, M. 1996 *Crustacea, Branchiopoda.* Fauna Iberica. 7. Madrid: Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Científicas. 486 pp.
- Botnariuc, N. & Orghidan, T. 1953 *Phyllopoda.* Fauna Republicii Populare Române. 4(2). Bucuresti: Academia Republicii Populare Române. 100 pp.
- Flössner, D. 1972 *Krebstiere, Crustacea, Kiemen- und Blattfusser, Branchiopoda, Fischlause, Branchiura.* Die Tierwelt Deutschlands. 60. 501 pp.
- Røen, U. I. 1995 *Gællefodder og karpelus.* Danmarks Fauna. 85. 358 pp.

Ostracoda

- Angel, M. V. 1993 *Marine Planktonic Ostracods.* Synopses of the British Fauna (New Series). 48. Shrewsbury: Field Studies Council. 240 pp.
- Athersuch, J., Horne, D. J. & Whittaker, J. E. 1989 *Marine and Brackish Water Ostracods (Superfamilies Cypridacea and Cytheracea). Keys and notes for the identification of the species.* Synopses of the British Fauna (New Series). 43. Leiden, New-York: E.J. Brill. 343 pp.

- Caraior, F. H. 1967 *Ostracoda. Fam. Cytheridae (Ostracode marine si salmistrole).* Fauna Republicii socialiste România. 4(10). Bucuresti: Academia Republicii socialiste România. 168 pp.
- Elofson, O. 1969 *Marine Ostracoda of Sweden with special consideration of the Skagerrak.* Jerusalem: Israël program for scientific translation. 286 pp.
- Klie, W. 1929 *Ostracoda.* Tierwelt der Nord- und Ostsee. 10b. s.l. 56 pp.
- Klie, W. 1938 *Krebstiere, oder Crustacea. III. Ostracoda Muschelkrebs.* Die Tierwelt Deutschlands. 34. Jena: G. Fischer. 230 pp.
- Müller, G. N. 1894 *Die Ostracoden des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte.* Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 21. Berlin: R. Friedländer & Sohn. 404 pp.
- Sywula, T. 1974 *Malzoraczki (Ostracoda).* Fauna Slodkowodna Polski. 24. Poznan: Panstwowe Wydawn. 314 pp.

Copepoda

- Brodskii, K. A. 1950 *Calanoidea of the Far Eastern seas of the USSR and the Arctic Basin.* Opredeliteli po faune SSSR. 35. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 442 pp.
- Brodskii, K. A., Vyshkvertseva, N. V., Kos, M. S. & Markhaseva, E. L. 1983 *Copepoda: Calanoidea of the USSR seas and adjacent waters.* Opredeliteli po faune SSSR. 135. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 357 pp.
- Carli, A. & Crisafi, P. 1983 *Copepodi lagunari.* Guide per il riconoscimento delle specie animale delle acque lagunari e costiere italiane. 11. Consiglio Nazionale delle Ricerche. 125 pp.
- Giesbrecht, W. 1892 *Systematik und faunistik der Pelagischen Copepoden des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte.* Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 19. Berlin: R. Friedländer & Sohn. 831 pp.
- Giesbrecht, W. 1899 *Die Asterocheriden des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte.* Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 25. Berlin: R. Friedländer & Sohn. 217 pp.
- Gotto, V. 1993 *Commensals and parasitic copepods associated with marine invertebrates (and whales).* Synopses of the British Fauna (New Series). 46. Oegstgeest: Universal Book Services / W. Backhuys. 264 pp.
- Huys, R., Gee, J. M., More, C. G. & Hamond, R. 1996 *Marine and brackish water harpacticoid copepods Part I.* Synopses of the British Fauna (New Series). 51. Leiden, New-York: E.J. Brill. 352 pp.
- Kabata, Z. 1979 *Parasitic Copepoda of British fishes.* London: The Ray Society. 469 pp.
- Kabata, Z. 1992 *Copepods parasitic on fishes.* Synopses of the British Fauna (New Series). 47. London: Universal Book Services / Dr. W. Backhuys. 264 pp.
- Oorde-de Lint van, G. M. & Schuurmans Stekhoven, J. H. 1936 *Copepoda parasitica.* Tierwelt der Nord- und Ostsee. 10c2. s.l. Pp. 73-198.
- Pesta, O. 1928 *Krebstiere, oder Crustacea. I. Ruderfüßer oder Copepoda (1. Calanoida, 2. Cyclopoida).* Die Tierwelt Deutschlands. 9. Jena: G. Fischer. 136 pp.
- Pesta, O. 1932 *Krebstiere, oder Crustacea. I. Ruderfüßer oder Copepoda. 3. Unterordnung Harpacticoida.* Die Tierwelt Deutschlands. 24. 164 pp.
- Pesta, O. 1934 *Krebstiere, oder Crustacea. I. Ruderfüßer oder Copepoda. 4. Monstrilloidea, 5. Notodelphyoida, 6. Caligoida, 7. Lernaeoida.* Die Tierwelt Deutschlands. 29. Jena: G. Fischer. 68 pp.
- Rose, M. 1933 *Copepodes pélagiques.* Faune de France. 26. Paris: P. Lechevalier. 374 pp.
- Shuvalov, V. S. 1980 *Cyclopeida of the family Oithonidae of the world's oceans.* Opredeliteli po faune SSSR. 125. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 197 pp.

Cirripedia

- Bassindale, R. 1964 British barnacles with keys and notes for the identification of the species. *Synopses of the British Fauna*. 14. London. 49 pp.
- Broch, H. 1924 *Cirripedia Thoracica von Norwegen und dem Norwegischen Nordmeer*. Skrifter, Videnskabs-Selskabet i Christiania, Matematisk-naturvidenskapelig Klasse 1924(17): 1-121.
- Høeg, J. & Lützen, J. 1985 *Crustacea Rhizocephala. Marine invertebrates of Scandinavia*. 6. Scandinavian University Press. 92 pp.
- Kolosvary, G. 1947 Die Balaniden der Adria. *Annales Historico Naturales Musei nationalis Hungarici* 39(1): 1-88.
- Kruger, P. 1927 *Cirripedia. Tierwelt der Nord- und Ostsee*. 10d. Leipzig: Akademische verlagsgesellschaft. 40 pp.
- Nilson-Cantell, C. A. 1978 *Cirripedia Thoracica and Acrothoracica. Marine invertebrates of Scandinavia*. 5. Oslo: Universitetsforlaget. 136 pp.
- Relini, G. 1980 *Cirripedi toracici. Guide per il riconoscimento delle specie animale delle acque lagunari e costiere italiane*. 2. Consiglio Nazionale delle Ricerche. 119 pp.
- Smith, G. 1906 *Rhizocephala. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte*. 29. Berlin: R. Friedländer & Sohn. 123 pp.
- Southward, A. J. & Crisp, D. J. 1963 *Les Cirripèdes des mers européennes. Catalogue des principales salissures marines (rencontrées sur les coques des navires dans les eaux européennes)*. Vol. 1: *Balanæ*. Paris: O.C.D.E. 46 pp.
- Stephensen, K. 1933 *Hovedderkopper (Pycnogonida) og rankefædder (Cirripedia)*. Danmarks Fauna. 38. Kobenhavn: GEC Gad. 158 pp.
- Tarasov, N. I. & Zevina, G. B. 1957 *Usonogieraki (Cirripedia Thoracica) morei SSSR. Fauna SSSR (novaia serii)*. 69. Moskva: Akademii nauk SSSR, Zoologicheskii institut. 267 pp.
- Young, P. S. 1998 *Cirripedia (Crustacea) from the "Campagne Biacores" in the Azores region including a generic revision of Verrucidae*. *Zoosystema* 20(1): 31-92.
- Zevina, G. B. 1981 *Cirripedia of the suborder Lepadomorpha (Cirripedia, Thoracica) of the world's oceans. Part I. Fam. Scalpellidae. Opredelitel po faune SSSR*. 127. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 406 pp.
- Zevina, G. B. 1982 Suborder Lepadomorpha of the world's oceans. Part 2. Opredelitel po faune SSSR. 133. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 224 pp.

Stomatopoda

- Balss, H. 1926 *Stomatopoda. Tierwelt der Nord- und Ostsee*. 10h1. Leipzig: Akademische verlagsgesellschaft. 8 pp.
- Froglio, C. & Manning, R. B. 1989 *Checklist and key to adult Mediterranean stomatopod Crustacea. Selected Symposia and Monographs*. 3. Mucchi: Unione Zoologica Italiana. Pp. 265-273.
- Giesbrecht, W. 1910 *Stomatopoden. Erster theil*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 33. Berlin: R. Friedländer & Sohn. 239 pp.
- Holthuis, L. B. 1950 *Decapoda (K IX) A Natantia, Macrura Reptantia, Anomura en Stomatopoda (K X)*. Fauna van Nederland. 15. Leiden: A.W. Sijthoff. 166 pp.
- Mauchline, J. 1984 *Euphausiid, stomatopod and leptostracan crustaceans. Keys and notes for the identification of the species. Synopses of the British Fauna (New Series)*. 30. London: E.J. Brill/Dr.W. Backhuys. 91 pp.
- Nobre, A. 1936 *Crustaceos, decapodes e stomatopodes marinhas de Portugal*. Fauna marinha de Portugal. 4. Barcelos: Companhia Editora do Minho. viii+213 pp.

Decapoda

- Adema, J. P. H. M. 1991 *De krabben van Nederland en België (Crustacea, Decapoda, Brachyura)*. Leiden: Nationaal Natuurhistorisch Museum. 244 pp.

- Allen, J. A. 1967 *The fauna of the Clyde Sea area: Crustacea: Euphausiaceae and Decapoda, with an illustrated key to the British species*. Millport: Scottish Marine Biological Association. 116 pp.
- Bacescu, M. C. 1967 *Decapoda*. Fauna Republicii Socialiste România. 4(9). Bucuresti: Academia Republicii Socialiste România. 356 pp.
- Balss, H. 1926 *Decapoda. Tierwelt der Nord- und Ostsee*. 10h2. Leipzig: Akademische verlagsgesellschaft. pp. 9-112.
- Balss, H. 1927 *Decapoda. Tierwelt der Nord- und Ostsee*. 10h3. Leipzig: Akademische verlagsgesellschaft. pp. 113-116.
- Bouvier, E.-L. 1940 *Décapodes marcheurs. Faune de France*. 37. Paris: P. Lechevalier. 404 pp.
- Burukovsky, R. N. 1985 *Key to shrimps and lobsters. Russian translations series*. 5. Rotterdam: A. A. Balkema. 188 pp.
- Christiansen, M. E. 1969 *Crustacea Decapoda Brachyura. Marine invertebrates of Scandinavia*. 2. Oslo: Universitetsforlaget. 143 pp.
- Christiansen, M. E. 1972 *Bestemmelsestabell over Crustacea Decapoda, tifotkrepes*. Oslo: Universitetsforlaget. 71 pp.
- Cottiglia, M. 1983 *Crostatei decapodi lagunari*. Guide per il riconoscimento delle specie animali delle acque lagunari e costiere italiane. 10. Genova: Consiglio Nazionale delle Ricerche. 1-148 pp.
- Crothers, J. & Crothers, M. 1983 A key to the crabs and crab-like animals of British inshore waters. *Field Studies* 5: 753-806.
- Falciai, L. & Minervini, R. 1992 *Guida dei crostacei decapodi d'Europa*. Padova: Franco Muzzio. 282 pp.
- Falciai, L. & Minervini, R. 1996 *Guide des homards, crabes, langoustes, crevettes et autres crustacés décapodes d'Europe*. Lausanne, Paris: Delachaux & Niestlé. 287 pp.
- Gonzales Perez, J. A. 1995 *Crustaceos decapodos de las Islas Canarias*. Santa Cruz de Tenerife: Turquesa. 282 pp.
- Guillén Neto, J. E. 1990 *Guía ilustrada de los crustáceos decápodos del litoral alicantino*. Alicante: Instituto de Cultura "Juan Gil-Albert". 316 pp.
- Gurriaran, E. G. & Mendez, M. 1985 *Crustaceos decapodos das costas de Galicia. I. Brachyura*. Cuadernos da Area de Ciencias Biologicas. Vol. 2. Coruna: Seminario de estudios Galegos. 242 pp.
- Holthuis, L. B. 1950 *Decapoda (K IX) A Natantia, Macrura Reptantia, Anomura en Stomatopoda (K X)*. Fauna van Nederland. 15. Leiden: A.W. Sijthoff. 166 pp.
- Holthuis, L. B., Heerebout, G. R. & Adema, J. P. H. M. 1986 *De Nederlandse Decapoda (Garnalen, kreeften en krabben)*. Wetenschappelijke mededelingen KNNV. 179. Hoogvliet: Koninklijke Nederlandse Natuurhistorische Vereniging. 66 pp.
- Ingle, R. W. 1980 *British crabs*. London, Oxford: British Museum (Natural History)/Oxford University Press. 222 pp.
- Ingle, R. W. 1991 *Larval stages of north east Atlantic crabs: an illustrated key*. Natural History Museum Publications. Chapman and Hall Identification Guides. 1. London, New-York: Chapman and Hall. 363 pp.
- Ingle, R. W. 1993 *Hermit crabs of the northeastern Atlantic Ocean and Mediterranean Sea. An illustrated key*. London: Chapman and Hall. 495 pp.
- Ingle, R. W. 1996 *Shallow water crabs*. Synopses of the British Fauna (New Series). 25. Shrewsbury: Field Studies Council. 245 pp.
- Ingle, R. W. 1997 *Crayfishes, lobsters and crabs of Europe: an illustrated guide to common and traded species*. London, New-York: Chapman & Hall. 281 pp.
- Makarov, V. V. 1938 *Anomura*. Fauna SSSR (novaia serii). 16. Moskva: Akademii nauk SSSR, Zoologicheskii institut. 324 pp.
- Nobre, A. 1931 *Crustaceos decapodes e stomatopodes marinhas de Portugal*. Porto: Imprensa Portuguesa. 307 pp.
- Nobre, A. 1936 *Crustaceos, decapodes e stomatopodes marinhas de Portugal*. Fauna marinha de Portugal. 4. Barcelos: Companhia Editora do Minho. viii+213 pp.
- Noë, P.-Y. 1992 *Clé préliminaire d'identification des Crustacea Decapoda de France et des principales autres*

- espèces d'Europe.* Patrimoines Naturels. 9. Paris: Muséum national d'Histoire naturelle. 146 pp.
- Paula, J. 1996 A key and bibliography for the identification of zoeal stages of brachyuran crabs from the Atlantic coast of Europe. *Journal of Plankton Research* 18(1): 17-27.
- Sandberg, L. & McLaughlin, P. A. 1998 *Crustacea, Decapoda, Paguridea.* Marine invertebrates of Scandinavia. 10: Scandinavian University Press. 120 pp.
- Schellenberg, A. 1928 *Krebstiere oder Crustacea. II. Decapoda, Zehnfüßler IV.* Die Tierwelt Deutschlands. 10. 146 pp.
- Smaldon, G., Holthuis, L. B. & Fransen, C. H. J. M. 1993 *Coastal shrimps and prawns. Keys and notes for the identification of the species.* Synopses of the British Fauna (New Series). 15. Shrewsbury: Field Studies Council. vii+142 pp.
- Stephensen, K. & Meinertz, N. T. 1910 *Storkrebs I. Skjoldkrebs.* Danmarks Fauna. 9. 193 pp.
- Zariquey Álvarez, R. 1968 *Crustáceos decápodos ibéricos.* 32. Barcelona: Investigación Pesquera. xv+510 pp.
- ### Mysidacea
- Bacescu, M. C. 1941 *Mysidacea.* Fauna Republicii Socialiste România. 4(3). Bucuresti: Academia Republicii popolare Romîne. 126 pp.
- Lagardère, J.-P. & Nouvel, H. 1980 Les Mysidacés du talus continental du golfe de Gascogne. II. Familles des Lophogastridae, Eucopiidae et Mysidae (tribu des Erythropini exceptée). *Bulletin du Muséum national d'Histoire naturelle, Paris, 4ème série, section A* 2(2): 375-412.
- Lagardère, J.-P. 1983 Les Mysidacés de la plaine abyssale du golfe de Gascogne. I. Familles des Lophogastridae, Eucopiidae et Petalophthalmidae. *Bulletin du Muséum national d'Histoire naturelle, Paris, 4ème série, section A* 5(3): 809-843.
- Makings, P. 1977 A guide to the British coastal Mysidacea. *Field Studies* 4: 575-595.
- Mauchline, J. 1980 *Biology of mysids and euphausiids.* Advances in Marine Biology. 18. London: Academic Press. 681 pp.
- Nouvel, H. 1950 Mysidacea. In *Fiches d'identification du zooplancton* (ed. P. Jespersen & F. S. Russell), pp. 18-27. Conseil permanent international pour l'exploitation de la mer.
- Nouvel, H. & Lagardère, J. P. 1976 Les Mysidacés du talus continental du golfe de Gascogne. I. Tribu des Erythropini (genre *Erythrops* excepté). *Bulletin du Muséum national d'Histoire naturelle, Paris, 3ème série, Zoologie* 414(291): 1243-1324.
- Tattersall, W. M. & Tattersall, O. S. 1951 *The British Mysidacea.* London: Ray Society. 460 pp.
- Tattersall, O. S. 1955 Mysidacea. "Discovery" Reports 28: 1-190.
- Zimmer, C. 1933 *Mysidacea.* Tierwelt der Nord- und Ostsee. 10g3. Leipzig: Akademische verlagsgesellschaft. Pp. 29-69.
- ### Amphipoda
- Barnard, J. L. & Karaman, G. S. 1991 The families and genera of marine gammaridean Amphipoda (except marine gammaroids). Part 1. *Records of the Australian Museum suppl.* 13: 1-418.
- Barnard, J. L. & Karaman, G. S. 1991 The families and genera of marine gammaridean Amphipoda (except marine gammaroids). Part 2. *Records of the Australian Museum suppl.* 13: 419-866.
- Bellan-Santini, D., Karaman, G., Krapp-Schickel, G., Ledoyer, M., Myers, A. A., Russo, S. & Schieke, U. 1982 The Amphipoda of the Mediterranean Part. 1 Gammaridea (Acanthonotozomatidae to Gammaridae), ed. S. Russo. *Mémoires de l'Institut océanographique de Monaco* 13: i-xiii+1-364.
- Bellan-Santini, D., Diviacco, G., Krapp-Schickel, G., Myers, A. A. & Russo, S. 1989 The Amphipoda of the Mediterranean Part. 2 Gammaridea (Haustoriidae to

- Lysianassidae), ed. S. Russo. *Mémoires de l'Institut océanographique de Monaco* 13: xiv-xix+365-576.
- Bellan-Santini, D., Karaman, G., Krapp-Schickel, G., Ledoyer, M. & S., R. 1993 The Amphipoda of the Mediterranean Part. 3 Gammaridea (Melphidippidae to Talitidae), Ingolfiellidae, Caprellidae, ed. S. Russo. *Mémoires de l'Institut océanographique de Monaco* 13: xx-xxv+577-813.
- Bellan-Santini, D., Karaman, G., Myers, A. A., Ledoyer, M., Russo, S. & Vader, W. 1998 The Amphipoda of the Mediterranean, Part. 4, ed. S. Russo. *Mémoires de l'Institut Océanographique de Monaco* 13: xxvii-xliv+815-959.
- Bulycheva, A. I. 1957 *Amphipoda - Talitroidea of the USSR seas and adjoining waters.* Opredelitel po faune SSSR. 65. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 186 pp.
- Chevreux, E. & Fage, L. 1925 *Amphipodes.* Faune de France. 9. Paris: P. Lechevalier. 488 pp.
- Dauvin, J.-C. & Bellan-Santini, D. 1988 Illustrated key to *Ampelisca* from Northeastern Atlantic. *Journal of the Marine Biological Association of the United Kingdom* 68: 659-676.
- Guryanova, E. F. 1951 *Amphipoda - Gammaridea of the seas of the USSR and adjoining waters.* Opredelitel po faune SSSR. 41. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 1032 pp.
- Harrison, R. J. 1944 *Caprellidea (Amphipoda, Crustacea).* Synopses of the British Fauna. 2. London: Linnean Society of London.
- Lincoln, R. J. 1979 *British marine Amphipoda: Gammaridea.* London: British Museum (Natural History). 658 pp.
- Mayer, P. 1882 *Die Caprelliden des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte.* Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 6. Leipzig: W. Engelmann. 201 pp.
- Mayer, P. 1890 *Die Caprelliden des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte.* Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 17. Berlin: R. Friedländer & Sohn. 157 pp.
- Palerud, R. & Vader, W. 1991 Marine Amphipoda Gammaridea in North-East Atlantic and Norwegian arctic. *Tromsø, Naturvitenskap* 68: 1-97.
- Schellenberg, A. 1942 *Krebstiere oder Crustacea. IV. Flohkrebse oder Amphipoda.* Die Tierwelt Deutschlands. 40. Jena: G. Fischer. 252 pp.
- Stephensen, K. 1929 *Amphipoda.* Tierwelt der Nord- und Ostsee. 10f. Leipzig: Akademische verlagsgesellschaft. 188 pp.
- Stephensen, K. 1935 The Amphipoda of N. Norway and Spitsbergen with adjacent waters. Part 1. *Tromsø Museums Skrifter* 3(1): 1-140.
- Stephensen, K. 1938 The Amphipoda of N. Norway and Spitsbergen with adjacent waters. Part 2. *Tromsø Museums Skrifter* 3(2): 141-278.
- Stephensen, K. 1940 The Amphipoda of N. Norway and Spitsbergen with adjacent waters. Part 3. *Tromsø Museums Skrifter* 3(3): 279-362.
- Stephensen, K. 1942 The Amphipoda of N. Norway and Spitsbergen with adjacent waters. Part 4. *Tromsø Museums Skrifter* 3(4): 363-526.
- Stephensen, K. & Meinertz, N. T. 1928 *Storkrebs II. Ringkrebs I. Tanglopper (Amfipoder).* Danmarks Fauna. 32. 399 pp.
- Valle, A. d. 1893 *Gammarini del Golfo di Napoli.* Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 20. Berlin: R. Friedländer & Sohn. 948 pp.
- Vasilenko, S. V. 1974 *Caprellidea of the USSR seas and adjacent waters.* Opredelitel po faune SSSR. 107. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 288 pp.
- Vinogradov, M. E., Volkov, A. F. & Semenova, T. N. 1996 *Hyperiid Amphipods of the world's oceans.* Opredelitel po faune SSSR. 132. Washington D.C.: Smithsonian Institution Libraries. xxvii+632 pp.

Cumacea

- Bacescu, M. C. 1951 *Cumacea*. Fauna Republicii Socialiste România. 4(1). Bucuresti: Academia Republicii popolare Romîne. 95 pp.
- Fage, L. 1951 *Cumacés*. Faune de France. 54. Paris: P. Lechevalier. 136 pp.
- Jones, N. S. 1958 *Cumacea*. Fiches d'identification du zooplancton. Pp. 71-80.
- Jones, N. S. 1976 *British cumaceans. Arthropoda: Crustacea. Keys and notes to the identification of the species. Synopses of the British Fauna (New Series)*. 7. London: Academic Press. 62 pp.
- Jones, N. S. 1984 The family Nannasticidae (Crustacea: Cumacea) from the deep Atlantic. *Bulletin of the British Museum (Natural History). Zoology Series* 46: 207-289.
- Lomakina, N. B. 1958 *Cumacea of the USSR seas*. Opredeliteli po faune SSSR. 66. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 302 pp.
- Zimmer, C. 1933 *Cumacea*. Tierwelt der Nord- und Ostsee. 10g4. Leipzig: Akademische verlagsgesellschaft. Pp. 70-120.

Tanaidacea

- Anonymous. 1988 *British shallow-water Tanaidacea*. Bangor: University College of North Wales. Taxonomic Workshop on the 'small crustacea'.
- Bacescu, M. C. 1961 Contribution à la connaissance des Tanaidacés de la Méditerranée orientale - I. Les Apseudidae et Kalliapseudidae des côtes d'Israël. *Bulletin of the Research Council of Israel, section B, Zoology* 10B(4): 137-170.
- Holdich, D. M. & Jones, J. A. 1983 *Tanaids. Keys and notes to the identification of the species. Synopses of the British Fauna (New Series)*. 27. Cambridge: Cambridge University Press. vii+98 pp.
- Holthuis, L. B. 1956 *Isopoda en Tanaidacea (KV)*. Fauna van Nederland. 16. Leiden: A.W. Sijthoff. 280 pp.
- Monod, T. 1923 Prodrome d'une faune des Tanaidacea et des Isopoda (excl. Epicaridea) des côtes de France (excl. Méditerranée). *Annales de la Société des Sciences Naturelles de Charente inférieure* 37(4): 19-126.
- Stephensen, K. & Meinertz, N. T. 1948 *Storkrebs IV. Ringkrebs 3. Tanglus (Marine Isopoder og Tanaider)*. Danmarks Fauna. 53. 187 pp.

Isopoda

- Birshtein, J. A. 1951 *Asellota. Crustaceans, VII (5)*. Fauna SSSR (novaia serii). 47. Leningrad. 143 pp.
- Bourdon, R. 1968 Les Bopyridae des mer européennes. *Mémoires du Muséum national d'Histoire naturelle. Nouvelle Série. Série A, Zoologie* 50: 77-424.
- Gruner, H. E. 1965 *Krebstiere oder Crustacea. V. Isopoda I*. Die Tierwelt Deutschlands. 51. Jena: G. Fischer Verlag. 149 pp.
- Gruner, H. E. 1966 *Krebstiere oder Crustacea. V. Isopoda 2*. Die Tierwelt Deutschlands. 53. Jena: Gustav Fischer Verlag. 380 pp.
- Guryanova, E. F. 1932 *Marine Arctic Isopoda*. Opredeliteli po faune SSSR. 4a. Leningrad: izdavaemye Zoologicheskim muzeem Akademii Nauk SSSR. 181 pp.
- Holthuis, L. B. 1956 *Isopoda en Tanaidacea (KV)*. Fauna van Nederland. 16. Leiden: A.W. Sijthoff. 280 pp.
- Huwae, P. H. M. 1977 *De isopoden van de Nederlandse kust*. Wetenschappelijke mededelingen KNNV. 118. Hoogwoud: Koninklijke Nederlandse Natuurhistorische Vereniging.
- Kusakin, O. G. 1979 *[Marine and brackish-water isopod crustaceans (Isopoda) of cold and temperate waters of the northern hemisphere. Vol.1. Suborder Flabellifera]* [in Russian]. Opredeliteli po faune SSSR. 122. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 479 pp.
- Kusakin, O. G. 1982 *[Marine and salt water isopod crustaceans (Isopoda) of cold and temperate waters of the Northern Hemisphere. Vol.2. Suborders Anthuridea,*

Microcerberidae, Valvifera, Tyloidea] [in Russian]. Opredeliteli po faune SSSR. 131. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 456 pp.

Kusakin, O. G. 1988 *[Marine and brackish isopods (Isopoda) of cold and temperate waters of the northern hemisphere. Volume 3. Suborder Asellota]* [in Russian]. Opredeliteli po faune SSSR. 152. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 500 pp.

Monod, T. 1923 Prodrome d'une faune des Tanaidacea et des Isopoda (excl. Epicaridea) des côtes de France (excl. Méditerranée). *Annales de la Société des Sciences Naturelles de Charente inférieure* 37(4): 19-126.

Naylor, E. 1972 *British marine isopods. Keys and notes for the identification of the species. Synopses of the British fauna (New Series)*. 3. London: Academic Press. 86 pp.

Stephensen, K. & Meinertz, N. T. 1948 *Storkrebs IV. Ringkrebs 3. Tanglus (Marine Isopoder og Tanaider)*. Danmarks Fauna. 53. 187 pp.

INSECTA

Freund, L. 1927 *Anoplura Pinnipediorum (Robbenläuse)*. Tierwelt der Nord- und Ostsee. 11d1. Leipzig: Akademische verlagsgesellschaft. 36 pp.

Hennig, W. 1935 *Thalassobionte und thalassophile Diptera Nematocera*. Tierwelt der Nord- und Ostsee. 11e3. Leipzig: Akademische verlagsgesellschaft. Pp. 85-102.

Karl, O. 1930 *Thalassobionte und thalassophile Diptera Brachycera*. Tierwelt der Nord- und Ostsee. 11e2. Leipzig: Akademische verlagsgesellschaft. Pp. 33-84.

Lengerken von, H. 1929 *Halophile und halobionte Coleoptera*. Tierwelt der Nord- und Ostsee. Lfg. 16. Leipzig: Akademische verlagsgesellschaft. Pp. 30-32.

Lindberg, H. 1935 *Halophile und halobionte Hemiptera*. Tierwelt der Nord- und Ostsee. 11e4. Leipzig: Akademische verlagsgesellschaft. Pp. 103-124.

Strenzke, K. 1955 *Collembola*. Tierwelt der Nord- und Ostsee. 11f2. Leipzig: Akademische verlagsgesellschaft. 52 pp.

PHORONIDA

Cori, C. J., Borg, F., Bückmann, A. & Michaelsen, W. 1930 *Muschellinge oder Molluscoidea und Manteltiere oder Tunicata. Kamptazoa Cori* (synonym Entoprocta Nitsche). Phoronidea. Moostierchen oder Bryozoa (Ectoprocta). Manteltiere oder Tunicata. Seescheiden oder Ascidiæ. Die Tierwelt Deutschlands. 17. Jena: G. Fischer. 188 pp.

Emig, C. C. 1979 *British and other phoronids. Keys and notes for the identification of the species. Synopses of the British Fauna (New Series)*. 13. London: Academic Press. 57 pp.

Selys-Longchamps de, M. 1907 *Phoronis*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 30. Berlin: R. Friedländer & Sohn. 280 pp.

Viéitez, J. M., Emig, C. C., Rodríguez Pablo, C. & García Carrascosa, A. M. 1987 Foronídeos de las costas de la Península Ibérica: Islas Baleares: sistemática, ecología, distribución y estados larvarios. Boletín del Instituto Español de Oceanografía 4(2): 63-78.

BRYOZOA

Cori, C. J., Borg, F., Bückmann, A. & Michaelsen, W. 1930 *Muschellinge oder Molluscoidea und Manteltiere oder Tunicata. Kamptazoa Cori* (synonym Entoprocta Nitsche). Phoronidea. Moostierchen oder Bryozoa (Ectoprocta). Manteltiere oder Tunicata. Seescheiden oder Ascidiæ. Die Tierwelt Deutschlands. 17. Jena: G. Fischer. 188 pp.

d'Hondt, J.-L. 1983 Tabular keys for identification of the recent ctenostomatous Bryozoa. Mémoires de l'Institut Océanographique de Monaco 14: 1-134.

Harmelin, J. G. 1976 *Le sous-ordre des Tubuliporina (Bryozoaires, Cyclostomes) en Méditerranée : écologie et systématique. Mémoires de l'Institut Océanographique de Monaco* 10: 1-326.

- Hayward, P. J.** 1985 Ctenostome bryozoans. Keys and notes for the identification of the species. *Synopses of the British Fauna (New Series)*. 33. London: E.J. Brill/Dr. W. Backhuys. 169 pp.
- Hayward, P. J. & Ryland, J. S.** 1979 British ascophoran bryozoans. Keys and notes for the identification of the species. *Synopses of the British Fauna (New Series)*. 14. London: Academic Press. 312 pp.
- Hayward, P. J. & Ryland, J. S.** 1985 Cyclostome bryozoans. Keys and notes for the identification of the species. *Synopses of the British Fauna (New Series)*. 34. London: E.J. Brill/Dr. W. Backhuys. 147 pp.
- Hayward, P. J. & Ryland, J. S.** 1998 Cheilostomatous Bryozoa. Part 1. Aeteoidea, Cribriolinae. *Synopses of the British Fauna (New Series)*. 10. Shrewsbury: Field Studies Council. viii+366 pp.
- Klyuge, G. A.** 1962 Bryozoans of the northern seas of the USSR. Opredelitel po faune SSSR. 76. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 584 pp.
- Lacourt, A. W.** 1978 De Nederlandse mariene mosdiertjes - Bryozoa (Entoprocta, Gymnolaemata). Wetenschappelijke mededelingen KNNV. 129. Hoogwoud: Koninklijke Nederlandse Natuurhistorische Vereniging. 21 pp.
- Lohmann, H.** 1911 Die Cyphonautes des nordischen Meeres. Nordisches Plankton 13: 31-40.
- Marcus, E.** 1933 Bryozoa. Tierwelt der Nord- und Ostsee. 7cl. Leipzig: Akademische verlagsgesellschaft. 100 pp.
- Marcus, E.** 1940 Mosdry (Bryozoa eller Polyzoa). Danmarks Fauna. 46. Kobenhavn: GEC, Gad. 401 pp.
- Nordgaard, O.** 1918 Bryozoa from the arctic regions. Tromsø Museums Aarshefte 1917 40(1): 1-99.
- Occipinti Ambrogi, A.** 1981 Brizoi lagunari. Guida per il riconoscimento delle specie animali delle acque lagunari e costiere italiane. 7. Consiglio Nazionale delle Ricerche. 147 pp.
- Prenant, M. & Bobin, G.** 1956 Bryozoaires. 1ère Partie. Entoproctes, Phylactolèmes, Ctenostomes. Faune de France. 60. Paris: P. Lechevalier. 398 pp.
- Prenant, M. & Bobin, G.** 1966 Bryozoaires. 2ème Partie. Chilostomes Anasca. Faune de France. 68. Paris: P. Lechevalier. 647 pp.
- Ryland, J. S.** 1974 A revised key for the identification of British intertidal Bryozoa. Field Studies 4: 77-86.
- Ryland, J. S. & Hayward, P. J.** 1977 British anascan bryozoans: Cheilostomata, Anasca. *Synopses of the British Fauna (New Series)*. 10. London, New-York: Academic Press. 188 pp.
- Zabala, M. & Maluquer, P.** 1988 Illustrated keys for the classification of Mediterranean Bryozoa. Treballs del Museu de Zoologia. 4. Barcelona: Museu de Zoologia. 294 pp.

BRACIOPODA

- Brunton, C. H. C. & Curry, G. B.** 1979 British brachiopods. Keys and notes for the identification of the species. *Synopses of the British Fauna (New Series)*. 17. London: Academic Press. 64 pp.
- Cori, C. J.** 1933 *Brachiopoda*. Tierwelt der Nord- und Ostsee. 7c3. Leipzig: Akademische verlagsgesellschaft. Pp. 133-150.
- Logan, A.** 1979 The recent Brachiopoda of the Mediterranean sea. *Bulletin de l'Institut Océanographique de Monaco* 72: 1-112.
- Nobre, A.** 1932 *Brachiotropes de Portugal*. Fauna marinha de Portugal. 6. Porto: Imprensa Portuguesa. 20 pp.

CHAETOGNATHA

- Grassi, B.** 1883 *I chetognati, anatomia e sistematica con aggiunte embriologiche*. Fauna und flora des Golfes von Neapel und der Angrenzenden Meeres-Abschnitte. 5. Leipzig: W. Engelmann. 126 pp.
- Kramp, P. L.** 1938 *Chaetognatha*. Zoology Iceland. 4(71). Copenhagen, Reykjavik: Levin & Munksgaard. 4 pp.

- Kuhl, W.** 1928 *Chaetognatha*. Tierwelt der Nord- und Ostsee. 7b. Leipzig: Akademische verlagsgesellschaft. 24 pp.
- Pierrot-Bults, A. C. & Chidgey, K. C.** 1988 *Chaetognatha. Synopses of the British Fauna (New Series)*. 39. Leiden, New-York. 66 pp.

HEMICORDATA

- Horst van der, C. J.** 1927 *Enteropneusta*. Tierwelt der Nord- und Ostsee. 7a1. Leipzig: Akademische verlagsgesellschaft. 12 pp.
- Horst van der, C. J.** 1927 *Pterobranchia*. Tierwelt der Nord- und Ostsee. 7a2. Leipzig: Akademische verlagsgesellschaft. Pp. 13-20.
- Spengel, J. W.** 1893 *Die Enteropneusten des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte*. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 18. Berlin: R. Friedländer & Sohn. 756 pp.

ECHINODERMATA

- Arndt, W., Broch, H., Krumbach, T., Pax, F. & Leiberkind, J.** 1928 *Porifera, Schwämme, Spongiens. Hydrozoen. Scyphozoa. Anthozoa. Ctenophora, Kammmallen oder Rippenquallen. Echinodermata, Stachelhäuter oder Echinodermen. Die Tierwelt Deutschlands*. 4. Jena: G. Fischer. 332 pp.
- Clark, A. M.** 1970 Echinodermata Crinoidea. Marine invertebrates of Scandinavia. 3. Oslo: Universitetsforlaget. 55 pp.
- Clark, A. M. & Downey, M. E.** 1992 Starfishes of the Atlantic. Natural History Museum Publications. Chapman and Hall Identification Guides. 3. London: Chapman and Hall. xxvi+794 pp.
- Clark, A. M. & Downey, M. E.** 1997 Starfishes of the Atlantic. Amsterdam: ETI, Netherlands CD (Mac) Date TBA CD (Windows). CD-ROM.
- Demetropoulos, A. & Hadjichristophorou, M.** 1976 Echinodermata of Cyprus. Fisheries Bulletin 4: 83 pp.
- Dyakonov, A. M.** 1933 Echinoderms of the northern seas. Opredelitel po faune SSSR. 8. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 166 pp.
- Dyakonov, A. M.** 1950 Sea stars in the seas of the USSR. Opredelitel po faune SSSR. 34. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 203 pp.
- Dyakonov, A. M.** 1954 Brittle-stars in the seas of the USSR. Opredelitel po faune SSSR. 55. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 136 pp.
- Engel, H.** 1932 Echinodermata. Fauna van Nederland. 6. Leiden: A.W. Sijthoff. 91 pp.
- Grieg, J. A.** 1900 Die Ophiuriden der Arktis. Fauna Arctica 1: 259-286.
- Koehler, R.** 1921 Echinoderms. Faune de France. 1. Paris: P. Lechevalier. 210 pp.
- Koehler, R.** 1924 Les échinodermes des mers d'Europe. 1. Paris: G. Douin. 362 pp.
- Koehler, R.** 1927 Les échinodermes des mers d'Europe. 2. Paris: G. Douin. 339 pp.
- Ludwig, H.** 1897 Die seesterne des Mittelmeeres. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 24. Berlin: R. Friedländer & Sohn. 491 pp.
- Madsen, F. J. & Hansen, B.** 1994 Echinodermata Holothuroidea. Marine invertebrates of Scandinavia. 9: Scandinavian University Press, Norway. 143 pp.
- Moreno, I. & Munar, J.** 1985 Equinoideos mediterráneos. Claves para la identificación de la fauna española. Monografías de la Universidad de Palma de Mallorca. 34 pp.
- Moreno, I. & Munar, J.** 1985 Osiuroideoes mediterráneos. Claves para la identificación de la fauna española. Monografías de la Universidad de Palma de Mallorca. 34 pp.
- Moreno, I. & Munar, J.** 1985 Astroideoes y Crinoideoes mediterráneos. Claves para la identificación de la fauna española. Monografías de la Universidad de Palma de Mallorca. 34 pp.

- Mortensen, T. 1924 Pighude (Echinodermer). Danmarks Fauna. 27. København: GEC Gad. 274 pp.
- Mortensen, T. 1927 Handbook of the echinoderms of the British Isles. Oxford: Oxford University Press. ix+471 pp.**
- Mortensen, T. 1928 Echinoderma. Tierwelt der Nord- und Ostsee. 8. Leipzig: Akademische verlagsgesellschaft. 128 pp.
- Nobre, A. 1931 Echinodermes de Portugal. Fauna Marinha de Portugal. 2. Porto: Instituto de Zoologia da Universidade do Porto. 176 pp.
- Pancucci-Papadopoulou, M.-A. 1996 The Echinodermata of Greece. Fauna Graeciae. 6. Athens: Hellenic Zoological Society. 162 pp.
- Paterson, G. L. J. 1985 The deep-sea Ophiuroidea of the North Atlantic ocean. Bulletin of the British Museum (Natural History) 49(1): 1-162.
- Pickton, B. E. 1986 The echinoderms of the British Isles; a colour guide. Ross-on-Wye: Marine Conservation Society. 75 pp.
- Pickton, B. E. 1993 A field guide to the shallow-water echinoderms of the British Isles. London: Immel Publishing. 96 pp.
- Southward, E. C. 1972 Keys for the identification of Echinodermata of the British Isles. Plymouth: Marine Biological Association
- Tortonese, E. 1965 Echinodermata. Fauna d'Italia. 6. Bologna: Edizioni Calderini. 424 pp.

TUNICATA

General and other than Ascidiacea

- Berrill, N. J. 1950 The Tunicata with an account of the British species. London: Ray Society. 354 pp.
- Bone, Q. (ed.) 1998 The Biology of Pelagic Tunicates. New York: Oxford University Press. 354 pp.
- Bückmann, A. 1926 Tunicata (Urochordata): Copepoda. Tierwelt der Nord- und Ostsee. 12a1. Leipzig: Akademische verlagsgesellschaft. 1-20 pp.
- Cori, J. C., Borg, F., Bückmann, A. & Michaelsen, W. 1930 Muschellinge oder Molluscoidea und Manteltiere oder Tunicata. Kamptozoa Cori (synonym Entoprocta Nitsche). Phoronidea. Moostierchen oder Bryozoa (Ectoprocta). Manteltiere oder Tunicata. Seescheiden oder Ascidae. Die Tierwelt Deutschlands. 17. Jena: G. Fischer. 188 pp.
- Fenau, R. 1966 Les Appendiculaires des mers d'Europe et du bassin Méditerranéen. Faune de l'Europe et du bassin Méditerranéen. 2. Paris: Masson & Cie. 117 pp.
- Fraser, J. H. 1981 British pelagic tunicates. Keys and notes for the identification of species. Synopses of the British Fauna (New Series). 20. London: Cambridge University Press. 57 pp.
- Harant, H. & Vernières, P. 1938 Tuniciers. Fasc. 2 : Appendiculaires et Thaliacés. Faune de France. 33. Paris: P. Lechevalier. 59 pp.
- Ihle, J. E. W. 1927 Thaliacea. Tierwelt der Nord- und Ostsee. 12a2. Leipzig: Akademische verlagsgesellschaft. Pp. 21-48.
- Ulanin, B. 1884 Die arten der gattung Doliolum im des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 10. Leipzig: W. Engelmann. 140 pp.

Ascidiacea

- Harant, H. & Vernières, P. 1933 Tuniciers. Fasc. 1 : Ascidiées. Faune de France. 27. Paris: P. Lechevalier. 101 pp.
- Huus, J. 1933 Ascidiacea. Tierwelt der Nord- und Ostsee. 12a3. Leipzig: Akademische verlagsgesellschaft. Pp. 49-115.
- Lafargue, F. 1968 Peuplements sessiles de l'Archipel des Glénans. Didemnidae (Ascidiées). Vie et Milieu 19(2): 353-446.
- Lafargue, F. & Wahl, M. 1987 The didemnid ascidian fauna of France. Annales de l'Institut océanographique de Monaco 63: 1-46.

- Lützen, J. 1967 Sækdyr. Danmarks Fauna. 75. København: GEC Gads. 267 pp.
- Millar, R. H. 1966 *Tunicata Ascidiacea. Marine invertebrates of Scandinavia. 1.* Oslo: Scandinavia University Books. 123 pp.
- Millar, R. H. 1970 *British ascidians. Tunicata: Ascidiacea. Keys and notes for the identification of the species. Synopses of the British Fauna. 1.* London: Academic Press for the Linnean Society of London. 99 pp.
- Monniot, C. 1965 Etude systématique et évolutive de la famille des Pyuridae (Ascidiacea). *Mémoires du Muséum national d'Histoire naturelle* 36: 1-203.
- Monniot, C. 1970 Les Molgulidae des mers européennes. *Mémoires du Muséum national d'Histoire naturelle, série 2* 41(5): 1131-1145.
- Monniot, C. 1975 ["1974"] Ascidiées littorales et bathyales récoltées au cours de la campagne Biaçores : Phlebobranches et Stolidobranches. *Bulletin du Muséum national d'Histoire naturelle, série 3* 251: 1327-1352.
- Monniot, C. & Monniot, F. 1973 Clé mondiale des genres d'ascidies. *Archives de Zoologie Expérimentale et Générale* 113(3): 311-367.
- Monniot, C. & Monniot, F. 1973 Ascidiées abyssales récoltées au cours de la campagne océanographique Biaçores par le "Jean Charcot". *Bulletin du Muséum national d'Histoire naturelle, série 3* 121: 389-475.
- Monniot, F. 1974 Ascidiées intersticielles des côtes d'Europe. *Mémoires du Muséum national d'Histoire naturelle, série A* 35: 1-154.
- Monniot, F. 1975 ["1974"] Ascidiées littorales et bathyales récoltées au cours de la campagne Biaçores : Aplousobranches. *Bulletin du Muséum national d'Histoire naturelle, série 3* 251: 1287-1325.
- Pickton, B. E. 1985 *Ascidians of the British Isles. A colour guide.* Ross-on-Wye: Marine Conservation Society
- Ramos Espla, A. A. 1991 *Ascidiás littorales del Mediterraneo ibérico. Faunística, ecología y biogeografía.* Universidad de Alicante. 405 pp.
- Romanov, V. N. 1989 *Kolonial'nye ascidií semejstva Didemnidae morej SSSR i sopredel'nyh vod [Colonial ascidiants of the family Didemnidae from the seas of the USSR and adjacent waters].* Fauna SSSR (novaia seriiia). 138. Leningrad: Akademiiia nauk SSSR, Zoologicheskii institut. 224 pp.
- Tursi, A. 1980 *Ascidiacei.* Guide per il riconoscimento delle specie animali delle acque lagunari e costiere italiane. 4. Genova: Consiglio Nazionale delle Ricerche. 85 pp.
- Vazquez-Otero, M. E. 1993 *Estudio faunístico autoecológico y biogeográfico de los tunicatos bentónicos de la Ría de Ferrol (Galicia).* Santiago de Compostella: Universidad de Santiago de Compostella. 443 pp.

CEPHALOCHORDATA

- Franz, V. 1927 *Branchiostoma.* Tierwelt der Nord- und Ostsee. 12b. Leipzig: Akademische verlagsgesellschaft. 46 pp.

VERTEBRATA

General

- Fischer, W., Schneider, M. & Bauchot, M.-L. 1987 Fiches FAO d'identification des espèces pour les besoins de la pêche. Méditerranée et Mer Noire, Zone de pêche 37. Volume 2. Vertébrés. Rome, Bruxelles: FAO/CEE. Pp. 761-1529.
- Martín Ferrero, M. P. 1979 Vertebrados marinos españoles (clase taxonómica). Cádiz: El autor. 212 pp.
- Nobre, A. 1935 Vertebrados (mamíferos, reptis e peixes). Fauna marinha de Portugal. I. Porto: Companhia Editoria do Minho. lxxxiv+21+5+574 pp.

Pisces

- Albuquerque, R. M. 1951-1956 Peixes de Portugal e ilhas adjacentes. Chavas para a sua determinação. *Portugaliae Acta Biologica, ser. B* 5: i-xvi+1-1167.

- Andriyashev, A. P. 1954 *Fishes of northern seas of the USSR.* Opredeliti po faune SSSR. 53. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 567 pp.
- Banarescu, P. 1964 *Pisces. Osteichthyes.* Fauna Republicii Socialiste România. 13. Bucuresti. 962 pp.
- Banarescu, P. 1969 *Cyclostomata si Chondrichthyes (Cyclostomi si Selaciensi).* Fauna Republicii Socialiste România. 12(1). Bucuresti: Editura Academiei Republicii Socialiste România. 102 pp.
- Barsukov, V. V. 1959 *Fam. Anarhichadidae. Fishes, V (5).* Fauna SSSR (novaia serii). 73. Leningrad: Akademiia nauk SSSR, Zoologicheskii institut. 174 pp.
- Batjakas, I. E. & Economakis, A. E. 1995 *Coastal Fishes of Greece.* Anixis, Attikes: Efsthadiadis Group. 131 pp.
- Bauchot, M. L. & Pras, A. 1980 *Guide des poissons marins d'Europe.* Neuchatel: Delachaux & Niestlé. 427 pp.
- Bauchot, M. L. & Pras, A. 1982 *Guia de los peces de mar de España y de Europa.* Barcelona: Omega. 432 pp.
- Bertolini, F., D'Ancona, U., Montalenti, G., Padoa, E., Ranzi, S., Sanzo, L., Sparta, A., Tortonese, E. & Vialli, M. 1973 *Uova, larve e stadi giovanili di Teleostei.* Fauna e flora del Golfo di Napoli. 38. Koenigstein: Otto Koeltz Antiquariat. 1064 pp.
- Bini, G. 1967 *Atlante dei pesci delle coste italiane. I Leptocardi, cyclostomi, selaci.* Milano: Mondo Sommerso. 206 pp.
- Bini, G. 1968 *Atlante dei pesci delle coste italiane. 5 Perciformi (percoidei).* Milano: Mondo Sommerso. 175 pp.
- Bini, G. 1968 *Atlante dei pesci delle coste italiane. 6 Perciformi (trichiuroidei...blennioidei).* Milano: Mondo Sommerso. 177 pp.
- Bini, G. 1968 *Atlante dei pesci delle coste italiane. 8 Pleuronettiformi, echeniformi, gobioesociformi, tetraodontiformi, loliformi.* Milano: Mondo Sommerso. 164 pp.
- Bini, G. 1968 *Atlante dei pesci delle coste italiane. 4 Perciformi (mugiloidei, percoidei).* Milano: Mondo Sommerso. 163 pp.
- Bini, G. 1969 *Atlante dei pesci delle coste italiane. 7 Perciformi (ofidioidei...dactylopteroidei).* Milano: Mondo Sommerso. 196 pp.
- Bini, G. 1970 *Atlante dei pesci delle coste italiane. 3 Notacantiformi...Zeiformi.* Milano: Mondo Sommerso. 229 pp.
- Bini, G. 1971 *Atlante dei pesci delle coste italiane. 2 Osteilli (aciopenseriformi, clupeiformi, siformi, anguilliformi).* Milano: Mondo Sommerso. 300 pp.
- Bini, G. 1972 *Atlante dei pesci delle coste italiane. 9 Introduzione. Parte generale. Aggiornamenti. Indici.* Milano: Mondo Sommerso. 176 pp.
- Cottiglia, M. 1980 *Pesci lagunari.* Guida per il riconoscimento delle specie animali delle acque lagunari e costiere Italiane. 1. Genova: Consiglio Nazionale delle Ricerche. 141 pp.
- Debelius, H. 1997 *Mediterranean and Atlantic Fish Guide. From Spain to Turkey, From Norway to South Africa.* Frankfurt: INKAN-Unterwasserarchiv. 305 pp.
- Demetropoulos, A. & Neocleous, D. 1969 The Fishes and Crustaceans of Cyprus. *Fisheries Bulletin* 1: 21 pp.
- Dipper, F. 1987 *British sea fishes.* London: Underwater World Publications. 194 pp.
- Drensky, P. 1951 *Ribite v Bulgariia.* Fauna na Bulgariia. 2. Sofia: Bulgarskata akademia na naukite. 270 pp.
- Duncker, G. *Teleostei Physoclisti: Scleroparei.* Tierwelt der Nord- und Ostsee. 12h2. Leipzig: Akademische verlagsgesellschaft. Pp. 61-84.
- Duncker, G. 1928 *Teleostei Physoclisti: Blenniformes.* Tierwelt der Nord- und Ostsee. 12g3. Leipzig: Akademische verlagsgesellschaft. Pp. 89-106.
- Duncker, G. 1928 *Teleostei Physoclisti: Gobiiformes.* Tierwelt der Nord- und Ostsee. 12g3. Leipzig: Akademische verlagsgesellschaft. Pp. 121-148.
- Ehrenbaum, E. 1927 *Chondrostei.* Tierwelt der Nord- und Ostsee. 12f1. Leipzig: Akademische verlagsgesellschaft. 8 pp.
- Ehrenbaum, E. 1927 *Elasmobranchii.* Tierwelt der Nord- und Ostsee. 12e. Leipzig: Akademische verlagsgesellschaft. 66 pp.
- Emery, C. 1880 *Le specie del genere Fierasfer nel Golfo di Napoli e regioni limitrofe.* Fauna und flora des Golfs von Neapel und der Angrenzenden Meeres-Abschnitte. 2. Leipzig: W. Engelmann. 76 pp.
- Fischer, W., Schneider, M. & Bauchot, M.-L. 1987 *Fiches FAO d'identification des espèces pour les besoins de la pêche. Méditerranée et Mer Noire, Zone de pêche 37. Volume 2. Vertébrés.* Rome, Bruxelles: FAO/CEE. Pp. 761-1529.
- Hartman, P., Groot & Edgell. 1997 *Field identification of coastal juvenile salmonids.* The Field Naturalist. 32 pp.
- Hureau, J.-C. 1994 *Fishes of the north eastern Atlantic and the Mediterranean.* Amsterdam, Paris: ETI/UNESCO. CD-ROM Macintosh.
- Hureau, J.-C. 1996 *Fishes of the north eastern Atlantic and the Mediterranean.* Amsterdam, Paris: ETI/UNESCO. CD-ROM Windows.
- Jennings, G. (in press) *Aegean, Greek and Eastern Mediterranean fishes: an identification guide.* Fishes of the Mediterranean Series. 250 pp.
- Jennings, G. (ed.) 1996 *Mediterranean gobies, blennies, wrasses and shore fishes. An identification guide to coastal and shallow water species.* Fishes of the Mediterranean Series.
- Jennings, G. (ed.) 1996 *Italian sea fishes. A guide to all recorded species of Italian and Adriatic marine fishes.* Fishes of the Mediterranean Series.
- Jennings, G. (ed.) 1996 *Maltese fishes. A guide to the fishes of the Maltese islands.* Fishes of the Mediterranean Series.
- Jennings, G. (ed.) 1996 *Sea fishes of the Spanish Mediterranean coast and islands.* Fishes of the Mediterranean Series.
- Jennings, G. 1996 *European Sea Fishes, The Pocketbook, Gibraltar to Norway. An Illustrated Identification Guide to Recorded Species.* European sea fishes. London: Calypso. 206 pp.
- Jennings, G. H. 1997 *Mediterranean Fishes: An Illustrated Identification Guide.* The Pocketbook. London: Calypso. 235 pp.
- Jonsson, G. 1992 *Íslenskir fiskar.* Reykjavik: Fjöldi. 568 pp.
- Kaspiris, P. 1986 *[Identification Key for Greek Fish] (in Greek).* Patras: Patras University, Section of Animal Zoology. 123 pp.
- Kyle, H. M. & Ehrenbaum, E. 1926 *Pisces, Allgemeiner Teil.* Tierwelt der Nord- und Ostsee. 12c. Leipzig: Akademische verlagsgesellschaft. 104 pp.
- Kyle, H. M. & Ehrenbaum, E. 1927 *Teleostei Physostomi: Clupeiformes, Apodes, Esociformes, Ostariophys.* Tierwelt der Nord- und Ostsec. 12f2. Leipzig: Akademische verlagsgesellschaft. Pp. 9-86.
- Lanfranco, G. G. 1996 *The fish around Malta: central Mediterranean.* Malta: Progress Press Co. xi+132 pp.
- Lozano y Rey, L. 1947 *Peces ganoides y fisostomos. Memorias de la Academia de Ciencias exactas, físicas y naturales de Madrid Serie Ciencias naturales 11: i-xv+1-839.*
- Lozano y Rey, L. 1952 *Peces fisoclistos, subserie toracicos. Segunda parte. Memorias de la Academia de Ciencias exactas, físicas y naturales de Madrid Serie Ciencias naturales 14: 379-705.*
- Lozano y Rey, L. 1952 *Peces fisoclistos, subserie toracicos. Primera parte. Memorias de la Academia de Ciencias exactas, físicas y naturales de Madrid Serie Ciencias naturales 14: i-xv+1-378.*
- Lozano y Rey, L. 1960 *Peces fisoclistos. Tercera parte subserie toracicos (ordenes equeniformes y gobiformes), pediculados y asimetricos. Memorias de la Academia de Ciencias exactas, físicas y naturales de Madrid Serie Ciencias naturales 14: 1-613.*
- Lythgoe, J. & Lythgoe, G. 1991 *Fishes of the sea. The north Atlantic and Mediterranean.* London: Blandford Press. 256 pp.

- Miller, P. & Loates, M. J. 1997 *Fish of Britain and Europe*. Collins Pocket Guide. The Field Naturalist. London: Harper Collins. 288 pp.
- Mohr, E. 1926 *Teleostei Physoclisti: Scombrids, Plectognathi, Ammodytiformes, Atheriniformes*. Tierwelt der Nord- und Ostsee. 12g1. Leipzig: Akademische verlagsgesellschaft. 44 pp.
- Mohr, E. 1927 *Teleostei Physoclisti: Labridae, Carangidae, Berycidae, Percidae, Scombridae*. Tierwelt der Nord- und Ostsee. 12h2. Leipzig: Akademische verlagsgesellschaft. Pp. 85-140.
- Mohr, E. 1928 *Teleostei Physoclisti: Trachinidae*. Tierwelt der Nord- und Ostsee. 12g3. Leipzig: Akademische verlagsgesellschaft. Pp. 107-120.
- Moreno, J. A. 1995 *Guía de los tiburones de aguas ibéricas, del Atlántico Nororiental y del Mediterráneo*: Pirámide, Spain. 310 pp.
- Muus, B. J. 1964 *Havfisk og fiskeri i Nordvesteuropa*. København: Gads forlag. 244 pp.
- Nijssen, H. & Groot, S. J. 1980 *Zeevissen van de Nederlandse kust*. Wetenschappelijke mededelingen KNNV. 143. Hoogwoud: Koninklijke Nederlandse Natuurhistorische Vereniging. 109 pp.
- Nobre, A. 1935 *Vertebrados (mamíferos, réptis e peixes)*. Fauna marinha de Portugal. 1. Porto: Companhia Editoria do Minho. Ixxxiv+21+5+574 pp.
- Notarbartolo di Sciara, G. & Bianchi, I. 1998 *Guida degli squali e delle razze del Mediterraneo*. 388 pp.
- Papakonstantinou, C. 1988 *Check-list of marine fishes of Greece*. Fauna Graeciae. 4. Athens: Hellenic Zoological Society. 257 pp.
- Pethon, P. 1994 *Aschehougs stor Fiskebok*. Oslo: Aschehoug forlag. 447 pp.
- Pipitone, C., Massari, F. & Thomas, M. 1995 *I pesci delle acque costiere italiane*. IPZS: L'Epos. 114 pp.
- Poll, M. 1947 *Poissons marins*. Faune de Belgique. Bruxelles: Musée Royal d'Histoire Naturelle de Belgique. 452 pp.
- Rauthier, M. 1925 *Die Syngnathiden des Golfs von Neapel*. Fauna e flora del Golfo di Napoli. 36. Berlin: R. Friedländer & Sohn. 365 pp.
- Redeke, H. C. 1948 *Pisces (Cyclostomi-Euichthyes) (I-I)*. Fauna van Nederland. 10. Leiden: A.W. Sijthoff. 331 pp.
- Russell, F. S. 1976 *The eggs and planktonic stages of British marine fishes*. London: Academic Press. xv+524 pp.
- Saemundsson, B. 1949 *Marine Pisces*. *Zoology of Iceland* 72(4): 1-50.
- Sanches, J. G. 1985 *Nomenclatura e diagnose dos principais peixes marinhos de Portugal: Cyclostomos, Selaceos e Holocefatos*. Publicações avulsas. 9. Lisbon: Instituto Nacional de Investigação das Pescas. 184 pp.
- Sanches, J. G. 1992 *Guia para identificação do pescado de Portugal submetido a tamanho mínimo de captura*. Publicações avulsas. 18. Lisbon: Instituto Nacional de Investigação das Pescas. 272 pp.
- Schnakenbeck, W. 1925 *Teleostei Physoclisti: Heterosomatidae*. Tierwelt der Nord- und Ostsee. 12h1. Leipzig: Akademische verlagsgesellschaft. 60 pp.
- Schnakenbeck, W. 1926 *Teleostei Physoclisti: Gadidae*. Tierwelt der Nord- und Ostsee. 12g2. Leipzig: Akademische verlagsgesellschaft. Pp. 45-88.
- Schnakenbeck, W. 1927 *Cyclostomi*. Tierwelt der Nord- und Ostsee. 12d. Leipzig: Akademische verlagsgesellschaft. 14 pp.
- Svetovidov, A. N. 1936 *Fam. Triglidae. Fishes, VI (9)*. Fauna SSSR (novaia seriia). 2. Leningrad: Akademiiia nauk SSSR, Zoologicheskii institut. 24 pp.
- Svetovidov, A. N. 1948 *Treskoobraznye*. Fauna SSSR (novaia seriia). 34. Moskva: Akademiiia nauk SSSR, Zoologicheskii institut. 221 pp.
- Svetovidov, A. N. 1963 *Fishes (Ryby)*. Volume 2, number 1 : *Clupeidae (Sel'devye)*. Fauna SSSR (novaia seriia). 48. Jerusalem: Israel Program for Scientific Translations. 428 pp.
- Svetovidov, A. N. 1964 *Fishes of the Black Sea*. Opredeliteli po faune SSSR. 86. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 552 pp.
- Tortonese, E. 1956 *Leptocardia, Cyclostoma, Selachii*. Fauna d'Italia. 2. Bologna: Edizioni Calderini. 334 pp.
- Tortonese, E. 1970 *Osteichthyes. Pesci ossei (parte 1)*. Fauna d'Italia. 10. Bologna: Edizioni Calderini. 565 pp.
- Tortonese, E. 1975 *Osteichthyes. Pesci ossei (parte 2)*. Fauna d'Italia. 11. Bologna: Edizioni Calderini. 636 pp.
- Vas, P. 1991 *A field guide to the sharks of the British coastal waters*. AIDGAP Guides. Shrewsbury: Field Studies Council. 36 pp.
- Wheeler, A. 1969 *The fishes of the British Islands and North West Europe*. London. 613 pp.
- Wheeler, A. 1978 *Key to the fishes of northern Europe*. London: Warne. xix+380 pp.
- Wheeler, A. & Newman, C. 1992 *The Pocket Guide to the Saltwater Fish of Britain and Europe*. 176 pp.
- Whitehead, P. J. P., Bauchot, M. L., Hureau, J. C., Nielsen, J. & Tortonese, E. 1986 *Fishes of the North-Eastern Atlantic and the Mediterranean*. 2. Paris: UNESCO. Pp. 521-1010.
- Whitehead, P. J. P., Bauchot, M. L., Hureau, J. C., Nielsen, J. & Tortonese, E. 1986 *Fishes of the North-Eastern Atlantic and the Mediterranean*. 3. Paris: UNESCO. Pp. 1010-1458.
- Whitehead, P. J. P., Bauchot, M. L., Hureau, J. C., Nielsen, J. & Tortonese, E. 1989 *Fishes of the North-Eastern Atlantic and the Mediterranean*. 1. Paris: UNESCO. 520 pp.
- Zyuganov, V. V. 1991 *Gasterosteidae of the world fauna. Fishes, V (5)*. Fauna SSSR (novaia seriia). 137. Akademiiia nauk SSSR, Zoologicheskii institut. 261 pp.

Tetrapoda

Reptilia

- Brongersma, L. D. 1967 *Guide for the identification of stranded turtles on British coasts*. Trustees of the British Museum (Natural History) 659. London: British Museum (Natural History). viii+23 pp.
- Brongersma, L. D. 1972 European Atlantic turtles. *Zoologische Verhandelingen* 121: 1-318.
- Casado, M. F. & et al. 1997 *Mamíferos marinos y tortugas del Mar de Alborán : guía d'especies marinas amenazadas*. Sevilla: Consejería de Medio Ambiente. 119 pp.
- Nobre, A. 1935 *Vertebrados (mamíferos, réptis e peixes)*. Fauna marinha de Portugal. 1. Porto: Companhia Editoria do Minho. Ixxxiv+21+5+574 pp.

Aves

- Dif, G. 1982 *Les oiseaux de mer d'Europe*. Paris: Arthaud. 445 pp.
- Grant, P. J. 1982 *Gulls. A guide to identification*. Calton: T. & A.D. Poyer. 280 pp.
- Kozlova, E. V. & Tugarinov, A. Y. 1947 *Colymbiformes, Procelariiformes, Pelecaniformes, Ciconiiformes, Phoenicopteriformes*. Birds, I (3). Fauna SSSR (novaia seriia). 33. Leningrad: Akademiiia nauk SSSR, Zoologicheskii institut. 317 pp.
- Kozlova, E. V. 1957 *Charadriiformes. Suborder Alcae. Birds, II (3)*. Fauna SSSR (novaia seriia). 65. Leningrad: Akademiiia nauk SSSR, Zoologicheskii institut. 144 pp.
- Kozlova, E. V. 1961 *Charadriidae. Suborder Limicolae. Birds, II (1), Part 2*. Fauna SSSR (novaia seriia). 80. Leningrad: Akademiiia nauk SSSR, Zoologicheskii institut. 501 pp.
- Kozlova, E. V. 1962 *Charadriidae. Suborder Limicolae. Birds, II (1), Part 3*. Fauna SSSR (novaia seriia). 81. Leningrad: Akademiiia nauk SSSR, Zoologicheskii institut. 433 pp.
- Lloyd, C. 1981 *Birdwatching on estuaries, coast and sea*. London: Severn House Naturalist's library. 160 pp.
- Olsen, K. M. & Larsson, H. 1995 *Terns of Europe and North America*. Princeton: Princeton University Press. 224 pp.
- Remmert, H. 1957 *Aves. Tierwelt der Nord- und Ostsee*. 12j. Leipzig: Akademische verlagsgesellschaft. 102 pp.

- Shtegman, B. K. 1933 *Birds of the USSR. Keys to families of the USSR birds.* Opredeliteli po faune SSSR. 14. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 14 pp.
- Tuck, G. & Heinzel, H. 1958 *Guide des oiseaux de mer.* Neuchâtel, Paris: Delachaux & Niestlé. 310 pp.
- Yudin, K. A. 1965 *Phylogeny and classification of Charadriiformes. Birds, II (I). Part. I.* Fauna SSSR (novaia serii). 91. Akademii nauk SSSR, Zoologicheskii institut. 262 pp.
- Mammalia**
- Cagnolaro, L., Di Natale, A. & Notarbartolo di Sciara, G. 1983 *Cetacei. Guide per il riconoscimento delle specie animali delle acque lagunari e costiere italiane.* 9. Roma. 185 pp.
- Casado, M. F. & et al.. 1997 *Mamíferos marinos y tortugas del Mar de Alborán : guía d'especies marinas amenazadas.* Sevilla: Consejería de Medio Ambiente. 119 pp.
- Duguy, R. & Robineau, D. 1973 Cétacés et phoques des côtes de France. *Annales de la Société des Sciences Naturelles de Charentes-Maritime, supplément:* 1-93.
- Duguy, R. & Robineau, D. 1982 *Guide des Mammifères marins d'Europe.* Neuchâtel: Delachaux & Niestlé. 200 pp.
- Duguy, R. & Robineau, D. 1987 *Guia de los mamíferos de Europa.* Barcelona: Omega. 198 pp.
- Duguy, R. & Robineau, D. (ed.) 1992 *Handbuch der Säugetiere Europas, 6(2), Robben.* Wiesbaden: Aula Verlag. 309 pp.
- Evans, P. G. H. 1996 *Guide to the identification of whales, dolphins and porpoises in European seas.* Edinburgh: Scottish Natural Heritage. 36 pp.
- Fraser, F. C. 1976 *British whales, dolphins and porpoises. A guide for the identification and reporting of stranded whales, dolphins and porpoises on the British coasts.* London: British Museum (Natural History). xi+34 pp.
- Freund, L. 1932 *Cetacea.* Tierwelt der Nord- und Ostsee. 12k1. Leipzig: Akademische verlagsgesellschaft. 64 pp.
- Freund, L. 1933 *Pinnipedia.* Tierwelt der Nord- und Ostsee. 12k2. Leipzig: Akademische verlagsgesellschaft. Pp. 65-83.
- Notarbartolo di Sciara, G. & Demma, M. 1994 *Guida dei mammiferi marini del Mediterraneo.* Padova: Franco Muzzio. 262 pp.
- Robineau, D., Duguy, R. & Klima, M. (ed.) 1995 *Handbuch der Säugetiere Europas, 6(1A&1B), Wale und Delphine.* Wiesbaden: Aula Verlag. 503 pp.
- Tomilin, A. G. 1962 *Cetacea in seas of the USSR.* Opredeliteli po faune SSSR. 79. Leningrad: izdavaemye Zoologicheskim institutom Akademii Nauk SSSR. 212 pp.

APPENDIX 1 - CONTACT DETAILS OF THE PERSONS AND ORGANISATIONS CONTRIBUTING TO ERMS

Person	Company, Department	Address	Telephone	Fax	E-mail address
Agatha, Sabine	Institut für Zoologie	Universitaet Salzburg Hellbrunnerstrasse 34 A-5020 Salzburg, Austria			Sabine.agatha@sbg.ac.at
Arvanitidis, Dr Christos	Institute of Marine Biology of Crete	PO Box 2214 Heraklion, GR-71003 Crete, Greece	+30-81-346860	+30-81-241882	arvanitidis@imbc.gr
Bachelet, Dr Guy	Laboratoire d'Océanographie Biologique	Station Marine D'Arcachon Université Bordeaux 1 2, rue du Professeur Jolyet F-33120 Arcachon, France	+33-556-223900	+33-556-835104	bachelet@biocean.m-bordeaux.fr
Bartsch, Dr Ilse	Forschungsinstitut und Naturmuseum Senckenberg	Notkestrasse 31 Hamburg, D-22607 Germany	+49-408-9693212	+49-408-9693115	bartsch@meeresforschung.de
Bellan, Gérard	Centre d'Océanologie de Marseille	Université de la Méditerranée, +33-491-041633 Station Marine d'Endoume Rue de la Batterie des Lions, F-13007 Marseille, France	+33-491-041633	+33-491-041635	gbellan@com.univ-mrs.fr
Bellan-Santini, Dr Denise	Centre d'Océanologie de Marseille	Université de la Méditerranée, +33-491-041633 Station Marine d'Endoume Rue de la Batterie des Lions, F-13007 Marseille, France	+33-491-041633	+33-491-041635	bellan@com.univ-mrs.fr
Bird, Dr Graham J.		8 Elthorne Villas, Edgar Road, West Drayton Middlesex, UB7 8HW, UK	+44-1895-448246		claire.cohen@brunel.ac.uk
Bisby, Prof. Frank A.	Centre for Plant Diversity and Systematics	School of Plant Sciences The University of Reading Reading RG6 6AS, UK	+44-118931 6437/6466 +44 118 931 6466	+44-118-9753676	F.A.Bisby@reading.ac.uk
Bouchet, Prof. Philippe	Laboratoire de Biologie des Invertébrés Marins & Malacologie	Muséum National d'Histoire Naturelle 55 rue Buffon F-75005 ParisFrance	+33-1-40793103	+33-1-40793089	pbouchet@mnhn.fr
Boxshall, Prof. Geoff	Department of Zoology	The Natural History Museum Cromwell Road London SW7 5BD, UK	+44-207-94-25749	+44-207-9425433	gab@nhm.ac.uk
Brandt, Dr Sue M.	Centre for Plant Diversity and Systematics	School of Plant Sciences The University of Reading Reading RG6 6AS, UK	+44-118931 6437/6466 +44 118 931 6466	+44-118-9753676	S.M.Brandt@reading.ac.uk
Brattegard, Dr Torleiv	Department of Fisheries and Marine Biology	P.O. Box 7800 N-5020 Bergen, Norway	+47-55-584414 +47-55-5846 27	+47-55-584450	Torleiv.Brattegard@ifm.uib.no
Bray, Dr Rod	Department of Zoology	The Natural History Museum Cromwell Road London SW7 5BD, UK	+44-207-94-25749	+44-207-9425433	r.bray@nhm.ac.uk
Castelli, Dr Alberto	Ist. Scienze Antropologiche	Corso Margherita di Savoia 15, Università di Sassari I-07100 Sassari, Italy	+39-79228922	+39-79228925	
Clipson, Dr Nick J.W.	Department of Industrial Microbiology	University College Dublin Belfield, Dublin 4, Ireland	+353-1-7061356	+353-1-7061183	nclipson@maccollamh.ucd.ie
Collier, Louise	Ecological Consultancy Services Ltd	I7 Rathfarnham Road Terenure, Dublin 6W, Ireland	+353-1-4903237	+353-1-4925694	lcollier@ecoserve.ie
Connor, David	Marine Information Team	Joint Nature Conservation Committee, Monkstone House, City Road, Peterborough PE1 1JY, UK	+44-1733-866837	+44-1733-555948	connor_d@jncc.gov.uk
Cornelius, Dr Paul	Department of Zoology	The Natural History Museum Cromwell Road London SW7 5BD, UK	+44-207-94-25749	+44-207-9425433	p.cornelius@nhm.ac.uk

Person	Company, Department	Address	Telephone	Fax	E-mail address
Costello, Dr Mark J.	Ecological Consultancy Services Ltd	17 Rathfarnham Road Terenure Dublin 6W, Ireland	+353-1-4903237	+353-1-4925694	mcostello@ecoserv e.ie
Dauvin, Dr Jean-Claude	Marine Station	B.P. 80, 62930 Wimereux France	+33-3-21992920	+33-3-21992901	
Dick, Prof. Michael	Centre for Plant Diversity and Systematics	School of Plant Sciences The University of Reading Whiteknights, Reading Berkshire RG6 6AS, UK			M.W.Dick@readin g.ac.uk
Dowse, Jenny	Ecological Consultancy Services Ltd	17 Rathfarnham Road Terenure Dublin 6W, Ireland	+353-1-4903237	+353-1-4925694	jdowse@ecoserve.i e
Emblow, Chris	Ecological Consultancy Services Ltd	17 Rathfarnham Road Terenure Dublin 6W, Ireland	+353-1-4903237	+353-1-4925694	cemblow@ecoserve .ie
Erséus, Prof. Christer	Department of Invertebrate Zoology	Swedish Museum of Natural History, Box 50007, S-104 05 Stockholm Sweden	+46-851954129	+46-851954125	christer.erseus@nr m.se
Faubel, Dr Anno	Institut fr Hydrobiologie u. Fischereiwissenschaft	Zeisweg 9, Hamburg D-22765, Germany	+49-40-4123-6690	+49-40-41236696	a.faub@rrz-cip- l.rrz.uni- hamburg.de
Garrido, Mateo	University of Las Palmas de Gran Canaria	Campus de Tafira Edif Ciencias Basicas B-116 35017 Las Palmas Canary Islands	+34-928-452851		Mgarrido@biologia .ulpgc.es
Gibson, Dr David	Department of Zoology	The Natural History Museum Cromwell Road London SW7 5BD, UK	+44-207-94-25749	+44-207-9425433	dig@nhm.ac.uk
Gofas, Serge	Laboratoire de Biologie des Invertébrés Marins & Malacologie	Muséum National d'Histoire Naturelle 55 rue Buffon F-75005 ParisFrance	+33-1-40793103	+33-1-40793089	gofas@mnhn.fr
Goodkov, Andrew	Biological Research Institute of St Petersburg State University	Oranienbaumskoye Sch 2 Stary Peterhof, St Petersburg 198904 Russia			good@AGI060.spb .edu
Gross, Dr Onno	Hegestrasse 46b	Universitaet Hamburg Hamburg, 20251 Hamburg Germany			onnogross@compu serve.com
Guiry, Prof. Michael D.	Department of Botany	Martin Ryan Institute National University of Ireland Galway, Ireland	+353-91-525005	+353-91-75002 +353-91-525005	mike.guiry@seawe ed.ie
Hansson, Hans G.	Tjaernoe Marine Biological Laboratory	96 Stroemstad SE-452 Sweden	+46-526-68636	+46-526-68607	Hans.G.Hansson@t mbl.gu.se
Harding, Pamela	DIVERSITAS - UNESCO	I rue Miollis, F-75732 Paris, Cedex 15, France	+33-1-45684067	+33-1-45685804	diversitas@unesco. org
Haslund, Ole- Henrik	Intergovernmental Oceanographic Commission	UNESCO, I rue Miollis F-75732 Paris, Cedex 15, France	+33-1-45684016	+33-1-45685812	o.haslund@unesco. org
Hayward, Dr Peter J.	School of Biological Sciences	University of Wales, Swansea, Singleton Park Swansea SA2 8PP, UK	+44-92-295360	+44-92-295447	P.J.Hayward@Sw ansea.ac.uk
Heip, Dr Carlo	Netherlands Institute of Ecology	Centrum voor estuariene en mariene oecologie Korringaweg 7, 4401 EA Yerseke, The Netherlands	+31-113-577 445/488	+31-113-573616	heip@cemo.nioo.kn aw.nl
Horne, Dr David J.	School of Earth and Environmental Sciences	University of Greenwich Chatham Maritime, Kent ME4 4TB, UK	+44-181-3319841	+44-181-3319805	d.j.horne@greenwi ch.ac.uk

Person	Company, Department	Address	Telephone	Fax	E-mail address
Hummel, Herman	Netherlands Institute of Ecology	Centrum voor estuariene en mariene oecologie Korringaweg 7, 4401 EA Yerseke, The Netherlands	+31-113-577 445/488	+31-113-573616	hummel@cemo.nio o.knaw.nl
Huys, Dr Rony	Department of Zoology	The Natural History Museum Cromwell Road London SW7 5BD, UK	+44-207-94-25749	+44-207-9425433	r.huys@nhm.ac.uk
Jankowski, Dr Anatol Wiktor		poste restante (ulica Tam- basova 32), post office 198259, St. Petersburg, Russia	+812-1360790		tatiana.kind@palo ma.spbu.ru
Jones, Dr Robert W.	B.P. Exploration Operating Company	4/5 Longwalk Road Stockley Park, Uxbridge, UB11 1B, Middlesex, UK.		+44-81-7500530	Jonesbob@bp.com
Karakassis, Dr Ioannis	Institute of Marine Biology of Crete	PO Box 2214 GR-71003 Heraklion, Crete, Greece	+30-81-346860	+30-81-241882	jkarak@imbc.gr
Karlsbakk, Egil	Department of Fisheries and Marine Biology	P.O. Box 7800 N-5020 Bergen, Norway	+47-55-584414 +47-55-584627	+47-55-584450	egil.karlsbakk@ifm .uib.no
Krapp, Dr Franz	Zoologisches Forschungsinstitut Museum	Alexander Koenig D-53113 Bonn, Adenamerallee 150-164, Germany	+49-228-9122294	+49-228-216979	Franz.Krapp.ZFMK @uni-bom.ie
Kuhnt, Prof. Wolfgang	Geologisch- Palaeontologische s Institut	Universitaet Kiel Olshausenstrasse 40 Kiel, 24118 Kiel Germany	+49-431-8802924		wk@gpi.uni-kiel.de
Landy, Eleanor	Department of Industrial Microbiology	University College Dublin Belfield, Dublin 4, Ireland	+353-1-7061356	+353-1-7061183	Eleanor.landy@ucd .ie
Lasserre, Dr Pierre	DIVERSITAS - UNESCO	1 rue Miollis, F-75732 Paris, Cedex 15, France	+33-1-45684067	+33-1-45685804	p.lasserre@unesco. org
Le Renard, Dr Jacques	Laboratoire de Biologie des Invertébrés Marins & Malacologie	Muséum National d'Histoire Naturelle 55 rue Buffon F-75005 Paris France	+33-1-40793103	+33-1-40793089	lerenard@mnhn.fr
Legakis, Dr Anastasios	Zoological Museum	University of Athens Panepistimiopolis, GR-15784 Athens, Greece	+301-724372	+301-724604	alegakis@atlas.uoa. gr
Los, Dr Wouter	ETI Biodiversity Center	Universiteit van Amsterdam, Mauritskade 61, 1092 AD, Amsterdam, The Netherlands	+31-20-5257239	+31-20-5257238	los@bio.uva.nl
Markham, John	Arch Cape Marine Laboratory	Arch Cape Oregon 97102-0105 USA	+1-503-436-2310		jmarkham@seasurf. com
McCrea, Mona	Ecological Consultancy Services Ltd	17 Rathfarnham Road Terenure Dublin 6W, Ireland	+353-1-4903237	+353-1-4925694	mona@ecoserve.ie
Meisterfeld, Dr Ralf	Institut fur Biologie II (Zoologie)	RWTH Aachen, Kopernikusstrasse 16, D-52074 Aachen, Germany			meisterfeld@rwth- aachen.de
Minelli, Prof Alessandro	Dept. of Biology	Università degli studi di Padova Viale Giuseppe Colombo 31-35121 Padova Italy	+39-49-8276303	+39-49-8276300	almin@civ.bio.unip d.it
Moestrup, Dr Ojvind	Bot. Inst Dept Mycology & Phycology	Kobenhavns Univ., Oster Farimagsgade 2D, Copenhagen K, DK-1353, Denmark	+45-35322290	+45-35322321	moestrup@bot.ku.d k

Person	Company, Department	Address	Telephone	Fax	E-mail address
Monniot, Claude	Laboratoire de Biologie des Invertébrés Marins & Malacologie	Muséum National d'Histoire Naturelle 55 rue Buffon F-75005 Paris, France	+33-1-40793103	+33-1-40793089	monniot@mnhn.fr
Mulisch, Dr Maria	Zoologisches Institut	Universitaet Koeln, Weyertal 119, 50923 Koeln Germany	+49-221-4703112	+49-221-4705171	mmulisch@novell. biolan.uni-koeln.de
Norena Janssen, Carlina	Museo Nacional de Ciencias Naturales	Jose Gutierrez Abascal 2, E-28006 Madrid, Spain			carolina.norena@t- online.de
Perez-Uz, Dr Blanca	Dept. Microbiologia III	Fac. C.C. Biologicas Universidad Complutense de Madrid, Madrid, Spain	+34-1-39449	+34-1-394-4964	perezuz@eucmax.si m.ucm.es
Radach, Dr Günther	Institut für Meereskunde	Universität Hamburg Tropowitzstr. 7, 22529 Hamburg, Germany	+49-40-42838 5656	+49-40-560-5724	radach@ifm.uni- hamburg.de
Ramos, Dr Maria A.	Dept. biodiversidad y biología evolutiva	Museo Nacional de Ciencias Naturales José Gutierrez Abascal 2 E-28006 Madrid, Spain	+34-91-4111328	+34-91-5645078	m.ramos@mncn.cs i.es
Rogerson, Andrew	Oceanographic Centre	Nova Southeastern University, 8000 N Ocean Drive, Dania, Florida 33004, USA			arogerso@ocean.no va.edu
Salvatori, Dr Arico	Head Jakarta Mandate Unit	World Trade Center, Suite 300, 393 St. Jacques St Montreal Quebec	+1-514-2877009	+1-514-2886588	salvatore.arico@bio div.org
Schmidl, Dr Gerhard	Department of Geological Sciences	Orton Hall 245, 155 South Oval Mall, The Ohio State University, Columbus, OH 43210, USA	+1-614-6884440	+1-614-2921496	Schmidl@geology .ohio-state.edu
Scholz, Joachim	Forschungsinstitut Senckenberg	Senckenberglage 25 D-60325 Frankfurt a. M., Germany	+49-69-752240	+49-69-746238	jscholz@sngkw.uni -frankfurt.de
Serrão Santos, Ricardo	Department of Oceanography and Fisheries	University of the Azores P-9900 Horta, Azores, Portugal	+351-92-292944	+351-92-292659	ricardo@dop.uac.pt
Sneli, Dr Jon- Ame	Department of Natural History	Trondhjem Biologiske Stasjon, Norwegian University of Science and Technology, N-7491 Trondheim , Norway	+47-73-591 586	+47-73-591597	jon.sneli@vm.ntnu. no
Southward, Dr Alan	Marine Biological Association of the United Kingdom	The Laboratory Citadel Hill, Plymouth PL1 2PB, UK			100721.3720@comp userve.com
Templado- Gonzalez, J.	Dept. biodiversidad y biología evolutiva	Museo Nacional de Ciencias Naturales José Gutierrez Abascal 2, E-28006 Madrid, Spain	+34-91-4111328	+34-91-5645078	j.templado@mncn.cs i.es
Türkay, Dr Michael	Forschungsinstitut Senckenberg	Senckenberglage 25 D-60325 Frankfurt a. M., Germany	+49-69-752240	+49-69-746238	mtuerkay@sng.uni- frankfurt.de
van der Land, Dr Jacob	Nationaal Naturhistorisch Museum	Darwinweg, P.O. Box 9517 2300 RA Leiden, The Netherlands	+31-71-5687654	+31-71-5687666	land@naturalis.nn m.nl
van Soest, Dr Rob	Institute for Systematics and Ecology (Zoologisch Museum)	University of Amsterdam Kruislaan 320, 1098 SM Amsterdam, The Netherlands			soest@bio.uva.nl

Person	Company, Department	Address	Telephone	Fax	E-mail address
Vincx, Dr Magda	Marine Biology Section, Biology Department	University Gent Ledegankstraat 35 B-9000, Gent Belgium	+32-9-2645210	+32-9-2645344	magda.vincx@rug.ac.be
Warwick, Dr Richard	Plymouth Marine Laboratory	Prospect Place West Hoe Plymouth PL1 3DH, UK	+44-1752-633 - 428 (-100)	+44-1752-633101	RMW@wpo.nerc.ac.uk, r.warwick@pml.ac.uk
Watling, Dr Les	School of Marine Sciences	Darling Marine Center University of Maine Walpole, ME 04573, USA	+1-207-563-3146- x248	+1-207-563-8407	watling@maine.maine.edu
White, Dr Richard J.	School of Biological Sciences	Bassett Crescent East University of Southampton Southampton SO16 7PX, UK	+44-2380-592021	+44-2380-594269	R.J.White@soton.ac.uk
Wirtz, Dr Peter	Department of Oceanography and Fisheries	University of the Azores P-9900 Horta, Azores, Portugal	+351-92-292944	+351-92-292659	peter@dop.uac.pt

INDEX TO ORDERS, FAMILIES AND GENERA

Order and family names are in **bold**, genus names are in *italic*.

- Aaptos*, 88
- Aartsenia*, 198
- Abarenicola*, 216
- Abditodentrix*, 70
- Abietinaria*, 116
- Abludomelita*, 304
- Abollifer*, 79
- Abothrium*, 146
- Abra*, 210
- Abralia*, 212
- Abraliopsis*, 212
- Absidia*, 15
- Abudesdus*, 371
- Abyla*, 121
- Abylidæ**, 121
- Abylopsis*, 121
- Abyssacidia*, 352
- Abyssidrilus*, 233
- Abyssinoe*, 219
- Abyssocucumis*, 348
- Abyssoninoe*, 219
- Abyssorchromene*, 301
- Acaenomolgus*, 258
- Acanella*, 105
- Acanthamoeba*, 53
- Acanthamoebidae**, 53
- Acanthaspididae*, 316
- Acanthella*, 89
- Acanthephyra*, 286
- Acanthurypon*, 91
- Acanthicolepis*, 223
- Acanthobothrium*, 146
- Acanthocardia*, 209
- Acanthocarpus*, 290
- Acanthocephala**, 176
- Acanthocephaloides*, 176
- Acanthocephalus*, 176
- Acanthochaetus*, 367
- Acanthochelius*, 174
- Acanthochitona*, 182
- Acanthochitonidae**, 182
- Acanthochondria*, 255
- Acanthochondrites*, 255
- Acanthocope*, 317
- Acanthocorbis*, 57
- Acanthocotyle*, 142
- Acanthocybium*, 373
- Acanthocythereis*, 250
- Acanthodasys*, 157
- Acanthodoris*, 202
- Acanthoeeca*, 57
- Acanthoecidae**, 57
- Acanthogorgia*, 104
- Acanthogorgiidae**, 104
- Acanthogyrus*, 176
- Acanthoica*, 76
- Acantholabrus*, 371
- Acantholaimus*, 161
- Acanthomacrostomum*, 127
- Acanthomunna*, 316
- Acanthomysis*, 293
- Acanthonchus**, 161
- Acanthonotozoma*, 296
- Acanthonotozomatidae**, 296
- Acanthonyx*, 290
- Acanthopharyngoides*, 161
- Acanthopharynx*, 161
- Acanthophora*, 32
- Acanthoscina*, 308
- Acanthostephia*, 305
- Acanthostomum*, 136
- Acanthostracion*, 374
- Acanthotrema*, 136
- Acanthotrochus*, 351
- Acanthuridae**, 373
- Acanthurus*, 373
- Acarinus*, 93
- Acarochelopodia*, 239
- Acarochelopodinae*, 239
- Acaromantis*, 241
- Acartia*, 263
- Acartiidae**, 263
- Acasta*, 282
- Acastinae**, 282
- Acaulidae**, 112
- Acaulis*, 112
- Acauloides*, 112
- Accacladocoelium*, 136
- Accacoelium*, 137
- Aceroides*, 305
- Acerotisa*, 128
- Acervulina*, 69
- Acervulinidae**, 69
- Acesta*, 207
- Aceste*, 348
- Aestoplana*, 128
- Acetabularia*, 23
- Acetabulastoma*, 250
- Achaeta*, 234
- Achaeus*, 291
- Achelia*, 241
- Acheliderma*, 93
- Achiridota*, 351
- Achlyonice*, 350
- Achoerus*, 125
- Achlooe*, 223
- Achradina*, 47
- Acidostoma*, 301
- Acinetospora*, 27
- Acipenser*, 360
- Acipenseridae**, 360
- Acipenseriformes*, 360
- Acirsa*, 186
- Aclididae**, 186
- Aclis*, 186
- Acmaea*, 182
- Acmaeidae**, 182
- Acmostomum*, 129
- Acoela*, 125
- Acoetidae**, 214
- Acontiophorus*, 259
- Acremonium*, 18
- Acrenhydrosoma*, 271
- Acriloscala*, 186
- Acobelione*, 322
- Acrocalanus*, 266
- Acrochaete*, 24
- Acrochaetiaceae**, 29
- Acrochaetales**, 29
- Acrochaetium*, 29
- Acrochalix*, 187
- Acrocirridae**, 214
- Acrocirrus*, 214
- Acrocytis*, 26
- Acrodiscus*, 35
- Acrorhynchides*, 132
- Acrorhynchus*, 132
- Acrosiphonaceae**, 21
- Acrosiphonia*, 21
- Acrosiphoniales**, 21
- Acrosorium*, 32
- Acrospongium*, 27
- Acrosympytaceae**, 35
- Acrosymphton*, 35
- Acrothamnion*, 30
- Acrothoracica*, 281
- Acrotrix*, 24
- Acrotichaceae**, 24
- Acrumena*, 130
- Acrumenidae**, 130
- Acryptolaria*, 115
- Actacarinae**, 239
- Actacarus*, 239
- Acteocina*, 199
- Acteon*, 198
- Acteonidae**, 198
- Acthelmis*, 107
- Actigia*, 113
- Actinarctus*, 236
- Actinauge*, 107
- Actinernidae**, 106
- Actinernus*, 106
- Actinia*, 106
- Actiniaria**, 106
- Actinichonidae**, 41
- Actiniidae**, 106
- Actiniscus*, 47
- Actinomucor*, 15
- Actinonema*, 161
- Actinoposthia*, 125
- Actinoscyphia*, 106
- Actinoscyphiidae**, 106
- Actinostola*, 106
- Actinostolidae**, 106
- Actinothoe*, 108
- Actinulida**, 112
- Aculeopsis*, 276
- Aculeorhynchidae**, 130
- Aculeorhynchus*, 130
- Adagnesia*, 352
- Adalaria*, 202
- Adamsia*, 107

- Addisonia*, 183
Addisoniidae, 183
Adelodrilus, 233
Adeloposthia, 125
Adelosina, 66
Adenocauda, 126
Adenoides, 47
Adenopea, 125
Adenopharynx, 133
Adenopleurellidae, 268
Adenorhynchus, 133
Adeonella, 326
Adeonellopsis, 326
Adeonidae, 326
Adeorbidae, 190
Adercodon, 215
Adercotryma, 62
Adeuchromadora, 161
Adeuomphalus, 184
Adinimonas, 47
Adinosoma, 137
Admete, 194
Adoncholaimus, 161
Adreus, 88
Adriogloia, 24
Advena, 260
Adyte, 223
Aedicira, 222
Aega, 319, 320
Aegaeon, 288
Aegialoalaimus, 161
Aegidae, 319
Aegina, 118
Aeginella, 296
Aeginidae, 118
Aeginina, 296
Aeginopsis, 118
Aeginura, 119
Aegires, 201
Aegiretidae, 201
Aegistidae, 268
Aegisthus, 268
Aeodes, 35
Aeolidia, 204
Aeolidiella, 204
Aeolidiidae, 204
Aeolosoma, 214
Aeolosomatidae, 214
Aepophilus, 324
Aepopsis, 323
Aepus, 323
Aequipecten, 207
Aequorea, 113
Aequoreidae, 113
Aeropsidae, 348
Aeropsis, 348
Aetea, 326
Aeteidae, 326
Aetheolineus, 153
Aethozoon, 332
Aethozontidae, 332
Aetideidae, 263
Aetideopsis, 263
Aetideus, 263
Affinetrina, 66
Afrocardium, 209
Afrolaophonte, 275
Afronta, 126
Afropinnotheres, 292
Agalma, 122
Agalmatidae, 122
Agamascaris, 174
Agamoneema, 174
Agardhiella, 37
Agaricaceae, 18
Agaricales, 18
Agathotanaini, 314
Agathotanais, 314
Agaue, 240
Agauides, 240
Agauopsis, 240
Agelas, 90
Agelasidae, 90
Agetus, 256
Aglaocypris, 247
Aglaja, 199
Aglajidae, 199
Aglantha, 119
Aglaophamus, 220
Aglaophenia, 113
Aglaopheniidae, 113
Aglaothamnion, 30
Agaura, 119
Agnatha, 358
Agnathiellidae, 151
Agnezia, 352
Agneziidae, 352
Agonidae, 369
Agonus, 369
Ahliesaurus, 363
Ahnfeltia, 29
Ahnfeltiaceae, 29
Ahnfeltiales, 29
Ahnfeltiopsis, 36
Aidablennius, 372
Aigondiceps, 278
Aiolocolax, 32
Aiptasia, 106
Aiptasiidae, 106
Aiptasiogeton, 106
Aka, 98
Akanthophoreinae, 314
Akanthophoreus, 314
Akentrogonida, 283
Akera, 201
Akeridae, 201
Akessonnia, 258
Akritogryra, 184
Aktedrilus, 233
Alaba, 185
Alabaminella, 72
Alabaminidae, 69
Alacia, 245
Alaimella, 161
Alaria, 28
Alariaceae, 28
Alataspora, 84
Alatasporidae, 84
Alaurina, 128
Albanemertes, 155
Albertorhynchus, 132
Albionella, 260
Albunea, 290
Albuneidae, 290
Alca, 375
Alcataenia, 146
Alcidae, 375
Alciopa, 214
Alciopidae, 214
Alciopina, 214
Alcyonacea, 104
Alcyonidiidae, 332
Alcyonium, 332
Alcyoniidae, 104
Alcyonium, 104
Alderella, 201
Alderia, 201
Alderina, 327
Aldisa, 202
Aldisidae, 202
Aldrovandia, 360
Alebion, 260
Alectis, 370
Alectona, 87
Alella, 260
Alentia, 223
Alentiana, 223
Alepas, 281
Alepes, 370
Alepiella, 56
Alepisauridae, 364
Alepisaurus, 364
Alepocephalidae, 361
Alepocephalus, 361
Alexandrium, 47
Alexandromenia, 181
Alexlutheria, 129
Algae, 21
Algarvia, 204
Aliceia, 196
Alicia, 106
Aliciidae, 106
Alima, 284
Alitropus, 319
Alkmaria, 215
Allantactis, 107
Allantion, 79
Allantogynus, 261
Alle, 375
Allescheria, 18
Alloanthura, 316
Allocyttus, 368
Alloeocarpa, 354
Allogramma, 211
Allogromia, 62
Allogromiidae, 62
Allogromiina, 62
Allomelita, 304
Allopatria, 341
Allopodocotyle, 137
Alloposidae, 213
Allopseudaxine, 142
Allopseudocolocytotrema, 137
Allosquilla, 284
Allostoma, 129
Alloteuthis, 212
Alluna, 125
Almaniella, 224
Alopias, 358
Alopiidae, 358
Alora, 186

- Alosa*, 361
Alpheidae, 287
Alpheopsis, 287
Alpheus, 287
Alsidium, 32
Alternaria, 18
Alteutha, 277
Althoffia, 356
Altiverruca, 282
Aluterus, 374
Alvania, 188
Alvinocarididae, 286
Amaeana, 230
Amage, 215
Amagopsis, 215
Amakusanhra, 316
Amallophora, 267
Amallothrix, 267
Amalosoma, 178
Amalthaea, 112
Amaryllis, 302
Amastigomonas, 57
Amathia, 332
Amathillopsidae, 297
Amathillopsis, 297
Amauropsis, 191
Ambasia, 302
Ambasiella, 302
Amblyops, 293
Amblyopsoides, 293
Amblyosyllis, 228
Amblyraja, 359
Ambunguipedidae, 268
Ambunguipes, 268
Ameira, 268
Ameiridae, 268
Ameiopsis, 269
Ameiopsyllus, 269
Amenophia, 279
Amidostomum, 174
Amigdascalpellum, 282
Ammatophora, 327
Ammobaculites, 64
Ammodiscidae, 62
Ammodiscus, 62
Ammodytes, 372
Ammodytidae, 372
Ammolagena, 62
Ammomassilina, 67
Ammonia, 69
Ammonicera, 196
Ammoniidae, 69
Ammopempix, 64
Ammosalaria, 63
Ammosphaeroidinidae, 62
Ammothea, 241
Ammotheidae, 241
Ammothella, 241
Ammotium, 63, 64
Ammotrochoides, 64
Ammotrypanella, 221
Amoebae, naked, 53
Amoebae, testate, 55
Amoebidae, 53
Amoebophyra, 47
Amoenoscopa, 57
Amonardia, 272
Amorphinopsis, 90
Amorphonemertes, 153
Ampelisca, 297
Ampeliscidae, 297
Amperima, 350
Ampharete, 215
Ampharetidae, 215
Amphianthus, 107
Amphiascoides, 272
Amphiascopis, 272
Amphiascus, 272
Amphibdella, 142
Amphibdelloides, 142
Amphiblestrum, 327
Amphicaryon, 121
Amphichaeta, 233
Amphichilus, 343
Amphicoryna, 75
Amficotide, 146
Amphicteis, 215
Amphidelus, 161
Amphidiniopsis, 47
Amphidinium, 47
Amphidiscophora, 102
Amphidoma, 47
Amphiduros, 218
Amphifrenestrella, 64
Amphiglena, 225
Amphilepididae, 344
Amphilepis, 344
Amphilina, 146
Amphilochidae, 297
Amphilochoides, 297
Amphilochopsis, 297
Amphilochus, 297
Amphimenia, 181
Amphimeniidae, 181
Amphimonas, 79
Amphimonhystera, 161
Amphinema, 118
Amphinome, 215
Amphinomidae, 215
Amphionides, 288
Amphionididae, 288
Amphiophiura, 345
Amphioplus, 343
Amphipolis, 343
Amphiplica, 183
Amphipoda, 296
Amphiporella, 154
Amphiporidae, 154
Amphiporus, 154
Amphirhynchus, 132
Amphirissoa, 189
Amphiro, 217
Amphiroa, 33
Amphisbetia, 116
Amphiscolops, 125
Amphisolenia, 47
Amphisorus, 69
Amphisphaeria, 18
Amphisphaeriaceae, 18
Amphissa, 194
Amphistegina, 69
Amphisteginidae, 69
Amphitantulus, 280
Amphitholina, 297
Amphithopsis, 299
Amphitremidae, 57
Amphitrite, 230
Amphitritides, 230
Amphiura, 343
Amphiuridae, 343
Amphiute, 101
Amphoriscidae, 101
Amphoriscus, 101
Ampithoe, 297
Ampithoidae, 297
Ampulla, 194
Amuleta, 317
Amygdalum, 206
Amylax, 47
Amylocarpus, 17
Amylocorticaceae, 18
Amythasides, 215
Anabathridae, 190
Anacetabulitrema, 137
Anachis, 194
Anadara, 206
Anadyomenaceae, 22
Anadyomene, 22
Anamenia, 182
Ananhra, 316
Anapagurus, 289
Anaperidae, 125
Anaperus, 125
Anapophysia, 277
Anarchias, 360
Anarhichadidae, 371
Anarhichas, 371
Anarthropora, 326
Anarthrura, 314
Anarthruridae, 314
Anarthrurinae, 314
Anarthrurini, 314
Anatanaini, 313
Anatidae, 375
Anatoma, 183
Anchialina, 293
Anchinia, 355
Anchistrocephalus, 146
Ancistargis, 223
Ancistrobasis, 184
Ancistrocheiridae, 212
Ancistrocheirus, 212
Ancistrocoma, 46
Ancistrocomidae, 46
Ancistrorhynchus, 131
Ancistrosyllis, 223
Ancistrosyrinx, 195
Ancistroteuthis, 212
Ancorabolidae, 269
Ancorabolus, 269
Ancorina, 85
Ancorinidae, 85
Ancula, 202
Ancylocoelium, 137
Ancyrocephalus, 142
Ancyrocotyle, 143
Ancyromonas, 79
Andaniella, 306
Andaniexis, 307
Andaniopsis, 307
Andaniotes, 307

- Andracantha*, 176
Andresia, 107
Andesiidae, 107
Andvakia, 107
Andvakiidae, 107
Anekes, 184
Anelasma, 281
Anelasmatidae, 281
Anelassorhynchus, 178
Anemonactis, 107
Anemonia, 106
Angelopsis, 122
Angliera, 319
Anguilla, 360
Anguillicola, 174
Anguillidae, 360
Anguilliformes, 360
Anguinella, 332
Angulorostrum, 246
Anidolyta, 201
Aniptodera, 16
Anisakis, 174
Anisarchus, 371
Anisarthrus, 322
Anisocladium, 137
Anisocoelium, 137
Anisocomides, 46
Anisocrella, 95
Anisodoris, 202
Anisogammaridae, 298
Anisolabis, 324
Anisomysis, 293
Anisonema, 59
Anisostagma, 16
Anisotylacanthea, 93
Annatiara, 118
Annectocyma, 333
Annectocymidae, 333
Annulonemertes, 155
Annularhynchus, 132
Anobothrus, 215
Anodontia, 207
Anoikostoma, 137
Anomalina, 69
Anomalinidae, 69
Anomalinoides, 73
Anomalocera, 267
Anomalohalacarinae, 239
Anomalohalacarus, 239
Anomalotrema, 137
Anomia, 207
Anomiidae, 207
Anomoclausia, 255
Anomoclausiidae, 255
Anomocora, 109
Anomopera, 354
Anomopoda, 243
Anomopsyllus, 257
Anomotaenia, 146
Anonymidae, 128
Anonymus, 128
Anonyx, 302
Anopla, 152
Anoplocopea, 320
Anoplodactylus, 242
Anoplodiera, 130
Anoplodiopsis, 130
Anoplodiscus, 143
Anoplodium, 130
Anoplogaster, 367
Anoplogasteridae, 367
Anoploneis, 218
Anoplosoma, 269
Anoplostoma, 161
Anopteridae, 364
Anopterus, 364
Anotrichium, 30
Anoxycalyx, 103
Ansates, 182
Anseriformes, 375
Anseropoda, 341
Antarcella, 55
Antedon, 338
Antedonidae, 338
Antennella, 201
Antennariidae, 366
Antennarius, 366
Antennella, 115
Antennuloniscus, 318
Anteucherites, 255
Anteucheridae, 255
Anthelia, 105
Anthelura, 316
Antheluridae, 316
Anthessiidae, 255
Anthessius, 255
Anthias, 369
Antho, 92
Anthobothrium, 146
Anthocephalum, 146
Anthocephalus, 146
Anthocotyle, 143
Anthohebella, 115
Anthomastus, 104
Anthopharynx, 133
Anthophiura, 345
Anthopleura, 106
Anthoptilidae, 105
Anthoptilum, 105
Anthosactis, 106
Anthosoma, 260
Anthosphaera, 76
Anthostomella, 18
Anthothela, 104
Anthothelidae, 104
Anthura, 316
Anthuridae, 316
Anthuridea, 316
Anticoma, 161
Anticomopsis, 161
Anticyathus, 161
Anticyclus, 161
Antigonaria, 125
Antigonariidae, 125
Antigonia, 368
Antillesoma, 178
Antimora, 365
Antipatharia, 109
Antipathes, 109
Antipathidae, 109
Antithamnion, 30
Antithamnionella, 30
Antomicron, 161
Antonbruunia, 215
Antonbruuniidae, 215
Antonietta, 204
Antromacrostomum, 127
Antroposthia, 125
Antroposthidae, 125
Antygomonas, 159
Antygomonidae, 159
Anurida, 324
Anuridella, 324
Anuropoda, 313
Aonides, 227
Aora, 298
Aoridae, 298
Apanthura, 316
Apanthuroides, 316
Apatemon, 137
Aperiovula, 191
Aphalloides, 137
Aphallus, 137
Aphanipathes, 109
Aphanius, 367
Aphanochaete, 23
Aphanocladia, 32
Aphanodomus, 259
Aphanopus, 373
Aphanostoma, 125
Aphanurus, 137
Apharyngostrigaea, 137
Apharyngtus, 217
Aphelochaeta, 216
Apherusa, 299
Aphia, 372
Aphotopontius, 260
Aphrocallistes, 103
Aphrocallistidae, 103
Aphrocera, 101
Aphrodita, 215
Aphroditidae, 215
Aphrogenia, 215
Aphropharynx, 217
Aphyonidae, 366
Aphyonus, 366
Apicomplexa, free-living, 40
Apionsoma, 178
Apiospora, 18
Apistobranchidae, 215
Apistobranchus, 215
Aplacophora, 181
Apletodon, 367
Aplidiopsis, 353
Aplidium, 353
Aplidium, 353
Aploparaksis, 146
Aplousina, 327
Aplousobranchia, 352
Aplysia, 201
Aplysiidae, 201
Aplysilla, 101
Aplysina, 101
Aplysinidae, 101
Aplysinopsis, 100
Aplysiopsis, 200
Apodida, 350
Apodomyzon, 262
Apodopsyllus, 277
Apoglossocolax, 32
Apoglossum, 32

- Apogon*, 369
Apogonidae, 369
Apolemia, 122
Apolemiidae, 122
Apomatus, 226
Aponema, 161
Aponuphis, 221
Aponurus, 137
Apophallus, 137
Apopriionospio, 227
Aporchis, 137
Aporhynchus, 146
Aporocotyle, 137
Aporrhaiidae, 190
Aporrhais, 190
Appendichordella, 16
Appendicularia, 356
Applanola, 275
Apristurus, 359
Apseudes, 313
Apseudidae, 313
Apseudinae, 313
Apseudomorpha, 313
Apterichtus, 360
Apusomonads, 57
Apygophora, 281
Arabella, 215
Arabellidae, 215
Arachnactidae, 108
Arachnactis, 108
Arachnanthus, 108
Aracbnida, 237
Arachnididae, 332
Arachnidium, 332
Arachniotus, 17
Arachnoidea, 332
Archonomysis, 293
Arachnophyllum, 32
Araeolaimus, 161
Araneum, 352
Araphura, 314
Arbacia, 346
Arbaciella, 346
Arbaciidae, 346
Arbacioida, 346
Arcia, 206
Arcella, 55
Arcellidae, 55
Arcellina, 55
Arcellinida, 55
Archactinoposthia, 125
Archaeobalanidae, 282
Archaeobalaninae, 282
Archaeognatha, 323
Archaeosphaerodiniopsis, 47
Archaias, 68
Archaphanostoma, 126
Archescolecithrix, 267
Archesola, 275
Archiascidia, 353
Archiconchoecia, 245
Archidistoma, 352
Archidorididae, 202
Archidoris, 202
Archileptastacus, 276
Archilina, 135
Archiloa, 135
Archilopsis, 135
Archimisopria, 262
Archimonocelis, 135
Archimonotresis, 129
Archisenia, 277
Architectonicidae, 196
Archocelis, 127
Archoplana, 135
Arcidae, 206
Arcitalitrus, 307
Arcopagia, 209
Arcoscalpellinae, 282
Arcoscalpellum, 282
Arctapodema, 119
Arctica, 210
Arcticidae, 210
Arcticomisopria, 262
Arctinula, 207
Arctogadus, 365
Arctolembos, 298
Arctonemertes, 154
Arctonula, 330
Arctopontius, 259
Arctozenus, 363
Arcturella, 321
Arcturidae, 321
Arcturinella, 321
Arcturopsis, 321
Arcturus, 321
Arctulia, 305
Arculus, 208
Arenariomyces, 16
Arenicola, 216
Arenicolidae, 216
Arenicolides, 216
Arenocaris, 276
Arenonemertes, 155
Arenoportia, 270
Arenopontiidae, 270
Arenosetella, 273
Arenotrocha, 217
Argentina, 361
Argentinidae, 361
Argestidae, 270
Argestigens, 270
Argilloecia, 251
Argis, 288
Argissa, 298
Argissidae, 298
Argonauta, 213
Argonautidae, 213
Argonotholca, 150
Argulidae, 243
Argulus, 243
Argyripnus, 362
Argyropelecus, 362
Argyrosomus, 370
Argyrotheca, 334
Arhaphuroides, 314
Arhodeoporos, 240
Arhythmorhynchus, 176
Aricidea, 222
Arietellidae, 263
Arietellus, 263
Ariosoma, 361
Aristaeopsis, 285
Aristeidae, 285
Aristeomorpha, 285
Aristeus, 285
Aristias, 302
Aristostomias, 363
Armadoskrjabinia, 146
Armandia, 221
Armatoplana, 128
Armaueria, 156
Armaueridae, 156
Armichona, 41
Armina, 204
Arminidae, 204
Armarella, 62
Armorhydra, 118
Armorhydridae, 118
Arnoglossus, 374
Arnola, 137
Aroui, 302
Arrhinopsis, 305
Arrhis, 305
Artacama, 230
Artediellus, 369
Artemisina, 92
Arthrochaete, 21
Arthrocladia, 26
Arthrocladiaceae, 26
Arthropoma, 331
Arthropsis, 18
Arthropsyllus, 269
Arthropyreniaceae, 15
Articularia, 66
Articulina, 66
Artotrogidae, 259
Artotrogus, 259
Asabellides, 215
Asbestopluma, 97
Ascalitis, 101
Asandra, 101
Ascarophis, 175
Aschemonella, 78
Ascidia, 352
Asciidiacea, 352
Ascidicola, 253
Ascidicolidae, 253
Ascidella, 352
Ascidiiidae, 352
Asclerocheilus, 226
Ascobolaceae, 18
Ascobolus, 18
Ascobulla, 200
Ascochyta, 18
Ascocotyle, 137
Ascolaimus, 161
Ascomorpha, 150
Ascomycota, 15
Asconema, 103
Ascophanus, 18
Ascophyllum, 28
Ascorhiza, 332
Ascorhynchus, 242
Ascosia, 326
Ascosiidae, 326
Ascothoracida, 283
Ascothoracidae, 283
Ascothorax, 283
Ascyssa, 102

- Asellopsis*, 275
Asellota, 316
Aselomaris, 117
Ashtoret, 290
Aslia, 348
Asparagopsis, 30
Aspella, 192
Aspelta, 150
Asperarca, 206
Aspergillus, 18
Asperococcaceae, 26
Asperococcus, 26
Aspersentis, 176
Asperspina, 201
Asperspinidae, 201
Aspidelectra, 329
Aspidiophorus, 156
Aspidochirotida, 349
Aspidoconcha, 251
Aspidodiadema, 346
Aspidodiadematidae, 346
Aspidoechia, 261
Aspidogastrea, 136
Aspidogastrida, 136
Aspidophiura, 345
Aspidosiphon, 178
Aspidosiphonidae, 178
Aspinatrium, 143
Asplanchna, 150
Asplanchnidae, 150
Asquamiceps, 362
Assiminea, 190
Assimineidae, 190
Assulina, 56
Astacilla, 321
Astacolus, 75
Astarte, 208
Astartidae, 208
Asterias, 342
Astericola, 256
Asterigerinata, 69
Asterigerinatidae, 69
Asteriidae, 342
Asterina, 341
Asterinidae, 341
Asterochères, 259
Asterocheridae, 259
Asterocolax, 32
Astroidea, 338
Asteromenia, 38
Asteromyces, 18
Asteronema, 27
Asteronychidae, 343
Asteronyx, 343
Asterope, 245
Asterorotalia, 75
Asteroschema, 343
Asteroschematidae, 343
Asterosmilia, 109
Asterostomatidae, 348
Asthelys, 184
Asthenognathus, 292
Astomonema, 161
Astrangia, 110
Astrodia, 343
Astrogeron, 343
Astroides, 110
Astronesthes, 363
Astrononion, 74
Astropecten, 338
Astropectinidae, 338
Astrophorida, 85
Astrorhiza, 62
Astrorbizidae, 62
Astrospartus, 343
Astrotorhynchus, 134
Astylinifer, 93
Astyra, 298
Astyridae, 298
Asymmetrione, 322
Asymphylodora, 137
Atactodea, 209
Atagema, 202
Ataxolepis, 368
Ataxophragmiidae, 63
Atelecrinidae, 338
Atelecrinus, 338
Atelecyclidae, 291
Atelecyclus, 291
Ateleopidae, 363
Ateleopodiformes, 363
Atelopsalis, 240
Atergatis, 292
Atergia, 88
Atergopedia, 277
Athanas, 287
Athecanephria, 235
Athelges, 322
Atherina, 367
Atherinidae, 367
Atheriniformes, 367
Atherinomorus, 367
Atherospius, 227
Athorybia, 122
Athorybiidae, 122
Atlanta, 192
Atlantapseudes, 313
Atlanterythrops, 293
Atlantidae, 192
Atlantidrilus, 233
Atlantiella, 65
Atolla, 111
Atollidae, 111
Atomiscala, 197
Atorella, 111
Atorellidae, 111
Atractophora, 37
Atriaster, 143
Atrina, 207
Atriofronta, 125
Atrionemertes, 155
Atrispinum, 143
Atrochromadora, 161
Atrophia, 257
Atylus, 299
Atys, 199
Aubignyna, 75
Auchenoplax, 215
Audouinella, 29
Auerbachia, 84
Auerbachiidae, 84
Augaptilidae, 264
Augaptilina, 264
Augaptilus, 264
- Augeneria*, 219
Augeneriella, 225
Aulactinia, 106
Auleta, 89
Aulocalycidae, 103
Aulocyathus, 109
Aulofolliculina, 41
Aulopharynx, 133
Aulopidae, 363
Aulopiformes, 363
Auloplax, 103
Aulopus, 363
Aurelia, 111
Aureliana, 107
Aurelianiidae, 107
Aurichona, 41
Auriculigerina, 187
Auriculinella, 205
Aurila, 248
Aurivillialepas, 281
Aurobogidiella, 298
Aurospio, 227
Austranema, 161
Austrobella, 234
Astrodecidae, 242
Astrodecus, 242
Astrognatharia, 151
Astrognathia, 151
Astrognathiidae, 151
Australaenilla, 224
Austroniscus, 319
Austrorhynchus, 132
Austrosyrrhoe, 307
Autolitus, 228
Autolytus, 228
Automate, 287
Autonoe, 298
Auxis, 373
Auxoarthron, 17
Avagina, 126
Avocettina, 361
Avrainvillea, 21
Axelella, 194
Axelsonia, 324
Axia, 129
Axiidae, 289
Axine, 143
Axinella, 89
Axinellidae, 89
Axingella, 90
Axinodon, 208
Axinophilus, 258
Axinopsida, 208
Axinus, 208
Axinyssa, 90
Axionice, 230
Axiothella, 219
Axius, 289
Axitelga, 131
Axonolaimus, 162
Azorinus, 210
Azygocypridina, 245
Bacciger, 137
Bacescomysis, 295
Bachelotia, 27
Baculellidae, 63
Baeolidia, 204

- Baeonectes*, 317
Baffinia, 230
Baffinicythere, 248
Bagginidae, 69
Baicallelia, 130
Bairdiidae, 247
Bairdioidea, 247
Bajacalifornia, 362
Balaena, 376
Balaenamertes, 156
Balaenamertidae, 156
Balaenidae, 376
Balaenophilidae, 270
Balaenophilus, 270
Balaenoptera, 376
Balaenopteridae, 376
Balanidae, 282
Balaniger, 76
Balaninae, 282
Balanoglossus, 336
Balanoidae, 282
Balanomorpha, 282
Balanophyllia, 110
Balanus, 282
Balbidiocolon, 316
Baldia, 216
Balgetia, 130
Balhochordaes, 356
Balistes, 374
Balistidae, 374
Balliella, 30
Balssia, 288
Baltalimania, 126
Baltoplana, 131
Bangia, 29
Bangiaceae, 29
Bangiiales, 29
Bankia, 210
Bantariella, 332
Baptodoris, 203
Barathrites, 366
Barathronus, 366
Barbantus, 362
Barbatia, 206
Barbonema, 162
Barcinididae, 114
Barcino, 114
Barentiidae, 179
Barentsia, 179
Bargmannia, 122
Barleeia, 190
Barleeidae, 190
Barnea, 210
Barrowgamma, 298
Barya, 17
Baseodiscus, 154
Basidiomycetes, 18
Basidiomycota, 18
Basilissopsis, 184
Basipodella, 280
Basipodellidae, 280
Basisulcata, 196
Bassia, 121
Basslerites, 250
Bassozetus, 366
Batheuchaeta, 263
Batheulima, 187
Bathophilus, 363
Bathothauma, 213
Bathoxiphus, 212
Bathyarca, 206
Bathybela, 196
Bathybiaster, 338
Bathycalanus, 266
Bathycampitus, 270
Bathychaetus, 221
Bathyconchoecia, 245
Bathycopea, 320
Bathycreadium, 137
Bathycrinicola, 187
Bathyocrinidae, 338
Bathycrinus, 338
Bathycura, 309
Bathydorididae, 201
Bathydoris, 201
Bathydorus, 103
Bathydrilus, 233
Bathyechiniscus, 236
Bathyeliasona, 224
Bathyepsilon, 162
Bathyfauvelia, 224
Bathygadus, 365
Bathygnathia, 321
Bathyhalacarus, 240
Bathyidia, 279
Bathylaceo, 362
Bathylagichthys, 361
Bathylagidae, 361
Bathylagus, 361
Bathyaimus, 162
Bathylophonte, 275
Bathylasma, 282
Bathylasmatidae, 282
Bathylasmatinae, 282
Bathylychnops, 361
Bathymacrostomum, 127
Bathymedon, 305
Bathymicrops, 363
Bathymodiolus, 207
Bathymonorchis, 137
Bathymysis, 293
Bathynectes, 291
Bathynocythere, 247
Bathynoe, 224
Bathyoncus, 354
Bathyonus, 366
Bathyopsurus, 317
Bathypalaemonella, 288
Bathypalaemonellidae, 288
Bathypathes, 109
Bathypectinura, 346
Bathyphellia, 107
Bathyphelliidae, 107
Bathyphotis, 298
Bathyphysa, 122
Bathyplotes, 349
Bathypolypus, 213
Bathypongia, 264
Bathypontiidae, 264
Bathyporeia, 306
Bathyprion, 362
Bathypammis, 277
Bathypterois, 363
Bathytyra, 354
Bathyraja, 359
Bathysaurus, 363
Bathysciadiidae, 182
Bathysciadium, 182
Bathysiphon, 62
Bathysolea, 374
Bathyspinula, 206
Bathysquilla, 284
Bathysquillidae, 284
Bathystyeloides, 354
Bathytaenassis, 315
Bathyteuthidae, 212
Bathyteuthis, 212
Bathytroctes, 362
Bathytyphlops, 363
Bathyuroconger, 361
Bathyvermilia, 226
Batillipedidae, 236
Batillipes, 236
Batophora, 23
Batrachoididae, 366
Batrachoidiformes, 366
Bdelloida, 151
Bdellonemertini, 156
Bdellouridae, 136
Beania, 326
Beaniidae, 326
Bebryce, 105
Beckerella, 35
Bedotella, 115
Beella, 73
Beklemisheviella, 134
Bela, 195
Belbolla, 162
Bellibos, 317
Bellottia, 366
Belomitra, 193
Belone, 367
Belonectes, 317
Belonidae, 367
Beloniformes, 367
Bemlos, 298
Benedenia, 143
Benedenipora, 332
Benedeniporidae, 332
Benthabella, 363
Bentharca, 206
Benthogennema, 285
Benthescymidae, 285
Benthescymus, 285
Bentheuphausia, 284
Bentheuphausiidae, 284
Benthobia, 194
Benthocometes, 366
Benthoctopus, 213
Benthodesmus, 373
Benthodytes, 350
Benthogone, 350
Benthomangelia, 195
Benthomisopria, 262
Benthonectes, 286
Benthonella, 189
Benthonellania, 189
Benthpecten, 339
Benthopectinidae, 339
Benthophiloides, 372
Benthophilus, 372

- Benthosema*, 364
Benthothuria, 349
Berghia, 204
Beringius, 193
Bermudella, 202
Beroe, 122
Beroida, 122
Beroidae, 122
Berthella, 201
Berthellina, 201
Bertiliella, 130
Bertiliellidae, 130
Berycidae, 367
Beryciformes, 367
Beryx, 367
Bhawania, 216
Biancolina, 298
Biancolinidae, 298
Bicellariella, 327
Bicellarina, 327
Biconiosporella, 18
Bicosoeca, 77
Bicosoecids, 77
Bicosta, 57
Bicrisia, 333
Biemna, 97
Biflua, 17
Bifurcaria, 27
Bifurgaster, 283
Bigelowina, 284
Bigenerina, 62
Biloculinella, 66
Bimeria, 117
Bipteria, 83
Biserramenia, 182
Bispira, 225
Bitectiporidae, 326
Bitias, 287
Bittium, 185
Bivagina, 143
Bivalvia, 205
Bivalvulida, 82
Blackfordia, 114
Blackfordiidae, 114
Blastophysa, 21
Bledius, 324
Blenniidae, 372
Blennius, 372
Blepharocysta, 47
Blidingia, 24
Boccardia, 227
Boccardiella, 227
Boderia, 62
Bodianus, 371
Bodo, 60
Bodotria, 309
Bodotriidae, 309
Bodotriinae, 309
Boecksinus, 302
Boergesenella, 32
Bogasonia, 199
Bogia, 183
Bogidiella, 298
Bogidiellidae, 298
Bolbocoleon, 21
Bolbosoma, 176
Bolinichthys, 364
Bolinopsidae, 123
Bolinopsis, 123
Bolinus, 192
Bolitaenidae, 213
Bolivina, 69
Bolivinella, 70
Bolivinidae, 69
Bolivinitidae, 70
Bolma, 185
Bolocera, 106
Boloceroididae, 107
Boltenia, 354
Bolteniopsis, 354
Bomolochidae, 255
Bomolochus, 255
Bonapartia, 362
Bonellia, 178
Bonelliidae, 178
Bonnemaisonaria, 30
Bonnemaisoniaceae, 30
Bonnemaisoniales, 30
Bonneviella, 119
Bonneviellidae, 119
Bonnierella, 301
Bonnierilla, 254
Boodea, 23
Boodeaceae, 22, 23
Boops, 370
Bopyridae, 322
Bopyrina, 322
Bopyrissa, 322
Bopyroides, 322
Bopyrus, 322
Bordnamonas, 78, 79
Boreadinium, 47
Boreocelis, 135
Boreocingula, 189
Boreogadus, 365
Boreohydra, 112
Boreohydridae, 112
Boreolimella, 270
Boreolithon, 33
Boreomicralaimoides, 162
Boreomysis, 293
Boreonymphon, 242
Boreopontia, 271
Boreotantulus, 280
Bornetia, 30
Bornia, 208
Borodinellaceae, 23
Boroecia, 245
Borostomias, 363
Borsonia, 195
Boschmaella, 283
Bosellia, 200
Boselliidae, 200
Bosmina, 243
Bosminidae, 243
Bosquetina, 250
Bostrychia, 32
Botachus, 254
Botellina, 63
Bothidae, 374
Bothitrema, 143
Bothrimonus, 146
Bothriocephalus, 146
Bothriomolus, 135
Bothus, 374
Botracnidifer, 108
Botrucnidiferidae, 108
Botryllophilidae, 253
Botryllophilus, 253
Botryllus, 354
Botrynema, 119
Botyocladia, 38
Botryphallus, 190
Botrytella, 27
Botrytis, 17
Botulus, 137
Bougainvillia, 117
Bougainvilliidae, 117
Bourgueticrinida, 337
Bourgueticrinidae, 337
Bouvierella, 299
Bovicornua, 16
Bowerbankia, 332
Boxshallia, 262
Brachidontes, 206
Brachiella, 260
Brachionidae, 150
Brachionis, 150
Brachioteuthidae, 212
Brachioteuthis, 212
Brachycalanus, 266
Brachycarpus, 288
Brachydeuterus, 370
Brachydiastylis, 309
Brachydinium, 47
Brachyenteron, 137
Brachylaima, 137
Brachylecithum, 137
Brachynotus, 292
Brachyphallus, 137
Brachyscelidae, 308
Brachyscelus, 308
Brada, 218
Bradya, 273
Bradyagae, 240
Bradybucca, 162
Bradycalanus, 266
Bradyelopsis, 273
Bradyetes, 263
Bradyidius, 263
Bradynectes, 127
Bradyponius, 259
Brama, 370
Bramidae, 370
Branchellion, 234
Branchiocerianthus, 112
Branchioglossum, 32
Branchiomaldane, 216
Branchiomma, 225
Branchiostoma, 357
Branchiostomidae, 357
Branchiosyllis, 229
Branchiura, 243
Brania, 229
Braunina, 137
Brederveldia, 134
Bregmaceros, 365
Bregmacerotidae, 365
Breizacanthus, 176
Brementia, 254
Bresilia, 286

- Bresiliidae**, 286
Bresslauilla, 130
Brevinucula, 205
Brianola, 271
Brinkmannia, 155
Brinkmanniella, 133
Brinkmanniidae, 155
Brisaster, 347
Brisinga, 342
Brisingella, 342
Brisingida, 342
Brisingidae, 342
Brissidae, 347
Brissopsis, 347
Brissus, 347
Brizalina, 70
Brocchinia, 194
Brongniartella, 32
Brookesena, 196
Brooksia, 355
Broscus, 323
Brosme, 365
Brotulotaenia, 366
Brumptiana, 234
Brunetia, 131
Bruzelia, 307
Brychiopontiidae, 259
Brychiopontius, 259
Bryobesia, 22
Bryocryptella, 327
Bryocryptellidae, 327
Bryopsidaceae, 21
Bryopsidales, 21
Bryopsisella, 21
Bryopsis, 21
Bryozoa, 326
Bubaridae, 90
Bubaris, 90
Bubaropsis, 92
Buccella, 71
Buccinidae, 193
Buccinulum, 193
Buccinum, 193
Bucephalopsis, 137
Bucephalus, 137
Buchneria, 330
Buenia, 372
Buergeriella, 156
Buergeriellidae, 156
Buffhamia, 26
Buffhamiaceae, 26
Buffonellaria, 328
Buglossidium, 374
Bugula, 327
Bugulella, 327
Bugulidae, 327
Bulbamphiascius, 272
Bulbella, 332
Bulbocephalus, 175
Bulbotopla, 135
Bulbus, 191
Bulimeulima, 187
Bulimina, 70
Buliminella, 75
Buliminidae, 70
Bulla, 199
Bullidae, 199
Bullowanthura, 316
Bulweria, 375
Bunocotyle, 137
Bunodactis, 106
Bunodeopsis, 107
Bunoderia, 137
Bunodosoma, 106
Buntonia, 250
Buproridae, 253
Buprorus, 253
Burrimysis, 293
Bursa, 192
Bursatella, 201
Bursidae, 192
Bursosaphia, 126
Bursovaginoidae, 151
Bushiella, 228
Buskea, 328
Buskia, 332
Buskiidae, 332
Byblis, 297
Bylgides, 224
Byrsophlebidae, 133
Byrsophlebs, 133
Byssochlamys, 16
Bythites, 366
Bythitidae, 366
Bythocaris, 287
Bythoceratina, 247
Bythocyprididae, 247
Bythocyparis, 247
Bythocythere, 247
Bythocytheridae, 247
Bythograeidae, 292
Bythotiara, 117
Caballerocotyla, 143
Caberea, 328
Cabestana, 192
Cachonina, 47
Cachonodium, 47
Cacospongia, 100
Cactosoma, 107
Cadenia, 143
Cadlina, 202
Cadulus, 212
Caecidae, 190
Caecijaera, 318
Caecitellus, 77
Caecognathia, 321
Caeconyx, 302
Caecosagitta, 335
Caecostenetroides, 318
Caecum, 190
Caelorinchus, 365
Caenopedia, 346
Caesaromyia, 293
Cafeteria, 77
Cainocreadium, 137
Calamyzas, 216, 229
Calamyzidae, 216
Calanidae, 264
Calanipedia, 267
Calanoida, 263
Calanoides, 264
Calanopia, 267
Calanticidae, 281
Calanus, 264
Calappa, 290
Calappidae, 290
Calastacus, 289
Calathura, 316
Calcarea, 101
Calcaronea, 101
Calceostoma, 143
Calceostomella, 143
Calcinea, 101
Calcinus, 289
Calcidinellum, 47
Calcituba, 68
Calescharidae, 327
Calicotyle, 143
Caligidae, 259
Caligodes, 259
Caligopsis, 259
Caligus, 259
Caliphylla, 201
Callanthias, 369
Callanthiidae, 369
Calliacantha, 57
Callactis, 107
Callianassa, 289
Callianassidae, 289
Callianira, 122
Calliapagurops, 289
Calliax, 289
Calliblepharis, 36
Calligyrus, 162
Callinectes, 291
Callinera, 152
Calliobdella, 234
Calliobothrium, 146
Callionymidae, 372
Callionymus, 372
Calliopea, 201
Calliopus, 299
Callianidae, 128
Callistoma, 184
Calliotropis, 184
Callipallene, 242
Callipallenidae, 242
Callista, 210
Callistochiton, 182
Callistocythere, 249
Callitetrarhynchus, 146
Callithamniella, 30
Callithamnion, 30
Callochiton, 182
Callocolax, 35
Callogorgia, 105
Callophyllis, 35
Callopora, 327
Calloporidae, 327, 330
Calloporina, 330
Callostracon, 198
Callumbonella, 184
Callyspongia, 98
Callyspongiidae, 98
Calma, 204
Calmella, 205
Calmidae, 204
Calocalanus, 266
Calocarides, 289
Calocarididae, 289
Calocaris, 289

- Calonectris*, 375
Calonemertes, 156
Caloria, 204
Calosiphonaceae, 36
Calosiphonia, 36
Calozodion, 313
Calpensia, 330
Calthropella, 85
Calvieriosoma, 346
Calverocheres, 260
Calverocheridae, 260
Calycella, 114
Calycodeles, 137
Calycophorae, 121
Calycopsidae, 117
Calyopsis, 117
Calymne, 347
Calymnidae, 347
Calypsinia, 258
Calyptoconcha, 191
Calyptomma, 293
Calyptotheca, 330
Calyptraea, 190
Calyptracidae, 190
Calyptromema, 162
Calyx, 98
Camacolaimoides, 162
Camacolaimus, 162
Camactognathus, 240
Camallanus, 175
Camarosporium, 19
Caminella, 86
Caminus, 86
Campalecium, 114
Campanoeca, 57
Campanopsis, 114
Campanularia, 119
Campanulariidae, 119
Campanulinidae, 114
Campascus, 56
Campeopea, 320
Campogramma, 370
Campomma, 114
Campula, 137
Campylaimus, 162
Campylaspis, 310
Campyloacantha, 57
Campyloderes, 159
Campylorhaphion, 187
Cancellaria, 194
Cancellariidae, 194
Cancellothyrididae, 334
Cancer, 291
Cancerilla, 260
Cancerillidae, 260
Cancricepon, 322
Cancridae, 291
Cancris, 69
Candacia, 264
Candaciidae, 264
Candelabridae, 112
Candelabrum, 112
Candidae, 328
Candidella, 105
Candonidae, 247
Canetellia, 130
Cantabrina, 87
Canthidermis, 374
Canthigaster, 374
Canthocamptidae, 270
Cantrainea, 185
Canuella, 271
Canuellidae, 271
Canuellina, 271
Canuellopsis, 271
Capillaria, 175
Capitata, 112
Capitella, 216
Capitellidae, 216
Caprella, 296
Caprellidae, 296
Caprellidea, 296
Caproidae, 368
Capronia, 15
Capros, 368
Capsala, 143
Capsaloides, 143
Capsella, 209
Capsosiphon, 23
Capsosiphonaceae, 23
Capulidae, 191
Capulus, 191
Carabidae, 323
Carangidae, 370
Carangoliopsidea, 298
Carangoliopsis, 298
Caranx, 370
Carapidae, 366
Carapus, 366
Carbarea, 329
Carbosphaerella, 16
Carcharhinidae, 359
Carcharhiniformes, 358
Carcharhinus, 359
Carcharias, 358
Carcharodon, 358
Carcharodorhynchus, 132
Carcinonemertes, 154
Carcinonemertidae, 154
Carcinophoridae, 324
Carcinoplacidae, 292
Carcinoplax, 292
Carcinus, 291
Cardiidae, 209
Cardiocephalus, 137
Cardiodesetes, 262
Cardioecia, 333
Cardiomya, 211
Cardita, 208
Carditidae, 208
Cardium, 209
Carencoilia, 135
Carenzia, 184
Careproctus, 369
Caretta, 375
Caridion, 287
Carinaria, 192
Carinariidae, 192
Carinesta, 152
Carinina, 152
Carinocythereis, 250
Carinoma, 152
Carinomidae, 152
Caristiidae, 370
Caristius, 370
Carlgrenia, 107
Carminodoris, 203
Carnivora, 376
Carobia, 222
Carpias, 318
Carpoapseudes, 313
Carpomitra, 29
Carybdea, 111
Carybdeidae, 111
Caryophyllaeus, 146
Caryophyllia, 109
Caryophylliidae, 109
Caspihalacarus, 240
Casiopoma, 372
Cassidae, 191
Cassidulina, 70
Cassidulinidae, 70
Cassidulinoidea, 70
Cassiduloida, 347
Cassiella, 185
Cassiopea, 111
Cassiopeidae, 111
Castrada, 134
Catablema, 118
Cataetyx, 366
Catapaguroides, 289
Cataphellia, 107
Catenella, 36
Catenulida, 127
Catharacta, 376
Cathariotrema, 143
Catherinum, 282
Catinia, 255
Catiniidae, 255
Catostylidae, 112
Catostylus, 112
Catriona, 205
Caudites, 249
Caulacanthaceae, 36
Caulacanthus, 36
Caulaster, 339
Caulerpa, 21
Caulerpaceae, 21
Caulerpales, 21
Caulieriella, 216
Caulophacidae, 103
Caulophacus, 103
Caulophryne, 366
Caulophrynidae, 366
Cauloramphus, 327
Causia, 64
Cavernularia, 106
Cavolinia, 200
Cavoliniidae, 200
Cecropidae, 260
Cecrops, 260
Cellana, 182
Cellaria, 328
Cellariidae, 328
Cellepora, 328
Celleporaria, 328
Celleporella, 330
Celleporidae, 328
Celleporina, 328
Celtia, 250

- Cemocotyle*, 143
Centracanthidae, 370
Centracanthus, 370
Centraugaptilus, 264
Centriscidae, 368
Centrobranchus, 364
Centroceras, 30
Centrocerocolax, 30
Centroderes, 159
Centroderidae, 159
Centroderma, 137
Centrodinium, 47
Centrodraco, 372
Centrolabrus, 371
Centrolophidae, 373
Centrolophus, 373
Centromedon, 302
Centropages, 265
Centropagidae, 265
Centrophoridae, 359
Centrophorus, 359
Centrophryne, 367
Centrophrynidae, 367
Centropyxidae, 55, 56
Centropyxiella, 56
Centropyxis, 55
Centrosyllium, 359
Centrosymnus, 359
Centrostephanus, 346
Cephalobaenida, 251
Cephalochordata, 357
Cephalodasys, 157
Cephalodella, 150
Cephalophanes, 266
Cephalopoda, 212
Cephalopyge, 203
Cephalorhyncha, 159
Cephalosporium, 19
Cephalothricidae, 152
Cephalothrix, 152
Cephalotrichella, 152
Cephalotrichum, 19
Cepheidae, 111
Cepola, 371
Cepolidae, 371
Cephus, 375
Ceractinomorpha, 89
Ceradocus, 304
Ceramaster, 339
Ceramiaceae, 30
Ceramiales, 30
Ceramium, 30
Ceramonema, 162
Cerapus, 301
Cerastoderma, 209
Ceratia, 190
Ceratias, 367
Ceratiidae, 367
Ceratium, 47
Ceratobothrium, 146
Ceratobulimina, 71
Ceratobuliminidae, 71
Ceratocancris, 71
Ceratocaulon, 105
Ceratocephale, 220
Ceratochona, 41
Ceratocolax, 36
Ceratocorys, 48
Ceratocuma, 309
Ceratocumatidae, 309
Ceratocymba, 121
Ceratomyxa, 83
Ceratomyxidae, 83
Ceratonereis, 220
Ceratonotus, 269
Ceratopera, 134
Ceratoperidinium, 48
Ceratoscopelus, 364
Ceratothoa, 320
Ceratotrochus, 109
Cerbaris, 90
Cerberilla, 204
Cerbussovicia, 136
Cercariooides, 137
Cercyra, 136
Cercyridae, 136
Cerebratulidae, 153
Cerebratulus, 153
Cerelasma, 78
Cerelasmidae, 78
Cereus, 108
Ceriantharia, 108
Cerianthidae, 108
Cerianthus, 108
Ceriosporopsis, 16
Cerithiella, 186
Cerithiidae, 185
Cerithiopsidae, 185
Cerithiopsis, 185
Cerithium, 185
Cernosvitoviella, 234
Ceroctena, 122
Cervera, 105
Cervinia, 271
Cerviniella, 271
Cerviniidae, 271
Cerviniopsis, 271
Cervonema, 162
Cestida, 122
Cestidae, 122
Cestoda, 146
Cestopagurus, 289
Cestoplana, 128
Cestoplanidae, 128
Cestum, 122
Cetacea, 376
Cetoconcha, 211
Cetomimidae, 368
Cetomimiformes, 368
Cetomimus, 368
Cetomurus, 365
Cetorhinidae, 358
Cetorhinus, 358
Cetostoma, 368
Chaceon, 292
Chaedofaudia, 16
Chaenophryne, 366
Chaetaster, 339
Chaetasteridae, 339
Chaetiliidae, 321
Chaetoderma, 181
Chaetodermatidae, 181
Chaetodon, 371
Chaetodontidae, 371
Chaetodoryx, 94
Chaetogaster, 233
Chaetognatha, 335
Chaetomastia, 17
Chaetomiaceae, 18
Chaetomium, 18
Chaetomorpha, 22
Chaetonema, 162
Chaetonotidae, 156
Chaetonotoidea, 156
Chaetonotus, 156
Chaetoparia, 222
Chaetopeltis, 24
Chaetophoraceae, 21
Chaetophorales, 21
Chaetopleura, 182
Chaetopteridae, 216
Chaetopterus, 216
Chaetosiphon, 21
Chaetosiphonaceae, 21
Chaetosphaeria, 18
Chaetosphaeriaceae, 18
Chaetostephanidae, 160
Chaetozone, 216
Chalarostylis, 309
Chalinidae, 98
Chalinopsilla, 98
Chalinula, 98
Chama, 208
Chamelea, 210
Chamidae, 208
Champia, 38
Champiaceae, 38
Chaperiidae, 328
Chaperiopsis, 328
Characella, 86
Characium, 22
Characodoma, 328
Charadriformes, 375
Chardezia, 56
Charistephane, 122
Charonia, 192
Charopinus, 260
Chartella, 329
Charybdis, 291
Chauhanea, 143
Chauliodus, 363
Chaunacidae, 366
Chaunax, 366
Chaunoteuthis, 212
Chauvetia, 193
Chauviniella, 214
Cheilopogon, 367
Cheilopora, 328
Cheioporina, 328
Cheioporinidae, 328
Cheilostomatida, 326
Cheiraster, 339
Cheirocratus, 304
Cheirodonta, 185
Cheironchus, 162
Cheiropotis, 300
Chelator, 317
Chelidonichthys, 368
Chelidonisis, 105
Chelidonura, 199
Cheliplana, 131

- Cheliplanilla*, 131
Chelon, 371
Chelonaplysilla, 101
Chelonia, 375
Chelonibia, 282
Cheloniidae, 282
Cheloniidae, 375
Chelophyes, 121
Chelura, 298
Cheluridae, 298
Chelyosoma, 352
Cherbonniera, 351
Chevreuxius, 298
Chevroderma, 181
Chiasmodon, 372
Chiasmodontidae, 372
Childia, 125
Childiidae, 125
Chilionema, 26
Chilodochona, 41
Chilodochonidae, 41
Chilodochonina, 41
Chilomonas, 59
Chilomycterus, 374
Chilophiurina, 345
Chilopoda, 323
Chilostomella, 74
Chilostomellidae, 71
Chimaera, 360
Chimaerarhynchus, 146
Chimaericola, 143
Chimaeridae, 360
Chimaeriformes, 360
Chimaerocestos, 146
Chimaerohemecus, 137
Chionoecetes, 291
Chiridiella, 263
Chiridius, 263
Chiridota, 351
Chiridotea, 321
Chiridotidae, 351
Chirimia, 219
Chirolophis, 371
Chirona, 282
Chironomidae, 324
Chirotomias, 363
Chirotylidae, 290
Chiroteuthidae, 213
Chiroteuthis, 213
Chirundina, 263
Chitinopoma, 226
Chiton, 182
Chitonidae, 182
Chitwoodia, 162
Chlamydoselachidae, 358
Chlamydoselachus, 358
Chlamys, 207
Chlidonia, 328
Chlidonias, 375
Chlidoniidae, 328
Chloeia, 215
Chloenopsis, 215
Chlopsidae, 360
Chlopsis, 360
Chloraster, 79
Chlorochytrium, 22
Chlorococcaceae, 22
Chlorococcales, 22
Chlorococcum, 22
Chlorocystidaceae, 22
Chlorocystidales, 22
Chloromyxidae, 84
Chloromyxum, 84
Chloronemella, 162
Chlorophthalmidae, 363
Chlorophthalmus, 363
Chlorophyceae, 21
Chlorophycota, 21
Chlorotocus, 287
Choanoeca, 58
Choanoflagellata, 57
Choanolaimus, 162
Cholidya, 279
Cholidyella, 279
Chondracanthidae, 255
Chondracanthodes, 255
Chondracanthus, 36, 255
Chondraster, 340
Chondria, 32
Chondrichthyes, 358
Chondrilla, 89
Chondrillidae, 89
Chondrocladia, 97
Chondrophellia, 107
Chondrophycus, 33
Chondrosia, 89
Chondrosida, 89
Chondrus, 36
Chondrymenia, 37
Chone, 225
Chonelasma, 102
Choniolaimus, 162
Choniorhiza, 261
Choniosphaera, 261
Choniostoma, 261
Chonosaurus, 41
Chonotrichia, 41
Chorda, 28
Chordaceae, 28
Chordaria, 24
Chordariaceae, 24
Chordariales, 24
Chordeumiidae, 253
Chordeumium, 253
Chordodasyidae, 158
Choreocolacaceae, 36
Choreocolax, 36
Choreonema, 33
Choricotyle, 143
Choristella, 183
Choristellidae, 183
Choristocarpaceae, 28
Choristocarpus, 28
Chorizopora, 328
Chorizoporidae, 328
Chorocaris, 286
Chrisomon, 137
Christerius, 234
Christianella, 146
Chromadora, 162
Chromadorella, 162
Chromadorina, 162
Chromadorissa, 163
Chromadorita, 163
Chromaspirina, 163
Chromis, 371
Chromodorididae, 202
Chromodoris, 202
Chromogobius, 372
Chroodactylon, 37
Chroomonas, 59
Chroothece, 38
Chrotella, 87
Chrysopodium, 19
Chrysallida, 197
Chrysaora, 111
Chrysochromulina, 76
Chrysogorgia, 104
Chrysogorgiidae, 104
Chrysopetalidae, 216
Chrysopetalum, 216
Chrysomenia, 38
Ctenopterygidae, 212
Ctenopteryx, 212
Chthamalidae, 282
Chthamalinae, 282
Chthamaloidea, 282
Chthamalophilidae, 283
Chthamalophilus, 283
Chthamalus, 282
Chuniella, 156
Chuniellidae, 156
Chuniphyes, 121
Chunomysis, 293
Chylocladia, 38
Cibicides, 69
Cibicididae, 71
Cibicidoides, 69
Cicerina, 130
Cicerinidae, 130
Cidaridae, 346
Cidaris, 346
Cidaroida, 346
Cigclisula, 331
Ciliata, 365
Ciliatocardium, 209
Cilicaea, 320
Ciliocincta, 104
Ciliopharyngiella, 133
Ciliopharyngiellidae, 133
Cilunculus, 242
Cima, 197
Cimidae, 197
Cimmerius, 309
Cinachyra, 87
Cinetorhynchus, 288
Cingula, 189
Cingulina, 197
Cingulopsidae, 188
Ciocalapata, 90
Ciocalypta, 90
Ciona, 352
Cionidae, 352
Circeis, 228
Circulus, 190
Cirkenneya, 137
Cirolana, 319
Cirolanidae, 319
Cirratulidae, 216
Cirratulus, 216
Cirrenalia, 19

<i>Cirrhinereis</i> , 217	Claviporidae , 332	<i>Cochlodesma</i> , 211
<i>Cirrifera</i> , 135	<i>Claviscala</i> , 186	<i>Cochlodinium</i> , 48
<i>Cirriformia</i> , 217	<i>Clavodesia</i> , 327	Codiaceae , 23
<i>Cirripedia</i> , 281	<i>Clavodorum</i> , 227	Codiales , 23
<i>Cirrophorus</i> , 222	<i>Clavopora</i> , 332	<i>Codium</i> , 23
<i>Cirroteuthidae</i> , 213	Clavoporidae , 332	<i>Codonellina</i> , 326
<i>Cirrothauma</i> , 213	<i>Clavopsella</i> , 117	<i>Codonocladium</i> , 58
<i>Cirsonella</i> , 184	<i>Clavularia</i> , 104	<i>Codonorchis</i> , 118
<i>Cirsotrema</i> , 186	Clavulariidae , 104	<i>Codosiga</i> , 58
<i>Cistacythereis</i> , 251	<i>Cleantis</i> , 321	Codosigidae , 58
<i>Citharidae</i> , 374	<i>Cleidochasma</i> , 328	<i>Coeliotrema</i> , 137
<i>Citharistes</i> , 48	Cleidochasmatidae , 328	<i>Coelocalypta</i> , 90
<i>Citharus</i> , 374	<i>Cleidochasmidra</i> , 328	<i>Coelocladia</i> , 26
<i>Ciureana</i> , 137	<i>Cleippides</i> , 299	<i>Coelogynopora</i> , 135
<i>Cladocarpus</i> , 114	<i>Cleithrarticus</i> , 143	Coelogynoporidae , 135
<i>Cladocera</i> , 243	<i>Clelandella</i> , 183	<i>Coelopleurus</i> , 346
<i>Cladocora</i> , 109	<i>Clementia</i> , 210	<i>Coelosphaera</i> , 94
<i>Cladocoryne</i> , 112	<i>Cleonardo</i> , 299	Coelosphaeridae , 94
<i>Cladocorynidiae</i> , 112	<i>Cleonardopsis</i> , 306	<i>Coelothrix</i> , 38
<i>Cladococe</i> , 98	<i>Clestobothrium</i> , 146	<i>Coenocyathus</i> , 109
<i>Cladonema</i> , 112	<i>Cletocampus</i> , 270	<i>Coenosimilia</i> , 109
<i>Cladonematidae</i> , 112	<i>Cletodidae</i> , 271	Coilodesmaceae , 26
<i>Cladophora</i> , 22	Cletopsyllidae , 271	<i>Coilodesme</i> , 26
<i>Cladophoraceae</i> , 22	<i>Clibanarius</i> , 289	<i>Colacodictyon</i> , 37
<i>Cladophorales</i> , 22	<i>Climacosorus</i> , 27	<i>Colegynopora</i> , 135
<i>Cladophoropsis</i> , 23	Clinidae , 372	Coleoptera , 323
<i>Cladopsammia</i> , 110	<i>Clinitrichus</i> , 372	<i>Colga</i> , 203
<i>Cladopyxis</i> , 48	<i>Clio</i> , 200	<i>Collarina</i> , 329
<i>Cladorhiza</i> , 97	<i>Cliona</i> , 87	<i>Collastoma</i> , 130
Cladorbizidae , 97	<i>Clione</i> , 200	Collembola , 324
<i>Cladosiphon</i> , 24	Clionidae , 87, 200	<i>Colletotrichum</i> , 19
<i>Cladosporium</i> , 19	<i>Cliopsis</i> , 200	<i>Collettea</i> , 314
<i>Cladostephus</i> , 28	<i>Cliothosa</i> , 88	<i>Colligocineta</i> , 46
<i>Clanculus</i> , 183	<i>Clipeodinium</i> , 48	<i>Collocheres</i> , 259
<i>Clangula</i> , 375	Clistosaccidae , 283	<i>Collotheca</i> , 151
<i>Claparedepelogenia</i> , 227	<i>Clistosaccus</i> , 283	<i>Collothecaceae</i> , 151
<i>Clathrella</i> , 197	<i>Clitellio</i> , 233	<i>Colobocephalus</i> , 199
<i>Clathria</i> , 92	<i>Cloacotaenia</i> , 146	<i>Coloboceras</i> , 240
<i>Clathrina</i> , 101	<i>Clunio</i> , 324	<i>Colobocerasides</i> , 240
Clathrinida , 101	<i>Clupea</i> , 361	<i>Colobomatooides</i> , 257
Clathrinidae , 101	Clupeidae , 361	<i>Colobomatus</i> , 257
<i>Clathrodiscus</i> , 26	Clupeiformes , 361	<i>Colobonema</i> , 119
<i>Clathrofrenella</i> , 185	<i>Clupeonella</i> , 361	Colomastigidae , 298
<i>Clathromangelia</i> , 195	<i>Cluthia</i> , 249	<i>Colomastix</i> , 298
<i>Clathromorphum</i> , 33	<i>Clydonella</i> , 54	<i>Colombia</i> , 320
<i>Clausia</i> , 255	<i>Clymenella</i> , 219	Coloniales , 179
Clausidiidae , 255	<i>Clymenopsis</i> , 219	Colossendeidae , 242
Clausiidae , 255	<i>Clymenura</i> , 220	Colossendeis , 242
<i>Clausinella</i> , 210	Clypeasteroida , 347	<i>Colpaster</i> , 342
Clausocalanidae , 265	<i>Clypeomorus</i> , 185	<i>Colpodaspis</i> , 199
<i>Clausocalanus</i> , 265	<i>Clyporhynchus</i> , 132	<i>Colpodella</i> , 40
<i>Clausophyes</i> , 121	<i>Clytemnestra</i> , 271	<i>Colpomenia</i> , 28
Clausophyidae , 121	Clytemnestridae , 271	<i>Colubraria</i> , 193
<i>Clava</i> , 117	<i>Clytia</i> , 119	<i>Columbella</i> , 194
<i>Clavagella</i> , 211	<i>Cnemidocarpa</i> , 354	Columbellidae , 194
Clavagellidae , 211	<i>Cobancythere</i> , 247	<i>Colurella</i> , 150
<i>Clavariopsis</i> , 19	<i>Cobia</i> , 163	<i>Colymbilepis</i> , 146
<i>Clavelina</i> , 352	<i>Cobbionema</i> , 163	<i>Comantenna</i> , 263
Clavelinidae , 352	<i>Coboldus</i> , 300	<i>Comarmondia</i> , 195
<i>Clavella</i> , 260	<i>Cocomyxha</i> , 22, 83	Comasteridae , 338
<i>Clavellisa</i> , 260	<i>Coccopigya</i> , 182	Comatulida , 338
<i>Clavellistes</i> , 260	<i>Coccarella</i> , 364	<i>Comesoma</i> , 163
<i>Clavelloides</i> , 260	<i>Coccotylus</i> , 36	<i>Commensodorum</i> , 227
<i>Clavellotis</i> , 260	<i>Cocculina</i> , 182	<i>Communoporus</i> , 154
<i>Claviceps</i> , 17	Cocculinidae , 182	<i>Comoplana</i> , 129
Clavicipitaceae , 17		
Clavidae , 117		

- Complexoburda*, 137
Compsonema, 28
Compsopogon, 33
Compsopogonaceae, 33
Compsopogonales, 33
Comsothamnion, 30
Conaea, 257
Conaperta, 126
Concentrotheca, 109
Conchoheres, 258
Conchodelphys, 254
Conchoderma, 281
Conchoecetta, 245
Conchoecia, 245
Conchoecilla, 245
Conchoecissa, 245
Conchyliurus, 255
Condylactis, 106
Condylanthidae, 107
Condylotyota, 137
Condyloderes, 159
Confluaria, 146
Conger, 361
Congericola, 260
Conglrophragmium, 63
Congridae, 361
Conica, 113
Conidae, 195
Conilera, 319
Coninckia, 163
Coniochaeta, 18
Coniochaetaceae, 18
Coniothyrium, 19
Connexochiton, 182
Conocara, 362
Conochilidae, 151
Conochilus, 151
Conocrinus, 337
Conocyema, 104
Conocyemidae, 104
Conopea, 282
Conopeum, 329
Conopleura, 196
Conorbella, 73
Contarinia, 35
Contracaecum, 175
Conus, 196
Convolutella, 125
Convoluta, 126
Convolutidae, 125
Cookeolus, 369
Coolia, 48
Copepoda, 253
Copiatestes, 137
Copidognathides, 240
Copidognathinae, 240
Copidognathus, 240
Copidozoum, 327
Copilia, 258
Coppatiidae, 87
Copulabyssia, 183
Corallicletodes, 270
Coralliidae, 104
Corallimorpharia, 108
Corallimorphidae, 108
Corallimorphus, 108
Corallina, 33
Corallinaceae, 33
Corallinales, 33
Coralliodrilus, 233
Coralliope, 292
Coralliphaga, 210
Coralliphila, 192
Coralliphilidae, 192
Corallistes, 87
Corallistidae, 87
Corallium, 104
Corallomyxa, 54
Corallophila, 30
Corambe, 201
Corambidae, 201
Corbula, 210
Corbulaseta, 275
Corbulidae, 210
Corycrogobius, 372
Cordagalma, 122
Cordylecladia, 38
Cordylophora, 117
Coregonus, 362
Corella, 352
Corellidae, 352
Corellopsis, 352
Corinnaeturris, 195
Coris, 371
Cornirostridae, 197
Cornucalanus, 266
Cornularia, 105
Cornulariidae, 105
Cornuloculina, 68
Cornulotrocha, 93
Cornulum, 93
Cornuspira, 66
Cornuspirella, 66
Cornuspiridae, 66
Cornuspiroides, 66
Corolla, 200
Corollospora, 16
Coronarctidae, 236
Coronarctus, 236
Coronaster, 343
Coronatae, 111
Coronellina, 327
Coronhelmis, 133
Coronida, 284
Coronididae, 284
Coronopharynx, 130
Coronopora, 333
Coronula, 282
Coronulidae, 282
Coronuloidea, 282
Corophiidae, 298
Corophium, 298
Corticium, 85
Cortinariaceae, 18
Cortinariales, 18
Corycaeidae, 256
Corycaeus, 256
Corydendrium, 117
Corymorphia, 112
Corymorphidae, 112
Corynactis, 108
Corynascidia, 352
Coryne, 113
Corynidae, 113
Corynocephalus, 214
Corynomorpha, 35
Corynomorphacea, 35
Corynophlaea, 25
Corynoporella, 327
Corynosoma, 176
Corygalops, 372
Coryphaena, 370
Coryphaenidae, 370
Coryphaenoides, 365
Coryphella, 205
Coryphoblemnus, 372
Corystes, 291
Corystidae, 291
Corythionella, 56
Corythodinium, 48
Corythostoma, 163
Coscinasterias, 342
Coscinopsis, 329
Coscinoderma, 100
Coscinospira, 68
Cosmetira, 115
Cosmocephalus, 175
Cosmoeca, 58
Cosmotriphora, 185
Cossura, 217
Cossuridae, 217
Costa, 251
Costaclis, 187
Costasiella, 201
Costasiellidae, 201
Costellariidae, 194
Cottidae, 369
Cottiniella, 32
Cottunculus, 369
Cottus, 369
Cotyloatlantica, 143
Cotylogaster, 136
Cotylorhiza, 112
Coullia, 275
Coxicerberus, 321
Coximedon, 302
Crambe, 94
Crambeidae, 94
Cranchiidae, 213
Crangon, 288
Crangonidae, 288
Craniata, 334
Craniida, 334
Craniidae, 334
Craniiformea, 334
Cranopsis, 183
Crasiella, 157
Craspedacusta, 118
Craspedotella, 48
Crassadoma, 207
Crassarietellus, 263
Crassatellidae, 209
Crassatina, 209
Crassicauda, 175
Crassimarginatella, 328
Crassolaimus, 163
Crassonemertes, 156
Crassophyllum, 105
Crassopleura, 194
Crassostrea, 207
Cratenemertidae, 154

- Crateristoma*, 46
Craterolophus, 112
Cratostigma, 354
Crebricoma, 46
Crella, 95
Crellastrina, 95
Crellidae, 95
Crellomima, 95
Crenella, 206
Crenidens, 370
Crenilabium, 198
Crenopharynx, 163
Crepidostomum, 137
Crepidula, 190
Crepis, 328
Creseis, 200
Cressa, 298
Cressidae, 298
Cressina, 299
Cribellopora, 330
Cribriolina, 329
Cribelinidae, 329
Cribrinopsis, 106
Criboelphidium, 72
Cribronion, 72
Cribropontius, 259
Cribrostomoides, 64
Cricolaimus, 163
Crimora, 202
Crinagera, 17
Crinoidea, 337
Crinolamia, 187
Crinolina, 58
Crinophtheiros, 187
Crisia, 333
Crisidia, 333
Crisiella, 333
Crisiidae, 333
Crisilla, 189
Cristacoxidae, 271
Cristapseudes, 313
Cristichona, 41
Cristitectus, 175
Crithiona, 64
Crithionina, 64
Critomolgus, 258
Crossaster, 341
Crossobothrium, 146
Crossota, 119
Crouania, 30
Crowcroacaecum, 137
Cruciloculina, 66
Crucispina, 58
Cruoria, 36
Cruoriaceae, 36
Cruoriopsis, 35
Crybelocephalus, 302
Cryodesma, 317
Cryothecononas, 79
Cryptenoplus, 163
Crypthecodinium, 48
Cryptelia, 118
Cryptocelides, 128
Cryptocelidae, 128
Cryptocelis, 128
Cryptocheles, 287
Cryptochiridae, 292
Cryptocoope, 315
Cryptocopinae, 315
Cryptocooides, 315
Cryptocotyle, 137
Cryptodifflugia, 56
Cryptodifflugiidae, 56
Cryptogaster, 283
Cryptogemmida, 41
Cryptolarella, 115
Cryptolaria, 115
Cryptonatica, 191
Cryptonemertes, 154
Cryptonemia, 35
Cryptonemiales, 34
Cryptopelta, 346
Cryptophytes, 59
Cryptopleura, 32
Cryptopontius, 259
Cryptopora, 334
Cryptoporidae, 334
Cryptopsaras, 367
Cryptosoma, 290
Cryptosula, 329
Cryptosulidae, 329
Cryptovalsa, 17
Crystalllogobius, 372
Crystallophyes, 121
Ctena, 207
Ctenella, 122
Ctenellidae, 122
Ctenocalanus, 265
Ctenochelidae, 289
Ctenochirichthys, 366
Ctenodiscidae, 339
Ctenodiscus, 339
Ctenodrilidae, 217
Ctenodrilus, 217
Ctenolabrus, 371
Ctenophora, 122
Ctenopoda, 244
Ctenosiphonia, 33
Ctenostomatida, 332
Cubiceps, 374
Cubomedusae, 111
Cuboza, 111
Cucullanus, 175
Cucumaria, 348
Cucumaridae, 348
Culeolus, 354
Culleopharynx, 133
Cumacea, 309
Cumanotidae, 204
Cumanotus, 204
Cumella, 310
Cumellopsis, 310
Cumoniscus, 280
Cumopsis, 309
Cumulospora, 19
Cuneata, 63
Cuneocythere, 247
Cuneocytheridae, 247
Cunina, 119
Cuninidae, 119
Cupuladria, 329
Cupuladriidae, 329
Curassanthura, 316
Curveulima, 187
Curvolaimus, 163
Cushmanideidae, 247
Cupidaria, 211
Cupidariidae, 211
Cupidella, 114
Cuthona, 205
Cutleria, 25
Cutleriaceae, 25
Cutleriales, 25
Cuvieria, 114
Cuvierina, 200
Cyamichona, 41
Cyamon, 91
Cyanea, 111
Cyaneidae, 111
Cyanophthalma, 155
Cyattonema, 163
Cyathidium, 337
Cyatholaimus, 163
Cyathura, 316
Cyclammina, 64
Cyclaspis, 309
Cyclaspoedes, 309
Cyclicoporidae, 329
Cyclophora, 177
Cyclocanna, 115
Cyclocaris, 302
Cyclocibicides, 71
Cyclocoyla, 143
Cycloforina, 66
Cycloleberis, 245
Cyclope, 193
Cyclopecten, 207
Cyclopetta, 253
Cyclopicina, 253
Cyclopidae, 253
Cyclopina, 254
Cyclopinella, 254
Cyclopiniidae, 253
Cyclopinooides, 254
Cycloplectanum, 143
Cyclopoida, 253
Cycloporella, 254
Cycloporus, 128
Cyclopteridae, 369
Cyclopteropsis, 369
Cyclopterus, 369
Cyclopyxis, 56
Cylchoragida, 159
Cyclorhiza, 257
Cyclosalpa, 355
Cycloscala, 186
Cyclostomatida, 333
Cyclostrematidae, 185
Cycloteuthidae, 212
Cycloteuthis, 212
Cyclothonae, 362
Cydippida, 122
Cyema, 361
Cyematidae, 361
Cyerce, 201
Cylichna, 199
Cylichnidae, 199
Cylichnina, 199
Cylichnium, 199
Cylicolaimus, 163
Cylindraxis, 37

- Cylindriscala*, 186
Cylindrocapon, 19
Cylindrocarpus, 25
Cylindrogullmia, 62
Cylindroleberidae, 245
Cylindromacrostomum, 127
Cylindronannopus, 277
Cylindroporella, 329
Cylindropsyllidae, 271
Cylindropsyllus, 271
Cylindrostoma, 129
Cylindrostomidae, 129
Cylindula, 271
Cymadoceaceae, 39
Cymadusa, 297
Cymatium, 192
Cymatocarpus, 137
Cymbalopora, 71
Cymbaloporetta, 71
Cymbaloporidae, 71
Cymbasoma, 262
Cymbium, 194
Cymbodinium, 48
Cymbomonas, 76
Cymbulia, 200
Cumbuliidae, 200
Cymodoce, 320, 321
Cymodocea, 39
Cyemonomidae, 290
Cyemonomus, 290
Cymopolia, 23
Cymothoa, 320
Cymothoidae, 320
Cynoglossidae, 374
Cynoglossus, 374
Cynoponticus, 361
Cynura, 163
Cyphocaris, 302
Cyphoderia, 56
Cyphoderiidae, 56
Cyphosaccus, 283
Cypraeidae, 191
Cyprideis, 247
Cypridina, 245
Cypridinidae, 245
Cypridoidea, 247
Cyprinodontidae, 367
Cyprinodontiformes, 367
Cyrtocrinida, 337
Cyrtodaria, 210
Cystidicola, 175
Cystiplana, 131
Cystiplanidae, 131
Cystiplex, 131
Cystirete, 131
Cysticidae, 194
Cystisella, 327
Cystisoma, 308
Cystisomatidae, 308
Cystocloniaceae, 36
Cystoclonium, 36
Cystodytes, 353
Cystonectae, 122
Cystoopsis, 175
Cystophora, 376
Cystoseira, 27
Cystoseiraceae, 27
- Cytaeididae**, 117
Cytæis, 117
Cytheromorula, 192
Cythere, 247
Cythereis, 247
Cytherella, 247
Cytherellidae, 247
Cytherelloidea, 247
Cytheretta, 247
Cytherettidae, 247
Cytheridae, 247
Cytheridea, 247
Cytherideidae, 247
Cytherideis, 247
Cytherois, 250
Cytheroma, 247
Cytheromatidae, 247
Cytheromorpha, 249
Cytheropteron, 248
Cytheropteron?, 248
Cytherura, 248
Cytheruridae, 248
Cytocystidae, 131
Cytocystis, 131
Cytopsis, 368
Dacydium, 206
Dactylamblyops, 293
Dactylerythrops, 293
Dactylocalyx, 102
Dactylochirotida, 349
Dactylogyrus, 143
Dactylopodamphiascopsis, 272
Dactylopodella, 277
Dactylopodola, 157
Dactylopodolidae, 157
Dactylopodopsis, 279
Dactylopteridae, 368
Dactylopterus, 368
Dactylopsia, 279
Dactylosioides, 279
Dactylosacus, 62
Dactylospora, 17
Dactylosporaceae, 17
Dactylostylis, 318
Dagda, 163
Daira, 292
Dairella, 308
Dairellidae, 308
Dalatias, 359
Dalatiidae, 359
Dalhousiella, 218
Dallina, 334
Dallinidae, 334
Dalmatogloia, 24
Dalophis, 360
Dalyelliida, 129
Dalyelliidae, 129
Damkaeria, 268
Danaella, 302
Dananemertes, 154
Danielssenia, 278
Danilia, 184
Danorhynchus, 132
Daontesia, 107
Daptonema, 163
Darcythompsonia, 272
Darcythompsoniidae, 272
- Dardanus*, 289
Darwinella, 101
Darwinellidae, 101
Dasmosmilia, 109
Dasya, 32
Dasyaceae, 32
Dasyatidae, 360
Dasyatis, 360
Dasybatotrema, 143
Dasybranchus, 216
Dasycladaceae, 23
Dasycladales, 23
Dasycladus, 23
Dasyella, 32
Dasynemella, 163
Dasynemoides, 164
Dasyrhynchus, 146
Dasyscyphus, 17
Dautzenbergia, 299
Davisia, 83
Deania, 359
Debeauchampiola, 129
Decapoda, 285
Decapterus, 370
Deepstaria, 111
Defrancia, 333
Degocheirocratus, 304
Deima, 350
Deimatidae, 350
Deiopea, 123
Deiopidae, 123
Delamarea, 26
Delamareaceae, 26
Delamarella, 276
Delectona, 88
Delectopecten, 207
Delesseria, 32
Delesseriaceae, 32
Delius, 266
Delphinapterus, 376
Delphinidae, 376
Delphinus, 376
Deltanema, 164
Deltentosteus, 372
Deltocyathoides, 110
Deltocyathus, 109
Democrinus, 337
Demoixys, 254
Demoleus, 261
Demonax, 225
Demonema, 164
Demospongiae, 85
Demoulia, 194
Dendoricella, 94
Dendoricellidae, 94
Dendrilla, 101
Dendrobeania, 327
Dendroceratida, 100
Dendrochirotida, 348
Dendrodasys, 157
Dendrodoa, 355
Dendrodorididae, 203
Dendrodoris, 203
Dendrogaster, 283
Dendrogastridae, 283
Dendrogastriidae, 283
Dendromonocotyle, 143

- Dendromunna*, 316
Dendronotidae, 203
Dendronotus, 203
Dendrophyllia, 110
Dendrophylliidae, 110
Dendrophyra, 63
Dendropodola, 157
Dendropoma, 190
Dendrotiidae, 316
Dendrotion, 316
Dendrotuba, 62
Dendrouterina, 146
Dendroxea, 98
Dendryphiella, 19
Dentaliidae, 211
Dentalina, 73
Dentalium, 211
Dentatonema, 164
Dentex, 370
Denticulella, 164
Dentigloborotalia, 73
Dentomuricea, 105
Deontolaimus, 164
Deontostoma, 164
Deoterthridae, 280
Depastridae, 112
Depastrum, 112
Derbesia, 21
Derhesiaceae, 21
Dercitancorina, 86
Dercitus, 86
Derecrema, 137
Derichthyidae, 361
Derichthys, 361
Dermaptera, 324
Dermateaceae, 17
Dermatocelis, 26
Dermatolithon, 33
Dermatomyzon, 259
Dermochelyidae, 375
Dermochelys, 375
Dermocorynus, 35
Dermomurex, 192
Demonema, 37
Dermophthirius, 143
Derocheilocarididae, 252
Derocheilocaris, 252
Derogenes, 137
Derogenoides, 137
Deropristis, 137
Desdemona, 225
Desmacella, 97
Desmacellidae, 97
Desmacididae, 94
Desmacidon, 94
Desmantidae, 87
Desmanthus, 87
Desmarella, 58
Desmarestia, 26
Desmarestiaceae, 26
Desmarestiales, 26
Desmidocerella, 175
Desmocapsa, 48
Desmodasys, 158
Desmodora, 164
Desmolaimus, 164
Desmophyes, 121
Desmophyllum, 109
Desmopteridae, 200
Desmopterus, 200
Desmoscolex, 164
Desmosoma, 317
Desmosomatidae, 316
Desmosomella, 317
Desmote, 130
Desmotrichum, 26
Desmoxyidae, 90
Detocarcinus, 292
Deutella, 296
Deuterammina, 65
Deuterobaris, 137
Devaleraea, 38
Developayella, 78
Devonia, 208
Dexamine, 299
Dexaminidae, 299
Dexiogonimus, 137
Dexiospira, 228
Dextrogaster, 354
Diacavolinia, 200
Dacria, 200
Diadema, 346
Diademataidae, 346
Diadematoida, 346
Diadumene, 107
Diadumenidae, 107
Diafolliculina, 41
Diaixidae, 265
Diaixis, 265
Diala, 185
Dialidae, 185
Diamysis, 293
Diaphana, 198
Diaphanidae, 198
Diaphanoeca, 58
Diaphorodoris, 202
Diaphus, 364
Diaporthales, 15
Diarthrodesella, 277
Diarthrodes, 279
Diaschistorchis, 138
Diascorhynchidae, 131
Diascorhynchides, 131
Diascorhynchus, 131
Diastoporidae, 333
Diastylidae, 309
Diastylis, 309
Diastyloides, 309
Diastylopsis, 309
Diazona, 352
Dibothriocephalus, 146
Dibranchus, 366
Dicarpa, 355
Dicata, 204
Dicentrarchus, 369
Diceratias, 366
Diceratiidae, 366
Dichelaspis, 281
Dichelesthiidae, 260
Dichelesthium, 260
Dichelopandulus, 287
Dichelyne, 175
Dichosporangium, 27
Dichromadora, 164
Diclidophora, 143
Diclidophoropsis, 143
Diclybothrium, 143
Dicodonium, 113
Dicoeladropora, 135
Dicologlossa, 374
Dicopia, 352
Dicoryne, 117
Dicranodromia, 290
Dicranophoridae, 150
Dicranotaenia, 147
Dicaspedella, 58
Dicrateria, 76
Dicrogaster, 138
Dicrotrichura, 280
Dicrumenia, 143
Dictyemera, 147
Dictyoceratida, 100
Dictyocotyle, 143
Dictyonella, 90
Dictyonellidae, 90
Dictyopteris, 26
Dictyosiphon, 26
Dictyosiphonaceae, 26
Dictyosiphonales, 26
Dictyosphaeria, 23
Dictyosporium, 19
Dictyota, 26
Dictyotaceae, 26
Dictyotales, 26
Dicyema, 104
Dicyemenna, 104
Dicyemidae, 104
Didelta, 164
Didemnidae, 352
Didemnum, 352
Didiadema, 130
Didiscus, 90
Didogobius, 372
Didymella, 15
Didymocystis, 138
Didymodiclinus, 138
Didymosphaeriaceae, 15
Didymosporangium, 22
Didymosulcus, 138
Didymozoon, 138
Diesingium, 147
Diexanthema, 261
Diffugia, 56
Diffugiidae, 56
Diffugina, 55
Digenea, 33, 136
Digitaria, 209
Digitatispora, 18
Diglotta, 323
Dignathodontidae, 323
Dihemistephanus, 138
Dikoleps, 184
Dilepis, 147
Dilsea, 35
Dimisophria, 262
Dimophyes, 121
Dinematomonas, 59
Dinemoura, 261
Dinobothrium, 147
Dinodasys, 158
Dinoflagellata, 47

- Dinonemertes*, 156
Dinonemertidae, 156
Dinophilus, 217
Dinophysis, 48
Dinopontiidae, 260
Dinopontius, 260
Dinosoma, 138
Dinurus, 138
Diadon, 374
Diadontidae, 374
Diadontolaimus, 164
Diodora, 183
Diogenes, 289
Diogenichthys, 364
Diogenidae, 289
Dionchus, 143
Diopatra, 221
Diopisthoporidae, 126
Diopisthoporus, 126
Dioplosyllis, 229
Diorchis, 147
Diosaccidae, 272
Diosaccopsis, 272
Diosaccus, 272
Dipetalonema, 175
Diphasia, 116
Diphtherostomum, 138
Diphyes, 121
Diphyidae, 121
Diphyllobothrium, 147
Diphylogaster, 259
Diplaconotum, 224
Diplastrella, 88
Diplecogaster, 367
Diplectanotrema, 143
Diplectanum, 143
Dipleurosoma, 114
Dipleurosomatidae, 114
Diplobrachia, 235
Diplocirrus, 218
Diplocotyle, 147
Diplocrinus, 338
Diplodasys, 157
Diplodonta, 208
Diplodus, 370
Diploeca, 58
Diplogonoporus, 147
Diplolaimella, 164
Diplolaimelloides, 164
Diplonema, 60
Diplopelta, 49
Diplopeltis, 164
Diplopeltoides, 164
Diplopeltula, 164
Diplophos, 362
Diplophrys, 78
Diplophryxus, 322
Diplopleura, 154
Diplopoda, 323
Diplopisthe, 147
Diplopsalis, 49
Diplopsalopsis, 49
Diplopteraster, 341
Diplosolen, 333
Diplosoma, 353
Diplospinus, 373
Diplostomum, 138
Diplothamnion, 30
Diplotheca, 58
Diplozoon, 143
Dipolydora, 227
Diporula, 330
Diptera, 324
Dipterosiphonia, 33
Dipturus, 359
Dipurena, 113
Diretmichthys, 367
Diretmidae, 367
Diretmoides, 367
Diretmus, 367
Dirivultidae, 260
Discalcis, 186
Discammina, 64
Discamminidae, 63
Discanomalina, 69
Dischides, 212
Disco, 265
Discoanomalina, 69
Discobothrium, 147
Discocelides, 128
Discocelis, 79
Discocoenchoecia, 246
Discocotyle, 143
Discodermia, 87
Discodorididae, 202
Discodoris, 202
Discoidae, 265
Discomedusa, 111
Disconectes, 317
Disconema, 164
Disconorbis, 71
Discoonchoecia, 246
Discopsis, 190
Discorbidae, 71
Discorbinella, 71
Discorbinellidae, 71
Discorbinoides, 71
Discorbis, 71
Discorhabdella, 94
Discospirina, 68
Discosporangium, 28
Discotectonica, 196
Disparella, 317
Dispilio, 227
Disporella, 333
Disseta, 265
Dissodinium, 49
Distansescharella, 329
Distaplia, 353
Distichoptilum, 106
Distioculus, 276
Distomus, 355
Distorsio, 192
Ditlevsenella, 164
Ditrichybothrium, 147
Ditrichocorycaeus, 256
Ditropichthys, 368
Ditrupa, 226
Diurodriliidae, 217
Diurodrilus, 217
Divaricella, 207
Dizoniopsis, 186
Dochmiotrema, 134
Doconesthes, 103
Dodecaceria, 217
Dohrnilla, 31
Dolabrilera, 201
Dolabriteridae, 201
Dolichodasys, 157
Dolichoenterum, 138
Dolichoiulus, 323
Dolicholagus, 361
Dolicholaimus, 164
Dolichomacrostomidae, 127
Dolichomacrostomum, 127
Dolichomastix, 76
Dolichopteryx, 361
Dolichosomatum, 164
Doliolaimus, 164
Dolioletta, 355
Doliolidae, 355
Doliolina, 355
Dolioloides, 355
Doliolum, 355
Doliopharynx, 134
Dolopichthys, 366
Doloria, 245
Domecia, 292
Donacidae, 209
Donacilla, 209
Donax, 209
Dondersia, 181
Dondersiidae, 181
Dondice, 204
Donsia, 42
Donsiella, 279
Dopseucythere, 247
Dorhynchus, 291
Doridicola, 258
Dorididae, 202
Doridunculus, 202
Doriopsilla, 203
Dorippidae, 290
Doris, 202
Doroixys, 254
Doropygella, 254
Doropygopsis, 254
Doropygus, 254
Dorothia, 66
Dorsiceratus, 269
Dorvillea, 217
Dorvilleidae, 217
Dorylaimopsis, 164
Dorymenia, 181
Doryphallophora, 280
Doryphallophoridae, 280
Doryplex, 87
Doryporella, 328
Dosima, 281
Dosinia, 210
Dothideales, 15
Dotidae, 203
Doto, 203
Dotona, 88
Drachiella, 32
Dracognomus, 165
Draconema, 165
Draconettidae, 372
Draculiciteria, 157
Dragmastra, 86
Dragmatella, 98

<i>Dragmatyle</i> , 91	<i>Echinocephalus</i> , 175	<i>Einarella</i> , 133
<i>Drepanilla</i> , 130	<i>Echinochasmus</i> , 138	<i>Einarhelmins</i> , 129
Drepanogigantidae , 155	<i>Echinocletodes</i> , 270	<i>Eiona</i> , 17
<i>Drepanogigas</i> , 155	<i>Echinocorambe</i> , 202	<i>Eirene</i> , 114
<i>Drepanomenia</i> , 181	<i>Echinocotyle</i> , 147	Eirenidae , 114
Drepanomeniidae , 181	<i>Echinocucumis</i> , 349	<i>Eisothistos</i> , 316
Drepanophoridae , 155	<i>Echinocyamus</i> , 347	<i>Ekleptostylis</i> , 309
<i>Drepanophoriella</i> , 155	<i>Echinocythereis</i> , 251	<i>Ekmania</i> , 348
<i>Drepanophorus</i> , 155	<i>Echinoderes</i> , 159	<i>Ektонодиастилис</i> , 309
<i>Drepanorchis</i> , 283	<i>Echinoderidae</i> , 159	<i>Ektonos</i> , 186
<i>Drepanorhynchides</i> , 131	<i>Echinodermata</i> , 337	<i>Elachisina</i> , 190
Drilliidae , 194	<i>Echinogammarus</i> , 300	Elachisinidae , 190
<i>Drilliola</i> , 195	<i>Echinoidea</i> , 346	<i>Elachista</i> , 25
<i>Drilonereis</i> , 215	<i>Echinolaophonte</i> , 275	Elachistaceae , 25
<i>Dromia</i> , 290	<i>Echinomacrurus</i> , 365	<i>Elagatis</i> , 370
Dromiidae , 290	<i>Echinomermella</i> , 175	<i>Elaphognathia</i> , 321
<i>Drymonema</i> , 111	<i>Echinoparyphium</i> , 138	<i>Elasipodida</i> , 350
<i>Dryodora</i> , 122	<i>Echinophallus</i> , 147	<i>Elasmopus</i> , 304
Dryodoridae , 122	<i>Echinoplana</i> , 128	Electridae , 329
<i>Dryosphaera</i> , 17	<i>Echinopleura</i> , 317	<i>Electrona</i> , 364
<i>Dubininolepis</i> , 147	<i>Echinopsyllus</i> , 270	<i>Eledone</i> , 213
<i>Dudresnaya</i> , 35	Echinorhinidae , 359	<i>Eleginus</i> , 365
<i>Dulichia</i> , 306	<i>Echinorhinus</i> , 359	<i>Eleutheria</i> , 113
<i>Dulichiopsis</i> , 306	<i>Echinorhynchus</i> , 176	Eleutheriidae , 113
<i>Dumontia</i> , 35	<i>Echinosigra</i> , 347	<i>Eleutherolaimus</i> , 165
Dumontiaceae , 35	<i>Echinostephilla</i> , 138	<i>Eleutheromenia</i> , 181
<i>Duosnemertes</i> , 154	<i>Echinostoma</i> , 138	<i>Elicodasia</i> , 223
<i>Duplominona</i> , 135	<i>Echinostylinos</i> , 94	<i>Elina</i> , 78
<i>Duploperaclistus</i> , 135	<i>Echinotheristus</i> , 165	<i>Eliotia</i> , 204
Duplorbidae , 283	Echinothuridae , 346	<i>Ellipinion</i> , 350
<i>Duplorbis</i> , 283	Echinothuroidea , 346	Ellipsolagenidae , 71
<i>Dussumieria</i> , 361	<i>Echinozone</i> , 318	<i>Ellisella</i> , 105
<i>Dynamena</i> , 116	<i>Echinus</i> , 346	Ellisellidae , 105
<i>Dynamene</i> , 320	<i>Echiodon</i> , 366	<i>Ellisina</i> , 328
<i>Dynamenella</i> , 320	<i>Echiostoma</i> , 363	Ellobiidae , 205
<i>Dyopedos</i> , 306	<i>Echiura</i> , 178	<i>Elminiinae</i> , 282
<i>Dysidea</i> , 100	Echiuridae , 178	<i>Elminius</i> , 282
Dysideidae , 100	<i>Echiurus</i> , 178	<i>Elofsonella</i> , 249
<i>Dysomma</i> , 360	<i>Echthrogaleus</i> , 261	<i>Elofsonia</i> , 249
<i>Dyspanopeus</i> , 292	<i>Ecklonia</i> , 28	Elopiformes , 360
<i>Dysponetus</i> , 216	<i>Eclyssipe</i> , 215	<i>Elphidiella</i> , 72
<i>Dysponius</i> , 259	<i>Ecteinascidia</i> , 352	Elphidiidae , 72
<i>Dystrogus</i> , 259	<i>Ectenurus</i> , 138	<i>Elphidium</i> , 72
<i>Dytaster</i> , 338	<i>Ectinosoma</i> , 273	<i>Elpidia</i> , 350
<i>Earlandammina</i> , 65	Ectinosomatidae , 273	Elpidiidae , 350
<i>Eastonia</i> , 209	<i>Ectinosomella</i> , 273	<i>Elstia</i> , 138
<i>Eatonina</i> , 188	<i>Ectinosomoides</i> , 273	<i>Elthusa</i> , 320
<i>Ebala</i> , 198	Ectocarpaceae , 27	<i>Elysia</i> , 200
<i>Ebalia</i> , 290	Ectocarpales , 27	Elysiidae , 200
Ebalidae , 198	<i>Ectocarpus</i> , 27	<i>Elzalia</i> , 165
<i>Ebinania</i> , 369	<i>Ectocotyla</i> , 135	<i>Emarginula</i> , 183
<i>Ebnerella</i> , 101	<i>Ectopleura</i> , 113	<i>Embletonia</i> , 205
<i>Ebria</i> , 79	<i>Ectyodoryx</i> , 94	<i>Emericella</i> , 16
<i>Ecliffeoegyra</i> , 186	<i>Edgertonia</i> , 63	<i>Emericellopsis</i> , 16
<i>Echelus</i> , 361	<i>Edwardsia</i> , 107	<i>Emetha</i> , 320
<i>Echeneibothrium</i> , 147	Edwardsiidae , 107	<i>Emiliania</i> , 76
Echeneidae , 369	<i>Eggerella</i> , 63	Emmelichthyidae , 370
<i>Echeneidocoelium</i> , 138	Eggerellidae , 63	<i>Emplectonema</i> , 154
<i>Echeneis</i> , 369	<i>Eggereilloides</i> , 63	Emplectonematidae , 154
<i>Echinaster</i> , 341	<i>Egmundella</i> , 114	<i>Emprostropharynx</i> , 129
Echinasteridae , 341	<i>Eguchipsammia</i> , 110	<i>Empruthotrema</i> , 143
<i>Echineulima</i> , 187	<i>Ehlersia</i> , 229	<i>Enalcyonium</i> , 256
Echinidae , 346	<i>Ehlersiella</i> , 230	<i>Enallopssammia</i> , 110
<i>Echiniscoides</i> , 236	<i>Ehrenbergina</i> , 70	<i>Enantia</i> , 128
Echiniscoididae , 236	<i>Eilohedra</i> , 72	Enantiidae , 128
<i>Echinobothrium</i> , 147	<i>Einara</i> , 362	<i>Enapteris</i> , 224

- Encentrum*, 150
Encheliella, 165
Enchelycore, 360
Enchelyopus, 365
Enchytraeidae, 234
Enchytraeus, 234
Encotylabe, 143
Endarachne, 28
Endectyon, 91
Endeidiidae, 242
Endeis, 242
Endocheres, 258
Endodinium, 49
Endosphaeraceae, 22
Engina, 193
Engraulidae, 361
Engraulis, 361
Enhydrosoma, 271
Enhydrosomella, 271
Enigmocoma, 46
Enipo, 224
Enneagonum, 121
Ennucula, 205
Enodiotrema, 138
Enopla, 154
Enoplocoyle, 144
Enoploides, 165
Enoplolaimus, 165
Enoplometopodidae, 288
Enoplometopus, 288
Enoplateuthidae, 212
Enoplus, 165
Ensaya, 302
Ensis, 209
Entalina, 212
Entalinidae, 212
Entalophoroecia, 333
Entelurus, 368
Enterocola, 254
Enterocolidae, 254
Enterocolides, 254
Enterognathidae, 254
Enterognathus, 254
Enteromorpha, 24
Enteropneusta, 336
Enteropsidae, 254
Enteropsis, 254
Enterostomula, 129
Enteroxenos, 187
Entobdella, 144
Entobiidae, 256
Entobius, 256
Entocladia, 22
Entocolax, 187
Entoconcha, 187
Entodictyon, 22
Entomobryidae, 324
Entomolepidae, 260
Entomopsyllus, 260
Entonomenia, 181
Entoprocta, 179
Entosolenia, 71
Enypniastes, 350
Eocuma, 309
Eophrixus, 322
Eoschizopera, 272
Eostichopus, 349
Epacanthion, 165
Ephesiella, 227
Ephippion, 374
Ephydra, 324
Ephydriidae, 324
Ephyrina, 286
Epiactis, 106
Epicalymma, 257
Epicaridea, 322
Epicladia, 24
Epicotyle, 144
Epidiopatra, 221
Epifolliculina, 42
Epigonichthys, 357
Epigonidae, 369
Epigonus, 369
Epilepton, 208
Epimeria, 299
Epimeriidae, 299
Epimolgus, 256
Epinephelus, 369
Epipractis, 106
Epipenaeon, 322
Eiphriixus, 322
Episiphon, 212
Epistomia, 329
Epistomiidae, 329
Epistominella, 72
Epistominidae, 72
Epitomapta, 350
Epitonidae, 186
Epitonium, 186
Epizoanthidae, 108
Epizoanthus, 108
Eponides, 72
Eponididae, 72
Epsiloninema, 165
Eranno, 219
Erato, 191
Eratoidae, 191
Ercolania, 201
Erenna, 122
Eretmochelys, 375
Eretmophorus, 365
Ergalatax, 192
Ergasilidae, 256
Ergasilus, 256
Ergasticus, 291
Ergenstrema, 144
Erginus, 182
Ergyne, 322
Ericiolus, 76
Eriechthonius, 301
Erignathus, 376
Eriocheir, 292
Eriochona, 41
Eriopisa, 304
Eriopisella, 304
Eriphia, 292
Eriphiidae, 292
Ernodesmis, 23
Erosaria, 191
Erpocotyle, 144
Errina, 118
Ersilia, 187
Erugosquilla, 284
Ervilia, 209
Erylus, 86
Erythrocladia, 35
Erythrocles, 370
Erythrocystis, 33
Erythrodermis, 36
Erythroglossum, 32
Erythropeltidae, 35
Erythropeltidales, 35
Erythrops, 293
Erythropsodinium, 49
Erythrotrichia, 35
Escharella, 331
Escharina, 331
Escharoides, 331
Eschrichtiidae, 376
Eschrichtius, 376
Esola, 275
Espegrendia, 134
Esperiopsis, 96
Eteone, 222
Ethmolaimus, 165
Ethmorhynchus, 130
Ethusa, 290
Ethusina, 290
Etmopterus, 359
Eualus, 287
Euandania, 307
Euapta, 350
Euarche, 214
Euaugaptilus, 264
Euborlasia, 153
Eubostrichus, 165
Eubothrium, 147
Eubrachiella, 260
Eubranchidae, 204
Eubranchus, 204
Eucalanidae, 265
Eucalanus, 265
Eucanuella, 271
Euchaeta, 265
Euchaetidae, 265
Euchaetomera, 294
Euchaetomeropsis, 294
Eucheilota, 114
Eucheilotidae, 114
Euchelipluma, 97
Euchirella, 263
Euchirograpsus, 292
Euchlanis, 150
Euchone, 225
Euchromadora, 165
Euclymene, 220
Eucodonium, 113
Euconchoecia, 246
Eucopia, 293
Eucopiidae, 293
Eucotyle, 138
Eucranta, 224
Eucrate, 292
Eucratea, 329
Eucrateidae, 329
Eucyliophora, 177
Eucythere, 248
Eucytheridae, 248
Eucytherura, 247, 248
Eudactylina, 260
Eudactylinella, 260

- Eudactylinidae**, 260
Eudactylopus, 279
Eudaronia, 184
Eudendriidae, 117
Eudendrium, 117
Eudesme, 24
Eudistoma, 353
Eudolium, 191
Eudorella, 310
Eudorellopsis, 310
Eodoxia, 121
Eodoxoides, 121
Eufolliculina, 42
Eugerda, 317
Eugerdella, 317
Euginoma, 328
Euglena, 59
 Euglenoids, heterotrophic, 59
Euglenozoa, 59
Euglypha, 56
Euglyphida, 56
Euglyphinidae, 56
Euglyphinopsis, 57
Eugomontia, 23
Eugotoea, 113
Eugymnanthea, 114
Eugyra, 354
Eugyrioides, 354
Eukrohnia, 335
Eukrohniidae, 335
Eulalia, 223
Eulalia, 223
Eulepethidae, 217
Eulima, 186
Eulimella, 197
Eulimidae, 186
Eulitoma, 187
Eumecynostomum, 126
Eumetula, 186
Eumicrotremus, 369
Eumida, 223
Eumorpholaimus, 165
Eumunida, 290
Eunereis, 220
Eunice, 217
Eunicella, 105
Eunicicola, 256
Eunicicolidae, 256
Eunicidae, 217
Euniphysidae, 218
Eunoë, 224
Euonyx, 302
Eupanthalis, 214
Euparthenia, 197
Eupasiphae, 286
Eupèlete, 277
Euphausia, 284
Euphausiacea, 284
Euphausiidae, 284
Euphellia, 108
Euphilomedes, 246
Euphrosome, 218
Euphrasinidae, 218
Euphrasyne, 218
Euphsya, 113
Euphsidae, 113
Euphsora, 113
Eupistella, 230
Euplectella, 103
Euplectellidae, 103
Euplokamis, 122
Eupogodon, 32
Eupolymnia, 230
Eupolymnophilus, 258
Eupolyodontes, 214
Euprosthiostomum, 128
Eupyrgidae, 351
Eupyrillus, 351
Euraphia, 282
Euraphiinae, 282
Eurete, 102
Euretidae, 102
Eurhamphaea, 123
Eurhamphaeidae, 123
Euritmia, 227
Euroiales, 16
Euryalina, 343
Eurycketodes, 270
Eurycope, 317
Eurycopidae, 317
Eurycyde, 242
Eurydice, 319
Eurylepta, 128
Euryleptidae, 128
Eurynebria, 323
Eurnome, 291
Euryozius, 292
Eurypharyngidae, 361
Eurypharynx, 361
Euryphoridae, 260
Euryphorus, 260
Euryplacidae, 292
Eurypon, 91
Erysilenum, 256
Eurysquilla, 284
Eurysquillidae, 284
Eurystomina, 165
Eurystrotos, 333
Eurysyllis, 229
Euryte, 253
Eurytemora, 268
Eurythenes, 302
Eurythoe, 215
Eusarsiella, 247
Eusirella, 299
Eusiridae, 299
Eusirogenes, 299
Eusiroides, 299
Eusirus, 299
Euspira, 191
Eustomias, 363
Eustrongylides, 175
Eusyllis, 229
Eusynonchus, 165
Eusynstyela, 355
Eusyrophoxus, 305
Eutaeniophorus, 368
Eutardigrada, 236
Euterpiña, 274
Euterpinidae, 274
Eutetrahynchus, 147
Euthalenessa, 227
Euthelepus, 230
Euthriostoma, 193
Euthynus, 373
Euthyonidiella, 348
Eutima, 114
Eutonina, 114
Eutrepiella, 59
Euxinia, 129
Euzetacanthus, 176
Euzonus, 221
Evadne, 244
Evansula, 271
Evermannella, 364
Evermannellidae, 364
Evoplosoma, 340
Exechonella, 331
Exilicrusta, 33
Exoclavarctus, 236
Exocetidae, 367
Exocoetus, 367
Exogemmida, 41
Exogone, 229
Exopenna, 262
Exosphaeroma, 320
Expansophria, 262
Exspina, 314
Exumella, 267
Fabespora, 83
Fabienna, 118
Fabricia, 225
Fabriciola, 225
Facciorella, 361
Facelinidae, 204
Facelinopsis, 204
Faerlea, 126
Fageapseudes, 313
Fagesia, 107
Falcidens, 181
Fallax, 334
Falsocythere, 251
Famelica, 196
Farrania, 265
Farranula, 256
Farrea, 102
Farreidae, 102
Farrella, 332
Fasciola, 138
Fasciolaria, 192
Fasciolariidae, 192
Fasciospongia, 100
Fauchea, 38
Faucheaceae, 38
Faucheria, 320
Fauvelia, 229
Fauveliopsidae, 218
Fauveliopsis, 218
Favorinidae, 204
Favorinus, 204
Favulina, 71
Fecampia, 130
Fecampiidae, 130
Fedora, 326
Fehria, 195
Feldmannia, 27
Feldmannophycus, 36
Felia, 246
Fellicola, 175
Fellogistomum, 138

- Fenestrolaimus*, 165
Fenestrulina, 330
Feresa, 376
Ferosagitta, 335
Fibulariidae, 347
Ficopomatus, 226
Fieldingia, 103
Figularia, 329
Filellum, 115
Filexilia, 269
Filichonidae, 41
Filicrisia, 333
Filifera, 117
Filinia, 150
Filipjeva, 166
Filipjevia, 166
Filitanais, 314
Filodinium, 49
Filograna, 226
Filogramula, 226
Filoncholaimus, 166
 Filose testate amoebae *incertae sedis*, 57
Filospermoidea, 151
Fimbriaria, 147
Fimbriariooides, 147
Fimbriofolium, 36
Fimbriosthenelais, 227
Finella, 185
Finmarchinella, 249
Fiona, 204
Fionidae, 204
Firolidae, 192
Firoloida, 192
Fischerina, 68
Fischerinidae, 66
Fissidentalium, 211
Fissurella, 183
Fissurellidae, 183
Fissurina, 71
Fissurisepta, 183
Fistulicola, 147
Flabellia, 23
Flabelliidae, 110
Flabellifera, 319
Flabelligella, 214
Flabelligera, 218
Flabelligeridae, 218
Flabellina, 204
Flabellinidae, 204
Flabelliphilus, 257
Flabellula, 53
Flabellullidae, 53
Flabellum, 110
Flaccisagitta, 335
Fladenia, 278
Flagellophora, 128
Flagellostomias, 363
Flaminolepis, 147
Flectichona, 41
Flexophora, 144
Flintina, 67
Flintinoides, 67
Florarctus, 236
Floriceps, 147
Floridosentis, 176
Flosculariaceae, 150
Flustra, 329
Flustrellidra, 332
Flustrellidridae, 332
Flustridae, 329
Foersteria, 115
Folia, 356
Folliculinidae, 41
Folliculinopsis, 42
Fontainella, 310
Foraminifera, 62
Foraminiferida, 62
Forcepia, 94
Forcipulatida, 342
Forskalia, 122
Forskaliidae, 122
Forticulcita, 138
Foslia, 26
Fosiella, 33
Fossaridae, 185
Fossarus, 185
Fosshageniella, 262
Foviella, 136
Fowlerina, 200
Foxtonia, 264
Fragilidium, 49
Francapseudes, 313
Fratercula, 375
Fratia, 254
Fratiidae, 254
Freyastera, 342
Freyella, 342
Freyellidae, 342
Fridericia, 234
Frieleiidae, 334
Frigidoalvania, 189
Frillagalma, 122
Fritillaria, 356
Fritillariidae, 356
Frondicularia, 74
Fryeria, 203
Fucaceae, 28
Fucales, 27
Fucus, 28
Fulmarus, 375
Fultonia, 270
Fulvia, 209
Funchalia, 286
Fungi, 15
Fungiacyathidae, 110
Fungiacyathus, 110
Funiculina, 105
Funiculinidae, 105
Furcellaria, 36
Furcellariaceae, 36
Furnestinia, 144
Furzenkoina, 70
Furzenkoinidae, 73
Fusarium, 19
Fuscapex, 187
Fusceulima, 187
Fusinus, 192
Fusiturris, 195
Fusonema, 166
Fustiaria, 212
Fustiariidae, 212
Gadella, 365
Gadiculus, 365
Gadidae, 365
Gadiformes, 365
Gadila, 212
Gadiliidae, 212
Gadilinidae, 212
Gadomus, 365
Gadus, 365
Gaetanus, 263
Gaeumannomyces, 17
Gaevskajatrema, 138
Gafrarium, 210
Gaidius, 263
Gaidropsarurus, 365
Gairleanema, 166
Galactosomum, 138
Galathea, 290
Galatheammina, 78
Galatheascus, 283
Galatheidae, 290
Galaxaura, 37
Galaxauraceae, 37
Galeocerdo, 359
Galeodea, 191
Galeoides, 371
Galeomma, 208
Galeommataidae, 208
Galeorhinus, 359
Galeus, 359
Galiteuthis, 213
Gamaster, 354
Gammanema, 166
Gammaracanthidae, 299
Gammaracanthus, 299
Gammarella, 304
Gammarellidae, 299
Gammarellus, 300
Gammaridae, 300
Gammaridea, 296
Gammarinema, 166
Gammaropsis, 300
Gammarus, 300
Gammogobius, 372
Ganesa, 184
Gangliopus, 261
Ganonema, 37
Ganymedebdella, 234
Gargarius, 46
Gari, 209
Garveia, 117
Gasterascidia, 355
Gasteropteridae, 200
Gasteropteron, 200
Gasterosteidae, 368
Gasterosteiformes, 368
Gasterosteus, 368
Gastrana, 209
Gastrochaena, 210
Gastrochaenidae, 210
Gastroclonium, 38
Gastrocotyle, 144
Gastrodelphyidae, 256
Gastrodelphys, 256
Gastrolecithus, 147
Gastropharynx, 129
Gastropoda, 182
Gastropodidae, 150
Gastroptychus, 290

- Gastrosaccus*, 294
Gastrotricha, 156
Gastroturia, 349
Gattyana, 224
Gaudryina, 66
Gaussia, 266
Gaussicia, 246
Gavelinellidae, 73
Gavelinopsis, 71
Gavia, 375
Gaviidae, 375
Gebiacantha, 289
Geeopsis, 278
Geitodoris, 202
Gelanemertes, 156
Gelaninospora, 18
Gelasinospora, 18
Gelidiaceae, 35
Gelidiales, 35
Gelidiella, 35
Gelidiocolax, 37
Gelidiopsis, 38
Gelidium, 36
Gelliodes, 98
Gemmellipora, 330
Gempylidae, 373
Gempylitrema, 144
Gempylus, 373
Genetyllis, 223
Genitoconia, 181
Genitocotyle, 138
Gennadas, 286
Genocidaris, 346
Genolinea, 138
Genostoma, 129
Geodia, 86
Geodiidae, 86
Geophilidae, 323
Geophilomorpha, 323
Geophilus, 323
Gephyroberyx, 367
Gephyrocrinus, 337
Gephyrotes, 329
Gephyrothuria, 349
Gerardia, 109
Gerlachius, 166
Geryon, 292
Geryonia, 119
Geryonidae, 291
Geryoniidae, 119
Gesiella, 224
Gianius, 233
Giardella, 255
Gibberula, 194
Gibbula, 183
Gieysztoria, 130
Gigantactinidae, 367
Gigantactis, 367
Gigantobilharzia, 138
Gigantocypris, 245
Gigantoporidae, 329
Gigartina, 36
Gigartinaceae, 36
Gigartinales, 36
Gilquinia, 147
Ginesia, 254
Ginglymostoma, 358
Ginglymostomatidae, 358
Giraudia, 26
Giraudiaceae, 26
Gitana, 297
Gitanopsis, 297
Glabratella, 73
Glabratellidae, 73
Glaciarcula, 334
Glandiceps, 336
Glandulactis, 106
Glandulina, 73
Glandulinidae, 73
Glannapontius, 259
Glans, 208
Glareolina, 254
Glaucidae, 205
GlauCUS, 205
Gleba, 200
Glenodinium, 49
Gleodinium, 49
Gliocladium, 19
Gliomastix, 19
Globicephala, 376
Globigerina, 73
Globigerinella, 73
Globigerinidae, 73
Globigerininita, 73
Globigerinoides, 73
Globivenus, 210
Globobulimina, 70
Globocassidulina, 70
Globopilumnus, 292
Globorotalia, 73
Globorotaliidae, 73
Globotextularia, 65
Globoturborotalia, 73
Globulina, 74
Gloiocladia, 38
Gloigullmia, 62
Gloiosiphonia, 35
Gloiosiphoniaceae, 35
Glomericirrus, 138
Glomospira, 62
Glossanodon, 361
Glossidae, 210
Glossobalanus, 336
Glossobius, 320
Glossodoris, 202
Glossus, 210
Glyceria, 218
Glycerella, 218
Glyceridae, 218
Glycinde, 218
Glycymerididae, 206
Glycymeris, 206
Glyphanostonum, 215
Glyphocrangon, 288
Glyphocrangonidae, 288
Glyphohesione, 218, 223
Glyphus, 286
Glyptelasma, 281
Glyptocephalus, 374
Glyptoxanthus, 292
Gnathia, 321
Gnathiidae, 321
Gnathophausia, 293
Gnathophausiidae, 293
Gnathophis, 361
Gnathophyllidae, 288
Gnathophyllum, 288
Gnathorhynchidae, 131
Gnathorhynchus, 131
Gnathostenetroides, 318
Gnathostenetroididae, 317
Gnathostomaria, 151
Gnathostomariidae, 151
Gnathostomula, 151
Gnathostomulida, 151
Gnathostomulidae, 151
Gnesiocerotidae, 128
Gnomonia, 15
Gnorimorhynchus, 131
Gnosonesimidae, 127
Gnosonesimila, 127
Gobiesocidae, 367
Gobiesociformes, 367
Gobiidae, 372
Gobius, 372
Gobiusculus, 372
Goesia, 301
Goezia, 175
Gofasia, 189
Golfingia, 178
Golfingiidae, 178
Golvanacanthus, 176
Gomontia, 22
Gomphopodarion, 254
Gonactinia, 107
Gonactiniidae, 107
Gonapodasmius, 138
Gonatidae, 212
Gonatus, 212
Goneplacidae, 292
Goneplax, 292
Goniada, 218
Goniadella, 218
Goniadidae, 218
Goniaeolididae, 204
Goniaeolis, 204
Goniaster, 340
Goniasteridae, 339
Gonichthys, 364
Gonilia, 209
Gonimocolax, 32
Gonimophyllum, 32
Goniocoma, 46
Goniodelphys, 254
Goniodoma, 49
Goniodorididae, 202
Goniodoris, 202
Goniolithon, 33
Goniomonas, 59
Gonionchus, 166
Gonionemus, 118
Goniotricaceae, 37
Goniotrichales, 37
Goniotrichiopsis, 38
Gonocerca, 138
Gonocercella, 138
Gononema, 27
Gononemertes, 155
Gonophysema, 258
Gonostoma, 362

- Gonostomatidae**, 362
Gonostomula, 129
Gonothyraea, 119
Gontrania, 24
Gonyaulax, 49
Goodalia, 209
Gorbunovia, 225
Gorgoniapolynoe, 224
Gorgoniidae, 105
Gorgonocephalidae, 343
Gorgonocephalus, 343
Gorgorhynchus, 176
Gossea, 118
Gotocotyla, 144
Gouania, 367
Gouldia, 210
Gourretia, 289
Gracilaria, 37
Gracilariaeae, 37
Gracilariales, 37
Gracilariosis, 37
Graffilla, 130
Grafillidae, 130
Grallatoria, 31
Grammaria, 115
Grammatostomias, 363
Grammicolepididae, 368
Grammicolepis, 368
Grammonus, 366
Grampus, 376
Grania, 234
Granigra, 184
Grantessa, 102
Grantia, 102
Grantiidae, 101
Granulina, 194
Graphis, 197
Graphonema, 166
Grapsicepon, 322
Grapsidae, 292
Grapsus, 292
Grateloupia, 35
Greeffiella, 166
Gregariella, 206
Gregarinidra, 329
Gregorioiscala, 186
Greuetodinium, 49
Griffithsia, 31
Grigelis, 74
Grillotia, 147
Grimpoteuthis, 213
Groenhiella, 18
Gromia, 56
Gromida, 56
Gromiidae, 56
Gronella, 301, 302
Grubea, 144
Grubeosyllis, 229
Gruberella, 54
Grubeulepis, 217
Gruvelialepas, 281
Gryphaeidae, 207
Gryphus, 334
Gyrophrynchus, 147
Guancha, 101
Guassicia, 246
Guentherus, 363
Guernea, 299
Guitarra, 98
Guitarriidae, 98
Gullmariella, 134
Gulsonia, 31
Gunenotophorus, 254
Gurjanovella, 154
Guttulina, 74
Guynia, 110
Guyniidae, 110
Gwynia, 334
Gwynioidea, 334
Gyge, 322
Gymnammodytes, 372
Gymnangium, 114
Gymnelus, 371
Gymnoascaceae, 17
Gymnoascus, 17
Gymnobela, 196
Gymnocanthus, 369
Gymnodinium, 49
Gymnodorididae, 203
Gymnogongrus, 36
Gymnognos, 113
Gymnolaemata, 326
Gymnomenia, 181
Gymnomeniidae, 181
Gymnophallus, 138
Gymnophycus, 31
Gymnorhynchus, 147
Gymnothamnion, 31
Gymnothorax, 360
Gymnura, 360
Gymnuridae, 360
Gynaecotyla, 138
Gypsina, 69, 74
Gyptis, 218
Gyratricella, 132
Gyratrix, 132
Gyrocolete, 147
Gyrocoeloides, 147
Gyrodactyloides, 144
Gyrodactylus, 144
Gyrodinium, 50
Gyroidina, 69
Gyroidinoides, 73
Gyrophillum, 105
Gyroscala, 186
Habrocidaris, 346
Hadriania, 192
Hadromerida, 87
Hadrothoe, 261
Hadwenius, 138
Haeckelia, 122
Haeckeliidae, 122
Haedropleura, 195
Haematocleptes, 216
Haemescharia, 36
Haemeschariaceae, 36
Haemobaphes, 262
Haemocera, 262
Haemulidae, 370
Hagiosynodos, 329
Haifameira, 269
Halacarellus, 240
Halacaridae, 239
Halacarininae, 240
Halacaropsis, 240
Halacarus, 240
Halaimus, 166
Halammohydra, 112
Halammohydridae, 112
Halammovortex, 129
Halanonchus, 166
Halaphanolaimus, 166
Halarachnion, 36
Halargyreus, 365
Halampa, 107
Halampella, 107
Halampidae, 107
Halampoides, 107
Halampoididae, 107
Haluriidae, 107
Halechiniscidae, 236
Halechiniscus, 236
Haleciidae, 114
Halecium, 114
Halectinosoma, 273
Haliactiidae, 107
Haliactis, 107
Halicardia, 211
Halice, 305
Halichaetonotus, 156
Halichoanolaimus, 166
Halichoerus, 376
Halichondria, 90
Halichondrida, 89
Halichondriidae, 90
Halichrysis, 38
Haliclona, 98
Halicystus, 112
Halicnemia, 90
Halicoides, 305
Halicometes, 88
Halicortex, 90
Halicreas, 119
Halicreatidae, 119
Halicreion, 305
Halicyptus, 160
Halicyclops, 253
Halidrys, 28
Haliella, 187
Halielloides, 187
Haligena, 16
Halimeda, 21
Halinema, 166
Haliopasma, 316
Haliotidae, 183
Haliotis, 183
Haliotrema, 144
Haliphron, 213
Haliplanella, 107
Haliplectus, 166
Halipteridae, 105
Halipteris, 105
Halipilon, 33
Halirages, 299
Haliragooides, 299
Halisarca, 101
Halisarcida, 101
Halisarcidae, 101
Halissera, 119
Halistemma, 122
Halitholus, 118

- Halitiara*, 118
*Halitrephe*s, 119
Halla, 221
Hallangia, 126
Hallangidae, 126
Halobatrachus, 366
Halobiotus, 236
Haloceras, 191
Haloceratidae, 191
Halocercus, 175
Halochlorococcum, 22
Halocladius, 324
Haloclavidae, 107
Halocoryne, 113
Halocorynidae, 113
Halocynthia, 354
Halocypria, 246
Halocyprididae, 245
Halocypris, 246
Halognathus, 18
Halonectria, 17
Halonema, 24
Halonympha, 211
Halophila, 38
Halophytophilus, 274
Halopithys, 33
Haloplanella, 134
Halopsis, 115
Halopterididae, 115
Halopteris, 29, 115
Haloptilus, 264
Halorhiza, 25
Halosacciocolax, 38
Halosarpeia, 16
Halosauridae, 360
Halosauropsis, 360
Halosaurus, 360
Halosbaena, 293
Halosbaenidae, 293
Haloschizopera, 272
Halosia, 31
Halosiphon, 28
Halosphaera, 76
Halosphaeria, 16
Halosphaeriaceae, 16
Halosphaeriales, 16
Halosphaeriopsis, 16
Halothrix, 25
Halothria, 16
Halteria, 43
Halteriidae, 43
Halurus, 31
Halvorsenius, 138
Halydictyon, 32
Halymenia, 35
Halymeniaceae, 35
Halyphysema, 63
Hamacantha, 98
Hamacanthidae, 98
Hamatoscalpellum, 282
Hamigera, 95
Hamingia, 178
Haminoea, 199
Haminoeidae, 199
Hancockia, 203
Hancockiidae, 203
Hangethelia, 130
Hanleya, 182
Hanleyidae, 182
Hannonia, 242
Hansenella, 298
Hansenium, 319
Hansenomysis, 295
Hanzawaia, 71
Hapalomus, 166
*Hapalospongidi*on, 25
Hapalotrema, 138
Haplocope, 314
Haplodiscus, 126
Haplognathia, 151
Haplognathiidae, 151
Haplogonaria, 126
Haplomesus, 318
Haplomunnidae, 318
Haploniscidae, 318
Haploniscus, 318
Hapoops, 297
Haplopharyngida, 127
Haplopharyngidae, 127
Haplopharynx, 127
Haplophragmoides, 63
Haplophragmoididae, 63
Haplophryne, 367
Haplopoma, 329
Haplopomidae, 329
Haploporus, 138
Haploposthia, 126
Haploposthiidae, 126
Haplosclerida, 98
Haplosplanchnus, 138
Haplospora, 29
Haplostatus, 254
Haplostoma, 254
Haplostomidae, 254
Haplostomides, 254
Haplostylus, 294
Haplosyllis, 229
Haploota, 330
Haplotestis, 127
Haplovejdovskya, 130
Haptenchelys, 360
Haptophytes, 76
Haraldia, 32
Haraldiophyllum, 32
Hardametopa, 307
Harmeria, 329
Harmocoma, 46
Harmothoe, 224
Harpacticidae, 274
Harpacticoida, 268
Harpacticus, 274
Harpinia, 305
Harpiniopsis, 306
Harrietella, 275
Harrimania, 336
Harrimaniidae, 336
Harriotta, 360
Hartlaubella, 119
Hartmannella, 53
Hartmannellidae, 53
Hartogia, 134
Harveyella, 36
Hastigerella, 274
Hastigerina, 73
Hastigerinidae, 73
Hathrometra, 338
Hatschekia, 260
Hatschekiidae, 260
Hauchellia, 230
Hauerinidae, 66
Haustoriidae, 300
Haustorius, 300
Havelockia, 348
Haynesina, 72
Hazelia, 340
Hebefustis, 319
Hebella, 115
Hecatonema, 26
Hedgpethia, 242
Hedingia, 351
Hedylopsidae, 201
Hedylopsis, 201
Heleobia, 190
Helgicirrha, 114
Heliacus, 196
Helicocranchia, 213
Helicolenus, 368
Helicometra, 138
Heliconema, 175
Helicosalpa, 355
Heligmus, 175
Heliochona, 41
Heliochonidae, 41
Helicotyle, 144
Heliodomidae, 329
Heliometra, 338
Helluoherpia, 181
Helminthocladia, 37
Helminthora, 37
Helotium, 18
Hemectyon, 92
Hemiaclis, 187
Hemiarthrus, 322
Hemiasster, 347
Hemiassterella, 88
Hemiasterellidae, 88
Hemiasteridae, 347
Hemibdella, 234
Hemicervinia, 271
Hemichaetonotus, 157
Hemichordata, 336
Hemicletodes, 270
Hemicyclopina, 254
Hemicyclopora, 331
Hemicyclops, 255
Hemicythere, 249
Hemicytheridae, 248
Hemicytherura, 248
Hemidasys, 157
Hemidinium, 50
Hemigaleidae, 359
Hemigellius, 98
Hemigrapsus, 292
Hemilamprops, 310
Hemilaophonte, 275
Hemilepton, 208
Hemimesochra, 270
Hemimycale, 95
Hemimysis, 294
Hemipera, 139
Hemipodus, 218

- Hemiptera**, 324
Hemiramphidae, 367
Hemiramphus, 367
Hemirhabus, 265
Hemiselmis, 59
Hemisphaerammina, 64
Hemistasia, 60
Hemitanarctus, 236
Hemithirididae, 334
Hemithriss, 334
Hemiurus, 139
Henia, 323
Henlea, 234
Henneguya, 84
Henricia, 341
Henryhowella, 251
Hepatoxylon, 147
Hepomadus, 285
Heptranchias, 358
Herbstia, 291
Herbstina, 254
Herceus, 95
Herdmania, 50, 354
Herklotischthys, 361
Hermadionella, 224
Hermaea, 201
Hermaeidae, 200
Hermodice, 215
Hero, 204
Heroidae, 204
Herponema, 27
Herpotrichiellaceae, 15
Herpyllobiidae, 256
Herpyllobius, 256
Herrmannella, 256
Hersiliodes, 255
Hertwigia, 103
Herwigia, 362
Hesione, 218
Hesionidae, 218
Hesionides, 218
Hesiorhynchus, 223
Hesiospina, 218
Heteralepadidae, 281
Heteralepas, 281
Heteramalla, 267
Heteramoeba, 54
Heteranomia, 207
Heteranthessius, 256, 258
Heteraxine, 144
Heteraxinoides, 144
Heterectya, 92
Heterenchelyidae, 360
Heterocapsa, 50
Heterocarpus, 287
Heterochaeta, 234
Heterochona, 41
Heteroconger, 361
Heterocotyle, 144
Heterocrypta, 291
Heterocyatholaimus, 166
Heterocyprideis, 247
Heterocythereis, 249
Heterodinium, 50
Heterodorididae, 204
Heterodoris, 204
Heterodrilus, 233
Heteroerythrops, 294
Heterogenys, 286
Heterokontophyta, 24
Heterokrohnia, 335
Heterolaophonte, 275
Heterolepidoderma, 157
Heteromastus, 216
Heteromesus, 318
Heteromyysis, 294
Heteromysoides, 294
Heteronchocotyle, 144
Heteronema, 59
Heteronemertini, 153
Heteronybelinia, 147
Heteronymphon, 242
Heteropale, 216
Heteropanope, 292
Heterophyes, 139
Heteropia, 102
Heteropiidae, 102
Heteropolynoe, 224
Heteropriacanthus, 369
Heteropsyllus, 270
Heteroptilus, 264
Heteropyramis, 121
Heterorhabdidae, 265
Heterorhabdus, 265
Heteroschismoides, 212
Heterosiphonia, 32
Heterospio, 219
Heterostegina, 74
Heterostigma, 354
Heterostylites, 265
Heterotanais, 314
Heterotardigrada, 236
Heterostrophyes, 139
Heteroteuthis, 212
Heteroxenotrichula, 157
Heteroxya, 90
Hexabothrium, 144
Hexactinella, 103
Hexactinellida, 102
Hexactinosa, 102
Hexadactylus, 355
Hexadella, 101
Hexanchidae, 358
Hexanchiformes, 358
Hexanchus, 358
Hexapleomera, 313
Hexaplex, 192
Hexarthra, 151
Hexasterophora, 102
Hexelasma, 282
Hexelasmatinae, 282
Hexostoma, 144
Hiatella, 210
Hiatellidae, 210
Higginsia, 90
Hildenbrandia, 37
Hildenbrandiaceae, 37
Hildenbrandiales, 37
Hiltermannicythere, 251
Himanthalia, 28
Himanthaliaceae, 28
Himantolophidae, 366
Himantolophus, 366
Himantura, 360
Himasthla, 139
Hincksia, 27
Hincksina, 329
Hincksinoflustra, 329
Hincksipora, 329
Hincksiporidae, 329
Hippaliosina, 329
Hippaliosinidae, 329
Hippasteria, 340
Hippellozoon, 330
Hippocampus, 368
Hippocrepina, 63
Hippocrepinidae, 63
Hippoglossoides, 374
Hippoglossus, 374
Hippolyte, 287
Hippolytidae, 287
Hippomedon, 302
Hippomenella, 329
Hippomonavella, 326
Hipponicidae, 190
Hipponix, 190
Hipponoe, 215
Hippopleurifera, 331
Hippopodiidae, 121
Hippopodina, 329
Hippopodinidae, 329
Hippopodius, 121
Hippoporella, 330
Hippoporidra, 330
Hippoporidridae, 329
Hippoporina, 326
Hippospongia, 100
Hippothoa, 330
Hippothoidae, 330
Hirondella, 302
Hirondellea, 302
Hirschmannia, 249
Hirudinea, 234
Hirudinella, 139
Hirundichthys, 367
Hispanirhynchia, 334
Histampica, 343
Histiobranchus, 360
Histioneis, 50
Histioteuthidae, 212
Histioteuthis, 212
Histodermella, 94
Histro, 366
Histriobdella, 219
Histriobdellidae, 219
Hoeglundina, 72
Hoferellus, 84
Hoffmaenneria, 166
Hofsteniidae, 126
Hofsteniola, 126
Holcomycteronus, 366
Holmsella, 37
Holobomolochus, 255
Holocentridae, 368
Holocoma, 46
Holognathidae, 321
Holopodidae, 337
Holorchis, 139
Holostephanus, 139
Holothuria, 349
Holothuriidae, 349

- Holothuroidea**, 348
Holoxea, 86, 87
Holtbyrnia, 362
Homalometron, 139
Homalopoma, 185
Homalorhagida, 160
Homarus, 288
Homaxinella, 90
Homoeognathia, 257
Homoeonema, 119
Homoeoscelis, 261
Homogammina, 78
Homola, 290
Homolidae, 290
Homolodromiidae, 290
Homologenus, 290
Homophiura, 345
Homoscleromorpha, 85
Homosclerophorida, 85
Homotrema, 73
Homotrematidae, 73
Hopkinsina, 75
Hopkinsinella, 75
Hoplangia, 109
Hoplaster, 339
Hoploneurmertini, 154
Hoplopera, 134
Hoploplana, 128
Hoplostethus, 367
Hopperia, 166
Hormathia, 107
Hormathiidae, 107
Hormiphora, 122
Hormosina, 63
Hormosinella, 63
Hormosinidae, 63
Hornera, 333
Horneridae, 333
Horsiella, 272
Hospitella, 62
Houartiella, 192
Howella, 369
Hubrechtella, 152
Hubrechtia, 152
Hubrechtidae, 152
Huilkalineus, 154
Humaria, 18
Huntemannia, 275
Huntemanniidae, 275
Huso, 360
Hyala, 190
Hyale, 300
Hyalidae, 300
Hyalinea, 74
Hyalinoecia, 221
Hyalocylis, 200
Halogyra, 197
Halogyrina, 197
Halogyrinidae, 197
Halonema, 102
Halonematidae, 102
Hyalopecten, 207
Hylopomatus, 226
Hylopontius, 261
Hyaloscypheaceae, 17
Hyalosphenia, 56
Hyalospheniidae, 56
Hyas, 291
Hyastenus, 291
Hyattella, 100
Hybocodon, 113
Hydatina, 198
Hydatinidae, 198
Hydractinia, 117
Hydractiniidae, 117
Hydrallmania, 116
Hydranthea, 114
Hydrasterias, 342
Hydrichthyidae, 118
Hydrichthys, 118
Hydrobates, 375
Hydrobia, 190
Hydrobiidae, 190
Hydrocharitaceae, 38
Hydrocharitales, 38
Hydroclathrus, 28
Hydrodendron, 115
Hydroides, 226
Hydrolagus, 360
Hydrolithon, 34
Hydroniscus, 318
Hydroschendyla, 323
Hydrozoa, 112
Hygophum, 364
Hygrosoma, 346
Hymedesmia, 95
Hymedesmiidae, 95
Hymenancora, 93
Hymenaster, 341
Hymeniacidon, 91
Hymenocephalus, 365
Hymenocotta, 139
Hymenodora, 286
Hymenomonas, 76
Hymenopenaeus, 286
Hymeraphia, 92
Hymerhabdia, 90
Hyocrinidae, 337
Hypalocletodes, 270
Hypania, 215
Hypaniola, 215
Hyperammina, 63
Hyperandrotrema, 139
Hypererythrops, 294
Hyperia, 308
Hyperiidae, 308
Hyperiidea, 308
Hyperioides, 308
Hyperiopsidae, 300
Hyperiopsis, 300
Hyperoche, 308
Hyperoglyphe, 374
Hyperoodon, 376
Hyperoplus, 372
Hypertrema, 139
Hyphalaster, 339
Hypeurochilus, 372
Hypnea, 36
Hypneaceae, 36
Hypneocolax, 36
Hypocoma, 46
Hypocomagalma, 46
Hypocomatidium, 46
Hypocomella, 46
Hypocomidae, 46
Hypocomides, 46
Hypocomidium, 46
Hypocomina, 46
Hypocreaceae, 17
Hypocreales, 17
Hypoderæum, 139
Hypodontolaimus, 166
Hypogastrura, 324
Hypogastruridae, 324
Hypoglossum, 32
Hypomenia, 181
Hypophorella, 332
Hypophorellidae, 332
Hyporhamphus, 367
Hypotrichinidae, 129
Hypseodoris, 202
Hypsibiidae, 236
Hypsicomus, 225
Hypsiechinus, 346
Hyrtios, 100
Hyssura, 316
Hyssuridae, 316
Hysterolecitha, 139
Hysterosomorpha, 139
Hysterothylacium, 175
Ialysus, 272
Ianiropsis, 318
Ianthopsis, 316
Icasterias, 342
Icelus, 369
Ichnopus, 302
Ichnusella, 277
Ichthyaniakis, 175
Ichthyascaris, 175
Ichthyodium, 157
Ichthyobronema, 175
Ichthyococcus, 362
Ichthyocotylurus, 139
Ichthyofilaria, 175
Ichthyomenia, 181
Ichthyotomidae, 219
Ichthyotomus, 219
Idas, 207
Idiacanthus, 363
Idmidronea, 333
Idomene, 278
Idotea, 321
Idoteidae, 321
Idunella, 301
Idusa, 320
Idyanthe, 279
Idyella, 279
Idyellospis, 279
Ihlea, 355
Imysis, 294
Ijimaia, 363
Ileraustroe, 307
Ilia, 290
Illex, 212
Illiosentis, 176
Ilyarachna, 318
Ilyarachnididae, 318
Ilyophis, 360
Illyphagus, 218
Immergentia, 332
Immergentiidae, 332

- Imogine*, 128
Inachus, 291
Inaloa, 135
Inanidrilus, 233
Inermidrilus, 233
Inermonephytis, 220
Inflatella, 94
Ingolfiella, 308
Ingolfiellidae, 308
Ingolfiellidea, 308
Innocuonema, 166
Inocybe, 18
Inopinodon, 198
Insecta, 323
Intercrusia, 278
Interleptomesochra, 269
Intoshia, 104
Intramolgidae, 256
Intramolgus, 256
Invenusta, 135
Inversochona, 41
Inversochonidae, 41
Ioanella, 72
Iolaea, 197
Ione, 322
Ionema, 166
Iophon, 93
Iophonidae, 93
Iopsiliidae, 219
Iosactiidae, 108
Iosactis, 108
Iothia, 182
Iotroata, 94
Iotrochotidae, 94
Iphimedia, 300
Iphimediidae, 300
Iphinoe, 309
Iphinopsis, 194
Iphitime, 217
Iphitus, 186
Ipnopidae, 363
Iravadiidae, 190
Ircinia, 100
Irciniidae, 100
Irenium, 114
Irenosyrinx, 195
Iridogorgia, 104
Irpa, 350
Irus, 210
Irvinea, 38
Isaea, 300
Isaeidae, 300
Isarachnanthus, 108
Isaurus, 109
Ischnocalanus, 266
Ischnochiton, 182
Ischnochitonidae, 182
Ischnomesidae, 318
Ischnomesus, 318
Ischyroceridae, 301
Ischyrocerus, 301
Ischyromene, 320
Isias, 265
Isidascus, 283
Isidella, 105
Isidiidae, 105
Isandiella, 70
Isobactrus, 241
Isochaetides, 234
Isochona, 41
Isochonidae, 41
Isocirrus, 220
Isocletopsyllus, 271
Isocomides, 46
Isocrinida, 338
Isocrinidae, 338
Isodictya, 98
Isodictyidae, 98
Isoedwardsia, 107
Isognomon, 207
Isognomonidae, 207
Isolda, 215
Isonomia, 207
Isophelliidae, 108
Isopoda, 316
Isops, 87
Isorropodon, 210
Isotomidae, 324
Isozoanthus, 109
Isthmoplea, 26
Istiophoridae, 373
Istiophorus, 373
Isurus, 358
Itaipusa, 131
Itaipusina, 131
Itasiella, 135
Itonoa, 36
Itunella, 270
Iwilsoniella, 16
Ixa, 290
Ixonema, 166
Jaculella, 63
Jaculina, 330
Jadammina, 65
Jaera, 318
Jakoba, 79
Janczewskia, 33
Jania, 34
Janicella, 286
Janickina, 54
Janira, 318
Janiralata, 318
Janirella, 318
Janirellidae, 318
Janiridae, 318
Janita, 226
Janolus, 204
Janthina, 186
Janthinidae, 186
Janua, 228
Janulum, 92
Janusion, 235
Japetella, 213
Japonacteon, 198
Jaschnovia, 263
Jasmineira, 225
Jaspis, 87
Jassa, 301
Javania, 110
Jaxea, 289
Jeanella, 258
Jensenia, 129
Joeropsidae, 318
Joeropsis, 318
Johania, 199
Johannsonia, 234
Johnstonia, 220
Johnstonmawsonia, 175
Jonesia, 247
Jonesiella, 278
Jonssonia, 27
Jorunna, 203
Josephella, 226
Joubinella, 306
Jugaria, 228
Jujubinus, 183
Julida, 323
Julidae, 323
Jullienella, 64
Kadosactidae, 108
Kadosactis, 108
Kainomatomyisis, 294
Kaiparapelta, 183
Kali, 372
Kalliapseudidae, 313
Kalliapseudinae, 313
Kallymenia, 35
Kallymeniaceae, 35
Kaloplocamus, 202
Kalyptorhynchia, 130
Kangarina, 248
Kantiella, 118
Karkinorhynchidae, 131
Karkinorhynchides, 131
Karkinorhynchus, 131
Karreriella, 63
Katadesmia, 205
Katamysis, 294
Kataplana, 135
Katerythrops, 294
Kathablepharids, 76
Kathablepharis, 76
Kathlania, 175
Katianira, 318
Katianiridae, 318
Katius, 302
Katodinium, 50
Katsuwonus, 373
Kefersteinia, 218
Kejdonia, 197
Kellia, 208
Kellicottia, 150
Kelliella, 210
Kelliellidae, 210
Kelliidae, 208
Kelliopsis, 208
Kemphyra, 287
Kenmackenzia, 139
Kentrochona, 41
Kentrochonopsis, 41
Kentrodorididae, 203
Kentrogonida, 283
Kentrosiga, 58
Keratella, 150
Keratoisis, 105
Kerguelenia, 302
Kesun, 221
Kiitokzia, 79
Kinetoplastids, 60
Kinetoskias, 327
Kinorhyncha, 159

- Kinorhynchus*, 160
Kirchenpaueria, 115
Kirchenpaueridae, 115
Kirnasia, 286
Kirschsteinothelia, 16
Kishinouyeidae, 112
Klebshormidiidae, 23
Kleionychocampoides, 275
Kleionychocampus, 275
Kleosoma, 274
Kliopsyllus, 277
Klugea, 166
Knipowitschia, 372
Knipowitschiatrema, 139
Kochlorine, 281
Koehlermetra, 338
Koellikeria, 139
Koellikerina, 117
Koellikeroides, 139
Kofoidinium, 50
Kogia, 376
Kogiidae, 376
Kohlmeyeriella, 16
Koinocystididae, 131
Kolga, 350
Kolkwitziella, 50
Kophobelemnidae, 105
Kophobelemnnon, 105
Korethraster, 341
Korethrasteridae, 341
Kornmannia, 23
Kornmanniaceae, 23
Koroga, 302
Korotnevella, 53
Kotorella, 147
Kowalevskia, 356
Kowalewskiidae, 356
Krachia, 186
Krampella, 115
Krampia, 166
Kraspedonema, 166
Kraussinidae, 334
Krithe, 249
Krithidae, 249
Krohnia, 214
Krohnitta, 335
Krohnittella, 335
Krohnittellidae, 335
Krohnittidae, 335
Kronborgia, 130
Kroyeria, 260
Kroyeriidae, 260
Kryptoperidinium, 50
Kryptos, 193
Ktariella, 144
Kuckuckia, 27
Kudoa, 84
Kudoidae, 84
Kuetzingiella, 27
Kuhnria, 144
Kukenthalia, 355
Kuma, 126
Kupellonura, 316
Kvaleyia, 34
Kyliniella, 38
Kyphosidae, 371
Kyphosus, 371
- Labidiasteridae**, 343
Labidocera, 267
Labidognathia, 151
Labidoplax, 350
Labioleanira, 227
Laboea, 43
Labratrema, 139
Labridae, 371
Labrostratus, 216
Labrospira, 63
Labrus, 371
Labyrinthomyxa, 77
Labyrinthula, 77
Labyrinthulids, 77
Laccaria, 18
Lacernidae, 330
Lachlanella, 66
Lacistorhynchus, 147
Lacuna, 188
Lacunidae, 188
Lacunovermis, 139
Lacydonia, 219
Lacydoniidae, 219
Laemonema, 365
Laemophiurina, 344
Laeocochlis, 186
Laetmatophilus, 306
Laetmogone, 350
Laetmogonidae, 350
Laetmonice, 215
Laevicardium, 209
Laevidentalina, 74
Laevidentalium, 212
Laevipeneroplis, 69
Laeviphitus, 190
Laevipilina, 182
Lafoea, 115
Lafoeidae, 115
Lafoeina, 114
Lafystiidae, 301
Lafystius, 301
Lagena, 73
Lagenammina, 64
Lagenidae, 73
Lagenipora, 328
Lagenodelphis, 376
Lagenorhynchus, 132, 376
Lagenosolenia, 71
Lagisca, 224
Lagocephalus, 374
Lagotia, 42
Lagynidae, 62
Lagynis, 62
Laimella, 166
Laingiidae, 118
Laingomedusae, 118
Lajonkairia, 210
Laleonectes, 291
Lamarckina, 71
Lamellaria, 191
Lamellariidae, 191
Lamellibrachia, 235
Lamellibrachiidae, 235
Lamellisabella, 235
Lamellisabellidae, 235
Lamellodiscus, 144
Laminaria, 28
- Laminariaceae**, 28
Laminariales, 28
Laminariocolax, 27
Laminarionema, 27
Lamippe, 256
Lamippella, 256
Lamippidae, 256
Lamippina, 256
Lamippulina, 256
Lamna, 358
Lamnidae, 358
Lamniformes, 358
Lampadena, 364
Lampadioteuthis, 212
Lampanyctus, 364
Lampea, 122
Lampeidae, 122
Lampetra, 358
Lampridae, 364
Lampriformes, 364
Lampris, 364
Lampritrema, 139
Lamprogrammus, 366
Lampropidae, 309
Lamprops, 310
Lanassa, 230
Lanceola, 308
Lanceolidae, 308
Lanceoporidae, 330
Landrumius, 267
Lanice, 230
Lanuginella, 103
Laodicea, 115
Laodiceidae, 115
Laomedea, 119
Laomediidae, 289
Laona, 199
Laonice, 227
Laonome, 225
Laophonte, 275
Laophontella, 278
Laophontidae, 275
Laophontina, 275
Laophontodes, 270
Laophontopsidae, 276
Laophontopsis, 276
Laothoes, 299
Laphania, 230
Laphystiopsidae, 301
Laphystiopsis, 301
Lappanella, 371
Laridae, 375
Larnacicus, 328
Larsenia, 190
Larus, 375
Laryngosigma, 74
Lasaea, 208
Lasaeidae, 208
Lasiognathus, 367
Lasiosphaeriaceae, 18
Lasiotocus, 139
Lateriporus, 147
Laternula, 211
Laternulidae, 211
Latiaxis, 192
Laticarinina, 71
Latiremidiae, 276

- Latiromitra*, 194
Latirus, 193
Latreillia, 290
Latreilliidae, 290
Latreutes, 287
Latronema, 166
Latrunculia, 88
Latrunculiidae, 88
Laubieria, 276
Laubierina, 191
Laubierinidae, 191
Laura, 283
Lauratonema, 166
Lauratonemoides, 166
Laurencia, 33
Laurida, 283
Lauridae, 283
Lautitia, 15, 16
Lautosporopsis, 17
Laxosuberites, 88
Laxus, 166
Leachia, 213
Leander, 288
Leanea, 230
Leanira, 227
Learedius, 139
Leathesia, 25
Leathesiaceae, 25
Lebbeus, 287
Lebetus, 373
Leblondiella, 26
Lebouraia, 50
Lecane, 150
Lecanidae, 150
Lecithaster, 139
Lecithobotrys, 139
Lecithochirium, 139
Lecithocladium, 139
Lecithodesmus, 139
Lecithoepitheliata, 127
Lecithophyllum, 139
Lecithostaphylus, 139
Leodovitia, 166
Leegaardiella, 44
Leguta, 131
Lehardyia, 131
Leiocapitella, 216
Leiochrides, 216
Leiodermatium, 87
Leiognathidae, 370
Leiognathus, 370
Leiopathes, 109
Leiophloea, 15
Leitoscoloplos, 222
Lejolisia, 31
Lekanesphaera, 320
Lekanorhynchus, 132
Lembos, 298
Lenopharynx, 134
Lensia, 121
Lenticulina, 75
Lentidium, 210
Leocrates, 218
Leodora, 228
Leonnates, 220
Leontocaris, 287
Leotiales, 17
Lepadella, 150
Lepadidae, 281
Lepadogaster, 367
Lepadomorpha, 281
Lepas, 281
Lepechinella, 301
Lepechinellidae, 301
Lepechinelloides, 301
Lepeophtheirus, 259
Lepeta, 182
Lepetella, 183
Lepetellidae, 182
Lepetidae, 182
Lepidapedon, 139
Lepidasthenia, 224
Lepidauchen, 139
Lepidepecreella, 302
Lepidepecreum, 302
Lepidion, 365
Lepidisis, 105
Lepidochelys, 375
Lepidochitona, 182
Lepidocybium, 373
Lepidodasyidae, 157
Lepidodasys, 157
Lepidoderma, 157
Lepidodeuterammina, 65
Lepidomenia, 181
Lepidomeniidae, 181
Lepidonotus, 224
Lepidophanes, 364
Lepidophyllum, 139
Lepidopleurus, 182
Lepidopora, 118
Lepidopus, 373
Lepidorhombus, 374
Lepidoteuthidae, 212
Lepidoteuthis, 212
Lepidotrigla, 368
Lepocreadium, 139
Leposphilus, 257
Lepraliella, 330
Lepraliellidae, 330
Lepralioides, 329
Leptacanthichthys, 366
Leptagonus, 369
Leptamphopus, 299
Leptanthuria, 316
Leptastacidae, 276
Leptastacus, 276
Leptasterias, 342
Leptinogaster, 255
Leptobothrium, 144
Leptocaris, 272
Leptocharias, 359
Leptochariidae, 359
Leptocheirus, 298
Leptochela, 286
Leptochelia, 314
Leptocheliidae, 314
Leptochilichthyidae, 362
Leptochilichthys, 362
Leptochiton, 182
Leptochitonidae, 182
Leptocletodes, 270
Leptoclinides, 353
Leptoclinus, 371
Leptocotyle, 144
Leptocythere, 249
Leptocytheridae, 249
Leptoderma, 362
Leptodiscus, 50
Leptofauchea, 38
Leptognathia, 314
Leptognathiella, 314
Leptognathiinae, 314
Leptognathoides, 314
Leptognathiopsis, 314
Leptogorgia, 105
Leptohyalis, 64
Leptolaimoides, 166
Leptolaimus, 166
Leptomesochra, 269
Leptometra, 338
Leptomysis, 294
Leptomyxidae, 53
Lepton, 208
Leptonematella, 25
Leptonemella, 167
Leptonidae, 208
Leptonotis, 190
Leptopentacta, 349
Leptophoxus, 306
Leptophyllus, 50
Leptophytum, 34
Leptoplana, 128
Leptoplanidae, 128
Leptopontia, 276
Leptopontidae, 276
Leptopsammia, 110
Leptopsyllus, 277
Leptorhynchoides, 176
Leptosiphonia, 33
Leptosomatides, 167
Leptosomatina, 167
Leptosomatum, 167
Leptosomella, 167
Leptosphaeria, 15
Leptosphaeriaceae, 15
Leptostomias, 363
Leptostylis, 309
Leptosynapta, 350
Leptotachidia, 278
Leptotheca, 83
Leptychaster, 339
Lequerrea, 254
Lernaescus, 257
Lernaeenicus, 262
Lernaeocera, 262
Lernaeodiscidae, 283
Lernaeodiscus, 283
Lernaeolophus, 262
Lernaeopoda, 260
Lernaeopodidae, 260
Lernaeopodina, 260
Lernanthropidae, 261
Lernanthropus, 261
Lernentoma, 255
Lesquerella, 57
Lesquereusiidae, 56
Lestidiops, 363
Lesueururia, 123
Lesueurigobius, 373
Lethenteron, 358

- Leucaltidae**, 101
Leucaltis, 101
Leucascidae, 101
Leucetta, 101
Leucia, 224
Leucilla, 101
Leuckartiara, 118
Leuclathrina, 101
Leucobrotula, 366
Leucocephalonemertes, 153
Leucochloridium, 139
Leucocryptos, 76
Leucon, 310
Leuconia, 102
Leuconidae, 310
Leucophloeus, 91
Leucopsacidae, 103
Leucopsacus, 103
Leucoraja, 359
Leucosia, 290
Leucosiidae, 290
Leucosolenia, 102
Leucosolenida, 101
Leucosoleniidae, 102
Leucosyrinx, 195
Leucothea, 123
Leucothoe, 301
Leucothoidae, 301
Levantocaris, 289
Leviapseudes, 313
Leviapseudinae, 313
Levinella, 101
Levinellidae, 101
Levinsenia, 222
Levinseniella, 139
Liagora, 37
Liagoraceae, 37
Libinia, 291
Lichenopora, 333
Lichenoporidae, 333
Lichia, 370
Lichomolgella, 256
Lichomolgidae, 256
Lichomolgides, 256
Lichomolgidium, 256
Lichomolgus, 256
Liebmannia, 24
Ligninicola, 17
Ligophorus, 144
Ligula, 147
Ligur, 287
Liliopsida, 38
Liljeborgia, 301
Liljeborgiidae, 301
Lilyopsis, 121
Lima, 207
Limacia, 203
Limacina, 200
Limacinidae, 200
Limacospora, 17
Limameira, 269
Limanda, 374
Limapontia, 201
Limapontiidae, 201
Limaria, 207
Limatula, 207
Limea, 207
Limidae, 207
Limifossoridae, 181
Limipolycestis, 132
Limirhynchus, 132
Limnactinia, 108
Limnactiniidae, 108
Limnocletodes, 271
Limnodriloides, 233
Limnodriloidinae, 233
Limnodrilus, 234
Limnomysis, 294
Limnopolyptae, 118
Limnoria, 320
Limnoriiidae, 320
Limopsidae, 206
Limopsis, 206
Linarea, 256
Linatella, 192
Lindia, 150
Lindiidae, 150
Lindra, 17
Lineidae, 153
Lineolia, 167
Lineopsella, 153
Lineosoma, 274
Lineus, 153
Linga, 207
Linguadactyla, 144
Linguella, 204
Lingulina, 74
Lingulopyrulinoides, 74
Linhomoeus, 167
Linopherus, 215
Linophryne, 367
Linophrynidae, 367
Linotaeniidae, 323
Lintonium, 139
Liocarcinus, 291
Liocarenus, 198
Liocranchia, 213
Liocyma, 210
Liomesus, 193
Lioniella, 134
Liosotomia, 198
Liparis, 369
Lipkea, 112
Lipkeidae, 112
Lipobranchius, 226
Lipomelum, 275
Lipomera, 317, 319
Liponema, 108
Liponematidae, 108
Lipophrys, 372
Liriope, 119
Liripora, 333
Lironeca, 320
Liropus, 296
Lissa, 291
Lissoclinum, 353
Lissodendoryx, 94
Lissomphalia, 184
Lisspecten, 207
Lissotesta, 184
Listea, 133
Listriella, 301
Lithistida, 87
Lithobubaris, 87
Lithoderma, 25
Lithodermataceae, 27, 28
Lithodes, 289
Lithodidae, 289
Lithoglyptidae, 281
Lithognathus, 370
Litholepis, 34
Lithonida, 102
Lithophaga, 206
Lithophyllum, 34
Lithoporella, 34
Lithothamnion, 34
Litigiella, 208
Litinium, 167
Litiopa, 185
Litiopidae, 185
Litocorsa, 223
Litoscalpellum, 282
Litosiphon, 26
Littorina, 188
Littorinidae, 188
Litucivis, 133
Lituolidae, 64
Lituotuba, 64
Lituotubidae, 64
Livonectus, 320
Liza, 371
Lizzella, 117
Lizza, 117
Lobata, 123
Lobianchia, 364
Lobiancopora, 332
Lobiancoporidae, 332
Lobiger, 200
Lobochona, 41
Lobochonidae, 41
Lobochonina, 41
Lobodelphys, 254
Lobophora, 26
Lobotes, 370
Lobotidae, 370
Loculicytheretta, 247
Loeblichopsis, 63
Lohmannella, 241
Lohmannellinae, 241
Lohmanniella, 44
Loimia, 230
Loimosina, 144
Loliginidae, 212
Loligo, 212
Lomanotidae, 203
Lomanotus, 203
Lomasoma, 139
Lomentaria, 38
Lomentariaceae, 38
Lonchoplanella, 134
Lonchotaster, 339
Longibrachium, 221
Longicyatholaimus, 167
Longiductotrema, 139
Longigammarus, 300
Longipedia, 276
Longipediidae, 276
Longithorax, 294
Longosomatidae, 219
Lopadorrhynchidae, 219
Lopadorrhynchus, 219

- Lophaster*, 341
Lophelia, 109
Lophiidae, 366
Lophiiformes, 366
Lophiostomataceae, 16
Lophiotrema, 16
Lophius, 366
Lophocladia, 33
Lophodolos, 366
Lophogaster, 293
Lophogastridae, 293
Lophorodoris, 202
Lophosiphonia, 33
Lophotaspis, 136
Lophothrix, 267
Lophotidae, 364
Lophotus, 364
Lophoura, 262
Loricifera, 160
Loricocercia, 246
Loripes, 207
Lottiidae, 365
Lottidae, 182
Loureirophonte, 275
Lourinia, 276
Louriniidae, 276
Lovenella, 115
Lovenellidae, 115
Loveniidae, 347
Loweina, 364
Loxoaula, 249
Loxoconcha, 249
Loxoconchidae, 249
Loxoconchidea, 249
Loxomespilon, 179
Loxosoma, 179
Loxosomatidae, 179
Loxosomella, 179
Loxostomina, 70
Lubbockia, 257
Lubbockiidae, 257
Lucernaria, 112
Lucernariidae, 112
Lucernariopsis, 112
Lucicutia, 265
Lucutiidae, 265
Lucifer, 286
Luciferidae, 286
Lucinella, 207
Lucinidae, 207
Lucinoma, 208
Luetkenia, 260
Luffisphaera, 79
Lugia, 223
Luidia, 339
Luidiidae, 339
Lulworthia, 17
Lumbricalus, 219
Lumbricillus, 234
Lumbriclymene, 220
Lumbrineridae, 219
Lumbrinerides, 219
Lumbrineropsis, 219
Lumbrineris, 219
Lumpenus, 371
Lunulogammarus, 300
Luria, 191
Luriculus, 130
Lusitanops, 196
Lutheriella, 134
Lutra, 376
Lutraria, 209
Luttrellia, 17
Luvaridae, 373
Luvarus, 373
Lycenchelys, 371
Lycodes, 371
Lycodonus, 371
Lycoteuthidae, 212
Lygdamis, 225
Lynchagalma, 122
Lyonsia, 211
Lyonsiella, 211
Lyonsiidae, 211
Lyrodus, 210
Lysianassa, 302
Lysianassidae, 301
Lysianella, 303
Lysibranchia, 218
Lysidice, 218
Lysilla, 230
Lysiosquilla, 284
Lysiosquillidae, 284
Lysiosquilloides, 284
Lysippe, 215
Lysippides, 215
Lysmata, 287
Lyssacinosa, 103
Lytocarpia, 114
Maasella, 105
Macandrevia, 334
Macandreviidae, 334
Macandrewia, 87
Macarorchestia, 307
Maccabeus, 160
Macellicephala, 224
Macellomenia, 181
Macellomeniidae, 181
Machaerus, 292
Machilidae, 323
Macoma, 209
Macrinella, 314
Macrochaeta, 214
Macrochiron, 257
Macrochironidae, 257
Macroclymene, 220
Macroconchoecia, 246
Macrocyprina, 251
Macrocyprididae, 251
Macrocypridina, 245
Macrocypridoidea, 251
Macrocyprina, 251
Macrocypris, 251
Macrocypridae, 157
Macrodasycidae, 157
Macrodasys, 157
Macromckenziea, 251
Macromphalus, 190
Macroparalepis, 363
Macrophoma, 19
Macrophthalmus, 292
Macropipus, 291
Macropodia, 291
Macroramphosus, 368
Macrorhynchia, 114
Macrosetella, 276
Macrostomida, 127
Macrostomidae, 127
Macrostomum, 127
Macrostylidae, 318
Macrostylis, 318
Macrouridae, 365
Macrouridophora, 144
Macrourus, 365
Macrurocyttidae, 368
Macruronus, 366
Mactra, 209
Mactridae, 209
Macvicaria, 139
Madracis, 110
Madrella, 204
Madrellidae, 204
Madrepore, 110
Maehrenthalia, 133
Maeotias, 118
Maera, 304
Maerella, 304
Maeropsis, 304
Magelona, 219
Magelonidae, 219
Magnaporthaceae, 17
Magnibursatus, 140
Magnisudis, 364
Magnoliophyta, 38
Maja, 291
Majidae, 290
Makaira, 373
Makrokylinurus, 309
Malacanthuria, 316
Malacobdella, 156
Malacobdellidae, 156
Malacocephalus, 365
Malacoceros, 227
Malacopsyllus, 269
Malacoraja, 360
Malacosaccus, 103
Malacosteus, 363
Malagazziidae, 115
Maldane, 220
Maldanella, 220
Maldanidae, 219
Malleidae, 207
Malletidae, 205
Malleus, 207
Mallotus, 362
Malmgreniella, 224
Mamiella, 76
Mamillocylichna, 200
Mammiphitime, 217
Manania, 112
Manayunkia, 225
Mangelia, 195
Manningia, 284
Manta, 360
Mantoniella, 76
Manunema, 167
Manzonia, 189
Marenzelleria, 227
Maresearsia, 121
Margareta, 330
Margarettidae, 330

- Margarites*, 184
Margelopsidae, 113
Margelopsis, 113
Marginaster, 340
Marginella, 194
Marginellidae, 194
Marginulina, 74
Marginulinopsis, 74
Magonema, 167
Margrethia, 362
Marguetta, 327
Marinella, 134
Marinobogidiella, 298
Marinosphaera, 18
Marinospora, 17
Marionia, 204
Marionina, 234
Mariplanella, 134
Marirhynchus, 131
Marisolaris, 17
Maritrema, 140
Marphysa, 218
Marrus, 122
Marseniella, 191
Marsenina, 191
Marsepulina, 62
Marshallora, 185
Marsipella, 63
Marsteinia, 276
Marsupenaeus, 286
Martensia, 303
Marthasterias, 342
Martinottiella, 63
Marycarmenia, 217
Maryllynnia, 167
Massarina, 16
Massilina, 67
Mastigiidae, 112
Mastigoteuthidae, 213
Mastigoteuthis, 213
Mastobranchus, 216
Mastocarpus, 36
Masturus, 374
Mathilda, 196
Mathildidae, 196
Matutidae, 290
Mauligobius, 373
Maulisia, 362
Maupasia, 219
Maurolicus, 362
Mawsonascaris, 175
Maxmuelleria, 178
Mayorella, 54
Mazocraeoides, 144
Mazocraes, 144
Meara, 128
Mecynocera, 266
Mecynoceridae, 266
Mecynoecia, 333
Mecynoeciidae, 333
Mecynostomidae, 126
Mecynostomum, 126
Mediaster, 340
Mediomastus, 216
Medorippe, 290
Megabalaninae, 283
Megabalanus, 283
Megacalanidae, 266
Megacalanus, 266
Megaclausia, 255
Megadasys, 157
Megadenus, 187
Megadesmolaimus, 167
Megalocercus, 356
Megalocotyle, 145
Megalocotyloides, 145
Megalocranchia, 213
Megalomma, 225
Megalomphalus, 190
Megalomycteridae, 368
Megalophallus, 140
Megalopidae, 360
Megalops, 360
Megalopidae, 304
Megalopus, 304
Megamorion, 127
Megamphopus, 301
Megamerilla, 221
Meganyctiphanes, 284
Megapontiidae, 261
Megapora, 328
Megaptera, 376
Megascolecidae, 234
Megastygarctides, 236
Megathiris, 334
Megathyrididae, 334
Megatrema, 282
Megatrematiniae, 282
Megaxinus, 207
Megavoldia, 206
Megerli, 334
Megeurystomina, 167
Meiocheta, 127
Meiodiscus, 38
Meiogymnophallus, 140
Meiopsyllus, 277
Meiorhopalon, 112
Melaenis, 224
Melamphaes, 367
Melamphaidae, 367
Melanella, 187
Melanitta, 375
Melanocetidae, 366
Melanocetus, 366
Melanochlamys, 199
Melanogastrales, 18
Melanogrammus, 365
Melanolagus, 361
Melanonidae, 365
Melanonus, 365
Melanostigma, 371
Melanostomias, 363
Melanotaenium, 18
Melarhaphe, 188
Melibe, 204
Melicertidae, 115
Melicertissa, 115
Melicertum, 115
Melicertus, 286
Melinna, 215
Melinnacheres, 258
Melinnampharete, 215
Melinopsis, 215
Melita, 304
Melitidae, 304
Mellonymptha, 103
Melobesia, 34
Melodichthys, 366
Melonanchora, 93
Melonchela, 93
Melonis, 69, 74
Melophysa, 122
Meloriastacus, 276
Meloscaphander, 200
Melphidippa, 305
Melphidippella, 305
Melphidippidae, 304
Melythasides, 215
Membranipora, 330
Membraniporella, 329
Membraniporidae, 330
Membranoptera, 32
Menestho, 198
Menigrates, 303
Menigratopsis, 303
Meningodora, 287
Mentodus, 362
Mercenaria, 210
Meredithia, 35
Merga, 118
Merhippolyte, 287
Meristotheca, 37
Merizocotyle, 145
Merlangius, 365
Merlia, 98
Merlucciidae, 366
Merlucciotrema, 140
Merluccius, 366
Merocryptus, 290
Meromenia, 181
Meromonakantha, 315
Merona, 117
Mertensia, 122
Mertensiidae, 122
Mesacanthion, 167
Mesacanthoides, 167
Mesacmaea, 107
Mesaikeras, 266
Mesaikokeratidae, 266
Mesalia, 185
Meseres, 349
Mesnilia, 255
Mesocalanus, 264
Mesochaetopterus, 216
Mesocheres, 259
Mesochordaeus, 356
Mesochra, 270
Mesocletodes, 270
Mesodasys, 157
Mesodesmatidae, 209
Mesoglicola, 257
Mesoglicolidae, 257
Mesogloia, 25
Mesognatharia, 151
Mesognathariidae, 151
Mesogobius, 373
Mesoikopleura, 356
Mesoixys, 254
Mesolamprops, 310
Mesometopa, 307
Mesometra, 140

- Mesonerilla*, 221
Mesophyllum, 34
Mesopleustes, 306
Mesoplodon, 376
Mesopodopsis, 294
Mesoporus, 50
Mesopsyllus, 270
Mesorchis, 140
Mesorhabdus, 265
Mesorynchus, 132
Mesosagattia, 335
Mesostygarctus, 236
Mesotanais, 314
Mesothuria, 349
Mesozoa, 104
Messevriella, 56
Messoplana, 134
Metacalanus, 263
Metachromadora, 167
Metacirolana, 320
Metaconchoecia, 246
Metacrangon, 288
Metacyatholaimus, 167
Metacyclopina, 254
Metacylicolaimus, 167
Metacyphocaris, 303
Metadasynemella, 167
Metadena, 140
Metadesmodora, 167
Metadesmolaimus, 167
Metafolliculina, 42
Metagathotanais, 314
Metahuntemannia, 275
Metalaimus, 168
Metalcyonidium, 332
Metalinhomoeus, 168
Metallogorgia, 104
Metambasia, 303
Metamblyops, 294
Metamicrocotyla, 145
Metamphiascopsis, 272
Metanema, 59
Metanematobothrium, 140
Metaonchoecia, 246
Metaparoncholaimus, 168
Metapenaeopsis, 286
Metapenaeus, 286
Metapeyssonnelia, 35
Metaphoxus, 306
Metapolycope, 246
Metapontius, 259
Metapseudidae, 313
Metapseudinae, 313
Metaraeolaimoides, 168
Metarhombognathus, 241
Metastrombidium, 43
Metasychis, 220
Metavermilia, 226
Metaverruca, 282
Metaxia, 185
Metechinorhynchus, 176
Meteoria, 366
Meteororometra, 338
Metepsilonema, 168
Meterythrops, 294
Metidae, 276
Metis, 276
- Metoncholaimus*, 168
Metopa, 307
Metopella, 307
Metopion, 79
Metorchis, 140
Metridia, 266
Metridiidae, 108
Metridinidae, 266
Metridium, 108
Metromonas, 79
Metroperiella, 326
Metschnikowia, 18, 100
Metschnikowiaceae, 18
Metzgeria, 194
Mexidrilus, 233
Meylia, 168
Micatuba, 62
Michelopagurus, 289
Michithyops, 294
Micippa, 291
Micoletzkyia, 168
Micracanthodinium, 50
Micralymma, 323
Micramphora, 56
Micranthodinium, 50
Micrella, 153
Micrenophrys, 369
Microarthridion, 278
Microascaceae, 17
Microascales, 17
Microascus, 17
Microbothrium, 145
Microcalanus, 265
Microcassiope, 292
Microcerberidae, 321
Microcerberidea, 321
Microcerberus, 321
Microcharon, 319
Microchirus, 374
Microchlamyiidae, 55
Microchlamys, 55
Microcionidae, 92
Microcionina, 91
Microcladia, 31
Microclymene, 220
Microcoryne, 25
Microcosmus, 354
Microcotyle, 145
Microcyema, 104
Microcythere, 250
Microcytheridae, 250
Microcytherura, 248
Microdajidae, 280
Microdqus, 280
Microdalyellia, 130
Microdeutopus, 298
Microdictyon, 22
Microdisseta, 265
Microdrillia, 195
Microgelidiopsis, 31
Microgloma, 206
Microhedyle, 201
Microhedyliidae, 201
Microhydrula, 118
Microhydrulidae, 118
Microichthys, 369
Microjaera, 318
- Microjanira*, 318
Microjassa, 301
Microlaimus, 168
Microlophilichthys, 366
Microloxoconcha, 249
Micromaldane, 220
Micromelo, 198
Micromenia, 181
Micromesistius, 365
Micrometula, 62
Micromonas, 76
Micronectriella, 16
Micronephthys, 220
Micronereis, 220
Micronerilla, 221
Microparasellidae, 319
Microphallus, 140
Microphthalmus, 218
Micropilina, 182
Micropisa, 291
Micropleurotoma, 195
Micropontiidae, 261
Micropontius, 261
Micropora, 330
Microporella, 330
Microporellidae, 330
Microporidae, 330
Microposthia, 125
Microprotopus, 301
Micropsammella, 56
Micropsammis, 278
Micropythia, 300
Microscaphidium, 140
Microsetella, 274
Microsomacanthus, 147
Microspio, 227
Microspongium, 25
Microspora, 23
Microsporaceae, 23
Microsporales, 23
Microsporangium, 25
Microstenothee, 307
Microstoma, 361
Microstomatidae, 361
Microstomidae, 128
Microstomum, 128
Microstomus, 374
Microstylifer, 88
Microthelia, 16
Microthriaceae, 16
Microthyrium, 16
Microvoluta, 194
Microxestoleberis, 251
Micrura, 153
Micrurides, 153
Micrurinella, 153
Mikro, 184
Mikroconchoecia, 246
Mikrosyphar, 27
Miliammina, 64
Miliolidae, 66
Miliolina, 66
Miliolinella, 67
Millericrinida, 337
Millerigobius, 373
Mimocalanus, 268
Mimonectes, 308

- Mimonectidae**, 308
Mimosella, 332
Mimosellidae, 332
Minchinellidae, 102
Minervella, 276
Miniacina, 73
Minicheviella, 201
Minicheviellidae, 201
Minictena, 122
Minipera, 354
Ministeria, 79
Minolaimus, 168
Minolia, 184
Minona, 135
Minuspio, 227
Minutanemertes, 155
Minyichthys, 368
Mirabilicoxa, 317
Miracia, 276
Miraciidae, 276
Mirandotanais, 315
Mirapinna, 368
Mirapinnidae, 368
Mirocaris, 286
Mirognathus, 362
Miroslavia, 268
Miscellania, 229
Misophria, 262
Misophriella, 262
Misophriidae, 262
Misophrioida, 262
Misophriopsis, 262
Mississippiinidae, 73
Mitosporic fungi, 18
Mitra, 194
Mitrella, 194
Mitridae, 194
Mitrocoma, 115
Mitrocomella, 115
Mitrocomidae, 115
Mitrocomium, 116
Mitrolumna, 195
Mitsukurina, 358
Mitsukurinidae, 358
Mixolineidae, 154
Mixolineus, 154
Mixtocalanus, 267
Mnemiidae, 123
Mnemiopsis, 123
Mobula, 360
Modeeria, 116
Modiolarca, 206
Modiolicola, 256
Modiolula, 207
Modiolus, 206
Moelleria, 185
Moelleriopsis, 184
Moerisia, 113
Moerisiidae, 113
Moevenbergia, 133
Mohnia, 193
Mola, 374
Molgolaimus, 168
Molgula, 354
Molgulidae, 354
Molguloides, 354
Molicola, 147
Molidae, 374
Mollia, 330
Mollicia, 246
Mollisia, 17
Mollusca, 181
Molpadia, 351
Molpadiida, 351
Molpadiidae, 351
Molva, 365
Momedossa, 317
Monacanthidae, 374
Monachocrinus, 338
Monachus, 376
Monacilla, 268
Monactis, 107
Monamphiura, 344
Monasaceae, 16
Monascus, 16, 140
Monhystera, 168
Monobiceros, 128
Monoblastiaceae, 16
Monobrachium, 118
Monobryozoon, 332
Monobryozontidae, 332
Monocelididae, 135
Monocelis, 135
Monocelopsis, 135
Monoceratina, 247
Monochirus, 374
Monocletodes, 275
Monocorycaeus, 256
Monocoryne, 112
Monocosta, 58
Monocotyle, 145
Monocrepidium, 90
Monoculodes, 305
Monoculopsis, 305
Monodaeus, 292
Monodanthura, 316
Monodella, 293
Monodellidae, 293
Monodhelmis, 140
Monodictys, 19
Monodon, 376
Monodontidae, 376
Monogenea, 142
Monognathidae, 361
Monognathus, 361
Monogononta, 150
Monolistra, 321
Monomitopus, 366
Monomyces, 110
Monoophorum, 129
Monophorus, 185
Monoplacophora, 182
Monoporella, 330
Monoposthia, 168
Monopylephorus, 233
Monorchis, 140
Monorygma, 147
Monosiga, 58
Monosporus, 31
Monostaechas, 115
Monostichoplana, 135
Monostilifera, 154
Monostroma, 23
Monostromataceae, 23
Monotheca, 116
Monoplana, 135
Monoplanidae, 135
Monogyma, 197
Monstrilla, 262
Monstrillidae, 262
Monstrilloida, 262
Monstrillopsis, 262
Montacuta, 208
Montacutidae, 208
Monterosatus, 209
Monticellina, 217
Mopalidae, 182
Mora, 365
Moridae, 365
Morlaixia, 168
Mormonilla, 262
Mormonillidae, 262
Mormonilloida, 262
Moroniidae, 369
Mortierella, 15
Mortierellaceae, 15
Morulaeplecta, 65
Mothocyia, 320
Muceddina, 254
Mucor, 15
Mucoraceae, 15
Mucorales, 15
Mucronalia, 187
Mucrosenia, 278
Muellerina, 249
Mugga, 215
Muggiae, 121
Mugil, 371
Mugilidae, 371
Mullidae, 371
Mullus, 371
Multicalyx, 136
Multipeniata, 129
Multivalvulida, 84
Munella, 316, 318
Munida, 290
Munidopsis, 290
Munna, 319
Munneurycope, 317
Munnidae, 319
Munnopsidae, 319
Munnopsis, 319
Munnopsurus, 317
Muraena, 360
Muraenesocidae, 361
Muraenesox, 361
Muraenidae, 360
Murchisonella, 197
Murex, 192
Murexiella, 192
Muriceides, 105
Muricidae, 192
Muricopsis, 192
Musculista, 207
Musculus, 206
Musellifer, 157
Mustelidae, 376
Mustelus, 359
Mya, 210
Mycale, 96
Mycalidae, 96

- Mycalina**, 96
Myceliospongia, 101
Mychophilus, 254
Mycophycophila, 17
Mycosphaerella, 16
Mycospbaerellaceae, 16
Mycteronastes, 145
Mycteroperca, 369
Myctophidae, 364
Myctophiformes, 364
Myctophum, 364
Myicolidae, 257
Myidae, 210
Myliobatidae, 360
Myliobatis, 360
Myodocopida, 245
Myoforceps, 206
Myonera, 211
Myosotella, 205
Myoxocephalus, 369
Myozona, 127
Myozonaria, 127
Myra, 290
Myriactula, 25
Myrianida, 229
Myriapora, 330
Myriaporidae, 330
Myriochele, 222
Myriocladia, 25
Myriogloea, 25
Myriogramme, 32
Myrionema, 25, 117
Myrionemataceae, 25
Myriotrichia, 26
Myriotrichiaceae, 26
Myriotrichidae, 351
Myriotrochus, 351
Myrmekiderma, 90
Myrtea, 207
Mysella, 208
Mysia, 210
Mysidacea, 293
Mysidae, 293
Mysideis, 294
Mysidella, 294
Mysidetes, 294
Mysidion, 261
Mysidobdella, 235
Mysidopsis, 294
Mysis, 294
Mystacocarida, 252
Mystides, 223
Mystrocentrus, 315
Mytilaster, 206
Mytilicola, 257
Mytilicolidae, 257
Mytilidae, 206
Mytilina, 150
Mytilus, 206
Myxaster, 342
Myxasteridae, 342
Myxicola, 225
Myxidiidae, 82
Myxidium, 82
Myxilla, 93
Myxiliidae, 93
Myxillina, 93
Myxine, 358
Myxinidae, 358
Myxiniformes, 358
Myxobilatus, 84
Myxobolidae, 84
Myxobolus, 84
Myxodium, 50
Myxomolgus, 258
Myxoproteus, 83
Myxosporea, 82
Myxotheca, 62
Myxotrichaceae, 17
Myxotrichum, 17
Myxozoa, 82
Myzomolgus, 255
Myzopontius, 259
Myzostoma, 220
Myzostomidae, 220
Myzotheridion, 258
Naccaria, 38
Naccariaceae, 37
Nacellidae, 182
Nadina, 127
Nadinidae, 127
Naiades, 214
Naididae, 233
Naineris, 222
Nais, 17, 233
Namalycastis, 220
Namanereis, 220
Nanalicidae, 160
Nanalaricus, 160
Nanaspididae, 261
Nanaspis, 261
Nanichthys, 367
Nannastacidae, 310
Nannastacus, 310
Nannobrachium, 364
Nannocalanus, 264
Nannocythere, 249
Nannoeca, 58
Nannolaimoides, 168
Nannolaimus, 168
Nannomesochra, 270
Nannoniscidae, 319
Nannoniscoides, 319
Nannonicus, 319
Nannonyx, 303
Nannopodella, 271
Nannopus, 275
Nannorhynchides, 130
Nannosquillidae, 284
Nannosquilloides, 284
Nanobalcis, 187
Nanocassiope, 292
Nanomia, 122
Nansenia, 361
Naobranchia, 260
Napoliplana, 135
Narcetes, 362
Narcissia, 340
Narcomedusae, 118
Narella, 105
Narrimania, 186
Nasicola, 145
Nassariidae, 193
Nassarius, 193
Natalolana, 320
Nathaniella, 271
Natica, 191
Naticidae, 191
Natonemertes, 156
Naucrates, 370
Naufragella, 17
Nausithoe, 111
Nausitboidae, 111
Nautosphaeria, 17
Nealotus, 373
Neanthes, 220
Neanuridae, 324
Neasellus, 319
Necora, 291
Nectantis, 108
Nectadamas, 121
Nectiopoda, 243
Nectochaeta, 224
Nectonema, 160
Nectonematida, 160
Nectonematidae, 160
Nectonemertes, 156
Nectonemertidae, 156
Nectopyramis, 121
Nectria, 17
Nectriella, 17
Neevea, 38
Negaprion, 359
Neilonella, 206
Neilonellidae, 206
Nemacystus, 25
Nemaliales, 37
Nemalion, 37
Nemanema, 168
Nemasomatidae, 323
Nemastoma, 36
Nemastomataceae, 36
Nematobothrium, 140
Nematobrachion, 284
Nematocarcinidae, 287
Nematocarcinus, 287
Nematocarpus, 114
Nematoda, free-living, 161
Nematoda, parasitic, 174
Nematodinium, 50
Nematomenia, 181
Nematomorpha, 160
Nematoneis, 218
Nematopagurus, 289
Nematophorus, 116
Nematoplana, 135
Nematoplanidae, 135
Nematorhynchidae, 132
Nematorhynchus, 132
Nematoscelis, 284
Nematoscella, 107
Nematothenais, 315
Nemella, 169
Nemertellina, 155
Nemertellopsis, 155
Nemertesia, 116
Nemertini, 152
Nemertoderma, 128
Nemertodermatida, 128
Nemertodermatidae, 128
Nemertopsis, 154

- Nemertoscolex*, 153
Nemesis, 260
Nemichthyidae, 361
Nemichthys, 361
Nemoderma, 28
Nemogullmia, 62
Nemopsis, 117
Neoamphitrite, 230
Neoarctus, 236
Neoargestes, 270
Neobisiidae, 237
Neobisium, 237
Neobrachiella, 260
Neobradya, 276
Neobradyidae, 276
Neocalanus, 264
Neocentrophyidae, 160
Neochromadora, 169
Neocnus, 348
Neocomatella, 338
Neoconorbina, 75
Neocrania, 334
Neocucumis, 348
Neocyclopina, 254
Neocyclops, 253
Neocytherideidae, 250
Neocytherideis, 250
Neocytthus, 368
Neodasyidae, 157
Neodasys, 157
Neodexiopira, 228
Neodiplostomum, 140
Neoechinophora, 278
Neoechinorhynchus, 176
Neoeponides, 71, 72
Neoergasilus, 256
Neogammarus, 300
Neogloboquadrina, 73
Neognathophausia, 293
Neognathorhynchus, 131
Neogobius, 373
Neogoniolithon, 34
Neohela, 298
Neohexostoma, 145
Neohyssura, 316
Neolagenipora, 331
Neolagisca, 224
Neolampadidae, 347
Neolampas, 347
Neoleanira, 227
Neolenticulina, 75
Neolepidapedon, 140
Neolepocreadium, 140
Neolepton, 208
Neoleptonidae, 208
Neolithodes, 289
Neomecynostomum, 126
Neomenia, 181
Neomeniidae, 181
Neomicrorbis, 228
Neomorphaster, 343
Neomorphasteridae, 343
Neomysis, 294
Neonchocotyle, 145
Neonesidea, 247
Neonesidea?, 247
Neonesthes, 363
Neonotoporus, 140
Neopalene, 242
Neoparamoeba, 54
Neoparvicapsula, 84
Neopentadactyla, 348
Neophasis, 140
Neoplilinidae, 182
Neoplanocera, 128
Neopleurotomoides, 196
Neopleustes, 306
Neopolycystis, 132
Neopolynoe, 224
Neopontius, 259
Neopseudocapitella, 216
Neopycnodonte, 207
Neoraja, 360
Neorhabdus, 265
Neorhadinorhynchus, 177
Neorossia, 212
Neorotalia, 74
Neosabellides, 215
Neoscalpellum, 282
Neoschizorhynchus, 132
Neoscolecidithrix, 268
Neoscopelidae, 364
Neoscopelus, 364
Neosimnia, 191
Neosiphonia, 31
Neosteganoderma, 140
Neostygarcetus, 236
Neotanaidae, 313
Neotanaidomorpha, 313
Neotanais, 313
Neotima, 114
Neotonchus, 169
Neoturris, 118
Neovermilia, 226
Nephasma, 178
Nephromonorchia, 140
Nephropidae, 288
Nephrops, 288
Nephropsis, 288
Nephroelmis, 76
Nephtyidae, 220
Nephtys, 220
Nepinnotheres, 292
Nepotilla, 196
Neptunea, 193
Nereia, 29
Nereicola, 257
Nereicolidae, 257
Nereididae, 220
Nereilinum, 235
Nereimyra, 219
Nereiospora, 17
Nereiphylla, 223
Nereis, 220
Nerilla, 221
Nerillidae, 221
Nerillidium, 221
Nerillidopsis, 221
Nerita, 183
Neritidae, 183
Nerocila, 320
Nerophis, 368
Nesiarchus, 373
Nessipus, 261
Nessorhamphus, 361
Nethea, 86
Nettastoma, 361
Nettastomatidae, 361
Neurocaulon, 36
Neverita, 191
Newmaniverra, 282
Nezumia, 365
Nia, 18
Niaceae, 18
Nicella, 105
Nicippe, 305
Nicolea, 230
Nicolla, 140
Nicomache, 220
Nicorhiza, 261
Nicothoe, 261
Nicotboidae, 261
Nidificaria, 228
Nigerrhynchus, 131
Nigmatullinus, 287
Nijhoffia, 169
Ninoe, 219
Niobia, 118
Niobiidae, 118
Nipergasilus, 256
Niphatidae, 98
Nipponnemertes, 154
Niso, 187
Nitokra, 269
Nitophyllum, 32
Nitschkiaceae, 18
Nitzschia, 145
Noanelia, 215
Noctiluca, 50
Nodellum, 62
Nodilittorina, 188
Nodipecten, 207
Nodobacularia, 68
Nodophthalmidium, 68
Nodosaria, 74
Nodosariidae, 73
Nodulus, 190
Noemiamea, 198
Noerrevangia, 197
Noetiidae, 206
Nolandella, 54
Nolella, 332
Nolleliidae, 332
Nameidae, 374
Nomeus, 374
Nonion, 74
Nonionella, 74
Nonionellina, 74
Nonionidae, 74
Nonionoides, 74
Noodtiella, 274
Noodtorthopsyllus, 271
Nordgaardia, 327
Normanella, 276
Normanelliidae, 276
Normanicypthere, 249
Normanion, 303
Normichthys, 362
Notacanthidae, 360
Notacanthiformes, 360

- Notacanthus*, 360
Notarchus, 201
Notaulax, 225
Nothobomolochus, 255
Notholca, 150
Nothria, 221
Nothrya, 221
Notobranchaea, 200
Notobranchaeidae, 200
Notocaryoplana, 136
Notocaryoturbella, 136
Notocelis, 127
Notochaetosoma, 169
Notocirridae, 128
Notocirrus, 216
Notocytulus, 140
Notodelphyidae, 254
Notodelphys, 254
Notolimea, 207
Notolychnus, 364
Notomastus, 216
Notommatidae, 150
Notomonophorum, 134
Notomyotida, 339
Notophyllum, 223
Notoplana, 128
Notoplanidae, 128
Notoplites, 328
Notoproctus, 216, 220
Notopterophorus, 255
Notopus, 290
Notopygus, 215
Notoscopelus, 364
Notoselenus, 59
Notospermus, 154
Notostomobdella, 235
Notostomus, 287
Notosudidae, 363
Nototanidae, 315
Nototeredo, 211
Nototriphora, 185
Nouria, 64
Nouriidae, 64
Novafabricia, 225
Novistrombidium, 43
Novocriniidae, 277
Novodinia, 342
Nubecularia, 68
Nubeculariella, 63
Nubeculariidae, 68
Nubeculina, 68
Nubiella, 117
Nucella, 192
Nucellicola, 257
Nucellicolidae, 257
Nucula, 205
Nuculana, 205
Nuculanidae, 205
Nuculidae, 205
Nudora, 169
Nummulitidae, 74
Nuttallides, 69
Nybelinella, 366
Nybelinia, 147
Nyctiphantes, 284
Nymphaster, 340
Nymphon, 242
- Nymphonella*, 242
Nymphonidae, 242
Obelia, 119
Obesotoma, 195
Obesula, 185
Obla, 370
Oblea, 50
Obtortionidae, 185
Obtusella, 189
Obtusoecia, 246
Occultocythereis, 251
Oceanapia, 98
Oceania, 117
Oceanida, 187
Oceanites, 375
Oceanitidae, 375
Oceanobdella, 235
Oceanodroma, 375
Ocenebra, 192
Ochetostoma, 178
Ochlochaete, 24
Ocinebrina, 192
Ocnus, 348
Ocostaspora, 17
Octacnemidae, 352
Octacnemus, 352
Octavella, 85
Octineon, 108
Octineonidae, 108
Octobranchus, 230, 231
Octocorallia, 104
Octodactylus, 145
Octogonade, 116
Octolabea, 145
Octolasmis, 281
Octonchus, 169
Octophialucium, 115
Octopicala, 257
Octopicalidae, 257
Octopodidae, 213
Octopoteuthidae, 212
Octopoteuthis, 212
Octopus, 213
Octotriara, 118
Oculina, 110
Oculinidae, 110
Oculosetella, 276
Ocyopode, 292
Ocyopodidae, 292
Ocyropsidae, 123
Ocyropsis, 123
Ocythoe, 213
Ocythoidae, 213
Odessa, 113
Odhneriella, 140
Odhnerium, 140
Odiidae, 305
Odiliacletodes, 270
Odius, 305
Odobenidae, 376
Odobenus, 376
Odondebuenia, 373
Odontaspidae, 358
Odontaspis, 358
Odontaster, 339
Odontasteridae, 339
Odonthalia, 33
- Odontomacrurus*, 365
Odontophora, 169
Odontophoroides, 169
Odontorhynchus, 131
Odontostomops, 364
Odontosyllis, 229
Odontozona, 286
Odostomella, 197
Odostomia, 197
Oecidiobranchus, 317
Oedalechilus, 371
Oedicerina, 305
Oediceroides, 305
Oediceropsis, 305
Oediceros, 305
Oedicerotidae, 305
Oenone, 221
Oenonidae, 221
Oenopota, 195
Oerstedia, 155
Oesophagocystis, 140
Ogocephalidae, 366
Ogdeniella, 56
Ogmogaster, 140
Ogyrides, 287
Ogyrididae, 287
Oidiodendron, 19
Oikopleura, 356
Oikopleuridae, 356
Oithona, 255
Oithonidae, 255
Okenia, 202
Olavius, 233
Olencira, 320
Oligobrachia, 235
Oligobrachiidae, 235
Oligochaeta, 233
Oligocladus, 128
Oligofilomorpha, 127
Oligognathus, 216
Oligotrichaea, aloricate, 43
Olinrias, 118
Olinriasidae, 118
Olivella, 194
Olividae, 194
Olssonium, 140
Omalogyra, 196
Omalogyridae, 196
Omalosecosa, 328
Omalostomum, 128
Omicronema, 169
Ommastrephes, 213
Ommastrephidae, 212
Ommatokoita, 261
Omosudidae, 364
Omosudis, 364
Omphalophyllum, 26
Oncaeaa, 257
Oncaeidae, 257
Onceroxenidae, 280
Onceroxenus, 280
Onchidella, 205
Onchidiidae, 205
Onchidiopsis, 191
Onchidorididae, 202
Onchidoris, 202
Onchium, 169

- Onchnesia*, 178
Onchobothrium, 148
Onchocalanus, 266
Onchocotyle, 145
Onchodia, 186
Oncholaimellus, 169
Oncholaimus, 169
Onchophora, 175
Oncopagurus, 289
Oncorhynchus, 362
Oncousoecia, 333
Oncousoeciidae, 333
Ondina, 198
Ondiniella, 17
Oneirodes, 366
Oneirodidae, 366
Oneirophanta, 350
Onesimoides, 303
Oniscidium, 277
Onisimus, 303
Onoba, 189
Onubactis, 106
Onuphidae, 221
Onuphis, 221
Onychiuridae, 324
Onychiurus, 324
Onychocamptus, 275
Onychocella, 330
Onychocellidae, 330
Onychocorycaeus, 256
Onychognathiidae, 151
Onychopoda, 244
Onychoteuthidae, 212
Onychoteuthis, 212
Onygenaceae, 17
Onygenales, 17
Onykia, 212
Onyx, 169
Oocorys, 191
Oolina, 71
Ooneides, 255
Oopsacas, 103
Opalia, 186
Opaliopsis, 186
Opeatogenys, 367
Opechona, 140
Opecoelina, 140
Opecoeloides, 140
Opercularella, 114
Ophelia, 221
Ophelicola, 258
Opheliidae, 221
Ophelina, 221
Ophiacantha, 344
Ophiacanthidae, 344
Ophiactidae, 343
Ophiactis, 343
Ophiambix, 344
Ophiarachnella, 346
Ophiaster, 76
Ophichthidae, 360
Ophichthus, 361
Ophicotyle, 145
Ophidiaster, 340
Ophidiasteridae, 340
Ophidiidae, 366
Ophidiiformes, 366
Ophidion, 366
Ophidocladus, 33
Ophiernus, 345
Ophieulima, 187
Ophinella, 115
Ophioblennius, 372
Ophiobolus, 16
Ophiocamax, 344
Ophiociton, 345
Ophiocnemis, 344
Ophiocoenis, 346
Ophiocrates, 345
Ophiocreas, 343
Ophioceten, 345
Ophiocentrella, 345
Ophiodelphys, 255
Ophioderma, 346
Ophiidermatidae, 346
Ophiodromus, 219
Ophiogerion, 343
Ophioika, 253
Ophiolamia, 187
Ophiolebes, 344
Ophiolepis, 345
Ophioleucidae, 345
Ophiolimna, 344
Ophiomedea, 344
Ophiomisidium, 345
Ophiomitra, 344
Ophiomitrella, 344
Ophiomusium, 345
Ophiomyces, 344
Ophiomyxa, 343
Ophiomyxidae, 343
Ophionereididae, 345
Ophionereis, 346
Ophiopholis, 343
Ophiophrixus, 343
Ophiophycis, 345
Ophiopleura, 345
Ophioplinthaca, 344
Ophioprium, 344
Ophiopsila, 344
Ophiospus, 343
Ophiopyren, 345
Ophioscolex, 343
Ophioseide, 255
Ophiosphalma, 345
Ophiothamnus, 344
Ophiothrix, 343
Ophiotalsa, 345
Ophiotoma, 344
Ophiotrema, 344
Ophiotrichidae, 343
Ophiotuba, 62
Ophiotypa, 345
Ophiozonella, 345
Ophiraphidites, 90
Ophisurus, 361
Ophiora, 345
Ophiurida, 343
Ophiuridae, 345
Ophiuroidea, 343
Ophiurolepis, 345
Ophiliataspongia, 93
Ophryocotyle, 148
Ophryotroca, 217
Ophryotrocha, 217
Ophthalmidiidae, 68
Opisa, 303
Opisthadena, 140
Opisthocystis, 132
Opisthodonta, 229
Opisthoproctidae, 361
Opisthoproctus, 361
Opisthorchis, 140
Opisthosyllis, 229
Opisthoteuthidae, 213
Opisthoteuthis, 213
Opistomum, 134
Oplophoridae, 286
Oplophorus, 287
Ophalmina, 68
Oradarea, 299
Orania, 192
Orbiculorhynchus, 131
Orbimyces, 19
Orbinia, 222
Orbiniidae, 222
Orbitestella, 197
Orbitstellidae, 197
Orbulina, 73
Orcadia, 17
Orchestia, 307
Orchidasma, 140
Orchistoma, 116
Orchistomella, 115
Orchistomidae, 116
Orchomene, 303
Orchomenella, 303
Orcinus, 376
Orcynopsis, 373
Orectolobiformes, 358
Oreosomatidae, 368
Oridorsalis, 72
Oriopsis, 225
Ornatocalpellum, 282
Ornithobilharzia, 140
Ornithocercus, 51
Oromiina, 254
Orophocotyle, 140
Orseis, 219
Orthagorisicola, 260
Orthoconchoecia, 246
Ortholinea, 83
Ortholineidae, 83
Orthometra, 338
Orthomorphina, 70
Orthonectida, 104
Orthopiana, 136
Orthopsyllidae, 277
Orthopsyllus, 277
Orthopyxis, 119
Orthoplanchus, 140
Orygmatobothrium, 148
Orzeliscus, 236
Osangularia, 69
Osangulariella, 69
Oscarella, 85
Oschmarinella, 140
Oscilla, 197
Osilinus, 183
Osmeridae, 362

- Osmeriformes**, 361
Osmerus, 362
Osmundaria, 33
Osmunda, 33
Osteichthyes, 360
Osteopelta, 183
Osteopeltidae, 183
Ostraciidae, 374
Ostracoblabe, 19
Ostracoda, 245
Ostrea, 207
Ostreidae, 207
Ostreobium, 23
Ostreola, 207
Ostreopsis, 51
Ostromovia, 113
Otideaceae, 18
Otina, 205
Otinidae, 205
Otobothrium, 148
Otocelididae, 127
Otocelis, 127
Otodistomum, 140
Otohydra, 112
Otohydridae, 112
Otomesostoma, 135
Otomesostomidae, 135
Otoplana, 136
Otoplarella, 136
Otoplanidae, 135
Otoplanidia, 136
Otopsis, 223
Ototyphlonemertes, 154
Ototyphlonemertidae, 154
Ourozeuktes, 320
Ovatella, 205
Ovulidae, 191
Ovulina, 56
Owenia, 222
Oweniidae, 222
Oxycordyla, 89
Oxygyrus, 192
Oxynaspidae, 281
Oxynaspis, 281
Oxynoe, 200
Oxynoidea, 200
Oxynotus, 359
Oxyonchus, 169
Oxyphysis, 51
Oxypolella, 153
Oxypolia, 153
Oxyrrhis, 51
Oxystomina, 169
Oxytonostoma, 235
Oxytoxum, 51
Oxyurichthys, 373
Ozobranchidae, 235
Ozobranchus, 235
Pachastrella, 86
Pachastrellidae, 86
Pachos, 258
Pachycara, 371
Pachycerianthus, 108
Pachychalina, 98
Pachychelium, 303
Pachycordyle, 117
Pachycreadium, 140
Pachydinium, 51
Pachygrapsus, 292
Pachylasma, 282
Pachylasmatidae, 282
Pachylasmatoidea, 282
Pachymatisma, 87
Pachypsolus, 140
Pachyptilus, 264
Pachypygus, 255
Pachysoeca, 58
Pachystomias, 363
Pachytrema, 140
Padina, 26
Padinopsis, 26
Paecilomyces, 19
Paedampharete, 215
Paedomecynostomum, 126
Paelopatides, 349
Paeon, 262
Pagellus, 370
Pagophila, 375
Pagrus, 370
Paguridae, 289
Paguristes, 289
Pagurus, 289
Palaemon, 288
Palaemonella, 288
Palaemonetes, 288
Palaemonidae, 288
Palaeonemertini, 152
Palaeophalacroma, 51
Palaeotropus, 348
Palaephalacroma, 51
Palazzia, 184
Paleanotus, 216
Palicidae, 292
Palicus, 292
Palinuridae, 289
Palinurus, 289
Pallenopsis, 242
Palliatus, 84
Palliolum, 207
Palmaria, 38
Palmariaceae, 38
Palmariales, 38
Palmellopsidaceae, 23
Palmenella, 247
Palmicellaria, 327, 328
Palmiskenea, 327
Palmocoンcha, 249
Palmophyllum, 23
Palmyreuphosyne, 218
Palola, 218
Palpiphitime, 217
Palpophria, 263
Palpophriidae, 263
Palposyllis, 229
Paludinella, 190
Pampus, 374
Pancolinae, 313
Pancolini, 313
Pandalidae, 287
Pandalina, 287
Pandalus, 287
Pandaridae, 261
Pandarus, 261
Pandeia, 118
Pandolaimus, 169
Pandora, 211
Pandoridae, 211
Panetela, 319
Pannychia, 350
Panolicella, 332
Panomya, 210
Panopea, 210
Panopeidae, 292
Panopeus, 292
Panopula, 140
Panousea, 225
Pantachogon, 119
Panthalis, 214
Pantoneura, 32
Pantopipetta, 242
Panturichthys, 360
Panulirus, 289
Papenfussiella, 25
Paphia, 210
Papia, 132
Pappomonas, 76
Papularia, 19
Papuliscala, 186
Parabalaenamertes, 156
Parablennius, 372
Parabothriocephalus, 148
Parabothrium, 148
Parabrotula, 366
Parabrotulidae, 366
Parabyrsophlebs, 133
Paracalanidae, 266
Paracalanus, 266
Paracalliaictis, 108
Paracallisoma, 303
Paracalyptrophora, 105
Paracandacia, 265
Paracanthocephaloides, 177
Paracanthonchus, 169
Paracapillaria, 175
Paracapitella, 216
Paracartia, 263
Paracassidulina, 71
Paraccacladium, 140
Paracentromedon, 303
Paracentrophyes, 160
Paracentrotus, 347
Paracerebratulus, 153
Parachevreuxiella, 303
Parachiridotea, 321
Parachordeumium, 253
Parachristianella, 148
Parachrorhynchus, 132
Paracicera, 131
Paraclione, 200
Paracomantenna, 263
Paracomesoma, 170
Paraconchoecia, 246
Paraconger, 361
Paracoryne, 113
Paracorynidae, 113
Paracoryphella, 205
Paracrenhydrosoma, 271
Paracryptogonimus, 140
Paractaea, 292
Paractinia, 108

- Paracuaria*, 175
Paracucumaria, 348
Paracyatholaimoides, 170
Paracyatholaimus, 170
Paracyathus, 109
Paracycloctyla, 145
Paracyclicolaimus, 170
Paracyphocaris, 303
Paracyprideis, 247
Paracypris, 247
Paracythere, 250
Paracytheridea, 247, 250
Paracytherideidae, 250
Paracytherois, 250
Paradactylopodia, 279
Paradanielssenia, 278
Paradasys, 157
Paradesmodora, 170
Paradexiospira, 228
Paradicranophorus, 150
Paradilepis, 148
Paradiopatra, 221
Paradiplectanotrema, 145
Paradiplozoon, 145
Paradisco, 265
Paradoneis, 222
Paradoris, 203
Paradoxoplana, 136
Paradoxostoma, 250
Paradoxostomatidae, 250
Paradraconema, 170
Paradrepanophoridae, 155
Paradrepanophorus, 155
Paradulichia, 306
Paradyte, 224
Paraedwardsia, 107
Paraegialoalaimus, 170
Paraergasilus, 256
Paraeuinophysa, 218
Parafissurina, 72
Paraflabellula, 53
Parafolliculina, 42
Parafondicularia, 75
Paragalea, 292
Paragaleus, 359
Paragathotanais, 314
Paragigantione, 322
Paragiopagurus, 289
Paraglypturus, 289
Paragnathia, 321
Paragnathiella, 151
Paragnathorhynchus, 131
Paragonaster, 340
Paragorgia, 105
Paragorgiidae, 105
Paragotoea, 113
Paragyliachen, 140
Paragymnomenia, 181
Parahaplogonaria, 126
Parahaplopisthia, 126
Parahedyle, 201
Parahemingwayella, 251
Parahemiurus, 140
Parahistioneis, 51
Parahyalinoecia, 221
Parahypocoma, 46
Parajassa, 301
Parakrithe, 249
Paraledrillus, 233
Paralacydonia, 222
Paralacydoniidae, 222
Paralaespira, 228
Paralamprops, 310
Paralaophonte, 275
Paralaophontodes, 270
Paralcyoniidae, 105
Paralcyonium, 105
Paraleiocapitella, 216
Paralepas, 281
Paralepidapedon, 140
Paralepididae, 363
Paralepis, 364
Paraleptastacus, 276
Paraleptomysis, 294
Paralibrotus, 303
Paralicella, 303
Paralineus, 153
Paralinhomoeus, 170
Paraliparis, 369
Paralipophrys, 372
Paralithodes, 289
Paralomis, 289
Paralongicyatholaimus, 170
Paralophogaster, 293
Paralycaea, 308
Paramblyops, 294
Paramecynostomum, 126
Parameiopsis, 269
Paramesacanthion, 170
Paramesochra, 277
Paramesochridae, 277
Paramesostoma, 134
Parametopa, 307
Paramicrolaimus, 170
Paraminutanemertes, 155
Paramisophria, 263
Paramoeba, 54
Paramoebidae, 53
Paramollicia, 246
Paramonohystera, 170
Paramonostomum, 140
Paramonotus, 135
Paramphiascella, 272
Paramphiascopsis, 272
Paramphilochoides, 297
Paramphinome, 215
Paramphithoe, 299
Paramphitrema, 57
Paramphitrite, 230
Paramphiura, 344
Paramunna, 319
Paramunnidae, 319
Paramunnopsis, 319
Paramuricea, 105
Paramysis, 295
Paramyxilla, 93
Paranaise, 233
Paranaitis, 223
Paranannopus, 278
Paranaperus, 125
Paranarthrura, 314
Paranemertes, 154
Paranemonia, 106
Paranerilla, 221
Paranesidea?, 247
Paraninoe, 219
Paranisakiopsis, 175
Paranisakis, 175
Paranthessius, 258
Paranthura, 316
Paranthuridae, 316
Paranthus, 106
Parantipathes, 109
Paranyphon, 242
Paraonidae, 222
Paraonides, 222
Paraonis, 222
Parapaguridae, 289
Parapagurus, 289
Parapasiphae, 286
Parapenaeus, 286
Paraphanostoma, 125
Paraphellia, 108
Paraphoxus, 306
Paraphrixus, 322
Paraphryotrocha, 217
Paraphyllina, 111
Paraphyllinidae, 111
Parapinnanema, 170
Parapionosyllis, 229
Paraplana, 136
Parapleustes, 306
Parapontella, 266
Parapontellidae, 266
Parapontophilus, 288
Parapriionospio, 227
Parapristipoma, 370
Paraprocerastea, 229
Paraprodilus, 217
Paraproporus, 125
Parapseudes, 313
Parapseudidae, 313
Parapseudoleptomesochra, 269
Parapseudomma, 295
Parapseudotanais, 315
Pararaeolaimus, 170
Pararenopontia, 270
Paragestes, 270
Parargissa, 300
Pararobertsonia, 272
Pararotaliidae, 74
Pararrhopalia, 181
Parartotrogus, 260
Parasaccocoelium, 141
Parasagitta, 335
Parasaveljevia, 170
Parascarophis, 175
Paraschizopera, 278
Paraschizorhynchoides, 132
Parascocotyle, 141
Parasicyonis, 106
Parasinelobus, 313
Parasmittina, 331
Paraspheerolaimus, 170
Paraspheerosyllis, 229
Paraspheophrya, 46
Parasquilla, 284
Parasquillidae, 284
Parastenhelia, 277
Parasteneliidae, 277
Parastenocarididae, 277

- Parastenocaris*, 277
Parastephos, 268
Parasterope, 245
Parasthenelais, 227
Parastichopus, 349
Parastilbe, 187
Parastrombidium, 44
Parastrophia, 190
Parastygarcetus, 236
Parataeniophorus, 368
Paratanaidae, 314
Paratanais, 314
Paratanarctus, 236
Parategastes, 278
Paratenerrhynchus, 131
Paraterschellingia, 170
Parathalassoalaimus, 170
Parathalestris, 279
Parathelepus, 230
Parathelges, 322
Paratiara, 118
Paratimea, 88
Paratimonia, 141
Paratomella, 127
Paratomellidae, 127
Paratricoma, 170
Parathyphosites, 303
Paratuerkiana, 170
Paraturbanella, 158
Paratyposyllis, 229
Paraugaptillus, 264
Parautelga, 132
Paravortex, 130
Paraxanthias, 292
Parazoanthidae, 108
Parazoanthus, 109
Pardalisca, 305
Pardalisella, 305
Pardaliscidae, 305
Pardaliscoides, 305
Pardia, 303
Parellisia, 328
Parepactophanes, 270
Parergodrilidae, 222
Parerythropodium, 104
Parerythrops, 295
Pareteone, 223
Pareuchaeta, 265
Pareudesmoscolex, 170
Pareugyrioides, 354
Pareurystomina, 170
Pareurythoe, 215
Parevansula, 269
Parexocoetus, 367
Parhalixodes, 241
Parhyale, 300
Parhyalella, 300
Pariambus, 296
Paricterotaenia, 148
Paridotea, 321
Parodontophora, 170
Paroedicerops, 305
Paroithona, 255
Paromalostomum, 127
Paromola, 290
Paronesimus, 303
Paronychocamptus, 276
Parophidion, 366
Paroriza, 349
Parotophana, 136
Parotoplanella, 136
Parotoplanina, 136
Parougia, 217
Paroxystomina, 170
Parrina, 67
Parthenope, 291
Parthenopea, 283
Parthenopidae, 291
Parunciola, 306
Parvamoeba, 54
Parvatrema, 141
Parvicapsula, 84
Parvicapsulidae, 84
Parvicardium, 209
Parvicorbicula, 58
Parvioris, 187
Parvipalpus, 296
Parvitaenia, 148
Parviturbo, 184
Parvocythere, 251
Parvotrochus, 351
Paryphostomum, 141
Pasiphaea, 286
Pasiphacidae, 286
Passerilepis, 148
Passeriniella, 15
Pasytheidae, 330
Patella, 182
Patellidae, 182
Patellina, 74
Patellinidae, 74
Pateoris, 68
Paulinella, 56
Paulinellidae, 56
Paulsenella, 51
Pavillardinium, 51
Pawsonia, 348
Paxillosida, 338
Payraudeautia, 191
Peachia, 107
Pecten, 207
Pectinaria, 222
Pectinariidae, 222
Pectinaster, 339
Pectinidae, 207
Pectinodonta, 182
Pectinodrilus, 233
Pectinura, 346
Pedicellaster, 342
Pedicellasteridae, 342
Pedicellina, 179
Pedicellinidae, 179
Pedicularia, 191
Pedicythere, 248
Pedicythere?, 248
Pedinidae, 346
Pedinoida, 346
Pedinosoma, 219
Pedipes, 205
Pedobesia, 21
Pedoculina, 296
Pedunculata, 281
Pegantha, 119
Pegea, 355
Pegusa, 374
Pelacarus, 241
Pelagia, 111
Pelagica, 156
Pelagiidae, 111
Pelagobia, 219
Pelagoalteria, 43
Pelagonema, 170
Pelagonemertidae, 156
Pelagopenaeus, 286
Pelagopleura, 356
Pelagostrobilidium, 44
Pelagothuridae, 350
Pelates, 369
Pelecanidae, 375
Pelecaniformes, 375
Pelecanus, 375
Pelecyophrya, 47
Pelichnibothrium, 148
Pelmatosphaera, 104
Pelmatosphaeridae, 104
Pelogenia, 227
Pelonaia, 355
Pelophila, 125
Pelosina, 63
Pelosphaera, 63
Pelseneeria, 187
Peltaster, 340
Peltidiidae, 277
Peltidium, 277
Peltisenia, 278
Peltobradya, 274
Peltocoxa, 297
Peltogaster, 283
Peltogastrella, 283
Peltogastridae, 283
Peltithestris, 279
Pelvetia, 28
Penaeidae, 286
Penaeopsis, 286
Penaeus, 286
Penares, 86
Pendonemertes, 156
Pendromidae, 184
Pendulomonas, 78
Peneroplidae, 68
Peneroplis, 69
Penetrantia, 332
Penetrantiidae, 332
Peniagone, 350
Penicillium, 19
Penicillus, 21
Peniculus, 262
Penilia, 244
Penilpida, 350
Pennaria, 113
Pennariidae, 113
Pennatula, 105
Pennatulacea, 105
Pennatulicola, 258
Pennatulidae, 105
Pennella, 262
Pennellidae, 262
Pentacoelum, 136
Pentamera, 348
Pentametrocrinidae, 338
Pentametrocrinus, 338

- Pentapharsodinium*, 51
Pentapora, 326
Pentastomida, 251
Penitherichthys, 366
Peponocephala, 376
Peponocyathus, 110
Peracle, 200
Peraclidae, 200
Peracreadium, 141
Peraeospinosus, 315
Perarella, 117
Percichthyidae, 369
Perciformes, 369
Percnon, 292
Percursaria, 24
Pereionotus, 305
Peresiella, 216
Peresime, 279
Periaptia, 186
Periboea, 219
Periclimenes, 288
Peridiniella, 51
Peridiniopsis, 51
Peridinium, 51
Perilinhamoeus, 170
Perinereis, 221
Peringiella, 190
Perioculodes, 305
Peripatagus, 348
Periphragella, 102
Periphylla, 111
Periphyllidae, 111
Periplegmatium, 22
Periplomatidae, 211
Peripygma, 62
Perisscope, 274
Perissopus, 262
Peristediidae, 368
Peristedion, 368
Perkinsiana, 225
Perkinsus, 40
Perna, 206
Perrina, 54
Peroderma, 262
Perophora, 352
Perophoridae, 352
Perrierella, 303
Persa, 119
Persephonaster, 339
Persicula, 194
Personidae, 192
Petalidium, 286
Petaliella, 134
Petalifera, 201
Petaloconchus, 190
Petalodinium, 51
Petalomonas, 59
Petalonia, 28
Petalophthalmidae, 295
Petalophthalmus, 295
Petaloprotus, 220
Petalosarsia, 310
Petasidae, 119
Petasiger, 141
Petasus, 119
Petitia, 229
Petrarca, 283
Petrarcidae, 283
Petricola, 210
Petricolidae, 210
Petrobius, 323
Petrocelidaceae, 36
Petroderma, 27
Petromica, 91
Petromyzon, 358
Petromyzontidae, 358
Petromyzontiformes, 358
Petrosia, 98
Petrosiidae, 98
Petrospongium, 25
Petta, 222
Pettiboneia, 217
Peyssonnelia, 35
Peyssonneliaceae, 35
Peziza, 17
Pezizales, 18
Phacellophora, 111
Phaenna, 266
Phaennidae, 266
Phaenocora, 134
Phaecystis, 76
Phaeonectriella, 18
Phaeophila, 23
Phaeophilaceae, 23
Phaeophilales, 23
Phaeophyceae, 24
Phaeophycota, 24
Phaeosphaeria, 16
Phaeosphaeriaceae, 16
Phaeostachys, 331
Phaeostroma, 27
Phaethon, 375
Phaethontidae, 375
Phagicola, 141
Phagus, 258
Phainogullmia, 62
Phakellia, 90
Phalacrocoracidae, 375
Phalacrocorax, 375
Phalacrophorus, 219
Phalacrostemma, 225
Phalaropodidae, 375
Phalaropus, 375
Phalium, 191
Phalldriliinae, 233
Phalldrilius, 233
Phallonemertes, 156
Phallonemertidae, 156
Phallusia, 352
Phallusiella, 256
Phanoderma, 170
Phanodermella, 171
Phanodermopsis, 171
Pharcidia, 16
Pharidae, 209
Pharodes, 257
Pharodidae, 257
Pharus, 209
Phascolion, 178
Phascolionidae, 178
Phascolosoma, 178
Phascolosomatidae, 178
Phaxas, 209
Phellia, 108
Phelliactis, 108
Pheopolykrikos, 51
Pheronema, 102
Pheronematidae, 102
Pherusa, 218
Pherusella, 332
Pherusellidae, 332
Phialellidae, 116
Phialidium, 119
Phialophorophoma, 19
Phialopsis, 114
Phidiana, 204
Phidolopora, 330
Phidoloporidae, 330
Philachoerusr, 125
Philactinoposthia, 125
Philichthyidae, 257
Philichthys, 257
Philine, 199
Philinidae, 199
Philinoglossa, 200
Philinoglossidae, 200
Philinopsis, 199
Philippia, 196
Philobrythos, 148
Philocelis, 127
Philocheras, 288
Philodinidae, 151
Philomecynostomum, 126
Philomedes, 245
Philomedidae, 246
Philometra, 175
Philometroides, 176
Philopthalmus, 141
Philorthagoriscus, 260
Philosyrtis, 136
Phipsia, 307
Phippsiella, 307
Phisidia, 230
Phlebobranchia, 352
Phliantidae, 305
Phloeodictyidae, 98
Phlyctaenopora, 97
Phlyctocythere, 249
Phoca, 376
Phocascaris, 176
Phoceana, 330
Phoceanidae, 330
Phocidae, 376
Phocoena, 376
Phocoenidae, 376
Pholadidae, 210
Pholadidea, 210
Pholadomya, 211
Pholadomyidae, 211
Pholas, 210
Pholeter, 141
Pholidae, 371
Pholis, 371
Pholoe, 222
Pholoidae, 222
Pholoides, 222
Phoma, 19
Phomatospora, 17
Phonorhynchella, 132
Phonorhynchus, 132

- Phorbas*, 96
Phorcus, 183
Phormosoma, 346
Phoronida, 325
Phoronis, 325
Phoronopsis, 325
Phosichthyidae, 362
Photis, 301
Photorectes, 363
Photostomias, 363
Photostylus, 362
Phoxichilidiidae, 242
Phoxichilidium, 243
Phoxocephalidae, 305
Phoxocephalus, 306
Phragmonemataceae, 38
Phronima, 308
Phronimidae, 308
Phrosina, 308
Phrosinidae, 308
Phryganella, 56
Phryganellidae, 56
Phryganellina, 56
Phrynocrinidae, 338
Phrynophiurida, 343
Phrynorhombus, 374
Phterichthys, 369
Phtisica, 296
Phycidae, 365
Phycis, 365
Phycodrys, 32
Phylactella, 331
Phyllangia, 109
Phyllaplysia, 201
Phyllariaceae, 28
Phyllariopsis, 28
Phyllidia, 203
Phyllidiidae, 203
Phyllidiopsis, 203
Phylliroe, 203
Phylliroiidae, 203
Phyllobothrium, 148
Phyllochaetopreus, 216
Phyllochona, 41
Phyllochonidae, 41
Phyllochoraceae, 18
Phyllochorales, 18
Phyllodicola, 258
Phyllodicolidae, 257
Phyllodictyon, 22
Phyllodistomum, 141
Phyllodoce, 223
Phyllodocidae, 222
Phyllognathopodidae, 277
Phyllognathopus, 277
Phyllomitus, 79
Phyllophora, 36
Phyllophoraceae, 36
Phyllophoridae, 348
Phyllophorus, 348
Phyllopodidae, 267
Phyllopodopsyllus, 278
Phyllopus, 267
Phyllorhinichthys, 367
Phyllorhiza, 112
Phyllothalestris, 279
Phyllothyreus, 262
Phylo, 222
Phymantidae, 108
Phymanthus, 108
Phymatolithon, 34
Phymorhynchus, 196
Physalia, 122
Physaliidae, 122
Physeter, 376
Physeteridae, 376
Physetocaris, 287
Physiculus, 365
Physonectae, 122
Physophora, 122
Physophoridae, 122
Pierrickia, 171
Pigromitus, 242
Pikea, 35
Pilalla, 62
Pilargidae, 223
Pilargis, 223
Pilayella, 27
Pilayellaceae, 27
Pileolaria, 228
Pilifera, 276
Pilinia, 27
Piliscus, 191
Pilosanthora, 316
Pilulina, 64
Pilumnidae, 292
Pilumnoides, 292
Pilumnoididae, 292
Pilumnopeus, 292
Pilumnus, 292
Pilus, 183
Pinctada, 207
Pinguinus, 375
Pinna, 207
Pinnidae, 207
Pinnotheres, 292
Pinnotheridae, 292
Pionosyllis, 229
Pirimela, 291
Pirimelidae, 291
Pirodrilus, 233
Piromis, 218
Pirsonia, 79
Pisa, 291
Pisania, 193
Pisanianura, 192
Pisanianuridae, 192
Piscicillaria, 176
Piscicola, 235
Piscicolidae, 234
Piseinotecidae, 205
Piseinotecus, 205
Pisidia, 290
Pisinna, 190
Pisone, 223
Pisionidae, 223
Pisodonophis, 361
Pisolamia, 187
Pista, 230
Pistana, 148
Pistella, 231
Pitar, 210
Pityrete, 103
Placida, 201
Placinolopha, 85
Placiphorella, 182
Placogorgia, 105
Placopsilina, 62, 75
Placopsilinella, 62
Placorhynchidae, 132
Placorhynchus, 132
Placospongia, 88
Placospongiidae, 88
Placostegus, 226
Placotrochides, 110
Placozoa, 103
Plagiobrissus, 347
Plagiocardium, 209
Plagioecia, 333
Plagioporus, 141
Plagiorchis, 141
Plagiorhynchus, 177
Plagioselmis, 59
Plagiospora, 35
Plagiostomidae, 129
Plagiostomula, 129
Plagiostomum, 129
Plagusia, 292
Plagystila, 190
Plakina, 85
Plakinastrella, 85
Plakinidae, 85
Plakortis, 85
Plakosyllis, 229
Planaxidae, 185
Planaxis, 185
Planctosphaera, 336
Planes, 292
Planifolliculina, 42
Planinspirinella, 68
Planinspirinoides, 67
Planktonemertidae, 156
Planocera, 128
Planoceridae, 128
Planodasyidae, 157
Planodinium, 51
Planogypsina, 74
Planophila, 23
Planopulvinulina, 72
Planorbulina, 74
Planorbulinidae, 74
Planoscalpellum, 282
Plantae, 38
Planulina, 69
Planulinidae, 74
Platichthys, 374
Platidia, 334
Platidiidae, 334
Platocystis, 141
Platoma, 37
Platonea, 333
Platorchestia, 308
Platyamoeba, 54
Platybdella, 235
Platybelone, 367
Platybothrium, 148
Platycephalidae, 368
Platycephalus, 368
Platyhelipus, 276
Platychilomonas, 59
Platycopoma, 171

- Platycomopsis*, 171
Platycopia, 262
Platycopida, 247
Platycopiidae, 262
Platycopioidea, 262
Platycetenida, 123
Platycetenidae, 123
Platydasys, 157
Platydorididae, 203
Platydoris, 203
Platyfolliculina, 42
Platyhedyle, 201
Platyhedyliidae, 201
Platylaophonte, 276
Platylepadidae, 282
Platylepas, 282
Platynereis, 221
Platypleura, 58
Platypodiella, 292
Platyscelidae, 308
Platyscelus, 308
Platyscolex, 148
Platysiphonia, 32
Platysporina, 84
Platysquilla, 284
Platysquilloides, 284
Platysympus, 310
Platytroctes, 362
Platytroctidae, 362
Plectanocotyle, 145
Plectanocotyloides, 145
Plectofrondiculariidae, 74
Plectorhinchus, 370
Plectroninia, 102
Plehnia, 128
Plejniidae, 128
Pleioplana, 128
Pleioplanidae, 128
Pleocola, 236
Pleodicyema, 104
Pleomassariaceae, 16
Pleonosporium, 31
Pleopsis, 244
Pleorchis, 141
Pleospora, 16
Pleosporaceae, 16
Pleraplysilla, 101
Plerocercoides, 148
Plerurus, 141
Plesiochorus, 141
Plesiadiadema, 346
Plesionika, 287
Plesiopenaeus, 285
Plesiothoa, 330
Pleurage, 18
Pleurasiga, 58
Pleurobrachia, 122
Pleurobrachiidae, 122
Pleurobranchaea, 201
Pleurobranchidae, 201
Pleurobranchus, 201
Pleurocladia, 27
Pleurocope, 319
Pleurocopidae, 319
Pleurocrypta, 322
Pleurocryptella, 322
Pleurodasys, 157
Pleurogoniidae, 319
Pleurogonium, 319
Pleurogonius, 141
Pleuromamma, 266
Pleuronectes, 374
Pleuronectidae, 374
Pleuronectiformes, 374
Pleuroprion, 321
Pleurostomella, 74
Pleurotomellidae, 74
Pleurotomella, 196
Pleustes, 306
Pleustidae, 306
Pleustomesus, 306
Pleusyntes, 306
Plexauridae, 105
Plexechinidae, 347
Plexechinus, 347
Plicastoma, 129
Plicobothrium, 148
Plinthaster, 340
Pliobothrus, 118
Pliophrixus, 322
Plocamiaceae, 38
Plocamiales, 38
Plocamiancora, 93
Plocamionida, 96
Plocamium, 38
Plocamopherus, 202
Ploeotia, 60
Ploesoma, 150
Ploima, 150
Plotocnide, 113
Plotohelmis, 214
Plotonemertes, 156
Plumaria, 31
Plumularia, 116
Plumulariidae, 116
Plutonaster, 339
Pneophyllum, 34
Pneumoderma, 200
Pneumodermatidae, 200
Pneumodermopsis, 200
Pocilloporidae, 110
Pocillorhynchus, 131
Podarke, 219
Podarkeopsis, 219
Podicipedidae, 376
Podicipediformes, 376
Podicipes, 376
Podoceridae, 306
Podocerus, 306
Podocidar, 346
Podocopida, 247
Podocryna, 117
Podocytyle, 141
Pododesmus, 207
Podolampas, 51
Podolampus, 51
Podon, 244
Podonidae, 244
Podoprion, 303
Podopriionella, 303
Podosphaeraster, 341
Podospongia, 88
Poecilancistrum, 148
Poecilasma, 281
Poecilasmatidae, 281
Poecillastra, 86
Poecilochaetidae, 223
Poecilochaetus, 223
Poecilosclerida, 91
Poecilostomatida, 255
Pogaina, 130
Pogonodon, 185
Pogonophora, 235
Pogotrichiaceae, 26
Pogotrichium, 26
Poirieria, 192
Polaruschakov, 224
Policordia, 211
Poliometra, 338
Poliopsis, 154
Politolana, 320
Pollachius, 365
Pollia, 193
Pollichthys, 362
Pollicipedidae, 281
Pollicipes, 281
Polyacanthonotus, 360
Polyandrocarpa, 355
Polyarthra, 150
Polybius, 291
Polybrachiidae, 235
Polybranchia, 200
Polycarpa, 355
Polycerata, 203
Polycerella, 203
Polyceridae, 203
Polychaeta, 214
Polycheles, 288
Polyhelidae, 288
Polycirrus, 231
Polycitor, 353
Polycitoridae, 353
Polycladida, 128
Polyclinella, 353
Polyclinidae, 353
Polyclinum, 353
Polycliphora, 145
Polyclithrum, 145
Polycope, 246
Polykopidae, 246
Polycope, 246
Polycyathus, 109
Poly cystididae, 132
Poly cystis, 132
Polydora, 227
Polyfibula, 58
Polygastrophora, 171
Polygordiidae, 223
Polygordius, 223
Polyidaceae, 37
Polyides, 37
Polyipnus, 362
Polykrikos, 51
Polylabris, 145
Polyleberis, 245
Polylekithum, 141
Polymastia, 89
Polymastiidae, 89
Polymerurus, 157
Polymetme, 363
Polymixia, 365

- Polymixiidae**, 365
Polymixiiformes, 365
Polymorphina, 74
Polymorphanidae, 74
Polymorphus, 177
Polynemidae, 371
Polyneura, 32
Poly noe, 224
Polynoidae, 223
Polyodontes, 214
Polyoeca, 58
Polyopthalmus, 222
Polyphysaceae, 23
Polyphysia, 226
Polyplacophora, 182
Polyplumaria, 116
Polyposthia, 128
Polyposthiidae, 128
Polyprion, 369
Polyprionidae, 369
Polysaccus, 283
Polysigma, 171
Polysiphonia, 31
Polysporoplasma, 83
Polystilifera, 155
Polystomammina, 66
Polystrata, 35
Polystyliphora, 136
Polystyliphoridae, 136
Polysyncraton, 353
Pomacentridae, 371
Pomadasys, 370
Pomatoceros, 226
Pomatodinium, 51
Pomatodrilidae, 224
Pomatodrilus, 224
Pomatoleios, 226
Pomatomidae, 369
Pomatomus, 369
Pomatoschistus, 373
Pomoriella, 56
Pompholyx, 151
Pomphorhynchus, 177
Pomponema, 171
Pontalaria, 132
Pontaster, 339
Pontella, 267
Pontellidae, 267
Pontellina, 267
Pontelopsis, 267
Pontinus, 368
Pontobdella, 235
Pontobdellina, 235
Pontoclausia, 255
Pontocletodes, 271
Pontocrates, 305
Pontocypris, 251
Pontocyprididae, 251
Pontocypridoidea, 251
Pontocypris, 251
Pontocythere, 247
Pontodora, 225
Pontodoridae, 224
Pontodrilus, 234
Pontoeciella, 262
Pontoeciellidae, 262
Pontogeneia, 15, 299
Pontogenia, 215
Pontochedyle, 201
Pontolineus, 153
Pontonema, 171
Pontonia, 288
Pontophilus, 288
Pontophonte, 276
Pontopolites, 275
Pontoporeia, 16, 306
Pontoporeiidae, 306
Pontoptilius, 264
Pontostratiotes, 271
Poracanthium, 141
Poralia, 111
Porania, 340
Poraciidae, 340
Poraniomorpha, 340
Porcellana, 290
Porcellanaster, 339
Porcellanasteridae, 339
Porcellanidae, 290
Porcellidiidae, 277
Porcellidium, 277
Porella, 327
Poria, 270
Porifera, 85
Poriocidaris, 346
Poroeponides, 72
Poromitra, 367
Poromya, 211
Poromyidae, 211
Porphyra, 29
Porphyridiaceae, 38
Porphyridiales, 38
Porphyridium, 38
Porphyrocrinidae, 338
Porphyrocrinus, 338
Porphyropsis, 35
Porphyrostromium, 35
Porpita, 113
Porpitidae, 113
Porrocaecum, 176
Porroecia, 246
Portatrocchamina, 66
Porterinema, 25
Portlandia, 206
Portunus, 291
Portunidae, 291
Portunus, 291
Posidonia, 39
Posidoniaceae, 39
Postaphanostoma, 126
Postbursovula, 136
Posterula, 331
Postgaardi, 79
Posthodiplostomum, 141
Postmecynostomum, 127
Potamethus, 226
Potamides, 185
Potamididae, 185
Potamilla, 226
Potamogotonales, 39
Potamothrrix, 234
Pothametus, 226
Pourtalecia, 347
Pourtaleciidae, 347
Pourtalecioida, 347
Pourtalosmilia, 109
Pozziella, 98
Prachynella, 303
Praeacanthonchus, 171
Praeanaperus, 125
Praeaphanostoma, 126
Praebursovula, 136
Praeconvoluta, 126
Praeglobobulimina, 70
Praeleptomesochra, 269
Prasinophytes, 76
Prasiola, 23
Prasiolaceae, 23
Prasiolales, 23
Pratoplana, 134
Praunus, 295
Praxillella, 220
Praxillura, 220
Praya, 121
Prayidae, 121
Prayola, 121
Predaea, 36
Preminona, 135
Premocyathus, 109
Preperidinium, 51
Presynaptiphilus, 258
Preussia, 16
Priacanthidae, 369
Priacanthus, 369
Priapocephalus, 148
Priapulida, 160
Priapulidae, 160
Priapulopsis, 160
Priapulus, 160
Primno, 308
Primnoa, 105
Primnoella, 105
Primnoidae, 105
Primovula, 191
Princaxelia, 305
Pringsheimiella, 22
Prionace, 359
Prionospio, 227
Prionotoleberis, 245
Priscillina, 306
Pristidae, 359
Pristiformes, 359
Pristigloma, 205
Pristiglomidae, 205
Pristis, 359
Proaeginina, 296
Proagnesia, 352
Proales, 150
Proameira, 269
Proaphanostoma, 126
Probalaenamertes, 156
Probolitrema, 141
Proboloides, 307
Proboscidactyla, 118
Proboscidactylidae, 118
Proboscoida, 119
Procamacolaimus, 171
Procampylaspis, 310
Procellariidae, 375
Procellariiformes, 375
Procephalothrix, 152
Proceraea, 229

- Procerastea*, 229
Procerodes, 136
Procerodidae, 136
Procerocetia, 246
Procerolagena, 74
Proceropharynx, 134
Processa, 288
Processidae, 288
Prochaetoderma, 181
Prochaetodermatidae, 181
Prochaetosoma, 171
Prochelator, 317
Prochromadora, 171
Prochromadorella, 171
Proclea, 231
Proclymene, 220
Proconvoluta, 126
Proctoeces, 141
Proctonotidae, 204
Proctonotus, 204
Proctophantastes, 141
Proctotrema, 141
Prodisco, 265
Prodistomum, 141
Profundivermis, 141
Progebiophilus, 322
Progeryon, 292
Prognathodes, 371
Prognathorhynchus, 131
Progoniada, 218
Progonus, 141
Proidotea, 321
Prolechithophora, 129
Proleptus, 176
Promesostoma, 133
Promesostomidae, 133
Promethichthys, 373
Promonotus, 135
Proneomenia, 181
Proneomeniidae, 181
Proneurotes, 154
Pronoictiluca, 51
Pronoidae, 308
Pronoprymna, 141
Prooncholaimus, 171
Proparvipyrum, 141
Propeamussium, 207
Prophioseides, 255
Propilidium, 182
Propontocypris, 251
Proporidae, 127
Proporus, 127
Prorocentrum, 51
Proschorizorhynchella, 132
Proschorizorhynchoides, 133
Proschorizorhynchus, 133
Proseriata, 135
Prosobothrium, 148
Prosorhochmidae, 155
Prosorhochmus, 155
Prosorhynchoides, 141
Prosorhynchus, 141
Prospaerolaimus, 171
Prostheceraeus, 128
Prosthiosomatidae, 128
Prosthiosomum, 128
Prosthogonimus, 141
Prostomatella, 155
Prostomiopsis, 155
Prosuberites, 88
Protacanthamoeba, 53
Protandrella, 133
Protankyra, 350
Protanthea, 107
Protasperococcus, 26
Protaspis, 78
Protatlanta, 192
Protectocarpus, 25
Proteocephalus, 148
Proterorhinus, 373
Proterospongia, 58
Protiera, 118
Protiaridae, 118
Protis, 226
Protoaricia, 222
Protoceratium, 52
Protochromadora, 171
Protocirrineris, 217
Protocotyle, 145
Protocuspidaaria, 211
Protodactylina, 260
Protoderma, 24
Protodorvillea, 217
Protodrilidae, 225
Protodriloides, 225
Protodrilus, 225
Protoglossus, 336
Protogoniceps, 279
Protogrammus, 372
Protoholozoa, 353
Protohydra, 113
Protohydridae, 113
Protolaeospira, 228
Protolamellodiscus, 145
Protolira, 184
Protomedieia, 301
Protomicrocotyle, 145
Protomolgula, 354
Protomonostroma, 24
Protomonotresidae, 129
Protomyctophum, 364
Protomystides, 223
Protomyszostomum, 220
Protopalyntha, 109
Protopelagonemertes, 156
Protopelagonemertidae, 156
Protoperidinium, 52
Protopsammotopa, 272
Protoptilidae, 106
Prototilum, 106
Protoschista, 64
Protosquilla, 284
Protosquillidae, 284
Protostyela, 355
Protubulanus, 152
Protula, 226
Provortex, 130
Provorticidae, 130
Proxenetes, 134
Prudhoeus, 141
Pruvotfolia, 204
Pruvotia, 181
Pruvotina, 181
Pruvotinidae, 181
Psalliora, 18
Psamathea, 276
Psammameira, 269
Psammaphiporus, 154
Psammascidia, 352
Psammastacus, 276
Psammechinus, 347
Psammettidae, 78
Psammina, 78
Psamminidae, 78
Psammis, 278
Psammobiidae, 209
Psammocampus, 270
Psammocythere, 251
Psammodasys, 157
Psammodrilidae, 225
Psammodriloides, 225
Psammodrilus, 225
Psammogammarus, 304
Psammohydra, 112
Psammomacrostomum, 128
Psammonobiotidae, 56
Psammonobiotus, 56
Psammopemma, 101
Psammopolycystis, 132
Psammopsyllidae, 277
Psammopsyllus, 277
Psammorhynchidae, 132
Psammorhynchus, 132
Psammoryctides, 234
Psammosphaera, 64
Psammostyela, 355
Psammotopa, 272
Psathyrocaris, 286
Psednos, 369
Pseliodinium, 52
Pselionema, 171
Psenes, 374
Pseudactinoposthia, 125
Pseudaephnidiogenes, 141
Pseudalataspora, 84
Pseudalcyonidium, 332
Pseudalius, 176
Pseudameira, 269
Pseudamphiascopsis, 272
Pseudamphistomum, 141
Pseudamussium, 207
Pseudanisakis, 176
Pseudanomotaenia, 148
Pseudanthessiidae, 258
Pseudanthessius, 258
Pseudanthobothrium, 148
Pseudanthocotyloides, 145
Pseudaphanostoma, 126
Pseudaphya, 373
Pseudarachna, 318
Pseudarchaster, 340
Pseudaugaptilus, 264
Pseudaxine, 145
Pseudaxinella, 90
Pseudectinosoma, 274
Pseudendoclonium, 22
Pseudeuchaeta, 263
Pseudevadne, 244
Pseudhaloptilus, 264
Pseudharpinia, 306
Pseudicyema, 104

- Pseudione*, 322
Pseudmecynostomum, 127
Pseudoanthocotle, 145
Pseudobacciger, 141
Pseudobaseodiscus, 154
Pseudobodo, 77
Pseudobolivina, 70
Pseudobothrium, 148
Pseudobradya, 274
Pseudobrania, 229
Pseudocalanus, 265
Pseudocaligus, 260
Pseudocapillaria, 176
Pseudocapitella, 216
Pseudocaranx, 370
Pseudocella, 171
Pseudoceros, 128
Pseudocerotidae, 128
Pseudochama, 208
Pseudocharopinus, 261
Pseudochirella, 263
Pseudochlorodesmis, 21
Pseudoclausia, 255
Pseudoclavulina, 64
Pseudocletoedes, 275
Pseudocoeculinidae, 183
Pseudoconchoecia, 246
Pseudocorticium, 85
Pseudocorynactis, 108
Pseudocorythion, 56
Pseudocotyle, 145
Pseudocuma, 310
Pseudocumatidae, 310
Pseudocyclopia, 267
Pseudocyclopidae, 267
Pseudocyclopiidae, 267
Pseudocyclopinoidea, 254
Pseudocyclops, 267
Pseudocycnidae, 262
Pseudocycnus, 262
Pseudocythere, 247
Pseudocytherura, 248
Pseudocyttus, 368
Pseudodactylogyrus, 145
Pseudodendoclonium, 24
Pseudodiaptomidae, 267
Pseudodiazona, 352
Pseudodifflugia, 56
Pseudodifflugiidae, 56
Pseudodosaccopsis, 272
Pseudodosaccus, 272
Pseudodiplectanum, 145
Pseudodistoma, 353
Pseudoeponides, 73
Pseudoeucanths, 255
Pseudoeurotium, 17
Pseudofabricia, 226
Pseudofabriciola, 226
Pseudoflustra, 326
Pseudogaudryinidae, 64
Pseudogerda, 317
Pseudograffilla, 130
Pseudogrillotia, 148
Pseudohalichondria, 96
Pseudohaplogonaria, 126
Pseudokuhnia, 145
Pseudolaophonte, 276
Pseudoleiocapitella, 216
Pseudolella, 171
Pseudoleptochelia, 314
Pseudoleptomesochrella, 269
Pseudolirius, 296
Pseudolithoderma, 25
Pseudolithophyllum, 34
Pseudolividae, 194
Pseudoloxoconcha, 249
Pseudolubbockia, 257
Pseudomalaxis, 196
Pseudomalletia, 206
Pseudomaritrema, 141
Pseudomassilina, 67
Pseudomastus, 216
Pseudomazocraes, 145
Pseudomelampus, 205
Pseudomesidae, 319
Pseudomesochra, 278
Pseudomesus, 319
Pseudometachromadora, 171
Pseudomma, 295
Pseudomonocelis, 135
Pseudomyicola, 257
Pseudomysis, 295
Pseudomystides, 223
Pseudoncholaimus, 171
Pseudonchus, 171
Pseudonereis, 221
Pseudonezumia, 365
Pseudoniphargus, 304
Pseudonotholca, 150
Pseudonotomastus, 216
Pseudonychocampus, 276
Pseudopallene, 242
Pseudoparamoeba, 54
Pseudoparanais, 315
Pseudoperanema, 60
Pseudophaeenna, 268
Pseudophichthys, 361
Pseudophilomedes, 245
Pseudoplumaria, 116
Pseudopolinices, 191
Pseudopolydora, 228
Pseudopolymorpha, 74
Pseudoposthia, 126
Pseudopotamilla, 226
Pseudopringsheimia, 22
Pseudoproleptus, 176
Pseudoprotella, 296
Pseudopsyllus, 255
Pseudopycnadena, 141
Pseudopythina, 208
Pseudorbis, 185
Pseudorca, 376
Pseudorthoplana, 136
Pseudosagitta, 335
Pseudosammocythere, 249
Pseudosarsameira, 269
Pseudoscalibregma, 226
Pseudoschizorhynchoides, 133
Pseudoscilla, 197
Pseudoscopelus, 372
Pseudoscorpiones, 237
Pseudoscourfieldia, 76
Pseudosetia, 190
Pseudosimnia, 191
Pseudosinella, 324
Pseudophyrapinae, 313
Pseudophyrapus, 313
Pseudospora, 31
Pseudosquillopsis, 284
Pseudosteineria, 171
Pseudostichopus, 349
Pseudostomella, 158
Pseudostomidae, 129
Pseudostomum, 129
Pseudosuberites, 88
Pseudosyllides, 229
Pseudosyllis, 229
Pseudosyrtis, 136
Pseudotachidiidae, 277
Pseudotachidius, 278
Pseudotanaidae, 315
Pseudotanainae, 315
Pseudotanais, 315
Pseudoterranova, 176
Pseudotharybis, 263
Pseudothyone, 349
Pseudotiron, 307
Pseudotolithus, 370
Pseudotorellia, 191
Pseudotorinia, 196
Pseudotrachya, 88
Pseudotriakidae, 359
Pseudotriakis, 359
Pseudotriloculina, 67
Pseudoturbanella, 158
Pseudovermidae, 205
Pseudovermilia, 226
Pseudovermis, 205
Pseudowebbinella, 65
Pseudozoogonoides, 141
Pseudupeneus, 371
Psilaster, 339
Psilochasmus, 141
Psilodens, 181
Psilomyzon, 259
Psilostomum, 141
Psiloteredo, 210
Psittacorhynchus, 131
Psolidae, 348
Psolidium, 348
Psolus, 348
Psychrolutes, 369
Psychrolutidae, 369
Psychroplanes, 350
Psychropotes, 350
Psychropotidae, 350
Psyllocampus, 269
Pteraclis, 370
Pteraster, 341
Pterastericola, 130
Pterastericolidae, 130
Pterasteridae, 341
Pteria, 207
Pteriacartia, 263
Pteriidae, 207
Pterinopsyllus, 254
Pterobdella, 235
Pterobothrium, 148
Pterohranchia, 336
Pterocirrus, 223
Pterocladia, 36

- Pterocladiella*, 36
Pterocladiophilaceae, 37
Pterocopus, 213
Pterocuma, 310
Pterodroma, 375
Pteroëides, 106
Pterognathia, 151
Pterognathiidae, 151
Pterognathus, 131
Pteromeris, 208
Pteromylaeus, 360
Pteropontis, 259
Pteropsyllus, 279
Pterosagitta, 335
Pterosagittidae, 335
Pterosiphonia, 33
Pterothamnion, 31
Pterorachea, 192
Pterrocirrus, 223
Pterycombus, 370
Pterygioteuthis, 212
Pterygocythereis, 251
Pterygonema, 171
Pterynotus, 192
Ptilocladiopsis, 31
Ptilocodiidae, 118
Ptilota, 31
Ptilothamnion, 31
Ptyalorhynchus, 131
Ptychobothrium, 148
Ptychodactiaria, 108
Ptychodactiidae, 108
Ptychodactis, 108
Ptychoderidae, 336
Ptychodiscus, 52
Ptychogastria, 119
Ptychogastriidae, 119
Ptychogena, 115
Ptychogonimus, 141
Ptycholaimellus, 171
Ptychomiliola, 67
Ptychopera, 134
Ptychostomella, 158
Puellina, 329
Puffinus, 375
Pullenia, 71
Pullularia, 19
Pulsellidae, 212
Pulsellum, 212
Punctaria, 26
Punctariaceae, 26
Punctiscala, 186
Punctodora, 171
Puncturella, 183
Pungitius, 368
Punnettia, 155
Purpuradusta, 191
Pusillargillus, 271
Pusillina, 188
Pusillotrocha, 217
Pussylineus, 153
Putzeysia, 184
Pycnadenoides, 141
Pycnanthus, 106
Pycnoclavella, 352
Pycnogonida, 241
Pycnogonidae, 243
Pycnogonum, 243
Pycnophyes, 160
Pycnophyidae, 160
Pyelosomum, 141
Pygidiopsis, 141
Pygophora, 281
Pygospio, 228
Pyragraphorus, 145
Pyramidellidae, 197
Pyramimonas, 76
Pyrenocollerma, 18
Pyrenolaceae, 18
Pyrenolales, 18
Pyrenopeziza, 17
Pyrgo, 67
Pyrgoella, 67
Pyrgomatidae, 282
Pyripora, 329
Pyrocystis, 52
Pyroleptomyysis, 295
Pyrophacus, 52
Pyrosoma, 355
Pyrosomatidae, 355
Pyrostephidae, 122
Pyrostremma, 355
Pyroteuthidae, 212
Pyroteuthis, 212
Pyrulina, 67
Pyrunculus, 199
Pythonaster, 342
Pyura, 354
Pyuridae, 354
Pyxidicula, 55
Quadrililia, 79
Quadicoma, 172
Quantanthura, 316
Quasillina, 89
Questa, 225
Questidae, 225
Quinquaophonte, 276
Quinqueloculina, 67
Quintoxilla, 93
Raabella, 46
Rabilimis, 251
Rachycentridae, 369
Rachycentron, 369
Racodiscula, 87
Racovitzanus, 267
Radiaster, 339
Radiasteridae, 339
Radicilingua, 32
Radicipes, 104
Radicula, 65
Radiicephalidae, 364
Radiicephalus, 364
Ragonula, 331
Raiarctus, 236
Raja, 360
Rajella, 360
Rajidae, 359
Rajiformes, 359
Rajonchocotide, 145
Ralfsia, 25
Ralfsiaceae, 25
Ramphonotus, 328
Ranella, 192
Ranellidae, 191
Raniceps, 365
Raninidae, 290
Ransonnia, 119
Ranzania, 374
Rapana, 192
Rapaniscus, 319
Raphidascaris, 176
Raphitoma, 195
Raricirrus, 217
Raspaciona, 92
Raspailia, 92
Raspailiidae, 91
Rastrelliger, 373
Rastrognathia, 151
Rastrognathiidae, 151
Ratania, 262
Rataniidae, 262
Rathkeia, 118
Rathkeidae, 118
Rectilabrum, 187
Rectobolivina, 70
Rectobuntonia, 251
Rectonychocella, 330
Rectuvigerina, 75
Recurvoidella, 66
Recurvoidea, 64
Redekea, 250
Regadrella, 103
Regalecidae, 364
Regalecus, 364
Reighardia, 251
Reighardiidae, 251
Reinhardtius, 374
Reisingeria, 129
Relichna, 199
Remanea, 277
Remaneica, 75
Remipedia, 243
Remispora, 17
Remora, 370
Remorina, 370
Renaudarctidae, 236
Renicola, 141
Reophax, 64
Repmanina, 62
Reptadeonella, 326
Reptantia, 155
Reteporella, 330
Reteporellina, 330
Retevirgula, 328
Reticulammina, 78
Reticulamoeba, 54
Reticulomyxidae, 54
Retigyra, 185
Retinometra, 148
Retronectes, 127
Retronectidae, 127
Retrotortina, 196
Retusa, 199
Retusidae, 199
Reussella, 75
Reussellidae, 75
Rhabdaminella, 63
Rhabdammina, 63
Rhabdamoeba, 54
Rhabderemia, 92
Rhabdermiidae, 92

- Rhabdeurypon*, 92
Rhabditis, 172
Rhabdobaris, 90
Rhabdochona, 176
Rhabdocoela, 129
Rhabdocoma, 172
Rhabdodemania, 172
Rhabdodictyum, 103
Rhabdomolgus, 351
Rhabdonia, 37
Rhabdoon, 113
Rhabdoplectella, 103
Rhabdopleura, 336
Rhabdopleuridae, 336
Rhabdoploca, 92
Rhabdopus, 258
Rhachotropis, 299
Rhadinema, 172
Rhadinesthes, 363
Rhadinorhynchus, 177
Rhamphobrachium, 221
Rhamphostomella, 327
Rhamphostomellina, 328
Rhaphidorus, 89
Rhaphidrilus, 217
Rhaptapagis, 118
Rhincalanus, 265
Rhincodon, 358
Rhincodontidae, 358
Rhinebothrium, 148
Rhinecotyle, 145
Rhinobatidae, 359
Rhinobatos, 359
Rhinochimaera, 360
Rhinochimaeridae, 360
Rhinoclama, 211
Rhinoclavis, 185
Rhindiaphana, 199
Rhinolabia, 303
Rhinomolgus, 255
Rhinoptera, 360
Rhipidocotyle, 141
Rhipidogammarus, 300
Rhipilia, 21
Rhips, 172
Rhithropanopeus, 292
Rhizammina, 63
Rhizamoeba, 53
Rhizangiidae, 110
Rhizaxinella, 88
Rhizocaulus, 119
Rhizocephala, 283
Rhizoclonium, 22
Rhizocrinus, 337
Rhizomonas, 79
Rhizoniera, 100
Rhizophyllidaceae, 35
Rhizophysa, 122
Rhizophysidae, 122
Rhizopriionodon, 359
Rhizopus, 15
Rhizorhagium, 117
Rhizorrhina, 261
Rhizostoma, 112
Rhizostomatidae, 112
Rhizostomeae, 111
Rhizothricidae, 278
Rhizothrix, 278
Rhodaliidae, 122
Rhodella, 38
Rhodichthys, 369
Rhodine, 220
Rhodinicola, 256
Rhodochaetaceae, 38
Rhodochaetales, 38
Rhodocheate, 38
Rhodochorton, 29
Rhodomela, 33
Rhodomelaceae, 32
Rhomonas, 59
Rhodope, 201
Rhodophyceae, 29
Rhodophycota, 29
Rhodophyllis, 36
Rhodophysema, 38
Rhodophysemataceae, 38
Rhodopidae, 201
Rhodopsis, 226
Rhodosoma, 352
Rhodosorus, 38
Rhodospora, 38
Rhodostethia, 375
Rhodothamniella, 38
Rhodothamniellaceae, 38
Rhodymenia, 38
Rhodymeniaceae, 38
Rhodymeniales, 38
Rhombifera, 353
Rhombognathides, 241
Rhombognathinae, 241
Rhombognathus, 241
Rhomboidella, 206
Rhombozoa, 104
Rhopalaea, 352
Rhopalomenia, 181
Rhopalomeniidae, 181
Rhopalonema, 119
Rhopalonematidae, 119
Rhopalophthalmus, 295
Rhopalura, 104
Rhopaluridae, 104
Rhopilema, 112
Rhumbleriella, 57
Rhumblerinella, 62
Rhyacodrilinae, 233
Rhynchmonas, 60
Rhynchobodo, 60
Rhynchobothrium, 148
Rhynchocinetidae, 288
Rhynchocoma, 46
Rhynchodida, 46
Rhynchogadus, 365
Rhynchogromia, 62
Rhynchohyalus, 361
Rhyncholagena, 272
Rhynchosmolidae, 258
Rhynchomyzon, 259
Rhynchonellata, 334
Rhynchonellida, 334
Rhynchonelliformea, 334
Rhynchonema, 172
Rhynchonereella, 214
Rhynchopharynx, 141
Rhynchopus, 60
Rhynchothalestris, 279
Rhynchothoracidae, 243
Rhynchothorax, 243
Rhynchozoon, 330
Rhytidodes, 142
Richardina, 286
Richtersia, 172
Ridgewayia, 267
Ridgewayiidae, 267
Ridleia, 89
Rigidomastix, 79
Rimicaris, 286
Rimostrombidium, 44
Ringicula, 199
Ringiculidae, 199
Rinoctes, 362
Rioselleolis, 205
Riseriellus, 154
Rissa, 375
Rissoa, 188
Rissoella, 37, 196
Rissoellaceae, 37
Rissoellidae, 196
Rissoidea, 188
Rissooides, 284
Rissoina, 190
Rissopsetia, 197
Ritterella, 353
Ritteriella, 356
Robertgurneyia, 273
Robertianella, 224
Robertina, 75
Robertinidae, 75
Robertinoides, 75
Robertsonia, 273
Robertsonites, 251
Roboastra, 203
Robphildolfusium, 142
Robustochelia, 314
Rochinia, 291
Rocinela, 319
Rodriguezella, 33
Rogneda, 132
Rokopella, 182
Rolandia, 204
Romancheinidae, 330
Rondeletia, 368
Rondeletiidae, 368
Rondeletiola, 212
Rosacea, 121
Rosalina, 71
Rosalinda, 113
Rosalindidae, 113
Rosalinidae, 75
Roscoffia, 52
Rosenbattichthys, 363
Rosenvingea, 28
Rosenvingiella, 23
Rosseliana, 330
Rossella, 103
Rossellidae, 103
Rossia, 212
Rossicotrema, 142
Rostanga, 203
Rostangidae, 203
Rostrocythere, 247
Rostroraja, 360

<i>Rotaliella</i> , 71	<i>Sagitta</i> , 335	<i>Scagelia</i> , 31
<i>Rotaliidae</i> , 75	<i>Sagittidae</i> , 335	<i>Scageliopsis</i> , 31
<i>Rotaliina</i> , 69	<i>Sagittiferidae</i> , 127	<i>Scagelothamnion</i> , 31
<i>Rotaria</i> , 151	<i>Sagmatocythere</i> , 250	<i>Scalibregma</i> , 226
<i>Rotifera</i> , 150	<i>Saharolana</i> , 320	<i>Scalibregmatidae</i> , 226
<i>Rotundiclipeidae</i> , 278	<i>Sahlingsia</i> , 35	<i>Scaliola</i> , 185
<i>Rotundiclipeus</i> , 278	<i>Sahnicythere</i> , 250	<i>Scaliolidae</i> , 185
<i>Rouleina</i> , 362	<i>Sahulina</i> , 65	<i>Scalpellidae</i> , 282
<i>Roundstonia</i> , 250	<i>Salacia</i> , 116	<i>Scalpellinae</i> , 282
<i>Roxania</i> , 199	<i>Salaria</i> , 372	<i>Scalpellomorpha</i> , 281
<i>Rozinante</i> , 299	<i>Saldidae</i> , 324	<i>Scalpellum</i> , 282
<i>Rudolphosetia</i> , 188	<i>Salenia</i> , 346	<i>Scambicornis</i> , 258
<i>Rugulina</i> , 184	<i>Saleniidae</i> , 346	<i>Scandia</i> , 115
<i>Rullierinereis</i> , 221	<i>Salenocidaris</i> , 346	<i>Scanorhynchus</i> , 132
<i>Runcina</i> , 199	<i>Salenoida</i> , 346	<i>Scaphander</i> , 200
<i>Runcinidae</i> , 199	<i>Salmacina</i> , 226	<i>Scaphocalanus</i> , 267
<i>Rupertina</i> , 73	<i>Salmincola</i> , 261	<i>Scaphodinium</i> , 52
<i>Rutiderma</i> , 246	<i>Salmo</i> , 362	<i>Scaphopoda</i> , 211
<i>Ruvettus</i> , 373	<i>Salmoneus</i> , 287	<i>Scaptognathus</i> , 241
<i>Ryocalanidae</i> , 267	<i>Salmonidae</i> , 362	<i>Scaptrella</i> , 172
<i>Ryocalanus</i> , 267	<i>Salmoniformes</i> , 362	<i>Scaridae</i> , 371
<i>Rythabis</i> , 268	<i>Salpa</i> , 356	<i>Scartella</i> , 372
<i>Rytiphloea</i> , 33	<i>Salpidae</i> , 355	<i>Sceptrintus</i> , 88
<i>Rzebakinidae</i> , 64	<i>Salpingoeca</i> , 58	<i>Schaudinnia</i> , 103
<i>Sabatieria</i> , 172	<i>Salpingoecidae</i> , 58	<i>Schedophilus</i> , 374
<i>Sabaudia</i> , 123	<i>Salvelinus</i> , 362	<i>Schendylidae</i> , 323
<i>Sabella</i> , 226	<i>Samus</i> , 87	<i>Scherochorella</i> , 64
<i>Sabellacheres</i> , 256	<i>Samytha</i> , 215	<i>Scherocumella</i> , 310
<i>Sabellaria</i> , 225	<i>Samythella</i> , 215	<i>Schikhobalotrema</i> , 142
<i>Sabellariidae</i> , 225	<i>Sapphirina</i> , 258	<i>Schilderia</i> , 191
<i>Sabellidae</i> , 225	<i>Sapphirinidae</i> , 258	<i>Schimmelmannia</i> , 35
<i>Sabellides</i> , 215	<i>Sarcacenaria</i> , 75	<i>Schistobrachia</i> , 261
<i>Sabelliphilidae</i> , 258	<i>Sarcodiaceae</i> , 37	<i>Schistocephalus</i> , 148
<i>Sabelliphilus</i> , 258	<i>Sarcodictyon</i> , 104	<i>Schistomerings</i> , 217
<i>Sabinea</i> , 288	<i>Sarcodiotheca</i> , 37	<i>Schistomyysis</i> , 295
<i>Sabinella</i> , 187	<i>Sarcotretes</i> , 262	<i>Schisturella</i> , 303
<i>Sabulodinium</i> , 52	<i>Sarda</i> , 373	<i>Schizammina</i> , 64
<i>Sabussowia</i> , 136	<i>Sardina</i> , 361	<i>Schizaster</i> , 347
<i>Saccaminidae</i> , 64	<i>Sardinella</i> , 361	<i>Schizasteridae</i> , 347
<i>Saccammina</i> , 65	<i>Sargassaceae</i> , 28	<i>Schizobrachiella</i> , 331
<i>Saccamoeba</i> , 53	<i>Sargassum</i> , 28	<i>Schizochilus</i> , 133
<i>Saccharomycetales</i> , 18	<i>Sargocentron</i> , 368	<i>Schizochytrium</i> , 78
<i>Saccocirridae</i> , 226	<i>Saroecka</i> , 58	<i>Schizocyathus</i> , 110
<i>Saccocirrus</i> , 226	<i>Sarostegia</i> , 102	<i>Schizomavella</i> , 326
<i>Saccocoelium</i> , 142	<i>Sarpa</i> , 370	<i>Schizopathes</i> , 109
<i>Saccodendron</i> , 65	<i>Sarsameira</i> , 269	<i>Schizopera</i> , 273
<i>Saccoglossus</i> , 336	<i>Sarsarietellus</i> , 264	<i>Schizoperopsis</i> , 273
<i>Sacconemertes</i> , 155	<i>Sarsia</i> , 113	<i>Schizoporella</i> , 331
<i>Saccopharyngidae</i> , 361	<i>Sarsicytheridea</i> , 247	<i>Schizoporellidae</i> , 331
<i>Saccopharyngiformes</i> , 361	<i>Sarsiella</i> , 113, 247	<i>Schizoproctus</i> , 253
<i>Saccopharynx</i> , 361	<i>Sarsiellidae</i> , 247	<i>Schizoretepora</i> , 330
<i>Saccopsidae</i> , 258	<i>Sarsiustra</i> , 329	<i>Schizorhamphus</i> , 310
<i>Saccorhiza</i> , 28, 63	<i>Sarsolectodes</i> , 268	<i>Schizorhynchidae</i> , 132
<i>Saccostrea</i> , 207	<i>Sarsonia</i> , 172	<i>Schizorhynchoides</i> , 133
<i>Sacculina</i> , 283	<i>Saturnius</i> , 142	<i>Schizorhynchus</i> , 133
<i>Sacculinidae</i> , 283	<i>Saundersella</i> , 25	<i>Schizotheca</i> , 330
<i>Sacodiscus</i> , 279	<i>Saurenchelys</i> , 361	<i>Schizothrix</i> , 276
<i>Sactosoma</i> , 178	<i>Saurida</i> , 363	<i>Schizotrema</i> , 310
<i>Saduria</i> , 321	<i>Sauvageaugloia</i> , 25	<i>Schizotricha</i> , 115
<i>Saduriella</i> , 321	<i>Saveljevia</i> , 172	<i>Schizymenia</i> , 37
<i>Saepicula</i> , 58	<i>Savignyella</i> , 331	<i>Schizymeniaceae</i> , 37
<i>Sagamichthys</i> , 362	<i>Savignyellidae</i> , 331	<i>Schleschia</i> , 200
<i>Sagamiella</i> , 277	<i>Savillea</i> , 58	<i>Schmitzia</i> , 36
<i>Sagartia</i> , 108	<i>Savoryella</i> , 18	<i>Schmitziella</i> , 29
<i>Sagartiidae</i> , 108	<i>Saxicavella</i> , 210	<i>Schottera</i> , 37
<i>Sagartiogeton</i> , 108	<i>Scacchia</i> , 208	<i>Schroederella</i> , 222
<i>Sagitella</i> , 231	<i>Scaeurgus</i> , 213	<i>Schulmania</i> , 83

- Sciadonus*, 366
Sciaena, 371
Sciaenidae, 370
Sciaenocotyle, 145
Sciophilus, 260
Scillaepas, 281
Scina, 308
Scinaia, 37
Scinidae, 308
Scissurella, 183
Scissurellidae, 183
Scleractinia, 109
Scleranthesia, 104
Sclerasterias, 342
Scleraulophoridae, 129
Scleraulophorus, 129
Sclerobregma, 226
Sclerocheilus, 226
Sclerochilus, 247
Sclerocrangon, 288
Sclerodistomum, 142
Sclerodoris, 202
Sclerolinidae, 235
Sclerolinum, 235
Scolanthus, 107
Scolaricia, 222
Scolecimorpha, 255
Scolecithricella, 267
Scolecithrix, 267
Scolecitrichidae, 267
Scolecobasidium, 19
Scolelepis, 228
Scolepis, 228
Scoletoma, 219
Scolex, 148
Scoliopharyngia, 133
Scoloplos, 222
Scomber, 373
Scomberesocidae, 367
Scomberesox, 367
Scomberomorus, 373
Scombridae, 373
Scombrolabracidae, 373
Scombrolabrax, 373
Scopalatum, 267
Scopalina, 90
Scopelarchidae, 363
Scopelarchus, 363
Scopelengys, 364
Scopeloberyx, 367
Scopelocheiropsis, 303
Scopelocheirus, 303
Scopelogadus, 367
Scopelosaurus, 363
Scophthalmidae, 374
Scophthalmus, 374
Scopulariopsis, 19
Scorpaena, 368
Scorpaenidae, 368
Scorpaeniformes, 368
Scorpaenodes, 368
Scotothuria, 349
Scottocalanus, 268
Scottocheres, 259
Scottolana, 271
Scottomyzon, 259
Scotopsyllus, 277
Scrippsiella, 52
Scripsiella, 52
Scrobicularia, 209
Scrobiculariidae, 209
Scruparia, 331
Scrupariidae, 331
Scrupocellaria, 328
Scupocellaria, 328
Scutellidium, 279
Scutopus, 181
Scyliorhinidae, 358
Scyliorhinus, 359
Scyllaea, 204
Scyllaeidae, 204
Scyllaridae, 289
Scyllarides, 289
Scyllarus, 289
Scymnodalatias, 359
Scymnodon, 359
Scyphidium, 103
Scyphopodium, 104
Scyphozoa, 111
Scyracepon, 322
Scytosiphon, 28
Scytosiphonaceae, 28
Scytosiphonales, 28
Searsia, 362
Seba, 306
Sebastapistes, 368
Sebastidae, 368
Sebdenia, 37
Sebdeniaceae, 37
Sebidae, 306
Securiflustra, 329
Segmentochona, 41
Segmentochonidae, 41
Segonzacia, 292
Segonzactis, 107
Seguenzia, 184
Seguenziidae, 184
Seila, 186
Seirospora, 32
Seison, 151
Seisonidae, 151
Seisonidea, 151
Selaginopsis, 116
Selar, 370
Selene, 370
Selenopsyllus, 271
Selimia, 130
Selioides, 257
Selius, 257
Semaeostomeae, 111
Semele, 209
Semelidae, 209
Semibalaninae, 282
Semibalanus, 282
Semicytherura, 248
Semivermilia, 226
Semnoderes, 160
Semnoderidae, 160
Sepia, 212
Sepietta, 212
Sepiidae, 212
Sepiola, 212
Sepiolidae, 212
Sergestes, 286
Sergestidae, 286
Sergia, 130, 286
Seriata, 135
Sericosura, 242
Seriocarpa, 355
Seriola, 370
Seritia, 130
Serolidae, 320
Serolidae i.s., 320
Serpula, 226
Serpulidae, 226
Serpulidicola, 258
Serpulidicolidae, 258
Serpulorbis, 190
Serranicotyle, 145
Serranidae, 369
Serranus, 369
Serrasentis, 177
Serratorhynchus, 133
Serratosagitta, 335
Serripes, 209
Serrivomer, 361
Serrivomeridae, 361
Sertularella, 116
Sertularia, 116
Sertulariidae, 116
Sertulipora, 330
Sessilia, 282
Sestropontius, 259
Setarches, 368
Setarchidae, 368
Setia, 188
Setoplectus, 172
Setosabatieria, 172
Setosella, 331
Setosellidae, 331
Setosellina, 329
Sextonia, 301
Shepareddella, 62
Sibitrema, 145
Siboglinidae, 235
Siboglinum, 235
Sicameira, 269
Sicyodelphys, 255
Sicyonia, 286
Sicyoniidae, 286
Sicyonis, 106
Sideractiidae, 108
Sideractis, 108
Sididae, 244
Sidneioides, 353
Sigalion, 227
Sigalionidae, 227
Sigambla, 223
Siganidae, 373
Siganus, 373
Sige, 223
Sigeheres, 257
Sigmatidium, 274
Sigmavirgulina, 73
Sigmoidea, 19
Sigmoidella, 75
Sigmoihauerina, 66
Sigmoilina, 68
Sigmoilinita, 68
Sigmoilopsis, 68

<i>Sigmophoranema</i> , 172	Smittinidae , 331	Spatangidae , 347
<i>Silhouettea</i> , 373	<i>Smittipora</i> , 330	Spatangoidea , 347
Siliquariidae , 185	<i>Smittoidea</i> , 331	<i>Spatangus</i> , 347
Sillaginidae , 373	<i>Sobolevianthus</i> , 148	<i>Spathipora</i> , 332
<i>Sillago</i> , 373	<i>Socarnes</i> , 303	Spathiporidae , 332
<i>Simenchelys</i> , 360	<i>Soestia</i> , 356	<i>Spathoderma</i> , 181
<i>Similiphora</i> , 185	<i>Sognocalanus</i> , 264	<i>Spathoteredo</i> , 211
<i>Simnia</i> , 191	<i>Solariella</i> , 184	<i>Spatoglossum</i> , 26
Simognathinae , 241	<i>Solaster</i> , 341	<i>Spatulodinium</i> , 52
<i>Simognathus</i> , 241	Solasteridae , 341	<i>Spectacturus</i> , 321
<i>Simplaria</i> , 228	<i>Solatia</i> , 194	<i>Spectrunculus</i> , 366
Simplicomorpha , 126	<i>Solatisonax</i> , 196	<i>Speleogobius</i> , 373
<i>Simrothiella</i> , 181	<i>Solea</i> , 374	<i>Speleonectes</i> , 243
Simrothiellidae , 181	<i>Solearhynchus</i> , 177	Speleonectidae , 243
<i>Simsagittifera</i> , 127	<i>Solecardia</i> , 208	<i>Speleophria</i> , 262
<i>Sinantenna</i> , 224	Solecurtidae , 210	Speleophriidae , 262
Sinelobinae , 313	<i>Solecurtus</i> , 210	<i>Speleophriopsis</i> , 263
<i>Sinezona</i> , 183	Soleidae , 374	<i>Spelotrema</i> , 142
<i>Siniotrochus</i> , 351	<i>Solemya</i> , 205	Spengelidae , 336
<i>Sinophysis</i> , 52	Solemyidae , 205	Spermatochnaceae , 25
<i>Sinum</i> , 191	<i>Solen</i> , 209	<i>Spermatochmus</i> , 25
<i>Sinuolinea</i> , 83	Solenidae , 209	<i>Spermosyllis</i> , 229
Sinuolineidae , 83	<i>Solenocera</i> , 286	<i>Spermothamnion</i> , 32
<i>Sinupharus</i> , 209	Solenoceridae , 286	<i>Sperosoma</i> , 346
Siphogenerinoidiae , 75	<i>Solenofilomorpha</i> , 127	<i>Sphacelaria</i> , 28
<i>Sipholagena</i> , 74	Solenolaimus , 172	Sphaelariaceae , 28
<i>Siphonactinopsis</i> , 107	Solenopharyngidae , 133	Sphaelariales , 28
<i>Siphonaptera</i> , 68	<i>Solenopharynx</i> , 134	<i>Sphaella</i> , 29
<i>Siphonaria</i> , 205	<i>Solenosmilia</i> , 109	Sphaerasteridae , 341
Siphonariidae , 205	<i>Soletellina</i> , 209	<i>Sphaerechinus</i> , 346
Siphonidiidae , 87	<i>Solidobalanus</i> , 282	<i>Sphaerifer</i> , 257
<i>Siphonidium</i> , 87	<i>Solidosagatta</i> , 335	<i>Sphaeriodiscus</i> , 340
<i>Siphonina</i> , 75	<i>Solieria</i> , 37	<i>Sphaerobothrium</i> , 148
Siphoninidae , 75	Solieriae , 37	<i>Sphaerocephalum</i> , 172
<i>Siphonochalina</i> , 98	<i>Solitaria</i> , 179	<i>Sphaerococcaceae</i> , 38
Siphonocladaeae , 23	<i>Solmaris</i> , 119	<i>Sphaerococcus</i> , 38
Siphonocladales , 23	Solmarisidae , 119	<i>Sphaerocoryne</i> , 113
<i>Siphonocladus</i> , 23	<i>Solmissus</i> , 119	Sphaerocorynidiae , 113
<i>Siphonodentalium</i> , 212	<i>Solmundella</i> , 119	Sphaerodoridae , 227
<i>Siphonoecetes</i> , 298	<i>Solostamenides</i> , 145	<i>Sphaerodoridium</i> , 227
<i>Siphonohydra</i> , 113	<i>Somateria</i> , 375	<i>Sphaerodoropsis</i> , 227
<i>Siphonolaimus</i> , 172	<i>Sommiosus</i> , 359	<i>Sphaerodororum</i> , 227
Siphonophora , 121	<i>Sopharynx</i> , 134	<i>Sphaerogypsina</i> , 69
<i>Siphonosoma</i> , 179	<i>Sophrosyne</i> , 303	<i>Sphaeroides</i> , 17
Siphonostomatoidea , 259	<i>Sorapion</i> , 27	<i>Sphaerolaimina</i> , 172
<i>Siphotextularia</i> , 65	<i>Sorbera</i> , 355	<i>Sphaeroma</i> , 321
<i>Sipuncula</i> , 178	Sorberacea , 355	Sphaeromatidae , 320
Sipunculidae , 179	<i>Sordaria</i> , 18	<i>Sphaeromyxa</i> , 82
<i>Sipunculus</i> , 179	Sordariaceae , 18	Sphaeromyxidae , 82
<i>Siriella</i> , 295	<i>Sordariales</i> , 18	Sphaeromyxina , 82
<i>Sirpus</i> , 291	<i>Sorites</i> , 69	Sphaeronectes , 122
<i>Situla</i> , 352	Soritidae , 69	Sphaeronectidae , 122
<i>Skeizia</i> , 90	<i>Sorocarpaceae</i> , 27	<i>Sphaeronella</i> , 261
<i>Skenea</i> , 184	<i>Sorosphaera</i> , 65	<i>Sphaerospora</i> , 83
Skeneidae , 184	<i>Sosane</i> , 215	Sphaerosporidae , 83
<i>Skeneoides</i> , 185	<i>Sosanopsis</i> , 215	<i>Sphaerostoma</i> , 142
Skeneopsidae , 188	<i>Southerniella</i> , 172	<i>Sphaerosyllis</i> , 229
<i>Skeneopsis</i> , 188	<i>Southwellina</i> , 177	<i>Sphaerotrichia</i> , 25
<i>Skogsbergia</i> , 245	<i>Spadella</i> , 335	<i>Sphaerotylus</i> , 89
<i>Skrjabinalius</i> , 176	Spadellidae , 335	<i>Sphaerozius</i> , 292
<i>Skrjabinolecithum</i> , 142	<i>Spanioplton</i> , 96	<i>Sphaerulina</i> , 16
<i>Skrjabinopsolus</i> , 142	<i>Sparicotyle</i> , 145	<i>Sphagmacrurus</i> , 365
<i>Smaragdia</i> , 183	Sparidae , 370	<i>Sphairogullmia</i> , 62
<i>Smaragdinella</i> , 199	<i>Sparidicola</i> , 261	<i>Sphenia</i> , 210
<i>Smilium</i> , 281	<i>Sparisoma</i> , 371	<i>Sphenophrya</i> , 47
<i>Sminthea</i> , 119	<i>Sparus</i> , 370	
<i>Smittina</i> , 331		

- Sphenophryidae**, 46
Sphenotrochus, 110
Sphinctrella, 86
Sphinter, 227
Sphaeroides, 374
Sphondylothamnion, 32
Sphyraena, 373
Sphyraenidae, 373
Sphyrapidae, 313
Sphyrapinae, 313
Sphyrapoides, 313
Sphyrapus, 313
Sphyriidae, 262
Sphyriocephalus, 148
Sphyrion, 262
Sphyrna, 359
Sphyrnidae, 359
Spicara, 370
Spicipes, 265
Spiliphera, 172
Spilophorella, 173
Spinachia, 368
Spinimuricea, 105
Spinitectoides, 176
Spinictectus, 176
Spinocalanidae, 268
Spinocalanus, 268
Spinoplagioporid, 142
Spinther, 227
Spintheridae, 227
Spinularia, 89
Spinulidae, 206
Spinulosida, 341
Spio, 228
Spiochaetopterus, 216
Spionidae, 227
Spiophanes, 228
Spiophanicola, 258
Spiophanicolidae, 258
Spiramphinema, 173
Spiraserpula, 226
Spirastrella, 88
Spirastrellidae, 88
Spiraulax, 52
Spiricella, 201
Spiridion, 233
Spirillina, 75
Spirillinidae, 75
Spirillinoides, 62
Spirinia, 173
Spirobranchus, 227
Spirodiscus, 227
Spirolaxis, 196
Spirolina, 69
Spiroloculina, 68
Spirontocaris, 287
Spiropagurus, 289
Spirophorida, 87
Spirophthalmidium, 68
Spiroplectammina, 65
Spiroplectamminidae, 65
Spiroplectinella, 65
Spirorbidae, 228
Spirorbis, 228
Spirorutilus, 65
Spilosigmoilina, 68
Spirostrombidium, 43
Spirotropis, 194
Spiroxys, 88
Spirula, 212
Spirulidae, 212
Spisula, 209
Splanchnotropidae, 258
Splanchnotropus, 258
Splitofolliculina, 42
Splitoiculina, 42
Spondylidae, 207
Spondyliosoma, 370
Spondylus, 207
Spongia, 100
Spongicolidae, 286
Spongicoloides, 286
Spongidiidae, 100
Sponginticola, 262
Sponginticolidae, 262
Spongicnizon, 262
Spongicnizontidae, 262
Spongionella, 101
Spongites, 34
Spongomorpha, 21
Spongonema, 27
Spongisorites, 91
Sporidesmium, 19
Sporochnaceae, 29
Sporochnales, 29
Sporochnus, 29
Sporolithaceae, 34
Sporolithon, 34
Sporomiaceae, 16
Sportella, 208
Sportellidae, 208
Spratelloides, 361
Sprattus, 361
Spurilla, 204
Spyridia, 32
Squalidae, 359
Squaliformes, 359
Squaliolus, 359
Squalonchocytyle, 145
Squalus, 359
Squamatoherpia, 181
Squatina, 359
Squatiniidae, 359
Squatiformes, 359
Squilla, 284
Squillidae, 284
Stainforthia, 70
Staphylinidae, 323
Staurocladia, 113
Staurodiscus, 115
Stauromedusae, 112
Staurosoma, 255
Staurostoma, 115
Steganoderma, 142
Steganodermatoidea, 142
Stegocephalidae, 306
Stegocephalina, 307
Stegocephaloidea, 307
Stegocephalopsis, 307
Stegocephalus, 307
Stegohornera, 333
Stegohorneridae, 333
Stegolaria, 116
Stegophiura, 345
Stegoplax, 297
Stegopoma, 116
Stegosoma, 356
Stegotricha, 46
Steineria, 173
Steineridora, 173
Stellanthura, 316
Stellella, 86
Stellicola, 256
Stelligera, 88
Stemphylium, 19
Stenakron, 142
Stenella, 376
Stenetriidae, 319
Stenetrium, 319
Stenelia, 273
Steno, 376
Stenocaris, 271
Stenocaropsis, 271
Stenocopia, 269
Stenocyathus, 110
Stenodus, 362
Stenogramme, 37
Stenohelia, 118
Stenolaemata, 333
Stenopleura, 299
Stenopleustes, 306
Stenopodidae, 286
Stenopus, 286
Stenorhynchus, 291
Stenostomidae, 127
Stenostomum, 127
Stenotanais, 314
Stenothocheres, 261
Stenothee, 307
Stenothoidae, 307
Stenofolliculina, 42
Stenula, 307
Stenurus, 176
Stephalia, 122
Stephanacantha, 58
Stephanasterias, 342
Stephanauge, 108
Stephanoberycidae, 367
Stephanoberyciformes, 367
Stephanocyathus, 109
Stephanoea, 58
Stephanolaimus, 173
Stephanolepis, 374
Stephanollona, 330
Stephanophyes, 122
Stephanostomum, 142
Stephidae, 268
Stephobruzelia, 307
Stephonyx, 303
Stephos, 268
Sterechmella, 330
Stercorariidae, 375
Stercorarius, 376
Stereales, 18
Stereobalanus, 336
Stereocidaris, 346
Stereoderma, 349
Stereomastis, 288
Stereomyxa, 54
Stereomyxidae, 54
Steringophorus, 142

<i>Steringotrema</i> , 142	<i>Strobilidiidae</i> , 44	<i>Stylocaulaceae</i> , 29
<i>Sterna</i> , 375	<i>Strobilidium</i> , 44	<i>Stylocaulon</i> , 29
<i>Sternaspidae</i> , 228	<i>Strobiligera</i> , 185	<i>Stylopodium</i> , 26
<i>Sternaspis</i> , 228	<i>Strobilocephalus</i> , 148	<i>Styracaster</i> , 339
<i>Sternoptychidae</i> , 362	<i>Strobopagurus</i> , 289	<i>Styraconyx</i> , 236
<i>Sternoptyx</i> , 362	<i>Stromateidae</i> , 374	<i>Subadyte</i> , 224
<i>Stetsonia</i> , 70	<i>Stromatella</i> , 22	<i>Suberites</i> , 89
<i>Sthenelais</i> , 227	<i>Stromateus</i> , 374	<i>Suberitidae</i> , 88
<i>Sthenoteuthis</i> , 213	<i>Strombidae</i> , 190	<i>Subeucalanus</i> , 265
<i>Stichaeidae</i> , 371	<i>Strombidiidae</i> , 43	<i>Subreophax</i> , 64
<i>Stichastrella</i> , 342	<i>Strombidinopsisidae</i> , 44	<i>Subulagera</i> , 133
<i>Stichococcaceae</i> , 23	<i>Strombidinopsis</i> , 44	<i>Subulella</i> , 314
<i>Stichococcus</i> , 23	<i>Strombidium</i> , 43	<i>Sudis</i> , 364
<i>Stichocotyle</i> , 136	<i>Strombus</i> , 190	<i>Suzukiella</i> , 57
<i>Stichocotylida</i> , 136	<i>Strongylacron</i> , 271	<i>Sula</i> , 375
<i>Stichopathes</i> , 109	<i>Strongylocentrotidae</i> , 347	<i>Sulculeolaria</i> , 121
<i>Stichopodidae</i> , 349	<i>Strongylocentrotus</i> , 347	<i>Sulidae</i> , 375
<i>Stichothamnion</i> , 33	<i>Strongylurella</i> , 314	<i>Sunamphitoe</i> , 297
<i>Stictelima</i> , 187	<i>Strophomenia</i> , 182	<i>Sunaristes</i> , 271
<i>Stictodora</i> , 142	<i>Strophomeniidae</i> , 182	<i>Superornatiremidae</i> , 278
<i>Stictyosiphon</i> , 26	<i>Struvea</i> , 23	<i>Superornatiremis</i> , 278
<i>Stiliger</i> , 201	<i>Stryphnus</i> , 86	<i>Svetovidovia</i> , 365
<i>Stiligeridae</i> , 201	<i>Styela</i> , 355	<i>Svratkina</i> , 69
<i>Stilomysis</i> , 295	<i>Styelicola</i> , 253	<i>Swiftia</i> , 105
<i>Stilophora</i> , 25	<i>Styelidae</i> , 354	<i>Sycandra</i> , 102
<i>Stilopsis</i> , 25	<i>Stygamoeba</i> , 54	<i>Sycetta</i> , 102
<i>Stilostomella</i> , 70	<i>Stygarctidae</i> , 236	<i>Sycettidae</i> , 102
<i>Stoecharthrum</i> , 104	<i>Stygarctus</i> , 236	<i>Sycodorus</i> , 102
<i>Stolidobranchia</i> , 354	<i>Stygiomedusa</i> , 111	<i>Sycon</i> , 102
<i>Stolonica</i> , 355	<i>Stygiopontius</i> , 260	<i>Sycozoa</i> , 353
<i>Stoloteuthis</i> , 212	<i>Stylocapitella</i> , 222	<i>Sykyssia</i> , 102
<i>Stomachetosella</i> , 331	<i>Stygocyclopia</i> , 267	<i>Sykidion</i> , 22
<i>Stomachetosellidae</i> , 331	<i>Stygoletta</i> , 128	<i>Syllidae</i> , 228
<i>Stomatolepas</i> , 282	<i>Stygotanellina</i> , 134	<i>Syllides</i> , 230
<i>Stomatopoda</i> , 284	<i>Stygotantulus</i> , 280	<i>Syllidia</i> , 219
<i>Stomatopora</i> , 333	<i>Stylactis</i> , 118	<i>Syllis</i> , 230
<i>Stomatoporina</i> , 333	<i>Stylaria</i> , 233	<i>Sylon</i> , 283
<i>Stomatorbina</i> , 73	<i>Stylaster</i> , 118	<i>Syltoidinium</i> , 52
<i>Stomias</i> , 363	<i>Stylasteridae</i> , 118	<i>Syltorhynchus</i> , 132
<i>Stomiidae</i> , 363	<i>Stylatula</i> , 106	<i>Symbibidae</i> , 177
<i>Stomiiformes</i> , 362	<i>Stylephoridae</i> , 364	<i>Symbiida</i> , 177
<i>Stomotoca</i> , 118	<i>Stylephorus</i> , 364	<i>Symbiodinium</i> , 52
<i>Stomphia</i> , 106	<i>Stylicletodes</i> , 271	<i>Symbion</i> , 177
<i>Storthosphaera</i> , 65	<i>Stylinos</i> , 91	<i>Symbolophorus</i> , 364
<i>Storthynghura</i> , 317	<i>Styliola</i> , 200	<i>Sympagella</i> , 103
<i>Strabax</i> , 255	<i>Stylissa</i> , 90	<i>Sympagurus</i> , 289
<i>Strangularia</i> , 25	<i>Stylocheiron</i> , 284	<i>Sympodus</i> , 371
<i>Stramenopiles incertae sedis</i> , 78	<i>Stylochidae</i> , 128	<i>Sympurus</i> , 374
<i>Stramonita</i> , 192	<i>Stylochona</i> , 41	<i>Sympycarpus</i> , 28
<i>Stratiopontotes</i> , 271	<i>Stylochonidae</i> , 41	<i>Sympyocladia</i> , 33
<i>Strauchia</i> , 310	<i>Stylochoplana</i> , 129	<i>Symplectoscyphus</i> , 116
<i>Streblocladia</i> , 33	<i>Stylochoplaniidae</i> , 128	<i>Symplocostoma</i> , 173
<i>Strebloides</i> , 71	<i>Stylochus</i> , 128	<i>Synagoga</i> , 283
<i>Streblonema</i> , 27	<i>Stylocidaridae</i> , 346	<i>Synagogidae</i> , 283
<i>Streblonemopsis</i> , 27	<i>Stylocordyla</i> , 89	<i>Synallactes</i> , 349
<i>Streblosoma</i> , 231	<i>Stylocordylidae</i> , 89	<i>Synallactidae</i> , 349
<i>Streblospio</i> , 228	<i>Stylocoronella</i> , 112	<i>Synalpheus</i> , 288
<i>Strepsithalia</i> , 25	<i>Styloceratidae</i> , 288	<i>Synamoeba</i> , 54
<i>Streptocara</i> , 176	<i>Stylocordylus</i> , 288	<i>Synaphobranchidae</i> , 360
<i>Streptocaulus</i> , 114	<i>Stylocladina</i> , 52	<i>Synaphobranchus</i> , 360
<i>Streptoleberis</i> , 246	<i>Stylocladina</i> , 181	<i>Synapseudes</i> , 313
<i>Streptosyllis</i> , 230	<i>Stylocladina</i> , 37	<i>Synapseudinae</i> , 313
<i>Striarca</i> , 206	<i>Stylopandalus</i> , 287	<i>Synapticolidae</i> , 258
<i>Striaria</i> , 26	<i>Stylopiton</i> , 94	<i>Synaptidae</i> , 350
<i>Striariaceae</i> , 26	<i>Styloptocuma</i> , 310	<i>Synaptiphilidae</i> , 258
<i>Strigamia</i> , 323	<i>Styloptygma</i> , 197	<i>Synaptiphilus</i> , 258
<i>Strigilla</i> , 209	<i>Stylostomum</i> , 128	<i>Synaptobothrium</i> , 142

- Synaptula*, 351
Synaptila, 374
Synasterope, 245
Synchaeta, 150
Synchaetidae, 150
Synchelidium, 305
Synchiropus, 372
Syncoryne, 22
Syndesmis, 130
Syndisyrinx, 130
Syneelmis, 223
Syneuroscope, 317
Syngastes, 278
Syngnathidae, 368
Syngnathiformes, 368
Syngnathiiformes, 368
Syngnathus, 368
Synidotea, 321
Synisoma, 321
Synnotum, 329
Synodontidae, 363
Synodontium, 173
Synodus, 363
Synoicum, 353
Synonchiella, 173
Synonchium, 173
Synonchus, 173
Synonema, 173
Synopiidae, 307
Syntheciidae, 116
Synthecium, 116
Syringammina, 79
Syringamminidae, 78
Syringoderma, 29
Syringodermataceae, 29
Syringodermatales, 29
Syringolaimus, 173
Syringonomus, 173
Syringopharynx, 46
Syrnola, 197
Syrrhoe, 307
Syrrhoites, 307
Syrticola, 276
Syscenus, 319
Systellaspis, 287
Taaningichthys, 364
Tachidiella, 279
Tachidiidae, 278
Tachidiopsis, 279
Tachidius, 278
Tachyrhynchus, 185
Tachytrypane, 222
Taenia, 148
Taeniacanthidae, 258
Taeniacanthus, 258
Taenioma, 32
Taeniura, 360
Talacalanus, 266
Talaromyces, 16
Talassia, 190
Talismania, 362
Talitridae, 307
Talitrus, 308
Talorchestia, 308
Talpina, 275
Tamarisca, 116
Tambja, 203
Tanabnormia, 314
Tanaella, 314
Tanaidacea, 313
Tanaidae, 313
Tanaidomorpha, 313
Tanainae, 313
Tanais, 313
Tanaisia, 142
Tanaissus, 315
Tanaopsis, 315
Tanarctus, 236
Tanganella, 332
Taningia, 212
Tantulacus, 280
Tantulocarida, 280
Tanystylum, 242
Taonia, 27
Taonius, 213
Tapes, 210
Taractes, 370
Taractichthys, 370
Taranis, 196
Tarasovium, 282
Tardigrada, 236
Tariscola, 258
Taringa, 203
Tarrhomos, 154
Tarvaia, 173
Tatria, 148
Taurida, 127
Taurididae, 127
Taurulus, 369
Tealidium, 106
Technitella, 65
Teclaia, 115
Tectidrilus, 233
Tectillaria, 356
Tectonatica, 191
Tectura, 182
Tedania, 93
Tedaniidae, 93
Tegastes, 278
Tegastidae, 278
Tegella, 328
Tegulaherpia, 181
Teinostoma, 190
Teissierella, 273
Telammina, 65
Teleaulax, 59
Telestula, 104
Tellamia, 22
Tellervotrema, 142
Tellina, 209
Tellinidae, 209
Telmatactis, 108
Telmatogeton, 324
Telonema, 79
Telopsammis, 278
Telorchis, 142
Teloscalpellum, 282
Telosentis, 177
Temnopleuridae, 346
Temnopleuroidea, 346
Temora, 268
Temoridae, 268
Temorites, 264
Temoropia, 268
Tenagodus, 185
Tenarea, 34
Tenedocythere, 249
Tenellia, 205
Teneriforma, 268
Tenerrhynchus, 132
Tensopharynx, 134
Tentacularia, 148
Tentorium, 89
Tenuilineus, 154
Terapon, 369
Terapontidae, 369
Teraterythrops, 295
Terebella, 231
Terebellidae, 230
Terebellides, 231
Terebratulida, 334
Terebratulidae, 334
Terebratulina, 334
Terebriporidae, 332
Teredinidae, 210
Teredo, 210
Teredoika, 258
Teredora, 210
Teredothrya, 211
Teretia, 196
Teretiopsis, 196
Tergestia, 142
Tergipedidae, 205
Tergipes, 205
Terminoflustra, 329
Terpios, 89
Terranova, 176
Terschellingia, 173
Tertia, 333
Terviidae, 333
Tessaradoma, 331
Tessaradomidae, 331
Tesserogastria, 119
Tesseropora, 282
Testudinella, 151
Testudinellidae, 150
Testudines, 375
Testulosiphon, 63
Tetanopsis, 274
Tethocyathus, 109
Tethya, 88
Tethyaster, 339
Tethydidae, 204
Tethyidae, 88
Tethylembos, 298
Tethys, 204
Tethysbaena, 293
Tethyspira, 90
Tetilla, 87
Tetillidae, 87
Tetrabothrius, 148
Tetracilia, 79
Tetracladium, 19, 142
Tetraclitidae, 282
Tetraclitinae, 282
Tetraclitoidea, 282
Tetractinomorpha, 85
Tetracytherura, 248
Tetragoniceps, 279
Tetragonicipitidae, 278
Tetragonuridae, 374

- Tetragonurus*, 374
Tetrameres, 176
Tetramitus, 54
Tetranchyroderma, 158
Tetranemertes, 154
Tetraodontidae, 374
Tetraodontiformes, 374
Tetraonchoides, 145
Tetraonchus, 145
Tetraplatia, 112
Tetraplatidae, 112
Tetrapoma, 114
Tetrapturus, 373
Tetrahyynchobothrium, 148
Tetrahyynchus, 148
Tetraselmis, 76
Tetrasporales, 23
Tetrastemma, 155
Tetrasystemmatidae, 155
Tetrochetus, 142
Tetrorchis, 120
Teuchopora, 331
Teuchoporidae, 331
Teuthowenia, 213
Textularia, 65
Textulariidae, 65
Textulariina, 62
Thaerocythere, 249
Thais, 192
Thalamita, 291
Thalamophaga, 62
Thalamophyllia, 109
Thalassarachna, 241
Thallasema, 178
Thalassematidae, 178
Thalassironus, 173
Thalassisobates, 323
Thalassoalaimus, 173
Thalassoascus, 15
Thalassobathia, 366
Thalassochaetus, 221
Thalassodrilidae, 233
Thalassodrilus, 233
Thalassoma, 371
Thalassometra, 338
Thalassometridae, 338
Thalassomonhystera, 173
Thalassomyia, 324
Thalassomysis, 295
Thalassomyxa, 54
Thalassoplarella, 135
Thalassoplana, 135
Thalassopterus, 200
Thalassosmittia, 324
Thalestridae, 279
Thalestris, 279
Thalia, 356
Thaliacea, 355
Thambema, 319
Thambematidae, 319
Thamnostoma, 117
Tharybidae, 268
Tharybis, 268
Tharyx, 217
Thaumastoderma, 158
Thaumastodermatidae, 157
Thaumastosoma, 317
Thaumaticthyidae, 367
Thaumatocotyle, 145
Thaumatocrinus, 338
Thaumatomastix, 78
Thaumatometra, 338
Thaumatomonads, 78
Thaumatopsyllidae, 255
Thaumatosyllus, 255
Thecacera, 203
Thecadinium, 52
Thecamoeba, 54
Thecamoebidae, 54
Thecanephria, 235
Thecocodium, 118
Thelepus, 231
Thelohanellus, 84
Themisto, 308
Thenea, 87
Theneidae, 87
Theonellidae, 87
Theragra, 365
Theristus, 173
Thermosbaenacea, 293
Therochaeta, 218
Thersitina, 256
Thesbia, 196
Thesea, 105
Theta, 196
Thetys, 356
Thia, 291
Thielavia, 18
Thiidae, 291
Thliptodon, 200
Thloarctus, 236
Tholosina, 65
Thompsonopia, 267
Thompsonula, 279
Thompsonulidae, 279
Thoosa, 88
Thor, 287
Thoracaster, 339
Thoracica, 281
Thoracosphaera, 52
Thoracostoma, 173
Thoracostomopsis, 174
Thoralus, 287
Thordisa, 203
Thorectandra, 100
Thorectidae, 100
Thoriella, 303
Thorogobius, 373
Thorybopus, 202
Thouarella, 105
Thracia, 211
Thraciidae, 211
Thraustochytrids, 78
Thraustochytrium, 78
Thrombidae, 87
Thrombus, 87
Thuiaria, 116
Thulinia, 142
Thunnus, 373
Thurammina, 65
Thuretella, 35
Thuridilla, 200
Thyasira, 208
Thyasiridae, 208
Thylacorhynchus, 133
Thymosia, 89
Thymosiopsis, 89
Thyone, 349
Thyonidium, 348
Thysanocardia, 178
Thysanoessa, 284
Thysanopoda, 284
Thysanote, 261
Thysanoteuthidae, 213
Thysanoteuthis, 213
Thysanozoon, 128
Tiaranna, 117
Tiarannidae, 116
Tiaropsidae, 117
Tiaropsidium, 116
Tiaropsis, 117
Tiberia, 197
Tibitin, 355
Tiffaniella, 32
Tigriopus, 274
Tillitaceae, 18
Tilopteridaceae, 29
Tilopteridales, 29
Tilopteris, 29
Tima, 114
Timarete, 217
Timea, 88
Timeidae, 88
Timoclea, 210
Timonia, 142
Timoniella, 142
Tindaria, 206
Tindariidae, 206
Tinerfe, 123
Tinogullmia, 62
Tiphotrecha, 66
Tirispora, 17
Tiron, 307
Tisbe, 279
Tisbidae, 279
Tisemus, 279
Titanoderma, 34
Tjaernoeia, 198
Tjaernoiidae, 198
Tmetonyx, 303
Todarodes, 213
Todaropsis, 213
Toia, 131
Tole, 318
Toledonia, 199
Tolypamma, 62
Tolypocladium, 19
Tomlinula, 198
Tomopteridae, 231
Tomopteris, 231
Tomura, 197
Tonicella, 182
Tonna, 191
Tonnidae, 191
Tontonia, 44
Topsentia, 91
Torellia, 191
Torkarlingia, 131
Tormopslus, 142
Tornidae, 190
Tornus, 190

- Torodinium*, 53
Torpedinidae, 359
Torpediniformes, 359
Torpedo, 359
Torpedospora, 17
Torreya, 214
Tortugaster, 283
Torwolia, 317
Toxopneustidae, 346
Toxorhysis, 115
Tracheleuglypha, 56
Trachelobdella, 235
Trachelosaccus, 283
Trachichthyidae, 367
Trachinidae, 372
Trachinotus, 370
Trachinus, 372
Trachipteridae, 364
Trachipterus, 364
Trachonurus, 365
Trachurus, 370
Trachycaris, 287
Trachyleberididae, 250
Trachymedusae, 119
Trachynema, 26
Trachyrincus, 365
Trachysalambria, 286
Trachyscorbia, 368
Trailia, 17
Trapania, 202
Trapezidae, 210
Trapeziidae, 292
Trapezium, 210
Traustedtia, 356
Travisia, 222
Travisiopsis, 231
Trebiiidae, 262
Trebius, 262
Trefusia, 174
Tregoubovia, 118
Tremaster, 341
Trematcia, 16
Tremoctopodidae, 213
Tremoctopus, 213
Treptoplax, 103
Tretodictyidae, 103
Tretodictyum, 103
Tretomphalus, 71
Triaenophorus, 148
Triakidae, 359
Trianguloscalpellum, 282
Triangulus, 283
Tribonema, 24
Tribonemataceae, 24
Tricellaria, 328
Tricharrhen, 142
Trichasterina, 103
Trichethmolaimus, 174
Tricheurypon, 92
Trichiuridae, 373
Trichiurus, 373
Trichobilharzia, 142
Trichobranchidae, 231
Trichobranchus, 231
Trichocerca, 150
Trichocercidae, 150
Trichocladium, 19
Trichocomaceae, 16
Trichoderma, 19
Trichogloeopsis, 37
Trichogypsia, 102
Trichohyalidae, 75
Tricholomataceae, 18
Trichomastix, 79
Trichometra, 338
Trichoplacidae, 103
Trichoplax, 103
Trichosida, 55
Trichosidae, 55
Trichosolen, 21
Trichosphaerium, 54, 55
Trichostemma, 89
Trichothecium, 19
Trichotheristus, 174
Trichotria, 150
Trichotropis, 191
Trichydra, 118
Trichydriidae, 118
Tricladida, 136
Tricleocarpa, 37
Tricolia, 185
Tricolliidae, 185
Tricoma, 174
Triconia, 257
Tricotyla, 145
Trididemnum, 353
Triebelina, 247
Trifarina, 75
Triflagellum, 79
Triforis, 186
Trigla, 368
Triglidae, 368
Triglops, 369
Triglopsis, 369
Trigonaspis, 76
Trigonocidaris, 346
Trigonocotyle, 149
Trigonolampa, 363
Trigonoporus, 128
Trigonopyxidae, 56
Trigonostomidae, 134
Trigonostomum, 134
Trileptium, 174
Trilobodrilus, 217
Trilocularia, 149
Triloculina, 68
Trilospora, 84
Trilosporidae, 84
Trimastix, 79
Trimusculidae, 205
Trimusculotrema, 145
Trimusculus, 205
Trinema, 57
Trinemataidae, 57
Triodontolaimus, 174
Triopella, 201
Triophidae, 202
Tripaphylus, 262
Triphoridae, 185
Tripleuchlanis, 150
Triplignatha, 151
Triploplana, 136
Triposolenia, 53
Tripterygiidae, 372
Tripterygion, 372
Triptolemus, 86
Tripyla, 174
Tripyloides, 174
Trisaccopharynx, 134
Trischizostoma, 304
Trisegmentina, 66
Trisopterus, 365
Trissonchulus, 174
Tristenium, 319
Tristoma, 145
Tristomella, 145
Tritaeta, 299
Tritaphros, 149
Tritaxis, 66
Triticella, 332
Triticellidae, 332
Triticellopsis, 332
Tritonia, 204
Tritoniidae, 204
Tritoniopsis, 204
Trivia, 191
Triviidae, 191
Trochaclididae, 184
Trochaclis, 184
Trochammina, 66
Trochamminidae, 65
Trochamminopsis, 66
Trochamus, 174
Trochicola, 257
Trochidae, 183
Trochochaeta, 231
Trochochaetidae, 231
Trochocyathus, 109
Trochoderma, 351
Trochodata, 351
Trochopella, 146
Trochopus, 146
Trochus, 183
Trogloarctus, 236
Troglocyclopina, 254
Troglophonte, 276
Tromikosoma, 346
Trophon, 192
Tropidomyia, 211
Troschelia, 193
Truncatella, 190
Truncatellidae, 190
Truncatoflabellum, 110
Trygaeus, 242
Trypanosyllis, 230
Trypetesa, 281
Trypetesidae, 281
Tryphana, 308
Tryphanidae, 308
Tryphoema, 278
Tryphosella, 304
Tryphosites, 304
Trypostega, 331
Trypostegidae, 331
Tsengia, 36
Tuba, 196
Tubbrevia, 188
Tubecularia, 19
Tuberapseudes, 313
Tuberculocythere, 250
Tuberoloxoconcha, 250

<i>Tubiclava</i> , 117	<i>Typhlotanais</i> , 315	<i>Uria</i> , 375
<i>Tubifex</i> , 233	<i>Typton</i> , 288	<i>Urias</i> , 318
Tubificidae , 233	<i>Tyrrhenia</i> , 142	<i>Uriopha</i> , 345
Tubificinae , 233	<i>Tyrrheniella</i> , 127	<i>Uristes</i> , 304
<i>Tubifiscoides</i> , 233	Tyrrheniellidae , 127	<i>Urobopyrus</i> , 322
Tubiluchidae , 160	<i>Tythocope</i> , 317	<i>Urocorycaeus</i> , 256
<i>Tubiluchus</i> , 160	<i>Uca</i> , 292	<i>Urocythereis</i> , 249
<i>Tubolaimoides</i> , 174	<i>Udonella</i> , 146	<i>Urodasys</i> , 157
Tubulanidae , 152	Udoteaceae , 21	<i>Uromunna</i> , 319
<i>Tubulanus</i> , 152	<i>Ulcina</i> , 369	<i>Uronema</i> , 22
<i>Tubularia</i> , 113	<i>Ulianinia</i> , 129	<i>Urophycis</i> , 365
Tubulariidae , 113	<i>Ulianiniidae</i> , 129	<i>Uroptychus</i> , 290
<i>Tubulipora</i> , 333	<i>Ulkenia</i> , 78	<i>Urosalpinx</i> , 192
Tubuliporidae , 333	Ulmaridae , 111	<i>Urospora</i> , 21
<i>Tuleariocaris</i> , 288	<i>Ulonema</i> , 25	<i>Urothoe</i> , 308
<i>Tulostoma</i> , 18	<i>Ulophysema</i> , 283	Urothoidae , 308
Tulostomataceae , 18	<i>Ulosa</i> , 97	Ursidae , 376
Tulostomatales , 18	<i>Ulothrix</i> , 23	<i>Ursus</i> , 376
<i>Tulpa</i> , 119	Ulotrichaceae , 23	<i>Urticina</i> , 106
<i>Tumidotubus</i> , 65	Ulotrichales , 23	<i>Uschakovia</i> , 327
Tunicata , 352	<i>Ulva</i> , 24	<i>Uschakovius</i> , 215
<i>Tunicatipsora</i> , 17	Ulvaceae , 24	Ustigmoyctetes , 18
<i>Tuoba</i> , 323	<i>Ulvalles</i> , 23	Ustilagiales , 18
<i>Tuphacheres</i> , 259	<i>Ulvaria</i> , 24	<i>Ute</i> , 102
<i>Turbanella</i> , 158	<i>Ulrella</i> , 22	<i>Utelga</i> , 132
Turbanellidae , 158	Ulvellaceae , 24	Uteriporidae , 136
<i>Turbellaria</i> , 125	<i>Umagilla</i> , 130	<i>Uteriporus</i> , 136
<i>Turbicellepora</i> , 328	Umagillidae , 130	<i>Utolineides</i> , 154
Turbinellidae , 194	<i>Umbellisyllys</i> , 230	<i>Utriculopsis</i> , 200
Turbinidae , 185	<i>Umbellula</i> , 106	<i>Uvigerina</i> , 75
Turbinoliidae , 110	Umbellulidae , 106	Uvigerinidae , 75
<i>Turbonilla</i> , 198	<i>Umbilicalis</i> , 187	<i>Vaginulina</i> , 75
<i>Turborotalita</i> , 73	<i>Umbonula</i> , 331	Vaginulinidae , 75
<i>Turcicula</i> , 184	Umbonulidae , 331	<i>Vahlkampfia</i> , 54
<i>Turnerella</i> , 37	Umbraculidae , 201	Vahlkampfiidae , 54
Turridae , 195	<i>Umbraculum</i> , 201	<i>Valbyteuthis</i> , 213
Turrilinidae , 75	<i>Umbrina</i> , 371	<i>Valdivianemertes</i> , 154
<i>Turrisipho</i> , 193	<i>Unanereis</i> , 221	<i>Valdiviella</i> , 263
<i>Turrisphaera</i> , 76	<i>Unantra</i> , 125	<i>Valenciennellus</i> , 362
<i>Turritella</i> , 185	<i>Uncimenia</i> , 182	<i>Valencinia</i> , 154
<i>Turritellella</i> , 62	<i>Uncinaria</i> , 233	Valenciniidae , 154
Turritellidae , 185	<i>Uncinorhynchus</i> , 131	<i>Valencinura</i> , 153
<i>Turritellopsis</i> , 196	<i>Uncinotarsus</i> , 298	<i>Valettietta</i> , 308
<i>Turritopsis</i> , 117	<i>Unciola</i> , 298	Valettiidae , 308
<i>Tursiops</i> , 376	<i>Uncioella</i> , 298	<i>Valettiopsis</i> , 308
<i>Turtonia</i> , 208	<i>Uncisudis</i> , 364	Valifera , 321
Turtoniidae , 208	<i>Undaria</i> , 28	<i>Valipora</i> , 149
<i>Tvaerminnea</i> , 133	<i>Undeuchaeta</i> , 263	<i>Vallicula</i> , 123
<i>Tylaster</i> , 341	<i>Undinella</i> , 268	<i>Vallisia</i> , 146
<i>Tylexocladus</i> , 89	<i>Undinula</i> , 264	<i>Valonia</i> , 23
<i>Tylobranchion</i> , 352	<i>Ungulina</i> , 208	Valoniaceae , 23
<i>Tylocephalum</i> , 149	Ungulinidae , 208	Valsaceae , 15
<i>Tyloodelphys</i> , 142	<i>Unicapsula</i> , 84	<i>Valvaelaimus</i> , 174
<i>Tylodina</i> , 201	Uniporidae , 155	Valvatida , 339
Tylochinidae , 201	<i>Uniporus</i> , 155	<i>Valvognathia</i> , 151
<i>Tylosurus</i> , 367	<i>Unitubulotestis</i> , 142	<i>Valvulineria</i> , 69
<i>Typhinellus</i> , 192	<i>Upeneus</i> , 371	Vampyroteuthidae , 213
<i>Typhlamphiascus</i> , 273	<i>Upogebia</i> , 289	<i>Vampyroteuthis</i> , 213
<i>Typhlocythere</i> , 251	Upogebiidae , 289	<i>Vanadis</i> , 215
<i>Typhlomangelia</i> , 195	Uranoscopidae , 372	<i>Vanbenedenia</i> , 261
Typhloplanida , 133	<i>Uranoscopus</i> , 372	<i>Vanhoeffenella</i> , 63
Typhloplanidae , 134	<i>Urasterias</i> , 342	Vanikoridae , 190
<i>Typhlopoly cystis</i> , 132	<i>Urastoma</i> , 129	<i>Vanneaugobius</i> , 373
<i>Typhlorhynchus</i> , 133	Urastomidae , 129	<i>Vannella</i> , 54
<i>Typhloscolecidae</i> , 231	<i>Urceolus</i> , 60	Vannellidiae , 54
<i>Typhloscolex</i> , 231	Urechinidae , 347	<i>Vannuccia</i> , 113, 135
<i>Typhlotanaidae</i> , 315	<i>Urechinus</i> , 347	<i>Vargula</i> , 245

- Variisporina**, 82
Variolepis, 149
Vasconiella, 208
Vasculonema, 174
Vasichona, 41
Vaucheria, 24
Vaucheriales, 24
Vaughanella, 110
Vaunthompsonia, 309
Vaunthompsoniinae, 309
Vectoriella, 257
Vejdovska, 130
Velamen, 122
Velatida, 341
Vellela, 113
Veleroninoides, 63
Veleropilina, 182
Velutina, 191
Velutinidae, 191
Venaficula, 361
Venericardia, 208
Veneridae, 210
Venerupis, 210
Ventricaria, 23
Ventriculinidae, 258
Ventromma, 115
Venus, 210
Veretillidae, 106
Veretillum, 106
Vermetidae, 190
Vermetus, 190
Vermiliopsis, 227
Verneuilina, 66
Verneuilinidae, 66
Verneuilinulla, 66
Verongida, 101
Verruca, 282
Verrucidae, 282
Verrucocythere, 251
Verrucomorpha, 282
Vertebralina, 68
Verticillium, 19
Verticordia, 211
Verticordiidae, 211
Verum, 282
Vesicomya, 210
Vesicomyidae, 210
Vesicularia, 332
Vesiculariidae, 332
Vestimentifera, 235
Vettoria, 258
Vetulonia, 185
Vexillifera, 54
Vexilliferidae, 54
Vexillum, 194
Vibilia, 308
Vibiliidae, 308
Vickersia, 32
Victorella, 332
Victorellidae, 332
Vigtoriella, 216
Villogorgia, 105
Viminella, 105
Vinciguerria, 363
Vinearia, 228
Virgularia, 106
Virgulariidae, 106
Virgulina, 73
Virgulopsis, 73
Viscosa, 174
Vitreledonella, 213
Vitreledonellidae, 213
Vitreolina, 188
Vitreotubus, 227
Vogtia, 121
Volutella, 19, 57
Volutellidae, 57
Volutidae, 194
Volutomitra, 194
Volutomitridae, 194
Volutopsis, 193
Volvarina, 194
Volvatellidae, 200
Volvocaceae, 24
Volvocales, 24
Volvulella, 199
Vorticeros, 129
Vosmaeria, 89
Vosmaeropsis, 102
Vulvulina, 65
Wahlia, 130
Wailesella, 56
Walkeria, 332
Walkeriidae, 332
Wardium, 149
Wardula, 142
Warnowia, 53
Watersipora, 332
Watersiporidae, 332
Webbinella, 65
Weberella, 89
Websterinereis, 221
Weelia, 356
Weinkauffia, 199
Wellmanellinella, 66
Weltneria, 281
Weltnerium, 282
Westbladiella, 133
Westwoodilla, 305
Weyprechtia, 300
Whoia, 317
Wieseria, 174
Wijnhoffella, 155
Willemoesia, 289
Willemstia, 272
Williamia, 205
Wilsoniella, 224
Wingstrandarctus, 236
Winkenthughesia, 146
Wirenia, 181
Witrockiellaceae, 23
Witrockiella, 23
Woloszynskia, 53
Womersleyella, 33
Wrangelia, 32
Wurdemannia, 37
Wurdemanniaceae, 37
Xaiva, 291
Xanthidae, 292
Xantho, 292
Xanthocalanus, 266
Xanthodaphne, 196
Xanthophyceae, 24
Xeniidae, 105
Xennella, 174
Xenobalanus, 282
Xenobranchion, 352
Xenocicerina, 131
Xenocoeloma, 259
Xenocoelomidae, 259
Xenodasys, 158
Xenodermichthys, 362
Xenodice, 306
Xenodonta, 182
Xenophora, 190
Xenophoridae, 190
Xenophyophora, 78
Xenoskenea, 197
Xenostrobus, 207
Xenosyllis, 230
Xenoplanula, 136
Xenotrichula, 157
Xenotrichulidae, 157
Xenoturbella, 205
Xenoturbellidae, 205
Xenuraega, 319
Xestoleberididae, 251
Xestoleberis, 251
Xestospongia, 98
Xiphias, 373
Xiphidae, 373
Xyala, 174
Xylariaceae, 18
Xylariales, 18
Xylodiscula, 196
Xylodisculidae, 196
Xylophaga, 211
Xylophagidae, 211
Xyloredo, 211
Xyloskenea, 185
Xyrichtys, 371
Yamagutisentis, 177
Yoldia, 206
Yoldiella, 206
Yoldiidae, 206
Ypsilothuria, 349
Ypsilothuriidae, 349
Yungia, 128
Zalerion, 19
Zalophotrema, 142
Zanardinia, 25
Zanclea, 113
Zancleidae, 113
Zariquieyon, 292
Zatsepinia, 215
Zaus, 274
Zausoscidia, 277
Zebina, 190
Zebrus, 373
Zeidae, 368
Zeidora, 183
Zeiformes, 368
Zelinkaderes, 160
Zelinkaderidae, 160
Zelinkiella, 151
Zenion, 368
Zenkevitchiella, 264
Zenopsis, 368
Zeppelinina, 217

<i>Zerotula</i> , 188	Zoarcidae , 371	Zosteraceae , 39
Zerotulidae , 188	<i>Zoarella</i> , 68	<i>Zosterisessor</i> , 373
<i>Zeuctocrinus</i> , 338	<i>Zonaria</i> , 27, 191	<i>Zosterocarpus</i> , 27
<i>Zeugopterus</i> , 374	<i>Zonorhynchus</i> , 131	<i>Zschokkella</i> , 83
<i>Zeus</i> , 368	<i>Zoobotryon</i> , 332	<i>Zu</i> , 364
<i>Zeuxo</i> , 314	<i>Zoogonoides</i> , 142	<i>Zygapikodinium</i> , 53
Ziphiidae , 376	<i>Zoogonus</i> , 142	<i>Zygomitus</i> , 22
<i>Ziphius</i> , 376	Zopfiaceae , 16	<i>Zygomolgus</i> , 256
<i>Zirfaea</i> , 210	<i>Zoroaster</i> , 342	Zygomycetes , 15
Zoanthidae , 109	Zoroasteridae , 342	Zygomycota , 15
Zoanthidea , 108	<i>Zosime</i> , 280	<i>Zygonemertes</i> , 154
<i>Zoarces</i> , 371	<i>Zostera</i> , 39	<i>Zygophylax</i> , 115

MUSÉUM NATIONAL D'HISTOIRE NATURELLE
SERVICE DU PATRIMOINE NATUREL,
INSTITUT D'ÉCOLOGIE ET DE GESTION DE LA BIODIVERSITÉ,
57, rue Cuvier - 75231 Paris Cedex 05



MUSÉUM NATIONAL D'HISTOIRE NATURELLE
SERVICE DES PUBLICATIONS SCIENTIFIQUES
Diffusion : Sandrine HOFFART MULLER
57, rue Cuvier - F 75005 Paris
Tel. : [33] 01 40 79 48 05
Fax : [33] 01 40 79 38 40
e-mail : diff.pub@mnhn.fr
<http://www.mnhn.fr/publication>




Date de distribution : le 21 Décembre 2001
Achevé d'imprimer en Décembre 2001
par EXPRESS TIRAGES à Bourg la Reine
Dépôt légal : Décembre 2001

PHOTOCOPIES

Les publications scientifiques du Muséum adhèrent au Centre Français d'Exploitation du Droit de Copie (CFC), 20, rue des Grands Augustins, 75006 Paris. Le CFC est membre de l'International Federation of Reproduction Rights Organisations (IFRRO). Aux Etats-Unis d'Amérique, contacter le Copyright Clearance Center, 27 Congress street, Salem, Massachusetts, 01970.


PHOTOCOPIES

The scientific publications of the Museum adhere to the Centre Français d'Exploitation du Droit de Copie (CFC), 20, rue des Grands Augustins, 75006 Paris. The CFC is a member of International Federation of Reproduction Rights Organizations (IFRRO). In USA, contact the Copyright Clearance Center, 27 Congress street, Salem, Massachusetts, 01970.



The foundation of biodiversity research and management is correctly identifying and naming species, but different names are used for the same species, and the same names for different species, in different parts of Europe. The present volume is the first coordinated attempt to list the marine species – plants, algae, invertebrates, vertebrates – from all over the European seas, from the Arctic and Iceland to the Canaries, the Mediterranean and the Baltic. The main part of the volume is a checklist of some 30,000 valid species, for certain groups supplemented by synonymies and basic distribution information. A bibliographical section points to ca. 700 identification guides. The volume is the collective work of 170 persons who contributed their expertise to the *European Register of Marine Species* (ERMS), a project part-funded by the European Commission (EC) Marine Science and Technology (MAST) research programme.

It is anticipated that the *European Register of Marine Species* will become a standard reference tool for marine biodiversity training, research and management in Europe. The present paper version is published in parallel with an electronic version available from the project web site <http://erms.biol.soton.ac.uk>.



Les noms scientifiques sont le véhicule de communication de tous les résultats de recherches et textes réglementaires sur la biodiversité, mais usages et habitudes d'emploi des noms varient encore d'un pays à l'autre. Pour la première fois, ce volume présente un répertoire complet et coordonné de toutes les espèces – plantes, algues, invertébrés, vertébrés – à l'échelle des mers d'Europe, de l'Océan Arctique et de l'Islande aux Canaries, à la Méditerranée et à la Baltique. Le résultat est une liste de près de 30 000 espèces valides, complétée pour certains groupes par des indications synonymiques et géographiques. Un index bibliographique de 700 titres oriente vers les principaux guides et outils de détermination de la flore et de la faune marines d'Europe. Ce volume est le travail collectif de près de 170 scientifiques dans le cadre du projet *European Register of Marine Species* (ERMS), soutenu par la Commission Européenne dans le cadre de son programme Marine Science and Technology (MAST).

Ce catalogue est destiné à devenir un outil de standardisation et de référence pour ceux qui sont impliqués dans la recherche, la formation et la gestion de la biodiversité marine en Europe. Cette version papier est publiée parallèlement à la liste consultable à l'adresse <http://erms.biol.soton.ac.uk>.

