

PROCEEDINGS
OF THE
ZOOLOGICAL SOCIETY
OF LONDON.



PART XXIII.

1855.

PRINTED FOR THE SOCIETY;
SOLD AT THEIR HOUSE IN HANOVER-SQUARE,
AND BY MESSRS. LONGMAN, BROWN, GREEN AND LONGMANS,
PATERNOSTER-ROW.

mis 3 planis, sequentibus convexiusculis, ultimo $\frac{2}{5}$ longitudinis subæquante, inflato; apertura obliqua, truncato-auriformi; plica columellari supera, nodiformi, parum torta; perist. fusco limbato, margine dextro subrecto, strictiusculo, columellari dilatato, appresso.

Long. $21\frac{1}{2}$, diam. 13 mill.

β. Nigra, pallide interrupte lineolata vel fasciata.

Hab. In insulis Sandwich (*Frick*).

25. *A. GLOBOSA*, Pfr. (Pl. XXX. fig. 25.) *A. testa subimperfurata, dextrorsa, conico-globosa, tenuiuscula, striatula, alba, lineis fuscis cingulata; spira brevi, convexiusculo-conica, subacuta; sutura levi, submarginata; anfr. 5, modice convexis, ultimo ventroso, spiram subæquante, basi fusco; apertura obliqua, subtetra-gono-ovali, intus margaritacea; plica columellari levi, vix torta; perist. acuto, fusco-limbato, intus albo-sublabiato, margine columellari incrassato, subadnato.*

Long. 17, diam. $11\frac{1}{2}$ mill.

Hab. In insulis Sandwich (*Frick*).

26. *A. CONSPERSA*, Pfr. (Pl. XXX. fig. 26.) *A. testa imperforata, dextrorsa, conico-ovata, tenuiuscula, conferte striata, saturate fusca, maculis sordide albidis, strigas angulosas formantibus conspersa; spira convexo-conica, apice acuta; sutura simplice; anfr. 6, superis planiusculis, penultimo convexiore, ultimo $\frac{3}{4}$ longitudinis subæquante, rotundato; apertura parum obliqua, sinuato-elliptica; plica columellari subbasali, compressa; perist. simplice, recto, margine columellari nullo.*

Long. 18, diam. $10\frac{1}{2}$ mill.

Hab. In insulis Sandwich (*Frick*).

Sect. LABIELLA, Pfr. in Malak. Bl. 1854, p. 142.

27. *A. DENTATA*, Pfr. (Pl. XXX. fig. 27.) *A. testa imperforata, conico-ovata, solida, striatula, nitida, sub epidermide fulvida alba; spira conica, obtusa; anfr. 7, planiusculis, ultimo $\frac{3}{4}$ longitudinis subæquante, basi rotundato; apertura fere verticali, sub-rhombea, basi angulata; plica columellari profunda, compressa; perist. albo-callosa, marginibus callo junctis, dextro recto, medio introrsum unidentato; columellari dilatato, adnato.*

Long. 12, diam. $5\frac{2}{3}$ mill.

Hab. In insulis Sandwich (*Dr. Newcomb*).

2. DESCRIPTIONS OF NINE NEW SPECIES OF LAND-SHELLS, IN THE COLLECTION OF H. CUMING, ESQ. BY DR. L. PFEIFFER.

1. *BULIMUS LATILABRIS*, Pfr. *B. testa imperforata, succinoidea, tenui, longitudinaliter subgranulato-rugulosa, flavida, strigas angulatis, castaneis subtessellata, vel ad suturam interrupte fasciata; spira brevi, conica, acutiusecula; anfr. 4, rapide accrescentibus, vix convexiusculis, ultimo fere $\frac{3}{4}$ longitudinis formante; columella ad apicem spiræ aperta; apertura subverti-*

cali, acuminato-ovali, intus margaritacea; perist. late expanso et reflexiusculo, marginibus callo tenui junctis, columellari arcuato, calloso, albo.

Long. 49, diam. 26 mill.

Hab. Santa Fé de Bogota.

2. *BULIMUS STRANGEI*, Pfr. *B. testa subcompressa umbilicata, fusiformi-oblonga, tenuiuscula, striata et strüs spiralibus levisime decussata, fulvida, ad suturam crenulatam albo-fasciata; spira elevato-conica, apice obtusa; anfr. 5, vix convexiusculis, ultimo spiram superante, antice breviter arcuatim ascendente, basi attenuato; apertura vix obliqua, elongato-ovali, intus vitellina, nitida; columella alba, longe plicata; perist. albo, subincrassato, margine dextro expanso et reflexo, superne arcuato, tum strictiusculo, columellari dilatato, plano, patente.*

Long. 46, diam. 17 mill.

Hab. Eddystone Island, Australian Seas.

3. *BULIMUS EDDYSTONENSIS*, Pfr. *B. testa imperforata, ovato-conica, tenuiuscula, confertim striata et suboblique malleata, sericea, saturate castanea; spira exacte conica, apice acuta; sutura levi, subcrenata; anfr. 6, convexiusculis, ultimo spira paulo longiore, basi rotundato; columella tenui, subplicata, recedente, aurantiaca; apertura parum obliqua, subangulato-ovali, intus margaritacea, virenti-livida; perist. simplice, vix expansiusculo, obtuso, sordide aurantiaco, margine columellari vix dilatato.*

Long. 74, diam. 34 mill.

Hab. Eddystone Island, Australian Seas.

4. *BULIMUS SIGNIFER*, Pfr. *B. testa perforata, subfusiformi-oblonga, tenui, confertim et leviter plicato-striata, albida, strigis spadiceis serratis irregulariter signata; spira inflato-conica, acutiuscula; anfr. 6, superis planiusculis, penultimo convexo, ultimo spiram subæquante, basi attenuato; columella torta, recedente; apertura vix obliqua, oblonga; perist. simplice, margine dextro sublato expanso, columellari dilatato, abrupte reflexo, patente.*

Long. 33, diam. $13\frac{1}{2}$ mill.

Hab. Venezuela?

5. *BULIMUS CERATACME*, Pfr. *B. testa profunde et anguste rimata, subfusiformi-oblonga, solidula, irregulariter striatula, parum nitente, carneo-albida; spira elongata, apice cornea, obtusula; anfr. $7\frac{1}{2}$, parum convexis, medianis remote et obsolete corneo-strigatis, ultimo $\frac{1}{3}$ longitudinis vix superante, basi subattenuato; columella recedente, leviter plicata; apertura vix obliqua, oblongo-ovali; perist. acuto, intus sublabiato, marginibus approximatis, callo tenui junctis, dextro expanso, superne arcuato, columellari patente.*

Long. $18\frac{1}{2}$, diam. 6 mill.

Hab. Peru?

6. *BULIMUS PEREGRINUS*, Pfr. *B. testa perforata, subfusiformi-turrita, solidiuscula, striatula, cerea; spira convexo-turrita, apice obtusula; sutura subcrenolata; anfr. 8½, vix convexiusculis, ultimo ⅓ longitudinis subæquante, basi subcompresso; apertura obliqua, oblonga, utrinque angulata; perist. simplice, recto, marginibus conniventibus, dextro arcuato, columellari fere stricto, a basi aperturæ dilatato, patente, planato.*

Long. 18, diam. $5\frac{2}{3}$ mill.

Hab. —?

7. *SPIRAXIS MANDARINA*, Pfr. *Sp. testa imperforata, cylindraceo-turrita, solidula, lævigata, nitida, lutescenti-cerea; spira elongata, apice attenuata, obtusa; sutura submarginata; anfr. 8, primis rotundatis, sequentibus modice convexis, ultimo ¼ longitudinis subæquante, basi rotundato; columella callosa, torta, subduplicata; apertura obliqua, elliptico-ovali; perist. simplice, recto, margine dextro superne antrorsum arcuato.*

Long. 23, diam. 6 mill.

Hab. China.

Allied to *Achatina erecta*, Bens., which is also a *Spiraxis*. *Achatina Dunkeri*, Pfr. must likewise be transferred to the same genus.

8. PUPA (ENNEA) *OBOVATA*, Pfr. *P. testa breviter arcuato-rimata, obovata, tenui, lævigata, hyalina; spira ventrosa, sursum turgida, obtusa; sutura lineari-marginata; anfr. 7, primis 4 convexis, sequentibus subplanis, ultimo ½ longitudinis vix attingente, attenuato, basi compresso, latere dextro scrobiculato; apertura verticali, subtriangulari, plica profunda columellari, lamina valida, intrante ad angulum parietis, denticulis 2 marginis dextri et singulo subbasali coarctata; perist. albo, breviter expanso.*

Long. 6, diam. supra medium 3 mill.

Hab. Liberia.

9. PUPA (ENNEA) *CEYLANICA*, Pfr. *P. testa breviter rimata, subcylindrica, striatula, pellucida, nitida, lutescenti-hyalina; spira versus apicem obtusum vix attenuata; sutura sub lente regulariter denticulata; anfr. 8½, convexis, ultimo ¼ longitudinis non attingente, latere constricto et antice profunde scrobiculato, basi gibbo; apertura subverticali, truncato-oblonga, plica profunda columellari, plica laminæformi juxta angulum parietis, dente valido in medio marginis dextri et minore basali coarctata; perist. albo, nitido, reflexiusculo.*

Long. $6\frac{2}{3}$, diam. 2 mill.

Hab. Ceylon (*E. L. Layard*).

3. NOTICE OF THE HORNS OF AN UNRECORDED SPECIES OF PRONG HORN (ANTILOCAPRA), IN THE COLLECTION OF THE DERBY MUSEUM, LIVERPOOL. BY DR. JOHN EDWARD GRAY, F.R.S., V.P.Z.S. ETC.

Some years ago the late Earl of Derby showed me a pair of horns attached together by the skin of the forehead, which he had then