

PROCEEDINGS
OF THE
ZOOLOGICAL SOCIETY
OF LONDON.



PART XXII.

1854.

PRINTED FOR THE SOCIETY ;
SOLD AT THEIR HOUSE IN HANOVER SQUARE,
AND BY MESSRS. LONGMAN, BROWN, GREEN, AND LONGMANS,
PATERNOSTER ROW.

dentibus 2 subæqualibus munito, columellari dilatato, umbilicum fere vel omnino claudente.

Long. 24, diam. max. 11, min. 9 mill.

Hab. Isle of Aneiteum, New Hebrides (*Macgillivray*).

2. **PYTHIA OVATA**, Pfr. *P. testa profunde rimata vel aperte umbilicata, compresse ovato-acuminata, solidula, striatula, pallida, livido-nebulosa et maculis castaneis marmorata; spira convexiusculo-conica, acuta; varicibus albis, macula æquilata nigro-castanea, suturam non attingente, marginatis; anfract. 10 planiusculis, antice descendantibus, ad suturam arcuato-plicatis, ultimo spiræ paulo longiore, basi subpresso; apertura verticali, intus albida; dentibus parietalibus 2, superiore triangulari, bicruri, altero valido, simplice, oblique descendente; plica columellari compressa, obliqua; perist. pallide livido, marginibus callo tenui, nitido junctis, dextro superne subsimplice, tum incrassato, reflexo, intus callo albo, inæqualiter 5-dentato munito, columellari per-dilatato, patente.*

Long. 31, diam. max. 26, min. 23½ mill.

Hab. —?

3. **PYTHIA CELEBENSIS**, Pfr. *P. testa anguste umbilicata, ovato-conica, solida, irregulariter striata, subunicolore saturate fulva; spira conica, acuta; varicibus fulvo-luteis; anfract. 10 planiusculis, antice descendantibus, ad suturam valide arcuato-plicatis, ultimo spiram subæquante, basi subpresso; apertura subobliqua, intus livido-carnea, nitida; dentibus parietalibus 2, superiore subtrigono, altero valido, compresso, subobliquo; plica columellari parvula, dentiformi, arcuatim ascendente; perist. livido-limbato, marginibus callo tenuissimo junctis, dextro superne subsimplice, tum breviter expanso et reflexiusculo, intus callo lato, carneo, inæqualiter 5-dentato munito, columellari crasso, sinuoso-patente.*

Long. 33, diam. max. 18, min. 15 mill.

Hab. Celebes.

3. DESCRIPTIONS OF EIGHTEEN NEW SPECIES OF CYCLOSTOMACEA, FROM MR. CUMING'S COLLECTION.
BY DR. L. PFEIFFER.

1. **CYCLOSTOMA CHINENSE** (*CYCLOTUS*?), Pfr. *C. testa umbilicata, convexo-depressa, solida, striata, sub epidermide fuscenti-fulva alba, obsolete fasciata; spira convexa, vix conoidea, apice subtili; anfract. 4 conveaxis, rapide accrescentibus, ultimo terete; umbilico conico, $\frac{1}{5}$ diametri occupante; apertura subverticali, circulari, intus margaritacea; perist. continuo, breviter adnato, duplice: interno haud porrecto, externo breviter patente, obtuso.—Operc. —?*

Diam. maj. 14, min. 11½, alt. 8 mill.

Hab. Hong Kong, China (*Mr. Fortune*).

2. CYCLOSTOMA MONTANUM (CYCLOTUS?), Pfr. *C. testa umbilicata, turbinato-depressa, solida, superne confertim spiraliter lirata, alba, fuscculo obsolete maculata et fasciata; spira turbinata, acutiuscula; anfract. 4 convexis, rapide accrescentibus, ultimo non descendente, subtus laevigato, in umbilico mediocri, profundo, spiraliter striato; apertura parum obliqua, angulato-rotundata; perist. simple, recto, breviter adnato, margine columellari reflexiusculo.—Operc. —?*

Diam. maj. $12\frac{1}{2}$, min. 10, alt. 8 mill.

Hab. Neilgherries.

3. CYCLOSTOMA BIROSTRE (CYCLOTUS), Pfr. *C. testa umbilicata, depressa, subdiscoidea, solidula, irregulariter striata, fulvolutea, fascia peripherica castanea et superne strigis latis angulatis ornata; spira plana, vertice subtili, vix prominulo; anfract. fere 5 convexiusculis, rapide accrescentibus, ultimo terete, subtus obsolete maculato, antice soluto et tubulo antrorum curvato munito; umbilico lato, anfractus omnes latiores et convexitores monstrante quam superne; apertura diagonali, circulari; perist. duplice, interno brevi, recto, externo expanso et inflexo, superne in rostrum apertum, anfractum ultimum tangentem, producto.—Operc. album, arctispiram, extus concavum.*

Diam. maj. 26, min. 19, alt. 7 mill.

Hab. Sarawak, Borneo.

4. CYCLOSTOMA PTEROCYCLOIDES (CYCLOTUS), Pfr. *C. testa umbilicata, depressa, subdiscoidea, solidula, striatula et striis spiralibus sub lente decussata, lutea, strigis angulatis castaneis picta; spira plana, vertice mucronatim prominulo; anfract. 5 rapide accrescentibus, parum convexis, ultimo depresso-rotundato, antice soluto et spiraculo antrorum vix ascendentem munito; umbilico profundo, $\frac{1}{3}$ diametri superante; apertura diagonali, circulari; perist. duplice, interno vix porrecto, superne leviter emarginato, externo dilatato, rectangule patente, limbo subinfuso, superne subangulatim producto.—Operc. —?*

Diam. maj. 25, min. 19, alt. 7 mill.

Hab. —?

5. CYCLOSTOMA FLORIDUM (CYCLOPHORUS), Pfr. *C. testa anguste umbilicata, globoso-turbinata, tenuiuscula, oblique levissime striatula, albida, maculis sagittiformibus fuscis, interdum in strigas angulosas confluentibus, conspersa; spira turbinata, acutiuscula, apice saepe nigro-cornea; anfract. 5 convexis, ultimo rotundato, infra peripheriam fascia castanea, deorsum dissoluta, tenuiato; apertura obliqua, subcirculari; perist. simple, breviter expanso, marginibus callo brevi junctis, sinistro subrefexo.—Operc. —?*

Diam. maj. 30, min. 24, alt. 20 mill.

Hab. —?

6. CYCLOSTOMA EXALTATUM (CYCLOPHORUS), Pfr. *C. testa*

umbilicata, globoso-turbinata, tenuiuscula, oblique striata, nigrocastanea, strigis angulosis pallidis notata; spira elevato-turbinata, acutiuscula; anfract. 5 regulariter accrescentibus, convexis, ultimo rotundato, peripheria pallide subfasciato; apertura vix obliqua, subcirculari, intus cærulescenti-albida; perist. simple, albo, marginibus callo brevi junctis, dextro perarcuato, breviter expanso, sinistro supra partem umbilici perangusti fornicatim patente.—Operc. —?

Diam. maj. 25, min. 21, alt. 19 mill.

Hab. Hong Kong, China (*Mr. Fortune*).

7. **CYCLOSTOMA SUMATRENSE (CYCLOPHORUS)**, Pfr. *C. testa umbilicata, depresso-turbinata, tenui, spiraliter tenuissime striata, nitidula, fusca, ad suturam maculis alternis castaneis et pallidis tæniata, infra peripheriam fasciis 2 articulatis ornata; spira breviter turbinata, acutiuscula; anfract. 5 rapide accrescentibus, superis convexiusculis, ultimo turgido, peripheria subcarinato; umbilico mediocri, infundibuliformi; apertura obliqua, subcirculari, intus margaritacea; perist. albo, simple, subcontinuo, breviter adnato, superne et ad dextram late expanso et revoluto, margine sinistro minus dilatato.—Operc. fulvum.*

Diam. maj. 45, min. 34, alt. 23 mill.

Hab. Pedang, Island of Sumatra (*Capt. A. Martin*).

8. **CYCLOSTOMA TÆNIATUM (CYCLOPHORUS)**, Pfr. *C. testa umbilicata, globoso-turbinata, solida, oblique confertim striata, parum nitente, albida, fasciis variis atrofuscis interruptis et maculatis tæniata; spira turbinata, acutiuscula; anfract. 5 convexis, ad suturam complanatis, minute maculatis, ultimo turgido, infra peripheriam carina funiformi, nonnullisque fasciis latioribus cincto, basi planiore; umbilico angusto, pervio; apertura obliqua, subcirculari, intus albida, fasciis perlucentibus; perist. simple, recto, continuo, breviter adnato, superne subangulato, margine dextro antrorsum dilatato, sinistro incrassato, subpatente.—Operc. rubello-fulvum.*

Diam. maj. 28, min. 24, alt. 18 mill.

Hab. Sumatra.

9. **CYCLOSTOMA CEREUM (CYCLOPHORUS ?)**, Pfr. *C. testa perforata, conoideo-globosa, solidula, leviter striata, parum nitente, virenti-cerea; spira conoidea, obtusula; anfract. 4 convexis, ultimo rotundato; apertura parum obliqua, subangulato-circulari; perist. continuo, breviter adnato, margine dextro recto, columellari subincrassato.—Operc. —?*

Diam. maj. 6 $\frac{2}{3}$, min. 6, alt. 5 mill.

Hab. —?

10. **CYCLOSTOMA DEPLANATUM (CYCLOPHORUS ?)**, Pfr. *C. testa late umbilicata, depressa, subdiscoidea, solida, confertim striatula, nitida, fusco-lutea vel castanea, fascia unica peripherica, lutea cineta; spira vix prominula, vertice subtili; sutura profunda;*

anfract. 5 rotundatis, ultimo terete, antice sensim descendente; apertura obliqua, subcirculari, intus alba; perist. continuo, in-crassato, expansiusculo, subduplicato, breviter adnato, margine columellari regulariter arcuato.—Operc. —?

Diam. maj. $27\frac{1}{2}$, min. 23, alt. 8 mill.

Hab. Koondah mountains, India (Mr. Pirrie).

11. **CYCLOSTOMA SEMICLAUSUM (LEPTOPOMA)**, Pfr. *C. testa perforata, globoso-turbinata, tenuiuscula, striis obliquis et spiralibus minute decussata, albida, flammis latis angulosis fusco-viridibus et fasciis inframedianis nonnullis saturatiorebus picta; spira elata, turbinata, acutiuscula; anfract. 5 convexis, rapide accrescentibus; apertura obliqua, fere circulari; perist. duplice: interno expansiusculo, incumbente, externo rectangule patente, ad perforationem reflexo, eam fere claudente.—Operc. virenti-fulvum.*

Diam. maj. 13, min. 10, alt. 10 mill.

Hab. Ceylon (Mr. Thwaites).

12. **CYCLOSTOMA SECHELLARUM (LEPTOPOMA?)**, Pfr. *C. testa anguste umbilicata, globoso-conica, tenuiuscula, radiato-striata, pluricarinata, fuscescenti-albida; spira turbinata, acutiuscula; anfract. 5 convexis, rapide accrescentibus, ultimo carinis 2-3 majoribus (maxima peripherica, aliis superis) lirisque spiralibus basi confertioribus munito; apertura vix obliqua, subcirculari; perist. simplice, breviter expanso, marginibus approximatis.—Operc. —?*

Diam. maj. 12, min. 10, alt. 10 mill.

Hab. Seychelles Islands.

13. **CYCLOSTOMA PŒCILUM (LEPTOPOMA)**, Pfr. *C. testa perforata, turbinata, tenuiuscula, oblique striata, virenti-lutea, flammis albidis et lineis spiralibus nigricantibus ornata; spira turbinata, apice acutiuscula, nigricante; anfract. 5 convexis, sensim accrescentibus, ultimo peripheria rotundato, basi planiusculo, circa perforationem punctiformem excavato; apertura parum obliqua, subcirculari; perist. simplice, recto, marginibus approximatis.—Operc. corneum.*

Diam. maj. 8, min. $6\frac{2}{3}$, alt. 6 mill.

Hab. Ceylon (Mr. Thwaites).

14. **CYCLOSTOMA KIENERI (CYCLOSTOMUS?)**, Pfr. *C. testa umbilicata, depressa, solida, radiatim conferte striata, superne liris sub 5 carinæformibus munita, parum nitida, rufa, albido-marmorata; spira brevi, subturbinata, apice nigra, acutiuscula; anfract. 5 convexis, celeriter accrescentibus, ultimo antice descendente, juxta suturam subexcavato, peripheria subobtuse carinato, basi lœvigato; umbilico profundo, $\frac{1}{3}$ diametri subæquante; apertura perobliqua, angulato-rotundata, intus ignea; perist. continuo,*

*subrecto, crasso, obtuso, superne angulatim producto, adnato.—
Operc. —?*

Diam. maj. $19\frac{1}{2}$, min. 16, alt. 9 mill.

Hab. Madagascar.

15. CYCLOSTOMA GRADATUM (CYCLOSTOMUS ?), Pfr. *C. testa perforata, globosa-conica, solidula, radiato-striata et spiraliter lirata (liris inæqualibus, juxta suturam subtilibus, tum 1-3 validis carinæformibus, reliquis obtusis), pallide lutescente; spira turbinata, gradata, acutiuscula; anfract. 5 angulato-convexus, ultimo basi inflato; apertura parum obliqua, subangulato-circulari; perist. albo, subincrassato, continuo, breviter adnato, vix expansiusculo.—Operc. —?*

Long. 7, diam. 5 mill.

Hab. Ceylon (Mr. Thwaites).

16. CATAULUS DUPLICATUS, Pfr. *C. testa subperforata, fusiformi-ovata, solida, striata striisque spiralibus obsolete decussata, virescenti-fusca, pallide nebulosa; spira curvilineari, elongata, apice acutiuscula; sutura levi, marginata; anfract. 7 $\frac{1}{2}$ vix convexiusculis, ultimo attenuato, antice subascendente; carina basali compressa, antrorsum vix dilatata; apertura circulari, basi axin vix excedente; perist. aurantiaco, duplice: interno breviter porrecto, basi profunde inciso, externo rectangule patente, basi ad canalem angustum vix protracto.*

Long. 18, diam. 7 mill.

Hab. Ceylon (Mr. Thwaites).

17. DIPLOMATINA MACGILLIVRAYI, Pfr. *D. testa sinistrorsa, breviter rimata, oblongo-ovata, tenui, diaphana, pallide cornea; spira turgida; anfract. 6 convexis, superis 2 apicem subpapillatum formantibus, sequentibus 2 distanter cordato-costulatis, penultimo latissimo, confertius costulato, ultimo antice ascende. $\frac{1}{3}$ longitudinis paulo superante, confertissime costulato, basi rotundato; apertura vix obliqua, subcirculari; perist. duplice: interno appresso, externo breviter et æqualiter expanso.*

Long. 6, diam. 3 mill.

β. *Violaceo-fusca, anfr. ultimo pallide unifasciato.*

Hab. Lord Howe's Island, Australian seas (Mr. Macgillivray).

18. DIPLOMATINA CAPILLACEA, Pfr. *D. testa sinistrorsa, breviter rimata, ovato-conica, tenui, capillaceo-striata, pellucida, haud nitente, pallide cornea; spira turgida, sursum conica, apice acutiuscula; anfract. 6 convexiusculis, penultimo latissimo, ultimo antice ascende. $\frac{1}{3}$ longitudinis paulo superante, basi rotundato; apertura subverticali, circulari; perist. albo, expanso, indistincte duplicato, marginibus callo junctis.*

Long. 5, diam. $2\frac{2}{3}$ mill.

Hab. Lord Howe's Island, Australian seas (Mr. Macgillivray).