

# PROCEEDINGS

OF THE

## SCIENTIFIC MEETINGS

OF THE

# ZOOLOGICAL SOCIETY OF LONDON

FOR THE YEAR

1865.



---

PRINTED FOR THE SOCIETY,  
AND SOLD AT THEIR HOUSE IN HANOVER SQUARE.

LONDON :

MESSRS. LONGMANS, GREEN, READER, AND DYER,  
PATERNOSTER ROW.

PARIS :

M. J. ROTHSCHILD,  
43, RUE SAINT ANDRÉ DES ARTS.

LEIPZIG :

M. J. ROTHSCHILD,  
34, QUERSTRASSE.

*columellaris ad marginem emersa; perist. continuum, undique breviter expansum.*

Long. 15-16, diam.  $3\frac{1}{2}$  mill.

*β. Peristome extus intusque albo-incrassato, inde apertura coarctata, lamella inferiore et plica columellari vix conspicuis.*

13. *PTERO CYCLOS WILSONI*, Pfr. (Pl. XLVI. fig. 12). *T. late umbilicata, subdiscoidea, solidula, striatula, nitida, fulva, castaneo dense fulgurata; spira medio vix elevata; anfr.  $4\frac{1}{2}$ , modice convexi, ultimus subdepresso-rotundatus, medio linea castanea cinctus; apertura diagonalis, subcircularis, intus violaceascens; perist. duplex; internum expansiusculum, adnatum, superne vix incisum, externum patens, ad anfractum contiguum productum, tum late inflexum, latere dextro angustius involutum. Operc.?*  
Diam. maj. 22, min. 18, alt.  $7\frac{1}{2}$  mill.

#### DESCRIPTION OF PLATE XLVI.

- Fig. 1. *Bulimus ineertus.*  
 2, 2a. — *swinhoei.*  
 3. — *spheroconus.*  
 4. *Helix mellea.*  
 5. — *shermani.*  
 6. — *swinhoei.*

- Fig. 7. *Helix formosensis.*  
 8. — *bacea.*  
 9. — *vesta.*  
 10. — *granti.*  
 11. *Clausilia swinhoei.*  
 12. *Pterocyclos wilsoni.*

#### 16. DESCRIPTIONS OF FIVE NEW SPECIES OF LAND-SHELLS, FROM THE COLLECTION OF THE LATE HUGH CUMING. BY DR. LOUIS PFEIFFER.

1. *BULIMUS AURIS*, Pfr. *T. profunde compresso-umbilicata, ovato-turrita, solidula, lœvigata, alba, strigis parcis undulatis castaneis signata; spira turrata, acutiuscula; anfr. 6, convexi, ultimus spiram subæquans, rotundatus, antice ascendens; apertura subverticalis, obverse auriformis, intus violaceo limbata; columella profunde et valide transverse plicata, violacea; perist. late expansum, marginibus approximatis, callo junctis, columellari flexuose elevato.*

Long. 39, diam.  $14\frac{1}{2}$  mill. Apert. cum perist. 20 mill. longa, medio 13 lata.

Hab. Venezuela.

2. *BULIMUS TENUILABRIS*, Pfr. *T. profunde umbilicata, ovato-turrita, tenuis, sublœvigata (sub lente minute spiraliter striata), nitida, alabastrina; spira turrata, apice acuto; anfr. 7, planiusculi, ultimus spira brevior, antice vix ascendens, basi compresso-rotundatus; apertura vix obliqua, truncato-oblonga; perist. simplex, margine dextro breviter expanso, columellari ad basin usque perditato, patente.*

Long. 30, diam. 12 mill. Apert. cum perist.  $14\frac{1}{2}$  mill. longa, 10 lata.

Hab. Venezuela.

3. **BULIMUS JUAREZI**, Pfr. *T. perforata, conico-ovata, tenuiuscula, striata vel plicatula, cerea, strigis linearibus olivaceis sparse notata; spira conica, acutiuscula; sutura subirregularis; anfr. 5½, convexiusculi, ultimus spiram superans, inflatus; apertura parum obliqua, angulato-subovalis, intus albido-margaritacea; perist. simplex, marginibus callo tenui vitreo junctis, dextro recto, columellari per dilatato, patente.*

Long. 31, diam. 18 mill. Apert. 19 mill. longa, 10-11 lata.

*Hab.* in provincia pacifica reipublicæ Mexicanæ.

4. **PSEUDACHATINA ELONGATA**, Pfr. *T. imperforata, oblongo-turrata, solida, irregulariter oblique plicata et supra suturam cingulo granulato munita, rubello-fulvida; spira turrata, apice obtusula; sutura albo marginata, crenata; anfr. 7½, superi convexi, sequentes subplanati, ultimus medio obtuse carinatus, ½ longitudinis superans; columella callosa, subtruncato-torta; apertura parum obliqua, oblongo-ovalis; perist. simplex, marginibus callo crasso albo junctis, dextro anguste expanso.*

Long. 84, diam. 33 mill. Apert. 37 mill. longa, 20 lata.

*β. Minor, nigro-castanea, pallide marmorata.*

Long. 70, diam. 28 mill.

*Hab.* ad fluvium Gaboon Africæ.

5. **ACHATINA CALABARICA**, Pfr. *T. oblongo-turrata, tenuiuscula, conferte striata, nitida, fusco-nigricans, interdum pallidius strigata; spira subconcavo-producta, vertice minuto; anfr. 9, modice convexi, summi costulati, ultimus ½ longitudinis subaequans, medio filocarinatus; columella arcuata, ad basin aperturæ subverticalis rhombo-ellipticæ oblique truncata; perist. simplex, rectum, marginibus callo tenui junctis.*

Long. 28, diam. 11 mill. Apert. 11 mill. longa, 6 lata.

*Hab.* Old Calabar Africæ.

#### 17. LIST OF ANIMALS COLLECTED AT MOHAMBO, MADAGASCAR, BY MR. W. T. GERRARD. BY ALFRED NEWTON, M.A., F.L.S., F.Z.S.

Specimens of animals from Madagascar are so very scarce in the museums of this country, that I make no apology for presenting to the Society a list of a small collection made in that remarkable island by Mr. W. T. Gerrard, a travelling naturalist, and lately received by me from Mr. Edward Newton. The collection does not, it is true, contain any species previously undescribed, and in this respect differs from all others from Madagascar that I have had an opportunity of examining; but several of the species are of undoubted rarity, and have perhaps never before been seen in England.

I am indebted to Mr. P. L. Sclater for being so good as to furnish me with the names of the Mammals; and as the Mammals of Ma-