

SOME NEW AND SOME PREVIOUSLY
UNREPORTED HYDROIDS, MAINLY
FROM THE CALIFORNIAN COAST

BY
C. McLEAN FRASER

UNIVERSITY OF CALIFORNIA PUBLICATIONS IN ZOOLOGY
Volume 28, No. 7, pp. 167-172, 7 figures in text

UNIVERSITY OF CALIFORNIA PRESS
BERKELEY, CALIFORNIA
1925

UNIVERSITY OF CALIFORNIA PUBLICATIONS

Notes.—The University of California Publications are offered in exchange for the publications of learned societies and institutions, universities and libraries. Complete lists of all the publications of the University will be sent upon request. For sample copies, lists of publications or other information address the **MANAGER OF THE UNIVERSITY PRESS, BERKELEY, CALIFORNIA, U. S. A.** All matter sent in exchange should be addressed to **EXCHANGE DEPARTMENT, UNIVERSITY LIBRARY, BERKELEY, CALIFORNIA, U. S. A.**

Publications of the University of California Press may be obtained from **THE CAMBRIDGE UNIVERSITY PRESS, FETTER LANE, LONDON, E. C. 4, ENGLAND**, to which orders originating in Great Britain and Ireland should be sent.

ZOOLOGY.—C. A. Kofoid and J. Grinnell, Editors.

This series contains the contributions from the Department of Zoology, from the Marine Laboratory of the Scripps Institution for Biological Research, at La Jolla, California, and from the California Museum of Vertebrate Zoology in Berkeley.

Cited as Univ. Calif. Publ. Zool.

Volume 1, 1902-1905, 317 pages, with 28 plates	\$3.50
Volume 2, 1904-1906, xvii + 382 pages, with 19 plates	\$3.50
Volume 3, 1906-1907, 383 pages, with 23 plates	\$3.50
Volume 4, 1907-1908, 400 pages, with 24 plates	\$3.50
Volume 5, 1908-1910, 440 pages, with 34 plates	\$3.50
Volume 6, 1908-1911, 478 pages, with 48 plates	\$3.50
Volume 7, 1910-1912, 446 pages, with 12 plates	\$3.50
Volume 8, 1911, 357 pages, with 25 plates	\$3.50
Volume 9, 1911-1912, 365 pages, with 24 plates	\$3.50
Volume 10, 1912-1913, 417 pages, with 10 plates	\$3.50
Volume 11, 1912-1914, 538 pages, with 26 plates	\$5.00
Volume 12, 1913-1916, 558 pages, with 22 plates	\$5.00
Volume 13, 1914-1916, 529 pages, with 39 plates	\$5.00
Volume 14, 1914-1918, 452 pages, with 60 plates	\$5.00
Volume 15, 1915-1916, 360 pages, with 38 plates	\$4.00
Volume 16, 1915-1917, 522 pages, with 46 plates	\$5.00
Volume 17, 1916-1918, 545 pages, with 24 plates	\$5.00
Volume 18, 1917-1919, 529 pages, with 20 plates	\$5.00
Volume 19, 1919-1920, 480 pages, with 33 plates	\$5.00
Volume 20, 1919-1924, 571 pages, with 48 plates	\$6.00
Vol. 21. 1. A Revision of the <i>Microtus californicus</i> Group of Meadow Mice, by Remington Kellogg. Pp. 1-42, 1 figure in text. December, 191850
2. Five New Five-toed Kangaroo Rats from California, by Joseph Grinnell. Pp. 43-47. March, 191965
3. Notes on the Natural History of the Bushy-tailed Wood Rats of California, by Joseph Dixon. Pp. 49-74, plates 1-3, 3 figures in text. December, 191925
4. Revision of the Avian Genus <i>Passerella</i> , with Special Reference to the Distribution and Migration of the Races in California, by H. S. Swarth. Pp. 75-224, plates 4-7, 30 figures in text. September, 1920	\$1.75
5. A Study of the California Jumping Mice of the Genus <i>Zapus</i> , by A. Brasler Howell. Pp. 225-238, 1 figure in text. May, 192015
6. Two New Rodents (Genera <i>Thomomys</i> and <i>Marmota</i>) from the Eastern Border of California, by Joseph Grinnell. Pp. 239-244, 6 figures in text. November, 192115
7. A Study of the Californian Forms of the <i>Microtus montanus</i> Group of Meadow Mice, by Remington Kellogg. Pp. 245-274, 25 figures in text.	
8. A Synopsis of the <i>Microtus mordax</i> Group of Meadow Mice in California, by Remington Kellogg. Pp. 275-302, plate 8, 29 figures in text. Nos. 7 and 8 in one cover. April, 192275
9. Behavior of the Leaf-nosed Snake, <i>Phyllorhynchus decurtatus</i> , by Sarah Rogers Atsath. Pp. 303-312. February, 192325
10. A Systematic List of the Mammals of California, by Joseph Grinnell. Pp. 313-324. January, 192325
11. The Systematic Status of the Mountain Lion of California, by Joseph Grinnell and Joseph Dixon. Pp. 325-332, plates 9-10. April, 192325
12. New Subspecies of Birds from Patagonia, by Alexander Wetmore. Pp. 333-337. June, 192325
13. Revision of the Genus <i>Lynx</i> in California, by Joseph Grinnell and Joseph Dixon. Pp. 339-354, plate 11, 1 figure in text. January, 192425

SOME NEW AND SOME PREVIOUSLY
UNREPORTED HYDROIDS, MAINLY
FROM THE CALIFORNIAN COAST

BY
C. McLEAN FRASER

In material collected by the United States Bureau of Fisheries Steamer "Albatross," and by the University of California, mainly from the Californian coast, sent to me for examination by Dr. C. A. Kofoid, there are seven species that appear to be new. Unfortunately, in no case was the gonosome present. Of necessity, then, some of the species can be only provisionally placed in the genus assigned. Five of the species were found in material from the Californian coast, one from off the Oregon coast, and one, rather out of place in such a collection, was taken at a "Fishhawk" station, off the Florida coast.

Besides these new species, four species, not previously reported from the Pacific coast of North America, were obtained, all from the San Francisco Bay region. As these will be described later, in a key to the hydroids of the Pacific coast of the United States, and as they have already been well described in one or more papers, only the distribution of the specimens in the collection is given here.

This paper is published with the permission of the United States Commissioner of Fisheries.

? *Coryne corrugata* sp. nov.

Figure 1

Trophosome.—Colony reaching a height of 3.5 cm., much branched, the branches coming off with a definite knee joint at a very acute angle with the stem; the secondary branches arise from the primary branches in the same manner; many of these secondary branches, while terminating in a hydranth, give off numerous branchlets that have no hydranths; perisarc thick, with deep annulations, that may be considered as corrugations, throughout the whole length of the

stem and branches; hydranths capable of great extension and usually appearing rather long and slender; tentacles 20–30, not arranged in very definite verticils.

Gonosome.—Unknown.

Distribution.—San Diego, near jetty.

Since no gonosome was present in the material examined it is not possible to say definitely whether the species is a *Coryne* or a *Syncoryne*. It bears considerable resemblance to *Coryne brachiata* Nutting, but differs from it materially in the mode of branching, and it lacks the special processes below the hydranth to which *C. brachiata* owes its name.

? *Bimeria pusilla* sp. nov.

Figure 2

Trophosome.—Colony small, straggling, less than 5 mm. high; from one to three main branches that are similar in size and appearance to the main stem; secondary branches vary in length and are irregularly placed; these may bear hydranths or may divide again to form pedicels for hydranths; the angle that the small branches make with the larger, and that the larger make with the stem, is very variable; the perisarc is thin, nowhere annulated or wrinkled, but there is a slight tendency to waviness that prevents complete smoothness; this waviness appears in the small branches as well as in the larger and the stem; hydranths with 12–14 tentacles.

Gonosome.—Unknown.

Distribution.—Lime Point, San Francisco Bay.


? *Bimeria tenella* sp. nov.

Figure 3

Trophosome.—Stem simple or with a slight tendency to fasciculation, reaching a height of 15 mm.; stem and branches slender, the branches making a wide angle with the stem, and the pedicels of the hydranths making a wide angle with the stem or branches; the pedicels attached to the stem in general much longer than those

Figures 1–7

1. *Coryne corrugata* sp. nov. Hydranths and mode of branching. $\times 5$.
2. *Bimeria pusilla* sp. nov. Main portion of colony. $\times 10$.
3. *Bimeria tenella* sp. nov. $\times 10$. *a* and *b*, two portions of colony.
4. *Campanularia castellata* sp. nov. $\times 10$. *a* and *b*, two hydrothecae.
5. *Lafæa adnata* sp. nov. $\times 10$. *a*, portion of colony showing network; *b*, separate portion of network showing two hydrothecae.
6. *Antennularia verticillata* sp. nov. Main stem showing arrangement of hydrocladia. $\times 10$.
7. *Halicornaria sinuosa* sp. nov. $\times 10$. *a*, portion of main stem; *b*, side view of hydrocladium; *c*, face view of hydrocladium.



For description see page 168.

attached to the branches; the main stem and branches mainly smooth, but there may be a wrinkling or even an annulation for short distances; pedicels annulated extensively at the base or more rarely annulated or wrinkled throughout the whole length; the perisarc surrounding the base of the hydranth is heavy and somewhat rough or wrinkled; hydranths with 10 tentacles.

Gonosome.—Unknown.

Distribution.—Several points near the entrance of San Francisco Bay in the neighborhood of Southampton Light, Angel Island, and Alcatraz Island. Depth, 9–22 fathoms.

This species bears close resemblance to *Bimeria vestita* Wright, the most important difference being in the much smaller number of tentacles in the hydranth, 16 in *B. vestita*. There is some resemblance to *B. gracilis* (Clark), also, but that species is strongly fascicled and the branches and pedicels pass out at a very narrow angle so that they are almost parallel with the stem.

? *Campanularia castellata* sp. nov

Figure 4

Trophosome.—Stem unbranched, forming the hydranth pedicel, from 0.8 mm. to 4.0 mm. in length, terminated below the hydrotheca in a ball-shaped joint, but otherwise only slightly annulated, if at all; in some cases there is a slight waviness near each extremity; stolon not annulated; hydrotheca large, 0.6 to 1.0 mm. in length and 0.4 to 0.6 mm. in greatest width, tapering but slightly from the margin to near the base, the base nearly hemispherical; the 12–14 teeth are deep, the same width throughout or slightly broader near the tip, which is just noticeably rounded; the hydrothecal wall is uniform in thickness, without lines or ridges; the space between the diaphragm and the base is shallow.

Gonosome.—Unknown.

Distribution.—Near Alcatraz Island, entrance of San Francisco Bay, 10–17 fathoms, growing on *Sertularia desmoides*.

This species bears close resemblance to *Campanularia grænlandica* Levinsen, but as *C. grænlandica* has such very distinct lines running vertically along the hydrotheca from the spaces between the teeth and these specimens have no indication of anything of the kind, they cannot well be included with that species.

Lafœa adnata sp. nov.

Figure 5

Trophosome.—Hydrothecae attached to a reticulate stolon, with no definite pedicels, although there is a definite constriction at the base of each; in nearly all cases the wall of the hydrotheca is attached

to the stolon for a part of its length but the proportion varies very materially; except for the tapering to the constriction the hydrotheca is of uniform diameter; there may or may not be a slight flare at the margin; the hydrotheca is always curved but the amount of curvature varies; in extreme cases the distal extremity is at right angles to the proximal; there is no diaphragm present; in the specimens examined the hydranth has 8 tentacles.

Gonosome.—Unknown.

Distribution.—Near Farallon Islands in 33–35 fathoms, on algae, bryozoa, and other hydroids.

***Antennularia verticillata* sp. nov.**

Figure 6

Trophosome.—A fragment of a stem 2.5 cm. long is stout and uniform, diameter 1.0 mm. The canaliculated appearance of the coenosarc is very regular, there being 24 vertical, parallel grooves showing at the surface; hydrocladia arranged in very regular whorls of six, the individual hydrocladia in one whorl opposite the spaces between the hydrocladia in the whorl above and the whorl below, so that there are really 12 vertical series, and thus two coenosarcial canals for each series; the basal hydrocladial internode is long, curved, and so much stouter than the remainder of the hydrocladium that it might be considered a process of the stem, bearing the hydrocladium, particularly, since it is at the distal extremity of the internode that the hydrocladium breaks off the most readily; this basal internode bears two nematophores; and next to it there is a hydrothecate internode, with one nematophore below the hydrotheca and two above it; a non-hydrothecate internode with two nematophores follows; the hydrothecate and non-hydrothecate internodes then alternate throughout the hydrocladium.

Gonosome.—Unknown.

Distribution.—31.7 miles N 75° E of Heeeta Head Light, Oregon coast, 84 fathoms.

In the regular and alternating whorls of hydrocladia and in the nature of the proximal internodes of the hydrocladia, this species is distinctly marked off from other species that have been described.

No species of the genus *Antennularia* has previously been reported from the Pacific coast of North America.

***Halicornaria sinuosa* sp. nov.**

Figure 7

Trophosome.—Colony unbranched, largest obtained 12 cm. in length; stem simple, divided into regular internodes, each of which bears two hydrocladia, placed on the one side of the stem at an angle of 135° or less, the one higher than the other on the internode, but the two on the one internode nearer together, in a vertical direction,

than the nearest two in succeeding internodes; hydrocladia divided into short internodes, all hydrothecate; hydrothecae with length and breadth nearly equal or slightly deeper than broad, but slightly curved, hence running almost parallel to the axis of the hydrocladium; the margin, which is sinuous, not toothed, making an angle of about 45° with the axis; the intrathecal ridge is strongly marked; mesial nematophore straight, just reaching the margin of the hydrotheca; the two supracalycine nematophores reaching the margin of the hydrotheca; two large cauline nematophores on the face of the stem at the base of each hydrocladium, and one, also large, at the back of the same process.

Gonosome.—Unknown.

Distribution.—Fishhawk Station 7511, $2\frac{1}{8}$ miles SSE of Fowey Rock Light, Gulf Stream off Cape Florida, 45 fathoms. With the hydroids *Aglaophenia rigida* and *Schizotricha tenella*.

Turritopsis nutricula McCrady

Oceania nutricula McCrady, Proceedings Elliott Soc., 1856, 1-56.

Turritopsis nutricula McCrady, Gymnoph. Charleston Har., 1857, 25.

Turritopsis nutricula Fraser, Hydroids of Beaufort, 1912, 345.

Distribution.—Oakland, California.

Pennaria tiarella McCrady

Pennaria tiarella McCrady, Gymnoph. Charleston Har., 1857, 51.

Pennaria tiarella Fraser, Hydroids of Beaufort, 1912, 355.

Distribution.—San Francisco Bay at entrance of San Pablo Bay, 4 fathoms.

Obelia bicuspidata Clarke

Obelia bicuspidata Clarke, Trans. Conn. Acad. Sci., 3, 1876, 58.

Obelia bicuspidata Fraser, Hydroids of Beaufort, 1912, 361.

Distribution.—Generally distributed throughout San Francisco Bay, from 7 to 12 fathoms; at the entrance to Carquinez Strait, at the entrance to San Pablo Bay, near Southampton Light and Alcatraz Island, and near Shag Rock in the lower bay.

Cryptolaria pulchella Allman

Cryptolaria pulchella Allman, Challenger Hyd., 23, 1888, 40.

Distribution.—Off Goat Island, San Francisco Bay, 10 fathoms.

UNIVERSITY OF BRITISH COLUMBIA,
DEPARTMENT OF ZOOLOGY,
Vancouver, B. C.

Transmitted September 24, 1925.

UNIVERSITY OF CALIFORNIA PUBLICATIONS—(Continued)

Vol. 22. 1. A Quantitative and Statistical Study of the Plankton of the San Joaquin River and Its Tributaries in and near Stockton, California, in 1913, by Winfred Emory Allen. Pp. 1-292, plates 1-12, 1 figure in text. June, 1920.	\$3.00
2. Variations in the Shell of <i>Teredo navalis</i> in San Francisco Bay, by Robert Cunningham Miller. Pp. 293-328, plates 13-17, 6 figures in text. November, 1922	.50
3. Quantitative Studies on Marine Phytoplankton at La Jolla in 1919, by Winfred Emory Allen. Pp. 329-347, 2 figures in text. November, 1922	.30
4. On the Effect of Low Salinity on <i>Teredo navalis</i> , by Harold Francis Blum. Pp. 349-368, 4 figures in text. December, 1922	.35
5. Quantitative Studies on Inshore Marine Diatoms and Dinoflagellates of Southern California in 1920, by Winfred Emory Allen. Pp. 369-378, 1 figure in text.	
6. <i>Stylarioides papillosa</i> , sp. nov., a New Annelid from the San Diego Region, by Christine E. Essenberg. Pp. 379-381, 8 figures in text. Nos. 5 and 6 in one cover. December, 1922	.25
7. The Digestion of Wood by <i>Teredo navalis</i> , by Walter H. Dore and Robert C. Miller. Pp. 388-400, plate 18. February, 1923	.25
8. Variations in the Pallets of <i>Teredo navalis</i> in San Francisco Bay, by Robert Cunningham Miller. Pp. 401-414, plates 19-20. February, 1923	.25
9. Some Tide-Water Collections of Marine Diatoms taken at Half-Hour Intervals near San Diego, California, by Winfred E. Allen. Pp. 413-416, 1 figure in text. June, 1923	.25
10. Preliminary Statistical Report on the Occurrence of Marine Copepoda in the Plankton at La Jolla, California, by Calvin O. Esterly. Pp. 417-433. June, 1923	.25
11. Studies on Marine Diatoms and Dinoflagellates Caught by Aid of the Kofoid Bucket in 1922, by Winfred Emory Allen. Pp. 435-445, 5 figures in text. October, 1923	.25
12. Statistical Studies of Marine Diatoms of the San Diego Region Collected by U.S.S. Pioneer in Midwinter of 1923, by Winfred Emory Allen. Pp. 445-448. November, 1923	.25
13. Life Cycle of <i>Oithona nana</i> Reared Experimentally, by Helen E. Murphy. Pp. 449-454, 5 figures in text. November, 1923	.25
14. Morphology of the Digestive Tract of <i>Teredo navalis</i> , by Edgar Locks Lazler. Pp. 455-474, plates 21-24. February, 1924	.35
Vol. 23. The Marine Decapod Crustacea of California, by Waldo L. Schmitt. Pp. 1-470, plates 1-50, 165 figures in text. May, 1921	5.00
Vol. 24. 1. A Geographical Study of the Kangaroo Rats of California, by Joseph Grinnell. Pp. 1-124, plates 1-7, 24 figures in text. June, 1922	1.75
2. Birds and Mammals of the Stikine River Region of Northern British Columbia and Southeastern Alaska, by H. S. Swarth. Pp. 125-314, plate 8, 34 figures in text. June, 1922	2.50
3. Birds and Mammals of the Skeena River Region of Northern British Columbia, by Harry S. Swarth. Pp. 315-394, plates 9-11, 1 figure in text. January, 1924	1.00
Vol. 25. A Bibliography of Eugenics, by Samuel J. Holmes. Pp. 1-514. January, 1924	5.00
Vol. 26. 1. On the Family of Achiridae or Broad-Soles, with description of a new species, <i>Achirus barnharti</i> , from California, by David Starr Jordan. Pp. 1-14, plate 1. December, 1923	.25
2. A Precipitin Test in Experimental Amoebic Dysentery in Cats, by Edna Hannibal Wagener. Pp. 15-20, plate 2. January, 1924	.25
3. On the Distinctions between <i>Endamoeba coli</i> and <i>Coenostimama tafleuri</i> , by Charles A. Kofoid, Olive Swezy, and John F. Kessel. Pp. 21-39, 24 figures in text. January, 1924	.45
4. The Boring Mechanism of <i>Teredo</i> , by Robert Cunningham Miller. Pp. 41-80, plates 3-6, 6 figures in text. May, 1924	.60
5. The Free-swimming Copepoda of San Francisco Bay, by Calvin O. Esterly. Pp. 81-129, 16 figures in text. May, 1924	.75
6. Fatal Effects of the Removal of the Micronucleus in <i>Euplores</i> , by Charles V. Taylor and William P. Farber. Pp. 131-144, plate 7. May, 1924	.25
7. Wood-boring Mollusks from the Hawaiian, Samoan, and Philippine Islands, by Robert Cunningham Miller. Pp. 145-158, plates 8-11.	
8. Wood-boring Crustacea from Hawaii and Samoa, by Robert Cunningham Miller. Pp. 159-164, plates 12-13. Nos. 7 and 8 in one cover. May, 1924	.45

UNIVERSITY OF CALIFORNIA PUBLICATIONS—(Continued)

Vol. 22. 1.	A Quantitative and Statistical Study of the Plankton of the San Joaquin River and Its Tributaries in and near Stockton, California, in 1913, by Winfred Emory Allen. Pp. 1-292, plates 1-12, 1 figure in text. June, 1920.	\$3.00
2.	Variations in the Shell of <i>Teredo navalis</i> in San Francisco Bay, by Robert Cunningham Miller. Pp. 293-328, plates 13-17, 6 figures in text. November, 1922	.50
3.	Quantitative Studies on Marine Phytoplankton at La Jolla in 1919, by Winfred Emory Allen. Pp. 329-347, 2 figures in text. November, 1922	.30
4.	On the Effect of Low Salinity on <i>Teredo navalis</i> , by Harold Francis Blum. Pp. 349-368, 4 figures in text. December, 1922	.35
5.	Quantitative Studies on Inshore Marine Diatoms and Dinoflagellates of Southern California in 1920, by Winfred Emory Allen. Pp. 369-378, 1 figure in text.	
6.	<i>Stylarioides papillosa</i> , sp. nov., a New Annelid from the San Diego Region, by Christine E. Essenberg. Pp. 379-381, 8 figures in text. Nos. 5 and 6 in one cover. December, 1922	.20
7.	The Digestion of Wood by <i>Teredo navalis</i> , by Walter H. Dore and Robert C. Miller. Pp. 383-400, plate 18. February, 1923	.30
8.	Variations in the Pallets of <i>Teredo navalis</i> in San Francisco Bay, by Robert Cunningham Miller. Pp. 401-414, plates 19-20. February, 1923	.35
9.	Some Tide-Water Collections of Marine Diatoms taken at Half-Hour Intervals near San Diego, California, by Winfred E. Allen. Pp. 413-416, 1 figure in text. June, 1923	.35
10.	Preliminary Statistical Report on the Occurrence of Marine Copepoda in the Plankton at La Jolla, California, by Calvin O. Esterly. Pp. 417-433. June, 1923	.95
11.	Studies on Marine Diatoms and Dinoflagellates Caught by Aid of the Kofoid Bucket in 1922, by Winfred Emory Allen. Pp. 435-445, 5 figures in text. October, 1923	.25
12.	Statistical Studies of Marine Diatoms of the San Diego Region Collected by U.S.S. Pioneer in Midwinter of 1923, by Winfred Emory Allen. Pp. 445-448. November, 1923	.25
13.	Life Cycle of <i>Oithona nana</i> Reared Experimentally, by Helen E. Murphy. Pp. 449-454, 5 figures in text. November, 1923	.25
14.	Morphology of the Digestive Tract of <i>Teredo navalis</i> , by Edgar Locke Latier. Pp. 455-474, plates 21-24. February, 1924	.55
Vol. 23.	The Marine Decapod Crustacea of California, by Waldo L. Schmitt. Pp. 1-470, plates 1-50, 165 figures in text. May, 1921	5.00
Vol. 24. 1.	A Geographical Study of the Kangaroo Rats of California, by Joseph Grinnell. Pp. 1-124, plates 1-7, 24 figures in text. June, 1922	1.75
2.	Birds and Mammals of the Stikine River Region of Northern British Columbia and Southeastern Alaska, by H. S. Swarth. Pp. 125-314, plate 8, 34 figures in text. June, 1922	2.50
3.	Birds and Mammals of the Skeena River Region of Northern British Columbia, by Harry S. Swarth. Pp. 315-394, plates 9-11, 1 figure in text. January, 1924	1.00
Vol. 25.	A Bibliography of Eugenics, by Samuel J. Holmes. Pp. 1-514. January, 1924	5.00
Vol. 26. 1.	On the Family of Achiridae or Broad-Soles, with description of a new species, <i>Achirus barnharti</i> , from California, by David Starr Jordan. Pp. 1-14, plate 1. December, 1923	.25
2.	A Precipitin Test in Experimental Amoebic Dysentery in Cats, by Edna Hannibal Wagener. Pp. 15-20, plate 2. January, 1924	.25
3.	On the Distinctions between <i>Endamoeba coli</i> and <i>Councilmania laffewi</i> , by Charles A. Kofoid, Olive Swezy, and John F. Kessel. Pp. 21-39, 24 figures in text. January, 1924	.45
4.	The Boring Mechanism of <i>Teredo</i> , by Robert Cunningham Miller. Pp. 41-80, plates 3-6, 6 figures in text. May, 1924	.30
5.	The Free-swimming Copepoda of San Francisco Bay, by Calvin O. Esterly. Pp. 81-129, 16 figures in text. May, 1924	.75
6.	Fatal Effects of the Removal of the Micronucleus in <i>Euplotes</i> , by Charles V. Taylor and William P. Farber. Pp. 131-144, plate 7. May, 1924	.25
7.	Wood-boring Mollusks from the Hawaiian, Samoan, and Philippine Islands, by Robert Cunningham Miller. Pp. 145-158, plates 8-11.	
8.	Wood-boring Crustacea from Hawaii and Samoa, by Robert Cunningham Miller. Pp. 159-164, plates 12-13. Nos. 7 and 8 in one cover. May, 1924	.45

UNIVERSITY OF CALIFORNIA PUBLICATIONS—(Continued)

9. The Cytology of <i>Endamoeba gingivalis</i> (Gros) Brumpt Compared with that of <i>E. dysenteriae</i> with Special Reference to the Determination of the Amoebas in Bone Marrow in Arthritis Deformans of Ely's Second Type, by Charles A. Kofoid and Olive Swezy. Pp. 165-198, plates 14-18, 1 figure in text. June, 1924	.50
10. Morphology and Mitosis of <i>Dinenympha fimbriata</i> sp. nov., by Harold Kirby, Jr. Pp. 199-220, plates 19-23. June, 1924	.35
11. <i>Karyamoeba falcata</i> , a New Amoeba from the Human Intestinal Tract, by Charles A. Kofoid and Olive Swezy. Pp. 221-242, plates 23-24, 2 figures in text. June, 1924	.35
12. Surface Catches of Marine Diatoms and Dinoflagellates made in 1923 by U.S.S. Pioneer between San Diego and Seattle, by Winfred Emory Allen. Pp. 243-248, plate 25. June, 1924	.25
13. Viability of Cysts of Human Intestinal Amoebas as determined by exposure to various substances and subsequent staining in Haemotoxylin, by Nathaniel Bercovitz, M.D. Pp. 249-261. September, 1924	.25
14. The Incomplete Digestive Tract of <i>Appendicularia sicula</i> , by Christine Essenberg. Pp. 263-265, 1 figure in text. September, 1924	.25
15. Experimental Amoebiasis in Cats from Acute and Chronic Human Cases, by Edna Hannibal Wagener and Margaret D. Thomson. Pp. 267-280. September, 1924	.25
16. Binary Fission in <i>Oxyrrhis marina</i> Dujardin, by Richard P. Hall. Pp. 281-324, plates 26-30, 7 figures in text. December, 1924	.35
17. A Larval Acanthocephalid, <i>Centrorhynchus californicus</i> sp. nov., from the Mesentery of <i>Hyla regilla</i> , by Baymund Millzner. Pp. 325-327, 3 figures in text.	
18. <i>Megalodiscus ranophilus</i> sp. nov., a Trematode from the Rectum of <i>Rana pipiens</i> , by Baymund Millzner. Pp. 328-330, 2 figures in text. Nos. 17 and 18 in one cover. November, 1924	.25
19. On the Number of Chromosomes and the Type of Mitosis in <i>Endamoeba dysenteriae</i> , by Charles A. Kofoid and Olive Swezy. Pp. 331-352, plates 31-33, 1 figure in text. January, 1925	.30
20. Studies on <i>Paramecium</i> . I. Experiments on the Action of Various Endocrine Substances, of Liver, and of Glycogen on the Division Rate of <i>Paramecium</i> , by Gordon H. Ball. Pp. 353-383, 6 figures in text.	
21. Studies on <i>Paramecium</i> . II. The Behavior of a Conjugating Race of <i>Paramecium caudatum</i> , by Gordon H. Ball. Pp. 385-433, 5 figures in text. Nos. 20 and 21 in one cover. March, 1925	1.35
22. <i>Karyamoebina</i> Substituted for <i>Karyamoeba</i> with a Note on its Occurrence in Man, by Charles A. Kofoid and Olive Swezy. Pp. 435-436.	
23. <i>Cryptotermes hermsi</i> sp. nov., a Termite from Fanning Island, by Harold Kirby, Jr. Pp. 437-441, 12 figures in text. Nos. 22 and 23 in one cover. April, 1925	.25
24. Improved Micromanipulation Apparatus, by Charles Vincent Taylor. Pp. 443-454, plate 34, 2 figures in text. May, 1925	.25
Vol. 27. A Synopsis of the Amphibia of California, by Tracy I. Storer. Pp. 1-342, plates 1-18, 42 figures in text. June, 1925	\$4.50
Vol. 28. 1. Mitochondria and Golgi Bodies in <i>Endamoeba gingivalis</i> (Gros) Brumpt, by David Causey. Pp. 1-18, plates 1-3.	
2. Mitochondria in <i>Leishmania brasiliensis</i> , by David Causey. Pp. 19-28, plate 4. Nos. 1 and 2 in one cover. November, 1925	.40
3. Mitosis in <i>Ceratium hirundinella</i> O. F. M., with Notes on Nuclear Phenomena in Encysted Forms and the Question of Sexual Reproduction, by Richard P. Hall. Pp. 29-64, plates 5-9, 5 figures in text. November, 1925.	.55
4. The Cultivation of <i>Endamoeba gingivalis</i> (Gros), by Beatrice Fay Howitt. Pp. 65-126, plates 10-12, 4 figures in text. November, 1925	.80
5. The Behavior of <i>Endamoeba dysenteriae</i> in Mixed Cultures with Bacteria, by Charles A. Kofoid and Edna Hannibal Wagener. Pp. 127-154, plates 13-15.	
6. Studies of the Effects of Certain Drugs upon <i>Endamoeba dysenteriae</i> in vitro, by Charles A. Kofoid and Edna Hannibal Wagener. Pp. 155-166. Nos. 5 and 6 in one cover. November, 1925	.46
7. Some New and Some Previously Unreported Hydroids, Mainly from the Californian Coast, by C. McLean Fraser. Pp. 167-172, 7 figures in text. December, 1925	.25